Identifying dysregulated regions in amyotrophic lateral sclerosis through DNA accessibility outliers

Muhammed Hasan Celik[1,2], Julien Gagneur[3,4,5,6], Ryan G Lim[7], Jie Wu[8], Leslie M. Thompson[7,8,9,10,11],

Xiaohui Xie[1]

Department of Computer Science, University of California Irvine, Irvine, CA, USA
Center for Complex Biological Systems, University of California Irvine, Irvine, CA, USA
Department of Informatics, Technical University of Munich, Garching, Germany
Helmholtz Association - Munich School for Data Science (MUDS), Munich, Germany
Institute of Human Genetics, School of Medicine, Technical University of Munich, Munich, Germany
Institute of Computational Biology, Helmholtz Center Munich, Neuherberg, Germany
Institute for Memory Impairments and Neurological Disorders, University of California Irvine, Irvine, CA 92697, USA
Department of Biological Chemistry, University of California Irvine, Irvine, CA, USA
UCI MIND, University of California Irvine, Irvine, CA, USA
Department of Psychiatry and Human Behavior and Sue and Bill Gross Stem Cell Center, University of California Irvine, Irvine, CA, USA
Department of Neurobiology and Behavior, University of California Irvine, Irvine, CA, USA

Abstract

The high heritability of ALS and similar rare diseases contrasts with their low molecular diagnosis rate post-genetic testing, pointing to potential undiscovered genetic factors. DNA accessibility assays quantify the activity of functional elements genome-wide, offering invaluable insights into dysregulated regions. In this research, we introduced EpiOut, a computational toolbox to identify outliers in DNA accessibility. These outliers represent dysregulated regions where DNA accessibility uniquely diverges from the population baseline in a single or few samples. Annotation of accessible regions with histone ChIP-seq and Hi-C indicates that outliers are concentrated in functional loci, especially among promoters interacting with active enhancers. Across different omics levels, outliers are robustly replicated, and DNA accessibility outliers are reliable predictors of gene expression outliers and aberrant protein levels. For example, 59% of gene expression outliers can be linked to epigenetic changes. A comparison of promoter activity against gene expression reveals if aberration in the molecule phenotype is linked to pre-transcriptional or post-transcriptional regulation. Our findings demonstrate that the outlier detection paradigm can uncover dysregulated regions in rare diseases. EpiOut is open-sourced and freely available at github.com/uci-cbcl/EpiOut.

Introduction

Amyotrophic lateral sclerosis (ALS) is a rare neuromuscular degenerative disease affecting 0.6 to 3.8 per 100,000 people with a poor survival prognosis without a cure[1,2]. ALS is a complex disease where a single gene or pathway cannot explain the disease phenotype due to the heterogeneity of genetic causes and over 30 genes associated with ALS[1,3]. Meta-analysis and twin studies estimate the heritability of ALS disease at 61% (with 38%-78% confidence intervals) in sporadic cases (sALS), patients without a history of the disease in the family[4]. Despite the high heritability of ALS, 11 to 21% of patients[5,6] receive a diagnosis after genetic testing. The gap between high heritability and low diagnostic rate implies the existence of many undiscovered ALS-related genes.

There are large-scale sequencing efforts to discover the genetic bases of ALS[7,8]. These studies utilized GWAS, QTL, and differential expression analysis from a large cohort of samples to detect aberrations in ALS patients compared to control samples[7,9–11]. These statistical approaches successfully detected the most common factors (variants, genes, and pathways) associated with disease phenotype, yet detecting rare genetic factors remains challenging due to low statistical power. Outlier detection is a complementary statistical approach that uncovers aberrations specific to one or a few patients. Applying the outlier detection paradigm to transcriptomics data has revealed the dysregulation of many novel splicing and gene expression outliers[12–18]. This approach offers a promising direction for enhancing diagnostic rates, as it effectively models the heterogeneous nature of rare diseases.

The outlier detection approach has recently been applied to proteomics, and robust replication of aberrations across multiple omics data demonstrates the reliability of the outliers for disease diagnostics[19]. Expanding the outlier detection approach to epigenetics could provide further insight into the dysregulation of functional regions and their impact on gene expression in disease. Because epigenetic mechanisms, such as chromatin accessibility and enhancer/promoter activity, control gene expression. Despite the widespread use of chromatin accessibility assays such as ATAC-seq and DNase-seq enabling genome-wide investigation of the epigenetic landscape[20], the outlier detection approach has not been applied to chromatin accessibility data to the best of our knowledge.

Controlling for both known and latent confounding factors is crucial to exclude outliers resulting from technical artifacts or confounding factors during outlier detection. A linear autoencoder (LR-AE) is an effective statistical method for obtaining latent confounders from high-dimensional data. For example, PCA is a specific case of an LR-AE that can detect and correct for latent confounders better than alternative statistical methods[21], such as PEER[22]. However, PCA has its limitations as well. PCA minimizes the Euclidean distance between measured and reconstructed counts; thus, it is suboptimal for discrete read counts of omics data with high dispersion and uncertainty due to low-coverage[23].

The LR-AE with negative binomial objective function proposed by OUTRIDER[24] is better suited for read
counts and has superior outlier detection performance. However, this approach is much more
computationally expensive to optimize than PCA. OUTRIDER can be applied to transcriptomics data
with a dimensionality ranging from ten to fifteen thousand expressed protein-coding genes. However,
the LR-AE implementation of OUTRIDER may not scale to DNA accessibility with a dimensionality of
hundreds of thousands of genomic regions. Nevertheless, DNA accessibility outlier detection with an
LR-AE has not been previously explored.

Here, we present EpiOut, a software developed for DNA accessibility outlier detection. Our proposed
method takes read alignment files and accessible regions as input, performs ultra-fast read counting
per accessible region, detects outliers using an LR-AE with a negative binomial objective function, and
annotates outlier regions using ChIP-seq and Hi-C. Optimization of the decoder layer and dispersion
parameters requires solving a large number of independent convex problems. We significantly
accelerated the LR-AE using TensorFlow[25] by utilizing a vectorized variation of backtracking line search
for the dispersion parameters and L-BFGS for the decoder layer.

We applied EpiOut to DNA accessibility data from ALS samples to identify epigenetic aberrations that
might be linked to the disease phenotype. EpiOut pinpoints a small number of sample-specific loci as
outliers. These outliers are significantly enriched for functional regions as indicated by histone ChIP-
seq experiments, and these outlier regions potentially interact with each other based on the Hi-C data.
The accessibility outliers are reliable predictors of outliers detected in transcriptomic and proteomic
analyses. Moreover, we observed a substantial depletion of high-impact rare coding variants in gene
expression outliers when the promoters of these genes are accessibility outliers. In contrast, the
enrichment of high-impact variants was detected in cases where their promoters were not outliers.
Our results demonstrate that comparing epigenetic and transcriptomic outliers can offer valuable
insights into whether aberrations in gene expression are influenced by pre-transcriptional or post-
transcriptional regulation.

Results

In this study, we explored aberrant DNA accessibility in ALS using ATAC-seq experiments from the
AnswerALS dataset (Methods). This dataset comprises paired genomics, transcriptomics, accessibility,
and proteomics experiments from 253 individuals. With our novel method, EpiOut, we pinpointed
regions with abnormal DNA accessibility and investigated the molecular impact of these outliers by
comparing paired experiments across different omics levels. Our results highlight the biological
relevance and robustness of the identified accessibility outliers.

Detection of accessible regions

Detection of aberrant accessibility requires read counts for a set of accessible regions consistent across the individuals. First, we merged ATAC-seq reads of all samples into one meta-sample, then performed joint peak calling with MACS2[26] using the meta-sample to obtain a consistent set of accessible regions. We detected a total of 858,268 peaks before any filtering. Next, we counted the number of ATAC-seq reads overlapping with each accessibility region. The counting read from a large sample cohort is computationally intensive[14]. Thus, we developed EpiCount, an efficient read counter designed for accessibility data. EpiCount is twice as fast as the state-of-the-art counting tool[27] (Methods), using ~ 50 times less memory (Extended Data Fig. 1a,b) and fifteen times faster than the reported runtime of commonly used counting methods[14]. The large memory footprint of the counting method is a limiting factor for the embarrassing parallelization of counting across the samples; thus, EpiCount is highly parallelizable across multiple CPU cores. Lastly, we filtered accessible regions based on the read counts because many accessible regions often do not replicate across samples. We imposed a replication filter to these peaks, ensuring that the accessible regions were observed in at least 50% of the samples with a minimum of two reads and exhibited high accessibility (100 reads) in at least one sample (Extended Data Fig. 2a). Applying these filters yielded 114,428 accessible regions replicated across samples.

Outlier detection and benchmark

We hypothesize that major covariation between samples is primarily due to confounding factors. Therefore, eliminating this major covariation is essential in the proposed methodology to spotlight the rare aberrations potentially associated with the disease phenotype. To substantiate this premise, we examined the relationships between the principal components of accessibility data and the disease status of the samples. Fig. 1a illustrates the lack of clustering between samples by disease phenotype using the top two principal components of DNA accessibility. The top 25 principal components account for approximately 94% of the DNA accessibility variation between samples. However, none of these principal components show a significant association with the disease phenotype (Extended Data Fig. 3a,b). These findings indicate that major covariations in accessibility are likely attributable to confounding factors.

To evaluate the performance of the outlier detection methods, we employed an artificial outlier injection procedure previously proposed for detecting aberrant gene expression[24]. To create ground truth, we injected large aberrations, called artificial outliers, to read counts of ALS samples and then benchmarked the performance of tools to classify those artificial outliers on the area under the precision-recall curve (Methods) (Fig. 1c). In naive negative binomial, we estimated the mean of the

negative binomial test as a sample mean of read count per ATAC-seq peaks (Methods) and estimated dispersion with maximum likelihood, then ranked predictions by p-value based on the negative binomial test. The naive negative binomial model performs poorly with auPRC of $1.8 \pm 1\%$ because the expected read counts of the naive negative binomial model are not sample-specific. As an alternative outlier detection method, expected read counts per peak and sample can be estimated by PCA (Methods), and ranking predicted outliers by Z-score based on the expected and observed read counts of the PCA model have the performance of auPRC $9.7 \pm 0.9\%$. Our proposed method estimates expected read counts using an LR-AE (Fig. 1b). Expected read counts are incorporated into a negative binomial test as the mean parameter, and the dispersion is estimated based on the expected and observed counts (Methods). In our implementation of the LR-AE, the weights of the encoder and decoder layers are initialized with the rotation matrix of PCA. Then, the dispersion parameter is initially estimated using expected counts based on initial weights, and weights of the decoder layer are updated to maximize negative binomial likelihood using initial dispersion estimation. After the decoder layer optimization, we recalculate the dispersion estimation and apply the negative binomial test to estimate outliers. Also, it is critical to account for read coverage differences between samples (Extend Data Fig. 2b); thus, read counts are normalized for size factors[28] (Methods). We chose the optimal bottleneck size of LR-AE with hyperparameter tuning on the validation set (Extended Data Fig. 4). This approach outperforms the previous two methods by achieving an auPRC of $50.3 \pm 0.2\%$. An alternative LR-AE-based method, OUTRIDER, outperforms PCA and performs similarly with EpiOut (auPRC of $48 \pm 0.2\%$). Both methods employ an LR-AE with a negative binomial, a more proper distribution to fit counts data than PCA (Methods). Those results show that the estimation of dispersion and gene/sample-specific accessibility expectation followed by the negative binomial test is critical for outlier detection.

DNA accessibility is higher dimensional than gene expression because accessibility data may contain hundreds of thousands of accessible genomic regions. In contrast, gene expression data only contains around ten to fifteen thousand expressed protein-coding genes. Thus, the scalability of the outlier detection approach is essential to apply the method to high dimensional DNA accessibility data. Although OUTRIDER and EpiOut have similar auPRC scores, the autoencoder implementation of OUTRIDER is significantly slower than our proposed method (Fig. 1d). For example, outlier detection with EpiOut (745 ± 49 seconds) is 60 times faster than OUTRIDER ($44,874 \pm 200$ seconds). Our implementation is faster due to a couple of reasons. Firstly, we do not optimize the encoder layer of the autoencoder because we observed that the initial estimation of encoder weights with PCA is close to optimal, so further training of the encoder is unnecessary. Also, EpiOut only performs one alternating optimization step to fit the decoder layer and estimate dispersion. In contrast, OUTRIDER

performs multiple alternative optimization steps to estimate dispersion and train the encoder and

decoder layers. Lastly, the optimization of the decoder layer and the estimation of dispersion

parameters necessitate solving a multitude of independent convex optimization problems. We

efficiently approach these using a vectorized L-BFGS for the decoder layer and a vectorized and

bounded backtracking line search for dispersion estimation, all implemented in TensorFlow

(Methods).

We further integrated known confounding factors such as batch ID, sex, reported race, and ethnicity

into outlier detection (Methods). The contribution of each feature for outlier detection is summarized

in Fig. 1e. The naive negative binomial test estimates 76.5 ± 614 outlier regions per sample, and the

inclusion of latent confounding factors for expected read count estimation significantly reduces the

number of outliers to 18.3 ± 24. Known confounding factors such as batch ID, sex, race, and ethnicity

further reduce the number of an outlier to 14.2 ± 13.6 out of $114,428$ accessible regions per sample.

Thus, our method pinpoints only a handful of aberrant regions per sample.

As another benchmark, we compared cross-correlation between samples. Samples are highly cross-

correlated based on the raw read counts (Fig. 1f), and samples do not cluster by the disease phenotype

but rather by potential latent cofounders. Correcting for latent and known confounding factors and

normalization of read counts decorrelate samples and eliminate clusters due to potential batch effects

(Fig. 1g). Samples have an average cross-correlation of 96.1% based on the raw read counts. In

comparison, there is a 0% average cross-correlation after the proposed correction method ($P = 0$

based on the paired Wilcoxon test, Fig. 1h).

189
Fig. 1: Proposed outlier detection methodology and its benchmark (a) Samples do not form distinct
clusters based on their disease phenotype according to the top two principal components of DNA
accessibility. **(b)** The architecture of the proposed methodology for outlier detection (EpiOut). The
approach employs the negative binomial test. The mean parameter of the negative binomial
distribution is predicted with an LR-AE, which uses latent confounders obtained from data in addition
to reported known confounding as features to predict sample-specific expected accessibility. The
dispersion parameter is fitted with MLE using the observed and expected counts. **(c)** A precision-recall
curve shows the performance of alternative outlier detection methods. Methods were benchmarked
based on the classification accuracy of the injected artificial outlier. Specific cutoffs of models are
indicated with cross marks (an absolute Z-score of 2 for PCA, a p-value of 0.05 for naive negative
binomial, OUTRIDER, and EpiOut). **(d)** Runtime benchmark of outlier detection methods. **(e)**
Contribution of each component (such as latent and known confounding factors) of the model to
reduce the number of outliers per sample. **(f)** A Cluster heatmap displaying the cross-correlation of
samples based on the raw accessibility counts and **(g)** cross-correlation of samples after correction of

counts. **(h)** Distribution of cross-correlation between sample pairs before and after correction of
accessibility reads.

Functional annotation of accessible regions and outliers

The biological relevance of accessibility outliers remains ambiguous unless annotated with epigenetic
marks. Thus, we developed EpiAnnot, which annotated accessible regions for histone marks and 3D
chromatin interactions based on ChIP-seq and Hi-C experiments. EpiAnnot integrates publicly
available ChIP-seq and Hi-C for cell lines/tissues available in Roadmap Epigenomics[29] and ENCODE[30] or
from custom data sources. We annotated the previously identified accessible regions and outliers
using the H3K4me3, H3K27ac, and H3K4me1 histone marks observed in motor neurons. These
neurons were derived from the iPSCs of clinically healthy individuals and ALS samples[31]. Based on
histone marks, accessible regions are further classified as a promoter if the region has a histone mark
of H3K4me3 and is within 200 base pair (bp) vicinity of the annotated transcript start site, an active
enhancer if the region has both H3K27ac and H3K4me1 mark, and a poised enhancer if only H3K4me1
signal is present while H3K27ac mark is lacking (Fig. 2a). The largest category of accessible regions by
ChIP-seq marks is active enhancers ($n = 39{,}559$) with H3K27ac and H3K4me1 followed by thirst
largest category of 18,817 poised enhancers regions with only H3K4me1 mark (Fig. 2b). There are
16,446 promoter-like accessible regions with all three H3K4me3, H3K27ac, and H3K4me1 histone
marks. Both over-accessibility and under-accessibility outliers are significantly enriched ($P =$
9.2×10^{-3} and 7.1×10^{119} respectively based on Fisher's exact test) for promoter-like regions
because 28% of the outliers are located in promoter regions while only 14.3% of accessible regions
are promoters (Fig. 2b-d). Active enhancers are the second largest category of outliers ($n = 951$)
based on the annotation and are not significantly enriched or depleted for outliers ($P = 0.06$ for
under-accessibility and $P = 0.7$ for over-accessibility based on Fisher's exact test). Furthermore,
under-accessibility outliers for both poised enhancer and unannotated regions strongly depleted for
outliers as expected ($P = 1.9 \times 10^{-2}$, 10^{-31} respectively based on Fisher's exact test). Significant
enrichment of outliers in functional regions indicates the potential importance of accessibility outliers
on the molecular phenotype of ALS disease.

Another interesting observation is that outliers co-occur in the vicinity of each other (Fig. 2e). For
example, 7% of outliers have a second outlier in the 10 kilo-bp vicinity with an odds ratio of 688 ($P =$
0 based on the Fisher's exact test), and 33% of outliers have a second outlier in 1 million bp with the
odds ratio of 33 ($P = 0$ based on the Fisher's exact test). We repeated enrichment analysis between
promoter outliers and active or poised enhancer outliers and again observed significant enrichment.
16% of promoter outliers have at least one active enhancer outlier in 100 kilo-bp vicinity
(odds ratio $= 403$, $P = 0$ based on the Fisher's exact test), and 33% of promoter outliers have an

active enhancer outlier in 1 million bp vicinity (odds ratio $= 74$, $P = 0$ based on the Fisher's exact test). The co-occurrence of the outliers indicates the potential interaction between them. Hi-C experiments from motor neurons provide further evidence for the potential interaction between outliers. For example, the relatively high Hi-C score between the outlier promoter of the *ZFP41* gene and a distal enhancer outlier located ~ 110 kilo-bp upstream of the gene suggests a potential interaction between outliers (Fig. 2f). We calculated the Hi-C score between pairs of accessible regions and categorized regions by outlier status. We observed that outlier pairs ($P = 0$ based on the Mann–Whitney U test), including promoter-active enhancer pairs ($P = 4 \times 10^{-44}$ based on the Mann–Whitney U test), have higher interaction scores compared to a baseline where at least one of the regions in the pair is not an outlier (Fig. 2g). Hi-C contact scores are distance-dependent and decay according to power-law with increasing genomic distance. Thus, a higher interaction score could be possibly confounded by the co-occurrence of outliers in the vicinity of each other. To avoid this potential bias, we fit power regression on Hi-C scores using the outlier status feature and distance control variable (Methods). Outlier pairs have higher interaction scores ($P = 0$ based on the t-test) even after controlling for distance with power regression (Extended Data Fig. 5a). Even when we restricted our analysis to region pairs at least 100,000 bp apart, the Hi-C contact scores of outlier pairs still surpass the baseline of non-outlier pairs (Extended Data Fig. 5b). Overall, the co-occurrence of outliers in the vicinity of each other and higher Hi-C contact scores between outlier pairs indicate a potential interaction between outliers.

Fig. 2: (a) Accessible regions are annotated using ChIP-seq marks and gene annotation. Then, accessible regions are annotated as promoters, active or poised, and proximal or distal enhancers. **(b)** An upset plot shows the overlap size between H3K4me3, H3K27ac, and H3K4me1 histone marks and accessible regions **(c)** and outliers. **(d)** Enrichment of over-accessibility and under-accessibility outliers in the promoter, poised or active enhancer, and unannotated regions in terms of the **l**og of the odds ratio. **(e)** Cumulative odds of observing the second outlier in a region given that there is an outlier in the region (top) and the cumulative percentage of outliers with a second outlier in the vicinity (bottom) based on the distance between regions and the annotation. **(f)** The interaction between the outlier promoter of ZFP41 and ∼100 kilo-bp apart outlier distal enhancer is highlighted by the Hi-C track containing contact score between 5 kilo-bp long genomic bins. Coverage tracks for H3K4me3, H3K27ac, and H3K4me1 histone marks are also included in the figure. Red boxes indicate the outlier status of the accessible regions. **(g)** Boxplot shows the Hi-C contact scores distribution of non-outlier, outlier, and promoter-enhancer pairs. P-values are calculated with the Mann–Whitney U test and corrected with the Bonferroni correction for multiple testing.

DNA accessibility outliers influence gene expression and protein levels

Aberration in accessibility can impact downstream molecular phenotypes such as gene expression and protein levels. For example, the promoter of the *LCMT1* gene is an under-accessibility outlier in an ALS sample (Fig. 3a-c). We investigated a similar aberration pattern in the gene expression and protein level of *LCMT1* from the same sample's RNA-seq and mass-spectrometry experiments. We observed that there is a significant decrease in gene expression (Z-score $= -11.1$, P $= 2.8 \times 10^{-20}$ based on the Negative Binomial test). Moreover, the reduction in protein levels (Z-score $= -5.4$) is potentially linked to the under-accessibility of the promoter (Fig. 3d,e).

To understand the relationship between accessibility outliers and gene expression, we compared the variations in promoter accessibility with variations in gene expression across all samples (Fig. 3f). We observed a significant correlation between the fold changes ($log_2(FC)$) in accessibility at genes' promoters and the respective expression levels of these genes (Spearman's correlation coefficient=66%, P $= 0$). The high correlation indicates that aberrations in promoter accessibility potentially influence the aberrations in gene expression.

Moreover, we demonstrated that gene expression outliers can be systematically predicted from the promoters, proximal, and distal enhancer accessibility (Fig. 3g). Ranking promoter outliers by their absolute log fold change ($|log_2(FC)|$) to predict gene expression outliers achieve an auPRC of 14%. If the promoter is an outlier, there is a 41% chance (precision) that its gene is an expression outlier, and 26% of gene expression outliers have an outlier promoter (recall). Similarly, proximal enhancer outliers are highly predictive of gene expression outliers. Specifically, 35% (the recall at 19% precision) of gene expression outliers have at least one outlier proximal enhancer. Ranking genes based on the absolute log fold change of their outlier proximal enhancers achieves the performance of auPRC of 7.8%. We also ranked genes based on the maximum absolute fold change of accessibility outliers in 100 kilo-bp vicinity regardless of annotation of outliers and achieved 3.2% auPRC. We weighted the absolute log fold change of distal enhancers by ABC score[32] and obtained a score for each gene (Methods). The score calculated from distal enhancer outliers is also a reliable predictor of gene expression outliers and achieves 5.9% auPRC. We trained an explainable boosting machine[33] to predict gene expression outliers by combining features from promoters, proximal, and distal enhancers (Methods). The machine learning model achieved 21.4% auPRC. This analysis shows that 59% of the gene expression outliers (with 1% precision) can be linked to aberrations in accessibility because those gene expression outliers have at least one promoter, proximal, or distal enhancer outliers in the vicinity.

We focused on gene expression outliers with outlier promoters and investigated aberrations in their protein levels (Fig. 3h). We observed that the gene expression outliers with overly accessible

308 promoters have higher protein levels (average Z-score = 2.4, P = 1.3×10^{-7} based on the Mann–

309 Whitney U test), and genes with under-accessible promoters have lower protein levels (average

310 Z-score = -2.6, P = 2×10^{-11} based on the Mann–Whitney U test). Overall, we present the

311 biological significance of accessibility outliers on molecular phenotype by replicating outliers from

312 multiple omics levels.

Fig. 3: Prediction of gene expression outliers and aberrant protein levels from DNA accessibility. (a)
The outlier promoter of the *LCMT1* gene (in red) has a much lower ATAC-seq read coverage than
promoters of non-outlier samples. **(b)** Expected and observed accessibility in the promoter of LCMT1
gene across samples. The outlier sample is indicated with a red dot in the figure panels. **(c)** QQ-plot of
expected and observed p-values of the promoter. P-values were calculated using the negative
binomial test, and adjusted p-values were obtained using the Benjamini-Yekutieli false-discovery rate
(FDR) method. **(d)** The volcano plot shows the aberrant gene expression of *LCMT1* across samples with
Z-score and adjusted p-values, which were calculated using a negative binomial test and the
Benjamini-Yekutieli false-discovery rate (FDR) method. **(e)** The rank plot of samples by the Z-score
displays the aberrant protein levels of *LCMT1* across samples. **(f)** The precision-recall curve reports
the performance of a range of predictors to estimate gene expression outliers. Those predictors are
the absolute log fold change of promoter outliers (orange), the absolute log fold change of proximal
enhancers (green), the maximum absolute fold change of any outlier in 100 kilo-bp vicinity (blue), the
absolute log fold change of distal enhancers weighted by ABC score (red), explainable boosting
machine (EBM) trained with promoter, proximal-enhancer, and distal-enhancers features (purple). **(g)**
The scatterplot shows the correlation between the absolute log fold change of accessibility and
expression outliers. **(h)** The boxplot shows the Z-score distribution of proteins, which are categorized
by the outlier status of the promoter that transcripts them.

A comparison of DNA accessibility and gene expression reveals if aberrations in molecular phenotype are linked to pre-transcriptional or post-transcriptional regulation

The transcriptional synthesis and degradation rate control the gene expression (Fig. 4a). Pre-transcriptional regulation, such as aberrant promoter or enhancer activity, can lead to up or down-regulation of gene expression by altering the synthesis rate. Alternatively, post-transcriptional processes disrupted by genetic variants may lead to the degradation of transcripts via a non-sense-mediated decay. The potential regulatory process associated with aberrant gene expression can be revealed by comparing accessibility outliers against gene expression outliers.

To demonstrate this point, we investigated the enrichment of rare high-impact variants in different outlier categories. Fig. 4b presents the frequency of high-impact rare variants, such as splice acceptor, donor, nonsense, and frameshift variants, based on their SnpEff[34] consequences by the outlier type of the affected gene. High-impact variants are observed rarely and appear in fewer than 0.1% of non-outlier genes. Compared to the baseline of all expression outliers, high-impact genetic variants are significantly more prevalent in gene expression outliers if their promoters are not accessibility outliers. The enrichment holds for splicing acceptor ($n = 11$, $P = 5.4 \times 10^{-4}$ based on the Mann–Whitney U test calculated with bootstrapping) and splicing donor ($n = 11, P = 3.8 \times 10^{-4}$), nonsense ($n = 39, P = 2.5 \times 10^{-7}$), and frameshift variants ($n = 52, P = 1.1 \times 10^{-7}$). Remarkably, none of the gene expression outliers with aberrant promoter activity contain high-impact variants, indicating a significant depletion pattern.

Further investigation on both moderate-impact missense and high-impact variants on the protein levels presents a similar trend of depletion of variants in genes with aberrant promoter accessibility (Fig. 4c). 28.7% of expression outlier genes ($n = 101$) with aberrant protein levels ($|\text{Z-score}| > 1$) contain at least one high-impact variant or missense variant. When genes with promoter outliers were excluded, the enrichment of variants rose to 34.1% ($n = 85, P = 0$ based on the Mann–Whitney U test calculated with bootstrapping). The remaining expression outlier genes ($n = 16$) have promoters with aberrant accessibility, and none of these genes contain genetic variants with high or moderate consequences. The substantial depletion of high-impact variants in these genes ($P = 0$) indicates that their aberrant protein levels are potentially linked to aberrant promoter accessibility rather than coding variants.

Splicing is another mechanism that regulates gene expression[35]. Thus, we further explored the impact of exonic or intronic splicing-disrupting variants prioritized by AbSplice[36]. We detected 43 gene expression outliers containing at least one splicing-disrupting variant across different thresholds (Fig. 4d). However, none of these genes has an outlier promoter regardless of the threshold choice. In

367 contrast, subsetting gene expression outliers without aberrant promoters increases the prevalence of

368 splicing disturbing variants for the subset.

369 The results underscore that comparing accessibility outliers against gene expression outliers and

370 aberrant protein levels can identify whether aberration in the molecular phenotype is tied to pre-

371 transcriptional or post-transcriptional regulation.

Fig. 4: Overlap between accessibility outliers, gene expression outliers, and high-impact genetic variants. (a) Aberration in gene expression can result from dysregulation in pre-transcriptional regulation, such as aberrant promoter or enhancer activity, or dysregulation in post-transcriptional regulation, such as splicing or nonsense-mediated decay. **(b)** Percentage of genes with high-impact genetic variants by outlier categories. Gene expression outliers without outlier promoters are enriched, and gene expression outliers with promoter outliers are depleted for variants in each category. **(c)** Percentage of protein with a missense or high-impact variant where protein levels are at least deviant by $|Z\text{-score}| > 1$ by outlier categories. **(d)** Percentage of genes containing splicing-disrupting variants predicted by AbSplice by outlier categories.

Discussion

In this study, we introduced a computational toolbox for detecting and annotating DNA accessibility outliers, which are characterized as large aberrations in a few regions specific to a single or few samples. We applied the toolbox to ATAC-seq data from a clinical health and ALS sample cohort. Our methodology employs a negative binomial test for detecting outliers with statistical significance. The mean parameter of the negative binomial is fitted using a linear autoencoder (LR-AE), and the dispersion parameter is inferred based on observed and expected counts using maximum likelihood estimation. Our software, EpiOut, is highly scalable and can handle unknown confounding factors. While primarily designed for ATAC-seq, our toolbox is readily compatible with other accessibility assays, such as DNase-seq.

Our toolbox includes EpiAnnot, which annotates accessible regions for ChIP-seq marks. Using EpiAnnot, we found that outliers are enriched in functional regions, particularly promoters and active enhancers. Interestingly, outlier pairs tend to occur nearby, with many promoter outliers tied to active enhancer outliers within one million bp vicinity. This observation is supported by the relatively high Hi-C scores for outlier pairs in the vicinity, indicating potential interactions between these outliers.

By examining multiple omics levels, we found consistent replication of outliers. Accessibility outliers are associated with downstream biological processes such as gene expression and protein levels. In particular, a significant proportion of the gene expression outliers can be predicted from the aberrant accessibility of the promoter, proximal, and distal enhancer regions. Similarly, aberrant promoter activity is correlated with up and down-regulation of protein levels.

Analyzing the interplay between accessibility and gene expression outliers yields insight into whether aberration in gene expression originates from pre-transcriptional regulation, such as increased synthesis rate via higher promoter activity, or post-transcriptional regulation, such as splicing or nonsense-mediated decay. We observed substantial depletion of high-impact variants in gene

expression outliers if promoters of these genes are an accessibility outlier and conversely observed
enrichment of high-impact variants if their promoter is not an outlier.

Our epigenetic outlier detection methodology offers a novel avenue for studying rare diseases. We
have successfully adapted the outlier detection approach for ATAC-seq data and underscored that
ATAC-seq is a beneficial complementary assay for rare disease diagnostics. Detected outliers in ALS
samples are highly robust and consistently replicated across multiple omics levels. The continued
development and integration of the outlier detection approach with disease gene discovery
methodologies may ultimately lead to a more comprehensive understanding of the genetic factors
contributing to ALS. Such advancements could finally bridge the gap between the disease heritability
and the known catalog of ALS disease genes.

Method

AnswerALS Dataset
The multi-omics dataset, which includes ATAC-seq, RNA-seq, proteomics, and whole genome
sequencing (WGS), for Amyotrophic Lateral Sclerosis (ALS) was downloaded from the Answer ALS
portal (dataportal.answerals.org). The data contains 245 individuals diagnosed with ALS and 45
samples from clinically healthy controls. All samples had corresponding ATAC-seq and RNA-seq
experiments. 253 samples also have whole genome sequencing (WGS) data. For the scope of our
study, we restricted our analysis to the 253 samples that had paired data across three omics levels.
Additionally, 204 of these samples have proteomics data. In the proteomics-related analysis, we
subset and only used these samples with the proteomics data.

Peak Calling
We performed joint peak calling on ATAC-seq data to detect accessible regions across all samples
using MACS2[26]. First, we merged bam files from every sample into a unified bam file utilizing
SAMtools[37], and subsequently filtered out reads with a mapping quality (MAPQ) below 10. Duplicate
reads were retained after merging reads across samples. The default arguments of MACS2 were
used, except for the duplicate read filter. ATAC-seq peaks contained in the narrow peak bed file
generated by MACS2 were used in the downstream analysis.

Read Counting
We implemented an ultra-fast read counting algorithm for ATAC-seq described in Supplementary
Algorithm-1. The counting algorithm is optimized based on two primary assumptions: ATAC-seq

peaks are not overlapping and are separated by at least a gap longer than a read length; moreover,

both peaks and reads are sorted by the genomic coordinates. We ensure the first assumption by

jointly calling peaks as described above and collapsing any peaks closer to each other than the

minimum gap distance (default 200 bp); thus, an ATAC-seq read never intersects with two peaks

simultaneously. Since the bam file format is pre-sorted by genomic coordinates, ATAC-seq peaks are

sorted to guarantee the second condition before the counting step. The counting algorithm creates

two stacks of sorted peaks and reads, iterates over reads and peaks, and tracks the number of

overlaps (Supplementary Algorithm-1). The resulting runtime complexity of the counting step is

$O(r + p)$ where r is the number of reads, and p is the number of peaks. In the preprocessing step,

peaks are sorted, so the overall complexity is $O(p\ log(p) + r)$. However, since the number of reads

is much larger than the number of peaks ($r > p$), the algorithm practically behaves in linear runtime

in terms of the number of reads. The memory complexity of counting is linear in terms of the

number of peaks ($O(p)$) because reads are fetched iteratively per chromosome due to the indexable

file format of BAM.

Filtering Peaks by Replication Rate

We applied two filters to the ATAC-seq peaks detected by MACS2 to ensure consistent peak

replication across samples. First, we eliminated peaks with low coverage, specifically those with

fewer than 100 reads across all samples. As a subsequent criterion, any peak must be supported by a

minimum of 2 reads in at least 50% of all samples.

Size Factor Normalization

To account for coverage differences across the ATAC-seq experiments, we performed size factor

normalization on the read counts, a method initially proposed by DESeq2. The size factor s_i for a

sample i is defined as:

$$k_i^g = \left(\prod_{j=1}^{n} k_{ij} \right)^{1/n} \tag{1.1}$$

$$s_j = \text{median}\,\frac{k_{ij}}{k_i^g} \tag{1.2}$$

where k_{ij} represents the number of reads mapped to region j in sample i and k_i^g is the geometric

mean of the reads across regions for sample i. The median of k_{ij} to k_i^g ratios is defined as the size

factor s_i of sample i.

Outlier Detection

After normalizing the read counts of peaks with size factor normalization, we log-transformed the

normalized counts and centered them around zero by subtracting the mean normalized counts:

$$x_{ij} = n_{ij} - \bar{n}_j \tag{2.1}$$

$$n_{ij} = \log\left(\frac{k_{ij} + 1}{s_j}\right) \tag{2.2}$$

474   Autoencoder is applied on x_{ij} to calculate the expectation of normalized read counts ($\hat{x}_{ij}$). f_e is the

475   encoder function of the autoencoder, which takes observed normalized counts (x_{ij}) as input and

476   calculates major covariates. Major covariates are concatenated with known confounders to obtain

477   latent representation (h), which is decoded back to expected normalized counts ($\bar{x}_{ij}$) with the

478   decoder function f_d:

$$h = [f_e(x_{ij})|c_{known}] \tag{3.1}$$

$$\hat{x}_{ij} = f_d(h) \tag{3}$$

481   We used linear encoder and decoder functions. The encoder model uses a principal component

482   analysis (PCA), where the encoder weights (W_e) are the rotation matrix of PCA. The decoder weights

483   (W_d) are initialized with linear regression and further trained with the negative binomial objective:

$$f_e(x) = W_e x \tag{4.1}$$

$$f_d(h) = W_d h \tag{4.2}$$

486   Log normalized counts (x_{ij}) is transformed back to the original natural scale:

$$\hat{k}_{ij} = e^{\hat{x}_{ij} + n_j} s_j - 1 \tag{5}$$

488   The dispersion parameter is optimized with maximum likelihood estimation (MLE) where the

489   likelihood function:

$$L(r_j) = \prod_i NB(k_{ij}|\hat{k}_{ij}, r_j) \tag{6}$$

Fitting the dispersion parameter requires solving an independent convex problem for each peak. To

solve a large number of independent convex problems quickly, we implemented a vectorized

backtracking line search algorithm using TensorFlow. The dispersion parameter range is set between

a lower bound of 0.01 and an upper bound of 1,000 to avoid numerical stability issues and

overfitting.

P-value (P_{ij}) of each peak and sample is calculated with a two-sided negative binomial test:

$$P_{ij} = \min\left\{\frac{1}{2}, \sum_{k=0}^{\hat{k}_{ij}} NB(k_{ij}|\hat{k}_{ij}, r_j), 1 - \sum_{k=0}^{\hat{k}_{ij}-1} NB(k_{ij}|\hat{k}_{ij}, r_j)\right\} \tag{7}$$

498   Finally, p-values are corrected for multiple testing corrections to control the false discovery rate with

499   the Benjamini-Yekutieli method.

500   In addition to p-values, we report the log of fold-change or the log ratio of observed read counts to

501   expected read counts calculated by:

502
$$l_{ij} = \log_2(FC) = \log_2\left(\frac{k_{ij} + 1}{\hat{k}_{ij} + 1}\right) \tag{8}$$

503   and Z-score based on fold changes defined as:

504
$$Z\text{-score}_{ij} = \frac{l_{ij} - \mu_j^l}{\sigma_j^l} \tag{9}$$

We consider peaks as outliers if their adjusted p-values are smaller than 5% ($P_{adj} < 0.05$), their

absolute log fold changes are greater than 50% ($|\log_2(FC)| > 0.5$) and either their read counts (k_{ij})

or expected read counts ($\hat{k}_{ij}$) are at least 50.

Injection of artificial outliers

In order to benchmark the performance of outlier detection methods, we conducted a simulation

experiment using artificial outliers. We created an outlier mask M_{ij} for sample-peak pairs. A sample-

region pair is categorized as either an over-accessibility or under-accessibility outlier with a

probability of 0.01%:

513
$$f(x) = \begin{cases} 1, & p = 5 \ 10^{-4} \\ -1, & p = 5 \ 10^{-4} \\ 0, & otherwise \end{cases} \tag{10}$$

514   Log-normalized counts are updated using an artificial outlier mask:

515
$$x_{ij}^{inj} = x_{ij} + \sigma_j e^{\mathcal{N}(3,1)} M_{ij} \tag{11}$$

where the deviation in accessibility outlier is simulated by scaling the standard deviation of the peak

(σ_j) using a value sampled from a log-normal distribution parameterized by a mean of 3 and a

standard deviation of 1. Eq. 5 transforms log-transformed normalized injected counts (x_{ij}^{inj}) back to

counts in the natural count scale (k_{ij}^{inj}).

Outlier detection methods are evaluated by benchmarking their capability to predict the outlier

mask (M_{ij}) from the injected counts (x_{ij}^{inj}). The primary benchmark metric for this evaluation is the

area under the precision-recall curve (auPRC). We generated ten outlier masks for testing and one

additional mask for the validation set. Hyperparameter tuning was performed on the validation set

to identify the optimal bottleneck size of autoencoder-based models. The evaluation procedure was

executed ten times on the test folds, and the average auPRC performance and its standard deviation

are reported.

We evaluated four methods: naive negative-binomial test, PCA, OUTRIDER, and EpiOut. During the

precision-recall calculation, predictions for each method were primarily ranked using p-values,

except for PCA, which was ranked by its Z-score. The performance of each method at a specific

cutoff (either at the adjusted p-values of 0.05 or the absolute Z-score of 2) is indicated on the

precision-recall curve.

For the naive negative binomial test, we averaged counts across samples per peak to obtain the

mean parameter of the negative binomial distribution, $NB\left(k_{ij}^{inj}|\bar{k}_{ij}^{inj}, r_j\right)$. The dispersion is

estimated with MLE (Eq. 6).

To evaluate the PCA model, we used PCA to estimate the expected read counts, $\hat{k}_j^{inj}$. These

expected read counts are the reconstructed read counts of PCA. The number of principal

components retained was determined through hyperparameter tuning on the validation set.

Predictions were subsequently ranked based on the Z-score, as formulated in Eq. 9, using the

expected counts from PCA. We did not compute dispersion or perform a negative binomial test.

We ran OUTRIDER and EpiOut with their default parameters. The bottleneck size of all autoencoder-

based methods was selected based on the hyperparameter tuning on the validation set.

Runtime Benchmark

To measure runtime to call outliers with OUTRIDER and EpiOut, we subset peaks into groups of

10000, 25000, 50000, and 100000 and calculated runtime for each input size. 8 CPU cores are used

during the benchmark. The analysis is repeated 10 times. The average runtime and standard

deviation of runtime were reported. We ran each tool with bottleneck sizes of 5, 10, 50, and 100 in

each iteration to avoid runtime differences resulting from hyperparameter tuning. We reported the

total runtime across different bottleneck sizes.

Functional annotation of accessible regions with ChIP-seq

We introduced EpiAnnot, a software tool for functional annotation of accessible genomic regions

using ChIP-seq marks. If a specific ChIP-seq for a cell line or tissue is present in the Roadmap

Epigenomics[29] or ENCODE[30] databases, EpiAnnot can retrieve ChIP-seq data from these public data

sources. Additionally, users can annotate their accessible regions using their own custom ChIP-seq

data.

EpiAnnot can also attribute accessible regions to genomic features using a gene transfer format

(GTF) file. For instance, it annotates accessible regions as 5' UTR, TSS, etc. In scenarios where an

accessible region overlaps with the H3K4me3 histone ChIP-seq mark and TSS of a gene, EpiAnnot

designates these regions as promoters. Conversely, regions intersecting with the H3K4me1 histone

ChIP-seq mark are labeled as enhancers. When the histone mark is available, enhancers overlapping

with H3K27ac are categorized as active; otherwise, they are poised. EpiAnnot also annotates

enhancers as proximal or distal based on their distance from genes. Specifically, enhancers within a

gene body or located up to 10,000 bp upstream or 2,000 bp downstream from a gene are tagged as

proximal. All others are marked as distal enhancers.

By using EpiAnnot with H3K4me3, H3K27ac, and H3K4me1 histone ChIP-seq marks from in vitro

differentiated motor neurons from clinically healthy and ALS samples[31], we annotated accessible

regions from ATAC-seq experiments and used the GTF file of GENCODE v38. The histone ChIP-seq

data was downloaded from ENCODE.

Enrichment of Outlier Pairs in the Proximity

To investigate the potential interaction between nearby outlier pairs, we quantified the pairs of

accessible regions within predefined distances, irrespective of the outlier status of regions. Then, we

applied Fisher's exact test to calculate the enrichment of outlier pairs, using a contingency table

structured with the outlier statuses of accessible regions in the pair. We repeated this analysis for

10,000, 100,000, 500,000, and 1,000,000 bp distances. The odds ratio was also calculated to

highlight the likelihood of the region being an outlier, given the outlier status of the nearby region.

Annotation of Interactions with Hi-C

To delve deeper into the interactions between outliers, we annotated their interactions based on Hi-

C contacts using EpiAnnot. We downloaded public Hi-C data from motor neurons differentiated from

iPSCs[31] from ENCODE and computed interaction scores between accessible regions using EpiAnnot.

For the interaction analysis, the genome is partitioned into bins of 5,000 bp. The interaction

between a first source region (s) and a target second region (t) is determined by the highest Hi-C

score between these bins or their immediate neighbors:

$$582 \qquad h_{s \to t} = \max\{HiC_{score}(bin_i, bin_j), i \in \{t-1, t, t+1\}, j \in \{s-1, s, s+1\}\} \qquad (12)$$

583 We utilized EpiAnnot to compute the ABC score, which signifies potential interactions between

584 regions informed by Hi-C scores. The ABC score is defined as:

$$585 \qquad \text{ABC}_{s \to t} = \frac{h_{s \to t} * k_t}{\sum_{i \in vicinity(s)} h_{s \to i} * k_i} \qquad (13)$$

586 where the numerator represents the Hi-C score between regions multiplied by the accessibility of

587 the target region ($k_t = \sum_i k_{ij}$) and the denominator normalizes this value for all region pairs

588 associated with the source region.

589 Aberrant Gene Expression Prediction from DNA Accessibility

Gene expression outliers are called with OUTRIDER using the DROP pipeline. Those gene expression

outliers constitute the ground truth labels. In order to predict gene expression outliers from

accessibility outliers, we trained an explainable boosting machine (EBM). The EBM model was

trained and tested with 5-fold cross-validation. The model aggregates features from promoters and

enhancers as input to predict the probability of a gene being categorized as an expression outlier.

For predicting a gene's outlier status, the model considers the following features: log fold change, p-

value, and outlier status of the promoter, maximum absolute log fold change of the proximal

enhancer outliers, and the ABC-score weighted absolute log fold change of the distal enhancer

outliers. The score for the weighted distal enhancer is computed as follows:

$$599 \qquad \text{enhancer}_{distal} = \sum_{enhancer \in g} |ABC_{promoter \to enhancer} * log_2(FC_{enhancer})| \qquad (14)$$

600 Depletion of high-impact variants in Genes with Promoter Outlier

We assessed the depletion of high-impact variants in the gene expression outliers where promoters

of these genes are also outliers. Variants are annotated with SnpEff v4.3[34] according to their

consequences for genes. Only rare variants with a frequency of 0.1% are in the analysis. Variant

frequencies are downloaded from the gnomAD database[38]. Consequence categories used for

depletion analysis include stop codon, frameshift, acceptor, donor site dinucleotides, and missense

variants. AbSplice was used to classify splicing-disrupting exonic or intronic variants into high,

medium, or low-impact categories, using thresholds of 0.2, 0.1, and 0.05, respectively, as advised by

the authors. We determined the proportion of genes affected by each variant category based on the

outlier status of the genes and their associated promoters. We repeated the enrichment analysis

100 times by randomly subsetting genes in each iteration for bootstrapping. We used the Mann–

Whitney U test to measure statistical significance based on the proportions calculated with

bootstrapping. Moreover, we analyzed proteomics data to quantify the proportion of proteins

impacted by high-impact or missense variants. Z-scores for the proteomics data were obtained with

PROTRIDER.

Source Code

All the analyses in the paper are implemented in the reproducible snakemake[39] format available at

github.com/uci-cbcl/ALS-accessibility-outliers-paper. Epiout, the Python package for outlier

detection and annotation, is available at **github.com/uci-cbcl/EpiOut**.

Acknowledgments

This project is funded by a grant from the Robert Packard Center for ALS Research at Johns Hopkins

and grants from the National Science Foundation [IIS-1715017, DMS-1763272].

References

1. Hardiman, O. *et al.* Amyotrophic lateral sclerosis. *Nat. Rev. Dis. Primer* **3**, 1–19 (2017).

2. Longinetti, E. & Fang, F. Epidemiology of amyotrophic lateral sclerosis: an update of recent
literature. *Curr. Opin. Neurol.* **32**, 771–776 (2019).

3. Masrori, P. & Van Damme, P. Amyotrophic lateral sclerosis: a clinical review. *Eur. J. Neurol.* **27**,
1918–1929 (2020).

4. Al-Chalabi, A. *et al.* An estimate of amyotrophic lateral sclerosis heritability using twin data. *J.*
Neurol. Neurosurg. Psychiatry **81**, 1324–1326 (2010).

5. Grassano, M. *et al.* Systematic evaluation of genetic mutations in ALS: a population-based study.
J. Neurol. Neurosurg. Psychiatry **93**, 1190–1193 (2022).

6. Renton, A. E., Chiò, A. & Traynor, B. J. State of play in amyotrophic lateral sclerosis genetics. *Nat.*
Neurosci. **17**, 17–23 (2014).

7. Van Rheenen, W. *et al.* Project MinE: study design and pilot analyses of a large-scale whole-
genome sequencing study in amyotrophic lateral sclerosis. *Eur. J. Hum. Genet.* **26**, 1537–1546
(2018).

8. Baxi, E. G. *et al.* Answer ALS, a large-scale resource for sporadic and familial ALS combining clinical
and multi-omics data from induced pluripotent cell lines. *Nat. Neurosci.* **25**, 226–237 (2022).

9. van Rheenen, W. *et al.* Common and rare variant association analyses in amyotrophic lateral
sclerosis identify 15 risk loci with distinct genetic architectures and neuron-specific biology. *Nat.*
Genet. **53**, 1636–1648 (2021).

10.   Gregory, J. M., Fagegaltier, D., Phatnani, H. & Harms, M. B. Genetics of Amyotrophic Lateral
Sclerosis. *Curr. Genet. Med. Rep.* **8**, 121–131 (2020).

11.   NeuroLINCS Consortium *et al.* An integrated multi-omic analysis of iPSC-derived motor
neurons from C9ORF72 ALS patients. *iScience* **24**, 103221 (2021).

12.   Frésard, L. *et al.* Identification of rare-disease genes using blood transcriptome sequencing
and large control cohorts. *Nat. Med.* **25**, 911–919 (2019).

13.   Mertes, C. *et al.* Detection of aberrant splicing events in RNA-seq data using FRASER. *Nat.*
Commun. **12**, 529 (2021).

14.   Yépez, V. A. *et al.* Detection of aberrant gene expression events in RNA sequencing data.
Nat. Protoc. **16**, 1276–1296 (2021).

15.   Li, X. *et al.* The impact of rare variation on gene expression across tissues. *Nature* **550**, 239–
243 (2017).

16.   Marwaha, S., Knowles, J. W. & Ashley, E. A. A guide for the diagnosis of rare and
undiagnosed disease: beyond the exome. *Genome Med.* **14**, 23 (2022).

17.   Jenkinson, G. *et al.* LeafCutterMD: an algorithm for outlier splicing detection in rare diseases.
Bioinformatics **36**, 4609–4615 (2020).

18.   Kremer, L. S. *et al.* Genetic diagnosis of Mendelian disorders via RNA sequencing. *Nat.*
Commun. **8**, 15824 (2017).

19.   Kopajtich, R. *et al.* Integration of proteomics with genomics and transcriptomics increases
the diagnostic rate of Mendelian disorders. 2021.03.09.21253187 Preprint at
https://doi.org/10.1101/2021.03.09.21253187 (2021).

20.   Yan, F., Powell, D. R., Curtis, D. J. & Wong, N. C. From reads to insight: a hitchhiker's guide to
ATAC-seq data analysis. *Genome Biol.* **21**, 22 (2020).

21.   Zhou, H. J., Li, L., Li, Y., Li, W. & Li, J. J. PCA outperforms popular hidden variable inference
methods for molecular QTL mapping. *Genome Biol.* **23**, 210 (2022).

22.   Stegle, O., Parts, L., Piipari, M., Winn, J. & Durbin, R. Using probabilistic estimation of
expression residuals (PEER) to obtain increased power and interpretability of gene expression
analyses. *Nat. Protoc.* **7**, 500–507 (2012).

23.   Townes, F. W., Hicks, S. C., Aryee, M. J. & Irizarry, R. A. Feature selection and dimension
reduction for single-cell RNA-Seq based on a multinomial model. *Genome Biol.* **20**, 295 (2019).

24.   Brechtmann, F. *et al.* OUTRIDER: A Statistical Method for Detecting Aberrantly Expressed
Genes in RNA Sequencing Data. *Am. J. Hum. Genet.* **103**, 907–917 (2018).

25.   Abadi, M. *et al.* TensorFlow: Large-Scale Machine Learning on Heterogeneous Distributed
Systems.

26.   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).

27.   Stovner, E. B. & Sætrom, P. PyRanges: efficient comparison of genomic intervals in Python.
Bioinforma. Oxf. Engl. **36**, 918–919 (2020).

28.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

29.   Kundaje, A. *et al.* Integrative analysis of 111 reference human epigenomes. *Nature* **518**, 317–
330 (2015).

30.   Dunham, I. *et al.* An integrated encyclopedia of DNA elements in the human genome. *Nature*
489, 57–74 (2012).

31.   Zhang, S. *et al.* Genome-wide identification of the genetic basis of amyotrophic lateral
sclerosis. *Neuron* **110**, 992-1008.e11 (2022).

32.   Fulco, C. P. *et al.* Activity-by-contact model of enhancer–promoter regulation from
thousands of CRISPR perturbations. *Nat. Genet.* **51**, 1664–1669 (2019).

33.   Nori, H., Jenkins, S., Koch, P. & Caruana, R. InterpretML: A Unified Framework for Machine
Learning Interpretability. Preprint at https://doi.org/10.48550/arXiv.1909.09223 (2019).

34.   Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff. *Fly (Austin)* **6**, 80–92 (2012).

35.   Daguenet, E., Dujardin, G. & Valcárcel, J. The pathogenicity of splicing defects: mechanistic
insights into pre-mRNA processing inform novel therapeutic approaches. *EMBO Rep.* **16**, 1640–
1655 (2015).

36.   Çelik, M. H. *et al.* Aberrant splicing prediction across human tissues. Preprint at
https://doi.org/10.1101/2022.06.13.495326 (2022).

37.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–
2079 (2009).
38.   Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in
141,456 humans. *Nature* **581**, 434–443 (2020).
39.   Köster, J. & Rahmann, S. Snakemake--a scalable bioinformatics workflow engine. *Bioinforma.*
Oxf. Engl. **28**, 2520–2522 (2012).