An amplicon panel for high-throughput and low-cost genotyping of Pacific

oyster

Ben J. G. Sutherland[1,2], Neil F. Thompson[3], Liam B. Surry[2], Krishna Reddy Gujjula[4], Claudio D.

Carrasco[4], Srinivas Chadaram[4], Spencer L. Lunda[5], Christopher J. Langdon[6], Amy M. Chan[7], Curtis A.

Suttle[7,8,9,10], Timothy J. Green[2]

[1] Sutherland Bioinformatics, Lantzville, BC, Canada V0R 2H0

[2] Faculty of Science and Technology, Vancouver Island University, Nanaimo, BC, Canada V9R 5S5

[3] Pacific Shellfish Research Unit, Agricultural Research Service, United States Department of

Agriculture, Hatfield Marine Science Center, Newport, OR 97365 USA

[4] ThermoFisher Scientific, 2130 Woodward Street, Austin, TX 78744 USA

[5] Oregon State University, Department of Microbiology, 226 Nash Hall, Corvallis, OR 97331

[6] Oregon State University, Coastal Oregon Marine Experiment Station, Hatfield Marine Science Center,

2030 SE Marine Science Dr., Newport, OR 97365

[7] Department of Earth, Ocean and Atmospheric Sciences, The University of British Columbia, Vancouver,

BC, Canada V6T 1Z4

[8] Department of Microbiology and Immunology, The University of British Columbia, Vancouver, BC,

Canada V6T 1Z3

[9] Department of Botany, The University of British Columbia, Vancouver, BC, Canada V6T 1Z4

[10] Institute for the Oceans and Fisheries, The University of British Columbia, Vancouver, BC, Canada,

V6T 1Z4

26 *Author for correspondence: BJGS

27 Current Address:

28 Sutherland Bioinformatics, Lantzville BC, Canada, V0R 2H0

29 Sutherland.Bioinformatics@protonmail.com

32 Running Title: *Pacific oyster amplicon panel*

Abstract

Maintaining genetic diversity in cultured shellfish can be challenging due to high variance in individual reproductive success, founder effects, and rapid genetic drift, but is important to retain adaptive potential and avoid inbreeding depression. To support broodstock management and selective breeding in cultured Pacific oysters (*Crassostrea* (*Magallana*) *gigas*), we developed an amplicon panel targeting 592 genomic regions and SNP variants with an average of 50 amplicons per chromosome. Target SNPs were selected based on elevated observed heterozygosity or differentiation in Pacific oyster populations in British Columbia (BC), Canada. The use of the panel for parentage applications was evaluated by genotyping three generations of oysters from a breeding program in BC (n = 181) and a set of families that were selected for Ostreid herpesvirus-1 (OSHV-1) resistance from the Molluscan Broodstock Program, a Pacific oyster breeding program in Oregon, USA (n = 136). Population characterization was evaluated using collections of wild, naturalized, farmed, or hatchery oysters sampled throughout the Northern Hemisphere (n = 190). Technical replicate samples showed high genotype concordance (97.5%; n = 68 replicates). Initial parentage analysis found instances of suspected pedigree and sample handling errors, demonstrating the panel's value for quality control in breeding programs. Suspected null alleles were identified in parentage datasets and were found to reduce assignment success. Null alleles were largely population dependent, suggesting population-specific variation impacts target amplification. By taking an iterative approach, null alleles were identified using existing data without the need for pedigree information, and once null alleles were removed, assignments increased to 93.0% and 86.0% of possible assignments in the two breeding program datasets. A pipeline for analyzing the amplicon sequence data from sequencer output, *amplitools*, is also provided.

Keywords: amplicon panel, aquaculture, *Crassostrea* (*Magallana*) *gigas*, genetic diversity, Pacific oyster, parentage assignment

Introduction

Sustainable aquaculture depends on the effective characterization, preservation, exchange and use of
genetic resources (Guo, 2009). High resolution genomic tools (e.g., high-density genetic maps, SNP chips,
chromosome-level reference genomes) enable genome-wide association studies to identify markers that can
be used for marker-assisted or genomic selection (Boudry et al., 2021). Low resolution genetic tools (e.g.,
amplicon panels) enable high-throughput genotyping at low cost (Meek & Larson, 2019). Lower-cost
applications can allow scalable use, such as in fisheries management through parentage-based tagging (e.g.,
Beacham et al., 2018), or in aquaculture to assign individuals back to putative family when mixed without
physical tagging (Allen et al., 2020). Amplicon panels can provide rapid turnaround time (Arbelaez et al.,
2019), and increased flexibility to use lower quality input DNA (Csernák et al., 2017); in general, these
panels have been described as reliable, scalable, moderately easy to analyze, and cost-effective (Meek &
Larson, 2019). Cost-effectiveness can bring genetic advances to small or medium sized operations not able
to afford high-density genotyping tools (Boudry et al., 2021; Delomas et al., 2023).

Pacific oyster *Crassostrea* (*Magallana*) *gigas* (Thunberg, 1793; Salvi & Mariottini, 2017), is one
of the most valuable aquaculture species globally (Botta et al., 2020). Oyster aquaculture is a significant
economic driver and food source for many developed and developing nations (Martínez-García et al., 2022).
However, production and growth of the industry outside of China has stagnated due to disease, regulatory
issues, and other factors (Botta et al., 2020). Large-scale mortality events have been a significant burden on
the industry, where disease may be caused by specific or combined protozoan, bacterial, or viral pathogens,
as well as by multifactorial or unexplained causes (King et al., 2019). Abiotic stressors such as salinity
shifts, elevated temperatures, or heat waves can interact with these biotic stressors, increasing negative
impacts (Green et al., 2019; King et al., 2019).

Environmental changes more generally are also challenging cultured and wild oysters. Ocean
acidification negatively impacts shell formation and maintenance (Barros et al., 2013; Barton et al., 2012;
Watson et al., 2009), and extreme weather has led to massive mortality events in recent years. For example,
an atmospheric river and freshwater influx into the San Francisco Bay (California, USA) coincided with a
near 100% mortality event of the highly abundant population of wild Olympia oyster *Ostrea lurida* (Cheng
et al., 2016). An unprecedented extreme heatwave in the Pacific Northwest during summer 2021 resulted
in large-scale negative impacts to wild and naturalized shellfish in the area (Raymond et al., 2022). With a
multitude of stressors and impediments to industry stability and growth, shellfish farmers in the Pacific
Northwest consider oyster health and selective breeding to be important scientific adaptation strategies to
increase crop resilience (Green et al., 2023; Thompson, 2023). As such, improvements to selective breeding
infrastructure and genomic resources are valuable to Pacific oyster breeders, as recently advocated for

eastern oyster *C. virginica* (Allen et al., 2020).

Maintenance of genetic diversity is essential to shellfish breeding. Although standing
polymorphism levels are very high in shellfish (Plough, 2016), where a SNP is expected every 40 bp in *C.*
gigas (Hedgecock et al., 2005; Sauvage et al., 2007), rapid reductions in genetic diversity can occur in
cultured lineages (Evans et al., 2004; Gurney-Smith et al., 2017; Hedgecock & Sly, 1990; Xiao et al., 2011),
with losses compounding over time, leading to significant concerns for breeders (e.g., Li et al., 2007). High
genetic load occurs through high fecundity and likely high mutation rates, which can produce severe
inbreeding depression affecting growth and survival (Evans et al., 2004). Although inbreeding depression
may be counteracted by crossing divergent lines to achieve heterosis (Hedgecock & Davis, 2007;
Hedgecock et al., 1995), long-term retention of genetic diversity, and maintenance of hatchery lineages is
important for broodstock programs (Carlsson et al., 2006; Gurney-Smith et al., 2017). Furthermore,
retaining standing genetic diversity is vital for rapid adaptation (Barrett & Schluter, 2008) and selective
breeding (Guo, 2009).

A rapid, inexpensive, automatable, and easily analyzed genotyping tool is needed for the Pacific
oyster. Existing genomic resources for the Pacific oyster include high quality reference genome assemblies
(Peñaloza et al., 2021; Qi et al., 2021) and genetic maps (Gutierrez et al., 2018; Li et al., 2018; Yin et al.,
2020), as well as SNP arrays of high (134K targets; Qi et al., 2017), medium (41K; Gutierrez et al., 2017),
and low density (384; Lapègue et al., 2014). However, due to significant costs and required expertise,
smaller or medium-scale breeding operations may not be able to benefit from these resources (Boudry et
al., 2021). Until present, there has been no low-density amplicon panel for the Pacific oyster. In addition to
lower cost per individual, amplicon panels can also be used to genotype non-target variants within an
amplicon window, potentially expanding its utility across populations and time, as long as primer sites
remain intact and functional. This longevity and malleability may be particularly advantageous to Pacific
oysters given their high genetic diversity (Hedgecock et al., 2005; Sauvage et al., 2007), global scale of
culture (Boudry et al., 2021; Martínez-García et al., 2022), including genetically diverse large-scale
commercial hatcheries (Sutherland et al., 2020), and temporal fluctuations in allele frequencies that occur
due to sweepstakes reproductive success (Hedgecock, 1994; Hedgecock & Pudovkin, 2011; Sun &
Hedgecock, 2017).

Here we describe the development of a 592 target SNP panel designed using previously identified
high heterozygosity and high differentiation markers from naturalized Pacific oysters in British Columbia
(BC), Canada, which are similar to other global populations derived from the Japan translocation lineage
(Sutherland et al., 2020). We test the panel on samples from Canada, France, Japan, and China, as well as
cultured populations from the United Kingdom, the United States, and Canada. We further evaluate and
demonstrated the panel's effectiveness for genetic parentage assignment using single SNP targets through

analyzing three generations of the Vancouver Island University (VIU) breeding program, and a closely
related set of families bred for viral resistance from Oregon State University's Molluscan Broodstock
Program (MBP). All design amplicons and target variant identities and sites are provided here, and the panel
is available via the commercial provider (ThermoFisher Scientific). All analytic code is available in
amplitools, a repository designed to move from sequencer output to parentage analysis (see *Data*
Availability). Details on the latest version of the panel's target file is available in the accompanying
repository *amplitargets* (see *Data Availability*), expected to be updated over time. Collectively, with these
outputs we aim to facilitate uptake and therefore bring rapid gains in Pacific oyster breeding and aquaculture.
131

Methods

Data sources and marker selection
Plink and Variant Call Format (VCF) outputs from Sutherland et al. (2020), a restriction-site associated
DNA sequencing (RAD-seq) study of Pacific oysters from the Northern Hemisphere, were obtained from
the authors and uploaded to FigShare (see *Data Availability*). Genotypes were read into R (R Core Team,
2023) using the read.PLINK function of adegenet (v.2.1.5; Jombart & Ahmed, 2011) and data were
converted from genlight format to genind, genepop, and hierfstat formats for analysis using adegenet and
hierfstat (v.0.5-10; Goudet & Jombart, 2022).

The pre-filtered dataset was limited to only include naturalized collections from British Columbia
(BC), the focal region of the study, which includes the locations Hisnit (HIS), Pendrell (PEN), Pipestem
(PIP), and Serpentine (SER); for details on collection sites see Sutherland et al. (2020). Within the BC
dataset, per locus global minor allele frequency (MAF) was recalculated, and SNP variants with MAF <
0.01 were removed. Per locus observed heterozygosity (H_{OBS}) and average F_{ST} (Weir & Cockerham, 1984)
were calculated using adegenet and pegas (v.0.12; Paradis, 2010), respectively. SNPs were excluded from
the RAD-seq dataset when $H_{OBS} > 0.5$ in an effort to avoid duplicated genomic regions, and remaining
SNPs with the highest H_{OBS} (n = 300), or the highest F_{ST} (n = 300) were selected for panel design. Several
additional SNP variants of interest were included based on previous detection as private alleles in Deep Bay,
BC (n = 15) or Guernsey, UK (n = 5) farmed populations (Sutherland et al., 2020). Nine SNPs were
redundant between the F_{ST} and H_{OBS} selected lists; 611 unique SNPs were identified for panel design (Table
S1).
152

Panel design and chromosomal locations
Flanking 200 bp sequences on either side of selected SNPs were obtained from the contig-level reference
genome (GCA_000297895.1; Zhang et al., 2012) that was used in marker discovery (Sutherland et al., 2020)

using a bed file derived from the marker discovery VCF and the bedtools function getfasta (Quinlan & Hall,

2010). All required code and instructions for this process are provided (see *Data Availability*,

ms_oyster_panel). Target amplicon window sequences (401 bp), variant positions, and reference and

alternate alleles (Additional File S1) were submitted to the AgriSeq primer design team for the Ion Torrent

system (ThermoFisher Scientific). Targets were passed through the AgriSeq quality control process

(ThermoFisher Scientific), using the GCA_000297895.1 genome. Those passing QC were then submitted

to the primer design phase. In the design phase, oligo candidates were generated, scored, filtered, and the

optimal oligo pair for each target was selected to be included in the final panel. The selected oligos were

checked *in silico* for specificity and sensitivity of the intended target regions using the *C. gigas* reference

genome GCA_000297895.1. The resultant amplicon panel is available commercially through

ThermoFisher Scientific (SKU A58237 AGRISEQ PACIFIC OYSTER PANEL).

As the marker discovery RAD-seq genotyping (Sutherland et al., 2020) used a contig-level genome

assembly (Zhang et al., 2012), the chromosomal positions of the targeted 401 bp sequences were

subsequently determined by mapping them against a chromosome assembly (GCF_902806645.1; Peñaloza

et al., 2021) using bowtie2 (Langmead & Salzberg, 2012) in end-to-end mode allowing multiple alignments

(-k 6). The target sequences were also aligned with bwa mem (Li, 2013). Amplicons were categorized as

mapping to a single or multiple positions, and those mapping to multiple positions were removed during

analysis. Positions of amplicon targets along chromosomes were visualized using ggplot2 (Wickham, 2016).

Panel testing

The panel was tested for use in characterizing different populations by genotyping oyster collections from

2017 and 2018, as described by Sutherland et al. (2020) and shown in Table 1, including from a farm (Deep

Bay, BC, DPB), commercial hatcheries (Guernsey, UK, GUR; and China, QDC), and wild (Japan, JPN;

China, CHN) or naturalized sources (Pendrell Sound, BC, PEN; France, FRA). Although these same

populations and collection years were used for marker discovery (Sutherland et al., 2020), the panel testing

was done on alternate individuals not used for discovery, except for JPN and CHN since additional samples

were not available. Notably, JPN and CHN were not used for marker selection, as described above.

The panel was tested for utility in parentage analysis by genotyping samples from three generations

of the Vancouver Island University (VIU) breeding program, and oyster families from two MBP generations

that were selected based on the presence or absence of a SNP marker on chromosome 8 (CHR8) for

resistance to the Ostreid herpesvirus-1 (OsHV-1) (Divilov et al., 2023). The breeding program samples were

provided with information about their respective generation, but pedigree information was not used until

after assignments were created and the initial analysis reviewed. These analyses are described below.

DNA from the VIU and MBP breeding program samples was extracted using the Monarch Genomic

DNA Purification kit (NEB) using the enzymatic cleanup tissue extraction protocol with proteinase K and

RNase A and quantified using a BioSpectrometer (Eppendorf). All other samples were extracted using the

BioSprint (QIAGEN) method and diluted to 25 ng/µl and stored at -20ºC. Samples were further normalized

to 10 ng/µl in 25 µl volumes and submitted to ThermoFisher Scientific laboratories in Austin, TX for

AmpliSeq library preparation using the AgriSeq™ HTS Library kit (ThermoFisher Scientific) with 16

cycles of amplification and IonCode barcode labelling. Libraries were sequenced using Ion 540 chips on

an Ion Torrent S5 (ThermoFisher Scientific). Some samples had low sequence yields, and so a portion of

the poorly amplifying libraries were synthesized again from undiluted source samples, then sequenced at

both diluted and supplied sample concentrations. Replicate samples with sufficient genotyping rates per

sample were used to evaluate repeatability of library preparation, sequencing, and genotyping (more details

provided below). The MBP samples were submitted for genotyping as described above but conducted at

the Institute de Biologie Intégrative et des Systèmes (IBIS) at Université Laval. At both facilities, target

variants were scored using the Torrent Suite software VariantCaller plugin (ThermoFisher Scientific) using

the original designed target SNP file (i.e., Cgig v.1.0 hotspots; see *Data Availability*). The custom pipeline

amplitools (see *Data Availability*) was used to filter multi-locus genotypes files for target variants and

convert to genepop format. Below, the population genetic samples and the VIU samples are referred to as

the 'pilot dataset', and the MBP CHR8 samples are referred to as the 'MBP CHR8 dataset'.

Repeatability, filtering, and population genetic analyses

The pilot dataset genepop file was imported into R using adegenet and analyzed using the custom pipeline

simple_pop_stats using instructions provided in *ms_oyster_panel* (see *Data Availability*). Technical

replicate pairs with at least a 50% genotyping rate per individual were used to determine the proportion of

concordant genotypes. Downstream analyses used the replicate individual with the highest genotyping rate.

Samples, then loci, were filtered to keep those with less than 30% missing genotypes. Monomorphic loci,

and loci that were previously determined to be align to more than one position in the genome (*see above*)

were also removed. Although markers were characterized for deviation from Hardy-Weinberg proportions

in each population using the *hw.test* of pegas, deviating markers were not removed, only flagged.

The dataset was converted to a genlight object using dartR (v.2.0.4; Gruber et al., 2018). Inter-

individual relatedness between individuals was estimated using the Ritland statistic (Ritland, 1996) as

implemented in related (v.1.0; Pew et al., 2015). First-degree relatedness values were estimated using

known full-sib pairs in the VIU dataset (*see below*). Pairs of individuals within a population that had

relatedness values greater than the lowest value for known first-degree relatives were subjected to a process

of elimination where one individual per pair was removed until no pairs remained above the threshold. The

purged first-degree relative dataset was then analyzed by principal components analysis (PCA) using the
function *glPca* in adegenet, retaining PC1-4, and population averaged F_{ST} (Weir & Cockerham, 1984) was
calculated for any population with more than 15 retained individuals using hierfstat (v.0.5-11; Goudet &
Jombart, 2022). The full dataset with putative first-degree relatives included was used to identify private
alleles in each genetic grouping using poppr (v.2.9.3; Kamvar et al., 2014), where Japan, France, and
Canada (Pendrell) were pooled together, and the three VIU generations were all pooled together. The
hatchery or farm collections were kept as separate groupings in this analysis. The full dataset was used to
calculate per locus mean F_{ST} using the *Fst* function of pegas, and per locus heterozygosity using the
summary function of adegenet. The data for the MBP CHR8 families was used as a test dataset and filtered
as above, removing individuals then loci with more than 30% missing genotypes, and removing
monomorphic loci. This dataset was used for parentage analysis to compare with the VIU parentage analysis
in order to assess the utility of the panel in other programs (*see below*).

Parentage assignment
The pilot dataset was reduced to only the VIU breeding program samples (now referred to as the VIU
dataset), filtered again to keep loci with MAF > 0.01 and to remove monomorphic loci, then converted from
genind format to rubias format (Moran & Anderson, 2019) using the function *genepop_to_rubias* of
simple_pop_stats (see *Data Availability*). Data preparation and analysis for parentage closely followed
vignettes of *close-kin mark recapture-sim* (CKMR-sim; Anderson, 2024). Using allele frequencies
estimated from the VIU dataset, a simulated set of parent-offspring, full-sib, half-sib, and unrelated
relationships were created to determine the power for the amplicon panel genotypes to resolve different
relationship types, and to estimate expected log likelihood values for the different relationship categories.
The dataset was checked for closely matching potential duplicate samples. Parent-offspring relationships
and full-sibling relationships in offspring and parents were evaluated using VIU samples (i.e., VIU_F2 vs.
VIU_F1). Siblings of true parents were included in the analysis to test the power of the amplicon panel to
resolve relationships in the present of closely related possible parents (e.g., 'aunts' or 'uncles').

Following analysis and grouping of individuals into putative families, pedigree information was
provided and compared against the genetic parentage analysis relationships. Offspring that failed to assign
to genotyped parents were tallied to assess false negatives, and incorrect assignments were tallied as false
positives. The number of putative parents was not limited to two, but rather all assignments over the log
likelihood cutoff (*logl* > 5) were retained in the results. This cutoff was based on the total number of
comparisons and expected false positive rates as per the CKMR-sim tutorials. The percentage of the total
possible assignments that could have been made was calculated for each family in both parentage analyses,
and the percentage of the total assignments made that were false positives was also calculated. Suspected

pedigree or handling errors were inspected alongside genetic sibship analyses and breeding records to

reconcile conflicting relationship information. These errors, when resolvable, were corrected.

Following the initial analysis described above, empirically identified trios (i.e., offspring and two

parents) with strong support were identified (i.e., both parents assigned with log likelihood values ≥ 10,

only two parental assignments with log likelihood scores above the set cutoff). When a set of parents were

supported by more than one trio, the trios were kept as the empirical trio results to be used to identify

unexpected genotypes potentially indicating null alleles. The empirical trios were used to infer expected

genotypes of offspring per locus and family, and the observed offspring genotypes were compared to

expected offspring genotypes to tally the number of unexpected genotypes per locus. Finally, the parentage

analysis with all samples was re-done as described above but with the following filtered datasets: (1) all

loci passing filters; (2) loci with fewer than four offspring with incompatible genotypes. Percent

completeness of assignments, and percent of assignments that were false positives were calculated for each

of the above three parentage approaches. This same approach was also carried out independently with the

MBP CHR8 dataset, and the problematic loci identified with each dataset were compared against each other

to determine whether the VIU and the MBP CHR8 dataset had the same loci that were exhibiting potential

null alleles or otherwise unexpected offspring genotypes.

274   **Results**

SNP selection for marker design and chromosomal positions

The input ddRAD-seq dataset (Sutherland et al., 2020) contained 366 Pacific oyster samples genotyped at

16,492 SNPs (single SNP retained per RAD-tag). It had been filtered to retain SNPs that were present in at

least 70% of individuals in all 15 populations, with global minor allele frequency (MAF) ≥ 0.01, and not

significantly out of Hardy-Weinberg equilibrium. The dataset was then limited to naturalized British

Columbia (BC) samples, retaining 122 individuals from four populations. The BC-only dataset was filtered

to keep SNPs with MAF ≥ 0.01, retaining 13,438 SNPs. Marker design candidates were selected for

elevated heterozygosity (n = 300; H_{OBS} = 0.41-0.5) or elevated F_{ST} between naturalized populations within

BC (n = 300; F_{ST} = 0.046-0.178), where nine markers were common to both lists. An additional 20 markers

were selected due to being previously identified as private alleles (see *Methods*; Figure S1; Additional File

S2). Of the 611 submitted SNPs for design, 19 were excluded, and 592 were designed into the panel (i.e.,

Cgig_v.1.0; see Additional File S3 for marker names, contig identifiers, positions and alleles).

Of the 592 design sequences, 499 (84.2%) aligned using bowtie2 to a single location in the

chromosome assembly, 55 aligned to multiple locations, 12 aligned singly to non-chromosome scaffolds,

and 45 did not map. However, when aligned with bwa mem, 25 of the design sequences aligned to multiple

locations, and these were different than the bowtie2 putative multi-mappers (see locus identifiers in

Additional File S4). Due to this uncertainty, the bowtie2 putative multi-mappers were removed from the

present analysis, but remain in the targets file. Each chromosome had an average (± s.d.) of 49.9 ± 26.3

single-mapping amplicons per chromosome (range: 15-83; Figure 1). Several significant gaps in

chromosome coverage with single mapping markers were observed (Figure 1).

Panel genotyping and repeatability

The pilot dataset comprised 543 samples including technical replicates and controls, or 370 unique samples

and 32 negative controls. Samples were genotyped at 592 amplicons across two sequencing runs (i.e., two

sequencing chips). Seven amplicons were uncalled in both chips, leaving 585 genotyped amplicons

remaining. The median of the average amplicon depth per individual was 172.5x and 180.1x for the two

chips, whereas negative controls were 0.90x and 5.3x, respectively (Figure S2). On the second chip, three

of the 19 negative control wells had mean marker coverage greater than 40x, suggesting some

contamination in these wells.

Technical replicate sample pairs (n = 68 pairs) showed average (± s.d.) genotype concordance of

97.5 ± 4.2% for SNPs genotyped in both samples (mean = 460 SNPs genotyped in both samples of the pair,

with a total of 31,303 genotypes). The most frequent discordant genotypes between technical replicates

were homozygous alternate and heterozygous genotypes (n = 552; 1.76%), and less frequent were

homozygous reference and heterozygous genotypes (n = 153; 0.49%) or least frequent as homozygous

reference and homozygous alternate genotypes (n = 4; 0.01%). Ten replicate sample pairs had more than

5% of the markers discordantly called; on average, these pairs had 359.5 loci typed in both samples, whereas

the remainder pairs on average had 477.7 loci. One replicate pair had genotype discordance above 10%,

and this pair was only genotyped at 342 loci. Therefore, samples with lower genotyping rates also had lower

concordance.

Filtering and population genetics

The 370 oysters in the pilot dataset were comprised of 181 oysters from the VIU breeding program, 71 from

aquaculture programs (i.e., hatcheries or farms), and 63 and 55 from naturalized and wild sources,

respectively (Table 1). After filtering samples for genotyping rate (GR > 70%), 312 samples were retained

(Table 1; Figure S3). Filtering loci for genotyping rate (GR > 70%) removed 111 loci (Figure S4) leaving

474 loci remaining. Removing monomorphic loci removed an additional 43 loci, resulting in the retention

of 431 loci. Any remaining of the 55 multi-mapping loci as determined by the bowtie2 alignment were also

removed at this stage, dropping 22 additional loci and retaining 409 loci. Retained markers showed a

bimodal H_{OBS} distribution (Figure 2A). The upper H_{OBS} grouping (H_{OBS} > 0.25; n = 246) has 221 loci that

were included in the panel due to high H_{OBS}. The lower H_{OBS} grouping (H_{OBS} < 0.25; n = 163) is mainly

comprised of loci included due to high F_{ST} (i.e., n = 132), as well as those included based on presence as

private alleles (n = 11). Per locus H_{OBS} and F_{ST} values for the 409 retained loci are shown in Additional File

S5. Loci Hardy-Weinberg proportions (HWP) were inspected for wild (JPN, CHN) and naturalized

populations (PEN, FRA) finding 15, 24, 26, and 26 variants out of HWP, respectively. This includes 11 loci

that were out of HWP in two populations, and one marker in three populations. These markers were not

removed from the analysis but can be referenced for future consideration in Additional File S6.

The filtered dataset (n = 312 individuals, 409 loci) was then inspected for highly related pairs of

individuals within each population (Figure S5). Outlier pairs with elevated relatedness were observed in

most populations. The known full-sibs in the largest cross event in the VIU population used (i.e., oyster

family run 5; OFR5) were used to calculate average relatedness values within each family, and the 14

families had an average pairwise relatedness value with the Ritland statistic of 0.43, and the average

minimum relatedness value per family was 0.26, which was used as a cutoff to remove highly related

individuals specifically to reduce family effects on the PCA and between population F_{ST}. This resulted in

the removal of most of samples from the hatchery or farm populations (i.e., VIU, DPB, and GUR) for PCA

and F_{ST} analyses. Populations with fewer than 15 individuals were not used for F_{ST} calculation.

A PCA with putative first-degree relatives removed found general overlap of all populations except

CHN, which were separated from the rest of the samples across PC1 (3.2% variance explained) and PC2

(2.2% variance explained), respectively (Figure S6). Similarly, the Japan translocation lineage (i.e., JPN,

PEN, FRA) had low F_{ST}, with the lower range of the 95% confidence intervals overlapping zero (i.e., no

differentiation) and the upper range reaching F_{ST} = 0.012 (Table S2). By contrast, CHN had higher

differentiation, for example JPN-CHN 95% C.I. F_{ST} = 0.023-0.048).

Private alleles were empirically identified in the genotyped dataset (with putative first-degree

relatives included) at the regional level (i.e., JPN lineage, VIU, DPB, GUR, CHN). In total, 20 private

alleles were observed, including some with high frequency. Three markers were included in the panel

because they contained private alleles in DPB (Sutherland et al., 2020); although different samples were

used here, two of these loci remained after filters and also were observed to be private alleles in this dataset,

including locus 39139 (eight occurrences), and locus 538043 (six occurrences). Similarly, locus 336452,

included in the panel as a private allele within GUR in Sutherland et al. (2020) was also private in this

dataset (nine occurrences). One high frequency private allele was observed in the VIU samples with eight

observations (locus 590965). However, this marker would have been designed based on presence in other

BC samples, and so is likely just a rare allele that has been enriched in the VIU broodstock program. A

private allele was also observed for each of CHN and JPN with five observations (see full list in Table S3).

Parentage assignment in the VIU dataset

The panel was tested on families from the oyster family run 5 (OFR5) spawning event of the Vancouver

Island University (VIU) Pacific oyster breeding program. OFR5 involved 15 crosses using 18 unique

parents that originated from cross events OFR1 and OFR2, including nine dams and nine sires, all of which

were genotyped. One OFR5 broodstock parent, BR22, was missing. In total, 91 OFR5 offspring (81.3%)

were retained after filters (Table 1, Table 2).

An initial analysis of the data found likely pedigree or handling errors. Four samples were identified

that were suspected handling errors in which the offspring had the incorrect family label. These were evident

using parent-offspring, or sibship analyses. Second, family OFR5-15 was found to be a mixed family, with

two separate sets of parents (i.e., the expected parents BR18 and BR10, or the parents of OFR5-14, BR24

and BR2; Table 2). Third, a pedigree error was identified where the sire of family OFR5-2 was BR11 and

not BR2 (family 1-12). These were all corrected before subsequent analyses.

To identify potential null alleles in the amplicon loci, reliable two-parent assignments were

identified and used to identify incompatible offspring genotypes based on the observed parental genotypes

(see Methods). Reliable assignments were defined as those trios with exactly two parents assigned above

an elevated cutoff. Putative families were estimated from the reliable assignments, keeping parental pairs

as a putative family when more than one observation of a parental pair occurred. These identified trios

included 37 offspring from 11 families. Within these trios, 36 loci were observed with at least four offspring

with incompatible genotypes, and 95 loci with at least two offspring with incompatible genotypes (Figure

S7). Parentage datasets were constructed to evaluate the impact of these loci exhibiting potential null alleles:

(i) keeping all filtered loci (n = 364 loci retained); or (ii) removing loci with four or more incompatible

genotypes (n = 328 loci retained). An improvement in assignments occurred and a reduction of false

positives by removing loci with at least four incompatible offspring genotypes, and so this approach was

taken (Table 3). In this dataset, the OFR5 broodstock was genotyped with an average of 313.2 loci per

individual (Table 2), and the offspring at an average of 307.0 loci per individual (Additional File S7).

Within the 328-loci dataset, 160 of 172 potential assignments were made (93.0% complete), and

eight false positives occurred (4.8% of assignments made; Table 3; Table 2). The greatest number of false

positives occurred in families F10 and F12, where maternal or paternal siblings were assigned in addition

to the correct assignment. All parent-offspring relationships are shown in Figure 3, and assignments with

accompanying log likelihood ratios are shown in Additional File S8.

Parentage assignment in the MBP CHR8 dataset

An independent dataset comprised of families from the Molluscan Broodstock Program (MBP CHR8) was

genotyped to further evaluate the panel performance for parentage assignment. This dataset was expected

to be more challenging for the panel given the longer captive breeding history and higher relatedness among

broodstock. The MBP spawn was comprised of 16 families from MBP cohort 27 (seventh generation of

selective breeding) including 25 unique broodstock (nine dams and 16 sires). All parents were genotyped,

but one parent (65_8F) was removed due to low genotyping rate. In total, 111 offspring (100.0%) were

retained after filtering (Table 1, Table 4). The dataset was comprised of 576 scored loci, but after filtering

the excess missing data (missing $\geq$ 30%), 519 loci were retained. An additional 174 monomorphic loci were

removed, leaving 345 loci. Removing the multi-mapping loci from the bowtie2 analysis resulted in an

additional 21 loci being removed, leaving 324 loci remaining. The MBP CHR8 family distributions of the

likelihoods of simulated relationships were similar to those of the VIU OFR5 families (Figure S8). Further,

the observed heterozygosity per locus of the VIU OFR5 family and MBP CHR8 shows many loci with

similarly elevated H_{OBS} in both datasets (Figure 2B).

An initial analysis found suspected handling errors, including three offspring with incorrect family

identifiers. Offspring were labeled as family/offspring number. Incorrect offspring identified were as

follows: 102/8 was from family 106 and not 102; 106/1 was from family 113 and not 106; and 113/8 was

from family 102 and not 113. These were sequential in DNA extraction and library preparation, and so the

series of unexpected assignments likely reflects a handling error in DNA extraction or library preparation

and not in sampling. These errors were corrected by providing new identifiers to the unexpected offspring

to match their correct families, and then the same approach to identify putative null alleles was taken as

described above for OFR5 (see above and Methods). Empirically determined trios with strong evidence

were identified in the MBP CHR8 families, including 38 offspring from nine different families. Using these

trios, 73 loci were identified that had at least two offspring with incompatible genotypes based on expected

genotypes inferred from parents, and 35 loci with at least four offspring with incompatible genotypes. The

two datasets (i.e., all filtered loci: 324 loci; remove when $\geq$ 4 incompatibilities: 289 loci), were used for

parentage analysis. The dataset with 289 loci resulted in the most assignments, but also the most false

positive assignments (Table 3, Table 4). In this dataset, the parent generation (potential broodstock) were

genotyped on average at 283.4 loci per individual (Table 4), and the offspring on average at 284.5 loci per

individual (Additional File S7). This dataset had 185 of 215 possible assignments (86.0% complete), and

17 false positives occurred (8.4% of assignments). All identified parent-offspring relationships are shown

in Figure 4, and assignment details in Additional File S8.

To determine whether a common set of loci were exhibiting genetic incompatibilities (i.e., putative

null alleles) in the VIU OFR5 and the MBP CHR8 families, overlap was inspected between the incompatible

loci. Of the 36 loci that were incompatible in VIU ($\geq$4 incompatibilities) and the 35 incompatibility loci in

MBP, only seven were found common in both datasets (i.e., 19% or 20% of the total VIU or MBP loci,

respectively). This low overlap occurred even though 35 of 36 of the incompatible VIU loci were present

in the MBP CHR8 dataset (before incompatibles were removed), and 28 of 35 of the incompatible MBP

CHR8 loci were present in the VIU dataset (before incompatibles were removed; Additional File S9). This

suggests that putative null alleles identified are largely population specific.

Discussion

This work demonstrates the effectiveness of the Pacific oyster Cgig_v.1.0 amplicon panel for characterizing

genetic variation and conducting parentage assignment, as well as provides the analysis framework to

support additional use of the panel, as well as future development of the panel in subsequent design

iterations. The unique characteristics of the Pacific oyster, including high levels of genetic polymorphism,

likely led to challenges faced in the analysis, such as the presence of numerous, population-specific null

alleles. Initial results were sufficient to detect suspected handling or pedigree errors, including in breeding

program samples from populations not part of the panel design, and with careful filtering steps, parentage

assignment success was able to be improved. We expect that the tools provided here will bring advances to

Pacific oyster breeding programs for both small and large operations.

Maintaining genetic diversity and supporting breeding program development

Several genetic approaches may be used to increase the resiliency of cultured Pacific oysters. First,

maintaining overall genetic diversity retains variation required for rapid adaptation (Barrett & Schluter,

2008) and avoids significant inbreeding depression that can impact growth and survival (Evans et al., 2004;

Wada & Komaru, 1994). Second, artificial selection methods including family-based selection (i.e.,

breeding families that show most favourable phenotypes as selection targets), marker-assisted selection (i.e.,

selecting broodstock based on genotypes at loci associated with a phenotype), and genomic selection (e.g.,

breeding based on multilocus genotype relationships derived from a training set of phenotyped individuals),

can be used to improve phenotypic targets of breeding populations (Boudry et al., 2021). Genomic or

marker-assisted selection cannot be undertaken without genotyping. Similarly, maintaining diversity or

separation of distinct breeding lines benefits from the ability to empirically measure diversity and confirm

family relationships.

Maintaining genetic diversity in Pacific oyster breeding populations is important as rapid reductions

in genetic diversity can occur (Hedgecock & Sly, 1990), as can small sustained reductions that accumulate

over time (Carlsson et al., 2006; Varney & Wilbur, 2020; Xiao et al., 2011). Using genetic markers, the

effective population size and other measures of diversity such as allelic richness can be directly monitored

in the breeding population (Gurney-Smith et al., 2017; Hedgecock & Sly, 1990). Allelic richness may be a

more appropriate measure to monitor diversity than observed heterozygosity or per-individual inbreeding

coefficients (Sutherland et al., 2023; Varney & Wilbur, 2020; Yu & Guo, 2004), in part because these other

metrics can be skewed by crossing divergent genetic lines for heterosis, an approach sometimes applied

within the bivalve aquaculture industry. When reduced diversity is detected, solutions may be needed to

replenish diversity (Li et al., 2007). Some countries, including Canada, have naturalized populations of

conspecifics that can be incorporated within breeding programs (Sutherland et al., 2020) to increase

diversity and avoid inbreeding. Having tools to empirically measure genetic diversity is important for

breeding programs (Gurney-Smith et al., 2017) for a range of industry scales (Boudry et al., 2021).

Family-based selection or crossing of divergent lines both require maintenance of separate genetic

groups, and benefit from confirming crosses and pedigree information to correct errors that may occur

during animal husbandry or data recording. This has been a subject of concern for many decades

(Hedgecock & Davis, 2007), and has led to the development of high-resolution melt curve genotyping

panels (Sun et al., 2015; Zhong et al., 2013). Pedigree errors have been identified in the present work and

elsewhere (Hedgecock & Pan, 2021) that would not have been identified without the use of a low-cost,

effective genotyping panel, and if or when left unchecked, could slow progress in family-based breeding

programs (Hedgecock & Pan, 2021). The amplicon panel presented here provides a rapid and lower-cost

tool to evaluate the parentage of crosses and correct pedigree errors, thereby increasing the precision of

family-based selection protocols.

There are many potential sources of error that can disrupt maintenance of separate genetic groups

and directional family-based selection. They may occur during spawning, larval rearing, or growth phases,

and can be caused by incorrect labeling, contaminating families (i.e., inadvertent mixing), as well as

handling and record keeping errors. Many of these errors disrupt pedigree relationships, and if left

uncorrected, can compound over time and impede directional selection progress (Hedgecock & Davis,

2007). As presented here, we identified several different sources of error: (1) mislabeled single offspring;

(2) a full-sib family contaminated with a second full-sib family (i.e., the preceding family); (3) handling or

recording errors for the sire of two families within one spawn event; (4) handling errors after sampling

during laboratory preparation. Importantly, as discussed by Hedgecock and Pan (2021), these errors could

impede selective breeding gains were they not detected, and may increase the mismatch between the genetic

breeding values and the phenotypic outcomes of trials.

The amplicon panel may bring additional benefits to aquaculture programs. Currently, the

application of genomic selection in shellfish aquaculture is limited by high genotyping costs per individual

(Delomas et al., 2023). One framework, described and recently evaluated *in silico* for Pacific oyster by

Delomas et al. (2023), is to use whole-genome resequencing of parents, lower-density amplicon panel

genotyping of offspring, and imputation to enable more cost-effective application of modern genomic

selection methods. Furthermore, field deployments for phenotyping trials can be conducted in common
garden replicates, rather than using separate nets that confound spatial and family effects, by using genetic
parentage analysis enabled by lower cost genetic marker panels (Allen et al., 2020). Finally, future versions
of the panels can be designed to include putatively functional markers for marker-assisted selection, for
example for shell colour or body size, or traits which can increase survival from thermal stress (Raymond
et al., 2022), or for field survival likely linked to disease resistance (e.g., OsHV-1; Divilov et al., 2023).
498
Future directions
The current panel performed well on the populations included in this analysis. Future work should continue
to evaluate the panel in other populations to determine any ascertainment bias and to assess how broadly
applicable the tool can be. As an example, in the present study, moving to different populations that were
not involved in design, such as MBP's CHR8 families, resulted in a specific loss of SNPs selected due to
high F_{ST} in BC, whereas the SNPs selected for high heterozygosity were more likely to pass filters. The
high F_{ST} markers are the most likely candidates to remove in future panel iterations. In this project, most
genotyped individuals originated from populations that were included in marker discovery and comprised
oysters from the Japan translocation lineage, China, and several commercial aquaculture hatcheries
(Sutherland et al., 2020). In new populations, other variants within the panel's amplified sequences (i.e.,
not the target designed SNP) may prove more powerful for parentage, and these could be characterized via
adjustments to the data analysis post sequencing, and not requiring any physical changes to the current
panel. This is beneficial because physical changes to the primer composition would be more laborious and
costly. This is a particular advantage for amplicon-based sequencing that is not possible in array or high-
resolution melt curve genotyping.
Using non-target variants from the existing amplicons may provide improvements to the panel for
parentage purposes without changes to the primer pool. This is apparent given that some markers within
Cgig_v.1.0 in the pilot study had low heterozygosity and minor allele frequency, including those selected
due to high differentiation potential or presence as private alleles. For technical replicates, low genotyping
rate pairs had the lowest percent concordance, suggesting that poor genotyping rate is not only an issue for
missing data, but the data that is present has a higher chance of miscalls. Therefore, emphasis should be
placed on retaining high quality throughout all phases of tissue collection, library preparation, and
sequencing to minimize the loss of data or the generation of poor quality data. End-users of the panel should
determine levels of missing data that are allowable for their uses. As missing data was more prevalent in
specific collections, it is likely that this is driven by DNA quality or quantity; other sample collections with
similar genetic backgrounds to the high failure collections had more consistent and complete genotyping,
suggesting that the missing data is not largely due to primer failures due to variants at binding sites but

rather due to DNA quality or quantity. Including higher value markers (i.e., higher MAF) may improve

resolution of secondary assignments to full sibs of the true parent (i.e., false positives) without a significant

increase in false negatives (i.e., missing assignments). Increased statistical power to assign genetic

relationships may also be realized through a microhaplotype approach (Baetscher et al., 2018), which is an

active area of research for the current panel.

Other advances to the panel are possible but would require modifying the primer suite by adding

new or removing existing amplicons. The uniformity of coverage across chromosomes could be improved,

filling in gaps observed in the current work, and thereby improving the potential for genomic selection

(Delomas et al., 2023). As discussed above, when additional variants with putatively functional roles (or

close linkage with causal variants) are identified, such as the recently identified SNP on chromosome eight

associated with increased field survival associated with OsHV-1 outbreaks (e.g., Divilov et al., 2023), these

will likely be valuable to add to the panel for breeding purposes. Although rigorous documentation and

standardization of locus names is required, the malleability of the amplicon panel is a major strength, as

loci can be added or dropped by changing the primer composition of the primer pool. Once it can be more

conclusively determined which loci are true multi-mapping loci, for example through detecting more than

two alleles in diploid individuals with microhaplotypes, these loci will be permanently removed from the

panel. Furthermore, removing loci that exhibit null alleles would strengthen the use of the panel, however,

the observation of the two different breeding programs largely showing different loci that had putative null

alleles negates the ability to screen for null allele loci across a broad geographic scale without including

this in an initial panel development protocol. In addition, given the high mutation rate of Pacific oysters,

new mutations are likely to regularly emerge that could disrupt primer binding sites, producing new null

alleles occurrences. Thus, users of the current panel are strongly encouraged to screen for null alleles within

the target population of interest as removing these loci improves genetic parentage assignment success, as

was demonstrated in the present work.

The amplicon panel provides a new tool for emerging Pacific oyster selection programs, but it is

also important to also consider the ease of which the data can be analyzed. Although the tab-delimited

genotype data produced by the genotyping platform may be easier to handle for those with less

bioinformatics experience than high volume sequence data, moving from this output to the formats required

for parentage or population genetic analysis does require moderate skills in data manipulation, quality

control, and data management. With this in mind, we developed *amplitools*, a repository that is designed to

make the use of this panel more accessible, including bash and R scripts for converting from raw variant

calling output from standard sequencer software to a standard format used by numerous genetic analysis

programs with a wide range of analysis types being available.

Some aspects of documentation within breeding programs will support the use of *amplitools*, such

as not including third generations when running parent-offspring analyses, given the potential for

grandparents to be miss-assigned as parents. Including parental sex information may also be beneficial for

analysis, although this has not yet been implemented in *amplitools*. Additionally, we developed the

associated repository *amplitargets* to keep and version control target files for genotyping, as well as any

notes regarding changes between versions of SNP target files. Although the primer panel, *amplitools* and

amplitargets are currently only implemented for the AgriSeq platform (ThermoFisher Scientific), the source

marker sequences and target variants are all provided here and so they could be used for future panel

development on other sequencing platforms, and the code could also be adapted and expanded to other data

processing platforms.

Conclusions

The Pacific oyster Cgig_v.1.0 amplicon panel is an effective tool for characterizing genetic variation,

genetic similarity among populations, and relatedness between individuals, including for genetic parentage

analysis applications. Combined with the *amplitools* workflow provided here, this amplicon SNP panel

could benefit oyster breeding programs by facilitating monitoring of genetic variation and confirming

pedigree relationships. The amplicon panel is a useful and lower-cost genotyping tool that has the ability to

easily evolve and improve in future iterations by targeting novel variants and microhaplotypes *in silico*, as

well as by more substantial revisions, such as including putatively adaptive loci, and achieving more

uniform coverage across the genome.

Acknowledgements

Thanks to Chen Yin Walker (Vancouver Island University), Claire Rycroft (University of British Columbia),

and the Molecular Genetics Lab of Fisheries and Oceans Canada for DNA extraction for the study. Thanks

to those who collected or provided samples for this project, including Dr. Kristi Miller (DFO), Rob Saunders

(RKS Labs), Prof. Nicolas Bierne and Dr. Ismaël Bernard (Eurêka Modélisation), Prof. Li Li and Dr. Sheng

Liu, David Howell, J.P. Hastey (Nova Harvest Ltd.), Keith Reid (Odyssey Shellfish), Brian Yip (Fanny Bay

Oysters), and Bruce Evans. Thanks to Geneviève Morin and Brian Boyle of IBIS (Université Laval) for

support in implementing the amplicon panel for genotyping. This work was funded by the Gordon and

Betty Moore Foundation (GBMF#5600) to CAS. This research used resources provided by the SCINet

project of the USDA Agricultural Research Service, ARS project number 0500-00093-001-00-D. Funding

for NFT was provided by USDA ARS CRIS project 2076-63000-005-000-D. Mention of trade names or

commercial products in this publication is solely for the purpose of providing specific information and does

not imply recommendation or endorsement by the U.S. Department of Agriculture.

Competing Interests

Ben Sutherland is affiliated with Sutherland Bioinformatics. The author has no competing financial interests to declare. Some authors affiliated with ThermoFisher Scientific have potential conflicts considering that the AgriSeq Targeted Genotyping by Sequencing solutions and associated Oligo panels that were designed and validated in the study are offered by ThermoFisher Scientific. However, the selection of markers, and data analysis was primarily conducted by other authors. The authors declare that the research was conducted in a scientific manner without any commercial considerations that could be construed as potential conflict of interest and further declare no other conflicts of interest. The other authors declare no competing interests.

Data Availability

The amplicon marker panel referred to here as Cgig_v.1.0 is available commercially from ThermoFisher Scientific as SKU A58237 AGRISEQ PACIFIC OYSTER PANEL. The input data required for marker selection (single SNP per locus; plink and VCF formats), the VariantCaller multilocus genotypes for the pilot study, alignment results of submitted sequences against the reference genome, and sample population annotation is available on FigShare, DOI: 10.6084/m9.figshare.23646471

Three repositories support this project:

Manuscript code repository: https://github.com/bensutherland/ms_oyster_panel

amplitools: https://github.com/bensutherland/amplitools

amplitargets: https://github.com/bensutherland/amplitargets

Additional Files

Additional File S1. Amplicon panel design file including target variant with the reference and alternate allele and flanking genomic sequence as well as genomic coordinates.

Additional File S2. Marker names from ddRAD-seq study, reason for selection in panel, and panel marker name.

Additional File S3. Target SNP file with Cgig_v.1.0 target sites used for variant calling.

Additional File S4. Target design windows that aligned more than once to the reference genome with bowtie2 or bwa mem.

Additional File S5. Per locus H_{OBS} and F_{ST} from the amplicon panel pilot study.

Additional File S6. Per locus Hardy-Weinberg equilibrium statistics for wild or naturalized populations.

Additional File S7. Number of loci genotyped per individual in the population genetic dataset before

626   filters, and in each parentage dataset (VIU OFR5 and MBP CHR8).

627   Additional File S8. Parent-offspring relationships for the VIU OFR5 and MBP CHR8 families.

628   Additional File S9. Per cross identities of null alleles and overlaps between the crosses.

630   Supplemental Results. Figure S1-S8 and Table S1-Table S3.

REFERENCES

631
632

Allen, SK, Rexroad, C, & Rheault, R. (2020). Improving production of the eastern oyster *Crassostrea*
virginica through coordination of genetic improvement programs, research, and technology
transfer activities. *Journal of Shellfish Research, 39*(2), 175-179, 175.
Anderson, EC. (2024). CKMRsim: Inference of pairwise relationships using likelihood ratios. . Retrieved
from https://github.com/eriqande/CKMRsim
Arbelaez, JD, Dwiyanti, MS, Tandayu, E, Llantada, K, Jarana, A, Ignacio, JC, et al. (2019). 1k-RiCA (1K-
Rice Custom Amplicon) a novel genotyping amplicon-based SNP assay for genetics and breeding
applications in rice. *Rice, 12*(1), 55. doi:10.1186/s12284-019-0311-0
Baetscher, DS, Clemento, AJ, Ng, TC, Anderson, EC, & Garza, JC. (2018). Microhaplotypes provide
increased power from short-read DNA sequences for relationship inference. *Molecular Ecology*
Resources, 18(2), 296-305. doi:https://doi.org/10.1111/1755-0998.12737
Barrett, RDH, & Schluter, D. (2008). Adaptation from standing genetic variation. *Trends in Ecology &*
Evolution, 23(1), 38-44. doi:10.1016/j.tree.2007.09.008
Barros, P, Sobral, P, Range, P, Chícharo, L, & Matias, D. (2013). Effects of sea-water acidification on
fertilization and larval development of the oyster *Crassostrea gigas*. *Journal of Experimental*
Marine Biology and Ecology, 440, 200-206. doi:https://doi.org/10.1016/j.jembe.2012.12.014
Barton, A, Hales, B, Waldbusser, GG, Langdon, C, & Feely, RA. (2012). The Pacific oyster, *Crassostrea*
gigas, shows negative correlation to naturally elevated carbon dioxide levels: Implications for
near-term ocean acidification effects. *Limnology and Oceanography, 57*(3), 698-710.
doi:https://doi.org/10.4319/lo.2012.57.3.0698
Beacham, TD, Wallace, C, MacConnachie, C, Jonsen, K, McIntosh, B, Candy, JR, et al. (2018).
Population and individual identification of Chinook salmon in British Columbia through
parentage-based tagging and genetic stock identification with single nucleotide polymorphisms.
Canadian Journal of Fisheries and Aquatic Sciences, 75(7), 1096-1105. doi:10.1139/cjfas-2017-
0168
Botta, R, Asche, F, Borsum, JS, & Camp, EV. (2020). A review of global oyster aquaculture production
and consumption. *Marine Policy, 117*, 103952. doi:https://doi.org/10.1016/j.marpol.2020.103952
Boudry, P, Allal, F, Aslam, ML, Bargelloni, L, Bean, TP, Brard-Fudulea, S, et al. (2021). Current status
and potential of genomic selection to improve selective breeding in the main aquaculture species
of International Council for the Exploration of the Sea (ICES) member countries. *Aquaculture*
Reports, 20, 100700. doi:https://doi.org/10.1016/j.aqrep.2021.100700
Carlsson, J, Morrison, CL, & Reece, KS. (2006). Wild and aquaculture populations of the eastern oyster
compared using microsatellites. *Journal of Heredity, 97*(6), 595-598. doi:10.1093/jhered/esl034
Cheng, BS, Chang, AL, Deck, A, & Ferner, MC. (2016). Atmospheric rivers and the mass mortality of
wild oysters: insight into an extreme future? *Proceedings of the Royal Society B: Biological*
Sciences, 283(1844), 20161462. doi:doi:10.1098/rspb.2016.1462
Csernák, E, Molnár, J, Tusnády, GE, & Tóth, E. (2017). Application of targeted next-generation
sequencing, TruSeq custom amplicon assay for molecular pathology diagnostics on formalin-
fixed and paraffin-embedded samples. *Applied Immunohistochemistry & Molecular Morphology,*
25(7), 460-466.
Delomas, TA, Hollenbeck, CM, Matt, JL, & Thompson, NF. (2023). Evaluating cost-effective genotyping
strategies for genomic selection in oysters. *Aquaculture, 562*, 738844.
doi:https://doi.org/10.1016/j.aquaculture.2022.738844
Divilov, K, Merz, N, Schoolfield, B, Green, TJ, & Langdon, C. (2023). Marker-assisted selection in a
Pacific oyster population for an antiviral QTL conferring increased survival to OsHV-1 mortality
events in Tomales Bay. *Aquaculture, 567*, 739291.
doi:https://doi.org/10.1016/j.aquaculture.2023.739291
Evans, F, Matson, S, Brake, J, & Langdon, C. (2004). The effects of inbreeding on performance traits of
adult Pacific oysters (*Crassostrea gigas*). *Aquaculture, 230*(1-4), 89-98.
Goudet, J, & Jombart, T. (2022). hierfstat: estimation and tests of hierarchical F-statistics.
Green, KM, Spalding, AK, Ward, M, Levine, A, Wolters, EA, Hamilton, SL, et al. (2023). Oregon
shellfish farmers: Perceptions of stressors, adaptive strategies, and policy linkages. *Ocean &*

Coastal Management, 234, 106475. doi:https://doi.org/10.1016/j.ocecoaman.2022.106475

Green, TJ, Siboni, N, King, WL, Labbate, M, Seymour, JR, & Raftos, D. (2019). Simulated marine heat
wave alters abundance and structure of Vibrio populations associated with the Pacific oyster
resulting in a mass mortality event. *Microbial Ecology, 77*(3), 736-747. doi:10.1007/s00248-018-
1242-9

Gruber, B, Unmack, PJ, Berry, OF, & Georges, A. (2018). dartr: An r package to facilitate analysis of SNP
data generated from reduced representation genome sequencing. *Molecular Ecology Resources,*
18(3), 691-699. doi:https://doi.org/10.1111/1755-0998.12745

Guo, X. (2009). Use and exchange of genetic resources in molluscan aquaculture. *Reviews in*
Aquaculture, 1(3-4), 251-259. doi:https://doi.org/10.1111/j.1753-5131.2009.01014.x

Gurney-Smith, HJ, Wade, AJ, & Abbott, CL. (2017). Species composition and genetic diversity of farmed
mussels in British Columbia, Canada. *Aquaculture, 466*, 33-40.
doi:https://doi.org/10.1016/j.aquaculture.2016.08.038

Gutierrez, AP, Bean, TP, Hooper, C, Stenton, CA, Sanders, MB, Paley, RK, et al. (2018). A genome-wide
association study for host resistance to Ostreid herpesvirus in Pacific oysters (*Crassostrea gigas*).
G3 Genes|Genomes|Genetics, 8(4), 1273-1280. doi:10.1534/g3.118.200113

Gutierrez, AP, Turner, F, Gharbi, K, Talbot, R, Lowe, NR, Peñaloza, C, et al. (2017). Development of a
medium density combined-species SNP array for Pacific and European oysters (*Crassostrea*
gigas and *Ostrea edulis*). *G3 Genes|Genomes|Genetics, 7*(7), 2209-2218.
doi:10.1534/g3.117.041780

Hedgecock, D. (1994). Does variance in reproductive success limit effective population sizes of marine
organisms. In A. R. Beaumont (Ed.), *Genetics and evolution of aquatic organisms* (1 ed., pp. 122-
134). London, UK: Chapman and Hall.

Hedgecock, D, & Davis, JP. (2007). Heterosis for yield and crossbreeding of the Pacific oyster
Crassostrea gigas. *Aquaculture, 272*, S17-S29.
doi:https://doi.org/10.1016/j.aquaculture.2007.07.226

Hedgecock, D, Gaffney, PM, Goulletquer, P, Guo, X, Reece, K, & Warr, GW. (2005). The case for
sequencing the Pacific oyster genome. *Journal of Shellfish Research, 24*(2), 429-441, 413.

Hedgecock, D, McGoldrick, DJ, & Bayne, BL. (1995). Hybrid vigor in Pacific oysters: an experimental
approach using crosses among inbred lines. *Aquaculture, 137*(1), 285-298.
doi:https://doi.org/10.1016/0044-8486(95)01105-6

Hedgecock, D, & Pan, FTC. (2021). Genetic divergence of selected and wild populations of Pacific
oysters (*Crassostrea gigas*) on the West Coast of North America. *Aquaculture, 530*, 735737.
doi:https://doi.org/10.1016/j.aquaculture.2020.735737

Hedgecock, D, & Pudovkin, AI. (2011). Sweepstakes reproductive success in highly fecund marine fish
and shellfish: a review and commentary. *Bulletin of Marine Science, 87*(4), 971-1002.

Hedgecock, D, & Sly, F. (1990). Genetic drift and effective population sizes of hatchery-propagated
stocks of the Pacific oyster, *Crassostrea gigas*. *Aquaculture, 88*(1), 21-38.
doi:https://doi.org/10.1016/0044-8486(90)90316-F

Jombart, T, & Ahmed, I. (2011). adegenet 1.3-1: new tools for the analysis of genome-wide SNP data.
Bioinformatics, 27(21), 3070-3071. doi:10.1093/bioinformatics/btr521

Kamvar, ZN, Tabima, JF, & Grünwald, NJ. (2014). poppr: an R package for genetic analysis of
populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ, 2*, e281.
doi:10.7717/peerj.281

King, WL, Jenkins, C, Seymour, JR, & Labbate, M. (2019). Oyster disease in a changing environment:
decrypting the link between pathogen, microbiome and environment. *Marine Environmental*
Research, 143, 124-140. doi:https://doi.org/10.1016/j.marenvres.2018.11.007

Langmead, B, & Salzberg, SL. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods, 9*(4),
357-359. doi:10.1038/nmeth.1923

Lapègue, S, Harrang, E, Heurtebise, S, Flahauw, E, Donnadieu, C, Gayral, P, et al. (2014). Development
of SNP-genotyping arrays in two shellfish species. *Molecular Ecology Resources, 14*(4), 820-830.
doi:https://doi.org/10.1111/1755-0998.12230

Li, C, Wang, J, Song, K, Meng, J, Xu, F, Li, L, et al. (2018). Construction of a high-density genetic map
and fine QTL mapping for growth and nutritional traits of *Crassostrea gigas*. *BMC Genomics,*

739   *19*(1), 626. doi:10.1186/s12864-018-4996-z

740   Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM,
741   arXiv:1303.3997. Retrieved from https://ui.adsabs.harvard.edu/abs/2013arXiv1303.3997L

742   Li, Q, Xu, K, & Yu, R. (2007). Genetic variation in Chinese hatchery populations of the Japanese scallop
743   (*Patinopecten yessoensis*) inferred from microsatellite data. *Aquaculture, 269*(1), 211-219.
744   doi:https://doi.org/10.1016/j.aquaculture.2007.04.017

745   Martínez-García, MF, Ruesink, JL, Grijalva-Chon, JM, Lodeiros, C, Arreola-Lizárraga, JA, de la Re-
746   Vega, E, et al. (2022). Socioecological factors related to aquaculture introductions and production
747   of Pacific oysters (*Crassostrea gigas*) worldwide. *Reviews in Aquaculture, 14*(2), 613-629.
748   doi:https://doi.org/10.1111/raq.12615

749   Meek, MH, & Larson, WA. (2019). The future is now: Amplicon sequencing and sequence capture usher
750   in the conservation genomics era. *Molecular Ecology Resources, 19*(4), 795-803.
751   doi:https://doi.org/10.1111/1755-0998.12998

752   Moran, BM, & Anderson, EC. (2019). Bayesian inference from the conditional genetic stock
753   identification model. *Canadian Journal of Fisheries and Aquatic Sciences, 76*(4), 551-560.
754   doi:10.1139/cjfas-2018-0016

755   Paradis, E. (2010). pegas: an R package for population genetics with an integrated–modular approach.
756   *Bioinformatics, 26*(3), 419-420. doi:10.1093/bioinformatics/btp696

757   Peñaloza, C, Gutierrez, AP, Eöry, L, Wang, S, Guo, X, Archibald, AL, et al. (2021). A chromosome-level
758   genome assembly for the Pacific oyster *Crassostrea gigas*. *GigaScience, 10*(3).
759   doi:10.1093/gigascience/giab020

760   Pew, J, Muir, PH, Wang, J, & Frasier, TR. (2015). related: an R package for analysing pairwise
761   relatedness from codominant molecular markers. *Mol Ecol Resour, 15*(3), 557-561.
762   doi:10.1111/1755-0998.12323

763   Plough, LV. (2016). Genetic load in marine animals: a review. *Current Zoology, 62*(6), 567-579.
764   doi:10.1093/cz/zow096

765   Qi, H, Li, L, & Zhang, G. (2021). Construction of a chromosome-level genome and variation map for the
766   Pacific oyster *Crassostrea gigas*. *Molecular Ecology Resources, 21*(5), 1670-1685.
767   doi:https://doi.org/10.1111/1755-0998.13368

768   Qi, H, Song, K, Li, C, Wang, W, Li, B, Li, L, et al. (2017). Construction and evaluation of a high-density
769   SNP array for the Pacific oyster (*Crassostrea gigas*). *PLOS ONE, 12*(3), e0174007.
770   doi:10.1371/journal.pone.0174007

771   Quinlan, AR, & Hall, IM. (2010). BEDTools: a flexible suite of utilities for comparing genomic features.
772   *Bioinformatics, 26*(6), 841-842. doi:10.1093/bioinformatics/btq033

773   R Core Team. (2023). R: A Language and Environment for Statistical Computing. In. Vienna, Austria.

774   Raymond, WW, Barber, JS, Dethier, MN, Hayford, HA, Harley, CDG, King, TL, et al. (2022).
775   Assessment of the impacts of an unprecedented heatwave on intertidal shellfish of the Salish Sea.
776   *Ecology, 103*(10), e3798. doi:https://doi.org/10.1002/ecy.3798

777   Ritland, K. (1996). Estimators for pairwise relatedness and individual inbreeding coefficients. *Genetics*
778   *Research, 67*(2), 175-185. doi:10.1017/S0016672300033620

779   Salvi, D, & Mariottini, P. (2017). Molecular taxonomy in 2D: a novel ITS2 rRNA sequence-structure
780   approach guides the description of the oysters' subfamily Saccostreinae and the genus Magallana
781   (Bivalvia: Ostreidae). *Zoological Journal of the Linnean Society, 179*(2), 263-276.
782   doi:10.1111/zoj.12455

783   Sauvage, C, Bierne, N, Lapègue, S, & Boudry, P. (2007). Single nucleotide polymorphisms and their
784   relationship to codon usage bias in the Pacific oyster *Crassostrea gigas*. *Gene, 406*(1), 13-22.
785   doi:https://doi.org/10.1016/j.gene.2007.05.011

786   Sun, X, & Hedgecock, D. (2017). Temporal genetic change in North American Pacific oyster populations
787   suggests caution in seascape genetics analyses of high gene-flow species. *Marine Ecology*
788   *Progress Series, 565*, 79-93.

789   Sun, X, Shin, G, & Hedgecock, D. (2015). Inheritance of high-resolution melting profiles in assays
790   targeting single nucleotide polymorphisms in protein-coding sequences of the Pacific oyster
791   *Crassostrea gigas*: Implications for parentage assignment of experimental and commercial
792   broodstocks. *Aquaculture, 437*, 127-139. doi:https://doi.org/10.1016/j.aquaculture.2014.11.009

793   Sutherland, BJG, Itoh, N, Gilchrist, K, Boyle, B, Roth, M, & Green, TJ. (2023). Genomic diversity of
794   wild and cultured Yesso scallop *Mizuhopecten yessoensis* from Japan and Canada. *bioRxiv*,
795   2023.2005.2031.540437. doi:10.1101/2023.05.31.540437

796   Sutherland, BJG, Rycroft, C, Ferchaud, AL, Saunders, R, Li, L, Liu, S, et al. (2020). Relative genomic
797   impacts of translocation history, hatchery practices, and farm selection in Pacific oyster
798   *Crassostrea gigas* throughout the Northern Hemisphere. *Evolutionary Applications, 13*(6), 1380-
799   1399. doi:10.1111/eva.12965

800   Thompson, K. (2023). *Pacific Coast Shellfish Growers Association (PCSGA) 2023 Research Priorities*.
801   Retrieved from https://pcsga.org/research-priorities/

802   Varney, RL, & Wilbur, AE. (2020). Analysis of genetic variation and inbreeding among three lines of
803   hatchery-reared *Crassostrea virginica* broodstock. *Aquaculture, 527*, 735452.
804   doi:https://doi.org/10.1016/j.aquaculture.2020.735452

805   Wada, KT, & Komaru, A. (1994). Effect of selection for shell coloration on growth rate and mortality in
806   the Japanese pearl oyster, *Pinctada fucata martensii*. *Aquaculture, 125*(1), 59-65.
807   doi:https://doi.org/10.1016/0044-8486(94)90282-8

808   Watson, S-A, Southgate, PC, Tyler, PA, & Peck, LS. (2009). Early larval development of the Sydney rock
809   oyster *Saccostrea glomerata* under near-future predictions of CO_2-driven ocean acidification.
810   *Journal of Shellfish Research, 28*(3), 431-437, 437.

811   Weir, BS, & Cockerham, CC. (1984). Estimating F-statistics for the analysis of population structure.
812   *Evolution, 38*(6), 1358-1370. doi:10.2307/2408641

813   Wickham, H. (2016). *ggplot2: elegant graphics for data analysis*: Springer-Verlag New York.

814   Xiao, J, Cordes, JF, Moss, JA, & Reece, KS. (2011). Genetic diversity in U.S. hatchery stocks of
815   *Crassostrea ariakensis* (Fujita, 1913) and comparison with natural populations in Asia. *Journal of*
816   *Shellfish Research, 30*(3), 751-760, 710.

817   Yin, X, Arias-Pérez, A, Kitapci, TH, & Hedgecock, D. (2020). High-density linkage maps based on
818   genotyping-by-sequencing (GBS) confirm a chromosome-level genome assembly and reveal
819   variation in recombination rate for the Pacific oyster *Crassostrea gigas*. *G3*
820   *Genes|Genomes|Genetics, 10*(12), 4691-4705. doi:10.1534/g3.120.401728

821   Yu, Z, & Guo, X. (2004). Genetic analysis of selected strains of eastern oyster (*Crassostrea virginica*
822   Gmelin) using AFLP and microsatellite markers. *Marine Biotechnology, 6*(6), 575-586.
823   doi:https://doi.org/10.1007/s10126-004-3600-5

824   Zhang, G, Fang, X, Guo, X, Li, L, Luo, R, Xu, F, et al. (2012). The oyster genome reveals stress
825   adaptation and complexity of shell formation. *Nature, 490*(7418), 49-54.
826   doi:10.1038/nature11413

827   Zhong, X, Li, Q, Yu, H, & Kong, L. (2013). Development and validation of single-nucleotide
828   polymorphism markers in the Pacific oyster, *Crassostrea gigas*, using High-resolution Melting
829   analysis. *Journal of the World Aquaculture Society, 44*(3), 455-465.
830   doi:https://doi.org/10.1111/jwas.12044

FIGURES

Figure 1. Positions of design sequences for the amplicon panel in the chromosome-level reference genome
(GCF_902806645.1; Peñaloza et al., 2021). Most design sequences mapped to a single location (n = 499;
red/lower), but some mapped to more than one position (teal green/upper, showing multiple locations for
single amplicons), and were subsequently removed from the analysis. Chromosome ends are shown by
hatched vertical lines, and the number of singly mapping design sequences per chromosome are shown to
the right of the plot. Abbreviated NCBI chromosome names are shown to the left of the plot (e.g.,
NC_047559.1 to NC_59).

Figure 2. (A) Per locus observed heterozygosity (H_{OBS}) vs. genetic differentiation (F_{ST}) in the pilot study dataset for all filtered loci. (B) Per locus H_{OBS} for the MBP CHR8 parentage dataset vs. per locus H_{OBS} for the VIU OFR5 and OFR1 parentage dataset for all retained loci in both datasets.

847
848   **Figure 3.** Parent/offspring (PO) relationship network for the VIU OFR5 cross showing all putative PO
849   relationships with log likelihood > 5 using 328 SNP loci. The offspring are denoted with "F" (e.g., F17-37)
850   and the potential parents with "BR" (e.g., BR9). For completeness, all putative linkages are shown,
851   including more than two per offspring individual.

Figure 4. Parent/offspring (PO) relationship network for the MBP CHR8 families showing all putative PO relationships with log likelihood > 5 using 289 SNP loci. The offspring are shown with numeric family identifiers followed by an individual identifier (e.g., 108_4). The putative parents are also shown this way but also including a sex identifier (e.g., 65_5M). For completeness, all putative linkages are shown, including more than two per offspring individual.

TABLES

Table 1. The sources and types of collections used in the study are shown alongside the number of unique
oysters genotyped per collection, and the number and percentage of samples retained after filtering. OFR1
and OFR2 are the broodstock for OFR5, and G0 are the broodstock for OFR1 and OFR2. Acronyms: VIU
= Vancouver Island University; MBP CHR8 = Molluscan Broodstock Program chromosome 8 families; BC
= British Columbia.

Collection	Type	Country of origin	Country collected	Genotyped (n)	Filtered (n)	Retained (%)
Pendrell (PEN)	Naturalized	Canada	Canada	31	29	93.5%
France (FRA)	Naturalized	France	France	32	31	96.9%
Japan (JPN)	Wild	Japan	Japan	22	22	100.0%
China (CHN)	Wild	China	China	33	33	100.0%
China (QDC)	Commercial hatchery	China	China	8	8	100.0%
Guernsey (GUR)	Farm / commercial hatchery	United Kingdom	BC, Canada	31	15	46.9%
Deep Bay (DPB)	Farm / commercial hatchery	*unknown*	BC, Canada	32	32	100%
VIU G0 (F0)	University hatchery	*unknown*	BC, Canada	16	11	68.8%
VIU OFR5 parents (F1)	University hatchery	*unknown*	BC, Canada	53	40	75.5%
VIU OFR5 offspring (F2)	University hatchery	*unknown*	BC, Canada	112	91	81.3%
MBP CHR8 F0	University hatchery	*unknown*	USA	25	24	96%
MBP CHR8 F1	University hatchery	*unknown*	USA	111	111	100%
			Totals	506	447	88.3%

Table 2. Parentage analysis of VIU OFR5 using filtered data (including incompatible genotype removal,
see Methods; n = 328 loci included). In total, 160 of 172 potential assignments were made (93.0% complete),
and eight false positives occurred (4.8% of calls). Per family, the individual identifier (with 'BR' prefix)
for dam and sire are shown alongside their originating family identifier (brackets) and their number of
retained genotyped markers (parentheses). Per family, the number of genotyped offspring is shown with the
number of correct two- or one-parent assignments, the percent completeness of the assignments (% ID),
and any incorrect assignments. All assignments above the applied cutoff were retained (see Methods).
Acronyms: pat. = paternal; mat. = maternal

OFR5 family	Dam [fam.] (loci)	Sire [fam.] (loci)	Off-spring	Two-parent assign.	One-parent assign.	% ID	Incorr. assign.
OFR5_2	BR19 [2-6] (323)	BR11 [1-14] (326)	6	6	-	100.0	
OFR5_3	BR14 [2-10] (306)	BR2 [1-12] (322)	9	8	1	94.4	
OFR5_4	BR14 [2-10] (306)	BR11 [1-14] (326)	6	6	-	100.0	
OFR5_5	BR3 [1-4] (317)	BR10 [1-18] (325)	6	5	1	91.7	1 x pat. sib
OFR5_7	BR17 [2-11] (323)	BR4 [1-10] (317)	7	7	-	100.0	
OFR5_8	BR3 [1-4] (317)	BR7 [2-11] (316)	8	7	1	93.8	
OFR5_9	BR9 [1-16] (318)	BR23 [2-6] (324)	8	8	-	100.0	
OFR5_10	BR25 [1-19] (322)	BR23 [2-6] (324)	7	5	2	85.7	3 x mat. sib
OFR5_12	BR14 [2-10] (306)	BR20 [1-19] (326)	7	7	-	100.0	4 x pat. sib
OFR5_13	BR22 [2-14] (0)	BR26 [1-9] (325)	3	*na*	3	100.0	
OFR5_14	BR24 [1-13] (318)	BR2 [1-12] (322)	8	8	-	100.0	
OFR5_15	BR18 [2-13] (320)	BR10 [1-18] (325)	2	2	-	100.0	1 x pat. sib
OFR5_16	BR22 [2-14] (0)	BR2 [1-12] (322)	7	*na*	4	57.1	
OFR5_17	BR21 [1-1] (322)	BR5 [2-21] (319)	8	5	2	75.0	

874 **Table 3.** Comparative parentage assignment for OFR5 and MBP CHR8 families with all filtered loci or
875 filtered loci with those with at least four incompatibilities observed removed. The number of loci present
876 in the dataset, the number of assignments and percentage of possible assignments, as well as the number of
877 false positives and percentage of assignments made as false positives are shown. The datasets with loci
878 having more than four incompatibilities removed were used for downstream analyses.

Cross event/ families	Locus filter	Num. loci	Num. assign.	Num. possible assign.	Percent assigned	Num. false positive (FP)	Percent assigns as FP
OFR5	all filtered loci	364	134	172	77.9%	11	7.6%
OFR5	Remove loci with ≥ 4 incompatibilities	328	160	172	93.0%	8	4.8%
MBP CHR8	all filtered loci	324	163	215	75.8%	7	4.1%
MBP CHR8	Remove loci with ≥ 4 incompatibilities	289	185	215	86.0%	17	8.4%

880 **Table 4.** Parentage analysis of MBP CHR8 families using filtered data (including incompatible genotype
881 removal, see Methods; n = 289 loci included). In total, 185 of 215 potential assignments were made (86.0%
882 complete), and 17 false positives occurred (8.4% of calls). See VIU OFR5 table for full table explanations.

CHR8 family	Dam [fam.] (loci)	Sire [fam.] (loci)	Off-spring	Two-parent assign.	One-parent assign.	% ID	Incorr. assign.
101	55_3F [55] (281)	55_16M [55] (284)	7	7	-	100.0	2 x [55] aunts
102	55_3F [55] (281)	55_14M [55] (284)	7	7	-	100.0	
103	55_45F [55] (283)	55_17M [55] (285)	7	7	-	100.0	6 x [55] uncles
105	55_27F [55] (286)	58_29M [58] (287)	6	6	-	100.0	1 x [58] uncle, 1 x [55] aunt
106	55_27F [55] (286)	58_20M [58] (284)	7	7	-	100.0	
107	55_45F [55] (283)	58_6M [58] (287)	7	1	6	57.1	3 x [58] uncle, 1 x [58] aunt
108	55_5F [55] (286)	65_5M [65] (281)	7	2	5	64.3	
109	55_5F [55] (286)	65_3M [65] (286)	7	4	3	78.6	
110	55_45F [55] (283)	65_11M [65] (286)	7	4	3	78.6	
111	55_2F [55] (255)	79_6M [79] (286)	6	4	2	83.3	2 x [58] unkn.
112	55_2F [55] (255)	79_4M [79] (281)	7	2	5	64.3	
113	55_45F [55] (283)	79_18M [79] (284)	7	6	1	92.9	
114	65_4F [65] (286)	58_33M [58] (287)	7	7	-	100.0	
115	65_8F [65] (0)	79_13M [79] (282)	7	*na*	7	100.0	1 x [65] uncle
116	55_41F [55] (287)	65_19M [65] (282)	7	4	3	78.6	
117	58_9F [58] (287)	79_1M [79] (284)	7	7	-	100.0	1 x [58] uncle