1
8   **A frontal cortical network is critical for language planning during spoken interaction**
9
10

11   Gregg A. Castellucci[1,2], Christopher K. Kovach[3], Farhad Tabasi[3], David Christianson[3], Jeremy
12   D.W. Greenlee[3,4], Michael A. Long[1,2]*§
13
14
15   [1]NYU Neuroscience Institute and Department of Otolaryngology, New York University Langone
16   Medical Center, New York, NY 10016 USA
17   [2]Center for Neural Science, New York University, New York, NY, 10003 USA
18   [3]Department of Neurosurgery, University of Iowa Hospitals and Clinics, Iowa City, IA, 52242
19   USA
20   [4]Iowa Neuroscience Institute, Iowa City, IA, 52242 USA

27   3 Figures
28   3 Extended Data Figures
29   2 Extended Data Tables

36   * To whom correspondence should be addressed: Michael A. Long (mlong@med.nyu.edu)
37   § Lead contact

Many brain areas exhibit activity correlated with language planning[1–9], but the impact of these dynamics on spoken interaction remains unclear. Here we use direct electrical stimulation to transiently perturb cortical function in neurosurgical patient-volunteers performing a question-answer task[10]. Stimulating structures involved in speech motor function evoked diverse articulatory deficits, while perturbations of caudal inferior and middle frontal gyri – which exhibit preparatory activity during conversational turn-taking – led to response errors. Perturbation of the same planning-related frontal regions slowed inter-speaker timing, while faster responses could result from stimulation of sites located in other areas. Taken together, these findings further indicate that caudal inferior and middle frontal gyri constitute a critical planning network essential for interactive language use[1].

Turn-taking is a central feature of conversation across languages and cultures[11–14]. This key social behavior requires numerous sensorimotor and cognitive operations[11,15,16] that can be organized into three general phases: comprehension of a partner's turn, preparation of a speaker's own turn, and execution of that turn. Using intracranial electrocorticography, we recently demonstrated that neural activity related to these phases is functionally distinct during both task-based and unconstrained turn-taking[1]. The networks active during the perceptual and articulatory stages of turn-taking consisted of structures known to be critical for speech-related sensory and motor function[17–26], while a separate region of frontal cortex was active when participants planned their turns. Specifically, we found planning dynamics were most regularly observed in caudal inferior frontal gyrus (cIFG) and middle frontal gyrus (cMFG).

Given these observations, we hypothesize that cIFG and cMFG are necessary for language-related

planning; therefore, perturbation of preparatory activity within these structures should result in

behavioral deficits selectively related to planning spoken output. In animal models, disruption of

planning dynamics in premotor regions can lead to slower[27] and incorrect[28,29] non-vocal responses

that are otherwise executed properly. Likewise, perturbations of white matter tracts and cortical

regions thought to be important for planning in humans can lead to slower and erroneous task-

based speech[8,30–34]. We therefore predict that manipulation of cIFG and cMFG should result in

similar deficits in the context of spoken interaction. Specifically, because these areas are

persistently active early during planning[1], we expect that disordering this activity would disturb

high-level preparatory processes (e.g., formulating semantic or lexical structure)[35] and thus lead to

response errors but not articulatory dysfunction. Additionally, because efficient planning is central

to achieving the rapid inter-speaker timing characteristic of conversational turn-taking[10,11], we

anticipate that perturbing this network should disrupt this coordination by protracting preparation

times.

Perturbation of neural activity during an interactive language task

To test our predictions, we used 50 Hz bipolar direct electrical stimulation (DES)[17–23,30,8,36–39] to

rapidly and reversibly perturb activity at 58 sites in 23 neurosurgical participants (Extended Data

Tables 1, 2). We stimulated broadly across both cortical hemispheres (48 sites located in the left

hemisphere) using both surface electrodes (n = 31) as well as intracerebral stereo-EEG depth

electrodes (n = 27), enabling us to assess the behavioral results of disrupting our putative planning

network (i.e., cIFG and cMFG; n = 22 sites) and a range of additional cortical loci (e.g., Fig. 1e).

Participants performed the critical information (CI) question-answer task[10,40], which was also used

in our electrocorticography study to isolate planning-related neural dynamics during interaction.

The CI paradigm relies on the same core processes as conversational turn-taking and specifically

requires speech perception, articulation, and all processes related to planning a one-word answer

(e.g., conceptual structure planning, retrieval of wordforms, phonological encoding)[4,41] to occur in

temporally defined epochs (Fig. 1a). At each stimulation site, participants answered a verbally

presented battery of 28 to 143 questions (65 ± 21, mean ± SD) as quickly as possible while DES

was randomly delivered on approximately half of the trials (49 ± 3%; range: 42% - 60%) (see

Methods). Stimulation intensity ranged from 7.5 to 24 V (15 ± 4 V) across sites, and DES was

applied for 3.1 ± 1.0 s during stimulated trials – corresponding to 85 ± 18% of total trial duration

and 91± 3% of planning period duration. Taken together, our perturbations should be capable of

manipulating preparatory and sensorimotor processes related to spoken interaction.

Diverse motor deficits can result from DES

We first observed that cortical stimulation could elicit various motor disturbances that prevented

performance of the CI task. For example, DES of two sites in participant 494L led to distinct

deficits in articulatory performance (Fig. 1b). Specifically, stimulation of a site in subcentral gyrus,

a region important for driving speech articulators[19,26], resulted in unintelligible speech (i.e.,

dysarthria; Fig. 1c). Stimulation of another site in anterior precentral gyrus induced speech arrest,

a well-documented DES-evoked motor deficit[19,22,23,39,42] proposed to arise from an inability to

initiate articulation[43]. In support of this interpretation, we found that answer onset occurred only

76 - 390 ms (191 ± 124 ms) following stimulation offset on arrested trials (Fig. 1d, Extended Data

Fig. 1), which is faster than single word planning times[44,45] and thus suggests that DES prevented

the output of a pre-planned answer. We observed that stimulation evoked similar disruptions to

Figure 1. DES evokes motor deficits. (**a**) Example critical information (CI) questions where the CI is presented early (top) or late (bottom) with behavioral phases indicated. (**b**) Left lateral cortical surface of participant 494L with two stimulation sites indicated. (**c**) Example control trial (left) and trial where stimulation of the posterior site in (b) resulted in articulatory disruption (right); spectrograms of participant response with formant frequencies and envelope amplitude indicated (bottom). (**d**) Schematics of all trials where the anterior site in (b) was stimulated (left) with latencies between stimulation and speech onset shown at right. (**e**) Canonical cortical surface depicting all sites across participants where stimulation evoked motor deficits; all 6 stimulation sites within non-surface structures did not display motor disruptions (Extended Data Table 2). For (e), the location of precentral and postcentral gyri is approximated in light red, and the regions of caudal inferior and middle frontal gyri displaying neural activity related to language planning[1] are approximated in light blue.

overt movement – such as involuntary voicing and gaze deviations – in 5 other participants (Fig.

1e; Extended Data Table 2), and sites exhibiting such effects were preferentially clustered to areas

important for orofacial motor function within precentral and postcentral gyri[17,22,39,46–48] (actual:

86%; mean shuffled: 26 ± 16%) (p = 0.0007; permutation test). Our findings therefore indicate

that these deficits resulted from selective perturbation of well-established motor networks[17–19,22,26]

that are distinct from regions performing other speech-related functions.

DES can evoke planning-related errors

We next tested our first prediction that perturbing planning-related areas would elicit non-motor

response errors, which are traditionally employed in DES studies as an indicator of disrupted

language-related cognitive function[8,18,22,36,37,39,42,49,50]. We observed that DES often resulted in

increased rates of semantic paraphasia (i.e., meaning-related errors; n = 11 sites) (Fig. 2a), anomia

(failure to generate a spoken answer; n = 11 sites) (Fig. 2b), and hesitations (i.e., "umm",

disfluencies, filler words, see Methods; n = 22 sites) (Fig. 2c). Only a single site displayed an

increased rate of neologisms (i.e., phonological errors) on stimulated trials (Extended Data Table

2), indicating that our perturbations largely affected high-level planning operations (e.g.,

formulating semantic and conceptual structure) as opposed to late-stage operations related to

programming phonological and phonetic structure which immediately precede speech onset[9,35,41].

Likewise, stimulation-induced errors did not exhibit gross articulatory disruptions (e.g., Fig. 2a)

and thus are unlikely to be related to motor-related dysfunction.

In our interactive paradigm, we observed that errors were typically infrequent at individual sites,

and their occurrence was variable across sites (Fig. 2d). Nevertheless, we found that DES

Figure 2. DES elicits planning-related response errors. (**a**) Preoperative magnetic resonance images depicting the stimulation site within the Basal Temporal Language Area (BTLA). At right, example control trial (top) and trial where BTLA stimulation resulted in semantic paraphasia (bottom); spectrograms and envelope amplitude of example responses are also presented (middle). (**b-c**) Example trials from sites where stimulation induced anomia (**b**) and hesitations (**c**); site locations indicated on participant cortical surfaces at left. (**d**) Cumulative distribution functions of error counts in control and stimulated trials across sites; inset scatterplots depict error rates. (**e**) The percentage of stimulation sites across participants displaying increased rates of one, two, or three error types (left), with corresponding locations depicted on canonical cortical surfaces (right); stimulation of sites in non-surface structures led to increases in either three error types (n = 1; BTLA site), one error type (n = 3), or no error types (n = 2) (Extended Data Table 2). Site displaying stimulation-induced neologisms is indicated with an arrow. For (e), the location of precentral and postcentral gyri is approximated in light red, and the regions of caudal inferior and middle frontal gyri displaying neural activity related to language planning[1] are approximated in light blue.

significantly increased error rates within frontal and temporal regions (Fig. 2e, Extended Data

Table 2), which is consistent with the findings of past neurosurgical studies that used non-

interactive paradigms (e.g., picture naming, repetition)[18,22,39,42,51]. For example, stimulating the

Basal Temporal Language Area (BTLA), an area of the fusiform gyrus thought to be critical for

semantic access during speech planning[8,37,52,53], resulted in semantic errors, anomia, and

hesitations (Fig. 2a; Extended Data Table 2). Aside from this site, we found that 54% of loci where

stimulation increased rates of any error type were located in cIFG (i.e., pars opercularis or pars

triangularis) or cMFG, which is significantly more than expected by chance (mean shuffled: 39%

± 7%) (p = 0.0251; permutation test); likewise, 64% of sites displaying stimulation-induced

increases in multiple error types were located within these regions. Therefore, while we found non-

motor speech errors only sporadically resulted from DES, stimulation sites exhibiting such deficits

were preferentially located in cIFG and cMFG, supporting our hypothesis that these regions are

essential for proper language-related planning.

Inter-speaker timing can be modulated by DES

We next sought to test our second prediction that perturbation of planning-related neural activity

could protract the time required to prepare an answer during the CI task and thus disrupt inter-

speaker temporal coordination. In particular, conversational turn-taking exhibits rapid transitions

between speakers, with inter-turn gaps (i.e., floor transfer offsets) regularly 200 ms in duration or

less[12,13]. Achieving these latencies requires speakers to simultaneously plan their responses while

perceiving their partner's speech[1,10,11], a behavioral strategy that is also possible when CI is

presented early in the question (e.g., Fig. 1a, top). During these early CI questions, participants

typically respond to the experimenter with sub-second gaps that are significantly shorter than trials

where the CI is presented late (Extended Data Fig. 2a), indicating that participants indeed began

planning earlier when provided the opportunity. We therefore focused our gap timing analysis on

early CI questions, and first tested how this metric was modulated by DES of an established speech

planning site (i.e., BTLA). We found that stimulation of the site within BTLA (Fig. 2a)

significantly lengthened gaps (Fig. 3a,b) ($p < 0.0001$, rank-sum test) without obviously affecting

articulation (e.g., Figs. 2a and 3a), indicating that perturbation of regions important for language-

related planning can measurably disrupt inter-speaker timing.

Across all sites, we observed that stimulation could result in significantly longer gaps ($n = 9$; Fig.

3c) or shorter gaps ($n = 3$ and an additional site in late CI trials only; Fig. 3d; Extended Data Table

2), with the remaining sites leading to no measurable changes ($n = 38$; Fig. 3e; Extended Data Fig.

2b). Because DES could increase or decrease gap duration, we tested whether either type of timing

alteration was associated with planning-related response errors (e.g., Fig. 2a-c). We found that all

locations where stimulation lengthened gap duration exhibited more anomia, semantic paraphasia,

and/or hesitations on stimulated trials than control trials, with all such sites displaying an increase

in at least one type of error (Extended Data Table 2) – significantly more than expected by chance

(actual: 100%; mean shuffled: $48 \pm 15\%$) ($p = 0.0006$; permutation test). Conversely, a consistent

trend for increased errors was not observed when stimulation resulted in shorter gaps (75%, $p =$

0.2795; permutation test) or no effect in gap duration (42%, $p = 0.9412$; permutation test)

(Extended Data Table 2). Therefore, perturbation of language-related planning via DES is likely

signaled by a combination of longer inter-turn gaps and qualitative behavioral deficits including

paraphasia, hesitations, and anomia.

Figure 3. DES-induced modulations of inter-turn gap duration indicate a planning network. (**a**) Example control trial (top left) and trial where the Basal Temporal Language Area (BTLA; see Fig. 2a) was stimulated (bottom left) with inter-turn gap durations indicated; spectrograms of participant response with envelope amplitude overlaid (right). (**b**) All trials related to stimulation of the site in (a) sorted by gap duration. (**c-e**) Cumulative distribution functions of gap duration for example sites where stimulation induced longer gaps (**c**), shorter gaps (**d**), or no statistical difference in gap duration (**e**); site location indicated on participant cortical surfaces at top. (**f-h**) The error rates in control and stimulated trials for all sites where stimulation resulted in longer gaps (**f**), shorter gaps (**g**), or no effect on gap duration (**h**). (**i**) Median and interquartile range of gap duration for all sites where stimulation significantly affected this measure (left); site where DES induced shorter gaps on late CI trials only indicated with an asterisk. Participant and site number indicated in leftmost column (i.e., participant #, site #). At right, canonical cortical surfaces depicting the effect of stimulation on gap duration in all sites across participants; stimulation of sites in non-surface structures either lengthened gaps (n = 2; i.e., BTLA and posterior cingulate gyrus) or did not affect gap duration (n = 4) (Extended Data Table 2). The location of precentral and postcentral gyri is approximated in light red, and the regions of caudal inferior and middle frontal gyri displaying neural activity related to language planning[1] are approximated in light blue. For (f) and (i), the site which displayed stimulation-induced perceptual deficits (see Extended Data Fig. 3c-g) is indicated with an arrow; note that this site was excluded from analyses of anomia in (f).

We then examined the anatomical distribution of stimulation sites across participants to test

whether DES-induced modulations in gap duration were spatially organized. We found that sites

where stimulation did not affect inter-turn gaps were located broadly across the left and right

hemispheres; in contrast, sites exhibiting stimulation-induced increases in gap duration were

primarily restricted to frontal cortical areas (Fig. 3i). Excluding the site in BTLA (Fig. 2a), 7 of

these 8 putative planning sites were located either in cIFG or cMFG – significantly more than

expected if these sites were randomly distributed (actual: 88%; mean shuffled: 39 ± 16%) (p =

0.004; permutation test) – while a single site was located in left posterior cingulate gyrus (Extended

Data Fig. 3a; Extended Data Table 2). Likewise, 32% of sites in cIFG and cMFG displayed

stimulation-induced increases in gap duration while only 3% of sites outside these regions

exhibited this behavioral deficit (shuffled mean: 14%). These results therefore confirm our second

prediction that activity within cIFG and cMFG is critical for maintaining rapid inter-speaker

coordination. Meanwhile, sites where DES resulted in shorter gaps (Extended Data Fig. 3b) were

located outside cIFG and cMFG and not clustered to any specific region of cortex (Fig. 2i),

suggesting this phenomenon does not arise from the same mechanism underlying stimulation-

induced gap lengthening.

Finally, because we are the first to examine the effects of DES on inter-turn gaps, we confirmed

that the observed increases in gap duration were unlikely to have arisen from factors other than

perturbed planning. For example, participants in this study were undergoing surgical treatment for

either epilepsy, brain tumors, Parkinson's disease, or essential tremor, and the constraints for these

procedures required DES to be delivered using three different methods (Extended Data Table 1).

However, we found no significant relationship between the probability of observing stimulation-

induced gap lengthening and the clinical condition of the participant (χ^2 vs. constant model: 0.965,

p = 0.617), the type of stimulator used (χ^2 vs. constant model: 0.468, p = 0.791), or stimulation

intensity (χ^2 vs. constant model: 1.370, p = 0.242), indicating that this putative planning deficit

did not arise from these unconstrained factors. Increased inter-turn gap durations were also

unlikely to have resulted from disrupted language comprehension, as perceptual deficits were

observed at only a single stimulation site examined in this study (Extended Data Fig. 3c-e).

Notably, DES of this particular locus in pars triangularis also evoked a combination of longer gaps,

semantic paraphasia, and hesitations (arrows in Fig. 3f,i; Extended Data Fig. 3f-g), indicating a

mixed function related to perception and planning. This finding is in agreement with our previous

intracranial recordings, which identified a subregion of cIFG overlapping with ventral pars

triangularis that was active during the preparatory and perceptual phases of the CI task[1].

Discussion

In summary, we confirm our predictions that perturbation of cIFG and cMFG would 1) evoke non-

articulatory response errors and 2) result in significantly longer inter-speaker gaps in an interactive

speech task. Our findings therefore demonstrate that these regions are essential to planning during

spoken communication and provide further evidence for the existence of a spatially and

functionally distinct language-related planning network[1]. Previous studies have reported that DES

of left cIFG (i.e., classical Broca's region[54,55]) can result in a range of deficits related to cognitive

and motor function during speech production[18,22,39,42,43,56], and activity within this region has been

linked to multiple linguistic and domain-general cognitive processes[3,5,9,54,57–63]. Therefore, while

cIFG likely represents a heterogenous, potentially multimodal structure[60,63], our results indicate

this region contains circuitry essential for planning spoken language[1]. Similarly, cMFG stimulation

has been observed to result in speech arrest and various non-motor errors[18,51,62], but these

behavioral effects often occur with less regularity when compared to DES of other frontotemporal

structures[18,22,39,42]. Although regions of cMFG are known to be important for several non-linguistic

functions[64–68] including generating eye movements[46,47] (e.g., Fig. 1e), its role in language has

remained unclear. However, a recent large-scale imaging study found that a subregion including

cMFG and adjacent precentral gyrus (i.e., Area 55b) displays activity during linguistic tasks and

exhibits strong connectivity to language-related circuitry[69]. Likewise, cMFG resection can lead to

transient[70] or long-lasting[71] deficits in verbal fluency, suggesting this area may contribute to speech

production. Our stimulation and intracranial recordings further these important findings by

demonstrating that cMFG is critically involved in the planning of spoken language[1]. While

additional research is required to determine the exact planning operations[2,35,72] performed in cMFG

and to define the boundaries of its language-related territory, its close proximity to speech areas in

cIFG[69,73], dorsal laryngeal motor cortex[20,38], and middle precentral gyrus[9] further motivate this

region as a key node within the human language network.

Our broad sampling of the brain with DES and targeted measurement of inter-speaker timing

revealed four sites where cortical stimulation resulted in significantly faster participant responses

(Extended Data Fig. 3b). Previous studies have reported that perturbation of cortical structures can

improve or speed up task behavior in humans and animal models, suggesting that this phenomenon

is generalizable across species and behaviors[74,75]. For example, speech rate could either increase

or decrease when language-related premotor regions were focally cooled[76]. We found that several

sites outside of regions related to action and speech elicited faster responses; therefore, disrupting

neural activity via DES may have modulated a variety of distinct processes both related and

unrelated to language – such as global arousal, impulsivity, planning, and/or language

comprehension – which ultimately led to faster responses. While the underlying mechanisms

remain unknown, these data provide promising initial evidence for cortical stimulation as a therapy

for augmenting cognitive function, perhaps in a manner analogous to the usage of deep brain

stimulation and transcranial magnetic stimulation to treat motor and nonmotor symptoms in a

range of disorders[77–79].

In conclusion, this study advances our understanding of the human language network by

establishing cMFG as a critical speech-related planning area and confirming that cIFG causally

contributes to the preparation of spoken language. Our results suggest these regions contribute to

generating the meaning of speech (e.g., conceptual or lexical content) rather than its lower-level

structure (e.g., phonological or phonetic content), as perturbing these regions elicited both slower

responses and high-level errors (i.e., semantic paraphasia, hesitations, anomia) but not neologisms

or articulatory errors. Therefore, cIFG and cMFG likely function as a key module in a larger

network of both language-specific circuitry, such as those involved in encoding phonetic and

phonological form[2,4,7,9,41,80–82] as well as areas involved in general task performance[5,60,61] – for

example, the longer inter-turn gaps resulting from posterior cingulate stimulation (Extended Data

Fig. 3a) may have resulted from disrupted cognitive control[83,84]. Further DES research combined

with sensitive, ethologically relevant behavioral metrics is well-suited to identify additional

components of this planning network and to delineate their precise contributions to typical and

disordered language generation[85–87].

REFERENCES

1. Castellucci, G. A., Kovach, C. K., Howard, M. A., Greenlee, J. D. W. & Long, M. A. A speech planning network for interactive language use. *Nature* **602**, 117–122 (2022).
2. Guenther, Frank H. *Neural Control of Speech*. (MIT Press, 2016).
3. Flinker, A. *et al.* Redefining the role of Broca's area in speech. *Proc Natl Acad Sci U A* **112**, 2871–5 (2015).
4. Kemmerer, D. From blueprints to brain maps: the status of the Lemma Model in cognitive neuroscience. *Lang. Cogn. Neurosci.* **34**, 1085–1116 (2019).
5. Basilakos, A., Smith, K. G., Fillmore, P., Fridriksson, J. & Fedorenko, E. Functional Characterization of the Human Speech Articulation Network. *Cereb. Cortex* **28**, 1816–1830 (2018).
6. Alario, F.-X., Chainay, H., Lehericy, S. & Cohen, L. The role of the supplementary motor area (SMA) in word production. *Brain Res.* **1076**, 129–143 (2006).
7. Mirman, D., Kraft, A. E., Harvey, D. Y., Brecher, A. R. & Schwartz, M. F. Mapping articulatory and grammatical subcomponents of fluency deficits in post-stroke aphasia. *Cogn. Affect. Behav. Neurosci.* **19**, 1286–1298 (2019).
8. Forseth, K. J. *et al.* A lexical semantic hub for heteromodal naming in middle fusiform gyrus. *Brain* **141**, 2112–2126 (2018).
9. Silva, A. B. *et al.* A Neurosurgical Functional Dissection of the Middle Precentral Gyrus during Speech Production. *J. Neurosci.* **42**, 8416–8426 (2022).
10. Bögels, S., Magyari, L. & Levinson, S. C. Neural signatures of response planning occur midway through an incoming question in conversation. *Sci. Rep.* **5**, (2015).
11. Levinson, S. C. & Torreira, F. Timing in turn-taking and its implications for processing models of language. *Front. Psychol.* **6**, (2015).
12. Stivers, T. *et al.* Universals and cultural variation in turn-taking in conversation. *Proc. Natl. Acad. Sci.* **106**, 10587–10592 (2009).
13. Heldner, M. & Edlund, J. Pauses, gaps and overlaps in conversations. *J. Phon.* **38**, 555–568 (2010).
14. Sacks, H., Schegloff, E. & Jefferson, G. A Simplest Systematics for the Organization of Turn-Taking for Conversation. *Language* **50**, 40 (1974).
15. Castellucci, G. A., Guenther, F. H. & Long, M. A. A Theoretical Framework for Human and Nonhuman Vocal Interaction. *Annu. Rev. Neurosci.* **45**, 295–316 (2022).
16. Kuhlen, A. K. & Abdel Rahman, R. Beyond speaking: neurocognitive perspectives on language production in social interaction. *Philos. Trans. R. Soc. B Biol. Sci.* **378**, 20210483 (2023).
17. Penfield, W. & Boldrey, E. SOMATIC MOTOR AND SENSORY REPRESENTATION IN THE CEREBRAL CORTEX OF MAN AS STUDIED BY ELECTRICAL STIMULATION. *Brain* **60**, 389–443 (1937).
18. Penfield, W. & Roberts, L. THE EVIDENCE FROM CORTICAL MAPPING. in *Speech and Brain Mechanisms* 119–137 (Princeton University Press, 1959).
19. Breshears, J. D., Molinaro, A. M. & Chang, E. F. A probabilistic map of the human ventral sensorimotor cortex using electrical stimulation. *J. Neurosurg.* **123**, 340–349 (2015).
20. Dichter, B. K., Breshears, J. D., Leonard, M. K. & Chang, E. F. The Control of Vocal Pitch in Human Laryngeal Motor Cortex. *Cell* **174**, 21-31.e9 (2018).
21. Hamilton, L. S., Oganian, Y., Hall, J. & Chang, E. F. Parallel and distributed encoding of speech across human auditory cortex. *Cell* **184**, 4626-4639.e13 (2021).
22. Tate, M. C., Herbet, G., Moritz-Gasser, S., Tate, J. E. & Duffau, H. Probabilistic map of critical functional regions of the human cerebral cortex: Broca's area revisited. *Brain* **137**, 2773–2782 (2014).
23. Breshears, J. D., Southwell, D. G. & Chang, E. F. Inhibition of Manual Movements at Speech Arrest Sites in the Posterior Inferior Frontal Lobe. *Neurosurgery* **85**, E496–E501 (2019).
24. Nourski, K. V., Steinschneider, M. & Rhone, A. E. Electrocorticographic Activation within Human Auditory Cortex during Dialog-Based Language and Cognitive Testing. *Front. Hum. Neurosci.* **10**, (2016).

25. Mesgarani, N., Cheung, C., Johnson, K. & Chang, E. F. Phonetic Feature Encoding in Human Superior
Temporal Gyrus. *Science* **343**, 1006–1010 (2014).

26. Bouchard, K. E., Mesgarani, N., Johnson, K. & Chang, E. F. Functional organization of human
sensorimotor cortex for speech articulation. *Nature* **495**, 327–332 (2013).

27. Churchland, M. M. & Shenoy, K. V. Delay of Movement Caused by Disruption of Cortical Preparatory
Activity. *J. Neurophysiol.* **97**, 348–359 (2007).

28. Li, N., Chen, T.-W., Guo, Z. V., Gerfen, C. R. & Svoboda, K. A motor cortex circuit for motor planning
and movement. *Nature* **519**, 51–56 (2015).

29. Li, N., Daie, K., Svoboda, K. & Druckmann, S. Robust neuronal dynamics in premotor cortex during
motor planning. *Nature* **532**, 459–464 (2016).

30. Chernoff, B. L., Sims, M. H., Smith, S. O., Pilcher, W. H. & Mahon, B. Z. Direct electrical stimulation
of the left frontal aslant tract disrupts sentence planning without affecting articulation. *Cogn.*
Neuropsychol. **36**, 178–192 (2019).

31. Klaus, J. & Hartwigsen, G. Dissociating semantic and phonological contributions of the left inferior
frontal gyrus to language production. *Hum. Brain Mapp.* **40**, 3279–3287 (2019).

32. Schuhmann, T., Schiller, N. O., Goebel, R. & Sack, A. T. Speaking of Which: Dissecting the
Neurocognitive Network of Language Production in Picture Naming. *Cereb. Cortex* **22**, 701–709
(2012).

33. Tremblay, P. & Gracco, V. L. Contribution of the pre-SMA to the production of words and non-speech
oral motor gestures, as revealed by repetitive transcranial magnetic stimulation (rTMS). *Brain Res.*
1268, 112–124 (2009).

34. Krieg, S. M. *et al.* Cortical distribution of speech and language errors investigated by visual object
naming and navigated transcranial magnetic stimulation. *Brain Struct. Funct.* **221**, 2259–2286 (2016).

35. Indefrey, P. & Levelt, W. J. M. The spatial and temporal signatures of word production components.
Cognition **92**, 101–144 (2004).

36. Ojemann, G., Ojemann, J., Lettich, E. & Berger, M. Cortical language localization in left, dominant
hemisphere. *J Neurosurg* **71**, (1989).

37. Schaffler, L., Luders, H. O. & Beck, G. J. Quantitative Comparison of Language Deficits Produced by
Extraoperative Electrical Stimulation of Broca's, Wernicke's, and Basal Temporal Language Areas.
Epilepsia **37**, 463–475 (1996).

38. Belkhir, J. R. *et al.* Direct electrical stimulation evidence for a dorsal motor area with control of the
larynx. *Brain Stimulat.* **14**, 110–112 (2021).

39. Chang, E. F. *et al.* Stereotactic probability and variability of speech arrest and anomia sites during
stimulation mapping of the language dominant hemisphere. *J. Neurosurg.* **126**, 114–121 (2017).

40. Bögels, S., Casillas, M. & Levinson, S. C. Planning versus comprehension in turn-taking: Fast
responders show reduced anticipatory processing of the question. *Neuropsychologia* **109**, 295–310
(2018).

41. Indefrey, P. The Spatial and Temporal Signatures of Word Production Components: A Critical Update.
Front. Psychol. **2**, (2011).

42. Sanai, N., Mirzadeh, Z. & Berger, M. S. Functional Outcome after Language Mapping for Glioma
Resection. *N. Engl. J. Med.* 10 (2008).

43. Ferpozzi, V. *et al.* Broca's Area as a Pre-articulatory Phonetic Encoder: Gating the Motor Program.
Front. Hum. Neurosci. **12**, 64 (2018).

44. Bates, E. *et al.* Timed picture naming in seven languages. *Psychon. Bull. Rev.* **10**, 344–380 (2003).

45. Levinson, S. C. & Torreira, F. Timing in turn-taking and its implications for processing models of
language. *Front. Psychol.* **6**, (2015).

46. Lobel, E. *et al.* Localization of human frontal eye fields: anatomical and functional findings of
functional magnetic resonance imaging and intracerebral electrical stimulation. *J. Neurosurg.* **95**, 804–
815 (2001).

47. Amiez, C. & Petrides, M. Anatomical organization of the eye fields in the human and non-human
primate frontal cortex. *Prog. Neurobiol.* **89**, 220–230 (2009).

48. Gordon, E. M. *et al.* A somato-cognitive action network alternates with effector regions in motor
cortex. *Nature* (2023) doi:10.1038/s41586-023-05964-2.
49. Forseth, K. J., Hickok, G., Rollo, P. S. & Tandon, N. Language prediction mechanisms in human
auditory cortex. *Nat. Commun.* **11**, 5240 (2020).
50. Ojemann, J. G., Ojemann, G. A. & Lettich, E. Cortical stimulation mapping of language cortex by
using a verb generation task: effects of learning and comparison to mapping based on object naming.
J. Neurosurg. **97**, 33–38 (2002).
51. Hazem, S. R. *et al.* Middle Frontal Gyrus and Area 55b: Perioperative Mapping and Language
Outcomes. *Front. Neurol.* **12**, 646075 (2021).
52. Lüders, H. *et al.* BASAL TEMPORAL LANGUAGE AREA. *Brain* **114**, 743–754 (1991).
53. Binder, J. R. *et al.* Temporal lobe regions essential for preserved picture naming after left temporal
epilepsy surgery. *Epilepsia* **61**, 1939–1948 (2020).
54. Tremblay, P. & Dick, A. S. Broca and Wernicke are dead, or moving past the classic model of language
neurobiology. *Brain Lang.* **162**, 60–71 (2016).
55. Broca, M. P. Remarques sur le siege de la faculté du langage articulé, suivies d'une observation
d'aphémie (perte de la parole). *Bull Mem Soc Anat Paris* **36**, 330–356 (1861).
56. Chang, E. F., Kurteff, G. & Wilson, S. M. Selective Interference with Syntactic Encoding during
Sentence Production by Direct Electrocortical Stimulation of the Inferior Frontal Gyrus. *J. Cogn.*
Neurosci. **30**, 411–420 (2018).
57. Sahin, N. T., Pinker, S., Cash, S. S., Schomer, D. & Halgren, E. Sequential Processing of Lexical,
Grammatical, and Phonological Information Within Broca's Area. *Science* **326**, 445–449 (2009).
58. Papoutsi, M. *et al.* From Phonemes to Articulatory Codes: An fMRI Study of the Role of Broca's Area
in Speech Production. *Cereb. Cortex* **19**, 2156–2165 (2009).
59. Grodzinsky, Y. & Santi, A. The battle for Broca's region. *Trends Cogn. Sci.* **12**, 474–480 (2008).
60. Fedorenko, E., Duncan, J. & Kanwisher, N. Language-Selective and Domain-General Regions Lie Side
by Side within Broca's Area. *Curr. Biol.* **22**, 2059–2062 (2012).
61. Novick, J. M., Trueswell, J. C. & Thompson-Schill, S. L. Broca's Area and Language Processing:
Evidence for the Cognitive Control Connection: Broca's Area and Language Processing. *Lang.*
Linguist. Compass **4**, 906–924 (2010).
62. Duffau, H. The error of Broca: From the traditional localizationist concept to a connectomal anatomy
of human brain. *J. Chem. Neuroanat.* **89**, 73–81 (2018).
63. Fedorenko, E. & Blank, I. A. Broca's Area Is Not a Natural Kind. *Trends Cogn. Sci.* **24**, 270–284
(2020).
64. Kumar, N., Sidarta, A., Smith, C. & Ostry, D. J. Ventrolateral Prefrontal Cortex Contributes to Human
Motor Learning. *eneuro* **9**, ENEURO.0269-22.2022 (2022).
65. Hosman, T. *et al.* Auditory cues reveal intended movement information in middle frontal gyrus
neuronal ensemble activity of a person with tetraplegia. *Sci. Rep.* **11**, 98 (2021).
66. Buchsbaum, B. R., Greer, S., Chang, W.-L. & Berman, K. F. Meta-analysis of neuroimaging studies
of the Wisconsin Card-Sorting task and component processes. *Hum. Brain Mapp.* **25**, 35–45 (2005).
67. Sierpowska, J. *et al.* Involvement of the middle frontal gyrus in language switching as revealed by
electrical stimulation mapping and functional magnetic resonance imaging in bilingual brain tumor
patients. *Cortex* **99**, 78–92 (2018).
68. Brass, M. The Role of the Frontal Cortex in Task Preparation. *Cereb. Cortex* **12**, 908–914 (2002).
69. Glasser, M. F. *et al.* A multi-modal parcellation of human cerebral cortex. *Nature* **536**, 171–178 (2016).
70. Morshed, R. A. *et al.* Functional outcomes after resection of middle frontal gyrus diffuse gliomas. *J.*
Neurosurg. **137**, 1–8 (2022).
71. Chang, E. F. *et al.* Pure Apraxia of Speech After Resection Based in the Posterior Middle Frontal
Gyrus. *Neurosurgery* **87**, E383–E389 (2020).
72. Dell, G. S. A spreading-activation theory of retrieval in sentence production. *Psychol. Rev.* **93**, 283–
321 (1986).
73. Catani, M. *et al.* Short frontal lobe connections of the human brain. *Cortex* **48**, 273–291 (2012).

74. Zagha, E., Ge, X. & McCormick, D. A. Competing Neural Ensembles in Motor Cortex Gate Goal-
Directed Motor Output. *Neuron* **88**, 565–577 (2015).
75. Guzulaitis, R., Godenzini, L. & Palmer, L. M. Neural basis of anticipation and premature impulsive
action in the frontal cortex. *Nat. Neurosci.* **25**, 1683–1692 (2022).
76. Long, M. A. *et al.* Functional Segregation of Cortical Regions Underlying Speech Timing and
Articulation. *Neuron* **89**, 1187–1193 (2016).
77. Lozano, A. M. *et al.* Deep brain stimulation: current challenges and future directions. *Nat. Rev. Neurol.*
15, 148–160 (2019).
78. Reis, J. *et al.* Noninvasive cortical stimulation enhances motor skill acquisition over multiple days
through an effect on consolidation. *Proc. Natl. Acad. Sci.* **106**, 1590–1595 (2009).
79. De Risio, L. *et al.* Recovering from depression with repetitive transcranial magnetic stimulation
(rTMS): a systematic review and meta-analysis of preclinical studies. *Transl. Psychiatry* **10**, 393
(2020).
80. Jorge, A. *et al.* Hyperdirect connectivity of opercular speech network to the subthalamic nucleus. *Cell*
Rep. **38**, 110477 (2022).
81. Mugler, E. M. *et al.* Differential Representation of Articulatory Gestures and Phonemes in Precentral
and Inferior Frontal Gyri. *J. Neurosci.* **38**, 9803–9813 (2018).
82. Hage, S. R. & Nieder, A. Dual Neural Network Model for the Evolution of Speech and Language.
Trends Neurosci. **39**, 813–829 (2016).
83. Leech, R., Braga, R. & Sharp, D. J. Echoes of the Brain within the Posterior Cingulate Cortex. *J.*
Neurosci. **32**, 215–222 (2012).
84. Pearson, J. M., Hayden, B. Y., Raghavachari, S. & Platt, M. L. Neurons in Posterior Cingulate Cortex
Signal Exploratory Decisions in a Dynamic Multioption Choice Task. *Curr. Biol.* **19**, 1532–1537
(2009).
85. Deger, K. & Ziegler, W. Speech motor programming in apraxia of speech. *J. Phon.* **30**, 321–335 (2002).
86. Jackson, E. S. *et al.* A fNIRS Investigation of Speech Planning and Execution in Adults Who Stutter.
Neuroscience **406**, 73–85 (2019).
87. Whillier, A. *et al.* Adults who stutter lack the specialised pre-speech facilitation found in non-stutterers.
PLOS ONE **13**, e0202634 (2018).
456
457

METHODS

Participants

Study participants were patient-volunteers undergoing surgical treatment at the University of Iowa Hospitals and Clinics for medically intractable epilepsy, brain tumors, or implantation of deep brain stimulation (DBS) electrodes. Data from participants with previous brain resections or infarcts were not included in this study. During the course of treatment, neural activity was recorded using electrocorticography (ECoG) electrodes and/or intracerebral stereo-EEG depth (sEEG) electrodes either chronically (i.e., seizure focus determination) or acutely (i.e., awake craniotomy for tumor removal/epilepsy treatment or during a deep-brain stimulation electrode implant procedure). In ten participants, language function was confirmed to be either left lateralized or bihemispherically distributed using Wada testing or functional magnetic resonance imaging; the remaining participants were right-handed (n = 11) or left-handed (n = 2) with unknown lateralization for language. Participants consented to research, and the University of Iowa Institutional Review Board approved all procedures. Finally, all participants were native speakers of English except for one (684L), who was a native speaker of Vietnamese; consequently, this participant was only screened for stimulation-induced sensorimotor effects and their behavioral data were not analyzed further. Additional demographic and clinical information is available in Extended Data Table 1.

Data Acquisition

For awake craniotomy patients, electrical signals resulting from direct electrical stimulation (DES) were recorded using subdural ECoG grids and/or strips manufactured by Ad-Tech Medical or PMT. Signals were amplified and sampled at 2034.5 Hz using a multichannel amplifier and digital

acquisition system (PZ2 or PZ5 preamplifier with an RZ2 processer; Tucker-Davis Technologies).

For chronically implanted epilepsy patients, electrical signals from subdural electrode grids, strips,

or sEEG intracerebral depth electrodes (Ad-Tech) were recorded at 2000 Hz with a multichannel

amplifier and digital acquisition system (Atlas system, Neuralynx). In both contexts, analog input

channels synchronized with the neural recordings additionally acquired the output of 1-3

microphone(s) which captured the speech acoustics of the experimenter and participant. Input

channels were typically sampled at 48,828 Hz by the TDT system and 16,000 Hz by the Neuralynx

system and downsampled offline. In addition to the electrical signals, a video of the participant

was also acquired at 24 fps for 20 of 23 experiments. The video was synced to the

electrophysiological data after the experiment and provided a secondary high-quality audio

recording channel, which was sampled at 48 kHz.

In a single awake craniotomy experiment, electrical signals were not recorded due to a recording

system error and consequently only a video of the experiment was recorded (641L; see Extended

Data Table 1). This video provided images of the stimulator and captured experiment acoustics;

thus, stimulation and behavioral timing was estimated using this data stream.

Direct Electrical Stimulation (DES)

DES was performed using a constant voltage stimulator (SD9, Grass Instruments). Charge

balanced, biphasic pulses (0.2 ms duration) at 50 Hz were applied to the brain using either: 1) a

handheld stimulator (Magstim BiPolar Probe; MVAP Medical Supplies, Inc., Thousand Oaks, CA)

in awake craniotomy for tumor resection, 2) subdural ECoG strip electrodes (Ad-Tech) in awake

craniotomies for deep brain stimulator implantation, or 3) subdural ECoG grid/strip electrodes or

sEEG electrodes (Ad-Tech) in chronic epilepsy monitoring procedures. Because the position of

the handheld stimulator could vary slightly from trial-to-trial (e.g., the stimulator could be removed

momentarily to apply saline to the brain), we considered stimulation locations that varied by less

than a centimeter to represent a single stimulation site. To titrate the stimulation voltage level, DES

was first applied starting at ~5V and increased in ~2.5V steps until the participant reported a

sensation, after-discharges were observed, or a maximum of 20-25V was reached. The highest

voltage not resulting in a sensation or after-discharges was used as the stimulation level for the

experiment. Occasionally, the stimulation voltage would be lowered during the course of an

experiment if delayed after-discharges were observed. The stimulation voltage at each site

included for analysis in this study is reported in Extended Data Table 2; in cases where the voltage

was lowered, the lowest intensity is reported.

Stimulation timing was manually controlled by the experimenter, who also presented the questions

to the participant in most cases – however, in some experiments, a separate experimenters

presented questions and delivered stimulation. The experimenter controlling the stimulator

typically began applying DES prior to CI presentation and stopped when the participant spoke

their response. However, considerable variability in stimulation timing was observed across trials,

and any trial where DES was applied between CI offset and answer onset was considered a

'stimulated' trial. In all cases, participants were not given any sensory cues related to the timing

of stimulation (e.g., visual or auditory).

Anatomical Reconstructions

Cortical surface reconstruction was performed using T1-weighted or magnetization-prepared

rapid gradient-echo (MPRAGE) images obtained during the clinical workup and the 'recon-all'

pipeline in FreeSurfer[1]. In cases of poor T1 imaging, FreeSurfer processing was repeated with

combined T1 and T2 images or T1 images with fluid-attenuated inversion recovery (FLAIR)

sequences to improve pial surface parcellation. White-matter segmentation errors resulting from

low image quality or tissue abnormalities were corrected manually.

Stimulation site localization in awake craniotomies for tumor resection

In one intraoperative tumor resection case (494L), stimulation site localization and coregistration

was performed using intraoperative photographs, which were aligned to reconstructed cortical

surface meshes via visual comparison of gyral anatomy by two raters (G.A.C. and J.D.W.G.). In

all other cases, intraoperative photographs along with pre- and post-operative magnetic resonance

(MR) images were used to localize both leads of the hand-held stimulator probe. Specifically, we

first localized the craniotomy by 1) aligning intraoperative photographs to reconstructed cortical

surfaces via visual inspection, 2) coregistering any burr holes and bone flap margins visible on the

postoperative MR images to preoperative images using the FLIRT tool in the FSL package[2], and

3) recording the coordinates from the Stealth neuro-navigation system (Medtronic, Minneapolis,

MN, USA) used for craniotomy planning. Stimulation position was then defined on cortical surface

renderings via visual comparison of gyral anatomy by two independent raters (F.T. and C.K.K)

and cross-verification with a third rater (J.D.W.G.).

Electrode localization in awake craniotomies for deep brain stimulator implantations

Localization of subdural ECoG strip electrodes (Ad-Tech, Racine, WI, USA) in patients

undergoing deep brain stimulation (DBS) surgery was performed using a combination of

intraoperative fluoroscopy, preoperative MR images, and pre- and post-implantation computed

tomography (CT) collected for clinical reasons. Because the ECoG strips were temporarily placed

during surgery and removed before closure, intraoperative fluoroscopy was required to record strip

locations for surface-based localization while CT and MR images were used to reconstruct cortical

surfaces and align the fluoroscopic images. First, all MR and CT images were converted from

DICOM (Digital Imaging and Communications in Medicine) to NIfTI (Neuroimaging Informatics

Technology Initiative) format using dsm2niix tool in MRIcroGL[3]. Preoperative CT scans were

then employed as a 3D frame for the coregistration of 2D fluoroscopic images. Specifically, the

'isosurface' function in MATLAB was used to compute the surface geometry of the preoperative

CT, and cortical surface meshes were then transformed into CT space. To coregister the

fluoroscopic image, multiple common control points were identified in the CT volume and 2D

fluoroscopic projection, including skull contours, the superior and inferior terminations of the

frontal sinuses, the glabella, and screws of the stereotactic frame (CRW; Integra, Inc., Princeton,

NJ). Because the view plane angle and field of view varied between fluorograms, control points

were individualized based on the visible skull anatomy. For each individual, a minimum of four

different control points were utilized. For cases with an insufficient number of plainly visible skull

landmarks, a post-implantation CT which displayed DBS leads and burr hole covers was

coregistered with the pre-implantation CT to provide an additional set of reference points. Next,

an orthographic projection from the CT image space to the imaging plane of the fluoroscope was

computed through constrained minimization of squared error between control points. This

procedure yielded an orthogonal affine projection relating control points in CT space to those in

the fluorograms. After this alignment, ECoG electrode locations were defined as the point at which

a line orthogonal to the fluoroscopy-aligned view plane, passing through the electrode shadow

visible in the fluorogram, intersected the cortical surface. Finally, the RAS coordinates of each

electrode were transferred to Montreal Neurological Institute (MNI) space.

Electrode localization in chronically implanted patients

Electrode localization in epilepsy patients undergoing chronic electrode implantation was

performed by identifying characteristic metallic-induced susceptibility artifacts and punctuate

radiodensities in post-implantation MR and CT images, respectively. Electrode coordinates were

then transferred to pre-implantation images via linear image coregistration, followed by manually

guided thin-plate spline warping to account for nonlinear imaging and structural distortions.

Control points for warping were determined by visually identifying corresponding landmark

coordinates in pre- and post-implantation imaging.

Stimulation site coregistration

All contact locations were first determined on cortical surface renderings in Right-Anterior-

Superior (RAS) coordinate space. Anatomical labeling of the electrodes was performed through

surface-based coregistration and segmentation for each participant using the Desikan-Killiany-

Tourville (DKT) atlas[4,5] as assigned by FreeSurfer[6]. After automatic parcellation, electrode

locations relative to DKT atlas labels were visually inspected by two raters (G.A.C and J.D.W.G)

and corrected if necessary. To provide a consistent operational boundary between rostral and

caudal middle frontal gyrus in line with our previous research[7], raters considered any stimulation

sites posterior to the extension of the anterior horizontal ramus of the inferior frontal gyrus as

within caudal middle frontal gyrus and those anterior to this boundary to be within rostral middle

frontal gyrus. Finally, preoperative T1 images were non-linearly coregistered to an MNI-aligned

template brain (CIT168 template)[8] using symmetric diffeomorphic registration in the ANTs

toolbox[9] to transfer electrode locations to MNI space.

Stimulation sites on the lateral cortical surface and superior temporal plane in individual

participants were rendered on coregistration plots (e.g., Fig. 1e) by plotting the center of the two

stimulation electrodes on the gyral surface of the MNI152 reference brain. The locations of sEEG

stimulation sites were rendered on canonical and individual cortical surfaces by plotting the

coregistered coordinates of the most superficial contact of the electrode shaft on the gyral surface

of the MNI152 brain or the location where the shaft penetrated the gyral surface, respectively.

SEEG stimulation sites located on the medial cortical surface, within insular cortex, or on the

ventral cortical surface (i.e., "non-surface sites"; n = 6) were not rendered on coregistration plots.

The coordinates for each pair of stimulated electrodes are reported in Extended Data Table 2.

Behavioral Task

Participants completed the Critical Information (CI) task, which required them to answer simple

questions as quickly as possible. The CI questions were read by the experimenter and adapted from

a stimulus set used in our previous work[7] that itself was adapted from an established Dutch

stimulus set[10,11]. Each CI question contains a single word (i.e., the critical information) which is

required to answer the question (Fig. 1a). In most experiments, questions required participants to

generate the antonym of common words; however, in a subset of experiments, questions could

also ask about animal sounds or body parts.

Questions were presented either in randomized order or in pseudorandomized order to avoid

repeating the same CI on subsequent trials. On approximately half of the trials, DES was applied.

In most experiments, stimulated and control (i.e., non-stimulated) trials were balanced such that

the same question would be presented twice – once as a control trial and once as a stimulated trial.

However, because of clinical considerations (e.g., time constraints in the operating room or to

avoid after-discharges), question order, stimulation status, and wording often deviated from the

planned protocol. In addition, most participants completed a block of the CI questions prior to the

stimulation experiment to familiarize them with the task; these data were not analyzed in the

present study. In one participant (460L), other tasks (e.g., repeating syllables, pressing a button)

were interleaved with trials of the CI task.

The audio acquired with the electrophysiological acquisition system and/or video camera was

annotated and timestamped by a trained phonetician (G.A.C.) to determine the onsets and offsets

of all experimenter questions, CI, and participant responses. All final timestamping was performed

blind in regard to stimulation timing. Questions which could not be accurately timestamped due to

background noise (e.g., clinical team talking in background, operating room door closing, medical

instrument alerts) were excluded from analysis. Likewise, any antonym question where the

participant gave an answer using "not" and the CI (e.g., Question: The opposite of hungry is what

word? Answer: Not hungry) were rejected to ensure participants performed the task by generating

a lexical item rather than repeating the CI. Aside from correct answers, any of the following were

noted:

Semantic paraphasia: 1) an incorrect answer (i.e., giving a synonym when the question required an antonym or giving an answer unrelated semantically to the CI) to a question which was answered correctly at least once or 2) answers related semantically to the CI and/or correct response were considered errors if the same correct answer at least twice during the experiment. In both cases, an answer would be considered a semantic error if the participant spontaneously corrected their answer prior to completing the next trial, except when a participant offered additional correct responses.

Phonological paraphasia: a neologism, or an unattested English word consisting of segments that are properly articulated at a gross level.

Anomia: questions that the participant failed to answer or could not respond for any reason, except for trials where the participant either reported not being able to hear due to background noise or appeared to fall asleep, which were excluded from further analysis. The stimulation site in 494L (Extended Data Fig. 3c) where perceptual dysfunction was reported was excluded from all analyses of anomia.

Hesitations: trials which the participant produced a hesitation (e.g., "uh", "um") or dysfluency (e.g., stuttering, repetition or prolongation of the first segment) prior to responding. In addition, because all CI questions could be answered with a single word, any answers where this word was preceded by filler words (e.g., Question: The opposite of hungry is what word? Answer: "The opposite of hungry is full.") were considered trials containing hesitations.

<u>Qualitative results</u>: any site where stimulation resulted in a motor effect (e.g., speech arrest,

gaze deviation, dysarthria, etc.) as defined by a neurosurgeon (J.D.W.G.). Such effects

were categorized as either 'speech motor' (i.e., speech arrest, orofacial tetanus, involuntary

orofacial movement, sensation of orofacial movement, dysarthria) or 'gaze motor' (i.e.,

involuntary head and/or eye movement). Any site which resulted in a sensation or

movement that indicated the delivery of DES to the participant was not included in

behavioral analyses. Descriptions of all qualitative motor effects are provided in Extended

Data Table 2.

Finally, any site displaying a higher rate of semantic paraphasia, anomia, hesitations, and/or

neologisms in stimulated versus control trials (minimum increase: 1%) was considered to display

an increase in errors.

Behavioral Analysis

Inter-turn gap duration was defined as the duration between question offset and answer onset

(excluding any hesitations) and was computed for all trials where an answer was produced (i.e.,

trials without anomia). Stimulus-response latency, defined as the duration between stimulation

offset and answer onset, was calculated for all stimulated trials.

Stimulation-induced differences in gap duration at each site were assessed in early and late CI

trials separately, as CI position significantly affects gap timing[7,10,11]. Statistical significance in gap

duration between control and stimulated trials was determined using a three-step process 1)

rejecting outliers in the gap duration distributions for control and stimulated trials separately (i.e.,

any gaps greater than 2 interquartile ranges above the 75th percentile or less than 2 interquartile

ranges below the 25th percentile), 2) identifying sites where the median gap duration in control and

stimulated trials differed by at least 100 ms, and 3) performing rank-sum tests ($\alpha = 0.05$) on the

outlier-rejected distributions of gap duration in control and stimulated trials. Behavioral analyses

were not performed unless there were at least 8 answered stimulated and 8 answered control trials

after outlier rejection. Due to restricted study times, we presented participants with more early

trials than late trials (at an ~2:1 ratio); consequently, many sites did not have sufficient late trials

for behavioral analysis. Trial numbers, gap duration summary statistics, and other behavioral data

for each stimulation site is presented in Extended Data Table 2.

Statistical Analyses

All analyses, including permutation tests (100,000 iterations), were performed in MATLAB

(Mathworks, Natick, MA) using custom scripts. Rank-sum tests were performed with the

'ranksum' function, correlation analyses were performed with the 'corr' function, and logistic

regression was performed using generalized linear modelling ('fitglm' function; binomial

distribution, logit linkage function, and maximum of 1000 iterations specified). For all logistic

regression analyses, the occurrence of stimulation-induced increases in gap duration was used as

the categorical response variable and predictor variables were defined as follows:

1) Stimulation intensity vs. increased gap duration: Stimulation voltage was used as a

706   single continuous predictor variable.

707   2) Clinical condition vs. increased gap duration: Stimulation sites were divided into three

708   groups according to whether participants were being treated for epilepsy, a brain tumor,

or essential tremor/Parkinson's disease (combined due to limited sample size)

(Extended Data Table 1). These groups were then used as categorical dummy variables,

with the combined essential tremor/Parkinson's disease group designated as the

reference category.

3) Stimulator type vs. increased gap duration: Stimulation sites were divided into three

groups according to whether DES was delivered via ECoG electrodes, a handheld

stimulator, or sEEG electrodes. These groups were then used as categorical dummy

variables, with the sEEG group designated as the reference category.

METHODS REFERENCES

1. Dale, A. M., Fischl, B. & Sereno, M. I. Cortical Surface-Based Analysis. *NeuroImage* **9**, 179–194 (1999).
2. Jenkinson, M., Bannister, P., Brady, M. & Smith, S. Improved Optimization for the Robust and Accurate Linear Registration and Motion Correction of Brain Images. *NeuroImage* **17**, 825–841 (2002).
3. Rorden, C. & Brett, M. Stereotaxic Display of Brain Lesions. *Behav. Neurol.* **12**, 191–200 (2000).
4. Klein, A. & Tourville, J. 101 Labeled Brain Images and a Consistent Human Cortical Labeling Protocol. *Front. Neurosci.* **6**, (2012).
5. Desikan, R. S. *et al.* An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage* **31**, 968–980 (2006).
6. Fischl, B. Automatically Parcellating the Human Cerebral Cortex. *Cereb. Cortex* **14**, 11–22 (2004).
7. Castellucci, G. A., Kovach, C. K., Howard, M. A., Greenlee, J. D. W. & Long, M. A. A speech planning network for interactive language use. *Nature* **602**, 117–122 (2022).
8. Tyszka, J. M. & Pauli, W. M. In vivo delineation of subdivisions of the human amygdaloid complex in a high-resolution group template: In Vivo Amygdala Subdivisions. *Hum. Brain Mapp.* **37**, 3979–3998 (2016).
9. Avants, B. B. *et al.* A reproducible evaluation of ANTs similarity metric performance in brain image registration. *NeuroImage* **54**, 2033–2044 (2011).
10. Bögels, S., Magyari, L. & Levinson, S. C. Neural signatures of response planning occur midway through an incoming question in conversation. *Sci. Rep.* **5**, (2015).
11. Bögels, S., Casillas, M. & Levinson, S. C. Planning versus comprehension in turn-taking: Fast responders show reduced anticipatory processing of the question. *Neuropsychologia* **109**, 295–310 (2018).

ACKNOWLEDGMENTS

Non-Author Contributions

We thank members of the Long laboratory as well as Taylor Abel, Frank Guenther, Elnaz Hozhabri, Jelena Krivokapić, Kenway Louie, Felix Moll, Caroline Niziolek, and Shy Shoham for comments on earlier versions of this manuscript. We also thank Joel Berger, Haiming Chen, Phillip Gander, Christopher Garcia, Matthew Howard III, My Hieu Kien Huynh, Kenji Ibayashi, Zahra Jourahmad, Hiroto Kawasaki, Mac McKay, Kirill Nourski, Ariane Rhone, and Andrea Rohl for help with data collection.

Funding

This research was supported by R01 DC019354 (M.A.L.), R01 DC015260 (J.D.W.G.), and Simons Collaboration on the Global Brain (M.A.L.).

AUTHOR CONTRIBUTIONS

Conceptualization: GAC, MAL

Data curation: GAC, CKK, FT

Formal analysis: GAC, CKK, FT, DC

Funding acquisition: JDWG, MAL

Investigation: GAC, DC, JDWG

Methodology: GAC, CKK, FT, DC, MAL, JDWG

Project administration: GAC, JDWG

Resources: JDWG, MAL

Software: GAC, CKK, FT, DC

768   Supervision: JDWG, MAL

769   Validation: GAC, CKK, JDWG

770   Visualization: GAC, MAL

771   Writing – original draft: GAC

772   Writing – review & editing: GAC, CKK, FT, DC, JDWG, MAL

774 **COMPETING INTERESTS**

775 The authors declare no competing interests.

777 **DATA AVAILABILITY**

778 Due to concerns about patient confidentiality, data are not publicly available; however, the

779 corresponding author will provide deidentified behavioral data upon request.

781 **CODE AVAILABILITY**

782 All MATLAB code used in this study will be provided upon request to the corresponding author.