Interrogating basal ganglia circuit function in Parkinson's disease and dystonia

S Sumarac, BEng[†1,2], KA Spencer, BSc[†1,2], LA Steiner, MD/PhD[†1,3,4], C Fearon, MD[1,5,6], E Haniff, BSN[1], AA Kühn, MD[3], M Hodaie, MD/MSc[1,7,8,9], SK Kalia, MD/PhD[1,8,9,10], AM Lozano MD/PhD[1,7,8,9], A Fasano MD/PhD[1,5,6,7,8], WD Hutchison, PhD[1,8,9,11],

L Milosevic, PhD[1,2,7,8,10]

1. Krembil Brain Institute, University Health Network, Canada.

2. Institute of Biomedical Engineering, University of Toronto, Canada.

3. Department of Neurology and Experimental Neurology, Charité-Universitätsmedizin Berlin, Germany.

4. Berlin Institute of Health (BIH), Germany.

5. Edmond J. Safra Program in Parkinson's Disease, Morton and Gloria Shulman Movement Disorders Clinic, Toronto Western Hospital, Canada.

6. Department of Neurology, University of Toronto, Canada.

7. Institute of Medical Sciences, University of Toronto, Canada.

8. Center for Advancing Neurotechnological Innovation to Application (CRANIA), Toronto, Canada.

9. Department of Surgery, University of Toronto, Canada.

10. KITE, University Health Network, Toronto, Canada.

11. Department of Physiology, University of Toronto, Canada.

† Indicates equal contributions

Correspondence: Luka Milosevic; 399 Bathurst St, 11MP301, Toronto, ON, Canada, M5T 2S8;

mailto:luka.milosevic@mail.utoronto.ca

Running title: Circuit function in Parkinson's and dystonia

Financial Disclosures / Conflicts of Interest: All authors have no financial disclosures / conflicts of interest to declare that are related to this work.

Funding: This project has been made possible with the financial support of Health Canada, through the Canada Brain Research Fund, an innovative partnership between the Government of Canada (through Health Canada) and Brain Canada, and of the Azrieli Foundation (L.M.); as well as a grant from the Banting Research Foundation in partnership with the Dystonia Medical Research Foundation (L.M.).

Abstract

Background: The dichotomy between the hypo- versus hyperkinetic nature of Parkinson's disease (PD) and dystonia, respectively, is thought to be reflected in the underlying basal ganglia pathophysiology.

Objective: Investigate differences in globus pallidus internus (GPi) neuronal activity, and short- and long-term plasticity of direct pathway projections.

Results: GPi neurons were slower, burstier, and less regular in dystonia. In PD, symptom severity positively correlated with the power of low-beta frequency spiketrain oscillations. In dystonia, symptom severity negatively correlated with firing rate, and positively correlated with neuronal variability and the power of theta frequency spiketrain oscillations. Dystonia was moreover associated with less long-term plasticity and slower synaptic depression.

Conclusion: We substantiated claims of hyper- versus hypofunctional GPi output in PD versus dystonia, and provided cellular-level validation of the pathological nature of theta and low-beta oscillations in respective disorders. Such circuit changes may be underlain by disease-related differences in plasticity of striato-pallidal synapses.

Keywords: basal ganglia, globus pallidus, Parkinson's disease, dystonia, neurophysiology, synaptic plasticity

Introduction

Parkinson's disease (PD) is a complex neurodegenerative disorder characterized by non-motor manifestations and motor symptoms including tremor, bradykinesia, rigidity, postural instability, and freezing of gait [1]. Dystonia is a movement disorder characterized by involuntary muscle contractions, repetitive movements, and abnormal fixed postures [2]. While PD is characterized as a collection of hypokinetic syndromes (apart from tremor and levodopa-induced dyskinesias), dystonia is considered a hyperkinetic movement disorder.

The dichotomy between the hypo- versus hyperkinetic nature of PD and dystonia, respectively, is thought to be reflected in the underlying basal ganglia pathophysiology. In particular, basal ganglia circuitry that involves "direct" (D1-mediated striatal projections to the internal globus pallidus; GPi) and "indirect" (D2-mediated striatal projections via the external globus pallidus and subthalamic nucleus to GPi) pathways has been implicated in these disorders [3]. In PD, activity in the indirect pathway is upregulated while the direct pathway is downregulated, both of which converge towards increased GPi-mediated inhibition of thalamocortical motor networks [4]. Conversely, dystonia has been hypothesized to occur as a result of hypofunctional indirect pathway activity [5,6] and/or hyperfunctional direct pathway activity [7], producing decreased GPi-mediated inhibition of thalamocortical motor networks. These circuit differences would thereby result in increased GPi firing rates in PD and decreased rates in dystonia; which has been substantiated by some [8,9] but not all [10] human single-neuron studies. In addition to potential differences in single-neuron firing properties, each of the disorders has been associated with aberrant synchronization at the neural aggregate level; namely, increased beta (13-30Hz) and theta (4-12Hz) local field potential (LFP) oscillations in PD [11] and dystonia [12], respectively.

Despite opposing neurocircuit pathologies, deep brain stimulation (DBS) of the GPi is a widely used treatment for the motor symptoms of both PD and dystonia. However, the timescales of symptomatic improvement after GPi-DBS and the reappearance of symptoms upon stimulation cessation are markedly longer in dystonia (minutes-hours-days) compared to PD (seconds-minutes) [13]. These differences in timescales may be the result of disease-specific dynamics of various forms of synaptic plasticity [14]. Intriguingly, high-frequency GPi stimulation has been shown to induce synaptic depression of inhibitory striato-pallidal (GPi) projections during stimulation (short-term plasticity effects), followed by an enduring potentiation of these

projections after stimulation cessation (long-term plasticity-like effects) in both patients with PD

[15] and dystonia [16]. However, the dynamics of these plastic effects have not been directly

compared between disorders.

In this work, we compared GPi single-neuron activities (rate and oscillation -based features)

between PD and dystonia, and investigated the relationships between electrophysiological

features and disease severity. Moreover, we investigated differences in the dynamics of short-

and long-term plasticity of the direct pathway projections.

Methods

Patients and clinical scores

For the single-neuron feature analysis, data were collected from 19 patients with non-genetic dystonia (135 included segments; see Supplementary Materials for information on subtypes) and 44 patients with PD (222 segments) during awake microelectrode-guided GPi-DBS surgeries [17]; after overnight withdrawal of antiparkinsonian medications in patients with PD. Corresponding preoperative clinical scores for PD using the Unified Parkinson's Disease Rating Scale Part III (UPDRSIII) and dystonia using either the Burke-Fahn-Marsden Dystonia Rating Scale (BFMDRS) or the Toronto Western Spasmodic Torticollis Rating Scale (TWSTRS) were also amalgamated for each patient. Given that UPDRSIII includes both hypokinetic and hyperkinetic symptoms of PD, we further sought to disaggregate the score by only considering items 23-26 in UPDRSIII, which assess hypokinetic symptoms of PD. To enable comparison across different dystonia scales, clinical scores were normalized (0-100%) with respect to the maximum severity per scale (108 for total UPDRSIII, 32 for hypokinetic only UPDRSIII, 85 for TWSTRS, and 120 for BFMDRS). For plasticity analyses, data were collected from a subset of 8 patients with dystonia (12 recording sites) and 10 patients with PD (13 recording sites). A data summary is available in Supplementary Table 1. Each patient provided informed consent and experiments were approved by the University Health Network Research Ethics Board and adhered to the guidelines set by the tri-council Policy on Ethical Conduct for Research Involving Humans.

Intraoperative data acquisition

Two independently driven microelectrodes (600µm spacing; 0.2–0.4MΩ impedances; ≥10kHz; sampling rate) were used for recordings; amplified using two Guideline System GS3000 amplifiers (Axon Instruments, Union City, USA) and digitized using a CED1401 data acquisition system (Cambridge Electronic Design, Cambridge, UK). For single-neuron data, units were sampled across the spatial extent of GPi as part of the standard of care neurophysiological mapping procedure; as previously described [17]. For plasticity data, microstimulation was delivered from one microelectrode using biphasic (cathodal followed by anodal) pulses (100µA, 150µs) from an isolated constant current stimulator (Neuro-Amp1A, Axon Instruments), while recording field evoked potentials (fEPs) using an adjacent microelectrode at the same depth; as

done previously [15,16,18]. Initially, ten low-frequency stimulation (LFS) pulses were delivered

at 1Hz to obtain baseline measurements of fEP amplitudes (i.e., functional readouts striato-

pallidal inhibitory efficacy). Then, a standard high-frequency stimulation (HFS) tetanizing

protocol was delivered (four 2s blocks of 100Hz, each separated by 8s) [16], after which another

set of LFS pulses was delivered to obtain measurements of post-HFS fEP amplitudes (plasticity

protocol depicted in Fig. 2A). Long-term plasticity analysis involved quantifying fEP changes

before versus after HFS. Short-term plasticity analysis involved quantification of successive fEP

amplitudes during HFS.

Offline analyses and statistics

GPi high-frequency discharge (HFD) neurons [17] of >4 signal-to-noise ratio and <1% interspike

interval violations were included in the single-neuron analyses. Segments were band-pass filtered

(300-3000Hz) and template matched, after which neurophysiological features were extracted

from each segment, including per patient median firing rate (FR), and the ninetieth percentile of

the burst index (BI), coefficient of variation (CV), and oscillatory power across the theta (4-

8Hz), alpha (8-12Hz), low-beta (12-21Hz), and high-beta (21-30Hz) frequency bands.

Oscillatory power was extracted using Lomb's periodogram, performed on the autocorrelation

function of single-neuron segments; as previously described [19]. The burst index was computed

by taking the ratio of the means from a two-component Gaussian mixture model applied to the

log interspike interval distribution, a modification of the previous mode-over-mean ISI method

[20]. Neuronal features were compared between PD and dystonia using 2-tailed Mann-Whitney

U (MWU) tests due to non-normality of features. Spearman correlations were employed to

investigate relationships between extracted features and clinical scores. For comparing

differences in single-neuron features between PD and dystonia, significant results were followed

up with post hoc multiple comparisons with a Bonferroni correction. For clinical correlations,

non-parametric Monte Carlo permutation tests were used, avoiding assumptions about data

distribution. The tested values were randomly shuffled 5,000 times to form a probability

distribution, with the p-value reflecting the original sample rank. All tests underwent adjustment

for multiple comparisons, controlling the false discovery rate (FDR) at an α-level of 0.05.

All fEP measurements for plasticity analyses were normalized with respect to the root mean

square amplitude of the rectified pre-stimulation LFP (low-pass filtered at 50Hz). For long-term

plasticity analyses, the median normalized fEP amplitudes pre- versus post-HFS at each

recording site were analyzed using a linear mixed model (LMM) with patient ID as a random

factor, normalized fEP amplitudes as the response variable, and epoch as a fixed effect, to

establish whether plasticity was elicited in each disorder. The percentage change from pre- to

post-HFS was also compared across disorders using a LMM with patient ID as a random factor.

For short-term plasticity analysis, the dynamic change of the first five fEP amplitudes during the

first train of HFS was assessed; normalized with respect to the first fEP amplitude for each train.

Given the relatively rapid rate of attenuation of fEP amplitudes in GPi with HFS [16,18], the first

five fEPs were fit with a double-exponential function, and the half-life of the faster decaying

exponential component was used as a readout of the rate of attenuation. Half-life measurements

were then compared across disorders using a using a LMM with patient ID as a random factor.

Data Sharing

The data that support the findings of this study are available in a public repository (see:

https://osf.io/nqzd2/)

Results

Differences in single-neuron features

In dystonia, FR was lower (MWU test; p=.0252) and neurons were burstier (higher BI; MWU test; p=.0024) and less regular (higher CV; MWU test; p<.0001) compared to PD (Fig. 1A). However, there were no significant differences in spiketrain oscillations in the delta (MWU test; p=.6265), theta (MWU test; p=.3649), low beta (MWU test; p=.1176), or high beta (MWU test; p=.4585) frequency bands (Fig. 1B). The post hoc Bonferroni correction (adjusted significance threshold to $p < 0.0071$) indicated that firing rate difference between PD and dystonia did not survive multiple comparisons (Supplementary Table 3). A neuronal feature summary is available in Supplementary Table 2.

Clinical correlations

For rate-based spiketrain features, no clinical correlations were found for PD. In dystonia, symptom severity negatively correlated with FR (Spearman's rho=-0.49, p=.0370) and positively correlated with CV (Spearman's rho=0.48, p=.0364; Fig. 1C). For oscillatory spiketrain features, symptom severity positively correlated with low beta in PD (Spearman's rho=0.30, p=.0426), with a marginally stronger correlation for PD hypokinetic symptoms only (items 23-26 of UPDRSIII, Spearman's rho=0.32, p=.0330; Supplementary Fig. 3), and theta in dystonia (Spearman's rho=0.51, p=.0242; Fig. 1C). However, none of the clinical correlations survived Benjamini-Hochberg FDR-correction for multiple comparisons (Supplementary Table 4).

Long-term plasticity-like effects

In PD, fEP amplitudes were significantly greater after compared to before HFS (LMM; p = .0075, effect size = 5.42 ± 1.79; Fig. 2C), while in dystonia, the increase approached but did not reach statistical significance (LMM; p = .0708, effect size = 2.82 ± 1.45; Fig. 2C). The percent change in fEP amplitude was significantly greater in PD compared to dystonia (LMM; p = .0396, effect size = 12.40 ± 5.41; Fig. 2D).

Short-term plasticity

Half-life values extracted from double exponential fits on the first five fEPs during HFS were significantly greater in dystonia compared to PD (LMM; p = 0.0139, effect size = 1.04 ± 0.37; Fig. 2G).

Discussion

In this work, we leveraged the intraoperative environment to characterize GPi spiketrain features and to record and manipulate the efficacy of inhibitory direct pathway striato-pallidal projections, in order to derive insights about disease-related features of basal ganglia circuit function in PD and dystonia. In particular, we found that GPi neurons exhibited lower firing rates, but greater burstiness and variability in dystonia compared to PD (Fig. 1A). While no differences were found in the power of spiketrain oscillations across disorders (Fig. 1B), we found that PD symptom severity positively correlated with the power of low-beta frequency spiketrain oscillations, whereas dystonia symptom severity positively correlated with the power of theta frequency spiketrain oscillations (Fig. 1C). Dystonia symptom severity moreover correlated negatively with firing rate, and positively with neuronal variability. These results are discussed in greater detail with respect to previous literature in the subsequent Discussion section entitled "*Neuronal correlates of PD and dystonia*." In response to electrical stimulation (protocol depicted in Fig. 2A), we found significant increases in the amplitudes of positive-going stimulation-evoked field potential amplitudes (considered to reflect striato-pallidal synaptic strength; as exemplified in Fig. 2B) before versus after HFS in both PD and dystonia (Fig. 2C); with recording sites in PD exhibiting significantly greater increases (Fig. 2D). While changes to evoked potential amplitude before versus after stimulation can be considered to be reflective of long-term plasticity [15,18], the dynamics of evoked potentials during HFS (as depicted in Fig. 2E) can be considered as reflective of short-term synaptic plasticity [18,21]. To this end, our findings are suggestive of faster latency synaptic depression in PD compared to dystonia (Fig. 2F/G). Plasticity findings are discussed in greater detail in the Discussion section entitled "*Direct pathway plasticity*."

Neuronal correlates of PD and dystonia

The basal ganglia circuit models of PD and dystonia have been suggested to be driven by an imbalance between direct and indirect pathways [22,23]. While hyperfunctionality of the indirect pathway and hypofunctionality of the direct pathway are associated with PD [24], the direct pathway has been suggested to be hyperfunctional in dystonia [7]. To this end, our findings (Fig. 1A) of lower firing rates in dystonia compared to PD may further substantiate [8,9] such claims, whereby greater inhibition of GPi would produce decreased inhibitory output, and thus under-

inhibition of thalamocortical motor networks, giving rise to hyperkinetic motor symptoms;

whereas the opposite would be true in PD [3]. These claims are further substantiated by findings

of an inverse relationship between firing rate and disease severity in dystonia (Fig. 1C; rate-

based features); although the opposite was not found in PD. Despite the lack of correlations with

firing rate in PD, our findings seem to align with those of Muralidharan and colleagues [25], who

showed that GPi neuronal firing rates may not directly correlate with motor severity but exhibit

variability across the disease severity continuum in parkinsonian non-human primates (initially

increasing, then decreasing, then increasing again at mild, moderate, and severe disease

manifestations, respectively). Thus, while GPi discharge rates may change in PD, such changes

may not be reflected by linear relationships with motor sign development and progression.

Indeed, variability in spike firing rates in PD may be reflected in the considerable overlap in

spiking activity between PD and dystonia (Fig. 1A), with many dystonia patients exhibiting

higher discharge rates compared to PD patients. While differences in discharge rates were

nevertheless observed between PD and dystonia, it may be that the combination of rate and

pattern (reflected in the BI and CV) changes best differentiates the two disorders.

From a neural synchronization standpoint, we recently showed that pathological oscillations at

the neural aggregate / LFP level are likely encoded by periodic oscillations in the firing rates of

single neurons [26]. In the current work, we did not find significant differences in the power of

spiketrain oscillations across disorders (Fig. 1B). Although previous research has reported

differences in the LFP power between PD and dystonia [27,28], a study in non-human primates

found no such differences in single-neuron oscillatory strength [8], as reflected in our findings.

However, despite a lack of difference in overall power across disorders, we were able to derive

disease/frequency-specific relationships with respect to clinical scores (Fig. 1C; oscillatory

features). In particular, we provided evidence of relationships between the power of spiketrain

oscillations in the theta and low beta frequency bands and the severity of motor symptoms in

dystonia and PD, respectively; providing a cellular level validation of previous frequency-

specific LFP clinical correlation studies in people with PD (beta oscillations) [11] and dystonia

(theta oscillations) [12]. Indeed, Muralidharan and colleagues [25] also showed linear group-

level relationships between low-beta frequency spiketrain oscillations and disease severity in

parkinsonian non-human primates, despite the lack of linear relationships with spike discharge

rates (as discussed above).

Direct pathway plasticity

We have previously shown that the amplitude of positive-going inhibitory fEPs is directly

associated with the duration of neuronal inhibition [15,29], thus serving as a viable substrate of

inhibitory synaptic transmission efficacy. These observations are supported by work in animals

demonstrating the reversal of both positive-going extracellular fields and neuronal inhibition

with the application of GABA antagonists [30,31]. Importantly, we previously [29] demonstrated

that low latency inhibitory fields recorded in the subthalamic nucleus (GPe-mediated) are far

smaller in amplitude and are resilient during HFS (i.e., minimal synaptic depression), whereas

high latency inhibitory fields recorded in substantia nigra pars reticulata and GPi [16] (striatum-

mediated) are far larger in amplitude and are subject to potent synaptic depression during HFS;

thus, enabling electrophysiological discernment of these distinct inhibitory projections in

humans.

To this effect, our findings may suggest disease-specific differences in mechanisms underlying

both short- [32] and long-term [33,34] forms of synaptic plasticity of direct pathway projections

(Fig. 2). Converging evidence from past animal and human studies suggests that dystonia is

associated with impaired synaptic function and abnormal synaptic plasticity [35–37]. Compared

to healthy controls, it has been shown that transcranial magnetic stimulation induced motor

evoked potentials (MEPs) are hyperexcitable in dystonia [38,39], and somatosensory and motor

cortical plasticity is greater [40]. Likewise, enhanced long-term potentiation at cortico-striatal

synapses has been shown in rodent models of dystonia [41,42]. While our finding that long-term

potentiation effects are greater in PD compared to dystonia (Fig. 2D) is difficult to corroborate

with this literature, one potential explanation can be that all of our PD patients are long-term

users of levodopa. We have previously shown that the intake of this antiparkinsonian

dopaminergic medication leads to potent increases in the magnitude of direct pathway plasticity

[15]. Although patients are 12hr withdrawn form antiparkinsonian medications for surgery, it

could be that striato-pallidal synapses are nevertheless chronically over-sensitized from

prolonged use of dopaminergic medication; which is a well-known hypothesis related to the

manifestation of levodopa-induced dyskinesias (a hyperkinetic feature) in PD [43]. Indeed, a lack

of depotentiation of striato-pallidal projections has previously been observed in patients with

levodopa-induced dyskinesias [44]. As such, excessive plasticity of these projections may

corroborate hyperkinetic features of dystonia and levodopa-induced dyskinesias in PD.

In addition, the variation in the modulation of these striato-pallidal projections to electrical

stimulation may also be reflective of differences in the mechanism by which DBS operates

across PD and dystonia, despite a common stimulation target. Clinical studies in dystonia have

shown that DBS leads to a more rapid improvement in the transient, dynamic muscle

contractions (phasic components) of the disorder when compared to the sustained, continuous

muscle contractions (tonic or fixed components) [45]. The early improvement of the phasic

component has been suggested to be associated with direct suppression of abnormal activity in

the GPi [46], whereas the improvement of tonic components has been associated with long-term

potentiation-like synaptic plasticity in TMS studies [14,47]. This contrast with PD, where the,

the maximal clinical response to DBS occurs within a much faster time course [13,48]. The

present study shows disease-specific differences in the dynamics associated with modulation of

direct pathway projections; thus, potentially providing a local synaptic rationale for the

differential time courses associated with the clinical effects of DBS in the same target.

Specifically, we found faster rates of synaptic depression during HFS and greater post-HFS long-

term plasticity-like effects in PD compared to dystonia, implying that striato-pallidal synapses in

dystonia are perhaps more resistant to change compared to PD. Our findings may therefore

explain the phenomenon that functional benefits of DBS require longer time courses in dystonia,

if the effects of DBS rely upon network reorganization, which involves adjustments in neural

connectivity or synaptic efficacy in response to stimulation [47]. Indeed, the benefits of GPi-

DBS in dystonia have also been shown to persist for long periods of time even after the cessation

of stimulation [14]. In PD, washout effects take longer with GPi-DBS compared to STN-DBS

[49], and long-term dyskinesia reduction has also been observed with GPi-DBS even when DBS

is off [50,51]. While further work is certainly required to better understand disease-related

differences in plasticity, our findings may nevertheless motivate the development of periodic

intermittent (ON/OFF) DBS strategies which periodically modulate synaptic plasticity for

therapeutic benefits which outlast stimulation delivery, as have recently been employed in

preclinical work [52,53].

Limitations and future directions

Important limitations of intraoperative/intracranial studies in humans include a lack of access to

healthy control data (hence comparison across disorders), the inability to use pharmacological

interventions to verify pathway specificity of elicited responses, and time constraints preventing

thorough scrutinization of time courses of long-term plasticity-like effects. Indeed, GPi receives

the greatest abundance of inhibitory inputs from striatum (direct pathway), but also it also

receives inhibitory inputs by way of GPe (indirect pathway). Although we can functionally

disaggregate these pathway-specific responses based on differences in morphology and dynamics

of GPe-mediated versus striatum-mediated inhibitory fEPs [21]; the possibility of compounded

effects cannot be completely ruled out. While our neuronal investigations provide cellular level

support for closed-loop targeting of disease-related neural oscillations [54,55], future

applications of DBS may also benefit from closed loop tuning of basal-ganglia-thalamo-cortical

circuit dynamics and plasticity through chronic monitoring of evoked potential responses [56].

Indeed, optogenetic studies in parkinsonian rodents have demonstrated the ability to achieve

lasting therapeutic efficacy via periodic activations of striatal direct pathway projections [52,53],

likely leveraging long-term plasticity-like mechanisms. An additional limitation of this study is

that we did not stringently monitor personal medication intake or levels of intraoperative

sedation; however, all patients with PD were asked to withdraw from medications the night

before surgery, and the analyses only included patients operated on awake. It is important to

consider that classifications of PD and dystonia as hypo- and hyperkinetic disorders can be

considered oversimplifications, as there can be contradictory comorbidities and drug- and DBS-

related effects [57,58]; though our results do not change substantially when only hypokinetic PD

features are considered (Supplementary Fig. 3). Additionally, our analyses involved pooling of

patients with various forms of dystonia. While we did not observe differences across dystonia

subtypes (Supplementary Fig. 1), future studies in larger patient cohorts would are warranted.

Finally, as many findings in Fig. 1 do not survive corrections for multiple comparisons, we

suggest interpretation of results with caution. Despite this, many of our findings related to

neuronal correlates are generally in line with previous literature, especially related to oscillatory

correlates of PD and dystonia.

Conclusion

345   We substantiated claims of hypo- versus hyperfunctional GPi output in PD and dystonia, while

346   furthermore providing cellular-level validation of the pathological nature of theta and low-beta

347   oscillations in respective disorders. Such circuit changes may be underlain by disease-related

348   differences in plasticity of striato-pallidal synapses, which are seemingly less plastic and/or

349   respond slower to change in dystonia compared to PD.

Acknowledgements

The authors thank the patients for their participation.

Author Roles

(1) Research project: A. Conception, B. Organization, C. Execution; (2) Statistical analysis: A. Design, B. Execution, C. Review and critique; (3) Manuscript preparation: A. Writing of the first draft, B. Review and critique.

S.S: 1C, 2A, 2B, 3A; K.A.S: 1C, 2A, 2B, 3A; L.A.S: 1C, 2A, 2C, 3A; C.F: 1C, 3B; E.H: 1C, 3A; A.A.K: 1B, 3B; M.H: 1B, 3B; S.K.K: 1B, 3B; A.M.L: 1B, 3B; A.F: 1B, 3B; W.D.H: 1A, 1B, 1C, 2B, 3B; L.M: 1A, 1B, 1A, 1C, 2B, 3B.

Fig 1 – GPi spiketrain feature analyses and clinical correlates of PD and dystonia. (A) With respect to (A) rate-
based spiketrain features, firing rate was greater in PD while burst index (BI) and coefficient of variation (CV) were
greater in dystonia; whereas no differences were found for (B) oscillatory spiketrain features for theta, alpha, low
beta, high beta frequencies. MWU statistical results depicted are not corrected for multiple comparisons; after
correction using the Bonferroni method, only CV and BI results remain significant (please see Supplementary Table
3). (C) In PD, the power of low beta spiketrain oscillations positively correlated (Spearman correlation) with
symptom severity; in dystonia, neuronal firing rate negatively correlated with symptom severity, whereas CV and
the power of theta spiketrain oscillations positively correlated with symptom severity. Depicted scatterplots are
results that were significant before correction for multiple comparisons; however, none of the results persist after
Benjamini-Hochberg correction for false discovery rate (please see Supplementary Table 4).

Fig 2 – Long-term and short-term effects of HFS on striato-pallidal plasticity in PD and dystonia. (A)
Schematic of the plasticity protocol to assess long-term plasticity via fEP amplitude comparisons pre- versus post-
HFS and short-term plasticity via fEP dynamics during HFS. (B) Highlights example fEP traces for measuring long-
term plasticity pre- versus post-HFS, with (C) displaying group-level fEP amplitudes pre- versus post-HFS across
diseases. (D) Illustrates the amount of plasticity (i.e., percentage change in fEP amplitudes pre- versus post-HFS) in
both PD and dystonia, with PD showing higher levels of plasticity. (E) Provides an example of fEP traces during
HFS for assessing short-term plasticity, with (F) depicting group-level decay rates of fEP amplitudes using an
exponential fit on the fEP amplitudes over the first 5 stimulus pulses across diseases. (G) Shows the half-life of the
fitted exponential (i.e., rate of attenuation of fEP amplitudes) between PD and dystonia, with PD demonstrating
faster fEP attenuation.

References

Jankovic J. Parkinson's disease: clinical features and diagnosis. *J Neurol Neurosurg Psychiatry*. 2008;79:368–76.

Albanese A, Bhatia K, Bressman SB, *et al.* Phenomenology and classification of dystonia: A consensus update. *Mov Disord*. 2013;28:863–73.

DeLong MR, Wichmann T. Circuits and Circuit Disorders of the Basal Ganglia. *Arch Neurol*. 2007;64:20–4.

DeLong MR. Primate models of movement disorders of basal ganglia origin. *Trends Neurosci*. 1990;13:281–5.

Naumann M, Pirker W, Reiners K, *et al.* Imaging the pre- and postsynaptic side of striatal dopaminergic synapses in idiopathic cervical dystonia: A SPECT STUDY Using [123I] epidepride and [123I] β-CIT. *Mov Disord*. 1998;13:319–23.

Simonyan K, Berman BD, Herscovitch P, *et al.* Abnormal Striatal Dopaminergic Neurotransmission during Rest and Task Production in Spasmodic Dysphonia. *J Neurosci*. 2013;33:14705–14.

Simonyan K, Cho H, Hamzehei Sichani A, *et al.* The direct basal ganglia pathway is hyperfunctional in focal dystonia. *Brain*. 2017;140:3179–90.

Starr PA, Rau GM, Davis V, *et al.* Spontaneous Pallidal Neuronal Activity in Human Dystonia: Comparison With Parkinson's Disease and Normal Macaque. *J Neurophysiol*. 2005;93:3165–76.

Tang JKH, Moro E, Mahant N, *et al.* Neuronal firing rates and patterns in the globus pallidus internus of patients with cervical dystonia differ from those with Parkinson's disease. *J Neurophysiol*. 2007;98:720–9.

Hutchison WD, Lang AE, Dostrovsky JO, *et al.* Pallidal neuronal activity: Implications for models of dystonia. *Ann Neurol*. 2003;53:480–8.

Neumann W-J, Degen K, Schneider G-H, *et al.* Subthalamic synchronized oscillatory activity correlates with motor impairment in patients with Parkinson's disease. *Mov Disord*. 2016;31:1748–51.

Neumann W-J, Horn A, Ewert S, *et al.* A localized pallidal physiomarker in cervical dystonia. *Ann Neurol*. 2017;82:912–24.

Stefani A, Cerroni R, Mazzone P, *et al.* Mechanisms of action underlying the efficacy of deep brain stimulation of the subthalamic nucleus in Parkinson's disease: central role of disease severity. *Eur J Neurosci*. 2019;49:805–16.

14 Ruge D, Cif L, Limousin P, *et al.* Shaping reversibility? Long-term deep brain stimulation in
dystonia: the relationship between effects on electrophysiology and clinical symptoms.
Brain. 2011;134:2106–15.

15 Milosevic L, Gramer R, Kim TH, *et al.* Modulation of inhibitory plasticity in basal ganglia
output nuclei of patients with Parkinson's disease. *Neurobiol Dis*. 2019;124:46–56.

16 Liu LD, Prescott IA, Dostrovsky JO, *et al.* Frequency-dependent effects of electrical
stimulation in the globus pallidus of dystonia patients. *J Neurophysiol*. 2012;108:5–17.

17 Hutchison WD, Lozano AM, Davis KD, *et al.* Differential neuronal activity in segments of
globus pallidus in Parkinson's disease patients. *Neuroreport*. 1994;5:1533–7.

18 Milosevic L, Kalia SK, Hodaie M, *et al.* Neuronal inhibition and synaptic plasticity of basal
ganglia neurons in Parkinson's disease. *Brain*. 2018;141:177–90.

19 Kaneoke Y, Vitek JL. Burst and oscillation as disparate neuronal properties. *J Neurosci
Methods*. 1996;68:211–23.

20 Hutchison WD, Lozano AM, Davis KD, *et al.* Differential neuronal activity in segments of
globus pallidus in Parkinson's disease patients. *Neuroreport*. 1994;5:1533–7.

21 Steiner LA, Kühn AA, Geiger JRP, *et al.* Persistent synaptic inhibition of the subthalamic
nucleus by high frequency stimulation. *Brain Stimulat*. 2022;15:1223–32.

22 Hallett M. The Neurophysiology of Dystonia. *Arch Neurol*. 1998;55:601–3.

23 Hallett M. Physiology of Basal Ganglia Disorders: An Overview. *Can J Neurol Sci*.
1993;20:177–83.

24 McGregor MM, Nelson AB. Circuit Mechanisms of Parkinson's Disease. *Neuron*.
2019;101:1042–56.

25 Muralidharan A, Jensen AL, Connolly A, *et al.* Physiological changes in the pallidum in a
progressive model of Parkinson's disease: Are oscillations enough? *Exp Neurol*.
2016;279:187–96.

26 Scherer M, Steiner LA, Kalia SK, *et al.* Single-neuron bursts encode pathological
oscillations in subcortical nuclei of patients with Parkinson's disease and essential tremor.
Proc Natl Acad Sci. 2022;119:e2205881119.

27 Silberstein P, Kühn AA, Kupsch A, *et al.* Patterning of globus pallidus local field potentials
differs between Parkinson's disease and dystonia. *Brain J Neurol*. 2003;126:2597–608.

28 Wang DD, de Hemptinne C, Miocinovic S, *et al.* Pallidal Deep-Brain Stimulation Disrupts
Pallidal Beta Oscillations and Coherence with Primary Motor Cortex in Parkinson's Disease.
J Neurosci Off J Soc Neurosci. 2018;38:4556–68.

29 Steiner LA, Kuehn AA, Geiger JR, *et al.* Persistent synaptic inhibition of the subthalamic
nucleus by high frequency stimulation. 2021. https://doi.org/10.1101/2021.10.20.465131

30 Yoshida M, Precht W. Monosynaptic inhibition of neurons of the substantia nigra by
caudatonigral fibers. *Brain Res*. 1971;32:225–8.

31 Precht W, Yoshida M. Blockage of caudate-evoked inhibition of neurons in the substantia
nigra by picrotoxin. *Brain Res*. 1971;32:229–33.

32 Prescott IA, Dostrovsky JO, Moro E, *et al.* Reduced paired pulse depression in the basal
ganglia of dystonia patients. *Neurobiol Dis*. 2013;51:214–21.

33 Ueki Y, Mima T, Kotb MA, *et al.* Altered plasticity of the human motor cortex in
Parkinson's disease. *Ann Neurol*. 2006;59:60–71.

34 Tamura Y, Ueki Y, Lin P, *et al.* Disordered plasticity in the primary somatosensory cortex in
focal hand dystonia. *Brain*. 2009;132:749–55.

35 Chen R, Udupa K. Measurement and modulation of plasticity of the motor system in humans
using transcranial magnetic stimulation. *Motor Control*. 2009;13:442–53.

36 Peterson DA, Sejnowski TJ, Poizner H. Convergent evidence for abnormal striatal synaptic
plasticity in dystonia. *Neurobiol Dis*. 2010;37:558–73.

37 Quartarone A, Pisani A. Abnormal plasticity in dystonia: Disruption of synaptic homeostasis.
Neurobiol Dis. 2011;42:162–70.

38 Chen R. Studies of human motor physiology with transcranial magnetic stimulation. *Muscle
Nerve*. 2000;23:S26–32.

39 Siebner HR, Tormos JM, Ceballos-Baumann AO, *et al.* Low-frequency repetitive
transcranial magnetic stimulation of the motor cortex in writer's cramp. *Neurology*.
1999;52:529–37.

40 Tamura Y, Ueki Y, Lin P, *et al.* Disordered plasticity in the primary somatosensory cortex in
focal hand dystonia. *Brain J Neurol*. 2009;132:749–55.

41 Köhling R, Koch U-R, Hamann M, *et al.* Increased excitability in cortico-striatal synaptic
pathway in a model of paroxysmal dystonia. *Neurobiol Dis*. 2004;16:236–45.

42 Martella G, Tassone A, Sciamanna G, *et al.* Impairment of bidirectional synaptic plasticity in
the striatum of a mouse model of DYT1 dystonia: role of endogenous acetylcholine. *Brain J
Neurol*. 2009;132:2336–49.

43 Calabresi P, Ghiglieri V, Mazzocchetti P, *et al.* Levodopa-induced plasticity: a double-edged
sword in Parkinson's disease? *Philos Trans R Soc B Biol Sci*. 2015;370. doi:
10.1098/rstb.2014.0184

44 Prescott IA, Liu LD, Dostrovsky JO, *et al.* Lack of depotentiation at basal ganglia output
neurons in PD patients with levodopa-induced dyskinesia. *Neurobiol Dis*. 2014;71:24–33.

45 Krauss JK. Deep Brain Stimulation for Dystonia in Adults. *Stereotact Funct Neurosurg*.
2002;78:168–82.

46 Barow E, Neumann W-J, Brücke C, *et al.* Deep brain stimulation suppresses pallidal low
frequency activity in patients with phasic dystonic movements. *Brain*. 2014;137:3012–24.

47 Ruge D, Tisch S, Hariz MI, *et al.* Deep brain stimulation effects in dystonia: Time course of
electrophysiological changes in early treatment. *Mov Disord*. 2011;26:1913–21.

48 Hammond C, Bergman H, Brown P. Pathological synchronization in Parkinson's disease:
networks, models and treatments. *Trends Neurosci*. 2007;30:357–64.

49 Follett KA, Weaver FM, Stern M, *et al.* Pallidal versus subthalamic deep-brain stimulation
for Parkinson's disease. *N Engl J Med*. 2010;362:2077–91.

50 Anderson VC, Burchiel KJ, Hogarth P, *et al.* Pallidal vs Subthalamic Nucleus Deep Brain
Stimulation in Parkinson Disease. *Arch Neurol*. 2005;62:554–60.

51 Bejjani B-P, Dormont D, Pidoux B, *et al.* Bilateral subthalamic stimulation for Parkinson's
disease by using three-dimensional stereotactic magnetic resonance imaging and
electrophysiological guidance. *J Neurosurg*. 2000;92:615–25.

52 Spix TA, Nanivadekar S, Toong N, *et al.* Population-specific neuromodulation prolongs
therapeutic benefits of deep brain stimulation. *Science*. 2021;374:201–6.

53 Mastro KJ, Zitelli KT, Willard AM, *et al.* Cell-specific pallidal intervention induces long-
lasting motor recovery in dopamine-depleted mice. *Nat Neurosci*. 2017;20:815–23.

54 Johnson V, Wilt R, Gilron R, *et al.* Embedded adaptive deep brain stimulation for cervical
dystonia controlled by motor cortex theta oscillations. *Exp Neurol*. 2021;345:113825.

55 Little S, Beudel M, Zrinzo L, *et al.* Bilateral adaptive deep brain stimulation is effective in
Parkinson's disease. *J Neurol Neurosurg Psychiatry*. 2016;87:717–21.

56 Versantvoort EM, Dietz BE, Mugan D, *et al.* Evoked compound action potential (ECAP)-
controlled closed-loop spinal cord stimulation in an experimental model of neuropathic pain
in rats. *Bioelectron Med*. 2024;10:2.

57 Reese R, Fasano A, Knudsen K, *et al.* Full Parkinsonian Triad Induced by Pallidal
High Frequency Stimulation in Cervical Dystonia. *Mov Disord Clin Pract*. 2014;2:99–101.

58 Esposito M, Dubbioso R, Peluso S, *et al.* Cervical dystonia patients display subclinical gait
changes. *Parkinsonism Relat Disord*. 2017;43:97–100.

516