1

Local organization of spatial and shape information during working memory in the

primate prefrontal cortex

Yunyi Sun[1], Wenhao Dang[2], Rye G. Jaffe[2], Christos Constantinidis[2,3,4]

1. Department of Biostatistics, Vanderbilt University Medical Center, Nashville TN
37203, USA.
2. Department of Biomedical Engineering, Vanderbilt University, Nashville TN 37235,
USA.
3. Neuroscience Program, Vanderbilt University, Nashville TN 37235, USA.
4. Department of Ophthalmology and Visual Sciences, Vanderbilt University Medical
Center, Nashville TN 37232, USA.
16
17

18 Abbreviated Title: PFC Organization

20 Address correspondence to:
21 Christos Constantinidis, PhD.
22 Christos.Constantinidis.1@vanderbilt.edu
23

24 The authors report no conflicts of interest during the research conducted.
25
26

ABSTRACT

While the current understanding of sensory and motor cortical areas has been defined

topographical maps across the surface of these areas, higher cortical areas, such as the

prefrontal cortex, seem to lack an equivalent organization, with only limited evidence of

functional clustering of neurons with similar stimulus properties. We sought to examine

whether neurons that represent similar spatial and object information are clustered in the

monkey prefrontal cortex and whether such an organization only emerges as a result of

training. We analyzed neurophysiological recordings from male macaque monkeys

before and after they were trained to perform cognitive tasks. Neurons with similar

spatial or shape selectivity were more likely than chance to be encountered at short

distances from each other. This pattern of organization was present even in naïve

animals, prior to any cognitive training. Our results reveal that prefrontal microstructure

automatically supports orderly representations of spatial and object information.

INTRODUCTION

The prefrontal cortex (PFC) has long been recognized as the seat of higher cognitive functions, based on a plethora of evidence, including lesion, anatomical, and neurophysiological experiments in non-human primates (Goldman-Rakic 1988). A population of prefrontal neurons generate persistent activity selective for the properties of stimuli held in memory (Constantinidis et al. 2018). Computational models typically simulate such persistent activity by networks of units with recurrent connections between units with similar tuning for spatial location; this organizations allows activity initiated by a stimulus to be maintained in the network during working memory (Wimmer et al. 2014; Hart and Huk 2020).

The prefrontal cortex lacks an obvious retinotopic organization (Kiani et al. 2015; Leavitt et al. 2017), or other type of systematic topography of sensory information that would be analogous to the orderly arrangement of units in these artificial networks – or in the sensory and motor cortices. Only some local organization has been previously described in the prefrontal cortex, for spatial information, with nearby neurons representing the same or adjacent spatial locations (Constantinidis et al. 2001; Arnsten 2013). It is unclear therefore how neurons with similar selectivity could organize into functional circuits.

Newer theories of working memory, motivated primarily by results of human imaging studies have proposed that working memory information is represented in the sensory cortex, instead (Harrison and Tong 2009; Albers et al. 2013; Ester et al. 2013; Xing et al. 2013; Sreenivasan et al. 2014). According to this viewpoint, prefrontal cortex

plays a control or supervisory role, perhaps highlighting the locations of remembered

stimuli, rather than maintaining their identity in working memory itself (Curtis and

D'Esposito 2003; Tong and Pratte 2012; D'Esposito and Postle 2015). It is possible,

therefore, that any local organization within the prefrontal cortex is limited to spatial

information, alone.

We were thus motivated to examine the local organization of prefrontal neurons

for the maintenance of spatial and object information in memory. We first replicated the

analysis of spatial information of prior studies (Constantinidis *et al.* 2001), from a

different set of experimental data. We then applied the same analysis to object

information. Additionally, we examined how this information is altered by training to

perform a working memory task, by analyzing data recorded from monkeys naïve to

training, and after they were trained to perform spatial and object working memory tasks.

METHODS

Animals: Data obtained from six male rhesus monkeys (*Macaca mulatta*), ages 5–9 years

old, weighing 5–12 kg, as previously documented (Riley et al. 2018), were analyzed in

this study. None of these animals had any prior experimentation experience at the onset

of our study. Monkeys were either single-housed or pair-housed in communal rooms with

sensory interactions with other monkeys. All experimental procedures followed

guidelines set by the U.S. Public Health Service Policy on Humane Care and Use of

Laboratory Animals and the National Research Council's Guide for the Care and Use of

Laboratory Animals and were reviewed and approved by the Wake Forest University

Institutional Animal Care and Use Committee.

Experimental setup: Monkeys sat with their heads fixed in a primate chair while viewing

a monitor positioned 68 cm away from their eyes with dim ambient illumination and were

required to fixate on a 0.2° white square appearing in the center of the screen. In order to

receive a liquid reward (typically fruit juice), the animals maintained fixation on the

square while visual stimuli were presented either at a peripheral location or over the

fovea. Any break of fixation immediately terminated the trial and no reward was given.

Eye position was sampled at 240 Hz, monitored on line, and recorded with an infrared

eye position scanning system (model RK-716; ISCAN, Burlington, MA). The visual

stimulus display, monitoring of eye position, and synchronization of stimuli with

neurophysiological data were performed with in-house software (Meyer and

Constantinidis 2005) implemented on the MATLAB environment (Mathworks, Natick,

MA), utilizing the Psychophysics Toolbox (Brainard 1997).

<u>Pre-training task:</u> Following a brief period of fixation training and acclimation to the

stimuli, monkeys were required to fixate on a center position while stimuli were

displayed on the screen. The stimuli shown in the pre-training passive spatial task were

white 2° squares, presented in one of nine possible locations arranged in a 3 × 3 grid with

10° distance between adjacent stimuli. The stimuli shown in the pre-training passive

feature task were white 2° geometric shapes drawn from a set comprising a circle,

diamond, the letter H, the hashtag symbol, the plus sign, a square, a triangle, and an

inverted Y-letter. These stimuli could also be presented in one of nine possible locations

arranged in a 3 × 3 grid with 10° distance between adjacent stimuli.

Presentation began with a fixation interval of 1 s where only the fixation point

was displayed, followed by 500 ms of stimulus presentation (referred to hereafter as cue),

followed by a 1.5 s "delay" interval (referred to hereafter as delay1) where, again, only

the fixation point was displayed. A second stimulus (referred to hereafter as sample) was

subsequently shown for 500 ms. For the spatial stimulus set, this second stimulus would

be either identical in location to the initial stimulus, or diametrically opposite the first

stimulus. For the feature set, this second stimulus would appear in the same location to

the initial stimulus and would either be an identical shape or the corresponding non-

match shape (each shape was paired with one non-match shape). Only one nonmatch

stimulus was paired with each cue, so that the number of match and nonmatch trials were

balanced in each set. In both the spatial and feature task, this second stimulus display was

followed by another "delay" period (delay2) of 1.5 s where only the fixation point was

displayed. Successful holding of fixation until the end of the second delay period resulted

in reward delivery. The location and identity of stimuli was of no behavioral relevance to

the monkeys during the "pre-training" phase, as fixation was the only necessary action

for obtaining reward.

<u>Post-training task:</u> Four of the six monkeys were trained to complete spatial and feature

working memory tasks. These tasks involved presentation of identical stimuli as the

spatial and feature tasks during the "pre-training" phase, but now monkeys were required

to remember the spatial location and/or shape of the first presented stimulus, and report

whether the second stimulus was identical to the first or not, via saccading to one of two

target stimuli (green for matching stimuli, blue for non-matching). Each target stimulus

could appear at one of two locations orthogonal to the cue/sample stimuli, pseudo-

randomized in each trial.

<u>Surgery and neurophysiology:</u> A 20 mm diameter craniotomy was performed over the

PFC and a recording cylinder was implanted over the site. The location of the cylinder

was visualized through anatomical magnetic resonance imaging (MRI) and stereotaxic

coordinates post-surgery. Electrode penetrations were aligned to the MRI image and

mapped onto the cortical surface.

<u>Neuronal recordings:</u> Neural recordings were carried before and after training in the

working memory tasks. The se of recordings analyzed here was previously used to

determine the properties of neurons before and after training in different PFC

subdivisions and more details about the properties of these neurons can be found in

previous reports (Meyer et al. 2011; Riley *et al.* 2018). Extracellular recordings were

performed with multiple microelectrodes that were either glass- or epoxylite-coated

tungsten, with a 250 μm diameter and 1–4 MΩ impedance at 1 kHz (Alpha-Omega

Engineering, Nazareth, Israel). A Microdrive system (EPS drive, Alpha- Omega

Engineering) advanced arrays of up to 8-microelectrodes, spaced 0.2–1.5 mm apart,

through the dura and into the PFC. The signal from each electrode was amplified and

band-pass filtered between 500 Hz and 8 kHz while being recorded with a modular data

acquisition system (APM system, FHC, Bowdoin, ME). Waveforms that exceeded a user-

defined threshold were sampled at 25 μs resolution, digitized, and stored for off-line

analysis. Neurons were sampled in an unbiased fashion, collecting data from all units

isolated from our electrodes, with no regard to the response properties of the isolated

neurons. A semi-automated cluster analysis relied on the KlustaKwik algorithm, which

applied principal component analysis of the waveforms to sort recorded spike waveforms

into separate units (Harris et al. 2000). To ensure a stable firing rate in the analyzed

recordings, we identified recordings in which a significant effect of trial sequence was

evident at the baseline firing rate (ANOVA, $p < 0.05$), e.g., due to a neuron disappearing

or appearing during a run, as we were collecting data from multiple electrodes. Data from

these sessions were truncated so that analysis was only performed on a range of trials

with stable firing rate. Less than 10% of neurons were corrected in this way. Identical

data collection procedures, recording equipment, and spike sorting algorithms were used

before and after training in order to prevent any confounds.

<u>Data analysis</u>. Analysis was implemented with the MATLAB computational environment

(R2020-2023, Mathworks, Natick, MA). To obtain tuning similarity analysis by optimal

stimuli, we first identified neurons with significant tuning for the stimulus set being

examined. A one-way ANOVA at the $\alpha=0.05$ significance level was employed,

separately for each epoch of the task (fixation, cue, delay1, sample, delay2 in Fig. 1).

The preferred stimulus in each task epoch was then defined for the neuron as the stimulus

(location/shape) that elicited the highest average firing rate in the corresponding epoch.

We examined pairs of nearby neurons recorded simultaneously from separate

microelectrodes, placed together at distances of 0.2 – 1.5 mm from each other. For each

task epoch, we identified such pairs both neurons of which were selective for the stimuli,

and determined if they had the same preferred (best) stimulus or not. The proportions of

such neuron pairs among all tuned neuron pairs were determined in each epoch and we

used a nonparametric test to test the statistical significance of this proportion. More

specifically, we randomly paired cells in our pool that were not recorded simultaneously

(from nearby locations), and then repeated the same analysis described above to

determine a simulated proportion of cells with the same preference. This random pairing

procedure was repeated 1000 times to get a full null distribution. We then counted the

number of occurrences in the null distribution for more extreme values compared to the

empirical value and divided that number by 1000 to get a probability measure, which

would be the p-value of the empirical measurement.

In addition to determining the best stimulus of each pair, we also analyzed the

tuning similarity of nearby neurons by quantifying the Euclidean distance defined by

their average response to all stimuli used in each task. More specifically, for a given

neuron, a trial-average firing rate vector was created by averaging the trials recorded with

each stimulus, resulting in a 9 by 1 vector for the spatial test, and an 8 by 1 vector for the

feature set. Each neuron's firing rate vector for a certain task epoch was then normalized

by the highest value in the vector, to control for firing rate difference between neurons.

For a given nearby neuron pair, and for a given task epoch, we calculated the Euclidean

distance in the 9- (or 8-) dimensional space using the built-in MATLAB function

"pdist2". If both cells in the pair displayed selectivity in more than one task epoch, the

averaged tuning distance for all task epochs in which significant selectivity was present

was used. We used a nonparametric test for testing the statistical significance of the

Euclidean distance measurement. We again randomly paired neurons from different

recordings, without having been recorded simultaneously. For a given neuron in this cell

pool, we randomly chose one of the four epochs to get data from. This is the first source

of our randomness introduced to the bootstrap analysis. Then we constructed a random

cell pair pool containing all possible combinations of cells in the pool. And lastly we

randomly draw the same number of pairs of cells as in the empirical pair dataset to

calculate one value of the mean Euclidean distance for the null distribution. We had 1000

such iterations to generate the full null distribution.

To visualize the representational similarity of different stimuli at the population

level, we used the MATLAB function "linkage" and "dendrogram". Specifically, we

created an nine by N matrix (or eight by N matrix for the shape stimuli), where N is the

number of unique neurons in the empirical neighboring-neuron pool as described above.

Each element in the matrix represents the average neural response to 9 (or 8) stimuli. We

then use the MATLAB function "pdist2" to create an 9 x 9 (or 8 by 8) pairwise distance

matrix representing the pairwise distance between the neuronal representation of the

stimuli. Lastly, the distance matrix served as the input to the MATLAB "linkage"

function to create a hierarchical cluster tree.

RESULTS

Database

Extracellular neurophysiological recordings were collected from the lateral PFC of six

male monkeys before and after they were trained to perform the match/nonmatch

working memory tasks (Meyer *et al.* 2011; Riley *et al.* 2018). The task required them to

view two stimuli appearing in sequence with an intervening delay period between them,

and after a second delay period to report whether or not the second stimulus was identical

to the first (Fig. 1). The two stimuli could differ in terms of their location (spatial task,

Fig. 1A-B) or shape (feature task, Fig. 1C-D). If the second stimulus was identical to the

first, monkeys were required to saccade towards a green choice target that appeared at the

end of the trial. Otherwise, they had to saccade to a blue target. The locations of the

choice targets varied randomly from trial to trial and the monkey could not plan a motor

response prior to their appearance.

We collected neurophysiological data before and after the monkeys learned to

perform these tasks. A total of 1617 cells from six monkeys were recorded while the

animals viewed the spatial stimuli passively prior to training, and 1493 cells from five

monkeys were recorded while the animals viewed the feature stimuli passively, prior to

training to perform the task ("pre-training" phase). A total of 1104 cells from three of the

same monkeys were collected after the monkeys were trained and were performing the

spatial working memory task. Another 1091 cells from two of the same monkeys were

collected after the monkeys were trained and performed the feature working memory

tasks ("post-training" phase).

Organization of Spatial Selectivity

We first identified neurons tuned for the spatial location of stimuli, evidenced by

significant selectivity for the nine spatial stimuli (1-way ANOVA, p<0.05). Percentages

of neurons that exhibited such significant selectivity at different time epochs of the task

(plus the baseline, fixation period, as a control) are shown in Fig. 2. Spatial selectivity

generally increased after training, as we have reported previously (Riley *et al.* 2018), with

approximately 15% of all neurons recorded exhibiting selectivity during any task period.

We relied on this dataset to examine how neurons selective for spatial stimuli are

arranged in local prefrontal circuits. We thus examined pairs of neurons recorded

simultaneously from different microelectrodes, spaced 0.2 – 1.5 mm apart from each

(mean distance = 0.77 mm). For 499 such pairs (222 recorded before and 277 recorded

after training), both neurons exhibited significant selectivity in the same task epoch. We

could thus test that neurons that had the same spatial preference (best response for the

same location) were encountered more often than would be expected by chance (Fig.2C),

which was evaluated using a bootstrap task, randomly pairing neurons recorded at

different sessions and random positions. We indeed determined that neurons with the

same preference were significantly higher than expected at multiple task epochs: cue, and

sample period, prior to training (p<0.001 in each case), and cue and delay period after

training (p=0.012 and p<0.001, respectively).

This analysis considers only the best location, however, the entire tuning function

of a neuron provides a more complete picture of neuronal tuning. We thus refined our

analysis to consider the vector defined by firing rates for all eight locations for each

neuron. We then computed the Euclidean distance between the vectors of two neurons,

examining again only task epochs in which both neurons exhibited significant spatial

selectivity. For pairs with significant selectivity in more than one task epoch, we

computed distances in each epoch and then averaged them to obtain a single distance

value for each neuron. To ensure that neurons with higher firing rates did not appear to

exhibit longer distances, we normalized firing rates before computing distances. The

mean distance computed in this manner was 1.02 prior to training and 1.11 after training

(Fig. 3). Both values were significantly lower than would be expected by chance

(permutation test, $p < 0.001$ in each case).

271   The pattern of responses to different stimuli also allowed us to determine how

different spatial locations are represented across neurons. A hierarchical cluster analysis

mostly captured the topology of stimulus location, with adjacent locations being

represented by nearby clusters (Fig. 4). Prior to training, the cluster corresponding to the

shorter distance in stimulus activity involved locations 4 and 5, which represent adjacent

stimuli on the screen (Fig. 1). Successive, higher-order clusters involved stimuli 6, 3, 9,

7, 1, 8, and 2, each one of which were adjacent with the preceding cluster of locations. A

similar ordering of stimulus locations was present after training, although the absolute

order of cluster distances changed somewhat.

Organization of Feature Selectivity

These results of the spatial analysis mostly replicated earlier studies which suggested a

local organization of spatial information across the surface of the prefrontal cortex, using

different spatial working memory tasks (Constantinidis *et al.* 2001). Since it has been

argued that spatial representations are fundamentally different than object working

memory (Curtis and D'Esposito 2003; Tong and Pratte 2012; D'Esposito and Postle

2015), we were interested to test whether neuronal selectivity for shapes exhibits some

organization or not.

We thus repeated our analysis of neuronal selectivity for shape information. The

percentage of neurons with significant selectivity for the eight geometric shapes of our

dataset evaluated with a 1-way ANOVA test ($p<0.05$) is shown in Fig. 5. Although shape

selectivity was lower compared to spatial selectivity, neurons selective for shape were

found both before and after training, with approximately 10% of all neurons recorded

exhibiting shape selectivity at any task epoch.

A total of 242 pairs of neurons were recorded simultaneously from separate

electrodes spaced 0.2 – 1.5 mm from each other, both neurons of which exhibited shape

selectivity in the same task epoch (Fig.5C). We again examined whether these neurons

were more likely to exhibit the same preferred stimulus than would be expected by

chance. For this dataset, a higher than chance same-shape preference was observed in the

cue period, after training (permutation test, $p=0.043$).

We compared the overall similarity of shape tuning between simultaneously

recorded neurons by computing their normalized Euclidean distance, as we did for the

spatial stimuli. The mean Euclidean distance was 0.86 prior to training and 0.88 after

training. These distances were significantly smaller than would be expected by chance

(permutation test, $p<0.001$, in each case). These results suggest that shape selectivity too,

is not represented randomly across the surface of the cortex. Prefrontal neurons tuned for

stimulus shape are present prior to any training as well as after training and organized in

patches of cortex with similar shape selectivity.

Finally, we determined how different shapes are represented across the population

of prefrontal neurons (Fig. 7). There is no obvious way to group the different shapes in

analogy to spatial locations. Clustering was generally similar before and after training,

although the absolute order of cluster distances changed somewhat after training.

DISCUSSION

Our analysis revealed a local organization of both spatial and non-spatial information

across the surface of the prefrontal cortex, with neurons 1.5 mm of each other

representing significantly more similar stimuli. These results are consistent with previous

studies that have suggested a local organization of spatial representations across the

prefrontal cortex, in monkeys trained to perform spatial working memory tasks

(Constantinidis *et al.* 2001; Leavitt *et al.* 2017). Moreover, we now reveal a similar

pattern of local organization for object representation, as well. Furthermore, we now

document that this organization is largely present even prior to any training to perform a

working memory task that required these stimuli to be recognized and maintained in

memory.

Organization of information across the prefrontal cortex

The prefrontal cortex is critical for higher cognitive functions, including the maintenance

of information in working memory (D'Esposito and Postle 2015; Riley and Constantinidis

2016). Specifically, this information is thought to be maintained through persistent

discharges in a network of brain areas centered on the prefrontal cortex (Constantinidis *et*

al. 2018). Computational models suggest that persistent activity is sustained in neural

circuits by virtue of re-entrant connections between neurons with similar tuning for

stimulus properties (Compte et al. 2000; Wimmer *et al.* 2014). Neural activity generated

by a stimulus presentation is maintained in the network, which behaves as a continuous

attractor (Seung et al. 2000). Until now however, the specific organization of these

neuronal networks across the surface of the prefrontal cortex has remained unresolved.

No obvious, retinotopic organization of visual space has been described across the

surface of the prefrontal cortex, or other higher-order cortical areas implicated in the

maintenance of working memory, such as the posterior parietal cortex (Qi et al. 2015).

Some indications exist for at least some local PFC organization. For example, electrode

penetrations that descent into the principal sulcus, parallel to its surface, suggest an

orderly progression of receptive field locations (Arnsten 2013). Our current results

suggest that selectivity for stimulus properties is an organizing principle in the neural

circuits of the prefrontal cortex, as speculated by early models of the cellular basis of

working memory (Goldman-Rakic 1995).

Other findings also suggest at least some coarse localization of sensory

information across domains. Whether object working memory is localized in a different

subdivision of the prefrontal cortex than spatial working memory (ventral vs. dorsal) has

been a matter of debate (Wilson et al. 1993; Rao et al. 1997), but at least a quantitative

difference seems to be evident, with spatial information more prevalent in the

dorsolateral prefrontal cortex than the ventrolateral prefrontal cortex (Meyer *et al.* 2011;

Constantinidis and Qi 2018). Some organization for the time course of activation has also

been revealed across cortical layers: neurons activated by visual stimuli are more likely to

be encountered in intermediate layers; neurons with delay period activity in superficial;

neurons with motor responses in deep, though this relative segregation is quantitative

rather than qualitative (Markowitz et al. 2015; Zhu et al. 2023). Finally, an anterior-

posterior gradient of stimulus properties has also been described so that posterior neurons

are more selective for the sensory attributes of stimuli no matter what their context,

whereas more anterior neurons are much more likely to become activated only in the

context of learned tasks (Riley et al. 2017; Riley *et al.* 2018).

Training effects

The activity of prefrontal neurons is highly plastic and responding to abstract variables of

the task being executed (Fuster et al. 2000; Wallis et al. 2001; Mante et al. 2013).

Neurophysiological studies of training have revealed that learning to perform cognitive

tasks generally increases responsiveness to stimuli that are relevant for the task execution

(Meyers et al. 2012; Constantinidis and Klingberg 2016). Prefrontal neurons, and

particularly those in more anterior areas, become responsive to stimuli that previously

failed to elicit a response (Qi et al. 2011; Riley *et al.* 2018). Recent human imaging

results parallel these findings, confirming appearance of stimulus selectivity for learned

stimuli after training (Miller et al. 2022). A variety of other changes have also been

reported, including in variability and correlation of prefrontal discharges (Qi and

Constantinidis 2012).

375   In view of these findings, it appeared possible that the local organization of the

prefrontal cortex with respect to stimulus selectivity might only emerge after training.

Contrary to this expectation, we found neurons with similar selectivity at nearby

locations in the cortex, even in naïve monkeys, prior to any training. This result is

consistent with previous findings suggesting that underlying network circuitry appears to

connect neurons with similar properties, and that these functional modules of neurons

remain fairly stable across different behavioral contexts or tasks (Kiani *et al.* 2015).

ACKNOWLEDGMENTS
384
Research reported in this paper was supported by the NIH National Eye Institute under

award number R01 EY017077 and by the National Institute of Mental Health under

award number R01 MH116675. We wish to acknowledge Chrissy Suell for technical

help; Junda Zhu, and Zhengyang Wang for helpful comments on the manuscript.

REFERENCES

Albers AM, Kok P, Toni I, Dijkerman HC, de Lange FP. 2013. Shared representations for
working memory and mental imagery in early visual cortex. Current biology : CB.
23:1427-1431.

Arnsten AF. 2013. The neurobiology of thought: the groundbreaking discoveries of
Patricia Goldman-Rakic 1937-2003. Cereb Cortex. 23:2269-2281.

Brainard DH. 1997. The Psychophysics Toolbox. Spat Vis. 10:433-436.

Compte A, Brunel N, Goldman-Rakic PS, Wang XJ. 2000. Synaptic mechanisms and
network dynamics underlying spatial working memory in a cortical network model.
Cereb Cortex. 10:910-923.

Constantinidis C, Franowicz MN, Goldman-Rakic PS. 2001. Coding specificity in
cortical microcircuits: a multiple electrode analysis of primate prefrontal cortex. J
Neurosci. 21:3646-3655.

Constantinidis C, Funahashi S, Lee D, Murray JD, Qi XL, Wang M, Arnsten AFT. 2018.
Persistent Spiking Activity Underlies Working Memory. J Neurosci. 38:7020-7028.

Constantinidis C, Klingberg T. 2016. The neuroscience of working memory capacity and
training. Nat Rev Neurosci. 17:438-449.

Constantinidis C, Qi XL. 2018. Representation of Spatial and Feature Information in the
Monkey Dorsal and Ventral Prefrontal Cortex. Front Integr Neurosci. 12:31.

Curtis CE, D'Esposito M. 2003. Persistent activity in the prefrontal cortex during working
memory. Trends in Cognitive Sciences. 7:415-423.

D'Esposito M, Postle BR. 2015. The cognitive neuroscience of working memory. Annu
Rev Psychol. 66:115-142.

Ester EF, Anderson DE, Serences JT, Awh E. 2013. A Neural Measure of Precision in
Visual Working Memory. Journal of Cognitive Neuroscience. 25:754-761.

Fuster JM, Bodner M, Kroger JK. 2000. Cross-modal and cross-temporal association in
neurons of frontal cortex. Nature. 405:347-351.

Goldman-Rakic PS. 1988. Topography of cognition: parallel distributed networks in
primate association cortex. Annu Rev Neurosci. 11:137-156.

Goldman-Rakic PS. 1995. Cellular basis of working memory. Neuron. 14:477-485.

Harris KD, Henze DA, Csicsvari J, Hirase H, Buzsaki G. 2000. Accuracy of tetrode spike
separation as determined by simultaneous intracellular and extracellular measurements. J
Neurophysiol. 84:401-414.

Harrison SA, Tong F. 2009. Decoding reveals the contents of visual working memory in
early visual areas. Nature. 458:632-635.

Hart E, Huk AC. 2020. Recurrent circuit dynamics underlie persistent activity in the
macaque frontoparietal network. Elife. 9.

Kiani R, Cueva CJ, Reppas JB, Peixoto D, Ryu SI, Newsome WT. 2015. Natural
grouping of neural responses reveals spatially segregated clusters in prearcuate cortex.
Neuron. 85:1359-1373.

Leavitt ML, Pieper F, Sachs AJ, Martinez-Trujillo JC. 2017. A Quadrantic Bias in
Prefrontal Representation of Visual-Mnemonic Space. Cereb Cortex.1-17.

Mante V, Sussillo D, Shenoy KV, Newsome WT. 2013. Context-dependent computation
by recurrent dynamics in prefrontal cortex. Nature. 503:78-84.

Markowitz DA, Curtis CE, Pesaran B. 2015. Multiple component networks support
working memory in prefrontal cortex. Proc Natl Acad Sci U S A. 112:11084-11089.

Meyer T, Constantinidis C. 2005. A software solution for the control of visual behavioral
experimentation. J Neurosci Methods. 142:27-34.

Meyer T, Qi XL, Stanford TR, Constantinidis C. 2011. Stimulus selectivity in dorsal and
ventral prefrontal cortex after training in working memory tasks. J Neurosci. 31:6266-
6276.

Meyers EM, Qi XL, Constantinidis C. 2012. Incorporation of new information into
prefrontal cortical activity after learning working memory tasks. Proc Natl Acad Sci U S
A. 109:4651-4656.

Miller JA, Tambini A, Kiyonaga A, D'Esposito M. 2022. Long-term learning transforms
prefrontal cortex representations during working memory. Neuron. 110:3805-3819
e3806.

Qi XL, Constantinidis C. 2012. Correlated discharges in the primate prefrontal cortex
before and after working memory training Eur J Neurosci. 36:3538-3548.

Qi XL, Elworthy AC, Lambert BC, Constantinidis C. 2015. Representation of
remembered stimuli and task information in the monkey dorsolateral prefrontal and
posterior parietal cortex. J Neurophysiol. 113:44-57.

Qi XL, Meyer T, Stanford TR, Constantinidis C. 2011. Changes in Prefrontal Neuronal
Activity after Learning to Perform a Spatial Working Memory Task. Cereb Cortex.
21:2722-2732.

Rao SC, Rainer G, Miller EK. 1997. Integration of what and where in the primate
prefrontal cortex. Science. 276:821-824.

Riley MR, Constantinidis C. 2016. Role of prefrontal persistent activity in working
memory. Front Syst Neurosci. 9:181.

Riley MR, Qi XL, Constantinidis C. 2017. Functional specialization of areas along the
anterior-posterior axis of the primate prefrontal cortex. Cereb Cortex. 27:3683–3697.

Riley MR, Qi XL, Zhou X, Constantinidis C. 2018. Anterior-posterior gradient of
plasticity in primate prefrontal cortex. Nat Commun. 9:3790.

Seung HS, Lee DD, Reis BY, Tank DW. 2000. Stability of the memory of eye position in
a recurrent network of conductance-based model neurons. Neuron. 26:259-271.

Sreenivasan KK, Vytlacil J, D'Esposito M. 2014. Distributed and Dynamic Storage of
Working Memory Stimulus Information in Extrastriate Cortex. Journal of Cognitive
Neuroscience. 26:1141-1153.

Tong F, Pratte MS. 2012. Decoding patterns of human brain activity. Annual review of
psychology. 63:483-509.

Wallis JD, Anderson KC, Miller EK. 2001. Single neurons in prefrontal cortex encode
abstract rules. Nature. 411:953-956.

Wilson FA, O Scalaidhe SP, Goldman-Rakic PS. 1993. Dissociation of object and spatial
processing domains in primate prefrontal cortex. Science. 260:1955-1958.

Wimmer K, Nykamp DQ, Constantinidis C, Compte A. 2014. Bump attractor dynamics
in prefrontal cortex explains behavioral precision in spatial working memory. Nat
Neurosci. 17:431-439.

Xing Y, Ledgeway T, McGraw PV, Schluppeck D. 2013. Decoding working memory of
stimulus contrast in early visual cortex. The Journal of neuroscience : the official journal
of the Society for Neuroscience. 33:10301-10311.

Zhu J, Hammond BM, Constantinidis C. 2023. Laminar pattern of adolescent
development changes in working memory neuronal activity. bioRxiv.
doi.org/10.1101/2023.07.28.550982.

   FIGURES

Figure 1. Task structure and stimuli used. A. Passive, spatial task. B. Active spatial task.
nine possible cue locations in a session shown in the inset. The peripheral locations are
numbered from 1-8, for reference in other figures. The animals were required to maintain
center fixation throughout both active and passive task trials. At the end of active tasks
trials however, monkeys were required to make a saccade to a green target if the stimuli
matched or to a blue target if the stimuli did not match. C-D. Passive and active shape
task. Eight possible shapes in a session are shown in the inset, numbered from 1-8, for
reference in other figures. Stimuli in both tasks extended 2 degree of visual angle.
495

498
499

500
501

502 **Figure 2**. Proportion of neurons exhibiting significant spatial selectivity at different task
503 epochs. A. Selective neurons prior to training. Sample size n=1661. B. Selective neurons
504 post-training (n=1185 neurons). C. Proportion of neurons with the same spatial selectivity
505 at different task intervals.

506
507

508
Figure 3. Normalized Euclidean distance of the representations of spatial stimuli among
pairs of neighboring prefrontal neurons. The box borders represent the interquartile range
(IQR) of the null distribution generated randomly. Whisker length indicates 1.5
interquartile range (IQR). Red crosses are outliers in the null distribution,. Black asterisks
represent empirical mean Euclidean distance values. The representational distance
between nearby neurons is smaller than the expected chance level.
515
516

518
519
520 **Figure 4**. Heat maps and dendrogram representing the representational distance between
521 any two stimuli used in the spatial task prior to training (top) and after training (bottom).
522 Stimulus numbers refer to spatial locations as indicated in Fig. 1. Deeper color in the heat
523 maps indicates less similarity between the neuronal responses elicited by two stimuli.
524 Distance numbers are in arbitrary units.
525

530 **Figure 5**. Proportion of neurons exhibiting significant shape selectivity at different task
531 epochs. A. Selective neurons prior to training. Sample size n=1493. B. Selective neurons
532 post-training (n=1091 neurons). C. Proportion of neurons with the same spatial selectivity
533 at different task intervals.
534

Figure 6. Normalized Euclidean distance of the representations of shape stimuli among pairs of neighboring prefrontal neurons. Conventions are the same as Fig. 3. In the Feature task, the representational distance between nearby neurons is smaller than the expected chance level.

543
544

Figure 7. Heat map and dendrogram representing the representational distance between
any two stimuli used in the feature task, before (top) and after training (bottom).
Conventions are the same as in Fig. 4.

550
551