Chloramphenicol reduces phage resistance evolution by suppressing bacterial cell surface mutants

Authors

Lavisha Parab*[1], Jordan Romeyer Dherbey[1], Norma Rivera[1], Michael Schwarz[1] and Frederic Bertels*[1]

[1]Department Microbial Population Biology, Max Planck Institute for Evolutionary Biology, Plön, Germany.

*Corresponding authors:

Lavisha Parab. Address: Max Planck Institute for Evolutionary Biology, August-Thienemann-Straße 2, 24306 Plön, Germany; email: parab@evolbio.mpg.de; phone: + 49 4522 763-229. ORCID 0000-0002-7626-3216.

Frederic Bertels. Address: Max Planck Institute for Evolutionary Biology, August-Thienemann-Straße 2, 24306 Plön, Germany; email: bertels@evolbio.mpg.de; phone: + 49 4522 763-222. ORCID 0000-0001-6222-4139.

Classification: Biological Sciences, Evolution

Keywords: antibiotic resistance, phage resistance, evolutionary rescue, fluctuation assay

Abstract

Bacteriophages infect Gram-negative bacteria by attaching to molecules present on the bacterial outer membrane, often lipopolysaccharides (LPS). Modification of the LPS can lead to phage resistance. LPS modifications also impact antibiotic susceptibility, allowing for phage-antibiotic synergism. The mechanism for these synergistic interactions is unclear. Here, we show that antibiotics affect the evolution of phage resistance using phage ΦX174 and *Escherichia coli* C wildtype. We use a collection of *E. coli* C LPS mutants, each of which is resistant to ΦX174, and has either a "rough" or "deep-rough" LPS phenotype. Growth of deep rough mutants is inhibited at subinhibitory chloramphenicol concentrations. In contrast, gentamicin has no major effect on growth. Treating *E. coli* C wildtype with ΦX174 and chloramphenicol eliminates deep rough mutants, and reduces phage resistance evolution. Our results show that differential survival of phage resistant mutants with antibiotics explains phage-antibiotic synergism in our model system.

Introduction

The rise of antibiotic resistant bacteria is a global threat to public health and food security (World Health Organization, 2014). To tackle this issue, many alternatives to antibiotics have been proposed (Czaplewski et al., 2016). Bacteriophages (phages) are among the most promising treatments and have been successfully applied to antibiotic resistant infections (Schooley et al., 2017; Chan et al., 2018; Abedon, 2019b; Pirnay et al., 2022). The efficacy of phage treatment can be improved by combining them with antibiotics (Lu and Collins, 2009; Kirby, 2012; Coulter et al., 2014; Kamal and Dennis, 2015; Chaudhry et al., 2017; Berryhill et al., 2021; Gordillo Altamirano et al., 2022; Pirnay et al., 2022). However, as with antibiotics the main reason for treatment failure is the evolution of resistance.

Specific combinations of phages and antibiotics can reduce resistance evolution (Coulter et al., 2014; Torres-Barceló et al., 2014; Chaudhry et al., 2017; Kebriaei et al., 2022). For example, phages NP1 and NP3 displayed synergistic effects with ceftazidime and ciprofloxacin when killing *P. aeruginosa* in biofilms (Chaudhry et al., 2017). Phages combined with antibiotics led to 100 to 1000 times higher killing of bacteria in biofilms than with phages alone. In a different study, combining phage ECA2 with a subinhibitory concentration of ciprofloxacin led to synergistic effects in both, eliminating *Escherichia coli* cells and reducing resistance evolution (Valério et al., 2017). But combinations with ampicillin, kanamycin, piperacillin, tetracycline and chloramphenicol were worse or comparable to phage or antibiotic monotherapy in liquid *in vitro* experiments. These effects strongly depend on antibiotic dosage among other factors. A lethal concentration of

ciprofloxacin prevented phage replication in a combination treatment, and no synergistic
effects were observed (Valério et al., 2017; Tagliaferri et al., 2019).

Some combinations of antibiotics and phages can also lead to increased levels of resistance
to treatment (Cairns et al., 2017). Similar observations to the antagonistic interaction
between ciprofloxacin and phage ECA2 were made in an experiment with phage SBW25Φ2
infecting *Pseudomonas fluorescens* SBW25. SBW25Φ2combined with subinhibitory
concentrations of streptomycin led to both, increased rates of phage resistance and
extinction of the phage population (Cairns et al., 2017). None of these studies have explored
the mechanism behind the evolution of phage resistance and how it is affected by antibiotics.

One common mechanism of phage resistance evolution is surface receptor modification
(Labrie et al., 2010; Egido et al., 2021). Surface receptors can be used by phages to infect
the cell. Common surface receptors include drug efflux pumps that help the cell to expel
antibiotics and allow bacteria to become resistant to antibiotics (Chan et al., 2018;
Burmeister et al., 2020). A deactivation of such a pump prevents phage infection but also
compromises the cell's ability to expel antibiotics and hence makes the cell susceptible to the
antibiotic (Burmeister et al., 2020). Conversely, if a phage uses an antibiotic uptake channel
to infect a cell, for example a porin, then the loss of such a receptor leads to increased
resistance to both the antibiotic and the phage (Kortright et al., 2021).

Apart from efflux pumps and porins, phages also infect the cell by attaching to
lipopolysaccharides (LPS) (Hancock and Reeves, 1976; Mutalik et al., 2020; Romeyer
Dherbey et al., 2023). LPS molecules are found at the outer membrane of all Gram-negative
bacteria and are the most common phage receptor of Gram-negative infecting phages
(Bertozzi Silva et al., 2016; Bio-Conversion Databank Foundation, 2021). In *E. coli*, most
strains produce a "smooth" LPS phenotype, consisting of a lipid A domain, a core
oligosaccharide (consisting of an inner and outer core) and a polysaccharide called the O-
antigen (Frirdich and Whitfield, 2005). Some strains, such as *E. coli* C, lack the O-antigen,
and produce a "rough" phenotype (Yethon et al., 1998). A deletion of both O-antigen and the
outer core oligosaccharide leads to a "deep rough" phenotype (Nikaido and Vaara, 1985).

Deep rough LPS phenotypes can affect antibiotic susceptibility. For example, in *E. coli* deep
rough mutants are more susceptible to membrane-targeting antibiotics such as colistin
(Ebbensgaard et al., 2018; Burmeister et al., 2020). LPS modifications also change
membrane permeability (Nikaido and Vaara, 1985; Nikaido, 2003), thereby affecting
resistance to many antibiotics that do not target the membrane. For example, *E. coli* K-12
strains carrying a deep rough LPS structure are more susceptible to hydrophobic molecules
such as novobiocin and spiramycin (Nikaido and Vaara, 1985; Nikaido, 2003). In contrast,
deep rough mutants caused by mutations in *galU* have led to increased antibiotic resistance

in *P. aeruginosa* (El'Garch et al., 2007; Dötsch et al., 2009). Combining an LPS-targeting
phage with an appropriate antibiotic and the appropriate target pathogen, could potentially
reduce the evolution of resistance.

Phage ΦX174 also uses the LPS structure to infect *E. coli* C (Michel et al., 2010). ΦX174 is
one of the oldest and best-known phage model systems (Sinsheimer, 1959; Sanger et al.,
1977; Smith et al., 2003). Surprisingly though, the resistance mechanism and how to
overcome phage resistance has only been explored recently (Romeyer Dherbey et al.,
2023). We have shown that *E. coli* C easily becomes resistant to phage infection by
modifying its LPS structure, and that ΦX174 can overcome this type of resistance. Yet, how
phage resistance evolution is affected by subinhibitory levels of antibiotics was not explored.

Here, we investigate the effects of antibiotics on ΦX174 resistance evolution in
Escherichia coli C. We first measure antibiotic susceptibility of 33 *E. coli* C strains resistant to
ΦX174 (Romeyer Dherbey et al., 2023), 32 of which have LPS modifications. We chose two
antibiotics: chloramphenicol and gentamicin. Chloramphenicol is a hydrophobic bacteriostatic
antibiotic without significant use in clinical settings and gentamicin is a more hydrophilic
bactericidal antibiotic, both inhibiting protein synthesis. We find that deep rough LPS mutants
do not grow at chloramphenicol concentrations where the wildtype grows densely. However,
there are only few deep rough mutants that cannot grow at subinhibitory concentration of
gentamicin. These results suggest that the phage-chloramphenicol combination treatment
should prevent the emergence of deep rough mutants but not the phage-gentamicin
treatment. We predict that without the emergence of deep rough mutants, chloramphenicol
will reduce the likelihood of phage resistance. Subsequent fluctuation experiments confirm
that deep rough mutants are completely absent from the phage-chloramphenicol treatment.
In contrast, most phage resistant mutants are deep rough in the phage-only treatment. Our
results show that we can predict phage resistance rates surprisingly accurately in
combination with subinhibitory levels of antibiotics.

Results

Deep rough LPS phenotypes are more susceptible to chloramphenicol

We studied how LPS modifications change antibiotic susceptibility using 33 *Escherichia coli*
C strains resistant to phage ΦX174 infection. The strains were generated in a previous study
through a fluctuation experiment (Romeyer Dherbey et al., 2023). Briefly, 33 independent
overnight cultures of *E. coli* C wildtype (WT) were plated together with ΦX174 high-titre
lysates on solid LB plates. After one day, a single colony was isolated at random from each
plate. Thirty-two of the 33 phage resistant strains carry at least one mutation in genes
involved in the LPS biosynthesis, assembly or regulation. Based on which LPS gene was

impacted and current knowledge on LPS biology (Yethon et al., 1998; Whitfield et al., 1999;
Amor et al., 2000; Frirdich and Whitfield, 2005), we predicted the LPS structure of each *E.
coli* C phage resistant strain. While we have previously shown that the exact predictions of
the LPS structure may be inaccurate, the categorisation into "rough" or "deep-rough" may be
useful (**Figure 1, Supplementary Table S1**).

Figure 1. Predicted LPS phenotypes of 33 phage resistant *E. coli* C strains. Strains are
categorised into "deep-rough" (outer core absent or inner core modified) and "rough" (outer
core present but modified and inner core unmodified) based on which gene the mutation is
located in even if the mutation does not delete or deactivate the gene (predicted structures
are likely incorrect). The vertical solid lines separate LPS categories. LPS phenotype for
strain R25 cannot be predicted as the mutated gene, *rfaH* controls the expression of the
entire *waa* LPS operon (Klein and Raina, 2019). While the presence or absence of LPS outer
core is commonly used to categorise LPS types, previous studies with gene deletion mutants
show that absence of phosphate molecules in the inner core also leads to a "deep rough"
phenotype, as in R6 and R11 (Yethon et al., 1998). Kdo: 3-deoxy-D-*manno*-octulosonic acid;
"?P": No information was found on the phosphorylation of Hep(I) in the absence of *waaF*; (P):
only 40% of the hexose phosphorylation was observed (Yethon et al., 2000); R#: the number
given to each *E. coli* C phage resistant strain; OS: oligosaccharide. Figure adapted from
Romeyer Dherbey et al., 2023.

LPS phenotypes may affect antibiotic susceptibility differently for different antibiotics. In our
first experiment we determined antibiotic susceptibility of LPS mutants for one bacteriostatic

antibiotic (chloramphenicol) and one bactericidal antibiotic (gentamicin). Both antibiotics
inhibit protein synthesis. Chloramphenicol binds to the 50S while gentamicin binds to the 30S
ribosomal subunit (Lin et al., 2018). To measure changes in antibiotic susceptibility, we
compared growth of phage resistant strains to growth of *E. coli* C wildtype in the presence of
these two antibiotics on solid agar plates.

With chloramphenicol, *E. coli* C WT and all rough mutants are inhibited at similar levels (4 to
6 µg/mL) with R2 and R7 being slightly more resistant than the wildtype. Whereas deep
rough mutants are inhibited at significantly lower levels (0.5 to 2 µg/mL, **Figure 2,**
Supplementary Table S2, Supplementary Figure S1). Among deep rough mutants, a
subset of predicted heptoseless LPS mutants (R8, R10, R18, R26) – possessing the most
truncated LPS structure – are the strains most susceptible to chloramphenicol. R25 (*rfaH*
mutant) has a susceptibility similar to rough LPS mutants, indicating that R25 indeed displays
rough LPS molecules. The higher chloramphenicol susceptibility of deep rough mutants may
be caused by different levels of membrane permeability. Previous studies show deep rough
LPS phenotypes have increased susceptibility to small hydrophobic molecules such as
chloramphenicol (Nikaido and Vaara, 1985; Chang et al., 2010; Wang et al., 2015; Pagnout
et al., 2019). Thus, deep rough mutants are (i) more susceptible to chloramphenicol than
rough mutants, and (ii) completely inhibited at a chloramphenicol concentration (2 µg/mL)
that is subinhibitory for wildtype *E. coli* C.

Figure 2. Deep rough LPS mutants are more susceptible to chloramphenicol. LPS
modifications do not affect gentamicin susceptibility much. The heatmaps show growth of
E. coli C WT and phage resistant strains (R#) at different concentrations of chloramphenicol
and gentamicin. Growth is measured in 8-bit pixel grayscale values from 0 to 255 (values in
Supplementary Table S2), and decreases from violet (uniform lawn) to yellow (no lawn,
colonies might be visible). Average grey scale values of larger than eight are defined as
growth (left of the bold black border) and values of less than eight are defined as no growth
(see **Methods** for details). Bacterial strains are ordered by their predicted LPS phenotype
and grouped as rough and deep rough LPS. White spaces are shown where we did not
measure bacterial growth. Wildtype minimum inhibitory concentration (MIC) is 4 µg/mL for
chloramphenicol, and 4 µg/mL for gentamicin. Higher bacterial yields in the presence of
subinhibitory concentrations of antibiotics in comparison to no antibiotics when grown on
plates and in liquid have been described previously (Frost et al., 2018; Mahrt et al., 2021;
Farr et al., 2023; Aduru et al., 2024).

Gentamicin susceptibility does not change much across different LPS phenotypes (**Figure**
2). Most phage resistant strains grow roughly similar to the wildtype (inhibited at 4 µg/mL to
6 µg/mL). Four rough LPS strains (R34, R1, R33, R20) and *E. coli* C R25 are inhibited at a
slightly higher gentamicin concentration (6 µg/mL) than the wildtype. Hence, one would
expect LPS mutants to occur in gentamicin resistance screens, which is indeed the case for
moderate levels of gentamicin (El'Garch et al., 2007; Ibacache-Quiroga et al., 2018).
However, some severely truncated heptoseless LPS and only one rough LPS strain (R29)
show increased susceptibility to gentamicin. A group of six *hldE/waaC* (heptose-less)
mutants (R8, R10, R18, R26, R31, R32) with the shortest LPS are inhibited at a lower
concentration (1.5 µg/mL to 2 µg/mL) than the wildtype MIC, but the difference to the
wildtype MIC is smaller than for chloramphenicol (**Figure 2, Supplementary Table S2,**
Supplementary Note S1).

Subinhibitory concentrations of chloramphenicol should reduce ΦX174 resistance
evolution

Our susceptibility assay suggests that subinhibitory levels of chloramphenicol should affect
phage resistance evolution, by reducing the emergence of deep rough mutants. We can test
our hypothesis by performing a fluctuation experiment with wildtype *E. coli* C grown with
phage alone and phage with antibiotics. A fluctuation experiment allows the measurement of
the resistance rate from the distribution of the number of phage and antibiotic resistant
mutants across replicates (Luria and Delbrück, 1943). If there are no interactions between
antibiotics and phages then we should be able to predict the outcome of the experiment
accurately, based only on our susceptibility data.

In our first model we assume that mutation rates do not differ significantly between
treatments. While this assumption is likely inaccurate as mutational bias can significantly
affect evolutionary outcomes (Lind et al., 2019), it is currently impossible to predict
mutational biases from the sequence alone. Furthermore, while the model may not be as
close to reality as a more complex model, the comparison between simple and complex
models provides insight into how much more understanding added complexity provides.
Additional experiments later on in the manuscript will enable us to take mutational biases in
our second model into account.

If mutation rates are constant across the genome, then mutational target size is proportional
to gene length. The larger the mutational target the likelier it is that a random mutation leads
to a resistant phenotype. Almost all LPS mutations we observed are deactivating mutations
that lead to resistance by introducing stop codons through frame shifts or nonsense
mutations. Because we do not know where exactly mutations that deactivate genes are
clustered, we assume a uniform distribution across genes.

The mutational target size can be inferred from the genes that contain mutations in the 33
phage resistant strains. Mutations in a total of 13 LPS genes can lead to phage resistant
mutants, with a combined length of 11,736 bp (**Table 1**). This mutational target size is only
available in the phage only treatment. In treatments with antibiotics the number of genes that
are available for the acquisition of resistance mutations is reduced because certain LPS
mutants are not viable in the presence of low levels of antibiotics (**Figure 2**). We can now
subtract the genes from the total length of the mutational target size to estimate the reduced
phage resistance rate (**Table 1**).

Table 1. Model to predict resistance rates for phage-antibiotic combinations

	Genes (previously observed)	Mutant survival, if mutation in that gene[$]				Gene coding region (bp)	Mutation probability (proportional to gene length)				
		CL 1 µg/mL	CL 2 µg/mL	GM 1 µg/mL	GM 2 µg/mL		Φ (entire gene)	Φ + CL 1 µg/mL	Φ + CL 2 µg/mL	Φ + GM 1 µg/mL	Φ + GM 2 µg/mL
R O U G H	yajC[§]	Y[§]	Y	Y	Y	333	333	333	333	333	333
	waaW	Y	Y	Y	Y	1026	1026	1026	1026	1026	1026
	galE	Y	Y	Y	Y	1017	1017	1017	1017	1017	1017
	waaT	Y	Y	Y	Y	996	996	996	996	996	996
	waaO	Y	Y	Y	M (6/7)	1017	1017	1017	1017	1017	872
D E E P	rfaH	Y	Y	Y	Y	489	489	489	489	489	489
	waaP	Y	M (1/2)	Y	Y	798	798	798	399	798	798
	waaG	Y	N	Y	Y	1125	1125	1125	0	1125	1125
R O U G	galU	Y	M (1/5)	Y	Y	909	909	909	182	909	909
	waaF	Y	N	Y	Y	1047	1047	1047	0	1047	1047
	gmhB	Y	N	Y	Y	573	573	573	0	573	573

H											
	hldE	M (3/6)	N	Y	M (1/6)	1434	1434	717	0	1434	244
	waaC	N	N	Y	N	972	972	0	0	972	0
	Total					11736	11736	10047	5459	11736	9428
	Relative resistance rate[#]						1.00	0.86	0.47	1.00	0.80

[$]Model is based on antibiotic susceptibilities of phage resistant strains. [§]"Y" indicates all strains with a mutation in that gene grow at that antibiotic concentration. "M (x/n)" indicates that x strains grow out of n strains with a mutation in that gene. "N" indicates none of the strains with a mutation in that gene grow at that concentration. The model assumes that mutations occur in genes with a probability proportional to their coding region length. Thus, in genes with a "Y", the mutational target size is the coding region length. For "M (x/n)", the mutational target size is (x/n) * (coding region length). For "N", the mutational target size is zero. [#]The predicted relative resistance rate is the ratio of the mutational target size of the treatment to that of ΦX174 alone. CL: chloramphenicol, GM: gentamicin. [§]*YajC* is the only gene that has no direct link with LPS assembly or synthesis.

The predicted resistance rate for chloramphenicol at 2 µg/mL is less than half (47%) of what is expected when treating the bacteria only with ΦX174 (**Table 1,** red dashes in **Figure 3B**). The predicted reduction in resistance rate is mostly the result of a lack of observed growth of deep rough mutants. Mutations in *waaG*, *waaF*, *gmhB*, *hldE* and *waaC* genes lead to complete inhibition of growth, while mutations in *waaP* and *galU* lead to a complete lack of growth in only some of the mutants (**Table 1**). In contrast, combination treatment with gentamicin reduces the predicted resistance by only 20% at the highest subinhibitory concentration (2 µg/mL). In gentamicin combination treatment only mutations in *waaC* are expected to completely inhibit growth and partial inhibition is expected for mutations in *hldE* and *waaO* (**Table 1**).

Fluctuation experiments qualitatively agree with predictions

In order to test whether our predictions hold, we performed a classic fluctuation experiment (Luria and Delbrück, 1943; Rosche and Foster, 2000). We treated *E. coli* C WT with ΦX174 in combination with either chloramphenicol or gentamicin at two subinhibitory concentrations or inhibitory concentration (**Figure 3A**, **Methods**). We counted colonies from 50 biological replicates for each treatment, and calculated the resistance rate from the distribution of colony numbers (Zheng, 2017).

Figure 3. Chloramphenicol reduces the rate of ΦX174 resistance. (A) Schematic of the

fluctuation experiment and expected results. Adding chloramphenicol at a subinhibitory

concentration is predicted to reduce spontaneous emergence of resistance when treating

E. coli C with ΦX174 due to inhibition of deep rough LPS phenotypes. Adding gentamicin,

however, is predicted to have little to no effect. **(B)** Resistance rates for each phage antibiotic

combination relative to the resistance rate of ΦX174 alone. Observed rates (black) are

qualitatively in line with the predicted rates from model 1 (gene length model, red, derived in

Table 1). Rates predicted by model 2 (gene proportion model inferred from the occurrence of

mutations in a phage-only fluctuation experiment (see below), yellow, derived in

Supplementary Table S6) are within 95% confidence intervals for gentamicin, and 2% away

from the 95% confidence interval in 2 µg/mL chloramphenicol. 1 µg/mL and 2 µg/mL are

subinhibitory concentrations for chloramphenicol (CL) and gentamicin (GM). Resistance

rates are not predicted at antibiotic concentrations inhibitory for wildtype (inhibitory

concentration (IC): 6 µg/mL for CL and 4 µg/mL for GM). **(C)** Absolute resistance rates (per

cell division), including gentamicin at wildtype MIC without ΦX174. Rates of ΦX174 with

antibiotic at inhibitory concentrations seem to be zero in (B) but are actually ~400-1000X

lower, as can be seen in (C). (Values are in **Supplementary Table S3**) Error bars

correspond to 95% confidence intervals. Resistance rate to chloramphenicol treatment

without ΦX174 could not be measured (see **Supplementary Note S2**).

Resistance rate of phage-chloramphenicol combination treatment is significantly lower than

phage alone treatment (Likelihood ratio test (LRT), p-value $<10^{-5}$ at both 1 µg/mL and

2 µg/mL, **Supplementary table S3**). The resistance rate of phage alone treatment is about

10^{-7} per cell division. Once chloramphenicol was added at a concentration of 1 µg/mL, the

resistance rate decreased to 2×10^{-8}, or 15% (95%CI: 12% to 20%) of the phage alone

treatment (baseline, **Figure 3B, Supplementary table S3**). This rate is about four times

lower than what we predicted with our simple gene model (model 1), where we predicted the

resistance rate to only decrease to 86%. At twice the chloramphenicol concentration of

2 µg/mL, the rate is predicted to decrease to 47% of our baseline, which is about twice of our

observation of 18% (95%CI: 14% to 23%).

Treating bacteria with ΦX174 and gentamicin at subinhibitory concentrations does not affect

phage resistance rate much. At 1 µg/mL gentamicin, the emergence of resistance is not

significantly different from the baseline rate (LRT, p-value ~ 0.53), as expected. The

resistance rate was predicted to be the same as the baseline (100%), which is inside the

95% confidence interval of the observed rate of 94% (95% CI: 81% to 106%). Adding

2 µg/mL gentamicin to ΦX174 significantly reduces resistance rate to 60% of the baseline

(LRT p-value ~ 10^{-5}, 95% CI: 50% to 70%), which is slightly lower than the predicted rate of

80% (**Figure 3B, Supplementary table S3**).

We also measured resistance rate for gentamicin treatment at wildtype MIC without ΦX174,

which is ten times higher than any treatment with ΦX174 (**Figure 3C, Supplementary table**

S3). Resistance rates to chloramphenicol treatment without ΦX174 could not be measured

due to the tiny size of most colonies (**Supplementary Note S2**). Any treatment with phage

only is at least ten times better at reducing resistance evolution than antibiotics at MIC. Of

course, at higher antibiotic concentrations we expect lower rates of antibiotic resistance.

We also combined phage treatment with inhibitory concentration of antibiotic and found

significantly lower rates of resistance than any individual treatment. When ΦX174 is

combined with either antibiotic at a concentration inhibitory for the wildtype (IC), resistance

rates decrease 1000 times below phage treatment alone (**Figure 3C, Supplementary table**

S3). Compared to gentamicin alone, resistance rates reduce 10,000-fold, hence, combining

phages and gentamicin synergistically (more than additive) reduces resistance rates.

In our first model, we predicted that the reduction in phage resistance rates is due to a

reduced mutational target size. We next explored whether the mutational target size, that is,

the genomic distribution of mutations, changes across different treatments.

Chloramphenicol skews mutations towards genes causing a rough LPS phenotype

Adding antibiotics to phage treatment reduces the available mutational target size for

resistance, as expected. Across five fluctuation experiments we genome sequenced a total

of 217 phage/antibiotic resistant mutants. Only four of these mutants (1.8%) did not contain

mutations in LPS related genes **(Supplementary Note S4, Supplementary Table S5)**.

Without antibiotic, a majority (86%) of the mutations are found in deep rough genes – genes

responsible for the assembly or modification of the inner core LPS structure (**Figure 4A and**

4B). In the following sections we will refer to these mutants as deep rough mutants. At

2 µg/mL of either antibiotic, this trend shifts towards rough genes – genes responsible for the

assembly and modification of the outer core, which we will refer to as rough mutants. At

2 µg/mL gentamicin, mutations are absent in four deep rough LPS genes (*gmhA*, *hldE*, *hldD*

and *waaC*). At 2 µg/mL chloramphenicol, mutations are not observed in any deep rough

genes or *rfaH* (**Figure 4A and 4B**).

Figure 4: Subinhibitory concentrations of antibiotics reduce the mutational target size. (A) Distribution of mutations across LPS genes. Red boxes show absence/rarity of mutations in a treatment where other treatments have many mutations. Genes that contribute to inner core synthesis are underlined and in red; loss of function mutations in these genes lead to a deep rough phenotype. *Gene *rfaH* is responsible for regulation of the entire LPS operon, so mutations in *rfaH* can lead to a mixture of rough and deep rough types (Pradel and Schnaitman, 1991; Klein and Raina, 2019). Large deletions (>100 bp) are indicated as solid horizontal lines. (Details about types of mutations and parallelism in **Supplementary Figure S2** and **Supplementary Note S3**) (B) Distribution of mutated LPS genes. We whole-genome sequenced mutants from each selective condition and identified the mutated genes. Graph shows proportion of mutants with a mutation in a given gene (n = 49 for phage only, n = 39 for phage with 1 µg/mL CL, n = 36 for phage with 2 µg/mL CL, n = 46 for phage with 1 µg/mL GM, n = 42 for phage with 2 µg/mL GM; CL: chloramphenicol, GM: gentamicin). Details of mutations are in **Supplementary Table S5**.

Combining phage treatment with antibiotics biases mutations towards rough LPS genes (**Figure 5A**). Mutations in strains resistant to phage treatment alone are predominantly found

in deep rough LPS genes (~86%), and only about ~12% are in rough LPS genes. As
expected from our predictions, adding chloramphenicol to phages enriches for rough
mutants. At 1 µg/mL chloramphenicol the proportion of rough mutants increases from 12% to
~62%, and the proportion of deep rough mutants decreases from 86% to ~36%. At a
concentration of 2 µg/mL chloramphenicol, all mutants are rough mutants. This result was
surprising, since we predicted to observe at least some deep rough mutants (~11%) (**Figure
5A**). Gentamicin treatment also leads to a smaller proportion of deep rough mutants, but as
expected from our predictions, the effect is much smaller. At a concentration of 1 µg/mL
gentamicin deep rough mutants remain largely stable at ~85% and about 15% of the mutants
are rough. At 2 µg/mL gentamicin, the ratio becomes more skewed towards a rough LPS with
about 50% deep rough mutants and about 43% rough mutants (**Figure 5A**).

Figure 5. Chloramphenicol skews mutations towards rough LPS genes. (A) Graphs show proportion of mutants that possess a mutation in either rough or deep rough LPS genes. Model 1 predictions (red, applies to all figures) assume that the probability of a mutation in a gene is proportional to the gene length (**Table 1**). In model 2 (yellow), the probability of a mutation occurring in a gene is based on the observed proportions in the phage only treatment (**Supplementary Table S6**). Predictions for antibiotic treatments are based on the susceptibility data. Mutations in *rfaH* or promoter regions of the *waa* operon can affect any number of inner or outer core genes, hence are in a separate category "*rfaH*"" (values in **Supplementary Table S8**). Error bars show the standard deviation for each LPS

type of 100 bootstrap replicates (resampling with replacement, see **Methods**). **(B)** Gene-wise

distribution of mutants resistant to phage treatment without antibiotics. In model 1, *hldD* and

gmhA were not accounted for because they were not observed by Romeyer Dherbey et al.,

2023. The proportions in model 2 differ from the proportions in the phage only treatment

because we add a pseudo-count of 1 to counts of each of the genes to account for the

absence of *gmhB* and *galE* from the phage only treatment (see **Methods**). Genes where

mutations are likely to lead to a rough or deep rough LPS are on the left and right side of the

dotted lines, respectively. "Non LPS" refers to genes that are not known to be involved in

LPS biosynthesis, assembly or regulation (four genes across all experiments,

Supplementary Note S4). Error bars show the standard deviation 100 bootstrap replicates

for each gene (**B** and **C**). **(C)** As per Model 2, in phage treatment with 2 µg/mL

chloramphenicol, mutants of *rfaH, galU* and *waaP* are predicted to survive but none are

observed. (See **Supplementary Figure S3** for model 3).

The discrepancies between predicted and observed mutation distributions across LPS genes

could be due to an ill fitted model. Hence, we examined whether discrepancies between

prediction and observation could be reduced by adjusting our model. If these adjustments do

not improve the predictions then it is possible that second order effects such as phage-

antibiotic interactions (Comeau et al., 2007) contribute to the discrepancies.

Before attempting to predict outcomes of combination treatments, our model should be able

to predict the phage alone treatment accurately. In the phage only treatment deep rough

mutants are significantly more common (86%) than we would predict from the mutational

target size of deep rough genes (58%) (**Figure 5A, Supplementary Table S8**). Perhaps, the

distribution of mutations in genes is not proportional to gene lengths, as we had previously

assumed, for example, due to mutational bias (Lind et al., 2019). Instead of using gene

length as proxy for mutational target size we simply used the proportion of mutants that we

observed in the phage only treatment as baseline. We also added a pseudo-count of 1 to

each gene, since two LPS genes which are potential targets for phage resistance were not

observed in the phage only treatment (*gmhB* and *galE*) but are present in combination

treatments (**Supplementary Table S6**). The adjusted proportions are shown in yellow in

Figure 5B (model 2). Adjusting the baseline predictions should in theory improve the

predictions for the combination treatments.

Adjusted model improves predictions and highlights potential phage-antibiotic interactions

Using model 2 improves predicted resistance rates (**Figure 3B**). For combinations with

2 µg/mL antibiotic, the predicted rates (25% for chloramphenicol, 61% for gentamicin) are

now almost within the 95% confidence interval for chloramphenicol and almost perfect for

gentamicin (18% and 60% respectively). However, the predicted distribution of rough and
deep rough mutants is worse than for model 1 (**Figure 5A**). The only treatment where the
LPS mutant prediction improved is at 1 µg/mL gentamicin, presumably because this
treatment is closest to the phage only treatment based on our initial susceptibility assay
(**Figure 5A**).

Model 2 qualitatively predicts the shift towards rough mutants, but not accurately (**Figure
5A**). Since some *waaP* and *galU* mutants grow at 2 µg/mL chloramphenicol in our
susceptibility assays, deep rough *waaP* and *galU* mutants are predicted to occur at a rate of
6% and 11% leading to a non-zero deep rough prediction (**Figure 5A and C,
Supplementary Table S6**). *RfaH* mutants are predicted to occur at a rate of 13%, because
they also grow well in our susceptibility assay. Yet, *waaP, galU* and *rfaH* mutants are never
observed at 2 µg/mL chloramphenicol (**Figure 5C**). Similarly, at 1 µg/mL chloramphenicol,
mutations are predicted in *rfaH*, *waaF*, *gmhB*, and *gmhA* but none are observed
(**Supplementary Figure S3C**). At 2 µg/mL gentamicin mutations are predicted in *hldE*, *gmhA
and gmhB*, all deep rough genes, but none are observed (**Supplementary Figure S3D**).
There are at least three possible reasons for these discrepancies between observation and
prediction. First, sampling only 40-50 mutants per treatment could under-sample the
proportion of mutated LPS genes. Sampling more mutants in the phage only treatment may
allow us to estimate the proportion of mutated LPS genes more accurately and prevent us
from using pseudo-counts in our models, leading to more accurate predictions. Second,
biological interactions between antibiotics and phages (Comeau et al., 2007; Gu Liu et al.,
2020) may lead to the overrepresentation of different LPS genes compared to the phage only
treatment. Third, it is possible that what we consider growth in our susceptibility assay is not
sufficient to be detected in the fluctuation experiment.

We are planning to test hypotheses one and two in future studies, hypothesis three can be
tested by adjusting the threshold for growth in our susceptibility assay. When we increase the
threshold for growth from 8 to 12 average greyscale value, phage resistance rate predictions
remain largely unchanged from model 2 to the new model 3, except for the 2 µg/ml
gentamicin treatment where the prediction becomes worse (**Supplementary Figure S3A**).
Predicted distributions across LPS genes do not change significantly for most treatments
(**Supplementary Figure S3B, Supplementary Table S7-S8**), but model 3 does indeed
improve the prediction for *waaG* at 2 µg/mL of chloramphenicol (**Supplementary Figure
S3E**). Interestingly, with the higher threshold *gmhB*, *gmhA* and *hldE* are not predicted to
grow anymore at 2 µg/mL gentamicin (**Supplementary Figure S3D**), which is also what we
observe in our experiment. However, none of the model adjustments can explain the
discrepancy between the predicted and observed phage resistance rates at 1 µg/mL
chloramphenicol or the discrepancy in the proportions of deep rough and rough mutants at

that concentration (**Supplementary Figure S3A-B**). This effect is likely the result of direct or
indirect interactions between phages and chloramphenicol.

Interactions between phages and chloramphenicol are also indicated by statistical
comparisons between observations and expectations according to different models (**Table**
2). If there is no statistical difference between model and observation, we assume that the
model performs well. As the difference between model and observation becomes more
significant, the model becomes worse. Model 1 is better at predicting gene-wise distributions
than the other two models, except for the phage with 1 µg/mL gentamicin treatment,
indicating that gene length is a good approximation for the likelihood of observing a mutation
in a certain gene (top of **Table 2**). Models 2 and 3 are better at predicting rough and deep
rough proportions (**Table 2** bottom) and phage resistance rates (**Figure 3B, Supplementary**
Figure S3). Hence, adjusting the mutation proportions to the phage only treatment leads to
greater errors for gene predictions but fewer errors for resistance rate and LPS type
predictions.

Table 2. Chi-squared tests to check goodness of fit between observed and predicted
mutant proportions
452

Gene-wise distribution (expected and observed proportions of mutations in each gene)			
Treatment	Model 1 (added genes)^	Model 2	Model 3
Phage	6.20E-05**	0.9981	0.9981
Phage + 1 ug/mL CL	1.69E-05	7.15E-07	7.15E-07
Phage + 2 ug/mL CL	0.0553	3.16E-08	3.51E-06
Phage + 1 ug/mL GM	0.0091	0.01022	0.0102
Phage + 2 ug/mL GM	0.0051	7.60E-05	0.0005

LPS type (expected and observed proportions of deep rough and rough mutants, Figure 5A)				
Treatment	Model 1 (added genes)	Model 2	Model 3	Model 1
Phage	0.0046	0.576	0.576	0.0005
Phage + 1 ug/mL CL	0.0368	6.35E-06	6.35E-06	0.0779
Phage + 2 ug/mL CL	0.0124	0.000388	0.01052	0.0124
Phage + 1 ug/mL GM	0.0084	0.429	0.429	0.0012
Phage + 2 ug/mL GM	0.7795	0.059	0.555	0.838

453
454   **Values shown are p-values obtained by comparing the observed distribution of mutants (gene-wise
455   or LPS type) with the distribution predicted by the given model. ^Before comparing model 1, two
456   missing genes *hldD* and *gmhA* were added to resistance genes based on the underlying gene length
457   assumption, to avoid zero expectation for any category.
458

Discussion

Here we find that phage-antibiotic combinations can reduce the evolution of phage
resistance by preventing the growth of certain mutants. We start by measuring the effect of
chloramphenicol and gentamicin on previously isolated phage resistant mutants. From these
measurements, we predict that certain classes of mutations will not emerge in a combination

treatment, and hence will reduce the rate of phage resistance evolution. We test these
predictions by performing an experiment where we expose *E. coli* C wildtype to antibiotics
and phage ΦX174. Phage resistance rates generally agree with predictions. Whole-genome
sequencing of phage resistant mutants shows that our predictions based on antibiotic
susceptibility assays also largely agree with the observed mutation distributions. Hence, we
show that antibiotics can reduce phage resistance evolution by preventing the growth of
deep rough resistance types. This effect is often referred to as an evolutionary trap (Baym et
al., 2016; Barbosa et al., 2017). Interestingly, first order effects – effects which do not take
interactions between phage and antibiotic into account – are largely sufficient to explain
reduced phage resistance evolution in combination treatments. Our predictions are based on
several assumptions, one of which may be violated.

The original model assumption for the fluctuation assay posits that the outcome of the assay
only depends on two parameters: the mutation rate and the mutational target size (Luria and
Delbrück, 1943). The mutation rate should be identical across all treatments, since mutations
occur during exponential growth in liquid culture in the absence of antibiotic and phages. The
second parameter, the mutational target size, is very close to our expectations for 1 µg/mL
chloramphenicol, and similar to the target size observed for 2 µg/mL gentamicin (**Figure 4A**).
Yet, for gentamicin the predicted phage resistance rate (61%) fits well with the observed
resistance rate (60%), while for chloramphenicol the rates differ significantly (64% predicted
vs. 15% observed, **Figure 3B**). Hence, our assumption on the target size is correct. Because
mutation rates are also identical between treatments there must be other factors that explain
our observations.

One possible explanation is that mutations do not guarantee resistance. It is possible that for
1 µg/mL chloramphenicol the mutations only provide a certain survival chance rather than
guaranteed survival. Previous studies have shown that subinhibitory concentrations of
chloramphenicol do not lead to stochastic population extinctions, but the combination with
bactericidal antibiotics does (Coates et al., 2018). Because phages can be considered
bactericidal agents stochastic population extinctions could play a role in our experiment. If
the stochastic extinction chance is about 50% then the expected resistance rate would
correspond to our observed resistance rate. This survival chance may be caused by local
fluctuation of chloramphenicol concentrations, the cell state or physiological interactions
between phage and chloramphenicol on the plate (Akiyama and Kim, 2021).

The combination between a bactericidal agent (phage) and a bacteriostatic antibiotic may
lead to a trade-off explaining why phage resistance rates at 1 µg/mL and 2 µg/mL
chloramphenicol are similar. At high subinhibitory chloramphenicol concentrations the
antibiotic can efficiently prevent the emergence of deep rough mutants but the efficiency of

killing by the phage may be compromised because the bacteria grow significantly lower at
subinhibitory chloramphenicol concentrations and phage infection is inhibited by slower
growing bacteria (Hadas et al., 1997; Coates et al., 2018). At slightly lower chloramphenicol
concentrations chloramphenicol may inhibit growth less, which will make phage infection
more efficient. These effects may balance each other out leading to similar phage resistance
rates at 1 µg/mL and 2 µg/mL chloramphenicol concentration.

Mutants resistant to phage only treatment are biased towards a deep rough LPS

In treatment with phage only, mutants with a deep rough LPS emerge at a higher proportion
than expected based on the assumption that mutations occur proportional to gene lengths.
We observed that 86% of all mutants were deep rough (compared to 58% expected based
on gene length, **Figure 5B**). It is unclear why deep rough mutants are overrepresented, but
one reason could be that mutation rates are not completely uniform throughout the genome.
Deep rough genes could have more loci with high mutation rates (mutational bias) (Moxon et
al., 1994; Lind et al., 2019; Bertels et al., 2021). Similarly, deep rough candidate genes might
have more mutational targets, that is, loci which produce a phage resistant LPS phenotype,
than in rough candidate genes. Alternatively, mutations in deep rough genes may be more
likely to produce robust resistance in the presence of phages alone. Robust resistance
means that the mutation almost guarantees the establishment of a colony, while for rough
mutations it is less likely that a resistant colony is successfully established.

Deep rough mutants are more abundant in phage resistant mutants, yet at least in *E. coli* are
more susceptible to certain antibiotics (Nikaido and Vaara, 1985; Wang et al., 2015). This
characteristic can be exploited by complementing phage therapy with antibiotics that
specifically affect bacterial strains with a deep rough LPS structure.

LPS modifications affect susceptibility to chloramphenicol and, to a lesser degree,
gentamicin

Chloramphenicol and gentamicin are similar in mechanism of action (Lin et al., 2018) and
resistance (Burns et al., 1989; Recht and Puglisi, 2001; Kehrenberg et al., 2005; El'Garch et
al., 2007), thus the reason chloramphenicol affects deep rough mutants differently may hinge
on its physicochemical properties. The antibiotics differ in their type of inhibition,
hydrophobicity and molecular size: chloramphenicol is a bacteriostatic, hydrophobic molecule
of 323 Da, gentamicin is a bactericidal, hydrophilic molecule of 477 Da. The key differences
are probably hydrophobicity and size, because small hydrophobic molecules can diffuse
through the LPS membrane, while hydrophilic molecules generally require active transport
for cell entry (Nikaido, 2003).

The increased susceptibility of deep rough LPS mutants to hydrophobic antibiotics is in line

with previous studies. Deleting deep rough LPS genes (*waaC*, *waaF*, *waaG* or *waaP*)

increases membrane hydrophobicity and permeability (Wang et al., 2015), can destabilise

the membrane, and increase the entry of small hydrophobic molecules (Nikaido and Vaara,

1985; Chang et al., 2010; Pagnout et al., 2019). Testing more antibiotics could reveal

whether it is molecular size, hydrophobicity or whether the antibiotic is

bacteriostatic/bactericidal that defines the change in susceptibility.

Combining phages with antibiotics to reduce the evolution of treatment resistance

Phage-antibiotic treatment success depends on the types of phage and antibiotic applied and

which bacterial species is targeted (Valério et al., 2017; Lopes et al., 2018; Abedon, 2019a;

Chang et al., 2019; Henriksen et al., 2019). For example, synergistic effects were observed

for chloramphenicol, gentamicin and tetracycline for some phage-bacteria combinations but

not others (Comeau et al., 2007; Kirby, 2012; Kamal and Dennis, 2015). Gu Liu and

colleagues tested many different antibiotics with two phages to test whether combinations

were synergistic, antagonistic, additive or dominated by one agent (Gu Liu et al., 2020).

Synergistic combinations were only observed with one of the two phages tested. Antibiotic

choice is key for treatment outcome, not only should they be more efficacious together (not

lead to phage extinction) but also reduce resistance evolution. Our results show varying

levels of synergetic effects for all antibiotic phage combinations in reducing phage resistance

evolution. It is possible that combining LPS targeting phages with antibiotics is more likely to

lead to synergetic effects than phages targeting specific cell membrane proteins like efflux

pumps or porins.

Implications for phage therapy research

ΦX174 is very specific in its receptor recognition: it only infects about 0.8% out of 783 *E. coli*

human and animal commensal strains (Michel et al., 2010), which currently makes ΦX174

unsuitable for therapeutic purposes. However, the host range of ΦX174 can be expanded

through evolution experiments (Romeyer Dherbey et al., 2023; Romeyer Dherbey and

Bertels, 2024) or genetic engineering (Yehl et al., 2019). Once the limitation of phage entry is

overcome, ΦX174 can replicate its DNA and lyse hosts successfully in different *E. coli* strains

and other species such as *Salmonella typhimurium* and *Enterobacter aerogenes* (Suzuki et

al., 1974; Mayr et al., 2005; Orta et al., 2023). If the host range of ΦX174 can be extended

then the knowledge of ΦX174 biology accumulated over the last decades could be utilized to

turn ΦX174 into an efficient therapeutic agent (Romeyer Dherbey and Bertels, 2024).

More generally our results show that phages in combination with certain antibiotics at

subinhibitory levels can reduce the evolution of phage resistance. This result indicates that

the continued use of antibiotics may be beneficial even when the target bacteria are antibiotic
resistant. Of course, further studies are required to test whether our results also hold for
higher, more clinically relevant antibiotic concentrations and resistance rates.

LPS modifications may also reduce pathogen virulence when treating with LPS-targeting
phages. LPS can increase virulence (Burns and Hull, 1998; Raetz and Whitfield, 2002; Pier,
2007) and elicit immune response (Alexander and Rietschel, 2001). If the evolution of phage
resistance leads to the truncation of LPS structures then by becoming resistant to phage
infection the pathogen may also become less virulent. Future experiments with clinically
relevant antibiotics and phages, should be conducted to allow us to understand, which LPS
targeting phage-antibiotic combinations are most beneficial in therapeutic settings.

Methods

Bacterial and phage strains

Ancestral *E. coli* C and coliphage ΦX174 (GenBank accession number AF176034) were
provided by H.A. Wichman (Program in Bioinformatics and Computational Biology, University
of Idaho, Moscow, Idaho 83844). The ancestral *Escherichia coli* strain C (wildtype) used in
this study differs from the NCBI *E. coli* C strain (GenBank accession number CP020543.1)
by 9 insertions and two nucleotide replacements, (details in Romeyer Dherbey et al., 2023).
All resistant strains used or generated in this study were derived from the same ancestral *E.*
coli C strain. *E. coli* C R1 to R35, used for antibiotic susceptibility measurements were
generated in fluctuation experiments in a previous study (Romeyer Dherbey et al., 2023),
while the new resistant strains have been generated in this study. The genomic sequence of
CP020543.1 was manually modified using Geneious Prime (version 2020.1.2) in a previous
study (Romeyer Dherbey et al., 2023). See **Data availability** for whole-genome sequences
and mutation details.

Media and growth conditions

Bacteria and phages were grown in 2.5% LB broth (Luria-Bertani Broth, Miller, Sigma-
Aldrich) supplemented with $CaCl_2$ and $MgCl_2$ at a final concentration of 5 mM and 10 mM
respectively, henceforth referred to as supplemented LB. Solid LB agar (2.5% Sigma LB
broth, 1.5% agar) was used to plate phages and bacteria. For bacterial overnight cultures, 5
mL supplemented LB in a 13 mL culture tube (Sarstedt) was inoculated with a single isolated
colony from a quadrant-streaked plate and incubated for 17 hours at 37°C with shaking at
250 rpm (New Brunswick™ Innova® 44, Eppendorf). For antibiotic agar plates, liquid LB agar
(2.5% Sigma LB broth, 1.5% agar) was supplemented with $CaCl_2$ and $MgCl_2$ at a final
concentration of 5 mM and 10 mM respectively, and chloramphenicol or gentamicin at

desired concentration (0.5 µg/mL, 0.75 µg/mL, 1 µg/mL, 1.5 µg/mL, 2 µg/mL, 3 µg/mL, 4

µg/mL or 6 µg/mL) before pouring plates inside a laminar flow hood. Plates were dried for 30

min before behind stored at 4°C for a maximum of two weeks. Chloramphenicol and

gentamicin stocks were made at 20 mg/mL in ethanol and Milli-Q water respectively and

stored at -20°C. ΦX174 stocks were prepared fresh every week and stored at 4°C

Determination of antibiotic susceptibility of bacterial mutants

Preliminary measurements of antibiotic susceptibility of *E. coli* C R1 to R35 were done with

E-test strips (bioMérieux, France, data not shown) on LB agar plates to narrow down the

range of concentrations to be tested for MIC determination. Overnight cultures of *E. coli* C

wildtype and phage resistant strains were diluted 10 times in Ringer's solution (Sigma). 50 µL

of each 10 times diluted overnight culture (5×10^6 cells) was spread, using 15 glass beads (4

mm), on a 90mm LB agar plate (supplemented with CaCl2 and MgCl2 at a final

concentration of 5mM and10mM respectively) containing chloramphenicol or gentamicin at

the concentration to be tested (0.5 µg/mL, 0.75 µg/mL, 1 µg/mL, 1.5 µg/mL, 2 µg/mL, 3

µg/mL, 4 µg/mL or 6 µg/mL). LB agar plates (without antibiotic or supplement) were used as

a control. Plates were incubated inverted at 37°C overnight (17 hours). The next day, growth

was assessed by imaging plates after they cooled to room temperature. If bacteria grew at all

the concentrations tested, new rounds of experiments were performed to determine the MIC.

Increasing antibiotic concentrations were tested until there was no detectable growth in three

replicates (CLSI, 2019).

Agar plates were imaged using Chemidoc imaging system (Bio-Rad) at the colorimetric

setting with an exposure of 1 second. Plate images were analysed using OMERO and

OMERO-imageJ plugin to quantify growth as lawn formation (Allan et al., 2012). For uniform

measurements, rectangular regions of interest with the same area (1157.85 mm²) were

marked on the agar plate images (without zooming or cropping). Growth values were

obtained by subtracting average grayscale value of a blank plate from the average grayscale

value of the sample plate. Growth was measured in 8-bit pixel grayscale values from 0 to

255. Uniform bacterial lawns corresponded to grayscale values of 30 to 80. No visible growth

or only a slight haze corresponded with grayscale values of 0 to 8. Average grayscale values

of larger than 8 are defined as growth and values of less than 8 are defined as no growth.

Median of growth values across 3-5 replicates was used to make **Figure 2**. The sensitivity of

the assay to the growth cut-off was tested by changing the growth cut-off to 12 in Model 3.

Fluctuation experiment and mutant isolation.

Mutation rates were measured using a revised Luria Delbrück fluctuation experiment (Luria

and Delbrück, 1943; Rosche and Foster, 2000) in nine treatments. Treatments were: (a)

ΦX174 with chloramphenicol at 1 μg/mL, 2 μg/mL or 6 μg/mL; (b) ΦX174 with gentamicin at

1 μg/mL, 2 μg/mL or 4 μg/mL; (c) ΦX174 only; (d) chloramphenicol only, at 6 μg/mL; (e)

gentamicin only, at 4 μg/mL. All treatment plates were made with LB agar supplemented with

5 mM CaCl2 and 10 mM MgCl2, and when needed, with chloramphenicol or gentamicin at 1

μg/mL, 2 μg/mL, 4 μg/mL or 6 μg/mL.

For each treatment, 50 complete independent single colonies of *E. coli* C WT, each started

from a single cell were inoculated into 100 μL of supplemented LB without antibiotic or

phage, in wells of a 96-well plate. Plates were incubated for 5 hours at 37°C (Eppendorf

Innova plate shaker) shaking at 750 rpm. Cultures were resuspended by pipetting at the 3-

hour and 4-hour timepoints to avoid settling from bacterial aggregation. After incubation, 50

μL of bacterial culture (cell density as below) was spread on the corresponding LB agar

plates using sterile glass beads. For treatments involving ΦX174, bacterial cultures were first

mixed with 50 μL of a high-titer lysate (3 x 10^8 PFU/mL) and immediately plated with sterile

beads on the corresponding LB agar plates. Plates were incubated for 17 hours at 37°C. The

plates with chloramphenicol at 6 μg/ml (with and without phage) were incubated further for

16 hours to account for small pinpricks which might turn into distinct colonies.

We altered the total number of cells spread on a selection plate according to the expected

resistance rate, spreading ten times more cells if the expected resistance rate was ten times

lower. This allowed us to measure rates ranging from 10^{-6} per cell division to 10^{-10} per cell

division. For treatments (a) and (b), 50 μL of undiluted bacterial culture (3 x 10^8 CFU/mL)

was mixed with 50μL of ΦX174 lysate; for (c), 50 μL of twice diluted bacterial culture was

mixed with 50 μL of ΦX174 lysate; for (d) and (e), 50 μL of 10 times diluted bacterial culture

was taken; the entire mixture was then plated on agar plates with the corresponding selective

condition using 15 glass beads. For obtaining total CFU counts, cultures were serially diluted

and plated on non-selective LB agar plates.

The next day, colonies were counted and one colony was randomly picked from each of the

50 plates for all treatments. Spontaneous resistant colonies were purified by re-streaking on

non-selective LB agar plates. Again, one colony was randomly picked from each plate and

was used to inoculate a fresh overnight culture in supplemented LB. One mL of each culture

was mixed with glycerol saline solution and frozen at -80°C.

Mutation rate calculation and statistical analysis

Resistant mutant frequencies, i.e., the number of resistant colonies across 50 independent

replicates were used to calculate mutation rate. Mutation rates were calculated by the Lea-

Coulson model following the method of Foster (Foster, 2006). We used this method because

our bacterial population all started from a single cell. The model assumes that growth rates of

mutants and nonmutants are the same in the absence of selection, and allows for partial
plating of cultures (Foster, 2006). Mutation rate and its 95% confidence interval were
estimated by Lea-Coulson ($\varepsilon<1.0$, i.e., partial plating) method of *webSalvador*, the web-
application of R package *rSalvador* (Zheng, 2017). Mutation rates were compared between
different selective conditions by Lea-Coulson ($\varepsilon<1.0$) likelihood ratio test from the package
rSalvador on R (Rstudio version 2022.02.1) (RStudio Team, 2019; R Core Team, 2020).

DNA extraction and whole-genome sequencing

Samples were prepared for whole genome resequencing from 500 µl of stationary-phase
culture. Genomic DNA of *E. coli* C mutants was extracted using DNeasy Blood & Tissue kit
(Qiagen). Quality of extracted DNA was confirmed before sequencing. Samples were
prepared as previously described (Picelli et al., 2014) and sequenced at the Max-Planck
Institute for Evolutionary Biology (Plön, Germany) with the NextSeq 500, using the NextSeq
500/550 High Output Kit v2.5 (300 Cycles) kit (cat. no. 20024908) (Illumina Inc.). The quality
of the sequencing outputs was controlled using *FastQC* version 0.11.9 (Simon, 2010), read-
trimmed using *Trimmomatic* (Bolger et al., 2014), mapped to the reference and analysed by
breseq (Deatherage and Barrick, 2014). All mutations other than SNPs were confirmed by
mapping reads to their corresponding modified genomes reconstructed using Geneious
Prime 2023.1.1 (https://www.geneious.com).

Predicting mutation rates

Model 1. The probability of a ΦX174 resistance mutation to occur in a particular gene is
assumed to be proportional to the length of the coding region of a resistance conferring gene
(**Table 1**). We assumed that only genes which were previously observed in Romeyer
Dherbey et al., 2023 can confer resistance to treatment with ΦX174. In case a strain
possessed more than one mutation, only the mutation affecting LPS structure was
considered, e.g. *waaO* for R29, *galE* for R1.

The ability of phage resistant strains to grow with antibiotics was used to predict resistance
rates and the genes that confer resistance in phage-antibiotic combination treatments(**Figure
2** and values in **Supplementary Table S2**). If out of "n" strains with a mutation in a gene,
only "x" strains grew with an antibiotic, (x/n) was assumed to be the probability that this gene
confers resistance in a combination treatment. Hence, resistance caused by a mutation in
this gene is proportional to (x/n)* gene length, which we call probability weight. For example,
out of five *galU* mutants, only one can grow at 2 µg/mL chloramphenicol. Thus, resistance
caused by a mutation in *galU* has a probability weight of (1/5*gene length) at 2 µg/mL
chloramphenicol.

Resistance rate is proportional to the sum of all probability weights of genes conferring
resistance. The predicted relative resistance rate for combination treatment is inferred as a
ratio of the total probability weights at each selective condition to that of ΦX174 alone (**Table**
1).

Model 2. The probability weights assigned to each gene were changed to the observed
number of mutations in the phage only treatment (**Supplementary Table S6**). Mutations in
two genes (*gmhB* and *galE*) are not observed in the phage only treatment, meaning a zero
frequency, and zero probability weight in model 2. A weight of zero incorrectly classifies
these genes as not conferring resistance to phage (Romeyer Dherbey et al., 2023) or
combination treatments (**Figure 2**) and will also predict zero probabilities for combination
treatments. We solved the issue by adding a pseudo-count. A count of one was added to all
genes including *galE* and *gmhB* (green highlight in **Supplementary Table S6**).

The inverse problem exists for *hldD* and *gmhA.* These genes have not been observed by
Romeyer Dherbey et al. and hence their antibiotic susceptibility has not been measured.
HldD is a bifunctional gene and mutations can cause two different LPS phenotypes, *waaC* or
waaF like (Coleman, 1983; Schnaitman and Klena, 1993). Resistance rates and LPS
distributions were calculated for both phenotypes. For final analyses, the model closest to
observations was used. *GmhA* mutants were assumed to grow like *gmhB* mutants, that is,
they can grow in all treatments except phage with 2 µg/mL chloramphenicol.

Model 3. The growth cut-off was increased from 8 to 12 (see **Determination of antibiotic**
susceptibility of bacterial mutants). Candidate genes and probability weights of the
baseline were the same as in model 2 (**Supplementary Table S7**).

Statistical Testing

T-test and Wilcoxon rank sum test were used to determine if antibiotic susceptibility for rough
and deep rough strains are significantly different. T-test was performed using the stats
package of R (R version 4.2.0, Rstudio version 2023.12.1) (RStudio Team, 2019; R Core
Team, 2020). A T-test can often be incorrect for discrete numerical variables, which is why a
Wilcoxon rank sum test was also performed to confirm the significance. Wilcoxon rank sum
test with an exact solution for tied datasets was performed using the EDISON WMW
calculator (Marx et al., 2016). Chi-squared tests to check goodness of fit of models were
performed using R (R version 4.2.0, Rstudio version 2023.12.1) (RStudio Team, 2019; R
Core Team, 2020). Observed distributions of mutations by gene and by LPS type were
compared with the distributions predicted by each model. Before comparing model 1, two
missing genes *hldD* and *gmhA* were added to resistance genes based on the underlying
gene length assumption, to avoid zero expectation for any category.

Bootstrap analysis

Mutants were resampled from the original observations with replacement 100 times for each treatment to calculate the expected variation in the sampled observation. To resample genes that have not been observed but are expected to provide resistance, a pseudo-count of one was added to each gene that could provide resistance to treatment. The standard deviation of the 100 samples was used to show the error bars in both gene-wise and LPS type distributions in **Figure 5** and **Supplementary Figure S3**.

Data visualisation

The heatmap (**Figure 2**) and histograms (**Figures 3B-C, 4B, 5, S2, S3**) were generated using Excel (version 16.75.2). All schematic drawings (**Figures 1, 2 and 3A**) were created with BioRender.com. Genomic distribution of mutation loci across treatments (**Figure 4A**) and boxplot of antibiotic susceptibilities (**Figure S1**) were visualised using R (Rstudio version 2022.02.1) (RStudio Team, 2019; R Core Team, 2020).

Data availability

Antibiotic susceptibility. Agar plate images and raw grayscale values used to generate **Figure 2** and **Supplementary Table S2** have been deposited in Edmond (doi: 10.17617/3.PIDVUT).

Whole-genome sequences. Complete genomes of *E. coli* C WT used as reference and ΦX174 WT, and raw sequencing reads of resistant strains isolated from the fluctuation experiments in this study (**Supplementary Table S5**) have been deposited in Edmond (doi: 10.17617/3.PIDVUT). Raw sequencing reads of phage resistant strains *E. coli* C R1 to R35 used for antibiotic susceptibility testing (**Supplementary Table S1**) have been deposited in Zenodo (doi: 10.5281/zenodo.6952399).

Fluctuation experiments. Statistical test results and raw colony counts are in **Supplementary Table S3 and S4.**

Acknowledgements

This work was generously supported by funds from the Max Planck Society (L.P.-F.B.). L.P. was supported by the International Max-Planck Research School for Evolutionary Biology (IMPRS EvolBio). The authors also thank the J. Barrick Lab for their fluctuation test protocols, Prof. Qi Zheng for his help with rSalvador, Dr. Carsten Fortmann-Grote for his help with image analysis and all members of the Microbial Population Biology department for their help with experiments.

Author Information

Affiliations: Max Planck Institute for Evolutionary Biology, Plön, Germany.

Contributions: L.P. and F.B conceived the study. L.P. and F.B. planned the experiments. L.P., N.R., J.R.D. and M.S. carried out the experiments. L.P. conducted the sequence analyses. L.P. and F.B interpreted the results. L.P. and F.B. wrote the original draft of manuscript, L.P., J.R.D., and F.B. reviewed and edited the manuscript. L.P. prepared all figures and tables. All authors reviewed and approved the manuscript.

Ethics declaration

The authors declare no competing interests.

References

Abedon, S. T. (2019a). Phage-Antibiotic Combination Treatments: Antagonistic Impacts of Antibiotics on the Pharmacodynamics of Phage Therapy? *Antibiotics* 8, 182. doi: 10.3390/antibiotics8040182

Abedon, S. T. (2019b). Use of phage therapy to treat long-standing, persistent, or chronic bacterial infections. *Adv. Drug Deliv. Rev.* 145, 18–39. doi: 10.1016/j.addr.2018.06.018

Aduru, S. V., Szenkiel, K., Rahman, A., Ahmad, M., Fabozzi, M., Smith, R. P., et al. (2024). Sub-inhibitory antibiotic treatment selects for enhanced metabolic efficiency. *Microbiol. Spectr.* 0, e03241-23. doi: 10.1128/spectrum.03241-23

Akiyama, T., and Kim, M. (2021). Stochastic response of bacterial cells to antibiotics: its mechanisms and implications for population and evolutionary dynamics. *Curr. Opin. Microbiol.* 63, 104–108.

Alexander, C., and Rietschel, E. T. (2001). Bacterial lipopolysaccharides and innate immunity. *J. Endotoxin Res.* 7, 167–202.

Allan, C., Burel, J.-M., Moore, J., Blackburn, C., Linkert, M., Loynton, S., et al. (2012). OMERO: flexible, model-driven data management for experimental biology. *Nat. Methods* 9, 245–253. doi: 10.1038/nmeth.1896

Amor, K., Heinrichs, D. E., Frirdich, E., Ziebell, K., Johnson, R. P., and Whitfield, C. (2000). Distribution of Core Oligosaccharide Types in Lipopolysaccharides from Escherichia coli. *Infect. Immun.* 68, 1116–1124.

Barbosa, C., Trebosc, V., Kemmer, C., Rosenstiel, P., Beardmore, R., Schulenburg, H., et al. (2017). Alternative Evolutionary Paths to Bacterial Antibiotic Resistance Cause Distinct Collateral Effects. *Mol. Biol. Evol.* 34, 2229–2244. doi: 10.1093/molbev/msx158

Baym, M., Stone, L. K., and Kishony, R. (2016). Multidrug evolutionary strategies to reverse antibiotic resistance. *Science* 351, aad3292. doi: 10.1126/science.aad3292

Berryhill, B. A., Huseby, D. L., McCall, I. C., Hughes, D., and Levin, B. R. (2021). Evaluating
the potential efficacy and limitations of a phage for joint antibiotic and phage therapy
of Staphylococcus aureus infections. *Proc. Natl. Acad. Sci.* 118, e2008007118. doi:
10.1073/pnas.2008007118

Bertels, F., Metzner, K. J., and Regoes, R. (2021). Convergent evolution as an indicator for
selection during acute HIV-1 infection. *Peer Community J.* 1, e4. doi:
10.24072/pcjournal.6

Bertozzi Silva, J., Storms, Z., and Sauvageau, D. (2016). Host receptors for bacteriophage
adsorption. *FEMS Microbiol. Lett.* 363. doi: 10.1093/femsle/fnw002

Bio-Conversion Databank Foundation (2021). *Bio-Convers. Databank Found. Portal.*
Available at: https://portal.bio-conversion.org/ (Accessed March 3, 2023).

Bolger, A. M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 30, 2114–2120. doi: 10.1093/bioinformatics/btu170

Burmeister, A. R., Fortier, A., Roush, C., Lessing, A. J., Bender, R. G., Barahman, R., et al.
(2020). Pleiotropy complicates a trade-off between phage resistance and antibiotic
resistance. *Proc. Natl. Acad. Sci.* 117, 11207–11216. doi: 10.1073/pnas.1919888117

Burns, J. L., Hedin, L. A., and Lien, D. M. (1989). Chloramphenicol resistance in
Pseudomonas cepacia because of decreased permeability. *Antimicrob. Agents*
Chemother. 33, 136–141.

Burns, S. M., and Hull, S. I. (1998). Comparison of Loss of Serum Resistance by Defined
Lipopolysaccharide Mutants and an Acapsular Mutant of Uropathogenic Escherichia
coli O75:K5. *Infect. Immun.* 66, 4244–4253.

Cairns, J., Becks, L., Jalasvuori, M., and Hiltunen, T. (2017). Sublethal streptomycin
concentrations and lytic bacteriophage together promote resistance evolution. *Philos.*
Trans. R. Soc. B Biol. Sci. 372, 20160040. doi: 10.1098/rstb.2016.0040

Chan, B. K., Turner, P. E., Kim, S., Mojibian, H. R., Elefteriades, J. A., and Narayan, D.
(2018). Phage treatment of an aortic graft infected with Pseudomonas aeruginosa.
Evol. Med. Public Health 2018, 60–66. doi: 10.1093/emph/eoy005

Chang, R. Y. K., Das, T., Manos, J., Kutter, E., Morales, S., and Chan, H.-K. (2019).
Bacteriophage PEV20 and Ciprofloxacin Combination Treatment Enhances Removal
of Pseudomonas aeruginosa Biofilm Isolated from Cystic Fibrosis and Wound
Patients. *AAPS J.* 21, 49. doi: 10.1208/s12248-019-0315-0

Chang, V., Chen, L.-Y., Wang, A., and Yuan, X. (2010). The Effect of Lipopolysaccharide
Core Structure Defects on Transformation Efficiency in Isogenic Escherichia coli
BW25113 rfaG, rfaP, and rfaC Mutants. 14, 7.

Chaudhry, W. N., Concepción-Acevedo, J., Park, T., Andleeb, S., Bull, J. J., and Levin, B. R.
(2017). Synergy and Order Effects of Antibiotics and Phages in Killing Pseudomonas
aeruginosa Biofilms. *PLOS ONE* 12, e0168615. doi: 10.1371/journal.pone.0168615

CLSI (2019). *Performance Standards for Antimicrobial Susceptibility Testing.* 29th ed.
Wayne, PA: Clinical and Laboratory Standards Institute.

Coates, J., Park, B. R., Le, D., Şimşek, E., Chaudhry, W., and Kim, M. (2018). Antibiotic-
induced population fluctuations and stochastic clearance of bacteria. *eLife* 7, e32976.
doi: 10.7554/eLife.32976

Coleman, W. G. (1983). The rfaD gene codes for ADP-L-glycero-D-mannoheptose-6-
epimerase. An enzyme required for lipopolysaccharide core biosynthesis. *J. Biol.*
Chem. 258, 1985–1990.

Comeau, A. M., Tétart, F., Trojet, S. N., Prère, M.-F., and Krisch, H. M. (2007). Phage-
Antibiotic Synergy (PAS): β-Lactam and Quinolone Antibiotics Stimulate Virulent
Phage Growth. *PLoS ONE* 2, e799. doi: 10.1371/journal.pone.0000799

Coulter, L. B., McLean, R. J. C., Rohde, R. E., and Aron, G. M. (2014). Effect of
Bacteriophage Infection in Combination with Tobramycin on the Emergence of
Resistance in Escherichia coli and Pseudomonas aeruginosa Biofilms. *Viruses* 6,
3778–3786. doi: 10.3390/v6103778

Czaplewski, L., Bax, R., Clokie, M., Dawson, M., Fairhead, H., Fischetti, V. A., et al. (2016).
Alternatives to antibiotics—a pipeline portfolio review. *Lancet Infect. Dis.* 16, 239–
251. doi: 10.1016/S1473-3099(15)00466-1

Deatherage, D. E., and Barrick, J. E. (2014). Identification of mutations in laboratory evolved
microbes from next-generation sequencing data using breseq. *Methods Mol. Biol.*
Clifton NJ 1151, 165–188. doi: 10.1007/978-1-4939-0554-6_12

Dötsch, A., Becker, T., Pommerenke, C., Magnowska, Z., Jänsch, L., and Häussler, S.
(2009). Genomewide Identification of Genetic Determinants of Antimicrobial Drug
Resistance in Pseudomonas aeruginosa. *Antimicrob. Agents Chemother.* 53, 2522–
2531. doi: 10.1128/aac.00035-09

Ebbensgaard, A., Mordhorst, H., Aarestrup, F. M., and Hansen, E. B. (2018). The Role of
Outer Membrane Proteins and Lipopolysaccharides for the Sensitivity of Escherichia
coli to Antimicrobial Peptides. *Front. Microbiol.* 9, 2153. doi:
10.3389/fmicb.2018.02153

Egido, J. E., Costa, A. R., Aparicio-Maldonado, C., Haas, P.-J., and Brouns, S. J. J. (2021).
Mechanisms and clinical importance of bacteriophage resistance. *FEMS Microbiol.*
Rev. 46, fuab048. doi: 10.1093/femsre/fuab048

El'Garch, F., Jeannot, K., Hocquet, D., Llanes-Barakat, C., and Plésiat, P. (2007).
Cumulative Effects of Several Nonenzymatic Mechanisms on the Resistance of
Pseudomonas aeruginosa to Aminoglycosides. *Antimicrob. Agents Chemother.* 51,
1016–1021. doi: 10.1128/AAC.00704-06

Farr, A. D., Pesce, D., Das, S. G., Zwart, M. P., and de Visser, J. A. G. M. (2023). The
Fitness of Beta-Lactamase Mutants Depends Nonlinearly on Resistance Level at
Sublethal Antibiotic Concentrations. *mBio* 14, e00098-23. doi: 10.1128/mbio.00098-
23

Foster, P. L. (2006). Methods for Determining Spontaneous Mutation Rates. *Methods*
Enzymol. 409, 195–213. doi: 10.1016/S0076-6879(05)09012-9

Frirdich, E., and Whitfield, C. (2005). Lipopolysaccharide inner core oligosaccharide structure
and outer membrane stability in human pathogens belonging to the
Enterobacteriaceae. *J. Endotoxin Res.* 11, 133–144. doi: 10.1179/096805105X46592

Frost, I., Smith, W. P. J., Mitri, S., Millan, A. S., Davit, Y., Osborne, J. M., et al. (2018).
Cooperation, competition and antibiotic resistance in bacterial colonies. *ISME J.* 12,
1582–1593. doi: 10.1038/s41396-018-0090-4

Gordillo Altamirano, F. L., Kostoulias, X., Subedi, D., Korneev, D., Peleg, A. Y., and Barr, J.
J. (2022). Phage-antibiotic combination is a superior treatment against Acinetobacter
baumannii in a preclinical study. *eBioMedicine* 80, 104045. doi:
10.1016/j.ebiom.2022.104045

Gu Liu, C., Green, S. I., Min, L., Clark, J. R., Salazar, K. C., Terwilliger, A. L., et al. (2020).
Phage-Antibiotic Synergy Is Driven by a Unique Combination of Antibacterial
Mechanism of Action and Stoichiometry. *mBio* 11, e01462-20. doi:
10.1128/mBio.01462-20

Hadas, H., Einav, M., Fishov, I., and Zaritsky, A. (1997). Bacteriophage T4 Development
Depends on the Physiology of its Host Escherichia Coli. *Microbiology* 143, 179–185.
doi: 10.1099/00221287-143-1-179

Hancock, R. E., and Reeves, P. (1976). Lipopolysaccharide-deficient, bacteriophage-
resistant mutants of Escherichia coli K-12. *J. Bacteriol.* 127, 98–108.

Henriksen, K., Rørbo, N., Rybtke, M. L., Martinet, M. G., Tolker-Nielsen, T., Høiby, N., et al.
(2019). P. aeruginosa flow-cell biofilms are enhanced by repeated phage treatments
but can be eradicated by phage-ciprofloxacin combination. *Pathog. Dis.* 77, ftz011.
doi: 10.1093/femspd/ftz011

Ibacache-Quiroga, C., Oliveros, J. C., Couce, A., and Blázquez, J. (2018). Parallel Evolution
of High-Level Aminoglycoside Resistance in Escherichia coli Under Low and High
Mutation Supply Rates. *Front. Microbiol.* 9, 427. doi: 10.3389/fmicb.2018.00427

Kamal, F., and Dennis, J. J. (2015). Burkholderia cepacia Complex Phage-Antibiotic Synergy
(PAS): Antibiotics Stimulate Lytic Phage Activity. *Appl. Environ. Microbiol.* 81, 1132–
1138. doi: 10.1128/AEM.02850-14

Kebriaei, R., Lev, K. L., Shah, R. M., Stamper, K. C., Holger, D. J., Morrisette, T., et al.
(2022). Eradication of Biofilm-Mediated Methicillin-Resistant Staphylococcus aureus
Infections In Vitro: Bacteriophage-Antibiotic Combination. *Microbiol. Spectr.* 10,
e00411-22. doi: 10.1128/spectrum.00411-22

Kehrenberg, C., Schwarz, S., Jacobsen, L., Hansen, L. H., and Vester, B. (2005). A new
mechanism for chloramphenicol, florfenicol and clindamycin resistance: methylation
of 23S ribosomal RNA at A2503. *Mol. Microbiol.* 57, 1064–1073. doi: 10.1111/j.1365-
2958.2005.04754.x

Kirby, A. E. (2012). Synergistic Action of Gentamicin and Bacteriophage in a Continuous
Culture Population of Staphylococcus aureus. *PLoS ONE* 7, e51017. doi:
10.1371/journal.pone.0051017

Klein, G., and Raina, S. (2019). Regulated Assembly of LPS, Its Structural Alterations and
Cellular Response to LPS Defects. *Int. J. Mol. Sci.* 20, 356. doi:
10.3390/ijms20020356

Kortright, K. E., Doss-Gollin, S., Chan, B. K., and Turner, P. E. (2021). Evolution of Bacterial
Cross-Resistance to Lytic Phages and Albicidin Antibiotic. *Front. Microbiol.* 12, 1423.
doi: 10.3389/fmicb.2021.658374

Labrie, S. J., Samson, J. E., and Moineau, S. (2010). Bacteriophage resistance mechanisms.
Nat. Rev. Microbiol. 8, 317–327. doi: 10.1038/nrmicro2315

Lin, J., Zhou, D., Steitz, T. A., Polikanov, Y. S., and Gagnon, M. G. (2018). Ribosome-
Targeting Antibiotics: Modes of Action, Mechanisms of Resistance, and Implications

for Drug Design. *Annu. Rev. Biochem.* 87, 451–478. doi: 10.1146/annurev-biochem-
062917-011942

Lind, P. A., Libby, E., Herzog, J., and Rainey, P. B. (2019). Predicting mutational routes to
new adaptive phenotypes. *eLife* 8, e38822. doi: 10.7554/eLife.38822

Lopes, A., Pereira, C., and Almeida, A. (2018). Sequential Combined Effect of Phages and
Antibiotics on the Inactivation of Escherichia coli. *Microorganisms* 6, 125. doi:
10.3390/microorganisms6040125

Lu, T. K., and Collins, J. J. (2009). Engineered bacteriophage targeting gene networks as
adjuvants for antibiotic therapy. *Proc. Natl. Acad. Sci.* 106, 4629–4634. doi:
10.1073/pnas.0800442106

Luria, S. E., and Delbrück, M. (1943). Mutations of Bacteria from Virus Sensitivity to Virus
Resistance. *Genetics* 28, 491–511.

Mahrt, N., Tietze, A., Künzel, S., Franzenburg, S., Barbosa, C., Jansen, G., et al. (2021).
Bottleneck size and selection level reproducibly impact evolution of antibiotic
resistance. *Nat. Ecol. Evol.* 5, 1233–1242. doi: 10.1038/s41559-021-01511-2

Marx, A., Backes, C., Meese, E., Lenhof, H.-P., and Keller, A. (2016). EDISON-WMW: Exact
Dynamic Programing Solution of the Wilcoxon–Mann–Whitney Test. *Genomics*
Proteomics Bioinformatics 14, 55–61. doi: 10.1016/j.gpb.2015.11.004

Mayr, U. B., Walcher, P., Azimpour, C., Riedmann, E., Haller, C., and Lubitz, W. (2005).
Bacterial ghosts as antigen delivery vehicles. *Adv. Drug Deliv. Rev.* 57, 1381–1391.
doi: 10.1016/j.addr.2005.01.027

Michel, A., Clermont, O., Denamur, E., and Tenaillon, O. (2010). Bacteriophage PhiX174's
Ecological Niche and the Flexibility of Its Escherichia coli Lipopolysaccharide
Receptor. *Appl. Environ. Microbiol.* 76, 7310–7313. doi: 10.1128/AEM.02721-09

Moxon, E. R., Rainey, P. B., Nowak, M. A., and Lenski, R. E. (1994). Adaptive evolution of
highly mutable loci in pathogenic bacteria. *Curr. Biol.* 4, 24–33. doi: 10.1016/S0960-
9822(00)00005-1

Mutalik, V. K., Adler, B. A., Rishi, H. S., Piya, D., Zhong, C., Koskella, B., et al. (2020). High-
throughput mapping of the phage resistance landscape in E. coli. *PLOS Biol.* 18,
e3000877. doi: 10.1371/journal.pbio.3000877

Nikaido, H. (2003). Molecular Basis of Bacterial Outer Membrane Permeability Revisited.
Microbiol. Mol. Biol. Rev. 67, 593–656. doi: 10.1128/MMBR.67.4.593-656.2003

Nikaido, H., and Vaara, M. (1985). Molecular Basis of Bacterial Outer Membrane
Permeability. *MICROBIOL REV* 49, 32. doi: 10.1128/mr.49.1.1-32.1985

Orta, A. K., Riera, N., Li, Y. E., Tanaka, S., Yun, H. G., Klaic, L., et al. (2023). The
mechanism of the phage-encoded protein antibiotic from ΦX174. *Science* 381,
eadg9091. doi: 10.1126/science.adg9091

Pagnout, C., Sohm, B., Razafitianamaharavo, A., Caillet, C., Offroy, M., Leduc, M., et al.
(2019). Pleiotropic effects of rfa-gene mutations on Escherichia coli envelope
properties. *Sci. Rep.* 9, 9696. doi: 10.1038/s41598-019-46100-3

Picelli, S., Björklund, A. K., Reinius, B., Sagasser, S., Winberg, G., and Sandberg, R. (2014).
Tn5 transposase and tagmentation procedures for massively scaled sequencing
projects. *Genome Res.* 24, 2033–2040. doi: 10.1101/gr.177881.114

Pier, G. B. (2007). Pseudomonas aeruginosa lipopolysaccharide: a major virulence factor,
initiator of inflammation and target for effective immunity. *Int. J. Med. Microbiol. IJMM*
297, 277–295. doi: 10.1016/j.ijmm.2007.03.012

Pirnay, J.-P., Ferry, T., and Resch, G. (2022). Recent progress toward the implementation of
phage therapy in Western medicine. *FEMS Microbiol. Rev.* 46, fuab040. doi:
10.1093/femsre/fuab040

Pradel, E., and Schnaitman, C. A. (1991). Effect of rfaH (sfrB) and temperature on
expression of rfa genes of Escherichia coli K-12. *J. Bacteriol.* 173, 6428–6431.

R Core Team (2020). *R: A Language and Environment for Statistical Computing.* Vienna,
Austria: R Foundation for Statistical Computing. Available at: https://www.R-
project.org/

Raetz, C. R. H., and Whitfield, C. (2002). Lipopolysaccharide Endotoxins. *Annu. Rev.*
Biochem. 71, 635–700. doi: 10.1146/annurev.biochem.71.110601.135414

Recht, M. I., and Puglisi, J. D. (2001). Aminoglycoside Resistance with Homogeneous and
Heterogeneous Populations of Antibiotic-Resistant Ribosomes. *Antimicrob. Agents*
Chemother. 45, 2414–2419. doi: 10.1128/AAC.45.9.2414-2419.2001

Romeyer Dherbey, J., and Bertels, F. (2024). The untapped potential of phage model
systems as therapeutic agents. *Virus Evol.* 10, veae007. doi: 10.1093/ve/veae007

Romeyer Dherbey, J., Parab, L., Gallie, J., and Bertels, F. (2023). Stepwise Evolution of E.
coli C and ΦX174 Reveals Unexpected Lipopolysaccharide (LPS) Diversity. *Mol. Biol.*
Evol. 40, msad154. doi: 10.1093/molbev/msad154

Rosche, W. A., and Foster, P. L. (2000). Determining Mutation Rates in Bacterial
Populations. *Methods San Diego Calif* 20, 4–17. doi: 10.1006/meth.1999.0901

RStudio Team (2019). *RStudio: Integrated Development Environment for R.* Boston, MA:
RStudio, Inc. Available at: http://www.rstudio.com/

Sanger, F., Air, G. M., Barrell, B. G., Brown, N. L., Coulson, A. R., Fiddes, J. C., et al. (1977).
Nucleotide sequence of bacteriophage φX174 DNA. *Nature* 265, 687–695. doi:
10.1038/265687a0

Schnaitman, C. A., and Klena, J. D. (1993). Genetics of lipopolysaccharide biosynthesis in
enteric bacteria. *Microbiol. Rev.* 57, 655–682. doi: 10.1128/mr.57.3.655-682.1993

Schooley, R. T., Biswas, B., Gill, J. J., Hernandez-Morales, A., Lancaster, J., Lessor, L., et
al. (2017). Development and Use of Personalized Bacteriophage-Based Therapeutic
Cocktails To Treat a Patient with a Disseminated Resistant Acinetobacter baumannii
Infection. *Antimicrob. Agents Chemother.* 61, e00954-17. doi: 10.1128/AAC.00954-17

Simon, A. (2010). FastQC A Quality Control tool for High Throughput Sequence Data.
Available at: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/ (Accessed
April 19, 2023).

Sinsheimer, R. L. (1959). Purification and properties of bacteriophage φX174. *J. Mol. Biol.* 1,
37-IN5. doi: 10.1016/S0022-2836(59)80005-X

Smith, H. O., Hutchison, C. A., Pfannkoch, C., and Venter, J. C. (2003). Generating a
synthetic genome by whole genome assembly: φX174 bacteriophage from synthetic
oligonucleotides. *Proc. Natl. Acad. Sci.* 100, 15440–15445. doi:
10.1073/pnas.2237126100

Suzuki, M., Kaneko-Tanaka, Y., and Azegami, M. (1974). Transfection of non-host bacterial
spheroplasts with bacteriophage ΦX174 DNA. *Nature* 252, 319–321. doi:
10.1038/252319a0

Tagliaferri, T. L., Jansen, M., and Horz, H.-P. (2019). Fighting Pathogenic Bacteria on Two
Fronts: Phages and Antibiotics as Combined Strategy. *Front. Cell. Infect. Microbiol.* 9,
22. doi: 10.3389/fcimb.2019.00022

Torres-Barceló, C., Arias-Sánchez, F. I., Vasse, M., Ramsayer, J., Kaltz, O., and Hochberg,
M. E. (2014). A Window of Opportunity to Control the Bacterial Pathogen
Pseudomonas aeruginosa Combining Antibiotics and Phages. *PLoS ONE* 9,
e106628. doi: 10.1371/journal.pone.0106628

Valério, N., Oliveira, C., Jesus, V., Branco, T., Pereira, C., Moreirinha, C., et al. (2017).
Effects of single and combined use of bacteriophages and antibiotics to inactivate
Escherichia coli. *Virus Res.* 240, 8–17. doi: 10.1016/j.virusres.2017.07.015

Wang, Z., Wang, J., Ren, G., Li, Y., and Wang, X. (2015). Influence of Core Oligosaccharide
of Lipopolysaccharide to Outer Membrane Behavior of Escherichia coli. *Mar. Drugs*
13, 3325–3339. doi: 10.3390/md13063325

Whitfield, C., Heinrichs, D. E., Yethon, J. A., Amor, K. L., Monteiro, M. A., and Perry, M. B.
(1999). Assembly of the R1-type core oligosaccharide of Escherichia coli
lipopolysaccharide. *J. Endotoxin Res.* 5, 151–156. doi:
10.1177/09680519990050030901

World Health Organization (2014). *Antimicrobial resistance: global report on surveillance.*
World Health Organization. Available at:
https://apps.who.int/iris/handle/10665/112642 (Accessed October 18, 2021).

Yehl, K., Lemire, S., Yang, A. C., Ando, H., Mimee, M., Torres, M. D. T., et al. (2019).
Engineering Phage Host-Range and Suppressing Bacterial Resistance through
Phage Tail Fiber Mutagenesis. *Cell* 179, 459-469.e9. doi: 10.1016/j.cell.2019.09.015

Yethon, J. A., Heinrichs, D. E., Monteiro, M. A., Perry, M. B., and Whitfield, C. (1998).
Involvement of waaY, waaQ, and waaP in the modification of Escherichia coli
lipopolysaccharide and their role in the formation of a stable outer membrane. *J. Biol.*
Chem. 273, 26310–26316. doi: 10.1074/jbc.273.41.26310

Yethon, J. A., Vinogradov, E., Perry, M. B., and Whitfield, C. (2000). Mutation of the
lipopolysaccharide core glycosyltransferase encoded by waaG destabilizes the outer
membrane of Escherichia coli by interfering with core phosphorylation. *J. Bacteriol.*
182, 5620–5623. doi: 10.1128/JB.182.19.5620-5623.2000

Zheng, Q. (2017). rSalvador: An R Package for the Fluctuation Experiment. *G3*
GenesGenomesGenetics 7, 3849–3856. doi: 10.1534/g3.117.300120

1062