Encyclopaedia of DNA polymerases localized in organelles: Evolutionary contribution of diverse bacteria including the proto-mitochondrion

Ryo Harada[1], Yoshihisa Hirakawa[2], Akinori Yabuki[3], Eunsoo Kim[4,5], Euki Yazaki[6], Ryoma Kamikawa[7], Kentaro Nakano[1], Marek Eliáš[8,*], Yuji Inagaki[1,9,*]

[1]Graduate School of Life and Environmental Sciences, University of Tsukuba, Japan
[2]Faculty of Life and Environmental Sciences, University of Tsukuba, Japan
[3]Deep-Sea Biodiversity Research Group, Research Institute for Global Change (RIGC), Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokosuka, Japan
[4]Division of EcoScience, Ewha Womans University, Seoul, South Korea
[5]Division of Invertebrate Zoology, American Museum of Natural History, New York, USA
[6]Research Center for Advanced Analysis, National Agriculture and Food Research Organization, Tsukuba, Japan
[7]Graduate School of Agriculture, Kyoto University, Kyoto, Japan
[8]Department of Biology and Ecology, Faculty of Science, University of Ostrava, Ostrava, Czech Republic
[9]Center for Computational Sciences, University of Tsukuba, Japan

*Authors for correspondence: marek.elias@osu.cz (M.E.), yuji@ccs.tsukuba.ac.jp (Y.I.)

Abstract

DNA polymerases (DNAPs) synthesize DNA from deoxyribonucleotides in a semi-
conservative manner and serve as the core of DNA replication and repair machineries. In
eukaryotic cells, there are two genome-containing organelles, mitochondria and plastids, that
were derived from an α-proteobacterium and a cyanobacterium, respectively. Except for rare
cases of genome-lacking mitochondria and plastids, both organelles must be served by
nucleus-encoded DNAPs that localize and work in them to maintain their genomes. The
evolution of organellar DNAPs has yet to be fully understood because of two unsettled
issues. First, the diversity of organellar DNAPs has not been elucidated in the full spectrum
of eukaryotes. Second, it is unclear when the DNAPs that were used originally in the
endosymbiotic bacteria giving rise to mitochondria and plastids were discarded, as the
organellar DNAPs known to date show no phylogenetic affinity to those of the extant α-
proteobacteria or cyanobacteria. In this study, we identified from diverse eukaryotes 134
family A DNAP sequences, which were classified into 10 novel types, and explored their
evolutionary origins. The subcellular localizations of selected DNAPs were further examined
experimentally. The results presented here suggest that the diversity of organellar DNAPs
has been shaped by multiple transfers of the PolI gene from phylogenetically broad bacteria,
and their occurrence in eukaryotes was additionally impacted by secondary plastid
endosymbioses. Finally, we propose that the last eukaryotic common ancestor may have
possessed two mitochondrial DNAPs, POP and a candidate of the direct descendant of the
proto-mitochondrial DNAP, rdxPolA, identified in this study.
48
49 Keywords: DNA polymerase, endosymbiosis, lateral gene transfer, LECA, mitochondria,
50 plastids

Introduction

Multiple endosymbiotic events played pivotal roles in the evolution of eukaryotes. The most
ancient endosymbiotic event that is currently recognizable likely occurred between the
primordial (proto)eukaryotic or archaeal cell (host) and an α-proteobacterium
(endosymbiont). This event gave rise to mitochondria, which operate metabolic pathways
that are critical for cell viability, such as oxidative energy production, amino acid synthesis,
β-oxidation of fatty acids, and Fe-S cluster assembly, in modern eukaryotic cells (Roger et al.
2017). Another bacterial endosymbiosis took place later than mitochondrial endosymbiosis in
eukaryotic evolution and yielded a photosynthetic organelle, the plastid. In primary plastid
endosymbiosis, the host and the endosymbiont were the common ancestor of
Archaeplastida and a cyanobacterium, respectively (Sibbald and Archibald 2020). The host
organism involved in the primary endosymbiosis then diverged into green algae plus land
plants (Chloroplastida), glaucophyte algae (Glaucophyta), and red algae (Rhodophyta) plus
their non-photosynthetic relatives, so that their plastids are termed primary plastids.
Photosynthesis further spread into phylogenetically diverged eukaryotes through multiple
"secondary endosymbiosis," in which a green or red alga was engulfed by a heterotroph and
transformed subsequently into a "complex plastid" (Sibbald and Archibald 2020). Green
alga-derived complex plastids have been found in two distantly related branches in the tree
of eukaryotes, namely Euglenophyceae and Chlorarachniophyta (ignoring a few
dinoflagellates also bearing chlorophyte-derived plastids (Kamikawa et al. 2015; Sarai et al.

2020; Matsuo et al. 2022)). On the other hand, the complex plastids in Ochrophyta,
Myzozoa, Haptophyta, and Cryptophyceae can be traced back to red algal endosymbionts.
Reflecting the bacterial origins of mitochondria and plastids, the two organelles
typically retain their own genomes, which are however highly reduced compared to their
free-living relatives (i.e., α-proteobacteria and cyanobacteria). Disregarding sporadic cases
related to the insertion of mobile genetic elements into mitochondrial (e.g., Burger et al.
1999; Nishimura et al. 2019) or plastid genomes (e.g., Kim et al. 2015), organellar genomes
do not encode proteins involved in DNA replication and repair. Hence, DNA maintenance in
organelles entirely relies on the set of proteins encoded in the nuclear genome, synthesized
in the cytosol, and then translocated into the corresponding organelle. DNA polymerases
(DNAPs), which catalyze the synthesis of the nascent DNA strand by referring to the
template strand, are the core components that are central to DNA replication.
Nucleus-encoded organellar DNAPs have been reported from diverse eukaryotes
(Klingbeil et al. 2002; Graziewicz et al. 2006; Moriyama et al. 2008; Mukhopadhyay et al.
2009; Moriyama et al. 2011; Moriyama and Sato 2014; Janouškovec et al. 2015; Hirakawa
and Watanabe 2019; Harada et al. 2020; Harada and Inagaki 2021). DNA-dependent
DNAPs are generally divided into six families (A, B, C, D, X, and Y) (Filée et al. 2002), and
organellar DNAPs belong to family A (famA) that is typified by the bacterial DNA polymerase
Ⅰ (PolⅠ). Bacterial PolⅠ is responsible for filling the gap between the Okazaki fragments
synthesized by PolⅢ (family C) during the lagging strand synthesis (Okazaki 2017). In
organelles, PolⅢ is absent, and PolⅠ-related (i.e., family A) DNAPs play a major role in
genome replication (Graziewicz et al. 2006; Parent et al. 2011). The known organellar
DNAPs can be subdivided into five phylogenetically distinct types, namely Polγ, POP, PREX,
PolIA, and PolIBCD+ (Harada and Inagaki 2021). POP has been found in phylogenetically
diverse eukaryotic lineages and can be targeted to mitochondria, plastids, or both organelles
in the same cell (i.e., dual-targeted) depending on the species (Kimura et al. 2002;
Christensen et al. 2005; Mori et al. 2005; Ono et al. 2007; Moriyama et al. 2008; Parent et al.
2011; Hirakawa and Watanabe 2019). The origin of POP remains uncertain, as it forms a
lineage of its own, lacking any specific relatives in famA DNAPs phylogenies presented so
far. In contrast, Polγ, PREX, PolIA, and PolIBCD+ appear to be lineage-specific. Polγ is the
mitochondrial DNAP found exclusively in opisthokonts. PREX has been reported only from
apicomplexans and closely related lineages (chrompodellids and squirmids), and works in
their plastids (Janouškovec et al. 2019). PolIA has been found in all three major classes
(Euglenoidea, Diplonemea, and Kinetoplastea) comprising the phylum Euglenozoa, and
PolIBCD+ is restricted to Diplonemea and Kinetoplastea (Harada et al. 2020; Harada and
Inagaki 2021). The origin of the different organellar DNAPs as elucidated by phylogenetic
analyses varies among them. PREX was acquired from a bacterial source, although the
specific bacterial group has not been resolved (Janouškovec et al. 2019). Polγ and
PolIBCD+ are derived from different bacteriophages (Filée et al. 2002; Harada and Inagaki
2021), and PolIA most likely shares the ancestry with two closely related nuclear famA
DNAPs, Polθ and Polν (Harada and Inagaki 2021). Importantly, none of the five types of
organellar DNAP has any phylogenetic affinity to the PolⅠ sequences of α-proteobacteria or
cyanobacteria, which are the origins of mitochondria and plastids, respectively.
While the recognition of the five aforementioned types of famA DNAP and their
taxonomic distribution among eukaryotes established by previous studies defines the
mechanistic basis for organellar genome replication and repair in the vast portion of the
eukaryote phylogenetic diversity, there are indications they do not constitute a full picture of

the organellar DNAP spectrum. Indeed, putative organelle-localized famA DNAPs unrelated
to any of the five well-defined types were identified in a single red alga (Moriyama et al.
2008) and a single species belonging to Discoba (Gray et al. 2020). In this study, we
conducted a comprehensive survey of famA DNAPs in eukaryotes, which led to the
discovery of several novel types of organellar DNAP, including a previously unnoticed DNAP
that appears to be a direct genetic legacy of the α-proteobacterial ancestor of the
mitochondrion.

125 Results & Discussion

126 Diversity and subcellular localizations of the novel types of famA DNAP
127 in eukaryotes

We surveyed various sequence databases collectively covering the whole breath of the
eukaryote phylogenetic diversity as captured by sequencing efforts so far. Phylogenetic
analyses and sequence curation enabled us to classify the retrieved sequences and
discriminate genuine eukaryotic genes from bacterial contaminants. We focused on
identifying putative organellar DNAPs in major eukaryote lineages that have not been
investigated in this regard in the past and on discovering possible novel famA DNAP types
unrelated to POP, Polγ, PREX, PolIA, or PolIBCD+. In this study, we identified 134 famA
DNAPs in diverse eukaryotes that were distantly related to any of the previously known
eukaryotic famA DNAPs. We prepared a "global famA DNAP" alignment by including diverse
bacterial and phage PolI sequences, the six previously known types of eukaryotic famA
DNAP sequences, and the famA DNAP sequences identified in this study. From this
alignment, we reconstructed the maximum likelihood (ML) tree and calculated ultra-fast
bootstrap supports (UFBPs) for bipartitions in the ML tree. Based on the global famA DNAP
phylogeny (Fig. 1), the novel types of famA DNAP coalesced into ten clades, which we
denote "alvPolA," "abanPolA," "acPolA," "rgPolA," "chlnmPolA," "eugPolA," "pyramiPolA,"
"chloroPolA," "cryptoPolA," and "rdxPolA" (Table 1). The putative domain structure of the
representative of each of the ten novel DNAPs is provided in Fig. S1. Importantly, the global
famA DNAP phylogeny demonstrated that none of the ten novel DNAPs appeared to be
related to any previously known famA DNAPs in eukaryotes (Fig. 1). For each novel famA
DNAP type introduced above, we described the phylogenetic distribution and evaluated the
subcellular localization primarily by considering experimental data. In the case of no
experimental data for a particular famA DNAP, we referred to the results from *in silico*
prediction tools (Supplementary Table S1).

151 alvPolA

Our survey revealed that certain representatives of Ciliophora, Dinoflagellata, Squirmidea,
Chrompodellida, and Apicomplexa, which all belong to Alveolata, possess previously
undescribed famA DNAP sequences that formed a clade supported by an UFBP of 100% in
the global famA DNAP phylogeny (Fig. 1). We hence denote the DNAP sequences found in
alveolates as "alvPolA." It is currently ambiguous whether alvPolA is ancestral for Alveolata
as a whole, as this particular DNAP type was not found in the sequence data available for
Colponemidia, a lineage sister to all other alveolates (Tikhonenkov et al. 2020). Furthermore,
this type of DNAP is sporadically found in this group. For instance, the alvPolA sequences
were found solely in two heterotricheans in Ciliophora, and only a single gregarine appeared

to possess alvPolA among the apicomplexans studied here (Table S1). Given the
occurrence of this DNAP type in plastid-lacking members of Alveolata, namely ciliates, it is
unlikely to be plastid-localized. Indeed, *in silico* analyses of the N-terminal pre-sequences of
alvPolA provided no strong evidence for organellar localization of this DNAP type (Table S1).
Combined, we regard alvPolA as nucleus-localized.

abanPolA

The global famA DNAP phylogeny reconstructed a clade of the novel DNAP sequences
identified in the members of three subclades of the eukaryotic "megagroup" Amorphea,
namely Apusomonadida, Breviatea, and Amoebozoa, as well as Nebulidia, a subgroup of
the supergroup Provora (hereafter designated as "abanPolA"). All the abanPolA-bearing
eukaryotes lack a plastid. Furthermore, abanPolA is present in the breviates *Pygsuia biforma*
and *Lensia limosa*, whose mitochondria are highly reduced with no genome (Table S1).
Thus, the phylogenetic distribution of abanPolA indirectly but strongly argues for this DNAP
type being involved in maintaining the nuclear genome. Consistent with the above notion,
little sign of a targeting signal for any organelle was suggested by the *in silico* analyses of
the full-length abanPolA sequences (Table S1).

acPolA

A group of previously unclassified DNAP sequences identified in two chrompodellids and 12
apicomplexans formed a clade with an UFBP of 100%. Prior to this study, the cellular
function of the *Plasmodium falciparum* representative was experimentally confirmed (Reesey
2017). The knockdown of the DNAP gene caused a decrease in the amount of mitochondrial
DNA, suggesting that this DNAP participates in DNA maintenance in the *P. falciparum*
mitochondrion. By extrapolating the result from the experiment on *P. falciparum* to other
apicomplexans and chrompodellids, the novel DNAPs discussed here are regarded as
mitochondrion-localized, although bioinformatic assessment of this notion is hampered by N-
terminal truncation in most of the available acPolA sequences. Recent studies placed a
newly recognized parasitic group, Squirmidea, at the base of Apicomplexa and
Chrompodellida (Janouškovec et al. 2019; Mathur et al. 2019; Salomaki et al. 2021; Yazaki
et al. 2021). We searched for putative mitochondrion-localized famA DNAP sequences in
Squirmidea and found only a single candidate, POP, in the transcriptome assembly of
Digyalum oweni. Thus, the novel famA DNAP group discussed here is most likely specific to
the organisms constituting the Apicomplexa + Chrompodellida clade and henceforth termed
"acPolA." In future, the phylogenetic distribution of acPolA should be examined rigorously by
surveys of famA DNAPs in high quality transcriptomes of squirmids.

rgPolA

A type of famA DNAP appeared to be shared among all the eight red algae and two
glaucophyte algae surveyed in this study. The ten DNAP sequences, together with the
previously known DNAP in the red alga *Cyanidioschyzon merolae* (Moriyama et al. 2008),
formed a clade with an UFBP of 100% in the global famA DNAP phylogeny (Fig. 1). We here
propose that the novel DNAPs were derived from a single ancestral DNAP. The plastid
localization of the *C. merolae* homolog was determined experimentally, suggesting that this
type of DNAP is localized in the plastids of red and glaucophyte algae in addition to the dual-
targeted POP (Moriyama et al. 2008). We designated this famA DNAP type as "rgPolA" to
reflect the restriction to Rhodophyta and Glaucophyta.

We found distinct famA DNAP types that were specific to Euglenophyceae,
Chlorarachniophyta, and Cryptophyceae and here termed "eugPolA," "chlnmPolA," and
"cryptoPolA," respectively. The global famA DNAP phylogeny united the four four eugPolA
sequences, chlnmPolA sequences, and five cryptoPolA into the individual clades supported
by UFBPs of 99-100% (Fig. 1).
The eugPolA sequence of the euglenid *Euglena gracilis* was detected specifically in
the plastid proteome (Novák Vanclová et al. 2020), suggesting that this DNAP type is
plastid-localized. Besides *E. gracilis*, the eugPolA homologs were found in the species
belonging to Euglenales and Eutreptiales in Euglenophyceae. *Euglena longa*, which is
heterotrophic but retains a non-photosynthetic plastid (Gockel and Hachtel 2000; Füssy et al.
2020), also possesses a eugPolA gene. No eugPolA sequence was detected in the
transcriptome of *Rapaza viridis*, which represents a lineage of Euglenophyceae sister to
Euglenales and Eutreptiales combined (Yamaguchi et al. 2012; Karnkowska et al. 2023). As
R. viridis is a kleptoplastidic phagotroph with no permanent plastid, we propose that eugPolA
co-occurs with the permanent Pyramimonadales-derived plastids in Euglenophyceae.
In Chlorarachniophyta and Cryptophyceae, four subcellular compartments contain
evolutionarily different genomes, i.e., a nucleus, a mitochondrion, a plastid, and a remnant
nucleus of endosymbiont (nucleomorph). The subcellular localization of chlnmPolA was
investigated by employing the model chlorarachniophyte *Amorphochlora amoebiformis*. As
the N-terminal region was absent in the *A. amoebiformis* chlnmPolA sequence reconstructed
from the transcriptome data (Table S1), we instead used the N-terminal amino acid residues
of the chlnmPolA sequence of the related species *Bigelowiella natans* and expressed it in *A.*
amoebiformis as a translational fusion with GFP. This N-terminal leader sequence navigated
GFP into the periplastidial compartment (PPC; Figs. 2 and S2). The PPC in
chlorarachniophytes harbors a nucleomorph derived from the green algal endosymbiont
taken up by the chlorarachniophyte ancestor. Thus, we conclude that chlnmPolA is
involved in the DNA maintenance of nucleomorphs in chlorarachniophyte cells.
As no cryptophyte transgene-expression system has been developed, we expressed
the N-terminal region of the *Guillardia theta* cryptoPolA as a translation fusion with GFP
heterogeneously in the model diatom *Phaeodactylum tricornutum*. This organism was
chosen because diatoms and cryptophytes share red alga-derived plastids with the same
membrane organization and the plastid protein-targeting mechanism (Nassoury and Morse
2005; Felsner et al. 2011). The green fluorescence signal appeared to overlap with the
plastid autofluorescence (Fig. 3A), indicating that the N-terminus of the *G. theta* cryptoPolA
was recognized as a plastid targeting signal (PTS) in the diatom cells. This is consistent with
the sequence features of N-terminus of the *G. theta* cryptoPolA protein, which includes a
predicted signal peptide (Fig. S3A). Phenylalanine residues appeared to be highly conserved
right after the predicted cleavage sites among the plastid-destined proteins in cryptophytes
(Gould et al. 2006; Patron and Waller 2007). Intriguingly, in the N-terminus of the *G. theta*
cryptoPolA, we found a phenylalanine residue after five amino acids away from the putative
cleavage site (Fig. S3A). The same features are conserved in other cryptoPolA proteins for
which a complete N-terminal sequence is available (Figs. S3B and S3C), suggesting that
cryptoPolA is targeted to the plastid in cryptophytes in general.
Besides cryptoPolA, *G. theta* possesses two distinct POP homologs (Hirakawa and
Watanabe 2019), raising the question of their function in the presence of cryptoPolA. Since
one of these POPs was a partial sequence and both gene models were potentially

inaccurate, we obtained the complete sequences of POP1 (JGI_79451) and POP2
(JGI_199174) from transcriptomic data based on RNA-seq data of PRJNA509798 and
PRJNA634446. We repeated the same experiments expressing the POP1 and POP2 N-
termini fused to GFP in the diatom cells. In the experiment assessing the localization of
POP1, the fluorescence of GFP and MitoTracker Orange, not the plastid autofluorescence,
overlapped each other, suggesting that the POP1 N-terminus functioned as a mitochondrial
targeting signal (MTS) in the diatom cells (Figs. 3B and 3C). On the other hand, we
observed two patterns of the subcellular localization of the green fluorescence signal in the
POP2 experiment. In the majority of the transformants, the green fluorescence was
observed as small blobs associated with plastids (Fig. 3D). As similar localization patterns
have been observed for the PPC proteins of diatoms (Gould et al. 2006; Moog et al. 2020),
POP2 may have been localized in the PPC of the diatom cells. In case of POP2 being a
genuine PPC protein of *G. theta*, this DNAP needs to be localized in the nucleomorph that is
a sole DNA-containing structure in the cryptophyte PPC. Nevertheless, in a small number of
the transformants examined, the green fluorescence overlapped with the plastid
autofluorescence (Fig. 3E), suggesting that the N-terminus of the *G. theta* POP2 can
function as PTS in the diatom cells. However, no typical signal peptide was predicted in the
N-terminus of the *G. theta* POP2 *in silico* (Fig.S3D). Although it is difficult to establish the
definite subcellular localization of POP2 based on the experiment expressing a cryptophyte
protein in a diatom, we propose POP2 as a nucleomorph protein rather than a plastid
protein. The subcellular localization of POP2 should be confirmed by electron microscopic
observations using an antibody against the DNAP of interest.

pyramiPolA and chloroPolA

We found two novel types of famA DNAP that are distributed in distinct sublineages of
Chloroplastida. First, at least four green algal species of Pyramimonadales possess a unique
type of famA DNAP, and these DNAP sequences formed a clade with an UFBP of 99% in
the global famA DNAP phylogeny (Fig. 1). The Pyramimonadales-specific famA DNAP is
termed as "pyramiPolA." All the pyamiPolA sequences were predicted to be localized in
mitochondria by PredSL (Table S1). Nevertheless, MitoFate and NommPred, the programs
designed to detect mitochondrial proteins exclusively, or PredAlgo, a program tuned for
green algal proteins, predicted no MTS in any of the four sequences (Table S1). We thus
hesitate to discuss further the subcellular localization of pyramiPolA until additional
information, preferentially experimental data, becomes available.
Another novel type of famA DNAP was detected in 27 species representing the class
Chlorophyceae and belonging to both its principal lineages known as CS and OCC clades.
An additional survey of the OneKP database containing transcriptome assemblies of a large
phylogenetically diverse set of green algal and plant species (Leebens-Mack et al. 2019)
strengthened the case for this novel DNAP type being restricted to Chlorophyceae, and thus
we here designate this DNA type as "chloroPolA." The Chlorophyceae-specific DNAP
sequences and cyanobacterial PolI sequences grouped together with an UFBP of 99% in
the global famA DNAP phylogeny (Fig. 1). We subjected 11 full-length chloroPolA
sequences to *in silico* prediction of subcellular localization, yet the predicted localizations
varied depending on the sequence analyzed or the program used (Table S1). We
additionally analyzed the same sequences by PredAlgo, a bioinformatic program that was
specifically designed to predict protein subcellular localization in green algae (Tardif et al.
2012). Intriguingly, five out of the 11 chloroPolA sequences were predicted as

mitochondrion-localized proteins. Thus, we tentatively regard chloroPolA as the DNAP
localized in mitochondria in chlorophycean green algae, albeit our conclusion needs to be re-
examined experimentally in the future.

rdxPolA

Gray et al. (Gray et al. 2020) reported from the jakobid *Andalucia godoyi* a novel famA
DNAP that was predicted mitochondrion-localized and was more closely related to bacterial
sequences than to any known previously characterized eukaryotic DNAP type. Our survey of
famA DNAP in phylogenetically diverse eukaryotes revealed that specific relatives of the *A.*
godoyi DNAP occur not only in other jakobids, but also heteroloboseans and the
phylogenetically unique flagellate *Tsukubamonas globosa*, all of which belong to the
supergroup Discoba. No putative orthologs were identified in the fourth major Discoba
lineage, Euglenozoa, but instead we found them in the two distantly related lineages,
Ancyromonadida (three species) and Malawimonadida (two species). All these DNAP
sequences formed a clade with an UFBP of 100% in the global famA DNAP phylogeny (Fig.
1). Henceforth, we designate the famA DNAP shared among Malawimonadida,
Ancyromonadida, and Discoba as "rdxPolA," where "rdx" is an abbreviation of the Latin word
radix (root). As discussed in the final section, rdxPolA is proposed as the direct descendant
of the PolI in the α-proteobacterial endosymbiont that gave rise to the mitochondrion.
We examined the subcellular localization of five rdxPolA proteins (representing all
three rdxPolA-possessing Discoba lineages plus a single ancyromonad and a single
malawimonad) by expressing their N-terminal regions fused to GFP in the yeast
Saccharomyces cerevisiae. Significantly, the green fluorescence signal appeared to overlap
with mitochondria labeled by MitoTracker Red (Figs. 4A-E), suggesting that the N-termini of
rdxPolA proteins were recognized as MTS by the translocon on the mitochondrial
membranes in yeast. We anticipate that the subcellular localization deduced from the yeast
experiments is applicable to rdxPolA proteins in general. The mitochondrial localization of
rdxPolA is independently supported by results of a previous proteomic study on the
heterolobosean *Naegleria gruberi* (Horváthová et al. 2021), which lists rdxPolA of this
species (XP_002679000.1) as a protein identified in the mitochondrial fraction.

Origins of the novel types of famA DNAP in eukaryotes

The global famA DNAP phylogeny hinted at no direct affinity between any of the eight novel
famA DNAP types and phage homologs (Fig. 1). In this section, we explore the origins of the
newly defined famA DNAP types in detail (summarized in Table 1). To this end, we prepared
eight famA DNAP alignments, each containing one or two novel organellar famA DNAP
types, together with a distinct set of bacterial homologs. The individual DNAP alignments
were then subjected to both ML and Bayesian phylogenetic analyses.
The origin of three famA DNAP types, acPolA, chlnmPolA, or alvPolA, could not be
elucidated with confidence. The monophyly of acPolA recovered with an ML bootstrap
support value (MLBP) of 93% and connected to a clade comprising the PolI sequences
identified in the bacterial species belonging to two phyla of the bacterial superphylum PVC,
Chlamydiae and Lentisphaerae. Nevertheless, the ML bootstrap analysis provided no
sufficient support to judge whether the affinity between acPolA and the PolI sequences in
Chlamydiae and Lentisphaerae is genuine (Fig. S4). The chlnmPolA homologs formed a
clade with the full MLBP, yet the clade showed no clear affinity to any of the bacterial PolI

sequences considered in the phylogenetic analysis (Fig. S5). The phylogenetic analysis
aiming to resolve the origin of alvPolA recovered a clade comprising 20 alvPolA sequences,
three PREX sequences, and PolI sequences of diverse bacteria with an MLBP of 100% (Fig.
S7. The alvPolA sequences formed a clade with an MLBP of 97%, albeit the analysis
recovered no specific affinity between the alvPolA clade and any other bacterial PolI or
PREX sequences. Thus, we leave the precise origin of alvPolA uncertain in this study.
In contrast, as individually discussed below, the origin of the remaining seven novel
famA DNAP types could be narrowed down to specific bacterial groups as the most likely
donors.

Planctomycetes origin of abanPolA

Twenty-seven abanPolA sequences formed a clade with an MLBP of 95% in the DNAP
phylogeny (Fig. S6). The abanPolA clade appeared to be a part of a larger, well-supported
clade enclosing the PolI sequences of bacteria belonging to Planctomycetes and those
belonging to Myxococcales (part of the traditional paraphyletic δ-proteobacteria). Within this
large clade, the abanPolA sequences showed a specific affinity to the PolI sequences of
planctomycetes *Novipirellula artificiosorum* and *Tautonia plasticadhaerens* (Fig. S6).
Altogether, we conclude that the origin of abanPolA is a laterally transferred PolI gene, and
the most plausible candidate for the gene donor is a Planctomycetes bacterium.

Rhodothermales origin of rgPolA

Moriyama et al. (Moriyama et al. 2008) demonstrated the specific phylogenetic affinity
between the rgPolA homolog in *Cyanidioschyzon merolae* and the PolI sequence of
Rhodothermus marinus, a bacterium belonging to the phylum Bacteroidota. We here
confirmed the origin of rgPolA proposed by Moriyama et al. (2008). In the DNAP phylogeny,
the ten rgPolA sequences including the *C. merolae* protein formed a clade with a moderate
MLBP, and the rgPolA clade was then grouped with the clade comprising the PolI
sequences of *Rhodothermus marinus, Rubricoccus marinus* and *Longimonas halophila*, all
of which belong to Rhodothermales in Bacteroidota, with an MLBP of 99% (Fig. 5A). The
most straightforward interpretation of the DNAP phylogeny is that the origin of rgPolA, which
participates in the DNA maintenance in plastids, can be traced back to a PolI gene of a
Rhodothermales(-related) bacterium.

Cyanobacterial origin of chloroPolA

The chloroPolA sequences, a type of famA DNAP found exclusively in the green algal class
Chlorophyceae, and the PolI sequences of cyanobacteria grouped together with an MLBP of
96% (Fig. 5B). In this clade, the cyanobacterial PolI sequences formed a subclade with an
MLBP of 94%, whereas the chloroPolA sequences were paraphyletic (Fig. 5B). We here
propose that chloroPolA emerged through lateral transfer of a cyanobacterial PolI gene to
the common ancestor of Chlorophyceae. In a strict sense, the former proposal demands
nesting the chloroPolA clade within the cyanobacterial PolI sequences and thus is
inconsistent with the tree topology shown in Fig. 5B. Nevertheless, the large difference in the
substitution rate between the cyanobacterial PolI and chloroPolA sequences likely biased
tree reconstruction, hindering the recovery of the genuine relationship between chloroPolA
and cyanobacterial PolI.

Red and glaucophyte algae possess primary plastids that can be traced back to the
"primary endosymbiont," a cyanobacterium taken up by the common ancestor of
Archaeplastida. Nevertheless, the PolI gene, which gave rise to chloroPolA, and that of
primary endosymbiont are most likely different. Despite the potential difficulty in elucidating
the precise position of the rapidly evolving chloroPolA homologs, the PolI of *Gloeomargarita*
lithophora, which has been proposed to be the closest relative of the primary endosymbiont
among the cyanobacteria known to date (Ponce-Toledo et al. 2017), was distantly related to
the chloroPolA in the DNAP phylogeny (green arrow in Fig. 5B). Moreover, chloroPolA
showed a highly restricted distribution within Archaeplastida. Thus, if chloroPolA is assumed
as the direct descendant of the PolI in primary endosymbiont, a potentially large number of
secondary losses of chloroPolA needs to be invoked after the diversification of the extant
members of Archaeplastida. Altogether, the aforementioned hypothesis on the emergence of
chloroPolA by lateral gene transfer (LGT) of a PolI gene from a cyanobacterial donor into
the common Chorophyceae ancestor appears more realistic.

eugPolA and pyramiPolA: γ-proteobacterial origin and endosymbiotic connection

The eugPolA and pyramiPolA sequences, of which a sister relationship was recovered in the
global famA DNAP phylogeny (UFBP of 86%; Fig. 1), were separately subjected to BLAST
surveys against the bacterial sequences and matched mainly with the PolI sequences of
diverse γ-proteobacteria. Thus, a single alignment containing both eugPolA and pyramiPolA
sequences was prepared and subjected to ML phylogenetic analyses. In the resultant ML
tree (Fig. 5C), the eugPolA and pyramiPolA sequences grouped together with an MLBP of
64% and were placed within the radiation of the γ-proteobacterial PolI sequences. The
current plastids in euglenophycean algae descended from a plastid of a green alga
belonging to the order Pyramimonadales via endosymbiosis or kleptoplasty (Turmel et al.
2009; Jackson et al. 2018; Karnkowska et al. 2023). Combining the phylogenetic affinity
between eugPolA and pyramiPolA (Fig. 5C) with the pyramimonadalean origin of
euglenophycean plastids, we here propose that the common ancestor of the extant
photosynthetic euglenophyceans acquired the gene encoding pyramiPolA from the
endosymbiont and then modified it into a plastid DNAP that corresponds to the ancestral
eugPolA. As the γ-proteobacterial PolI sequences enclosed both eugPolA and pyramiPolA
sequences in the DNAP phylogeny (Fig. 5C), we here additionally propose the γ-
proteobacterial PolI origin of pyramiPolA (and indirectly of eugPolA). We are currently
uncertain about the subcellular localization of pyramiPolA (see above). However, this
uncertainty is unlikely to spoil the aforementioned hypothesis, as the ancestor of
photosynthetic euglenophyceans may have changed the subcellular localization of the
endosymbiotically acquired DNAP. It is still possible, though less parsimonious, that the
origin of eugPolA is unrelated to the DNAP of the pyramimonadalean endosymbiont that was
transformed into the plastid. If so, two distinct γ-proteobacteria donated their PolI genes
separately to the common ancestor of the extant photosynthetic euglenophyceans and the
common ancestor of Pyramimonadales.

cryptoPolA and rdxPolA originated from separate α-proteobacterial PolI genes

The cryptoPolA and rdxPolA sequences appeared to share a phylogenetic affinity to the PolI
sequences of α-proteobacteria in BLAST surveys. Hence, we prepared and analyzed a
single alignment including both rdxPolA and cryptoPolA. Significantly, the rdxPolA and

cryptoPolA sequences grouped together with the α-proteobacterial PolI sequences with an
MLBP of 97% by excluding the PolI sequences of other bacteria (Fig. 5D). Within this clade,
the rdxPolA, cryptoPolA, and α-proteobacterial PolI sequences coalesced into three
subclades with full statistical support, and the sister relationship between the cryptoPolA and
α-proteobacterial PolI subclades was recovered with an MLBP of 83% (Fig. 5D). In the
phylogenetic tree of eukaryotes, cryptophytes are included in Cryptista, which further forms
the "CAM" clade with *Microheliella maris* and Archaeplastida (Yazaki et al. 2022), and
distantly related to any of the rdxPolA-bearing lineages. Thus, if we assume that cryptoPolA
and rdxPolA share a common ancestry, such hypothetical famA DNAP may have been lost
secondarily in potentially a large number of the lineages that intervene Cryptophyceae,
Malawimonadida, Ancyromonadida, and Discoba in the tree of eukaryotes. Notably,
cryptoPolA type is present in multiple distantly related members of Cryptophyceae but is also
missing from the Goniomonadea, the immediate non-photosynthetic (plastid-lacking) sister
lineage of Cryptophyceae represented by *Goniomonas* spp. and *Hemiarma marina* (Yazaki
et al. 2022). It thus seems that cryptoPolA was established in the common ancestor of
Cryptophyceae and its emergence coincides with the establishment of the secondary plastid
in the evolution of the Cryptista lineage, in agreement with the plastidial localization of this
DNAP (Fig. 3A). This provides an extra argument for the separate origins of cryptoPolA and
rdxPolA. Thus, although the mutual relationship of cryptoPolA and rdxPolA cannot be settled
with the present data, the DNAP phylogeny can be interpreted as cryptoPolA and rdxPolA
being traced back to distinct α-proteobacterial DNAPs.

Evolution of organellar famA DNAPs in eukaryotes

This study expanded our knowledge of famA DNAPs in eukaryotes. From now on, we will
focus on the evolution of organellar famA DNAPs and thus omit the two newly discovered
putative nuclear DNAPs, alvPolA and abanPolA, from the discussion below. As summarized
in Fig. 6, 16 types of famA DNAP (three types of POP with distinct subcellular localizations
counted as one) have been identified in phylogenetically diverse eukaryotes, with 12 of them
known or predicted to function in organelles. In this section, we combine the updated
diversity of organellar famA DNAPs with the current view on the global eukaryotic phylogeny
to paint an integrated picture of the evolutionary history of organellar DNAPs in eukaryotes,
dividing the discussion according to the major segments of the eukaryote phylogenetic tree.

Mitochondria in Diaphoretickes

Diaphoretickes is a designation of a large-scale taxonomic assemblage ("megagroup")
comprised of Archaeplastida (including Rhodelphidia and Picozoa), Pancryptista, Haptista,
Telonemea, and SAR (Adl et al. 2019), recently shown to embrace also Hemimastigophora
and Provora (Lax et al. 2018; Tice et al. 2021; Tikhonenkov et al. 2022). We detected POP
proteins with predicted mitochondrial localization in representatives of several deeply
diverged Diaphoretickes taxa where the nature of the mitochondrial DNAP had not been
defined before, including Telonemea, Centrohelea, Microhelida, Rhodelphidia, Picozoa, and
Provora (Fig. 6; Table S1). Taken together, all members of Diaphoretickes except
Apicomplexa appeared to utilize POPs as the principal DNAP type for DNA replication/repair
in mitochondria (some plastid-bearing lineages use a single POP that localizes both
mitochondria and plastids; see below for the details) (Fig. 6). Thus, we predict that the
common ancestor of Diaphoretickes used POP for DNA maintenance in mitochondria.

In apicomplexan mitochondria, acPolA, instead of POP (completely missing from the
group), most likely works as the principal DNAP in their mitochondria (Fig. 6). Some of us
previously demonstrated that chrompodellids, the closest relatives of Apicomplexa, possess
a mitochondrion-localized POP (Hirakawa and Watanabe 2019), albeit acPolA was
additionally identified in chrompodellids in this study (Fig. 6). Intriguingly, as mentioned
above our survey detected a mitochondrion-localized POP but provided no evidence for
acPolA in Squirmidea, a lineage sister to Apicomplexa and Chrompodellida combined. We
thus infer that in the mitochondrion of the common ancestor of the clade of Apicomplexa,
Chrompodellida, and Squirmidea (ACS clade; (Mathur et al. 2023)), like other alveolates
including their sister group, Dinozoa, DNA maintenance used to be governed by POP. A
novel type of mitochondrial DNAP, acPolA, was then introduced in the common ancestor of
Apicomplexa and Chrompodellida, with chrompodellids keeping POP along with acPolA.
Nevertheless, the secondary loss of POP having occurred on the branch leading to
Apicomplexa suggests acPolA could fully take over the ancestral role of POP in this group.
We revealed that chlorophycean green algae share a unique type of a putative
mitochondrion-localized famA DNAP, chloroPolA (Fig. 6). It is possible that another unique
mitochondrial famA DNAP, pyramiPolA, was independently introduced to Pyramimonadales,
although the bioinformatic evidence for the specific organellar localization is less clear in this
case (Supplementary Table S1). Although experimental confirmation is required, chloroPolA,
together with POP, is likely involved in DNA maintenance in chlorophycean mitochondria,
and it cannot be ruled out that a similar duality exists in Pyramimonadales. Interestingly, the
chloroPolA gene in *Chlamydomonas reinhardtii* (Cre17.g736150) was found to be expressed
specifically in the zygote, with very limited expression detected in vegetative cells (Joo et al.
2017), pointing to functional specialization of this DNAP.
Two distinct types of DNAP thus appeared to be localized in a mitochondrion in the
mitochondria of some eukaryotes (e.g., POP and acPolA in the chrompodellid mitochondria).
Unfortunately, the sequence data we analyzed here are insufficient to conclude whether the
two co-existed DNAPs are functionally differentiated or redundant in the mitochondrial DNA
maintenance. To address this issue appropriately, the function of each DNAP of interest
needs to be examine experimentally.

Plastids in Diaphoretickes
Diaphoretickes contains multiple plastid-bearing lineages and the repertories of the DNAPs
involved in plastid DNA maintenance vary in lineage-specific manners. Chloroplastids,
ochrophytes (plastid-bearing species in Stramenopiles), and haptophytes all seem to
operate "dual-target POPs," which can manage DNA maintenance in both mitochondria and
plastids (Kimura et al. 2002; Christensen et al. 2005; Mori et al. 2005; Ono et al. 2007;
Parent et al. 2011; Hirakawa and Watanabe 2019) (Fig. 6). The red alga *Cyanidioschyzon*
merolae was also demonstrated to have a dual-targeted POP, yet supplemented with a
plastid-localized rgPolA (Moriyama et al. 2008). Our study has expanded the phylogenetic
distribution of the pair of the (presumably) dual-targeted POP and the plastid-localized
rgPolA from a single red alga to multiple red algae as well as glaucophytes (Fig. 6). The
organismal phylogenies inferred from nucleus-encoded proteins constantly united
Chloroplastida and Glaucophyta (Tice et al. 2021; Irisarri et al. 2022; Yazaki et al. 2022),
while those inferred from plastid proteins often recovered the sister relationship between
Chloroplastida and Rhodophyta (Rodríguez-Ezpeleta et al. 2005; Mackiewicz and Gagat
2014; Figueroa-Martinez et al. 2019). Regardless of the two competing scenarios for the

phylogeny of Archaeplastida, the common ancestor of this group may have possessed both
dual-targeted POP and plastid-localized rgPolA. The latter was lost not only in Rhodelphidia
(retaining a genome-less plastid (Gawryluk et al. 2019)) and Picozoa (that lost the plastid
completely (Schön et al. 2021)) but also Chloroplastida. In the case of both POP and rgPolA
being localized in a plastid, we need experimental data to clarify the precise functions of the
two DNAPs in the plastid DNA maintenance.
The situation concerning plastid DNAPs is specific also in Myzozoa, the alveolate
clade combining Dinozoa and ACS clade (Fig. 6). Those dinozoans that have retained a
plastid genome (i.e., "core" dinoflagellates) employ plastid-targeted POPs that are distinct
from mitochondrion-targeted POPs (Hirakawa and Watanabe 2019). In contrast, the
members of ACS clade, i.e., apicomplexans, chromopodellids, and squirmids, exhibit PREX
for DNA maintenance in their plastids (Hirakawa and Watanabe 2019; this study), whereas
POPs in chromopodellids and squirmids were predicted to be mitochondrion-localized
bioinformatically (Fig. 6). At this moment, we are uncertain what type (or types) of DNAP
worked in the plastid of the ancestral myzozoan. If the ancestral myzozoan used a single
type of a plastid DNAP, either a switch from POP to PREX or that from PREX to POP
occurred after the separation of Dinozoa and ACS clade. Alternatively, the ancestral
myzozoan may have had both POP and PREX for plastid DNA maintenance. In the latter
scenario, differential losses of one of the two types of DNAP need to be invoked on the
branches leading to Dinozoa and ACS clade.
Chlorarachniophytes operate distinct POPs localized in the mitochondrion and the
plastid (Hirakawa and Watanabe 2019). In addition, we identified an additional famA DNAP,
chlnmPolA, which most likely participates in DNA maintenance in the nucleomorph. Besides
chlnmPolA, the chlorarachniophyte *Bigelowiella natans* was reported to use an unrelated
(family B) DNAP of viral origins for the replication of the nucleomorph genome, in addition to
a homolog of DNA polymerase eta (family Y DNAP) presumably involved in nucleomorph
genome repair (Suzuki et al. 2016). The ancestral chlorarachniophyte may thus have
modified the original green alga-derived machinery for DNA maintenance in the
nucleomorph by integrating components with exogenous origins.
The members of another algal nucleomorph-bearing lineage, Cryptophyceae,
possess a similar repertory of famA DNAPs to that in chlorarachniophytes. *G. theta* as the
reference cryptophyte species specifically has two POPs plus a lineage-specific famA DNAP
(cryptoPolA) (Fig. 6). Considering the fluorescent protein tagging in diatom cells, *G. theta*
likely uses one of the two types of POP (i.e., POP1) and the lineage-specific famA DNAP
(i.e., cryptoPolA) for the DNA maintenance in the mitochondrion and plastid, respectively.
Interestingly, the nucleomorph is the best estimate for the subcellular localization of *G. theta*
POP2 (see above). Notably, except for the closely related species *Hanusia phi*, our
investigation of the genome and/or transcriptome assemblies available for other cryptophyte
species did not indicate the presence of the second POP paralog as in *G. theta*. At face
value, the restricted distribution of POP2 among cryptophyte algae suggests the recruitment
of a POP polymerase for a nucleomorph function only after the divergence of this algal
group. However, an alternative interpretation exists: the POP2 transcripts may have been
overlooked in at least some of the other cryptophytes. In general, eukaryotic genomes are
replicated by family B DNA polymerases (Guilliam and Yeeles 2020), and the cryptophyte
nucleomorph genome effectively is a reduced and divergent eukaryotic (specifically red
algal) genome. Thus, we speculate that analogously to the presumed role of chlnmPolA in
the chlorarachniophyte nucleomorph, the *G. theta* POP2 protein plays an auxiliary role in
maintaining the nucleomorph genome in this species (e.g., DNA repair).

Amorphea plus CRuMs

A large taxonomic assemblage, Amorphea, comprises Opisthokonta, Apusomonadida, Breviatea, and Amoebozoa. Recent phylogenomic studies united collodictyonids, rigifilids, and mantamonads into a clade called CRuMs with high statistical support and further nominated this assemblage as a candidate for the sister branch of Amorphea (Brown et al. 2018). A putative mitochondrion-localized POP has been previously detected in Apusomonadida and Amoebozoa (Hirakawa and Watanabe 2019), and we now found the same for CRuMs, whereas opisthokonts have been known to use Polγ for DNA maintenance in their mitochondria (Fig. 6). Breviates possess severely modified mitochondria (mitochondrion-related organelles) adapted to the microaerophilic/anaerobic lifestyle and all probably lack a genome (confirmed for the most extensively studied representative *Pygsuia biforma*; (Stairs et al. 2014)), so it is no surprise that no mitochondrial DNAP candidate was found in this group. By mapping the types of mitochondrial DNAP on the tree of Amorphea, we propose that the common ancestor of this taxonomic assemblage possessed a mitochondrion-localized POP but secondary losses of POP occurred in the two descendant lineages, Opisthokonta and Breviatea. The ancestral opisthokont had discarded the pre-existing POP after the emergence of a lineage-specific mitochondrial DNAP, Polγ. Breviates simply lost the POP due to no use for any mitochondrion-localized DNAP. If CRuMs and Amorphea are truly the closest relatives among eukaryotes, their common ancestor had likely used POP for mitochondrial DNA maintenance.

Ancyromonadida, Discoba, Malawimonadida, and LECA

Prior to this study, the information on mitochondrial DNAPs in Discoba has been available only for euglenozoans and a single jakobid, *Andalucia godoyi*. In euglenozoans, three types of mitochondrion-localized DNAP, namely POP, PolIA, and PolIBCD+, have been identified (Klingbeil et al. 2002; Harada et al. 2020; Harada and Inagaki 2021), and *A. godoyi* was proposed to use a novel type of famA DNAP for the mitochondrion (Gray et al. 2020). The current study revealed that all the discobids except euglenozoans possess rdxPolA, typified by the *A. godoyi* protein, for mitochondrial DNA maintenance. In the phylogenetic tree of Discoba, jakobids and then *Tsukubamonas globosa* were sequentially separated prior to the split of heteroloboseans and euglenozoans. Thus, we regard that rdxPolA was present prior to the diversification of Discoba as the mitochondrial DNAP and a switch from rdxPolA to POP or PolIA/PolIBCD+ occurred on the branch leading to euglenozoans. The introduction of eugPolA in the ancestor of photosynthetic members of Euglenophyceae should have coincided with the plastid acquisition through green algal endosymbiosis (see above).

Significantly, the origin of rdxPolA may be much earlier than the emergence and diversification of Discoba, as this DNAP is shared by ancyromonads and malawimonads. To elucidate the origin and evolution of rdxPolA, understanding the phylogenetic relationship among the three rdxPolA-bearing lineages is crucial. The situation would be simplest if Malawimonadida, Ancyromonadida, and Discoba constituted a monophyletic group among eukaryotes. Intriguingly, all three taxa do share some morphological/ultrastructural features. Malawimonadida and Discoba include species with typical excavate ultrastructures, such as a ventral suspension feeding groove lined by multi-microtubular roots associated with basal bodies (Simpson 2003). Although ancyromonads lack excavate-like feeding grooves and have not been regarded as typical excavates, some features in the flagellar apparatus of typical excavates were identified in the ancyromonad *Ancyromonas sigmoides* (Heiss et al. 2011). A set of ultrastructural traits specific to typical excavates is also found in many

metamonads (particularly free-living members; (Simpson 2003)), but all metamonads known
to date have severely reduced, genome-lacking versions of mitochondria as the result of
adopting microaerophilic/anaerobic environments (Leger et al. 2017; Stairs et al. 2021) and
hence lack any mitochondrial DNAP due to secondary loss. Combined, the monophyly of the
rdxPolA-bearing lineages (with/without Metamonada) is arguable from the
morphological/ultrastructural viewpoint and would nominate rdxPolA as a candidate
synapomorphy for this hypothetical clade.
In contrast to the morphological traits shared among the rdxPolA-bearing lineages
plus Metamonada, no phylogenomic study has ever recovered their monophyly (e.g., (Tice
et al. 2021; Tikhonenkov et al. 2022; Yazaki et al. 2022). We examined the abovementioned
issue by analyzing a phylogenomic alignment comprising 340 proteins with no outgroup
taxon (Figs. 7A & 7B). In the strict sense, the unrooted phylogenetic analyses presented
here are inappropriate to assess whether the rdxPolA-bearing lineages are genuinely
monophyletic. However, the lineages of our interest need to form a clan (i.e., be at least
potentially monophyletic in the unrooted phylogeny) if they are to form a genuine clade of
eukaryotes. Another point to be noted is that all the taxa included in the alignment belong to
Metamonada or lineages primarily using either POP or rdxPolA as mitochondrial DNAPs.
Thus, the monophyly of the rdxPolA-bearing lineages is synonymous with the monophyly of
POP-bearing assemblages (i.e., Diaphoretickes and Amorphea plus CRuMs), with
Metamonada (with an unknown ancestral state as of the type of the mitochondrial DNAP)
allowed to be placed anywhere in the tree. As anticipated from the previously published
works, the resultant unrooted tree of eukaryotes failed to recover the monophyly of the
rdxPolA-bearing lineages (with or without Metamonada) or that of POP-bearing
assemblages. In the ML tree, the branch connecting Diaphoretickes and Amorphea plus
CRuMs was interrupted by two distinct branches, one uniting Discoba and Metamonada and
the other uniting Malawimonadida and Ancyromonadida (Fig. 7A). We subjected the ML tree
and six alternative trees to an approximately unbiased (AU) test (Shimodaira 2002).
Significantly, the AU test rejected one of the alternative trees, in which Diaphoretickes and
Amorphea plus CRuMs were forced to be monophyletic, at a 1% α-level (Tree 7; Fig. 7B).
Thus, we disfavor the monophyly of the rdxPolA-bearing lineages regardless of the
inclusion/exclusion of Metamonada. As discussed below, this conclusion has important
implications for reconstructing details of mitochondrial DNA maintenance in the earliest
phases of eukaryote evolution.

Perspective for the mitochondrial DNAP in pre-LECA

It is widely accepted that mitochondrial endosymbiosis was indispensable for
eukaryogenesis and thus an essentially modern mitochondrion containing a reduced
genome was most likely present in the common ancestor of all extant eukaryotes (the last
eukaryotic common ancestor or LECA). Considering the α-proteobacterial origin of
mitochondria, it is rational to assume that the DNA maintenance had been carried out by the
α-proteobacterial DNAPs including PolI in the proto-mitochondrion (a primitive
mitochondrion in the pre-LECA stem lineage). Is it then possible that rdxPolA is a direct
descendant of the putative proto-mitochondrial PolI? The most recent studies suggested
that the proto-mitochondrion represented a lineage phylogenetically sister to essentially all
the α-proteobacteria known to science so far (Martijn et al. 2022; Muñoz-Gómez et al. 2022).
Intriguingly, our analysis found the intimate affinity of the rdxPolA sequences to, yet outside,
the radiation of α-proteobacterial PolI sequences (Fig. 5D), being compatible with the

phylogenetic position of the putative bacterial lineage from which the proto-mitochondrion
emerged. In the first scenario, multiple secondary losses of this DNAP must have occurred
after the divergence of major branches in the tree of eukaryotes and POP supplemented the
secondary loss of rdxPolA. Alternatively, the sporadic distribution of rdxPolA in eukaryotes
can be explained by lateral gene transfer. This second scenario was not positively supported
by the rdxPolA phylogeny, in which the relationship among the discobid, malawimonad, and
ancyromonad sequences remains unresolved (Fig. 5D). Significantly, regardless of how the
phylogenetic distribution of rdxPolA has been shaped, the proto-mitochondrion may be the
plausible candidate for the origin of rdxPolA. The rdxPolA-containing lineages have been
predicted to be early branches of eukaryotes [e.g., Derelle and Lang (2012); see below].
Thus, the proto-mitochondrial origin of rdxPolA is well inclusive to both scenarios. Overall,
we prefer the scenario assuming the vertical inheritance of rdxPolA rather than the other
involving eukaryote-to-eukaryote LGT but the two scenarios should be reevaluated when an
additional rdxPolA-containing lineage is found in the future.
To examine whether rdxPolA is derived from the proto-mitochondrion in the LECA, it
would be useful to know the type of mitochondrion-localized DNAP in the earliest eukaryotes
including LECA, which can in principle be deduced historical reconstructions guided by the
rooted eukaryote phylogeny. Unfortunately, most studies so far carried out to explore the
eukaryotic root by using a non-eukaryotic outgroup are by itself not sufficient to infer the
early evolution of mitochondrion-localized DNAP, as they included only Discoba of all the
three rdxPolA-containing lineages (Derelle and Lang 2012; He et al. 2014; Al Jewari and
Baldauf 2023a; Al Jewari and Baldauf 2023b). However, when the root positions suggested
by most of these analyses are mapped onto taxonomically comprehensive unrooted
eukaryote phylogenies as recently inferred from multigene datasets (Lax et al. 2018; Tice et
al. 2021; Tikhonenkov et al. 2022), the rdxPolA-containing taxa are found in both principal
clades defined by the root, with Discoba in one of them the Malawimonadida plus
Ancyromonadida in the other. This is consistent with the outgroup-based taxon-rich rooting
analysis conducted by Derelle et al. (2015), which splits Discoba and Malawimonadida into
the two major clades separated by the inferred root (ancyromonads were missing from the
analysis). On the other hand, the analysis by Cerón-Romero et al. (2022), who took a gene
tree–species tree reconciliation approach (taking into account gene duplications and losses)
to infer the eukaryotic root, does not support rdxPolA as a LECA-resident protein. However,
the results reported by these authors need to be interpreted with caution, given the fact that
the root position most favoured by their analysis was inside opisthokonts (between fungi and
all other eukaryotes), which is at odds with solid evidence for opisthokont monophyly and
would require major reconsideration of early eukaryote evolution.
Although the position of the eukaryotic root has yet to be resolved with confidence,
the distribution of rdxPolA mapped on the current tree of eukaryotes certainly demands
multiple secondary losses of this DNAP. Interestingly, as proposed by previous studies
(Moriyama et al. 2011; Hirakawa and Watanabe 2019; Harada and Inagaki 2021) and further
reinforced by our expanded sampling, the phylogenetic distribution of POP in eukaryotes
implies that the LECA also possessed POP as another mitochondrion-localized DNAP, with
the POP absence in the current rdxPolA-containing lineages, metamonads, opisthokonts,
and other POP-lacking taxa resulting from secondary loss. Altogether, in light of the results
from our comprehensive survey of famA DNAPs in eukaryotes, we propose that both
rdxPolA and POP were established in the pre-LECA stage. Such ancestral "redundancy" of
the original (i.e., α-proteobacterial) and a newly acquired mitochondrial component, removed
by subsequent differential loss of one or the other, would not be without a precedent. An

interesting similar example is the history of mitochondrial RNA polymerases (RNAPs), with
jakobids possessing the ancestral multisubunit RNAP (encoded by the mitochondrial
genome) and all other eukaryotes utilizing a novel nucleus-encode phage-derived single-
subunit RNAP (Roger et al. 2017; Gray et al. 2020). Here the co-existence of both RNAPs in
the LECA followed by their differential loss also seems to best fit their modern distribution
and the phylogenetic position of jakobids among eukaryotes.
712

713 Conclusions

Our analyses uncovered a series of previously unnoticed types of eukaryotic family A DNA
polymerases that each occur in multiple representatives of broader protist taxa and trace
their origin to evolutionary events at least hundreds of millions of years ago. Two of these
DNAP types, not analysed in detail here (alvPolA and abanPolA), may be novel components
of the nuclear machinery of DNA maintenance and deserve specific attention in a separate
future study. The remaining eight newly defined DNAP types fundamentally enrich our
understanding of the organellar biology. The origin of four of them appear to be directly
linked to the emergence of the primary (rgPolA) or different types of secondary (chlnmPolA,
eugPolA, cryptoPolA) plastids and their identification adds so far missing pieces into the
mosaic of plastid endosymbioses. Possibly most interesting is the discovery of rdxPolA,
which might represent a previously unknown ancestral (α-proteobacterial) mitochondrial
component established pre-LECA and retained by only a minor fraction of extant eukaryote
lineages. Further growth of sequence data from eukaryotes, prokaryotes as well as viruses
will certainly help to refine our current interpretation of the origin and evolutionary history of
the different DNAPs and will resolve open questions such as the nature of the mitochondrial
DNAP in Hemimastigophora. Functional genomics approaches need to be employed to more
precisely define the role of the functionally uncharacterized DNAPs and to resolve why some
mitochondria and plastids seem to possess two different DNAP types. A salient unanswered
question is whether organellar genomes may depend on DNAPs from families other than the
family A. Except for family B DNAPs serving in the chlorarachniophyte nucleomorph, no
such cases have been described, but a systematic study like we have carried out here for
the family A is lacking. Clearly, a lot is yet to be learned about how mitochondria and plastids
deal with their DNA replication and repair at the paneukaryotic scale.
737

738 Materials and Methods

739 Cell cultures and RNA-seq analyses of *Tsukubamonas globosa* and
740 *Fabomonas tropica*

Tsukubamonas globosa NIES-1390 was purchased from Microbial Culture Collection at the
National Institute for Environmental Studies (https://mcc.nies.go.jp/). *Fabomonas tropica*
SRT902 was isolated from seawater sampled at Enoshima Aquarium (Enoshima,
Kanagawa, Japan) on October 30th, 2019. The species identification of the strain was
supported by its 18S rRNA gene being 99% identical to the the respective gene sequence
(JQ340336.1) of the type strain *F. tropica* nyk4. The culture of *T. globosa* NIES-1390 was
maintained in the laboratory in the 1,000× diluted LB medium containing soil extract and a
vitamin mix at 20 °C. The culture of *F. tropica* SRT902 was maintained in the laboratory in
seawater containing soil extract and vitamin mix at 20 °C. We harvested the cells from the

laboratory cultures of *T. globosa* and then extracted total RNA using the TRIzol reagent. The
T. globosa RNA sample was subjected to a cDNA library construction using NEBNext Ultra II
Directional RNA Library Prep Kit and then to RNA-seq analysis by using the Illumina HiSeq
platform at a biotech company (AZENTA Japan Corp., Tokyo, Japan). The experimental
procedures for the RNA-seq analysis of *F. tropica* were principally the same, albeit SMART-
Seq v4 Ultra Low Input RNA Kit, Nextera DNA Flex Library Prep Kit, and the Illumina HiSeq
X platform were applied for cDNA synthesis, cDNA library construction, and sequencing,
respectively (the sequencing has been done by Macrogen Japan Corp., Tokyo, Japan). We
obtained 64,305,977 and 204,000,750 paired-end reads from the RNA-seq analyses of *T.*
globosa and *F. tropica*, respectively. The raw sequence reads were trimmed with fastp
v0.23.2 (Chen et al. 2018) with the -q 20 -u 80 option and then assembled with Trinity
v2.13.2 (Grabherr et al. 2011).

Phylogeny of famA DNAPs identified in bacteria, phages, and eukaryotes

We searched for novel famA DNAPs using HMMER searches (Eddy 2011) against EukProt3
database (Richter et al. 2022) with a profile HMM generated from the seed alignment of the
Pfam family DNA_Pol_A (pfam00476). Significant hits (up to the inclusion threshold) were
gathered and preliminary phylogenetic analyses were carried out to identify potentially novel
eukaryotic DNAP types. Additional databases were then searched with blastp or tblastn
(Camacho et al. 2009) to identify additional members of the putative new types, including the
resources provided by GenBank (Sayers et al. 2021), JGI genome portal
(https://genome.jgi.doe.gov/portal/), the MMETSP (Keeling et al. 2014) and OneKP
transcriptome sequencing projects (Leebens-Mack et al. 2019), and individual sequence
datasets deposited in Dryad as supplemental data to publications reporting on the
generation of genome or transcriptome assemblies (further details on the sequence sources
are provided in Table S1). In addition, we searched the transcriptome assmeblies from *T.*
globosa and *F. tropica* generated as part of this study, and we also obtained a rdxPolA
sequence from a transcriptome assembly from the jakobid *Ophirina amphinema* (the whole
dataset to be published elsewhere) (Yabuki et al. 2018). The famA DNAP sequences
potentially contaminated from non-eukaryotic sources and those belonging to the previously
described eukaryotic famA DNAP types, namely Polγ, POP, Polθ, PolIA, PolIBCD+, and
PREX, were discarded based on reciprocal blast analyses and preliminary phylogenetic
analyses. The sequences used for further analyses were manually curated to identify and fix
possible mistakes in the respective gene models and to evaluate their completeness
(especially concerning the N-terminus critical for correct prediction of subcellular
localization). In several cases a complete or at least extended sequence was obtained by
manually joining partially assembled sequence fragments. We eventually retained 134
sequences as the candidates for previously undescribed famA DNAPs, which are listed with
relevant details in Table S1.
For phylogenetic analyses we additionally retrieved a set of reference sequences
representing the diversity of PolI in bacteria and phages from the refseq_select_prot
database in NCBI as of March 5, 2021 [the detailed procedure was the same as described in
Harada and Inagaki (2021)]. The representative sequences of Polγ, POP, Polθ, PolIA,
PolIBCD+, and PREX were selected from the alignment generated by some of us before
(Harada and Inagaki 2021). The 134 eukaryotic sequences of the candidates for novel famA

DNAPs, the PolII sequences of bacteria and phages, and the sequences of the previously
known famA DNAP in eukaryotes were aligned by MAFFT v7.490 with the L-INS-i model
(Katoh and Standley 2013). The aa sequences encoding the polymerase domain is the only
part of the protein that is shared among all the sequences analyzed. Further, ambiguously
aligned positions were excluded manually and gap-containing positions were trimmed by
trimAl v1.4 (Capella-Gutiérrez et al. 2009) from the initial alignment, leaving 355
unambiguously aligned aa positions in 488 famA DNAPs ("global DNAP" alignment). We
then subjected the global DNAP alignment to the ML tree reconstruction by using IQ-TREE
v.2.2.0 with the LG+C60+F+R10 model (Minh et al. 2020) that was selected by ModelFinder
with "-mset LG+C60" option (Kalyaanamoorthy et al. 2017). Statistical support for each
bipartition in the ML tree was calculated by 1000-replicate an ultrafast bootstrap
approximation (Hoang et al. 2018).

Phylogenetic analyses assessing the origins of novel types of famA DNAPs in eukaryotes

The origins of the ten novel types of famA DNAPs in eukaryotes—acPolA, rgPolA,
chlnmPolA, eugPolA, pyramiPolA, chloroPolA, rdxPolA, cryptoPolA, abanPolA, and
alvPolA—were assessed by separate phylogenetic analyses. Six of the DNAPs (acPolA,
rgPolA, chlnmPolA, chloroPolA, abanPolA, and alvPolA) were each analyzed as part of a
separate alignment, whereas the other four were combined in pairs (eugPolA plus
pyramiPolA, rdxPolA plus cryptoPolA) into two separate alignments based on the fact the
DNAPs in pairs were closely related in the global DNAP phylogeny. Each of the eight
alignments was supplemented by a set of bacterial sequences selected by targeted
searches to include the putative closest relatives of the eukaryotic DNAPs analyzed.
Specifically, the bacterial PolI sequences were retrieved by running a BLASTP search
against the GenBank Refseq_selec_prot database with the eukaryotic famA DNAP
sequences of interest as queries. We kept approximately the top 1,000 bacterial PolI
sequences matched to the queries in each of the BLAST surveys. The redundancy in the set
of bacterial PolI sequences retrieved from the database was removed by using CD-HIT (Li
and Godzik 2006). The E-value cutoffs for the BLAST surveys and parameters for CD-HIT
are listed in Table S2. The query sequences (one or two types of the novel famA DNAPs in
eukaryotes) and the bacterial PolI sequences retrieved from the corresponding BLAST
survey were aligned by using MAFFT with the L-INS-i model. We then polished the eight
alignments by manual trimming of ambiguously aligned positions and exclusion of gap-
containing positions by trimAl. As a result, the alignments built for acPolA, chlnmPolA,
chloroPolA, abanPolA, and alvPolA plus the respective sets of bacterial PolI sequences
were comprised of unambiguously aligned positions in the polymerase domain only,
whereas the final alignments made for rgPolA, eugPolA/pyramiPolA, and rdxPolA/cryptoPolA
plus their bacterial homologs were longer for including also unambiguously aligned positions
in the 5′-3′ exonuclease and the 3′-5′ exonuclease domains. The eight alignments were
subjected to ML tree search and 100-replicate nonparametric ML bootstrap analysis by using
IQ-TREE. The substitution models applied for the tree searches were selected by
ModelFinder. For the nonparametric ML bootstrap analysis, we applied the same model as
the corresponding tree search but incorporated PMSF (posterior mean site frequencies) that
were calculated over the ML tree as the guide (Wang et al. 2018). The four alignments for
rgPolA, eugPolA/pyramiPolA, rdxPolA/crytoPolA, and chloroPolA were also subjected to

Bayesian analysis with the CAT+GTR+Γ model using PhyloBayes v1.8 (Lartillot et al. 2013).
For each alignment, two Markov chain Monte Carlo runs were run to calculate the
consensus tree and Bayesian posterior probabilities. Details on the eight alignments (i.e., the
numbers of the DNAP sequences and positions used for the phylogenetic analyses) and the
substitution models selected for the phylogenetic analyses are provided in Table S2.

In silico prediction of subcellular localization and functional domains of
famA DNAPs

Seventy-five out of the 134 sequences representing the novel famA DNAP types identified in
this study were considered to have complete and accurately defined N-termini. We subjected
their N-terminal regions to *in silico* prediction of the targeting potential to subcellular
compartments (i.e., endoplasmic reticulum, mitochondria, and plastids) using a suite of tool,
namely MitoFates (Fukasawa et al. 2015), NommPred (Kume et al. 2018), DeepLoc-1.0
(Almagro Armenteros et al. 2017), TargetP-1.1 (Emanuelsson et al. 2000), PredSL
(Petsalaki et al. 2006), and TMHMM-2.0 (Krogh et al. 2001). Additionally, we subjected the
full-length pyramiPolA and chloroPolA sequences to PredAlgo (Tardif et al. 2012), which was
developed for predicting the subcellular localization of green algal proteins. Conserved
functional domains in the 134 famA DNAPs were predicted using InterProScan v5.55 with
the Pfam database (Jones et al. 2014; Paysan-Lafosse et al. 2023).

Experiments assessing the subcellular localizations of the selected famA
DNAPs

rdxPolAs

The N-terminal amino acid sequences predicted as mitochondrial targeting signals were
fused to green fluorescent protein (GFP), and the fusion proteins were expressed in yeast
(*Saccharomyces cerevisiae*). In the five rdxPolA sequences of *Ophirina amphinema*,
Gefionella okellyi, *Naegleria gruberi*, *F. tropica*, and *T. globosa*, the first 44, 26, 95, 71, and
76 amino acid residues were predicted as MTS by MitoFates, respectively. Double-stranded
DNA (dsDNA) fragments corresponding to regions encoding the putative MTS aa sequences
of the five rdxPolA proteins were synthesized with two modifications described below. First,
the codon usage, which may be specific to the five eukaryotic genomes, was optimized for
expression in yeast. Second, the synthesized nucleotide sequences were flanked with the
sequence recognized by *Eco*RI (5′-GAATTC-3′) and that by *Bam*HI (5′-GGATCC-3′) at the 5′
and 3′ termini, respectively. The synthesized dsDNA fragments were inserted into pYX142-
mtGFP vector (Westermann and Neupert 2000) via the restriction sites mentioned above.
The dsDNA synthesis and vector construction were done by a biotech company (AZENTA
Japan Corp, Japan). The vectors were transformed in the BY4742 strain of yeast
(Brachmann et al. 1998). Yeast transformations were performed according to the lithium
acetate procedure (Schiestl and Gietz 1989), and transformants were selected on a
synthetic dextrose minimal medium lacking lysine, histidine, and uracil. Mitochondria in yeast
transformants were stained with MitoTracker Red at a final concentration of 0.5 μM and were
observed with an Olympus BX51 fluorescent microscope (Olympus) equipped with an
ORCA-3CCD color camera (Hamamatsu Photonics).

Bigelowiella natans chlnmPolA

To confirm the subcellular localization of the *B. natans* chlnmPolA, a GFP-tagged protein was heterologously expressed in the chlorarachniophyte *Amorphochlora amoebiformis*. The *B. natans* chlnmPolA was predicted to have an N-terminal extension containing a signal peptide (the first 24 amino acid residues) by TargetP 2.0 (Almagro Armenteros et al. 2019). We constructed a plasmid vector expressing GFP fused with the first 100 aa residues of *B. natans* chlnmPolA. Firstly, total RNA was extracted from *B. natans* CCMP621 cells using TRIzol reagent (Invitrogen), and cDNA was synthesized using SuperScript IV Reverse Transcriptase (Invitrogen) with an oligo(dT) primer. Secondly, the cDNA fragment encoding chlnmPolA was amplified by PCR with a pair of specific primers (5′-ATGATCAGGAAGCAATATATGTTGAG-3′ and 5′-CACCAGAGAGTTAGATCCCAT-3′), and inserted at the 5' of the GFP gene in the plasmid vector pLaRGfp (Hirakawa et al. 2009) using GeneArt Seamless Cloning and Assembly Enzyme Mix (Invitrogen). The plasmid was introduced into *Escherichia coli* DH5α, and the inserted sequence was verified by Sanger sequencing. To express the GFP fusion protein, *A. amoebiformis* CCMP2058 cells were transformed with the plasmid using a Gene Pulser Xcell electroporation system (Bio-Rad), as described previously (Fukuda et al. 2020). One day after transformation, GFP fluorescence was observed under an inverted Zeiss LSM 510 laser scanning microscope (Carl Zeiss).

Guillardia theta cryptoPolA and POPs

To confirm the subcellular localization of the *G. theta* cryptoPolA and the two POPs, the N-terminal amino acid sequences of the three DNAPs fused to GFP were heterologously expressed in the diatom *Phaeodactylum tricornutum* UTEX642. For the three DNAPs, dsDNA fragments of the region from the start codon until the beginning of the first functional domain were synthesized by (AZENTA Japan Corp, Japan). Each of the synthesized DNA fragments was amplified by PCR and inserted at the 5' of the GFP gene in the pPha-NR vector (NovoPro Bioscience) using NEBuilder HiFi DNA Assembly (New England BioLabs). The pPha-NR constructs were introduced into *P. tricornutum* using a NEPA21 gene gun (NEPAGENE) (Miyahara et al. 2013). Actively growing *P. tricornutum* transformants in a zeocin-based selection medium were observed with an Olympus BX51 fluorescent microscope (Olympus) equipped with an Olympus DP72 CCD color camera (Olympus). Mitochondria of transformants transfected with fragments of POP1 were stained with MitoTracker Orange at a final concentration of 0.1 µM.

Phylogenomic analysis of eukaryotes

We updated the 351-gene phylogenomic alignment used for Yazaki et al. (2022). Each of the 351 single-gene alignments was updated by adding the homologous sequences of *T. globosa* and *F. tropica* SRT902 that were retrieved from the RNA-seq data, followed by separate inspections by preliminary ML phylogenetic analyses. Among the 351 alignments, we selected 340 single-gene alignments for the final ML phylogenetic inferences. The 340 updated alignments were concatenated into a single "phylogenomic" alignment comprising 97 taxa and 107,862 unambiguously aligned amino acid positions. We analyzed the 340-gene phylogenomic alignment with the ML method by using IQ-TREE (Minh et al. 2020) with the LG+C60+F+Γ model. The statistical support for each bipartition was calculated by 100-replicate nonparametric ML bootstrap analysis. For the bootstrap analysis, we applied the

LG+C60+F+Γ+PMSF model (the ML tree inferred with the LG+C60+F+Γ model was used as
the guide).
We tested the possibility that the rdxPolA-bearing lineages form a monophylum
(synonymous with the monophyly of the POP-bearing assemblages) using an AU test
(Shimodaira 2002). Six alternative trees were generated from the ML tree (see above) by
pruning and regrafting the Amorphea plus CRuMs clade from the original position to six
alternative positions (Fig. 7B). We calculated the site-wise log-likelihoods (site-lnLs) over the
ML and six alternative trees (the substitution model used for the ML tree search was applied).
The resultant site-lnL data were then subjected to an AU test. IQ-TREE (Minh et al. 2020)
was used for all the calculations required for the test described above.

Data availability

936 All the alignments, treefiles inferred from the alignments, and the fasta files containing
937 transcriptome assemblies generated in this study are available at
938 https://doi.org/10.5061/dryad.b5mkkwhjz. The transcriptome data of *T. globose* NIES-1390
939 and *F. tropica* SRT902 were deposited in GenBank/EMBL/DDBJ Sequence Read Archive
940 under accession no. PRJNA932580.

Acknowledgments

We would like to thank the Enoshima Aquarium (Kanagawa, Japan) for providing the
seawater samples and Dr. Takashi Shiratori (University of Tsukuba) for establishing the *F.*
tropica strain. The work was supported by the Japan Society for Promotion of Sciences
projects 18KK0203, 19H03280, 23H02535, and BPI05044 (to Y.I.), 22J11104 (to R.H.),
19H03274 (to R.K.), and the Czech Science Foundation grant no. 21-19664S (to M.E.). *T.*
globosa (NIES-1390) and *S. cerevisiae* BY4742 strain was provided by the NIES and YGRC
through the National Bio-Resource Project (NBRP) of the Ministry of Education, Culture,
Sports, Science and Technology (MEXT), Japan, respectively. Computations were partially
performed on the NIG supercomputer at ROIS National Institute of Genetics. The
phylogenetic analyses conducted in this work have been carried out partially under the
'Interdisciplinary Computational Science Program' in the Center for Computational Sciences,
University of Tsukuba.

Literature Cited

Adl SM, Bass D, Lane CE, Lukeš J, Schoch CL, Smirnov A, Agatha S, Berney C, Brown MW,
Burki F, et al. 2019. Revisions to the classification, nomenclature, and diversity of
eukaryotes. *J. Eukaryot. Microbiol.* 66:4–119.

Al Jewari C, Baldauf Sandra L. 2023. Conflict over the eukaryote root resides in strong
outliers, mosaics and missing data sensitivity of site-specific (CAT) mixture models.
Syst. Biol. 72:1–16.

Al Jewari C, Baldauf Sandra L. 2023. An excavate root for the eukaryote tree of life. *Sci. Adv.*
9:eade4973.

Almagro Armenteros JJ, Salvatore M, Emanuelsson O, Winther O, Heijne G von, Elofsson A,
Nielsen H. 2019. Detecting sequence signals in targeting peptides using deep
learning. *Life Sci. Alliance* 2:e201900429.

Almagro Armenteros JJ, Sønderby CK, Sønderby SK, Nielsen H, Winther O. 2017.
DeepLoc: prediction of protein subcellular localization using deep learning.
Bioinformatics 33:3387–3395.

Brachmann BC, Davies A, Cost GJ, Caputo E, Li J, Hieter P, Boeke JD. 1998. Designer
deletion strains derived from Saccharomyces cerevisiae S288C: A useful set of
strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast*
14:115–132.

Brown MW, Heiss AA, Kamikawa R, Inagaki Y, Yabuki A, Tice AK, Shiratori T, Ishida K-I,
Hashimoto T, Simpson AGB, et al. 2018. Phylogenomics places orphan protistan
lineages in a novel eukaryotic super-group. *Genome Biol. Evol.* 10:427–433.

Burger G, Saint-Louis D, Gray MW, Lang BF. 1999. Complete sequence of the mitochondrial
DNA of the red alga *Porphyra purpurea*: Cyanobacterial introns and shared ancestry
of red and green algae. *Plant Cell* 11:1675–1694.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.
BLAST+: architecture and applications. *BMC Bioinformatics* 10:421.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25:1972–
1973.

Cerón-Romero MA, Fonseca MM, de Oliveira Martins L, Posada D, Katz LA. 2022.
Phylogenomic analyses of 2,786 genes in 158 lineages support a root of the
eukaryotic tree of life between opisthokonts and all other lineages. *Genome Biol.*
Evol. 14:evac119.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 34:i884–i890.

Christensen AC, Lyznik A, Mohammed S, Elowsky CG, Elo A, Yule R, Mackenzie SA. 2005.
Dual-domain, dual-targeting organellar protein presequences in *Arabidopsis* can use
non-AUG start codons. *Plant Cell* 17:2805–2816.

Derelle R, Lang BF. 2012. Rooting the eukaryotic tree with mitochondrial and bacterial
proteins. *Mol. Biol. Evol.* 29:1277–1289.

Derelle R, Torruella G, Klimeš V, Brinkmann H, Kim E, Vlček Č, Lang BF, Eliáš M. 2015.
Bacterial proteins pinpoint a single eukaryotic root. *Proc. Natl. Acad. Sci.* 112:E693–
E699.

Eddy SR. 2011. Accelerated profile HMM searches. *PLOS Comput. Biol.* 7:e1002195.

Emanuelsson O, Nielsen H, Brunak S, von Heijne G. 2000. Predicting subcellular
localization of proteins based on their N-terminal amino acid sequence. *J. Mol. Biol.*
300:1005–1016.

Felsner G, Sommer MS, Gruenheit N, Hempel F, Moog D, Zauner S, Martin W, Maier UG.
2011. ERAD components in organisms with complex red plastids suggest recruitment

of a preexisting protein transport pathway for the periplastid membrane. *Genome Biol.*
Evol. 3:140–150.

Figueroa-Martinez F, Jackson C, Reyes-Prieto A. 2019. Plastid genomes from diverse
glaucophyte genera reveal a largely conserved gene content and limited architectural
diversity. *Genome Biol. Evol.* 11:174–188.

Filée J, Forterre P, Sen-Lin T, Laurent J. 2002. Evolution of DNA polymerase families:
evidences for multiple gene exchange between cellular and viral proteins. *J. Mol.*
Evol. 54:763–773.

Fukasawa Y, Tsuji J, Fu S-C, Tomii K, Horton P, Imai K. 2015. MitoFates: improved
prediction of mitochondrial targeting sequences and their cleavage sites. *Mol. Cell.*
Proteomics 14:1113–1126.

Fukuda K, Cooney EC, Irwin NAT, Keeling PJ, Hirakawa Y. 2020. High-efficiency
transformation of the chlorarachniophyte *Amorphochlora amoebiformis* by
electroporation. *Algal Res.* 48:101903.

Füssy Z, Záhonová K, Tomčala A, Krajčovič J, Yurchenko V, Oborník M, Eliáš M. 2020. The
cryptic plastid of *Euglena longa* defines a new type of nonphotosynthetic plastid
organelle. *mSphere* 5:10.1128/msphere.00675-20.

Gawryluk RMR, Tikhonenkov DV, Hehenberger E, Husnik F, Mylnikov AP, Keeling PJ. 2019.
Non-photosynthetic predators are sister to red algae. *Nature* 572:240–243.

Gockel G, Hachtel W. 2000. Complete gene map of the plastid genome of the
nonphotosynthetic euglenoid flagellate *Astasia longa*. *Protist* 151:347–351.

Gould SB, Sommer MS, Hadfi K, Zauner S, Kroth PG, Maier U-G. 2006. Protein targeting
into the complex plastid of cryptophytes. *J. Mol. Evol.* 62:674–681.

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,
Raychowdhury R, Zeng Q, et al. 2011. Full-length transcriptome assembly from
RNA-Seq data without a reference genome. *Nat. Biotechnol.* 29:644–652.

Gray MW, Burger G, Derelle R, Klimeš V, Leger MM, Sarrasin M, Vlček Č, Roger AJ, Eliáš
M, Lang BF. 2020. The draft nuclear genome sequence and predicted mitochondrial
proteome of *Andalucia godoyi*, a protist with the most gene-rich and bacteria-like
mitochondrial genome. *BMC Biol.* 18:22.

Graziewicz MA, Longley MJ, Copeland WC. 2006. DNA polymerase γ in mitochondrial DNA
replication and repair. *Chem. Rev.* 106:383–405.

Guilliam TA, Yeeles JTP. 2020. An updated perspective on the polymerase division of labor
during eukaryotic DNA replication. *Crit. Rev. Biochem. Mol. Biol.* 55:469–481.

Harada R, Hirakawa Y, Yabuki A, Kashiyama Y, Maruyama M, Onuma R, Soukal P,
Miyagishima S, Hampl V, Tanifuji G, et al. 2020. Inventory and evolution of
mitochondrion-localized family A DNA polymerases in Euglenozoa. *Pathogens* 9:257.

Harada R, Inagaki Y. 2021. Phage origin of mitochondrion-localized family A DNA
polymerases in kinetoplastids and diplonemids. *Genome Biol. Evol.* 13:evab003.

He D, Fiz-Palacios O, Fu C-J, Fehling J, Tsai C-C, Baldauf SL. 2014. An alternative root for
the eukaryote tree of life. *Curr. Biol.* 24:465–470.

Heiss AA, Walker G, Simpson AGB. 2011. The ultrastructure of Ancyromonas, a eukaryote
without supergroup affinities. *Protist* 162:373–393.

Hirakawa Y, Nagamune K, Ishida K. 2009. Protein targeting into secondary plastids of
chlorarachniophytes. *Proc. Natl. Acad. Sci.* 106:12820–12825.

Hirakawa Y, Watanabe A. 2019. Organellar DNA polymerases in complex plastid-bearing
algae. *Biomolecules* 9:140.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: improving the
ultrafast bootstrap approximation. *Mol. Biol. Evol.* 35:518–522.

Horváthová L, Žárský V, Pánek T, Derelle R, Pyrih J, Motyčková A, Klápšťová V, Vinopalová
M, Marková L, Voleman L, et al. 2021. Analysis of diverse eukaryotes suggests the
existence of an ancestral mitochondrial apparatus derived from the bacterial type II
secretion system. *Nat. Commun.* 12:2947.

Irisarri I, Strassert JFH, Burki F. 2022. Phylogenomic insights into the origin of primary
plastids. *Syst. Biol.* 71:105–120.

Jackson C, Knoll AH, Chan CX, Verbruggen H. 2018. Plastid phylogenomics with broad
taxon sampling further elucidates the distinct evolutionary origins and timing of
secondary green plastids. *Sci. Rep.* 8:1523.

Janouškovec J, Paskerova GG, Miroliubova TS, Mikhailov KV, Birley T, Aleoshin VV,
Simdyanov TG. 2019. Apicomplexan-like parasites are polyphyletic and widely but
selectively dependent on cryptic plastid organelles. *eLife* 8:e49662.

Janouškovec J, Tikhonenkov DV, Burki F, Howe AT, Kolísko M, Mylnikov AP, Keeling PJ.
2015. Factors mediating plastid dependency and the origins of parasitism in
apicomplexans and their close relatives. *Proc. Natl. Acad. Sci.* 112:10200–10207.

Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell
A, Nuka G, et al. 2014. InterProScan 5: genome-scale protein function classification.
Bioinformatics 30:1236–1240.

Joo S, Nishimura Y, Cronmiller E, Hong RH, Kariyawasam T, Wang MH, Shao NC, El Akkad
S-E-D, Suzuki T, Higashiyama T, et al. 2017. Gene regulatory networks for the
haploid-to-diploid transition of *Chlamydomonas reinhardtii*. *Plant Physiol.* 175:314–
332.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder:
fast model selection for accurate phylogenetic estimates. *Nat. Methods* 14:587–589.

Kamikawa R, Tanifuji G, Kawachi M, Miyashita H, Hashimoto T, Inagaki Y. 2015. Plastid
genome-based phylogeny pinpointed the origin of the green-colored plastid in the
dinoflagellate *Lepidodinium chlorophorum*. *Genome Biol. Evol.* 7:1133–1140.

Karnkowska A, Yubuki N, Maruyama M, Yamaguchi A, Kashiyama Y, Suzaki T, Keeling PJ,
Hampl V, Leander BS. 2023. Euglenozoan kleptoplasty illuminates the early
evolution of photoendosymbiosis. *Proc. Natl. Acad. Sci.* 120:e2220100120.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol. Biol. Evol.* 30:772–780.

Keeling PJ, Burki F, Wilcox HM, Allam B, Allen EE, Amaral-Zettler LA, Armbrust EV,
Archibald JM, Bharti AK, Bell CJ, et al. 2014. The Marine Microbial Eukaryote
Transcriptome Sequencing Project (MMETSP): illuminating the functional diversity of
eukaryotic life in the oceans through transcriptome sequencing. *PLOS Biol.*
12:e1001889.

Kim JI, Yoon HS, Yi G, Kim HS, Yih W, Shin W. 2015. The plastid genome of the
cryptomonad *Teleaulax amphioxeia*. *PLOS ONE* 10:e0129284.

Kimura S, Uchiyama Y, Kasai N, Namekawa S, Saotome A, Ueda T, Ando T, Ishibashi T,
Oshige M, Furukawa T, et al. 2002. A novel DNA polymerase homologous to
Escherichia coli DNA polymerase I from a higher plant, rice (*Oryza sativa* L.).
Nucleic Acids Res. 30:1585–1592.

Klingbeil MM, Motyka SA, Englund PT. 2002. Multiple mitochondrial DNA polymerases in
Trypanosoma brucei. *Mol. Cell* 10:175–186.

Krogh A, Larsson B, von Heijne G, Sonnhammer ELL. 2001. Predicting transmembrane
protein topology with a hidden Markov model: application to complete genomes. *J.*
Mol. Biol. 305:567–580.

Kume K, Amagasa T, Hashimoto T, Kitagawa H. 2018. NommPred: prediction of
mitochondrial and mitochondrion-related organelle proteins of nonmodel organisms.
Evol. Bioinforma. 14:1176934318819835.

Lartillot N, Rodrigue N, Stubbs D, Richer J. 2013. PhyloBayes MPI: phylogenetic
reconstruction with infinite mixtures of profiles in a parallel environment. *Syst. Biol.*
62:611–615.

Lax G, Eglit Y, Eme L, Bertrand EM, Roger AJ, Simpson AGB. 2018. Hemimastigophora is a
novel supra-kingdom-level lineage of eukaryotes. *Nature* 564:410–414.

Leebens-Mack JH, Barker MS, Carpenter EJ, Deyholos MK, Gitzendanner MA, Graham SW,
Grosse I, Li Z, Melkonian M, Mirarab S, et al. 2019. One thousand plant
transcriptomes and the phylogenomics of green plants. *Nature* 574:679–685.

Leger MM, Kolisko M, Kamikawa R, Stairs CW, Kume K, Čepička I, Silberman JD,
Andersson JO, Xu F, Yabuki A, et al. 2017. Organelles that illuminate the origins of
Trichomonas hydrogenosomes and *Giardia* mitosomes. *Nat. Ecol. Evol.* 1:1–7.

Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* 22:1658–1659.

Mackiewicz P, Gagat P. 2014. Monophyly of Archaeplastida supergroup and relationships
among its lineages in the light of phylogenetic and phylogenomic studies. Are we
close to a consensus? *Acta Soc. Bot. Pol.* 83:263–280.

Martijn J, Vosseberg J, Guy L, Offre P, Ettema TJG. 2022. Phylogenetic affiliation of
mitochondria with Alpha-II and Rickettsiales is an artefact. *Nat. Ecol. Evol.* 6:1829–
1831.

Mathur V, Kolísko M, Hehenberger E, Irwin NAT, Leander BS, Kristmundsson Á, Freeman
MA, Keeling PJ. 2019. Multiple independent origins of apicomplexan-like parasites.
Curr. Biol. 29:2936-2941.e5.

Mathur V, Salomaki ED, Wakeman KC, Na I, Kwong WK, Kolisko M, Keeling PJ. 2023.
Reconstruction of plastid proteomes of apicomplexans and close relatives reveals the
major evolutionary outcomes of cryptic plastids. *Mol. Biol. Evol.* 40:msad002.

Matsuo E, Morita K, Nakayama T, Yazaki E, Sarai C, Takahashi K, Iwataki M, Inagaki Y.
2022. Comparative plastid genomics of green-colored dinoflagellates unveils parallel
genome compaction and RNA editing. *Front. Plant Sci.* 13:918543.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear
R. 2020. IQ-TREE 2: new models and efficient methods for phylogenetic inference in
the genomic era. *Mol. Biol. Evol.* 37:1530–1534.

Miyahara M, Aoi M, Inoue-Kashino N, Kashino Y, Ifuku K. 2013. Highly efficient
transformation of the diatom *Phaeodactylum tricornutum* by multi-pulse
electroporation. *Biosci. Biotechnol. Biochem.* 77:874–876.

Moog D, Nozawa A, Tozawa Y, Kamikawa R. 2020. Substrate specificity of plastid
phosphate transporters in a non-photosynthetic diatom and its implication in evolution
of red alga-derived complex plastids. *Sci. Rep.* 10:1167.

Mori Y, Kimura S, Saotome A, Kasai N, Sakaguchi N, Uchiyama Y, Ishibashi T, Yamamoto T,
Chiku H, Sakaguchi K. 2005. Plastid DNA polymerases from higher plants,
Arabidopsis thaliana. Biochem. Biophys. Res. Commun. 334:43–50.

Moriyama T, Sato N. 2014. Enzymes involved in organellar DNA replication in
photosynthetic eukaryotes. *Front. Plant Sci.* 5:480.

Moriyama T, Terasawa K, Fujiwara M, Sato N. 2008. Purification and characterization of
organellar DNA polymerases in the red alga *Cyanidioschyzon merolae. FEBS J.*
275:2899–2918.

Moriyama T, Terasawa K, Sato N. 2011. Conservation of POPs, the plant organellar DNA
polymerases, in eukaryotes. *Protist* 162:177–187.

Mukhopadhyay A, Chen C-Y, Doerig C, Henriquez FL, Roberts CW, Barrett MP. 2009. The
Toxoplasma gondii plastid replication and repair enzyme complex, PREX.
Parasitology 136:747–755.

Muñoz-Gómez SA, Susko E, Williamson K, Eme L, Slamovits CH, Moreira D, López-García
P, Roger AJ. 2022. Site-and-branch-heterogeneous analyses of an expanded
dataset favour mitochondria as sister to known Alphaproteobacteria. *Nat. Ecol. Evol.*
6:253–262.

Nassoury N, Morse D. 2005. Protein targeting to the chloroplasts of photosynthetic
eukaryotes: getting there is half the fun. *Biochim. Biophys. Acta - Mol. Cell Res.*
1743:5–19.

Nishimura Y, Shiratori T, Ishida K, Hashimoto T, Ohkuma M, Inagaki Y. 2019. Horizontally-
acquired genetic elements in the mitochondrial genome of a centrohelid *Marophrys*
sp. SRT127. *Sci. Rep.* 9:4850.

Novák Vanclová AMG, Zoltner M, Kelly S, Soukal P, Záhonová K, Füssy Z, Ebenezer TE,
Lacová Dobáková E, Eliáš M, Lukeš J, et al. 2020. Metabolic quirks and the colourful
history of the *Euglena gracilis* secondary plastid. *New Phytol.* 225:1578–1592.

Okazaki T. 2017. Days weaving the lagging strand synthesis of DNA — A personal
recollection of the discovery of Okazaki fragments and studies on discontinuous
replication mechanism —. *Proc. Jpn. Acad. Ser. B* 93:322–338.

Ono Y, Sakai A, Takechi K, Takio S, Takusagawa M, Takano H. 2007. *NtPolI-like1* and
NtPolI-like2, bacterial DNA polymerase I homologs isolated from BY-2 cultured
tobacco cells, encode DNA polymerases engaged in DNA replication in both plastids
and mitochondria. *Plant Cell Physiol.* 48:1679–1692.

Parent J-S, Lepage E, Brisson N. 2011. Divergent roles for the two PolI-like organelle DNA
polymerases of Arabidopsis. *Plant Physiol.* 156:254–262.

Patron NJ, Waller RF. 2007. Transit peptide diversity and divergence: A global analysis of
plastid targeting signals. *BioEssays* 29:1048–1058.

Paysan-Lafosse T, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar GA, Bileschi ML,
Bork P, Bridge A, Colwell L, et al. 2023. InterPro in 2022. *Nucleic Acids Res.*
51:D418–D427.

Petsalaki EI, Bagos PG, Litou ZI, Hamodrakas SJ. 2006. PredSL: a tool for the N-terminal
sequence-based prediction of protein subcellular localization. *Genomics Proteomics*
Bioinformatics 4:48–55.

Ponce-Toledo RI, Deschamps P, López-García P, Zivanovic Y, Benzerara K, Moreira D.
2017. An early-branching freshwater cyanobacterium at the origin of plastids. *Curr.*
Biol. 27:386–391.

Reesey E. 2017. Characterization of the mitochondrial DNA polymerase in *Plasmodium*
falciparum. Available from:
https://www.proquest.com/docview/1984381231/abstract/2682E909B503458DPQ/1

Richter DJ, Berney C, Strassert JFH, Poh Y-P, Herman EK, Muñoz-Gómez SA, Wideman
JG, Burki F, de Vargas C. 2022. EukProt: A database of genome-scale predicted
proteins across the diversity of eukaryotes. *Peer Community J.* 2:e56.

Rodríguez-Ezpeleta N, Brinkmann H, Burey SC, Roure B, Burger G, Löffelhardt W, Bohnert
HJ, Philippe H, Lang BF. 2005. Monophyly of primary photosynthetic eukaryotes:
green plants, red algae, and glaucophytes. *Curr. Biol.* 15:1325–1330.

Roger AJ, Muñoz-Gómez SA, Kamikawa R. 2017. The origin and diversification of
mitochondria. *Curr. Biol.* 27:R1177–R1192.

Salomaki ED, Terpis KX, Rueckert S, Kotyk M, Varadínová ZK, Čepička I, Lane CE, Kolisko
M. 2021. Gregarine single-cell transcriptomics reveals differential mitochondrial
remodeling and adaptation in apicomplexans. *BMC Biol.* 19:77.

Sarai C, Tanifuji G, Nakayama T, Kamikawa R, Takahashi K, Yazaki E, Matsuo E, Miyashita
H, Ishida K, Iwataki M, et al. 2020. Dinoflagellates with relic endosymbiont nuclei as
models for elucidating organellogenesis. *Proc. Natl. Acad. Sci.* 117:5364–5375.

Sayers EW, Beck J, Bolton EE, Bourexis D, Brister JR, Canese K, Comeau DC, Funk K, Kim
S, Klimke W, et al. 2021. Database resources of the National Center for
Biotechnology Information. *Nucleic Acids Res.* 49:D10–D17.

Schiestl RH, Gietz RD. 1989. High efficiency transformation of intact yeast cells using single
stranded nucleic acids as a carrier. *Curr. Genet.* 16:339–346.

Schön ME, Zlatogursky VV, Singh RP, Poirier C, Wilken S, Mathur V, Strassert JFH,
Pinhassi J, Worden AZ, Keeling PJ, et al. 2021. Single cell genomics reveals plastid-
lacking Picozoa are close relatives of red algae. *Nat. Commun.* 12:6651.

Shimodaira H. 2002. An approximately unbiased test of phylogenetic tree selection. *Syst.*
Biol. 51:492–508.

Sibbald SJ, Archibald JM. 2020. Genomic insights into plastid evolution. *Genome Biol. Evol.*
12:978–990.

Simpson AGB. 2003. Cytoskeletal organization, phylogenetic affinities and systematics in
the contentious taxon Excavata (Eukaryota). *Int. J. Syst. Evol. Microbiol.* 53:1759–
1777.

Stairs CW, Eme L, Brown MW, Mutsaers C, Susko E, Dellaire G, Soanes DM,
van der Giezen M, Roger AJ. 2014. A SUF Fe-S cluster biogenesis system in the
mitochondrion-related organelles of the anaerobic protist *Pygsuia. Curr. Biol.*
24:1176–1186.

Stairs CW, Táborský P, Salomaki ED, Kolisko M, Pánek T, Eme L, Hradilová M, Vlček Č,
Jerlström-Hultqvist J, Roger AJ, et al. 2021. Anaeramoebae are a divergent lineage
of eukaryotes that shed light on the transition from anaerobic mitochondria to
hydrogenosomes. *Curr. Biol.* 31:5605-5612.e5.

Suzuki S, Ishida K-I, Hirakawa Y. 2016. Diurnal transcriptional regulation of
endosymbiotically derived genes in the chlorarachniophyte *Bigelowiella natans.*
Genome Biol. Evol. 8:2672–2682.

Tardif M, Atteia A, Specht M, Cogne G, Rolland N, Brugière S, Hippler M, Ferro M, Bruley C,
Peltier G, et al. 2012. PredAlgo: a new subcellular localization prediction tool
dedicated to green algae. *Mol. Biol. Evol.* 29:3625–3639.

Tice AK, Žihala D, Pánek T, Jones RE, Salomaki ED, Nenarokov S, Burki F, Eliáš M, Eme L,
Roger AJ, et al. 2021. PhyloFisher: A phylogenomic package for resolving eukaryotic
relationships. *PLOS Biol.* 19:e3001365.

Tikhonenkov DV, Mikhailov KV, Gawryluk RMR, Belyaev AO, Mathur V, Karpov SA,
Zagumyonnyi DG, Borodina AS, Prokina KI, Mylnikov AP, et al. 2022. Microbial
predators form a new supergroup of eukaryotes. *Nature* 612:714–719.

Tikhonenkov DV, Strassert JFH, Janouškovec J, Mylnikov AP, Aleoshin VV, Burki F, Keeling
PJ. 2020. Predatory colponemids are the sister group to all other alveolates. *Mol.*
Phylogenet. Evol. 149:106839.

Turmel M, Gagnon M-C, O'Kelly CJ, Otis C, Lemieux C. 2009. The chloroplast genomes of
the green algae *Pyramimonas*, *Monomastix*, and *Pycnococcus* shed new light on the
evolutionary history of prasinophytes and the origin of the secondary chloroplasts of
euglenids. *Mol. Biol. Evol.* 26:631–648.

Wang H-C, Minh BQ, Susko E, Roger AJ. 2018. Modeling site heterogeneity with posterior
mean site frequency profiles accelerates accurate phylogenomic estimation. *Syst.*
Biol. 67:216–235.

Westermann B, Neupert W. 2000. Mitochondria-targeted green fluorescent proteins:
convenient tools for the study of organelle biogenesis in *Saccharomyces cerevisiae.*
Yeast 16:1421–1427.

Yabuki A, Gyaltshen Y, Heiss AA, Fujikura K, Kim E. 2018. *Ophirina amphinema* n. gen., n.
sp., a new deeply branching discobid with phylogenetic affinity to jakobids. *Sci. Rep.*
8:16219.

Yamaguchi A, Yubuki N, Leander BS. 2012. Morphostasis in a novel eukaryote illuminates
the evolutionary transition from phagotrophy to phototrophy: description of *Rapaza*
viridis n. gen. et sp. (Euglenozoa, Euglenida). *BMC Evol. Biol.* 12:29.

Yazaki E, Miyata R, Chikami Y, Harada R, Kawakubo T, Tanifuji G, Nakayama T, Yahata K,
Hashimoto T, Inagaki Y. 2021. Signs of the plastid: Enzymes involved in plastid-
localized metabolic pathways in a eugregarine species. *Parasitol. Int.* 83:102364.

Yazaki E, Yabuki A, Imaizumi A, Kume K, Hashimoto T, Inagaki Y. 2022. The closest lineage
of Archaeplastida is revealed by phylogenomics analyses that include *Microheliella*
maris. Open Biol. 12:210376.

1264

1265

1266 Figure legends

Figure 1. Global phylogeny of family A DNA polymerases sampled from
bacteria, phages, and eukaryotes.

The ML tree was inferred from an alignment comprising 488 family A DNA polymerases
(famA DNAPs). The terminal branches colored in orange, green, and blue represent the
famA DNAPs originated from bacteria, phages, and eukaryotes, respectively. The branches
of 15 types of famA DNAP found in eukaryotes are shaded in blue. We present the ultra-fast
bootstrap support (UFBP) value for the node uniting the paraphyletic chloroPolA sequences
and cyanobacterial PolI sequences on the corresponding node. Otherwise, the UFBP value
for the monophyly of each type of eukaryotic famA DNAP is shown in the shade.

Figure 2. Subcellular localization of GFP fused with the N-terminus of
chlnmPolA when expressed in chlorarachniophyte cells.

Green fluorescent protein (GFP) fused with the first 100 amino acid residues of the
chlorarachniophyte *Bigelowiella natans* chlnmPolA was expressed in the chlorarachniophyte
Amorphochlora amoebiformis. The red color corresponds to chlorophyll autofluorescence.
The green signal represents the GFP localization. The GFP signal was found to be
associated tightly with but distinct from plastids, suggesting the protein is localized in the
periplastidial compartment. The scale bars are 10 µm.

Figure 3. Subcellular localizations of GFP fused with the N-termini of the
three organellar family A DNA polymerases found in the cryptophyte
Guillardia theta when expressed in diatom cells.
GFP fused with the N-terminal 332 amino acid residues of *G. theta* cryotoPolA (A), 100
residues of *G. theta* POP1 (B & C), and 516 residues of *G. theta* POP2 (D & E) were
heterogeneously expressed in the diatom *Phaeodactylum tricornutum* UTEX642. In A, C, D,
and E, the red color corresponds to chlorophyll autofluorescence and the green signal
represents GFP. In B, the red color indicates both chlorophyll autofluoresence and
mitochondria stained by MitoTracker Orange and the green signal represents GFP. The
scale bars are 10 µm.

Figure 4. Subcellular localizations of GFP fused with the N-termini of five
rdxPolAs when expressed in yeast cells.
GFP fused with the N-terminal amino acid residues predicted as the mitochondrial targeting
signal (MTS) of rdxPolA proteins from three members of Discoba (*Ophirina amphinema*,
Tsukubamonas globosa, and *Naegleria gruberi*; A-C), the malawimonad *Gefionella okellyi*
(D), and the ancyromonad *Fabomonas tropica* (E). The green signal corresponds to GFP.
The red color indicates mitochondria stained by MitoTracker Red. The results presented
here demonstrate that the mitochondrial translocons in yeast recognize the N-termini of the
five rdxPolAs as the MTS. The scale bars are 5 µm.

Figure 5. Phylogenies that clarified the origins of six novel types of family
A DNA polymerases localized in organelles.
The phylogenies assessing the origins of rgPolA (panel A), chloroPolA (panel B),
pyramiPolA and eugPolA (panel C), and rdxPolA and cryptoPolA (panel D). The trees shown
here were inferred with the ML phylogenetic method. ML nonparametric bootstrap support
values are presented only for nodes key to the origins of the DNA polymerases of interest. In
addition, the key nodes that received Bayesian posterior probabilities equal to or greater
than 0.95 are marked by dots. The clades/branches are color-coded, as shown in the inset.
The PolI of the cyanobacterium *Gloeomargarita lithophora* is marked with a green allow.

Figure 6. Summary of organellar family A DNA polymerases and their
phylogenetic distributions
This figure summarizes the diversity of organellar family A DNA polymerases (DNAPs) and
their phylogenetic distributions over the tree of eukaryotes (schematically shown on the left).
The presence of each type of organellar DNAP is indicated by a dot colored in blue
(mitochondrion-localized; MT), red (plastid-localized; PL), or green (nucleomorph-localized;
NM), respectively. pyramiPolA found in green algae of the order Pyramimonadales is shown
in a gray dot, as its exact organellar (mitochondrial or plastid) localization remains
inconclusive. None of the currently known members of Breviatea or Metamonada, which
bear highly reduced mitochondria whose own genomes had been discarded completely, is
anticipated to retain any organellar DNAP, and the corresponding rows were left blank. We
found no candidate for mitochondrial DNAP in the transcriptomes of hemimastigophorans
and thus commented so ("NO DATA AVAILABLE").

Figure 7. Phylogenomic analyses of eukaryotes.

(A) The phylogenetic relationship among 97 eukaryotes was inferred from a phylogenomic
alignment comprising 340 proteins by using the ML method. Major clades are shown by
triangles. All the nodes, except for those labeled as a-e, received ultra-fast bootstrap support
(UFBP) values of 100%. The UFBP values for nodes a-e were summarized in the inset table.
The assemblage/lineages/species that use rdxPolA as mitochondrion-localized DNA
polymerase are marked by stars. (B) Alternative topologies of the eukaryotic phylogeny
assessed in an approximately unbiased test. In the ML tree topology (Tree 1), Amorphea
plus CRuMs branch at the node labeled as 1 in the schematic tree. Amorphea plus CRuMs
were pruned and regrafted to six alternative positions labeled as 2-7 to generate alternative
trees (Trees 2-7). For each alternative tree (Trees 2-7), the difference in log-likelihood from
the ML tree ($\Delta \log L$) and the P-value from the test are summarized in the table.

1337
1338

1339   Table 1. List of family A DNA polymerases (DNAPs) in Eukaryota known to date.

DNAP	Phylogenetic distribution	Subcellular localization	Proposed origin
POP	Diverse eukaryotes	mitochondrion & plastid	uncertain
Polθ/ν	Diverse eukaryotes	nucleus	uncertain
Polγ	Opisthokonta	mitochondrion	a T-odd number phage (Filée et al. 2002)
PREX	Apicomplexa Chlompodellids Squirmidea	plastid	an uncertain bacterium (Janouškovec et al. 2019)
PolIA	Euglenozoa	mitochondrion	Polθ/ν (Harada and Inagaki 2021)
PolIBCD+	Diplonemea Kinetoplastea	mitochondrion	an autographivirus (Harada and Inagaki 2021)
acPolA	Apicomplexa Chlompodellids	mitochondrion	uncertain
rdxPolA	Malawimonadida Ancyromonadida Discoba*	mitochondrion	an α-proteobacterium
rgPolA	Rhodophyta Glaucophyta	plastid	a Rhodothermales bacterium (Moriyama et al. 2008)
eugPolA	Plastid-bearing members of Euglenophyceae	plastid	the endosymbiont green alga that gave rise to the euglenophycean plastids (i.e., *Pyramimonas* or their close relative)
pyramiPolA	Pyramimonadales	uncertain	a γ-proteobacterium
chloroPolA	Chlorophyceae	mitochondrion	a cyanobacterium
cryptoPolA	Cryptophyceae	plastid	an α-proteobacterium
chlnmPolA	Chlorarachniophyta	nucleomorph	an uncertain bacterium
abanPolA	Apusomonadida Breviatea Amoebozoa Nebulidia	nucleus	a Planctomycetes bacterium
alvPolA	Alveolata	nucleus	uncertain

1340   * Discoba comprises Jakobida, *Tsukubamonas*, Heterolobosea, and Euglenozoa, but
1341   rdxPolA appeared absent in Euglenozoa.

bacterial famA DNAP (PolI)
phage famA DNAP
famA DNAP in eukaryotes

100 rgPolA

99
chloroPolA
Cyanobacteria

100 rdxPolA
100 cryptoPolA
99 eugPolA
99 pyramiPolA
100 chlnmPolA

100 PolIA & Polθ/ν

100 POP

100 acPolA

100 PREX

100 alvPolA

100 abanPolA

100 PolIBCD+

100 Polγ

0.5

Harada et al. Fig. 1

Bright Field **Chl** **GFP** **Merge**

Harada et al. Fig. 2

Harada et al. Fig. 3

Bright Field	GFP	MitoTracker Red	Merge

Harada et al. Fig. 4

A

rgPolA

Chondrus crispus
Porphyra umbilicalis
Galdieria sulphuraria
Rhodella violacea
Compsopogon caeruleus
Porphyridium purpureum
Rhodosorus marinus
Cyanidioschyzon merolae
Cyanophora paradoxa
Gloeochaete wittrockiana
Longimonas halophila
Rhodothermus marinus
Rubricoccus marinus

Rhodothermales PolI

67
58
99
89

0.5 substitutions/site

Eukaryota
Alphaproteobacteria
Betaproteobacteria
Gammaproteobacteria
Deltaproteobacteria
Cyanobacteria
Firmicutes
Other bacteria

B

chloroPolA

Golenkinia longispicula
Fritschiella tuberosa
Stigeoclonium helveticum
Hormotilopsis gelatinosa
Chaetopeltis orbicularis
Chlamydomonas schloesseri
Chlamydomonas incerta
Chlamydomonas reinhardtii
Gonium pectorale
Gonium pectorale
Chloromonas rosa
Chloromonas tughillensi
Pediastrum duplex
Monoraphidium neglectum
Scenedesmus dimorphus
Scenedesmus sp. NREL 46B-D3
Chlamydomonas noctigama
Chlamydomonas sp. UWO 241
Chlamydomonas eustigma
Carteria obtusa
Oogamochlamys gigantea
Dunaliella primolecta
Dunaliella tertiolecta
Dunaliella salina
Pleurastrum insigne
Chloromonas perforata
Microspora cf. tumidula

0.5 substitutions/site

C

pyramiPolA

79
64
96

Pyramimonas parkeae NIES254 & CCMP726
Polyblepharides amylifera NIES251
Cymbomonas tetramitiformis
Euglena gracilis
Euglena longa
Euglena sp. M2084
Eutreptiella gymnastica

eugPolA

0.5 substitutions/site

D

cryptoPolA

Guillardia theta
Hanusia phi
Baffinella frigidus
Geminigera cryophila
Rhodomonas lens

Naegleria gruberi
Naegleria fowleri
Willaertia magna
Barbeliidae sp. AE-1
Percolomonas cosmopolitus WS
Neovahlkampfia damariscottae
Tsukubamonas globosa
Pharyngomonas kirbyi
Heterolobosea sp. BB2
Ancyromonas sigmoides
Nutomonas longa
Fabomonas tropica SRT902
Andalucia godoyi
Ophirina amphinema
Reclimonas americana
Gefionella okellyi
Malawimonas californiana

rdxPolA

0.5 substitutions/site

Harada et al. Fig. 5

Harada et al. Fig. 6

Harada et al. Fig. 7