Limb regeneration is blastema dependent in a ladybird beetle, *Cheilomenes sexmaculata* (Fabricius)

Saumya Rawat[1], Shriza Rai[1], Geetanjali Mishra[2]*

1. **Saumya Rawat:** M Sc. Student, Ladybird Research Laboratory, Department of Zoology, University of Lucknow, Lucknow- 226007, email: rsaumya810@gmail.com

   Contact no.: +91 9129183389

2. **Shriza Rai**: Research Scholar, Ladybird Research Laboratory, Department of Zoology, University of Lucknow, Lucknow- 226007, India; email: shrizarai@gmail.com

   Contact no.: +91 9648072790

3. **Geetanjali Mishra*:** Professor, Ladybird Research Laboratory, Department of Zoology, University of Lucknow, Lucknow- 226007, India; email: geetanjalimishra@gmail.com

   Contact no.: +91 9415956692

Saumya Rawat and Shriza Rai are joint first authors

*Corresponding author email: geetanjalimishra@gmail.com

Short running title: Blastema dependent limb regeneration in ladybirds.

Acknowledgements

GM acknowledges SERB DST (F.No. CRG/2020/001095 via letter no. BP-2020-20-3308 dated December 30, 2020. SR acknowledges UGC Fellowship by University Grants Commission, New Delhi, India (F.No.16-9(June 2019)/ 2019(NET/CSIR)) dated August 09, 2019.

Authors Contributions

SR and GM conceived the presented idea and designed the experiment; SR and SaR collected the sample and conducted the experiment; SaR analysed the data; GM supervised the complete study along with SR who also contributed in writing of the manuscript. All authors contributed significantly to the draft and gave final approval for publication.

Conflict of Interest

27   All authors are in agreement with the content of the manuscript and have no conflict of interest.

28   **Ethical Approval Statement**: All the ethics regarding the insect model system have been

29   followed in conducting the experiment.

Limb regeneration is blastema dependent in a ladybird beetle, *Cheilomenes sexmaculata* (Fabricius)

Abstract

Holometabolous insects undergoes metamorphosis which involves an intercalated pupal

stage between larva and adult. The body plan of the adult is established during pupal stage and

larval systems are de-differentiated and reorganized in insects undergoing complete

metamorphosis. In ladybird beetles, limbs amputated in larval stages are regenerated in adults.

This occurs during pupation. Given that changes in pupa are akin to embryogenesis, does the

lost limbs are redeveloped as a part of metamorphosis or has some pre-patterning initiated prior

pupation? To study this we hypothesised that limb regeneration in a holometabolous ladybird

beetles, *Cheilomenes sexmaculata* is the result of the recapitulation of the embryonic gene

programs in the pupa. To test this, we exposed third larval stages of *C. sexmaculata* to

amputation and prior to amputation the tissues on the amputated site was scraped off every

24hrs. It was found that limb regeneration does not occur in the treatment where scrapping was

done. Assuming that these epidermal cells correspond to blastema, for limb regeneration in

ladybird beetles blastema is essential and does not occur in its absence.

Keywords: Amputation, leg development, holometabolous, pupa, metamorphosis

Introduction

More than 80% of insects undergo complete metamorphosis or holometaboloy with an intercalated pupal stage between larva and adult (Engel and Grimaldi, 2005). The pupal stage, which is largely immobile, involves extensive remodelling of organs and tissues, resulting in rebuilding of entire body plan (Rolff et al., 2019). Many metabolic and developmental genes have been reported to revert to embryonic-like state during metamorphosis. Ozerova and Gelfand (2022), hypothesised that there is recapitulation of the embryonic expression program during metamorphosis and demonstrated an increased similarity of pupa to embryonic stage via metanalyses of numerous datasets of holometabolous insects.

Ladybird beetles are holometabolous insects that undergoes limb regeneration (Wang et al. 2015, Saxena et al. 2016, Michaud et al., 2020). An amputated limb is regenerated when larva metamorphosizes into adult. The capacity of limb regeneration has been reported to be broadly conserved in coccinellids with 15 out of 16 species of ladybird beetles exhibiting limb regeneration with delayed pupal duration. Regeneration neither occurred from one moult to another nor the amputated legs in adults showed regeneration (Wang et al., 2015; Saxena et al., 2016) and was evident only during pupal development. An activation of genes responsible for regeneration probably occurred in pupal stage (Michaud et al., 2020; Zhong et al., 2023). Possibly the cells initiate differentiation during pupation forming imaginal primordia, contributing to the development of adult structures and maintain pluripotency similar to stem cells (Truman and Riddiford, 2002; McClure and Schubiger, 2007). In *Drosophila,* regeneration occurs via imaginal disc formation while in *Gryllus* regeneration involves blastema formation; however, the signalling involved during regeneration is similar in both (Das. 2015). Having said that it is yet to be explored if redevelopment of lost limbs in pupa is a part of metamorphosis or involves the process of regeneration via blastema formation. Since pupal stage involves whole-body reorganization; most larval tissues and organs are respecified

including epidermis (Riddiford 1978) and muscles (Lubischer et al 1999), it was hypothesized

that limb regeneration in holometabolous insects like ladybird beetles, is the embryonic

recapitulation of the gene programs in the pupa and is therefore independent of blastema.

Therefore, this experiment will help us to understand if the ability to form a blastema is directly

related to the ability to regenerate lost legs or is a part of metamorphosis.

Ladybird beetles are known globally for their biological potential. During mass rearing, it

is not uncommon for larvae to incur nonlethal injuries including appendage loss owing to fierce

cannibalism exhibited by these beetles (Michaud 2003). Current study will be helpful in

increasing our understanding of limb regeneration in non-model insects like ladybirds and offer

better insight into the evolution of regeneration.

Materials and Methods

Study Organism

Cheilomenes sexmaculata (Fabricius) was selected for this study because it exhibits limb

regeneration (Saxena et al., 2016) and is widely distributed in the local agricultural fields of

India.

Stock Maintenance

Adults of *C. sexmaculata* (n=20) were collected from the fields of Lucknow, India

(26°50'N, 80°54'E), and were reared for two generations in the laboratory. They were provided

with an *ad libitum* supply of *Aphis nerii* Boyer de Fonscolombe (Hemiptera: Aphididae) on the

host plant *Calotropis procera* (Family: Apocynaceae) commonly known as milkweed aphid

which is found locally in abandoned lands. Stock culture of beetles was done in a 1 litre plastic

beaker and kept in temperature-controlled room set to 27±2°C, 65±5% R.H. Mated females

were removed from beakers and transferred to plastic Petri dishes (9.0×2.0 cm) and

provisioned *ad libitum* aphids. The eggs laid were collected every 24h, and observed for

hatching and the first instar larvae were used in the experiment.

Experimental Design

120 newly hatched larvae from the stock were placed in two sets of 40 each and kept

singly in Petri dishes (9.0×2.0 cm). In the first set, amputation was performed in third larval

instar (L3) stage. At first, the larvae were chilled individually for 5 min in a refrigerator and

were amputated at the proximal end of the femur of the right foreleg. Amputation was

performed under a stereomicroscope (Magnus)with the magnification 16X with the help of a

micro-scalpel. After 24hrs of amputation, the epidermal growth at the proximal position of

amputation site was scraped off using a micro-scalpel under the stereomicroscope (Magnus).

This was repeated after every 24hrs till the larva underwent pupation. In the second set there

was amputation in L3 stage but no scraping off was performed while in third set there was no

amputation and was called the control. All three sets were maintained on *ad libitum* supply of

A. nerii, until the emergence of the adult. Petri dishes were cleaned every 24 hours throughout

the experiment to maintain the prey quality and to avoid any microbial contamination. Larvae

were examined daily to record their larval stage developmental durations, pupal duration and

post amputation developmental duration, for amputated (scraped), amputated (unscraped) and

control beetles.

Statistical Analysis

Data sets were first tested for normality using the Kolmogorov-Smirnov test and were

found to be non-normal for larval stage developmental durations, pupal duration, and post

amputation developmental duration of Control, Scraped and Unscraped beetles as treatments.

On finding the data non-normal, data were subjected to Generalized Linear Model (GLM) with

Gamma log link. Treatments were taken as fixed factors and larval stage developmental

durations, pupal duration, and post amputation developmental duration were taken as

dependent factors. This analysis was followed by a comparison of means using Wald Chi-

Square test. The analysis was done using IBM SPSS version 20 statistical software.

Results and Discussion

Out of a total of 40 replicates, only 6 regenerated the amputated leg, while the rest did not regenerate (Figure 1). The everyday scraping off of the tip of the regenerating leg corresponds to removal of blastema. Thus, our results demonstrated the importance of blastema for the regeneration of lost limbs in ladybird beetles. The 6 individuals that showed some level of regeneration of the leg amputated in the larval stage, could be the failure to completely scrape off the blastema. Thus, even some remaining cells of blastema initiated regeneration but were not enough to lead to complete development of the lost leg.

The proximal part of the amputated site, gets covered by a scab of epidermal cells. The removal of this scab probably resulted in the removal of blastema because of which there was no regeneration in the treatment that involved scraping. This demonstrates the blastema dependency of limb regeneration in *C. sexmaculata.* Das (2015) reported that limb regeneration in insects is initiated via scab formation, followed by migration of epidermal cells at the site of amputation and these epidermal cells known as blastema, acted as the precursors of the regenerating cells. Various cellular processes and signalling unfold in blastema ultimately specifying the cells to give rise to the formation of lost structures (Zhong et al., 2023). There is a combination of genetic factors forming a 'prepattern', which presages the final differentiated structure. For example, several genes like *dachshund* (Dac) aid in regeneration and their absence resulted in abnormally shortened legs (Lee et al., 2013). Dac silenced in amputated larval leg, affected the patterning of the adult leg, indicating that larval appendage morphology is inherited by the adult appendages.

Amputation of foreleg led to delayed post amputation development duration with maximum delay in unscraped treatment (Table 1). Pupal duration was found to be maximum in unscraped treatment followed by scraped treatment and control. Interestingly, the pupal

duration in scraped treatment was similar to control. This difference from previous studies

where the maximum delay of amputated beetles was reported in the pupal stage (Wang et al.,

2015; Saxena et al., 2016; Michaud et al., 2020; Rai et al., 2023), can be attributed to the

absence of blastemal cells for the initiation of regeneration and hence no regeneration occurred

in pupa, therefore no delay. While delay in the developmental duration of the fourth larval

stage of the amputated treatment demonstrated the attempt to form the blastema via cellular

proliferation, with no blastema in the larvae entering pupation there was no initiation of

regeneration. Since, regeneration in ladybird beetles occurs during pupal duration (Wang et al.,

2015; Saxena et al., 2016; Michaud et al., 2020; Zhou et al., 2021), with no regeneration taking

place in the pupa, the amputated larvae of scraped off treatment showed no delay in the pupa.

This explains that the delay in the pupa is indicative of regenerative processing occurring

inside the pupa.

Our hypothesis that the legs amputated in the larval stages, are redeveloped during

metamorphosis as a part of embryonic recapitulation of the gene programs in the pupa, was

rejected. This study concluded that (i) scraping of scab formed at the proximal position of the

amputation site, prevented the formation of blastema, (ii) the ability to form a blastema is

directly related to the insects' ability to regenerate lost legs, and (iii) the delay in pupa is

indicative of undergoing regeneration processes. Thus, limb regeneration in ladybird beetle *C.*

sexmaculata is blastema dependent and the lost structure is reconstructed from blastema.

Further studies are needed to understand the role of biotic and abiotic factors on blastema

development. Transcriptome analysis of pupa with blastema and without blastema is needed to

further understand the molecular aspect of limb regeneration in ladybird beetles.

References

Engel MS, Grimaldi D. (2005). *Evolution of the insects*. Cambridge, UK: Cambridge

University Press

Lee, A.K., Sze, C.C., Kim, E.R. *et al.* (2013). Developmental coupling of larval and adult

stages in a complex life cycle: insights from limb regeneration in the flour

beetle, *Tribolium castaneum. EvoDevo.* 4, 20.

Lubischer JL, Verhegge LD, Weeks JC. (1999) Respecified larval proleg and body wall

muscles circulate hemolymph in developing wings of *Manduca sexta* pupae. *J. Exp.*

Biol. 202, 787-796.

McClure, K. D. & Schubiger, G. (2007). Transdetermination: Drosophila imaginal disc cells

exhibit stem cell-like potency. *Int. J. Biochem. Cell Biol*. 39, 1105–1118.

Michaud JP, Nedved O, Bayoumy M, Abdelwahab A, Torres J, Saxena S, Omkar, de Santos-

Cividanes TM, Sakaki S, Rasekh A, Tomoli Torfi E. (2020). The Capacity for Limb

Regeneration During Metamorphosis is Broadly Conserved in the Coccinellidae. *Annals*

of the Entomological Society of America. 113(6):452-60.

Michaud, J. P. (2003). A comparative study of larval cannibalism in three species of

ladybird. *Ecological entomology* 28, (1): 92-101.

Ozerova, A.M., Gelfand, M.S. (2022). Recapitulation of the embryonic transcriptional program

in holometabolous insect pupae. *Sci Rep.* 12, 17570.

Rai S, Singh A, Omkar O, Mishra G (2023). Effect of larval thermal conditions on limb

regeneration in a ladybird beetle, *Cheilomenes sexmaculata. Journal of Experimental*

Zoology Part A: Ecological and Integrative Physiology. 339(9), 825-837.

Riddiford LM. (1978). Ecdysone-induced change in cellular commitment of the epidermis of the tobacco hornworm, *Manduca sexta*, at the initiation of metamorphosis. *Gen. Comp. Endocrinol.* 34, 438-446.

Rolff Jens, Johnston Paul R. and Reynolds Stuart (2019).Complete metamorphosis of insects *Phil. Trans. R. Soc.* B3742019006320190063.

Saxena, S., & Mishra, G. (2016). Does regeneration ability influence reproductive fitness in *Menochilus sexmaculatus* (Coleoptera: Coccinellidae)? *Journal of Asia-Pacific Entomology.* 19, 829–834.

Das, S (2015). Morphological, Molecular, and Hormonal Basis of Limb Regeneration across Pancrustacea, *Integrative and Comparative Biology*, 55(5), 869–877.

Truman, J. W. & Riddiford, L. M. (2002). Endocrine insights into the evolution of metamorphosis in insects. *Annu. Rev. Entomol*. 47, 467–500.

Wang, S., Tan, X. L., Michaud, J. P., Shi, Z. K., & Zhang, F. (2015). Sexual selection drives the evolution of limb regeneration in *Harmonia axyridis* (Coleoptera: Coccinellidae).

Zhong, J., Jing, A., Zheng, S. *et al.* (2023). Physiological and molecular mechanisms of insect appendage regeneration. *Cell Regen* 12, 9.

Zhou H, Ma Z, Wang Z, Yan S, Wang D, Shen J. (2021). Hedgehog signaling regulates regenerative patterning and growth in *Harmonia axyridis* leg. *Cellular and Molecular Life Sciences*. 78, 2185-97.

217   **Table 1:** Results of GLM with Gamma log link depicting the effect of amputation on Post
218   amputation development duration (PADD), larval development durations (L_3 and L_4) and pupal
219   duration in *Cheilomenes sexmaculata,* along with their mean values.

Parameters	Wald-Chi square	P- value	df	Mean values	
PADD	304.790	0.000	2	Unscraped	(2.472±0.022) days
				Scraped	(2.206±0.022) days
				Control	(1.935±0.022) days
L_3 developmental duration	38.797	0.000	2	Unscraped	(1.021±0.059) days
				Scraped	(0.655±0.059) days
				Control	(0.516±0.059) days
L_4 developmental duration	107.745	0.000	2	Unscraped	(1.322±0.043) days
				Scraped	(1.386±0.043) days
				Control	(0.811±0.043) days
Pupal duration	399.633	0.000	2	Unscraped	(1.672±0.022) days
				Scraped	(1.147±0.022) days
				Control	(1.107±0.022) days

Figure 1: Effect of scraping off of proximal portion of amputation site on regeneration in comparison to the unscraped larvae of *Cheilomenes sexmaculata*.

223
224

225