Pervasive nuclear envelope ruptures precede ECM signaling and disease onset without

activating cGAS-STING in Lamin-cardiomyopathy mice

Atsuki En[1,2], Hanumakumar Bogireddi[1], Briana Thomas[1], Alexis Stutzman[3], Sachie Ikegami[3], Brigitte

LaForest[3], Omar Almakki[3], Peter Pytel[4], Ivan P. Moskowitz[3,4,5], Kohta Ikegami[1,6,7]

[1]Division of Molecular Cardiovascular Biology, Cincinnati Children's Hospital Medical Center, Cincinnati,

OH, USA

[2]Graduate School of Nanobioscience, Yokohama City University, Yokohama, Kanagawa, Japan

[3]Department of Pediatrics, The University of Chicago, Chicago, IL, USA

[4]Department of Pathology, The University of Chicago, Chicago, IL, USA

[5]Department of Human Genetics, The University of Chicago, Chicago, IL, USA

[6]Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH, USA

[7]Corresponding author

SUMMARY

Nuclear envelope (NE) ruptures are emerging observations in Lamin-related dilated cardiomyopathy, an adult-onset disease caused by loss-of-function mutations in Lamin A/C, a nuclear lamina component. Here, we tested a prevailing hypothesis that NE ruptures trigger pathological cGAS-STING cytosolic DNA-sensing pathway, using a mouse model of Lamin-cardiomyopathy. Reduction of Lamin A/C in cardiomyocytes of adult mice caused pervasive NE ruptures in cardiomyocytes, preceding inflammatory transcription, fibrosis, and fatal dilated cardiomyopathy. NE ruptures were followed by DNA damage accumulation without causing immediate cardiomyocyte death. However, cGAS-STING-dependent inflammatory signaling remained inactive. Deleting *cGas* or *Sting* did not rescue cardiomyopathy. The lack of cGAS-STING activation was likely due to the near absence of cGAS expression in adult cardiomyocytes at baseline. Instead, extracellular matrix (ECM) signaling was activated and predicted to initiate pro-inflammatory communication from Lamin-reduced cardiomyocytes to fibroblasts. Our work nominates ECM signaling, not cGAS-STING, as a potential inflammatory contributor in Lamin-cardiomyopathy.

INTRODUCTION

Lamin A/C (*LMNA*) are nuclear lamina proteins that provide structural integrity to the nuclear envelope [1–4]. Mutations in *LMNA* cause a spectrum of degenerative disorders, collectively called laminopathies, including frequent dilated cardiomyopathy (*LMNA*-related DCM) [5–10]. *LMNA*-related DCM is prevalent adult-onset DCM [11,12] that accompanies cardiac conduction disease, fibrosis, heart failure, and mortality [13–16]. *LMNA*-related DCM is predominantly caused by heterozygous loss-of-function *LMNA* mutations that cause Lamin A/C protein reduction most strongly in cardiomyocytes [16–20]. Activation of stress response-related signaling, such as MAPK signaling [21–24], mTOR signaling [25], and DNA damage response signaling [26,27], has been reported in *LMNA*-related DCM models. However, the direct pathological alterations caused by Lamin A/C reduction in cardiomyocytes remain undefined.

Reduction of nuclear lamins can cause ruptures of the nuclear envelope (NE ruptures) [28–31,33–35]. Indeed, NE ruptures have been observed in patients [17,18,32,36,37], animal models [30,31,38–42], and cell culture models [30,35] of *LMNA*-related DCM and laminopathies. However, the extent of NE ruptures in *LMNA*-related DCM and the contribution of this event to the pathogenesis of DCM remain undefined.

A prevailing hypothesis for *LMNA*-related DCM pathogenesis implicates the cGAS-STING pathway as a link between NE rupture, inflammatory signaling, and DCM pathogenesis [39,40,43–45]. cGAS-STING is a cytosolic DNA-sensing innate immune mechanism, in which cGAS binds to cytosolic DNA and activates cell-autonomous STING-mediated interferon transcription [46,47]. cGAS-STING pathway activation has been observed in a cellular model of progeria [48] and *Lmna*-knockout developing hearts [27]. cGAS-STING activation has also been reported in mouse models of other heart diseases [49–52]. However, a direct link between NE ruptures and the cGAS-STING activation in *LMNA*-related DCM remains undefined. In this study, we report pervasive NE ruptures prior to DCM development and provide evidence against the hypothesis that cGAS-STING activation contributes to *LMNA*-related DCM. Instead, our study nominates a role for extracellular matrix signaling from cardiomyocytes in this disease.

RESULTS

Lamin A/C reduction in cardiomyocytes causes dilated cardiomyopathy in adult mice

We generated a mouse model of *LMNA*-related DCM using cardiomyocyte-specific deletion of *Lmna* (*Lmna*^{F/F};*Myh6-MerCreMer*; *Lmna*^{CKO} hereafter) and littermate control wild-type mice (*Lmna*^{+/+};*Myh6-MerCreMer*) (**Fig. S1A, B**). *Lmna* deletion in cardiomyocytes was induced by tamoxifen administration to adult mice at 6-8 weeks of age (**Fig. 1A**). Lamin A and Lamin C proteins were 47% and 43% reduced, respectively, in cardiomyocytes of *Lmna*^{CKO} mice at 2 weeks post tamoxifen, consistent with the 4-week half-life of Lamin A/C in mouse hearts [53] (**Fig. 1B; Fig. S1C**). The reduced, but persistent, Lamin A/C in individual cardiomyocytes was confirmed at 1, 2, and 3.5 weeks post tamoxifen (**Fig. 1C**). Thus, *Lmna*^{CKO} mice reproduced Lamin A/C protein insufficiency, not elimination, observed in cardiomyocytes of *LMNA*-related DCM patients [18,20].

Lmna^{CKO} mice developed progressive DCM as reported previously (**Fig. 1D, E; Fig. S1D**)[38]. Hearts of *Lmna*^{CKO} mice were indistinguishable from wild-type hearts in gross morphology and contractile activity until 2 weeks post tamoxifen. However, at 3 weeks post tamoxifen, left ventricular ejection fraction, an indicator of cardiac systolic function, significantly diminished (**Fig. 1E**). At 3.5 weeks post tamoxifen, *Lmna*^{CKO} hearts became severely dilated, with a significant reduction of ventricular wall thickness and loss of systolic activity. *Lmna*^{CKO} mice invariably died between 3.5 weeks and 4.5 weeks post tamoxifen (**Fig. 1F**). Thus, the modest reduction of Lamin A/C in cardiomyocytes was sufficient to cause DCM that progressed to heart failure, recapitulating a crucial aspect of human *LMNA*-related DCM.

Lamin A/C reduction in cardiomyocytes activates inflammation-related transcription

To identify the onset of molecular changes preceding the functional decline of *Lmna*^{CKO} hearts, we profiled the transcriptome of the heart at Days 7, 11, and 14 post tamoxifen by RNA-seq (**Table S1**). At Days 7 and 11, the transcriptome of *Lmna*^{CKO} hearts was almost identical to that of wild-type hearts (**Fig. 1G**). However, at Day 14, *Lmna*^{CKO} hearts exhibited strong transcriptional upregulation (1,751 genes) and modest downregulation (351 genes). The upregulated genes were highly overrepresented for gene ontology (GO) terms related to inflammatory responses (**Fig. 1H**). These upregulated genes included innate immune-related genes such as *Cxcl10*, *Ifit1*, and *Irf7* (**Fig. 1I**), which can be activated by the cGAS-STING pathway. The downregulated genes were overrepresented for gene pathways for cardiomyocyte metabolism and function (**Fig. 1H**).

We examined whether the upregulation of inflammation-related genes reflected an inflammatory response in *Lmna*^{CKO} hearts. We observed that the CD45+CD68+ macrophage population increased in *Lmna*^{CKO} hearts at 2 weeks post tamoxifen (**Fig. 1J; Fig. S1E**). We also observed increased interstitial collagen deposition at 2 weeks post tamoxifen, which developed into extensive fibrosis by 3.5 weeks post tamoxifen (**Fig. 1K**). Thus, Lamin A/C reduction in cardiomyocytes resulted in extensive upregulation of

inflammation-related genes, macrophage expansion, and initial fibrotic response at 2 weeks post
tamoxifen, preceding the morphological and functional changes reflective of clinical DCM (**Fig. 1L**).

***Lmna^CKO* cardiomyocytes develop pervasive localized nuclear envelope ruptures**
We examined whether Lamin A/C reduction caused nuclear envelope (NE) ruptures in cardiomyocytes.
Strikingly, we observed protrusion of DNA from one or multiple nuclei inside cardiomyocytes in *Lmna^CKO*
heart sections at 2 weeks post tamoxifen, while this event was absent in wild-type hearts (**Fig. 2A**; of
note, 90% of cardiomyocytes are binucleated in adult mice [54]). Nuclei with protruded DNA were positive
for Lamin A/C staining, indicating that partial reduction of Lamin A/C was sufficient to cause DNA
protrusion (**Fig. 2A**). Electron microscopy revealed strong condensation of protruded DNA that appeared
devoid of the surrounding nuclear membrane, suggesting NE ruptures (**Fig. 2B; Fig. S2A**). We isolated
cardiomyocytes from hearts at 2 weeks post tamoxifen, immediately fixed them, and examined the
nuclear envelope. Lamin A/C signals were specifically lost at the tips of nuclei from which DNA protruded
in *Lmna^CKO* cardiomyocytes (**Fig. 2C**). The Lamin A/C-lost locations also lost staining for PCM1, a
perinuclear matrix protein localized at the cytoplasmic side of the outer nuclear membrane [55] (**Fig. 2D**).
The loss of Lamin A/C and PCM1 occurred specifically at the tips of the elongated nuclei positioned along
the longitudinal axis of the cardiomyocytes (**Fig. 2A–D; Fig. S2A**). The specific loss of nuclear envelope
proteins Lamin A/C and PCM1 at nuclear tips suggested localized NE ruptures in *Lmna^CKO*
cardiomyocytes.

To determine whether NE ruptures occurred in *Lmna^CKO* cardiomyocytes, we examined the
retention of nuclear-localized tdTomato. We expressed nuclear localization signal (NLS)-fused tdTomato
in cardiomyocytes from a transgene *in vivo* and investigated tdTomato localization immediately after
cardiomyocyte isolation at 2 weeks post tamoxifen. NLS-tdTomato was localized exclusively to the
nucleus in wild-type cardiomyocytes as well as to the intact nuclei in *Lmna^CKO* cardiomyocytes (**Fig. 2E;**
Fig. S2B). However, NLS-tdTomato signals significantly diminished in nuclei with local PCM1 loss in
Lmna^CKO cardiomyocytes (**Fig. 2E, F**). Reciprocally, NLS-tdTomato signals increased in the cytoplasm of
Lmna^CKO cardiomyocytes with PCM1-lost nuclei. NLS-tdTomato leakage from damaged nuclei indicated
that the nuclear envelope had ruptured in *Lmna^CKO* cardiomyocytes.

To quantify NE rupture events, we expressed GFP-tagged catalytically-inactive cGAS (icGAS) in
cardiomyocytes *in vivo* using muscle-tropic adeno-associated virus (MyoAAV) [56] (**Fig. 2G**). icGAS is
widely used as a "NE rupture marker" owing to the cGAS's affinity to cytoplasmic DNA [57,58]. We observed
strong icGAS punctum at every PCM1-lost site on the nuclear envelope in *Lmna^CKO* cardiomyocytes at 2
weeks post tamoxifen (**Fig. 2H; Fig. S2C**). We found that 66% of *Lmna^CKO* cardiomyocytes have at least
one nucleus with icGAS puncta on nuclear envelopes, while icGAS puncta were virtually absent in wild-
type cardiomyocytes (**Fig. 2I**). Thus, NE ruptures were pervasive in *Lmna^CKO* cardiomyocytes at 2 weeks
post tamoxifen.

To determine whether NE ruptures preceded the transcriptional change, we quantified NE
ruptures at Day 11 post tamoxifen, when no transcriptional change was detected (**Fig. 1G**). We used a
local loss of PCM1 with concomitant cytoplasmic DNA protrusion as a proxy for NE ruptures, as this
feature strongly correlated with icGAS localization. We observed that 31% of *Lmna*[CKO] cardiomyocytes
had at least one ruptured nucleus at Day 11 (**Fig. 2J; Fig. S2D, E**). At Day 14, the NE rupture frequency
was 54% based on this method, which was comparable to 66% based on the sensitive icGAS-based
quantification (**Fig. 2J**). Thus, about one third of *Lmna*[CKO] cardiomyocytes developed NE ruptures as
early as Day 11 post tamoxifen, preceding the earliest transcriptional changes, with more than 50% of
nuclei presenting ruptures by Day 14.

We investigated whether the pervasive NE ruptures in *Lmna*[CKO] cardiomyocytes accompanied
DNA damage accumulation (**Fig. 2K; Fig. S2F**). We did not find a statistically significant increase of
gamma-H2AX-positive nuclei indicative of DNA double-strand break in *Lmna*[CKO] hearts at 2 weeks post
tamoxifen. In addition, there was no indication of increased cell death assessed by TUNEL at this time
point (**Fig. 2L; Fig. S2G**). However, at 3.5 weeks post tamoxifen, gamma-H2AX-positive nuclei
significantly increased. Thus, pervasive NE ruptures did not cause immediate DNA damage accumulation
or cell death at the time of strong inflammation-related transcriptional upregulation. Instead, DNA damage
accumulated when the heart structure and function began to deteriorate (**Fig. 2M**).

***Lmna*[CKO] cardiomyocytes do not activate cGAS-STING-related transcription**
Given the pervasive NE ruptures, we tested whether the cGAS-STING cytosolic DNA sensing pathway
was activated within *Lmna*[CKO] cardiomyocytes. In the cGAS-STING pathway, cGAS detection of
cytoplasmic DNA activates STING-mediated interferon transcription within cells bearing the cytoplasmic
DNA [46,47]. This pathway has been implicated in Lamin A/C-based cardiomyopathy in a previous study [27].
cGAS and STING proteins were expressed in the heart, but interestingly, their expression within
cardiomyocytes was very low in both wild-type and *Lmna*[CKO] backgrounds (**Fig. 3A; Fig. S3A, B**). To test
whether cGAS-STING-downstream genes were upregulated within cardiomyocytes of *Lmna*[CKO] hearts,
we distinguished cardiomyocyte and non-cardiomyocyte transcriptomes in the heart using SLAM-IT-seq
[59]. SLAM-IT-seq labels transcripts in *Cre*-positive cardiomyocytes with 4-thiouracil (4sU) and
distinguishes them from unlabeled transcripts from non-cardiomyocyte cells (**Fig. S3C**). As expected,
cardiomyocyte-specific genes were specifically 4sU-labeled in wild-type hearts (**Fig. S3D**). We then
assessed the labeling state of the 1,020 upregulated genes in *Lmna*[CKO] hearts and found that 135 genes
were 4sU-labeled (i.e. cardiomyocyte-derived) and 885 genes were not 4sU-labeled (i.e. non-
cardiomyocyte-derived) in *Lmna*[CKO] hearts at 2 weeks post tamoxifen (**Fig. 3B**). The cardiomyocyte-
originated upregulated genes were not overrepresented for inflammation-related GO terms (**Fig. 3B**) and
did not include well-established cGAS-STING downstream genes (**Fig. S3E**; **Table S2**). Instead, these
genes were overrepresented for the ECM-related GO terms such as "Proteoglycans in cancer". On the

other hand, non-cardiomyocyte-originated upregulated transcripts were strongly enriched for

inflammation-related GO terms (**Fig. 3B**). These data suggested that the cGAS-STING pathway was not

activated within *Lmna*^{CKO} cardiomyocytes and that the inflammation-related transcriptional activation

originated from non-cardiomyocytes in *Lmna*^{CKO} hearts.

We hypothesized that the near absence of cGAS expression prevented cGAS-STING activation

within *Lmna*^{CKO} cardiomyocytes. We tested this hypothesis by overexpressing wild-type cGAS in

cardiomyocytes *in vivo* with MyoAAV (**Fig. 3C**). We first confirmed that endogenous cGAS was not

detectable at NE rupture sites (**Fig. 3D**) and that the well-established cGAS-STING downstream genes,

Cxcl10, *Ifit3*, and *Ifnb1* [60–63], remained transcriptionally silent in cardiomyocytes of *Lmna*^{CKO} mice

transduced with control luciferase MyoAAV (**Fig. 3E**). However, cGAS overexpression in *Lmna*^{CKO}

cardiomyocytes resulted in strong cGAS accumulation at NE rupture sites (**Fig. 3D**). Consistently, *Cxcl10*,

Ifit3, and *Ifnb1* transcripts were 10-, 128-, and 6-fold upregulated, respectively, upon cGAS

overexpression (**Fig. 3E**). In wild-type mice, cGAS overexpression did not upregulate cGAS-downstream

genes, confirming the NE rupture-dependency of cGAS-downstream activation upon cGAS

overexpression. Taken together, these data suggested that *Lmna*^{CKO} cardiomyocytes did not activate the

cGAS-STING pathway despite pervasive NE ruptures and that the lack of cGAS-STING activation was

likely due to the low cGAS expression in adult cardiomyocytes.

***Cgas* and *Sting* are not required for inflammation-related gene expression and DCM in *Lmna*^{CKO}**

mice

We examined the genetic requirement for *Cgas* and *Sting*, the essential mediators of the cGAS-STING

pathway, to the inflammation-related transcriptional activation and DCM in *Lmna*^{CKO} hearts. We

introduced the germ-line *Cgas*$^{−/−}$ allele [64] into *Lmna*^{CKO} mice or control *Lmna*^{Het} mice

(*Lmna*$^{F/+}$*;Myh6MerCreMer*) and validated the absence of cGAS protein in the heart of *Cgas*$^{−/−}$ carriers

(**Fig. S3A**). We then performed RNA-seq in whole hearts at 2 weeks post tamoxifen. We first confirmed

the expected strong upregulation of inflammation-related genes in *Cgas*$^{+/+}$;*Lmna*^{CKO} hearts compared

with *Cgas*$^{+/+}$;*Lmna*^{Het} hearts (**Fig. 3F; Fig. S3F; Table S1**). We then compared *Cgas*$^{−/−}$;*Lmna*^{CKO} hearts

with *Cgas*$^{+/+}$;*Lmna*^{CKO} hearts. There was almost no difference in gene expression between them (**Fig.**

3F). Representative cGAS-STING-downstream genes did not show statistically significant differences in

gene expression between *Cgas*$^{−/−}$;*Lmna*^{CKO} and *Cgas*$^{+/+}$;*Lmna*^{CKO} hearts (**Fig. 3G; Fig. S3G**). These data

suggested that the inflammation-related transcription in *Lmna*^{CKO} hearts was independent of cGAS.

We next combined germ-line *Sting*$^{−/−}$ mice [65] with *Lmna*^{CKO} mice (**Fig. S3B**) and investigated the

heart phenotype. By 3.5 to 4 weeks post tamoxifen, *Sting*$^{−/−}$;*Lmna*^{CKO} mice showed as severely reduced

left ventricular ejection fraction (**Fig. 3H**) and extensive fibrosis (**Fig. 3I**) as the littermate control

Sting$^{+/+}$;*Lmna*^{CKO} mice did, whereas *Sting*$^{+/+}$;*Lmna*^{Het} and *Sting*$^{−/−}$;*Lmna*^{Het} mice maintained normal hearts.

Likewise, *Cgas*$^{−/−}$;*Lmna*^{CKO} hearts developed severe fibrosis (**Fig. S3H**). Moreover, both *Sting*$^{−/−}$;*Lmna*^{CKO}

mice and $Cgas^{-/-};Lmna^{CKO}$ mice died as early as $Sting^{+/+};Lmna^{CKO}$ or $Cgas^{+/+};Lmna^{CKO}$ mice did, whereas

no lethality was observed in $Sting^{-/-};Lmna^{Het}$ and $Cgas^{-/-};Lmna^{Het}$ mice (**Fig. 3J; Fig. S3I**). Taken

together, our data suggested that the cGAS-STING pathway did not mediate inflammatory transcription

and the pathophysiologic features of DCM in $Lmna^{CKO}$ hearts (**Fig. 3K**).

Cardiomyocytes, fibroblasts, and immune cells are transcriptionally altered in $Lmna^{CKO}$ hearts

We explored alternative mechanisms by which Lamin A/C reduction in cardiomyocytes caused DCM. To

identify intercellular signaling initiated by $Lmna^{CKO}$ cardiomyocytes, we defined the cell type-resolved

transcriptional changes using single-nucleus (sn) RNA-seq (**Table S3**). We obtained transcriptomes for

14,111 nuclei from either wild-type or $Lmna^{CKO}$ hearts at 2 weeks post tamoxifen (n=3; **Fig. 4A**). These

nuclei were grouped into cardiomyocytes (CM1 and CM2), fibroblasts (Fib1 and Fib2), macrophages

(Mac1 and Mac2), T cells, endothelial cells (coronary, lymphatic, and other), pericytes, or neurons, based

on their transcriptomes (**Fig. S4A**). Among these, CM2, Fib2, Mac2, and T-cells were highly

overrepresented by $Lmna^{CKO}$ heart-derived cells (**Fig. 4B; Fig. S4B**). CM2 was characterized by high

Rtn4 expression, a feature also reported in *Lmna*-knockout myotubes [66] and Lamin A/C-independent

heart disease [67–70] (**Fig. S4A**). Fib2 was characterized by myofibroblast marker *Postn* [71], Mac2 by

circulating monocytic receptor *Ccr2* [72], and T cells by T cell adaptor *Skap1* [73]. We found extensive

differential gene expression within cardiomyocytes (CM1 and CM2 combined), fibroblasts (Fib1 and Fib2

combined), and immune cells (Mac1, Mac2, T cell combined) while other cell types exhibited almost no

expression differences (**Fig. 4C, D; Fig. S4C; Table S4**). The upregulated genes in cardiomyocytes were

not overrepresented for cytosolic DNA sensing pathway, did not include cell death-related genes, but

included some DNA damage repair-related genes, as expected (**Fig. S4D–F**).

Interestingly, upregulated genes in $Lmna^{CKO}$ cardiomyocytes and those in fibroblasts were both

most strongly overrepresented for ECM-related pathways (**Fig. 4D**). The upregulated ECM-related genes

included collagen and integrin genes (in cardiomyocytes and fibroblasts) and ECM proteinase and

crosslinker genes (specifically in fibroblasts) (**Fig. 4E**). On the other hand, fibroblasts and immune cells,

but not cardiomyocytes, commonly upregulated immune-related genes including cytokine, cytokine

receptor, and cytokine transcription factor genes (**Fig. 4D, F**). These results suggested potential signal

crosstalk between cardiomyocytes and fibroblasts and between fibroblasts and immune cells in $Lmna^{CKO}$

hearts.

ECM-mediated signaling from $Lmna^{CKO}$ cardiomyocytes is predicted to activate fibroblasts

We investigated intercellular signal crosstalk in $Lmna^{CKO}$ hearts by applying the CellChat program [74] to

the snRNA-seq data. This analysis predicted ECM-receptor signaling from CM2 to Fib2 as the only

strongly upregulated signaling originating from CM2, the population predominantly consisted of $Lmna^{CKO}$

cardiomyocyte nuclei (**Fig. 4G, Box a**). Collagens and laminins were predicted to mediate this CM2-to-

Fib2 signaling (**Fig. 4H**). CM2 were not predicted to signal directly to immune cell populations. Instead,

Fib2 were predicted to send ECM-mediated signals to CM2 (**Box b**), Fib2 (**Box c**), and T cells (**Box d**),

via collagens, laminins, thrombospondins, and fibronectins (**Fig. 4G, H**). Fib2 were also predicted to send

secretion-mediated signals to Fib2 (**Box e**), and Mac2 (**Box f**), via ANGPTL (angiopoietin-like), TGF beta,

BMP, and SEMA3 (semaphorin3). Fib2 might therefore act as a central signaling hub, receiving signals

from CM2 and sending signals to immune cells (**Fig. 4I**). Finally, Mac2 were predicted to send secretion-

mediated signals to Mac2 themselves (**Box g**) via CCL (chemokine ligands), SPP1 (osteopontins), and

complement factors. This signaling might mediate the macrophage infiltration observed in $Lmna^{CKO}$

hearts (**Fig. 1J**). Taken together, our results suggested that $Lmna^{CKO}$ cardiomyocytes activated

fibroblasts via ECM-mediated signaling, not cytokine-mediated signaling such as cGAS-STING, and that

fibroblasts activated and recruited immune cells through both ECM signaling and secretion signaling to

orchestrate inflammatory signaling in the heart (**Fig. 4J**).

DISCUSSION

We report that frequent localized NE ruptures precede transcriptional changes, DNA damage accumulation, and heart functional decline in *Lmna*^{CKO} hearts. Given our observation that a 50% reduction of Lamin A/C proteins is sufficient to induce pervasive NE ruptures in cardiomyocytes, NE ruptures might be more prevalent than previously anticipated in *LMNA*-related DCM. NE ruptures might have been overlooked due to the specific location of localized NE ruptures at the tips of elongated nuclei within cardiomyocytes. The mechanisms underlying NE ruptures and the causal role of NE ruptures in DCM remain to be investigated.

Our data indicate that the cGAS-STING pathway does not contribute to *LMNA*-related DCM in the adult mouse model investigated herein. This suggests that pharmacologically reducing cGAS-STING activity for treating *LMNA*-related DCM, an idea previously discussed [27,75], may be ineffective. Our conclusion diverges from a previous report that *Cgas* deletion can delay early postnatal cardiomyopathy caused by *Lmna* deletion in embryonic hearts [27]. This difference may be due to that the near absence of cGAS expression prevents cGAS-STING activation in adult cardiomyocytes, whereas the basal cGAS-STING activity may be higher in early postnatal cardiomyocytes [76,77]. The lack of robust cGAS expression may reduce pathogen-derived DNA sensing in adult cardiomyocytes, similar to hepatocytes that also lack functional cGAS-STING [78]. Many reports suggest an association between DNA damage and cGAS-STING activation [27,79–82]. Our data that *Lmna*^{CKO} cardiomyocytes accumulate DNA damage without activating cGAS-STING suggest that DNA damage and cGAS-STING can be uncoupled.

Our study nominates ECM signaling as a mechanism by which Lamin A/C-reduced cardiomyocytes initiate an inflammatory response by activating fibroblasts. A recent study reported a similar observation in which NE rupture induces ECM remodeling through unknown mechanisms that require DNA damage [83]. Further study on the potential roles of ECM signaling in *LMNA*-related DCM is required.

Limitations of the study

We reduced Lamin A/C exclusively within the cardiomyocytes of adult mice. Whether Lamin A/C reduction in other cell types or at other stages causes NE ruptures and contributes to DCM through cGAS-STING activation remains an open question.

ACKNOWLEDGEMENT

We thank core facilities for animal care, microscopy, genomics, and pathology at Cincinnati Children's

and the University of Chicago for their assistance. This work is supported by NIH grant R21/R33

AG054770 (K.I.), Cincinnati Children's Research Innovation and Pilot grant (K.I.), NIH grants R01

HL163523, R01 HL124836, and R01 HL126509 (I.P.M).

AUTHOR CONTRIBUTION

Conceptualization, A.E. and K.I.; Methodology, A.E., I.P.M., and K.I.; Formal Analysis, A.E. and K.I.;

Investigation, A.E., H.B., B.T., A.S., S.I., B.L., O.A., P.P., K.I.; Resources, I.P.M.; Data Curation, A.E. and

K.I; Writing – Original Draft, A.E., and K.I.; Writing – Review & Editing, A.E., I.P.M., and K.I.; Visualization,

A.E. and K.I.; Supervision, K.I.; Project Administration, K.I.; Funding Acquisition, K.I. and I.P.M.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURE LEGENDS

Figure 1. Lamin A/C reduction in cardiomyocytes causes dilated cardiomyopathy in adult mice

A) Tamoxifen (Tam) induces *Lmna* deletion in cardiomyocytes in adult mice. See **Fig. S1** for related analyses.

B) Lamin A/C immunoblot in isolated cardiomyocytes (top) with signal quantification (bottom).

C) Lamin A/C immunohistochemistry (brown) with hematoxylin counterstaining (blue) in mouse heart tissues. Scale bar: 20 µm.

D) Hematoxylin-eosin staining of hearts. Scale bar: 1 mm.

E) Left ventricular ejection fraction with interquartile range (box) measured by echocardiography.

F) Kaplan-Meier survival analysis.

G) Volcano plot comparing RNA-seq read counts in *Lmna*^{CKO} versus WT hearts. Up, upregulated gene count. Down, downregulated gene count.

H) The top 10 Gene Ontology terms overrepresented among differentially expressed genes.

I) Normalized RNA-seq gene expressed levels for immune-related genes (bar, mean).

J) Immunofluorescence of heart tissue sections for CD45 (pan-leukocyte, red) and CD68 (macrophage, green). Scale bar: 20 µm.

K) Masson's trichrome staining of heart sections. Arrow, collagen deposition. Scale bar: 20 µm.

L) Summary

Figure 2. Lamin A/C reduction causes localized nuclear envelope rupture in cardiomyocytes

A) Lamin A/C immunohistochemistry in heart tissue. Arrow, DNA protruded from nuclei. Scale bar: 20 µm. See **Fig. S2** for related analyses.

B) Top: Transmission electron micrograph of heart sections focusing on cardiomyocytes. Bottom: Close-up image of the area indicated by rectangle in upper panel. Arrow, protruded chromatin. Scale bar: 1 µm.

C) Immunofluorescence for Lamin A/C in isolated cardiomyocytes. Phalloidin stains F-actin. Arrowhead, local loss of Lamin A/C with protruded DNA. Scale bar: 20 µm.

D) Same as **(C)**, but for PCM1.

E) NLS-tdTomato signals in isolated cardiomyocytes co-stained for PCM-1. Scale bar: 20 µm.

F) NLS-tdTomato signal intensity in nucleus and cytoplasm of cardiomyocytes (box, interquartile range). *Lmna*^{CKO} cardiomyocytes are stratified by the presence or absence of local PCM-1 loss at the nuclear envelope. WT n=51, *Lmna*^{CKO} n=53 (PCM1-loss 34, intact 19) cardiomyocytes.

G) MyoAAV-mediated GFP-icGAS expression in cardiomyocytes *in vivo*.

H) GFP-icGAS signals in isolated cardiomyocytes co-stained for PCM1. Scale bar: 20 µm.

I) Percentage of cardiomyocytes with GFP-icGAS punctum at nuclear tip (bar, mean).

J) Percentage of cardiomyocytes with DNA protrusion from PCM1-lost nuclear tip (bar, mean) at 11 days (left) and 14 days (right) post tamoxifen.

K) Percentage of gamma-H2AX-positive nuclei in heart section (bar, mean).

L) Number of TUNEL-positive cells in heart section (bar, mean).

M) Summary

Figure 3. *Cgas* and *Sting* are not required for inflammation-related gene expression and dilated
cardiomyopathy in *Lmna^{CKO}* mice

A) Normalized cGAS and STING abundance quantified by immunoblot shown in **Fig. S3A, B** (bar, mean).

B) Left: Upregulated genes in *Lmna^{CKO}* hearts at 2 weeks post tamoxifen are categorized into
cardiomyocyte-derived or non-cardiomyocyte-derived based on 4sU labeling in SLAM-IT-seq. Right: GO
terms enriched among each class of upregulated genes.

C) MyoAAV-mediated GAS overexpression in cardiomyocytes *in vivo*.

D) Top: Anti-cGAS and Lamin A/C immunofluorescence in isolated cardiomyocytes. Bottom: Close-up
image of the area indicated by rectangle in upper panel. Arrowhead, GFP-icGAS localization at nuclear
tip. Scale bar: 20 µm.

E) Transcript abundance in isolated cardiomyocytes quantified by quantitative PCR and normalized by
Actb transcripts (bar, mean).

F) Volcano plot comparing RNA-seq read counts in *Cgas$^{-/-}$*; *Lmna^{Het}* (n=4) versus *Cgas$^{+/+}$*; *Lmna^{Het}* (n=6)
(left) and in *Cgas$^{-/-}$*; *Lmna^{CKO}* (n=4) versus *Cgas$^{+/+}$*; *Lmna^{CKO}* (n=6) (right).

G) Normalized RNA-seq gene expressed levels for cGAS-STING-related genes (bar, mean).

H) Left ventricular ejection fraction measured by echocardiography (box, interquartile range).

I) Masson's trichrome staining of heart sections. Scale bar: 20 µm.

J) Kaplan-Meier survival analysis.

K) Summary

Figure 4. ECM-mediated signaling from *Lmna^{CKO}* cardiomyocytes is predicted to activate
fibroblasts

A) snRNA-seq UMAP plot for 14,111 nuclei from either WT or *Lmna^{CKO}* hearts, colored by cell-type
annotation. CM, cardiomyocyte. Fib, fibroblast. Mac, macrophage. EC, endothelial cell. CEC, coronary
EC. LEC, lymphatic EC. PC, pericyte. See **Fig. S4** for additional analyses.

B) Count of WT and *Lmna^{CKO}* heart-originating nuclei within each cell type cluster. Cell types combined
in later analyses are indicated below graph.

C) Pseudo-bulk volcano plot comparing sbRNA-seq read counts in cells of *Lmna^{CKO}* hearts versus cells
of WT hearts within each cell type. Red, upregulated genes. Blue, downregulated genes.

D) GO terms enriched among upregulated genes within indicated cell type. Red arrows, ECM-related GO
term. Blue arrow, inflammation-related GO term.

E) Pseudo-bulk transcript abundance of ECM-related genes. Point, mean of normalized read count per

replicate. Bar, mean across replicates.

F) Same as (E), but for immune-related genes.

G) Predicted signaling strength between sender cells (x-axis) and receiver cells (y-axis). Color indicates

the extent of signaling gains in $Lmna^{CKO}$ hearts relative to WT hearts.

H) Predicted signaling mediators for the signaling indicated by box in (**G**) with alphabetical labels.

I) Predicted intercellular signaling (indicated in **G** by alphabet labels) in $Lmna^{CKO}$ hearts.

J) Model

 SUPPLEMENTARY FIGURE LEGENDS

Figure S1. Characterization of adult mice with cardiomyocyte-specific *Lmna* deletion

A) Immunofluorescence for PCM1, NLS-tdTomato, and COX4 in heart sections. PCM1 and COX4 are

used to stain cardiomyocyte nuclei and the cytoplasm, respectively. Scale bar: 20 µm.

B) Left: Schematic for the *Lmna* LoxP allele and the expected recombined allele upon Cre expression.

Arrow, PCR primer location. Right: Amplification of the wild-type, floxed, and recombined *Lmna* alleles

by PCR using primers indicated in the left panel. The recombination efficiency (%) was calculated by the

intensity of the 153-bp recombined band over the intensity of the 598-bp floxed band.

C) The whole immunoblot for **Fig. 1B**, with Coomassie staining of the original gel.

D) Echocardiography of left ventricular (LV) geometry. P, Wilcoxon test.

E) Quantification of CD45 and CD68-positive cells in heart sections. Representative image in **Fig. 1J**.

Figure S2. Characterization of nuclear envelope ruptures in *Lmna^{CKO}* cardiomyocytes

A) Top: Additional transmission electron micrographs of heart sections focusing on cardiomyocytes,

related to **Fig. 2B**. Bottom: Enlarged images of the area indicated by yellow box in the upper panels.

Arrow, protruded chromatin. Scale bar: 1 µm.

B) Additional immunofluorescence images for PCM-1 with detection of native NLS-tdTomato. Scale

bar: 20 µm.

C) Immunofluorescence for PCM-1 with detection of exogenous GFP-icGAS in isolated cardiomyocytes,

related to **Fig. 2H**. Phalloidin stains F-actin. Arrowhead, GFP-icGAS localization at nuclear tips. Scale

bar: 20 µm.

D) Immunofluorescence images for PCM1 in isolated cardiomyocytes at Day 11 post tamoxifen.

Arrowhead, local loss of PCM1 at nuclear tips. Scale bar: 20 µm.

E) Fraction of cardiomyocytes with DNA protrusion from PCM1-lost nuclear tips. Cardiomyocytes are

stratified by the number of ruptured nuclei per cell. Only multinucleated cardiomyocytes are analyzed.

F) Immunofluorescence for γH2AX and desmin in heart tissue sections. Desmin stains cardiomyocytes.

Arrowhead, γH2AX-stained nuclei. Scale bar: 20 µm.

G) TUNEL assay for cell death detection in heart tissues. Arrowhead, TUNEL-positive cells. Scale

bar: 20 µm.

Figure S3. Analysis of the cGAS-STING pathway in *Lmna^{CKO}* hearts

A) cGAS and GAPDH immunoblots, and Coomassie staining of original gel, in hearts and isolated

cardiomyocytes, related to **Fig. 3A**.

B) Same as (A), but for STING.

C) Principle of SLAM-IT-seq. Transcripts in cardiomyocytes are labeled by 4-thiouracil (4sU).

D) SLAM-IT-seq data for all genes in WT hearts at 2 weeks post tamoxifen, with the expression level on the X axis and the 4sU labeling frequency on the Y axis. Select genes known to be preferentially expressed in cardiomyocytes (red) and those not known to be preferentially expressed in cardiomyocytes (black) are indicated.

E) SLAM-IT-seq data for the 1,020 upregulated genes at 2 weeks post tamoxifen, with the rank order of the fold change of upregulation on the X axis and the 4sU labeling frequency in $Lmna^{CKO}$ hearts on the Y axis. All cytokine genes among the upregulated genes, which are all among unlabeled genes (blue), are indicated. Highly labeled genes (Log_{10} P-value > 3.5) are also indicated (orange).

F) The top 10 Gene Ontology terms overrepresented among differentially expressed genes in $Cgas^{-/-}$; $Lmna^{Het}$ versus $Cgas^{+/+}$; $Lmna^{Het}$.

G) Normalized RNA-seq gene expressed levels for additional cGAS-STING-related genes (bar, mean).

H) Masson's trichrome staining of heart sections. Scale bar: 20 μm.

I) Kaplan-Meier survival analysis.

Figure S4. Single-nucleus RNA-seq analysis of $Lmna^{CKO}$ hearts

A) Expression (sum of single-nucleus normalized read count across 3 mice within cell type) of maker genes used to classify single nuclei to the cell types indicated along the Y axis, related to **Fig. 4A.**

B) snRNA-seq UMAP plot for 14,111 nuclei, colored by sample genotype. Inset, UMAP colored by cell type for reference.

C) GO terms enriched among downregulated genes within indicated cell type.

D) Enrichment for select GO terms relevant to cytosolic DNA sensing among upregulated genes.

E) Expression (mean of single-nucleus normalized read count within cell type) of cell death-related genes. Dot, mean within individual mice. Bar, mean across three mice.

F) Same as (C), but for DNA damage repair-related genes.

SUPPLEMENTARY TABLES

Table S1.

Differentially expressed genes identified by bulk RNA-seq, related to **Figures 1 and 3**. The table includes differentially expressed genes between wild-type and $Lmna^{CKO}$ hearts at 1 week post tamoxifen, 11 days post tamoxifen, and 2 weeks post tamoxifen, and for $Cgas/Lmna$ double-knockout experiment.

Table S2.

Normalized expression values, C-to-T conversion rates, and beta-binomial test P-values for SLAM-IT-seq data, related to **Figure 3**.

Table S3.

Indexed oligonucleotides used in snRNA-seq, related to **Figure 4**, including indexed RT primers, indexed

ligation primers, and indexed P7 PCR primers.

Table S4.

Differentially expressed genes between cells of wild-type and $Lmna^{CKO}$ hearts, derived from pseudo-bulk

analysis of snRNA-seq data, related to **Figure 4**.

METHODS

Resource availability

Lead contact

Kohta Ikegami (email: Kohta.Ikegami@cchmc.org).

Materials availability

All mice are available at the sources specified in **Key Resources Table**. Plasmids produced in this study are provided upon request from the lead contact after material transfer agreements. Any information required to reanalyze the data reported in this paper is available from the lead contact.

Data availability

High-throughput sequencing data associated with this version of the manuscript are limited to reviewers and journal editors. All data will become available to the public after peer review.

Method details

Mouse genetics and treatment

Lmna-LoxP mice [84] were provided by Dr. Yixian Zheng at Carnegie Institution. *Myh6-MerCreMer* (Myh6MCM) transgene mice [85] (JAX stock No: 005657), $Rosa26^{CAG-LSL-tdTomato}$ ("Ai75") mice [86] (JAX stock No: 25106), *Cgas-null* mice [64] (JAX stock No: 026554), and *CAG-Lox-Stop-Lox-Uprt* transgene mice [87] ($Uprt^{Tg}$; JAX stock No: 021469) were obtained from the Jackson Laboratory. *Sting-null* mice were generated by breeding *Sting-flox* mice [65], obtained from the Jackson Laboratory (JAX stock No: 031670), with mice carrying the *Mef2C-AHF-Cre* transgene allele [88], provided by Dr. Brian L. Black at the University of California San Francisco. For cardiomyocyte-specific *Lmna* knockout experiments, $Lmna^{F/F};Myh6MCM^{Tg/0}$ ($Lmna^{CKO}$) and $Lmna^{+/+};Myh6MCM^{Tg/0}$ (wild-type control) mice were used. For the analysis of nuclear tdTomato intensity, $Lmna^{F/F};Myh6MCM^{Tg/0};Rosa26^{CAG-LSL-tdTomato/+}$ and $Lmna^{+/+};Myh6MCM^{Tg/0};Rosa26^{CAG-LSL-tdTomato/+}$ mice were used. For SLAM-IT-seq experiments, $Lmna^{F/F};Myh6MCM^{Tg/0};Uprt^{Tg/0}$, $Lmna^{+/+};Myh6MCM^{Tg/0};Uprt^{Tg/0}$, $Lmna^{F/F};Myh6MCM^{Tg/0};Uprt^{0/0}$, and $Lmna^{+/+};Myh6MCM^{Tg/0};Uprt^{0/0}$ mice were used. For *Cgas* deletion experiments, $Cgas^{-/-};Lmna^{F/F};Myh6MCM^{Tg/0}$ mice, $Cgas^{+/+};Lmna^{F/F};Myh6MCM^{Tg/0}$, $Cgas^{+/+};Lmna^{F/+};Myh6MCM^{Tg/0}$, and $Cgas^{-/-};Lmna^{F/+};Myh6MCM^{Tg/0}$ mice were used. For *Sting* deletion experiments, $Sting^{-/-};Lmna^{F/F};Myh6MCM^{Tg/Tg}$ mice, $Sting^{+/+};Lmna^{F/F};Myh6MCM^{Tg/Tg}$, $Sting^{+/+};Lmna^{F/+};Myh6MCM^{Tg/Tg}$, and $Sting^{-/-};Lmna^{F/+};Myh6MCM^{Tg/Tg}$ mice were used. All mice used in experiments were administered with tamoxifen (Sigma, T5648) at 6-8 weeks of age via intraperitoneal injections (100 µL of 4 mg/mL solution per day for 4 consecutive days) dissolved in corn oil (Sigma, C8267). All mice were used in mixed genetic backgrounds. All mouse experiments were approved by the Institutional Animal Care and Use Committee

493   (IACUC) at Cincinnati Children's Hospital under IACUC protocol 2021-0014 or at University of Chicago
494   under IACUC protocol 71730-10. All procedures were performed in compliance with institutional and
495   governmental regulations under PHS Animal Welfare Assurance number D16-00068 (Cincinnati
496   Children's) or D16-00322 (University of Chicago).
497

Mouse genotyping
For genotyping of the *Lmna* alleles, genomic DNA of isolated cardiomyocytes was extracted with
Phenol/Chloroform/Isoamyl alcohol (Invitrogen,15593-031), treated with RNaseA and Proteinase K, and
purified. Floxed, recombined, and wild type *Lmna* alleles were detected by PCR using the primers KI279
and KI280 (**Key Resources Table**). Based on the agarose gel (1.5%) analysis of PCR-amplified
products, recombination efficiency (%) was calculated by 153-bp band (Recombined) over 598-bp band
(Floxed) intensities. Genotyping of *Myh6-MerCreMer*, *Cgas*, *Sting*, *Uprt*, and *Rosa26::CAG-LSL-*
tdTomato alleles was performed at Transnetyx, Inc.
506

Echocardiography
The transthoracic echocardiography was performed using a Vevo 3100 LT (FUJI FILM VisualSonics) and
a transducer of 50-MHz MX-700. Parasternal long-axis view and short-axis view at the papillary muscle
level were imaged. Two-dimensional M-mode tracing was recorded at three or more consecutive cardiac
cycles. Data were analyzed using Vevo LAB Software Package V3.2.6. Echocardiographic parameters
between two groups were compared using Wilcoxon rank sum tests, and P-values were adjusted for
multiple testing using the Benjamini-Hochberg procedure in R.
514

515   **Survival analysis**
516   Kaplan-Meier survival analyses were performed using the *survival* package and visualized using the
517   *survminer* package in R. Log-rank tests were performed using the *survdiff* function in the survival
518   package.
519

520   **Histological staining**
521   Hearts were perfused with 100 mM KCl and fixed in 10% Formalin (Fisherbrand, 245-685). Fixed hearts
522   were embedded in paraffin and sectioned to a thickness of 5 μm. Heart sections were stained with
523   hematoxylin and eosin (H&E) or a Masson's Trichrome Kit (Thomas Scientific LLC, KTMTRPT) according
524   to the manufacturer's protocol.
525

Cardiomyocyte isolation
Mice were administered with heparin (100 Units, NDC 25021-400-10), anesthetized with Isoflurane, and
euthanized by cervical dislocation. Hearts were perfused with 100 mM KCl and moved to Tyrode's

solution (10 mM glucose, 5 mM HEPES, 5.4 mM KCl, 1.2 mM $MgCl_2$, 150 mM NaCl and 2 mM sodium

pyruvate, pH 7.35). Excised hearts were cannulated to the Langendorff retrograde perfusion system

through the aorta and perfused with Base solution (Tyrode's solution with 10 mM taurine and 12 mM

2,3-butanedione monoxime) and then digested with prewarmed Digestion buffer (Base solution with 180

U/mL Collagenase Type II (Worthington, LS004177) and 25 µM $CaCl_2$) for 20 minutes at 37^{o}C. Heart

tissue was then isolated, gently minced in Base solution containing 5 mg/mL BSA. Cell suspension was

filtrated through a 240 µm mesh. Cardiomyocytes settled to the bottom of conical tubes by gravity were

used in experiments.

Immunohistochemistry

Deparaffinized heart tissue sections, that had been fixed with 10% formalin, were subjected to heat-

induced antigen retrieval in Target Retrieval Solution (S2367, DAKO) for 5 minutes. Antigen-retrieved

sections were incubated with anti-Lamin A/C antibody Ab26300 (rabbit, Abcam) (1:400) for 1 hour at

room temperature, followed by biotinylated anti-rabbit IgG antibody for 30 minutes at room temperature.

The antigen-antibody binding was detected by Elite kit (PK-6100, Vector Laboratories) and DAB (DAKO,

K3468) system. Sections were counterstained by hematoxylin.

Immunofluorescence on tissue sections

Deparaffinized heart tissue sections, that had been fixed with 10% formalin, were subjected to heat-

induced antigen retrieval in 10 mM Tris EDTA buffer (pH 9.0) for 5 minutes. Antigen-retrieved sections

were incubated overnight at 4 °C with goat anti-CD45 antibody (R&D Systems, AF114-SP) and rabbit

anti-CD68 antibody (Cell Signaling, 97778), or rabbit anti-γ-H2A.X antibody (Cell Signaling, 9718) and

goat anti-desmin antibody (Invitrogen, PA5-19063), or rabbit anti-PCM1 antibody (Sigma, HPA023370),

goat anti-TdTomato antibody (OriGene, AB8181), and mouse anti-Cox4 antibody (R&D Systems,

MAB6980), followed by Alexa fluorophore-conjugated secondary antibodies for 1 hour at 37°C. Cells

were counterstained with DAPI (4′,6-diamidino-2-phenylindole), submerged in 0.25% Sudan Black B

(Electron Microscopy Slides, #21610) in 70% Isopropanol for 10 minutes, and then mounted with ProLong

Glass Antifade Mountant (Invitrogen, P36984). Fluorescence signals and imaging were acquired using a

Nikon A1R laser-scanning confocal microscope. CD45 and CD68 quantification of whole heart tissue was

performed using Nikon NIS-Elements; each genotype had five biological replicates. CD45/CD68 positive

cells were identified based on the overlap of the DAPI signal with CD45/CD68. Signal intensities between

two groups were compared using Wilcoxon rank sum tests in R. For γ-H2A.X quantification, nuclei with

DAPI were counted and those with at least one γ-H2A.X focus were quantified using CellProfiler. Five to

eight images per mouse (374 x 374 µm^2 per image) were used for the quantification.

Immunofluorescence on isolated cardiomyocytes

Isolated cardiomyocytes were fixed in 4% paraformaldehyde (Electron Microscopy Slides, #15710) in PHEM buffer (60 mM PIPES pH7.5, 25 mM HEPES pH7.5, 10 mM EGTA, 4 mM $MgSO_4$) for 10 minutes at 37°C, then washed with PBS and attached to coverslips with Cell-Tak Cell Adhesive (Sigma Aldrich, 354240). Cells were blocked and permeabilized with in a buffer containing 5% normal donkey serum (Jackson ImmunoResearch, 017-000-121), 1% non-fat milk, and 0.1% Triton X-100 in PBS for 1 hour at 37°C. Permeabilized cells were incubated overnight at 4°C with mouse anti-Lamin A/C antibody (Santa Cruz, sc-376248), rabbit anti-PCM-1 antibody (Sigma, HPA023370), mouse anti-cGAS antibody (Cell Signaling, D3O8O). Cells were washed and incubated with Alexa fluorophore-conjugated secondary antibodies and Alexa Fluor Plus 647 Phalloidin for 1 hour at 37°C. Cells were counterstained with DAPI and then mounted with ProLong Glass Antifade Mountant. Fluorescence signals were detected on Nikon A1R laser-scanning confocal microscope or Yokogawa CSU-W1 Sora spinning disk confocal microscope.

Plasmid construction

The expression vectors used in this study were derived from the pAAV:cTNT::Luciferase vector (gift from William Pu; Addgene plasmid # 69915) [89]. For cloning of pAAV:cTnT::cGAS and pAAV:cTnT::GFP vectors, we digested the pAAV:cTNT::Luciferase vector with *NheI* and *NotI*, and the luciferase gene was replaced with the cGAS or EGFP sequences using NEBuilder HiFi DNA Assembly Master Mix (NEB E2621). The pAAV:cTnT::GFP-icGAS vector was generated by inserting the catalytically-inactive cGAS (icGAS) sequence (DNA fragment ID: EA086, **Key Resources Table**) into the pAAV:cTnT::GFP vector digested with *NotI*. The icGAS sequence was derived from the human cGAS protein sequence with an E225A/D227A amino acid substitution that abolishes enzyme activity and interferon production, but retains DNA binding ability and functions, used as a NE rupture marker previously [57]. The DNA fragments for mouse wild-type *Cgas* (NM_173386.5) and *icGAS* were synthesized using gBlocks Gene Fragment synthesis service (Integrated DNA Technologies). *EGFP* with a flexible GS linker (GGGGS) at the C-terminus [90] was amplified from the pLJM1-EGFP vector (gift from David Sabatini; Addgene plasmid #19319) [91] using the EA062 and EA064 primers (**Key Resources Table**).

MyoAAV production and *in vivo* transduction

For myoAAV production, we used a published protocol with modifications [92]. We co-transfected the pAAV-cTnT-transgene vector (see above), the pHelper vector (GenBank: AF369965.1), and the pRep/Cap 1A-MYO capsid vector [56] into AAVpro 293T cells (TaKaRa, 632273) using polyethylenimine (PEI) reagent. The 1A-MYO capsid vector allows production of muscle-tropic AAV9 derivative (MyoAAV) [56]. The transfected cells were cultured in the OptiPRO™-SFM medium (Thermo Fisher, 12309019) for 72 hours at 37°C to produce virus. Cells and the culture media were collected for AAV purification. AAV particles were extracted with chloroform, precipitated with polyethylene glycol, purified with

DNase/RNase digestion followed by chloroform extraction, and concentrated in PBS with Amicon filters
(Millipore), as detailed in a previous report [92]. The viral genome copy numbers (vg) were estimated by
qPCR using primers amplifying the cTnT promoter region (primer ID KI466 and KI467, **Key Resources**
Table). MyoAAV particles were administered to mice through retro-orbital venous sinus injection at a
dose of 2×10^{11} vg per mouse at 4 weeks of age, which was 14-15 days prior to tamoxifen.

Image quantification related to nuclear envelope ruptures

For quantification of nuclei with PCM1-lost nuclear tips and DNA protrusion, we used immunofluorescent
images of isolated cardiomyocytes stained for PCM1 and counter-stained with DAPI (DNA) and Phalloidin
(F-actin). Rod-shape cardiomyocytes that had a well-defined sarcomere structure based on Phalloidin
staining and two or more nuclei were used for quantification. We quantified instances of DAPI signal
protrusion from a nuclear envelope site at which PCM1 signals were specifically discontinued ("local
PCM1 loss"). The percentage of cells with one or more nuclei with a local PCM1 loss and the number of
such nuclei per cell were quantified manually post imaging. We obtained the percentage from >50
cardiomyocytes per animal with a total of 3 animals per genotype. The percentages of cardiomyocytes
with PCM1-lost nuclei were compared between genotypes using unpaired one-tailed Welch's t-tests in
R.

For quantification of nuclei with icGAS puncta, we used isolated cardiomyocytes from wild-type
or $Lmna^{CKO}$ mice treated with MyoAAV-GFP-icGAS. The percentage of cells with icGAS puncta was
quantified manually post imaging. We obtained the percentage from >50 cardiomyocytes per animal with
a total of 3 animals per genotype. The icGAS-positive cardiomyocyte percentages were compared
between genotypes using an unpaired one-tailed Welch's t-test in R.

For quantification of NLS-tdTomato intensities, we used $Lmna^{F/F};Myh6MCM^{Tg/0};Rosa26^{CAG-LSL-}$
$^{tdTomato/+}$ and $Lmna^{+/+};Myh6MCM^{Tg/0};Rosa26^{CAG-LSL-tdTomato/+}$ mice. Isolated cardiomyocytes were fixed,
immunostained for PCM1, and counterstained with DAPI. Quantification of NLS-tdTomato signals was
performed in the Fiji image analysis software. Cardiomyocytes that had two or more nuclei were used for
quantification. Nuclei with nuclear envelope ruptures were identified as described above. A freehand-
drawn area was placed inside a nucleus for quantification of the nuclear tdTomato signal. A nucleus was
selected for quantification using the following algorithm: If a cell had only ruptured nuclei, one ruptured
nucleus was randomly selected for quantification; if a cell had both ruptured and unruptured nuclei, one
ruptured nucleus was randomly selected for quantification; if a cell had only unruptured nuclei, one
unruptured nucleus was randomly selected for quantification. For each nucleus, whether it had a local
PCM1 loss with DNA protrusion was manually determined as described above. A second quantification
area of the same size as the nuclear area was placed in the cytoplasm for cytoplasmic tdTomato signal
quantification. For each of the two areas, a mean NLS-tdTomato pixel intensity was measured using the
ROI manager function in Fiji. We performed this quantification process for three mice per genotype. In

total, we had quantification information for 99 wild-type cardiomyocytes (all intact PCM1), 90 *Lmna^CKO*
cardiomyocytes with local PCM1 loss, and 36 *Lmna^CKO* cardiomyocytes with intact PCM1. Signal
intensities between two groups were compared using Wilcoxon rank sum tests, and P-values were
adjusted for multiple testing using the Benjamini-Hochberg procedure in R.
640

TUNEL assay
TUNEL assays were performed on deparaffinized heart sections using a Click-iT™ Plus TUNEL Assay
Kits for In Situ Apoptosis Detection (Invitrogen, C10617), according to the manufacturer's instruction. The
samples were then co-stained with Alexa Fluor Plus 647 Phalloidin (Invitrogen, A30107) for 1 hour at
37°C to visualize cells in tissue sections. For positive controls, sample tissues were treated with DNase
I prior to the TUNEL assay. The number of cells with positive staining in the TUNEL assay (TUNEL$^+$ cells)
was determined using sections from three hearts per genotype. TUNEL positive cells in 2-3 images per
mouse (257 x 257 µm^2 per image) were counted manually and compared between two groups using
Wilcoxon rank sum tests in R.
650

Immunoblot
Snap-frozen hearts or isolated cardiomyocytes were lysed in the RIPA buffer for cGAS and STING
immunoblotting (20 mM Tris-HCl, 150 mM NaCl, 1% IGEPAL, 0.5% sodium deoxycholate, 0.1% SDS, 10
mM dithiothreitol, protease and phosphatase inhibitors) or the Urea buffer for Lamin A/C immunoblotting
(20 mM HEPES pH 7.4, 1 M NaCl, 8 M urea, protease and phosphatase inhibitors). Proteins were
extracted using pestle homogenization and sonication. Cell lysates were centrifuged at 13,000 rpm for 5
minutes at 4°C, and proteins in the supernatant were separated by SDS-PAGE. Proteins were transferred
to a PVDF membrane. Membranes were blocked with nonfat milk. The primary antibodies were rabbit
anti-Lamin A/C antibody (Santa Cruz, sc-20681, 1:1000), rabbit anti-cGAS antibody (Cell Signaling,
#31659, 1:500), rabbit anti-STING antibody (Cell Signaling, #13647, 1:500), and rabbit anti-GAPDH
antibody (ABclonal, AC001, 1:1000); Secondary antibodies were anti-rabbit or anti-mouse IgG (H+L)
DyLight 680 and 800 (Cell Signaling, 1:5000). Signals were detected and quantified in the Odyssey CLx
Imager (LI-COR). Gels after protein transfer were counter-stained with Coomassie to evaluate the loaded
protein amount.
665

Electron microscopy
Heart tissues were dissected and fixed in 2.5% glutaraldehyde, post-fixed in 1% OsO4 for 1 hour at 4°C,
rinsed, dehydrated in ethanol, and infiltrated overnight. The semithin Epon sections were screened by
light microscopy to select areas with longitudinal orientation of myocytes that were subsequently
processed to prepare thin sections. The sections were stained with 1% uranyl acetate, followed by lead

citrate. The sections were imaged using Hitachi Model H-7650 Transmission Electron Microscope at

Cincinnati Children's Integrated Pathology Research Facility.

Quantitative reverse-transcriptase PCR (RT-PCR)

For RT-PCR analysis, we used wild-type treated with MyoAAV-Luciferase (n=5) or MyoAAV-cGAS (n=3)

and *Lmna^{CKO}* mice treated with MyoAAV-Luciferase (n=4) or MyoAAV-cGAS (n=5) at 2 weeks post

tamoxifen. Total RNAs of isolated cardiomyocytes were extracted with Trizol LS (Invitrogen, 10296010),

treated with DNaseI, and purified. Quantitative RT-PCR was conducted with the Luna Universal One-

Step RT-qPCR Kit (NEB, E3005) on a Bio-Rad CFX96 Real-Time PCR Detection System (Bio-Rad) in

accordance with the manufacturer's instructions. The primers for mouse genes were: EA096 and EA097

for *Cxcl10*; EA098 and EA099 for *Ifnb1*; EA100 and EA101 for *Ifit3*; and KI444 and KI445 for *Actb* (**Key**

Resources Table). The delta-delta Ct method was used to quantify mRNA abundance with *Actb* as a

normalization control. mRNA levels between two groups were compared using Wilcoxon rank sum tests,

and P-values were adjusted for multiple testing using the Benjamini-Hochberg procedure in R.

RNA-seq

For RNA-seq comparing wild-type mice with *Lmna^{CKO}* mice at different post tamoxifen times, we used

wild-type (n=5) and *Lmna^{CKO}* mice (n=4) at 1 week post tamoxifen, wild-type (n=3) and *Lmna^{CKO}* mice

(n=2) at 11 days post tamoxifen, and wild-type (n=7) and *Lmna^{CKO}* mice (n=5) at 2 weeks post tamoxifen.

For RNA-seq analysis of *Lmna/Cgas* double knockout mice, we used *Cgas$^{-/-}$*;*Lmna$^{F/F;Myh6MCM}$* mice (n=4),

Cgas$^{+/+}$;*Lmna$^{F/F;Myh6MCM}$* (n=6), *Cgas$^{+/+}$*;*Lmna$^{F/+;Myh6MCM}$* (n=6), and *Cgas$^{-/-}$*;*Lmna$^{F/+;Myh6MCM}$* (n=4) for

experiments. Hearts were perfused with 100 mM KCl and removed for RNA extraction. Total RNAs were

extracted with Trizol LS (Invitrogen, 10296010), treated with DNaseI, and purified. For tamoxifen time

course experiments, mRNA sequencing libraries were generated using the poly-A selection module in

the NEBNext UltraII Directional RNA Library Prep Kit (NEB E7760) and sequenced on the Illumina HiSeq

2500 sequencer with single-end 50 cycles. For *cGas/Lmna* double knockout experiments, libraries were

generated using QuantSeq 3' mRNA-Seq Library Prep Kit FWD (Lexogen, K01596) and sequenced on

the Illumina NovaSeq 6000 sequencer with single-end 100 cycles.

RNA-seq analysis

High-throughput sequencing reads were aligned to the mouse mm39 reference genome with the

Gencode vM27 basic gene annotation using STAR version 2.7.9 [93] with the default alignment parameters

except using "clip3pAdapterSeq AGATCGGAAGAGCACACGTCTGAACTCCAGTCA". From raw read

counts per transcript, TPMs (transcripts per million) were calculated in R as follows: TPM=10^6 x

RPK/sum(RPK), where RPK=read count/transcript length in kp, and sum(RPK) is the sum of RPKs for

all transcripts. Raw read counts were used for differential gene expression analyses using DESeq2 [94] in

R. Protein-coding or lncRNA genes with adjusted P-value < 0.05 and absolute $\log_2$ fold change > 1 (NEB

Directional RNA-seq) or with adjusted P-value < 0.05 and absolute $\log_2$ fold change > 0 (QuantSeq 3'

mRNA-seq) were considered differentially expressed genes (**Table S1**). Differentially expressed genes

were analyzed for the enrichment of GO Biological Processes, GO Molecular Functions, GO Cellular

Components, and KEGG pathway terms using Metascape [95] with the default enrichment parameters.

Single-nucleus RNA-seq

For single-nucleus RNA-seq, we used wild-type (n=3) and *Lmna*[CKO] mice (n=3) at 2 weeks post

tamoxifen. Combinatorial indexing-based single-nucleus RNA-seq was conducted according to the

optimized sci-RNA-seq3 protocol developed previously [96]. Briefly, mice were anesthetized with

isoflurane, euthanized by cervical dislocation, and then hearts were perfused with 100 mM KCl. Removed

hearts were diced into small pieces in cold PBS containing Diethyl pyrocarbonate (DEPC, Sigma D5758).

Fresh heart pieces were dounce-homogenized to liberate nuclei in Hypotonic lysis buffer B (7.7 mM

Na_2HPO_4, 4.5 mM NaH_2PO_4, 1.8 mM KH_2PO_4, 2.7 mM KCl, 10.3 mM NaCl, 3 mM $MgCl_2$, 0.08% BSA,

0.025% Igepal CA-630, 1% DEPC) and passed through cell strainers. Nuclei were fixed in 1 mg/mL

dithiobis(succinimidyl propionate) (DSP, Thermo Fisher 22585) for 15 min on ice, washed, and stored at

–80°C until use. For each mouse, 50,000 nuclei were placed in each of 8 wells of a 96-well plate (total

48 wells for 6 mice). The remaining 48 wells were filled with similarly processed mouse heart nuclei with

comparable sample quality from an unrelated project. Nuclei were subjected to reverse transcription with

96-indexed poly-T RT primers that contained a unique molecular identifier (UMI) sequence

(3lvl_mRNA_RT_plate_1; **Table S3**). Nuclei from all wells were combined and redistributed evenly to 96

wells. Redistributed cells were subjected to ligation with 96-indexed ligation primers in 96 wells

(3lvl_mRNA_Lig_plate_1; **Table S3**). Nuclei from all wells were combined again and redistributed evenly

to 96 wells at 1,000 nuclei per well. Redistributed cells underwent second strand synthesis, protease

digestion, tagmentation using N7-loaded Tn5 transposase, and PCR-amplified using 96-indexed PCR

P7 primers (PCR_P7_plate1; **Table S3**) and the unindexed P5 primer, for 16 cycles. PCR amplicons

were combined and size-selected for DNA fragments between 250 bp and 600 bp. Purified library DNA

were sequenced for 50 bases on the paired-end mode (34 cycles on Read1 to sequence the ligation

index, UMI, and the RT barcode; 10 cycles on Index1 to sequence the P7 index; and 48 cycles on Read2

to sequence the cDNA) using an Illumina NovaSeq6000 sequencer.

Single-nucleus RNA-seq analysis

Pre-processing of the single-nucleus RNA-seq data was performed using the published sci-RNA-seq3

pipeline [96], available in Github (https://github.com/JunyueC/sci-RNA-seq3_pipeline). The sci-RNA-seq3

pipeline takes fastq files separated by the P7 index as input, aligns reads to the genome, and returns a

gene-by-cell read count matrix. First, raw reads were trimmed to remove poly A tails, aligned to the mm39

mouse reference genome using STAR, and filtered for MAPQ greater than or equal to 30. Duplicate reads

were removed based on UMI. Read counts per gene were computed for each cell defined by the unique

combination of the RT index, ligation index, and P7 index. The gene-by-cell count matrix produced by

the sci-RNA-seq3 pipeline was further processed in the *SingleCellExperiment* package framework in R

[97]. We used the *scDblFinder* package for doublet detection and the *scuttle* package for mitochondrial

RNA quantification in R. We obtained 14,111 nuclei after filtering for nuclei with the *scDblFinder* doublet

score less than 0.25 and mitochondrial RNA contamination percentage less than 2% of total RNAs per

cell. Read counts for a gene were normalized by per-cell read depth and then log-transformed to obtain

normalized expression values ("logcounts"). Logcounts were used to plot expression values in graphs.

Principal component analysis (PCA) was performed on logcounts for 2,000 highly variable genes across

cells using the *fixedPCA* function in the *scran* package. The top 50 principal components were used in

the *runUMAP* function in the *scater* package with n_neighbors=20 and min_dist=1 parameters to obtain

UMAP dimensions. To cluster cells, logcounts were processed in the *clusterCells* function in the *scran*

package using the top 50 principal components, with the number of nearest neighbors k=12. Cell clusters

were annotated manually with cell type names based on marker gene expression. Per-gene read counts

were summed within cell groups and processed by the *DESeq2* package [94] to perform pseudo-bulk

differential gene expression analysis. Genes with adjusted P-values smaller than 0.05 were defined as

differentially expressed genes (**Table S4**). GO analysis of differentially expressed genes was performed

as described in the RNA-seq analysis section. For the analysis for the enrichment within selected GO

and KEGG terms, all genes associated with the selected terms were obtained using the R packages

GO.db and *KEGGREST*, and enrichment of differentially expressed genes among these genes was

computed by Fisher's exact test with P-values adjusted by the Benjamini-Hochberg procedure. Cell-cell

communication analysis was performed using the *CellChat* package in R [74] using the 12 cell type

annotations and the default mouse database with a minimum of ten cells for the analysis. Positive

communications were selected by the *subsetCommunication* function with threshold P-value less than

0.05, and the reported interaction counts were used to plot data.

SLAM-IT-seq

For SLAM-IT-seq experiments, we used $Lmna^{F/F};Myh6MCM^{Tg/0};Uprt^{Tg/0}$ (n=6) and

$Lmna^{+/+};Myh6MCM^{Tg/0};Uprt^{Tg/0}$ (n=8) mice as experimental groups, and $Lmna^{F/F};Myh6MCM^{Tg/0};Uprt^{0/0}$

(n=4) and $Lmna^{+/+};Myh6MCM^{Tg/0};Uprt^{0/0}$ (n=3) mice as negative control groups that would not incorporate

4-thiouracil (4sU) due to the absence of the *Uprt* allele. All mice were administered with tamoxifen and

used at 2 weeks post tamoxifen treatment. SLAM-IT-seq was conducted according to a previously

established protocol [53]. Briefly, mice were intraperitoneally injected with 4-thiouracil (4sU) dissolved in a

DMSO/corn oil (1:3) solution at a dose of 10 mg per gram of body weight. Six hours after the injection,

mice were sacrificed, and hearts were perfused with 100 mM KCl and removed. RNAs from the left

ventricular free wall were extracted by Trizol LS, purified, and DNaseI-treated using Directzol RNA
miniprep kit (Zymo, R2052). Purified RNAs (1.5 µg) were subjected to alkylation in the presence of 12
mM Iodoacetamide (IAA, Sigma I1149). High-throughput sequencing libraries were generated from the
alkylated RNAs using Quant-seq 3' mRNA-seq FWD kit (Lexogen K01596) and sequenced 100 bases
on the single-end mode on an Illumina NovaSeq6000 sequencer.
784

SLAM-IT-seq analysis
We analyzed SLAM-IT-seq data according to the published SLAM-IT-seq data analysis pipeline that uses
the *slamdunk* package [98], available on GitHub (https://github.com/t-neumann/slamdunk), and R functions
[59]. First, raw reads were processed by the *slamdunk all* function in the the *slamdunk* package with the
following alignment parameters: --trim-5p 12 --topn 100 --multimap --max-read-length 101 with default --
var-fraction (0.8) and default -c (1). This function aligned reads to 3' UTRs of Gencode vM27 genes (total
208,307 UTRs from 37,065 unique genes) in the mm39 mouse reference genome. Aligned reads were
processed by the *alleyoop utrrates* function in the *slamdunk* package with default parameters to quantify
C-to-T conversion events. UTRs with at least one read coverage for all 21 samples were retained for
further analyses (47,717 UTRs from 14,250 unique genes). Statistically significant C-to-T conversion
events in each transcript (i.e. the likelihood of cardiomyocyte-derived transcripts) were identified by
comparing the conversion events in mice carrying the *Uprt* allele with those in mice not carrying this allele
using beta-binomial test (*bbtest* function) in R. P-values of the beta binomial test were adjusted with the
Benjamini-Hochberg procedure for multiple tests (**Table S2**). A transcript with the lowest adjusted P-
value was chosen to represent a gene if there were multiple transcripts per gene. Genes with the adjusted
P-values smaller than 0.05 were defined as originating from cardiomyocytes, whereas genes with
adjusted P-values greater than or equal to 0.05 were defined as not originating from cardiomyocytes.
802

803 **Quantification and statistical analysis**
804 Statistical details of experiments and sample sizes are described in Method sections and indicated in
805 figures and/or figure legends. Quantitative data were shown as means (bar) of biological replicates (dots).
806 P values < 0.05 were considered statistically significant. P-values are indicated in the figures as
807 appropriate.
808

809 **Image graphics**
810 Graphical abstract and the study summary (**Fig.4J**) were created with BioRender.com.
811

REFERENCES

1. Aebi, U., Cohn, J., Buhle, L., and Gerace, L. (1986). The nuclear lamina is a meshwork of intermediate-type filaments. Nature *323*, 560–564.

2. Goldman, A.E., Maul, G., Steinert, P.M., Yang, H.Y., and Goldman, R.D. (1986). Keratin-like proteins that coisolate with intermediate filaments of BHK-21 cells are nuclear lamins. Proc. Natl. Acad. Sci. U. S. A. *83*, 3839–3843.

3. Röber, R.A., Weber, K., and Osborn, M. (1989). Differential timing of nuclear lamin A/C expression in the various organs of the mouse embryo and the young animal: a developmental study. Development *105*, 365–378.

4. Dechat, T., Pfleghaar, K., Sengupta, K., Shimi, T., Shumaker, D.K., Solimando, L., and Goldman, R.D. (2008). Nuclear lamins: major factors in the structural organization and function of the nucleus and chromatin. Genes Dev. *22*, 832–853.

5. Fatkin, D., MacRae, C., Sasaki, T., Wolff, M.R., Porcu, M., Frenneaux, M., Atherton, J., Vidaillet, H.J., Jr, Spudich, S., De Girolami, U., et al. (1999). Missense mutations in the rod domain of the lamin A/C gene as causes of dilated cardiomyopathy and conduction-system disease. N. Engl. J. Med. *341*, 1715–1724.

6. Bonne, G., Di Barletta, M.R., Varnous, S., Bécane, H.M., Hammouda, E.H., Merlini, L., Muntoni, F., Greenberg, C.R., Gary, F., Urtizberea, J.A., et al. (1999). Mutations in the gene encoding lamin A/C cause autosomal dominant Emery-Dreifuss muscular dystrophy. Nat. Genet. *21*, 285–288.

7. Eriksson, M., Brown, W.T., Gordon, L.B., Glynn, M.W., Singer, J., Scott, L., Erdos, M.R., Robbins, C.M., Moses, T.Y., Berglund, P., et al. (2003). Recurrent de novo point mutations in lamin A cause Hutchinson-Gilford progeria syndrome. Nature *423*, 293–298.

8. Chen, L., Lee, L., Kudlow, B.A., Dos Santos, H.G., Sletvold, O., Shafeghati, Y., Botha, E.G., Garg, A., Hanson, N.B., Martin, G.M., et al. (2003). LMNA mutations in atypical Werner's syndrome. Lancet *362*, 440–445.

9. De Sandre-Giovannoli, A., Bernard, R., Cau, P., Navarro, C., Amiel, J., Boccaccio, I., Lyonnet, S., Stewart, C.L., Munnich, A., Le Merrer, M., et al. (2003). Lamin a truncation in Hutchinson-Gilford progeria. Science *300*, 2055–2055.

10. Cao, H., and Hegele, R.A. (2000). Nuclear lamin A/C R482Q mutation in canadian kindreds with Dunnigan-type familial partial lipodystrophy. Hum. Mol. Genet. *9*, 109–112.

11. Parks, S.B., Kushner, J.D., Nauman, D., Burgess, D., Ludwigsen, S., Peterson, A., Li, D., Jakobs, P., Litt, M., Porter, C.B., et al. (2008). Lamin A/C mutation analysis in a cohort of 324 unrelated patients with idiopathic or familial dilated cardiomyopathy. Am. Heart J. *156*, 161–169.

12. Hershberger, R.E., and Morales, A. (2008). LMNA-Related Dilated Cardiomyopathy. In GeneReviews®, M. P. Adam, H. H. Ardinger, R. A. Pagon, S. E. Wallace, L. J. H. Bean, K. Stephens, and A. Amemiya, eds. (University of Washington, Seattle).

13. Taylor, M.R.G., Fain, P.R., Sinagra, G., Robinson, M.L., Robertson, A.D., Carniel, E., Di Lenarda, A., Bohlmeyer, T.J., Ferguson, D.A., Brodsky, G.L., et al. (2003). Natural history of dilated cardiomyopathy due to lamin A/C gene mutations. J. Am. Coll. Cardiol. *41*, 771–780.

14. Sébillon, P., Bouchier, C., Bidot, L.D., Bonne, G., Ahamed, K., Charron, P., Drouin-Garraud, V., Millaire, A., Desrumeaux, G., Benaïche, A., et al. (2003). Expanding the phenotype of LMNA mutations in dilated cardiomyopathy and functional consequences of these mutations. J. Med.

Genet. *40*, 560–567.

15. van Rijsingen, I.A.W., Nannenberg, E.A., Arbustini, E., Elliott, P.M., Mogensen, J., Hermans-van
Ast, J.F., van der Kooi, A.J., van Tintelen, J.P., van den Berg, M.P., Grasso, M., et al. (2013).
Gender-specific differences in major cardiac events and mortality in lamin A/C mutation carriers.
Eur. J. Heart Fail. *15*, 376–384.

16. Charron, P., Arbustini, E., and Bonne, G. (2012). What Should the Cardiologist know about Lamin
Disease? Arrhythm Electrophysiol Rev *1*, 22–28.

17. Verga, L., Concardi, M., Pilotto, A., Bellini, O., Pasotti, M., Repetto, A., Tavazzi, L., and Arbustini,
E. (2003). Loss of lamin A/C expression revealed by immuno-electron microscopy in dilated
cardiomyopathy with atrioventricular block caused by LMNA gene defects. Virchows Arch. *443*,
664–671.

18. Arbustini, E., Pilotto, A., Repetto, A., Grasso, M., Negri, A., Diegoli, M., Campana, C., Scelsi, L.,
Baldini, E., Gavazzi, A., et al. (2002). Autosomal dominant dilated cardiomyopathy with
atrioventricular block: a lamin A/C defect-related disease. J. Am. Coll. Cardiol. *39*, 981–990.

19. Kato, K., Ohno, S., Sonoda, K., Makiyama, T., Ozawa, T., and Horie, M. (2020). Splice site
mutation of LMNA causes severe dilated cardiomyopathy via strong dominant reduction of total
lamin expression. Eur. Heart J. *41*. 10.1093/ehjci/ehaa946.0333.

20. Narula, N., Favalli, V., Tarantino, P., Grasso, M., Pilotto, A., Bellazzi, R., Serio, A., Gambarin, F.I.,
Charron, P., Meder, B., et al. (2012). Quantitative expression of the mutated lamin A/C gene in
patients with cardiolaminopathy. J. Am. Coll. Cardiol. *60*, 1916–1920.

21. Muchir, A., Pavlidis, P., Decostre, V., Herron, A.J., Arimura, T., Bonne, G., and Worman, H.J.
(2007). Activation of MAPK pathways links LMNA mutations to cardiomyopathy in Emery-Dreifuss
muscular dystrophy. J. Clin. Invest. *117*, 1282–1293.

22. Muchir, A., Shan, J., Bonne, G., Lehnart, S.E., and Worman, H.J. (2009). Inhibition of extracellular
signal-regulated kinase signaling to prevent cardiomyopathy caused by mutation in the gene
encoding A-type lamins. Hum. Mol. Genet. *18*, 241–247.

23. Wu, W., Muchir, A., Shan, J., Bonne, G., and Worman, H.J. (2011). Mitogen-Activated Protein
Kinase Inhibitors Improve Heart Function and Prevent Fibrosis in Cardiomyopathy Caused by
Mutation in Lamin A/C Gene. Circulation *123*, 53–61.

24. Muchir, A., Wu, W., Choi, J.C., Iwata, S., Morrow, J., Homma, S., and Worman, H.J. (2012).
Abnormal p38α mitogen-activated protein kinase signaling in dilated cardiomyopathy caused by
lamin A/C gene mutation. Hum. Mol. Genet. *21*, 4325–4333.

25. Choi, J.C., Muchir, A., Wu, W., Iwata, S., Homma, S., Morrow, J.P., and Worman, H.J. (2012).
Temsirolimus activates autophagy and ameliorates cardiomyopathy caused by lamin A/C gene
mutation. Sci. Transl. Med. *4*, 144ra102.

26. Chen, S.N., Lombardi, R., Karmouch, J., Tsai, J.-Y., Czernuszewicz, G., Taylor, M.R.G., Mestroni,
L., Coarfa, C., Gurha, P., and Marian, A.J. (2019). DNA Damage Response/TP53 Pathway Is
Activated and Contributes to the Pathogenesis of Dilated Cardiomyopathy Associated With LMNA
(Lamin A/C) Mutations. Circ. Res. *124*, 856–873.

27. Cheedipudi, S.M., Asghar, S., and Marian, A.J. (2022). Genetic Ablation of the DNA Damage
Response Pathway Attenuates Lamin-Associated Dilated Cardiomyopathy in Mice. JACC Basic
Transl Sci *7*, 1232–1245.

28. Davidson, P.M., and Lammerding, J. (2014). Broken nuclei--lamins, nuclear mechanics, and
disease. Trends Cell Biol. *24*, 247–256.

29. Chen, N.Y., Yang, Y., Weston, T.A., Belling, J.N., Heizer, P., Tu, Y., Kim, P., Edillo, L., Jonas, S.J.,
Weiss, P.S., et al. (2019). An absence of lamin B1 in migrating neurons causes nuclear membrane
ruptures and cell death. Proc. Natl. Acad. Sci. U. S. A. *116*, 25870–25879.

30. Cho, S., Vashisth, M., Abbas, A., Majkut, S., Vogel, K., Xia, Y., Ivanovska, I.L., Irianto, J., Tewari,
M., Zhu, K., et al. (2019). Mechanosensing by the Lamina Protects against Nuclear Rupture, DNA
Damage, and Cell-Cycle Arrest. Dev. Cell *49*, 920-935.e5.

31. Earle, A.J., Kirby, T.J., Fedorchak, G.R., Isermann, P., Patel, J., Iruvanti, S., Moore, S.A., Bonne,
G., Wallrath, L.L., and Lammerding, J. (2020). Mutant lamins cause nuclear envelope rupture and
DNA damage in skeletal muscle cells. Nat. Mater. *19*, 464–473.

32. Sylvius, N., Bilinska, Z.T., Veinot, J.P., Fidzianska, A., Bolongo, P.M., Poon, S., McKeown, P.,
Davies, R.A., Chan, K.-L., Tang, A.S.L., et al. (2005). In vivo and in vitro examination of the
functional significances of novel lamin gene mutations in heart failure patients. J. Med. Genet. *42*,
639–647.

33. Hatch, E.M., and Hetzer, M.W. (2016). Nuclear envelope rupture is induced by actin-based nucleus
confinement. J. Cell Biol. *215*, 27–36.

34. Vargas, J.D., Hatch, E.M., Anderson, D.J., and Hetzer, M.W. (2012). Transient nuclear envelope
rupturing during interphase in human cancer cells. Nucleus *3*, 88–100.

35. De Vos, W.H., Houben, F., Kamps, M., Malhas, A., Verheyen, F., Cox, J., Manders, E.M.M.,
Verstraeten, V.L.R.M., van Steensel, M.A.M., Marcelis, C.L.M., et al. (2011). Repetitive disruptions
of the nuclear envelope invoke temporary loss of cellular compartmentalization in laminopathies.
Hum. Mol. Genet. *20*, 4175–4186.

36. Siu, C.-W., Lee, Y.-K., Ho, J.C.-Y., Lai, W.-H., Chan, Y.-C., Ng, K.-M., Wong, L.-Y., Au, K.-W., Lau,
Y.-M., Zhang, J., et al. (2012). Modeling of lamin A/C mutation premature cardiac aging using
patient-specific induced pluripotent stem cells. Aging *4*, 803–822.

37. Fan, Y., Tan, D., Zhang, X., Song, D., Chang, X., Wang, S., Yan, H., Ge, L., Yang, H.,
Bönnemann, C., et al. (2020). Nuclear Factor-κB Pathway Mediates the Molecular Pathogenesis of
LMNA-Related Muscular Dystrophies. Biochem. Genet. *58*, 966–980.

38. Chai, R.J., Werner, H., Li, P.Y., Lee, Y.L., Nyein, K.T., Solovei, I., Luu, T.D.A., Sharma, B.,
Navasankari, R., Maric, M., et al. (2021). Disrupting the LINC complex by AAV mediated gene
transduction prevents progression of Lamin induced cardiomyopathy. Nat. Commun. *12*, 4722.

39. Chen, R., Buchmann, S., Kroth, A., Arias-Loza, A.-P., Kohlhaas, M., Wagner, N., Grüner, G.,
Nickel, A., Cirnu, A., Williams, T., et al. (2023). Mechanistic Insights of the LEMD2 p.L13R
Mutation and Its Role in Cardiomyopathy. Circ. Res. *132*, e43–e58.

40. Kim, P.H., Chen, N.Y., Heizer, P.J., Tu, Y., Weston, T.A., Fong, J.L.-C., Gill, N.K., Rowat, A.C.,
Young, S.G., and Fong, L.G. (2021). Nuclear membrane ruptures underlie the vascular pathology
in a mouse model of Hutchinson-Gilford progeria syndrome. JCI Insight *6*.
10.1172/jci.insight.151515.

41. Ross, J.A., Arcos-Villacis, N., Battey, E., Boogerd, C., Avalos Orellana, C., Marhuenda, E.,
Swiatlowska, P., Hodzic, D., Prin, F., Mohun, T., et al. (2023). Lem2 is essential for cardiac
development by maintaining nuclear integrity. Cardiovasc. Res. 10.1093/cvr/cvad061.

42. Zhang, Y., Ramirez-Martinez, A., Chen, K., McAnally, J.R., Cai, C., Durbacz, M.Z., Chemello, F.,
Wang, Z., Xu, L., Bassel-Duby, R., et al. (2023). Net39 protects muscle nuclei from mechanical
stress during the pathogenesis of Emery-Dreifuss muscular dystrophy. J. Clin. Invest. *133*.
10.1172/JCI163333.

43. Eissenberg, J.C., and Gonzalo, S. (2020). Pushing the limit on laminopathies. Nat. Mater. *19*, 378–
380.

44. Gauthier, B.R., and Comaills, V. (2021). Nuclear Envelope Integrity in Health and Disease:
Consequences on Genome Instability and Inflammation. Int. J. Mol. Sci. *22*.
10.3390/ijms22147281.

45. Miller, K.N., Victorelli, S.G., Salmonowicz, H., Dasgupta, N., Liu, T., Passos, J.F., and Adams, P.D.
(2021). Cytoplasmic DNA: sources, sensing, and role in aging and disease. Cell *184*, 5506–5526.

46. Yum, S., Li, M., Fang, Y., and Chen, Z.J. (2021). TBK1 recruitment to STING activates both IRF3
and NF-κB that mediate immune defense against tumors and viral infections. Proceedings of the
National Academy of Sciences *118*, e2100225118.

47. Decout, A., Katz, J.D., Venkatraman, S., and Ablasser, A. (2021). The cGAS–STING pathway as a
therapeutic target in inflammatory diseases. Nat. Rev. Immunol. *21*, 548–569.

48. Kreienkamp, R., Graziano, S., Coll-Bonfill, N., Bedia-Diaz, G., Cybulla, E., Vindigni, A., Dorsett, D.,
Kubben, N., Batista, L.F.Z., and Gonzalo, S. (2018). A Cell-Intrinsic Interferon-like Response Links
Replication Stress to Cellular Aging Caused by Progerin. Cell Rep. *22*, 2006–2015.

49. King, K.R., Aguirre, A.D., Ye, Y.-X., Sun, Y., Roh, J.D., Ng, R.P., Jr, Kohler, R.H., Arlauckas, S.P.,
Iwamoto, Y., Savol, A., et al. (2017). IRF3 and type I interferons fuel a fatal response to myocardial
infarction. Nat. Med. *23*, 1481–1487.

50. Cao, D.J., Schiattarella, G.G., Villalobos, E., Jiang, N., May, H.I., Li, T., Chen, Z.J., Gillette, T.G.,
and Hill, J.A. (2018). Cytosolic DNA Sensing Promotes Macrophage Transformation and Governs
Myocardial Ischemic Injury. Circulation *137*, 2613–2634.

51. Hu, D., Cui, Y.-X., Wu, M.-Y., Li, L., Su, L.-N., Lian, Z., and Chen, H. (2020). Cytosolic DNA sensor
cGAS plays an essential pathogenetic role in pressure overload-induced heart failure. Am. J.
Physiol. Heart Circ. Physiol. *318*, H1525–H1537.

52. Yan, M., Li, Y., Luo, Q., Zeng, W., Shao, X., Li, L., Wang, Q., Wang, D., Zhang, Y., Diao, H., et al.
(2022). Mitochondrial damage and activation of the cytosolic DNA sensor cGAS-STING pathway
lead to cardiac pyroptosis and hypertrophy in diabetic cardiomyopathy mice. Cell Death Discov *8*,
258.

53. Hasper, J., Welle, K., Swovick, K., Hryhorenko, J., Ghaemmaghami, S., and Buchwalter, A. (2024).
Long lifetime and tissue-specific accumulation of lamin A/C in Hutchinson-Gilford progeria
syndrome. J. Cell Biol. *223*. 10.1083/jcb.202307049.

54. Patterson, M., Barske, L., Van Handel, B., Rau, C.D., Gan, P., Sharma, A., Parikh, S., Denholtz,
M., Huang, Y., Yamaguchi, Y., et al. (2017). Frequency of mononuclear diploid cardiomyocytes
underlies natural variation in heart regeneration. Nat. Genet. *49*, 1346–1353.

55. Srsen, V., Fant, X., Heald, R., Rabouille, C., and Merdes, A. (2009). Centrosome proteins form an
insoluble perinuclear matrix during muscle cell differentiation. BMC Cell Biol. *10*, 28.

56. Tabebordbar, M., Lagerborg, K.A., Stanton, A., King, E.M., Ye, S., Tellez, L., Krunnfusz, A.,

Tavakoli, S., Widrick, J.J., Messemer, K.A., et al. (2021). Directed evolution of a family of AAV
capsid variants enabling potent muscle-directed gene delivery across species. Cell *184*, 4919-
4938.e22.

57. Denais, C.M., Gilbert, R.M., Isermann, P., McGregor, A.L., te Lindert, M., Weigelin, B., Davidson,
P.M., Friedl, P., Wolf, K., and Lammerding, J. (2016). Nuclear envelope rupture and repair during
cancer cell migration. Science *352*, 353–358.

58. Raab, M., Gentili, M., de Belly, H., Thiam, H.R., Vargas, P., Jimenez, A.J., Lautenschlaeger, F.,
Voituriez, R., Lennon-Duménil, A.M., Manel, N., et al. (2016). ESCRT III repairs nuclear envelope
ruptures during cell migration to limit DNA damage and cell death. Science *352*, 359–362.

59. Matsushima, W., Herzog, V.A., Neumann, T., Gapp, K., Zuber, J., Ameres, S.L., and Miska, E.A.
(2019). Sequencing cell-type-specific transcriptomes with SLAM-ITseq. Nat. Protoc. *14*, 2261–
2278.

60. Kuhl, N., Linder, A., Philipp, N., Nixdorf, D., Fischer, H., Veth, S., Kuut, G., Xu, T.T., Theurich, S.,
Carell, T., et al. (2023). STING agonism turns human T cells into interferon-producing cells but
impedes their functionality. EMBO Rep. *24*, e55536.

61. Liu, Y., Xu, P., Rivara, S., Liu, C., Ricci, J., Ren, X., Hurley, J.H., and Ablasser, A. (2022). Clathrin-
associated AP-1 controls termination of STING signalling. Nature *610*, 761–767.

62. Ikushima, H., Negishi, H., and Taniguchi, T. (2013). The IRF family transcription factors at the
interface of innate and adaptive immune responses. Cold Spring Harb. Symp. Quant. Biol. *78*,
105–116.

63. Della Corte, C.M., Sen, T., Gay, C.M., Ramkumar, K., Diao, L., Cardnell, R.J., Rodriguez, B.L.,
Stewart, C.A., Papadimitrakopoulou, V.A., Gibson, L., et al. (2020). STING Pathway Expression
Identifies NSCLC With an Immune-Responsive Phenotype. J. Thorac. Oncol. *15*, 777–791.

64. Schoggins, J.W., MacDuff, D.A., Imanaka, N., Gainey, M.D., Shrestha, B., Eitson, J.L., Mar, K.B.,
Richardson, R.B., Ratushny, A.V., Litvak, V., et al. (2014). Pan-viral specificity of IFN-induced
genes reveals new roles for cGAS in innate immunity. Nature *505*, 691–695.

65. Jin, L., Hill, K.K., Filak, H., Mogan, J., Knowles, H., Zhang, B., Perraud, A.-L., Cambier, J.C., and
Lenz, L.L. (2011). MPYS is required for IFN response factor 3 activation and type I IFN production
in the response of cultured phagocytes to bacterial second messengers cyclic-di-AMP and cyclic-
di-GMP. J. Immunol. *187*, 2595–2601.

66. Cohen, T.V., Gnocchi, V.F., Cohen, J.E., Phadke, A., Liu, H., Ellis, J.A., Foisner, R., Stewart, C.L.,
Zammit, P.S., and Partridge, T.A. (2013). Defective skeletal muscle growth in lamin A/C-deficient
mice is rescued by loss of Lap2α. Hum. Mol. Genet. *22*, 2852–2869.

67. Ortega, A., Roselló-Lletí, E., Tarazón, E., Molina-Navarro, M.M., Martínez-Dolz, L., González-
Juanatey, J.R., Lago, F., Montoro-Mateos, J.D., Salvador, A., Rivera, M., et al. (2014).
Endoplasmic reticulum stress induces different molecular structural alterations in human dilated
and ischemic cardiomyopathy. PLoS One *9*, e107635.

68. Sasagawa, S., Nishimura, Y., Okabe, S., Murakami, S., Ashikawa, Y., Yuge, M., Kawaguchi, K.,
Kawase, R., Okamoto, R., Ito, M., et al. (2016). Downregulation of GSTK1 Is a Common
Mechanism Underlying Hypertrophic Cardiomyopathy. Front. Pharmacol. *7*, 162.

69. Fan, P., Zhang, L., Cheng, T., Wang, J., Zhou, J., Zhao, L., Hua, C., and Xia, Q. (2021). MiR-590-
5p inhibits pathological hypertrophy mediated heart failure by targeting RTN4. J. Mol. Histol. *52*,

955–964.

70. Wehrens, M., de Leeuw, A.E., Wright-Clark, M., Eding, J.E.C., Boogerd, C.J., Molenaar, B., van
der Kraak, P.H., Kuster, D.W.D., van der Velden, J., Michels, M., et al. (2022). Single-cell
transcriptomics provides insights into hypertrophic cardiomyopathy. Cell Rep. *39*, 110809.

71. Snider, P., Standley, K.N., Wang, J., Azhar, M., Doetschman, T., and Conway, S.J. (2009). Origin
of cardiac fibroblasts and the role of periostin. Circ. Res. *105*, 934–947.

72. Lavine, K.J., Epelman, S., Uchida, K., Weber, K.J., Nichols, C.G., Schilling, J.D., Ornitz, D.M.,
Randolph, G.J., and Mann, D.L. (2014). Distinct macrophage lineages contribute to disparate
patterns of cardiac recovery and remodeling in the neonatal and adult heart. Proc. Natl. Acad. Sci.
U. S. A. *111*, 16029–16034.

73. Liu, C., Raab, M., Gui, Y., and Rudd, C.E. (2023). Multi-functional adaptor SKAP1: regulator of
integrin activation, the stop-signal, and the proliferation of T cells. Front. Immunol. *14*, 1192838.

74. Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.-H., Myung, P., Plikus,
M.V., and Nie, Q. (2021). Inference and analysis of cell-cell communication using CellChat. Nat.
Commun. *12*, 1088.

75. Zhang, H., Ren, L., and Wu, J.C. (2022). New Insights Into the Therapy for Lamin-Associated
Dilated Cardiomyopathy. JACC Basic Transl Sci *7*, 1246–1248.

76. Long, Z.-J., Wang, J.-D., Xu, J.-Q., Lei, X.-X., and Liu, Q. (2022). cGAS/STING cross-talks with cell
cycle and potentiates cancer immunotherapy. Mol. Ther. *30*, 1006–1017.

77. Yang, H., Wang, H., Ren, J., Chen, Q., and Chen, Z.J. (2017). cGAS is essential for cellular
senescence. Proc. Natl. Acad. Sci. U. S. A. *114*, E4612–E4620.

78. Thomsen, M.K., Nandakumar, R., Stadler, D., Malo, A., Valls, R.M., Wang, F., Reinert, L.S.,
Dagnaes-Hansen, F., Hollensen, A.K., Mikkelsen, J.G., et al. (2016). Lack of immunological DNA
sensing in hepatocytes facilitates hepatitis B virus infection. Hepatology *64*, 746–759.

79. Li, T., and Chen, Z.J. (2018). The cGAS-cGAMP-STING pathway connects DNA damage to
inflammation, senescence, and cancer. J. Exp. Med. *215*, 1287–1299.

80. Willaume, S., Rass, E., Fontanilla-Ramirez, P., Moussa, A., Wanschoor, P., and Bertrand, P.
(2021). A Link between Replicative Stress, Lamin Proteins, and Inflammation. Genes *12*.
10.3390/genes12040552.

81. Rouhi, L., Auguste, G., Zhou, Q., Lombardi, R., Olcum, M., Pourebrahim, K., Cheedipudi, S.M.,
Asghar, S., Hong, K., Robertson, M.J., et al. (2022). Deletion of the Lmna gene in fibroblasts
causes senescence-associated dilated cardiomyopathy by activating the double-stranded DNA
damage response and induction of senescence-associated secretory phenotype. J Cardiovasc
Aging *2*. 10.20517/jca.2022.14.

82. Liu, Y., Chen, X., Zhao, Y., Wang, X.-Y., Luo, Y.-W., Chen, L., Wang, W., Zhong, S., Hu, M., Dai,
Z., et al. (2023). Small cytosolic double-stranded DNA represses cyclic GMP-AMP synthase
activation and induces autophagy. Cell Rep. *42*, 112852.

83. Nader, G.P. de F., Agüera-Gonzalez, S., Routet, F., Gratia, M., Maurin, M., Cancila, V., Cadart, C.,
Palamidessi, A., Ramos, R.N., San Roman, M., et al. (2021). Compromised nuclear envelope
integrity drives TREX1-dependent DNA damage and tumor cell invasion. Cell *184*, 5230-5246.e22.

84. Kim, Y., and Zheng, Y. (2013). Generation and characterization of a conditional deletion allele for

Lmna in mice. Biochem. Biophys. Res. Commun. *440*, 8–13.

85. Sohal, D.S., Nghiem, M., Crackower, M.A., Witt, S.A., Kimball, T.R., Tymitz, K.M., Penninger, J.M.,
and Molkentin, J.D. (2001). Temporally regulated and tissue-specific gene manipulations in the
adult and embryonic heart using a tamoxifen-inducible Cre protein. Circ. Res. *89*, 20–25.

86. Daigle, T.L., Madisen, L., Hage, T.A., Valley, M.T., Knoblich, U., Larsen, R.S., Takeno, M.M.,
Huang, L., Gu, H., Larsen, R., et al. (2018). A Suite of Transgenic Driver and Reporter Mouse
Lines with Enhanced Brain-Cell-Type Targeting and Functionality. Cell *174*, 465-480.e22.

87. Gay, L., Miller, M.R., Ventura, P.B., Devasthali, V., Vue, Z., Thompson, H.L., Temple, S., Zong, H.,
Cleary, M.D., Stankunas, K., et al. (2013). Mouse TU tagging: a chemical/genetic intersectional
method for purifying cell type-specific nascent RNA. Genes Dev. *27*, 98–115.

88. Verzi, M.P., McCulley, D.J., De Val, S., Dodou, E., and Black, B.L. (2005). The right ventricle,
outflow tract, and ventricular septum comprise a restricted expression domain within the
secondary/anterior heart field. Dev. Biol. *287*, 134–145.

89. Lin, Z., von Gise, A., Zhou, P., Gu, F., Ma, Q., Jiang, J., Yau, A.L., Buck, J.N., Gouin, K.A., van
Gorp, P.R.R., et al. (2014). Cardiac-specific YAP activation improves cardiac function and survival
in an experimental murine MI model. Circ. Res. *115*, 354–363.

90. Chen, X., Zaro, J.L., and Shen, W.-C. (2013). Fusion protein linkers: property, design and
functionality. Adv. Drug Deliv. Rev. *65*, 1357–1369.

91. Sancak, Y., Peterson, T.R., Shaul, Y.D., Lindquist, R.A., Thoreen, C.C., Bar-Peled, L., and
Sabatini, D.M. (2008). The Rag GTPases bind raptor and mediate amino acid signaling to
mTORC1. Science *320*, 1496–1501.

92. Negrini, M., Wang, G., Heuer, A., Björklund, T., and Davidsson, M. (2020). AAV Production
Everywhere: A Simple, Fast, and Reliable Protocol for In-house AAV Vector Production Based on
Chloroform Extraction. Curr. Protoc. Neurosci. *93*, e103.

93. Frankish, A., Diekhans, M., Jungreis, I., Lagarde, J., Loveland, J.E., Mudge, J.M., Sisu, C., Wright,
J.C., Armstrong, J., Barnes, I., et al. (2021). GENCODE 2021. Nucleic Acids Res. *49*, D916–D923.

94. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. *15*, 550.

95. Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A.H., Tanaseichuk, O., Benner, C., and
Chanda, S.K. (2019). Metascape provides a biologist-oriented resource for the analysis of
systems-level datasets. Nat. Commun. *10*, 1523.

96. Martin, B.K., Qiu, C., Nichols, E., Phung, M., Green-Gladden, R., Srivatsan, S., Blecher-Gonen, R.,
Beliveau, B.J., Trapnell, C., Cao, J., et al. (2022). Optimized single-nucleus transcriptional profiling
by combinatorial indexing. Nat. Protoc. 10.1038/s41596-022-00752-0.

97. Amezquita, R.A., Lun, A.T.L., Becht, E., Carey, V.J., Carpp, L.N., Geistlinger, L., Marini, F., Rue-
Albrecht, K., Risso, D., Soneson, C., et al. (2020). Orchestrating single-cell analysis with
Bioconductor. Nat. Methods *17*, 137–145.

98. Herzog, V.A., Reichholf, B., Neumann, T., Rescheneder, P., Bhat, P., Burkard, T.R., Wlotzka, W.,
von Haeseler, A., Zuber, J., and Ameres, S.L. (2017). Thiol-linked alkylation of RNA to assess
expression dynamics. Nat. Methods *14*, 1198–1204.

1103

1104 **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-Lamin A/C	Abcam	Cat # Ab26300
Anti-Lamin A/C	Santa Cruz	Cat # sc-376248
Anti-Lamin A/C	Santa Cruz	Cat # sc-20681
Anti-CD45 antibody	R&D Systems	Cat # AF114-SP
Anti-CD68 antibody	Cell Signaling	Cat # 97778
Anti-γ-H2A.X antibody	Cell Signaling	Cat # 9718
Anti-desmin antibody	Invitrogen	Cat # PA5-19063
Anti-PCM-1 antibody	Sigma	Cat # HPA023370
Anti-tdTomato antibody	OriGene	Cat # AB8181
Anti-COX4 antibody	R&D Systems	Cat # MAB6980
Anti-cGAS antibody	Cell Signaling	Cat # 31659
Anti-STING antibody	Cell Signaling	Cat # 13647
Anti-GAPDH antibody	ABclonal	Cat # AC001
Alexa Fluor Plus 647 Phalloidin	ThermoFisher	Cat # A30107
Bacterial and virus strains		
Chemicals, peptides, and recombinant proteins		
Tamoxifen	Sigma	Cat # T5648
Collagenase Type II	Worthington	Cat # LS004177
Polyethylenimine, Linear, MW 25000, Transfection Grade (PEI 25K™)	Fisher Scientific	Cat # 23966-100
Target Retrieval Solution	DAKO	Cat # S2367
Corn oil	Sigma	Cat # C8267
Phenol/Chloroform/Isoamyl alcohol	Invitrogen	Cat # 15593031
Trizol LS	Invitrogen	Cat # 10296010
Heparin	Sagent Pharmaceuticals.	Cat # 25021-400-10
VectaStain ELITE ABC Kits	Vector Laboratories	Cat # PK-6100
DAB	DAKO	Cat # K3468
ProLong Glass Antifade Mountant	Invitrogen	Cat # P36984
Normal donkey serum	Jackson ImmunoResearch	Cat # 017-000-121
Sudan Black B	Electron Microscopy Slides	Cat # 21610
NEBuilder HiFi DNA Assembly Master Mix	NEB	Cat # E2621
Dulbecco's modified Eagle medium	Fisher Scientific	Cat # 11965-092
penicillin/streptomycin	Invitrogen	Cat # 105727
OptiPRO™-SFM	Thermo Fisher	Cat # 12309019
Fetal Bovine Serum	Sigma	Cat # F1051
Critical commercial assays		
Masson's Trichrome Kit	Thomas Scientific LLC	Cat # KTMTRPT
Click-iT™ Plus TUNEL Assay Kits for In Situ Apoptosis Detection; Alexa Fluor™ 488	Invitrogen	Cat # C10617
Luna Universal One-Step RT-qPCR Kit	New England Biolabs	Cat # E3005
NEBNext UltraII Directional RNA Library Prep Kit	New England Biolabs	Cat # E7760
Quant-seq 3' mRNA-seq FWD kit	Lexogen	Cat # K01596
Deposited data		

Experimental models: Organisms/strains		
Lmna-LoxP mice	Jackson Laboratory	JAX stock No: 26284
Myh6-MerCreMer transgene mice	Jackson Laboratory	JAX stock No: 005657
Rosa26$^{CAG-LSL-tdTomato}$ mice	Jackson Laboratory	JAX stock No: 25106
Cgas-null mice	Jackson Laboratory	JAX stock No: 026554
CAG-Lox-Stop-Lox-Uprt transgene mice	Jackson Laboratory	JAX stock No: 021469
Sting-flox mice	Jackson Laboratory	JAX stock No: 031670
Mice carrying the *Mef2C-AHF-Cre* transgene allele	Verzi *et al.*, 2005	DOI: 10.1016/j.ydbio.2005.08.041
Sting-null (generated from *Sting-LoxP* mouse)	This study	N/A
AAVpro 293T Cell	TaKaRa	Cat # 632273
Oligonucleotides		
DNA oligonucleotide KI279 for *Lmna* genotype primer: AACCCAGCCTCAGAAACTGGTGGATG	This paper	Ikegami lab ID: KI279
DNA oligonucleotide KI280 for *Lmna* genotype primer: GACAGCTCTCCTCTGAAGTGCTTGGA	This paper	Ikegami lab ID: Ki280
DNA oligonucleotide EA062 for EGFP cloning: AGTACTTAATACGACTCACTATAGGCTAGCGCCACCATGGTGAGC AAGGGCGAGGA	This paper	Ikegami lab ID: EA062
DNA oligonucleotide EA064 for EGFP with GS linker cloning: TTATCATGTCTGGATCCGCGCGGCCGCGCTGCCCCCGCCTCCCTT GTACAGCTCGTCCATGC	This paper	Ikegami lab ID: EA064

| DNA oligonucleotide EA086 for icGAS (human cGAS with E225A/D227A amino acid substitutions): AAGGGAGGCGGGGGCAGCGCGGCCGCAGACTACAAAGACGAT GACGACAAGATGCAGCCCTGGCACGGTAAAGCAATGCAACGGG CTAGTGAAGCAGGAGCTACTGCTCCAAAGGCCAGTGCTAGGAAT GCCAGGGGCGCTCCCATGGATCCAAATGAAAGCCCCGCAGCACC CGAAGCAGCTCTGCCTAAAGCAGGTAAGTTTGGCCCTGCTCGGA AGTCTGGTTCCAGACAAAAGAAATCAGCTCCTGATACTCAAGAG AGGCCTCCAGTAAGGGCTACCGGAGCTAGGGCCAAGAAAGCAC CCCAGAGAGCCCAGGATACCCAGCCTAGTGATGCAACTTCAGCC CCAGGCGCAGAGGGGCTGGAACCTCCCGCCGCACGAGAGCCCG CCCTTTCACGAGCCGGATCCTGCCGCCAACGAGGTGCCAGATGC AGCACAAAACCCAGGCCCCCTCCCGGTCCATGGGATGTACCCAG CCCCGGTCTTCCCGTAAGTGCACCAATATTGGTTCGAAGGGACG CCGCACCAGGGGCCTCTAAACTGCGGGCCGTGCTGGAGAAGCT GAAGTTGTCTCGGGATGATATCAGCACTGCCGCTGGCATGGTTA AAGGGGTTGTTGACCATCTGCTTCTGCGACTGAAATGCGACTCA GCTTTTAGGGGTGTAGGGCTTCTGAACACTGGATCCTATTACGA GCACGTCAAGATTAGTGCTCCCAACGCCTTCGCAGTCATGTTTAA GCTGGAAGTTCCTAGAATTCAGCTGGAAGAGTATTCTAACACTCG AGCTTATTATTTTGTCAAATTTAAGCGAAACCCAAAGGAGAATCA CCTGTCTCAGTTCTTGGAAGGGGGAGATACTGTCCGCCTCTAAGAT GCTGTCTAAGTTCAGAAAGATTATAAAGGAGGAAATTAATGATA TCAAAGATACTGACGTGATTATGAAAAGGAAGCGCGGAGGTTCT CCTGCAGTGACTCTGCTCATTAGCGAGAAGATATCTGTTGACATA ACCCTCGCACTCGAATCCAAGTCTAGCTGGCCTGCCTCTACCCAG GAGGGCTTGAGAATTCAAAACTGGTTGTCTGCAAAAGTAAGGAA ACAACTTCGACTGAAGCCATTTTATCTCGTCCCAAAGCACGCAAA AGAGGGCAATGGTTTCCAAGAAGAAACTTGGCGCTTGTCCTTTTC ACACATTGAGAAGGAGATTCTGAACAATCACGGAAAGTCCAAAA CTTGCTGCGAGAATAAGGAAGAAAAGTGCTGTCGAAAGGACTG CTTGAAGTTGATGAAGTATCTCCTGGAACAGTTGAAGGAACGAT TTAAAGATAAGAAACACTTGGACAAATTTAGTTCCTACCACGTGA AGACAGCCTTCTTTCACGTGTGTACCCAGAATCCCCAGGATTCAC AGTGGGATAGAAAGGACCTCGGACTCTGCTTTGACAATTGCGTC ACTTATTTTCTTCAGTGTTTGCGAACTGAGAAGCTGGAAAACTAT TTTATCCCCGAATTTAATTTGTTCTCATCAAACCTGATTGACAAGA GATCTAAAGAATTCCTCACAAAACAGATCGAGTACGAACGCAAC AACGAATTCCCCGTCTTTGATGAATTTTGAGCGGCCGCGCGGATC CAGACATGATAA | This paper | Ikegami lab ID: EA086 |

DNA oligonucleotide EA087 for mouse cGAS: AGTACTTAATACGACTCACTATAGGCTAGCGCCACCATGGACTAC AAAGACGATGACGATAAGATGGAGGATCCTCGCAGGCGCACCA CAGCTCCACGCGCAAAGAAACCATCTGCTAAGAGAGCACCTACC CAGCCTTCTAGGACAAGAGCCCATGCCGAGTCCTGCGGACCACA ACGAGGCGCCAGGAGTCGGCGGGCAGAACGGGACGGAGACAC AACCGAGAAACCTAGGGCTCCTGGCCCTAGAGTGCACCCAGCAC GCGCTACAGAGCTGACTAAAGATGCACAGCCCAGTGCAATGGAT GCAGCTGGTGCCACCGCACGCCCAGCCGTGCGGGTTCCACAACA GCAGGCTATATTGGACCCCGAACTTCCTGCTGTAAGGGAACCTCA ACCCCCCGCAGATCCCGAAGCTAGAAAAGTAGTACGAGGACCTT CTCACCGACGAGGGGCTCGATCAACCGGCCAGCCACGGGCTCCC AGGGGGTCACGCAAAGAACCAGATAAGCTCAAGAAAGTCCTGG ATAAGTTGCGGTTGAAAAGAAAAGACATCAGTGAAGCTGCTGAA ACAGTTAACAAAGTCGTCGAGAGACTGTTGCGGCGCATGCAAAA AAGGGAATCTGAATTCAAAGGAGTCGAACAACTCAACACCGGTT CATACTACGAGCATGTTAAAATCTCAGCCCCAAATGAGTTTGATG TGATGTTTAAGCTTGAAGTGCCTCGCATTGAACTGCAGGAGTACT ATGAGACTGGAGCCTTTTACCTCGTAAAGTTTAAGCGAATTCCTA GGGGTAATCCCTTGTCTCATTTTCTCGAAGGTGAAGTCTTGTCAG CTACCAAAATGCTGAGTAAATTTAGGAAGATAATAAAAGAGGAG GTTAAGGAAATAAAGGACATCGACGTTTCAGTGGAGAAGGAGA AGCCCGGTTCTCCAGCAGTTACACTCCTCATCAGGAATCCCGAGG AAATTAGCGTAGACATCATTCTCGCTTTGGAAAGCAAGGGTAGTT GGCCCATTTCTACCAAAGAAGGACTTCCCATTCAGGGTTGGCTCG GTACCAAAGTACGCACCAATCTCCGAAGGGAACCTTTCTACCTGG TTCCTAAAAATGCTAAAGATGGAAACTCTTTCCAGGGCGAGACTT GGCGCCTCTCCTTTTCCCACACAGAGAAGTACATCCTTAACAACC ACGGAATCGAAAAAACCTGCTGCGAAAGTTCCGGTGCCAAGTGT TGTCGAAAGGAATGCCTTAAACTGATGAAATACCTTTTGGAGCA GCTCAAAAAGGAATTCCAAGAGCTCGACGCTTTTTGCTCTTACCA CGTTAAGACCGCTATTTTCCACATGTGGACACAGGACCCACAAGA TTCTCAATGGGACCCTAGGAATCTCAGCTCATGTTTCGATAAGTT GTTGGCTTTTTTCCTCGAATGTCTGCGCACCGAGAAGTTGGATCA TTACTTCATTCCCAAGTTCAACCTCTTCTCACAGGAACTGATAGAT CGAAAGTCTAAGGAATTCTTGTCTAAAAAAATAGAGTATGAGCG AAACAATGGTTTCCCCATTTTTGACAAGTTGTGAGCGGCCGCGCG GATCCAGACATGATAA	This paper	Ikegami lab ID: EA087
DNA oligonucleotide EA096 for *Cxcl10* qPCR primer: GCCGTCATTTTCTGCCTCA	This paper	Ikegami lab ID: EA096
DNA oligonucleotide EA097 for *Cxcl10* qPCR primer: CGTCCTTGCGAGAGGGATC	This paper	Ikegami lab ID: EA097
DNA oligonucleotide EA098 for *Ifnb1* qPCR primer: CCTGGAGCAGCTGAATGGAA	This paper	Ikegami lab ID: EA098
DNA oligonucleotide EA099 for *Ifnb1* qPCR primer: TGGATGGCAAAGGCAGTGTAA	This paper	Ikegami lab ID: EA099
DNA oligonucleotide EA100 for *Ifit3* qPCR primer: CCTACATAAAGCACCTAGATGGC	This paper	Ikegami lab ID: EA100
DNA oligonucleotide EA101 for *Ifit3* qPCR primer: ATGTGATAGTAGATCCAGGCGT	This paper	Ikegami lab ID: EA101
DNA oligonucleotide KI444 for *Actb* qPCR primer: CTTTGCAGCTCCTTCGTTGC	This paper	Ikegami lab ID: KI444
DNA oligonucleotide KI445 for *Actb* qPCR primer: CCTTCTGACCCATTCCCACC	This paper	Ikegami lab ID: KI445

DNA oligonucleotide KI466 for AAV titering primer: GCGGATCCAGACATGATAAG	This paper	Ikegami lab ID: KI466
DNA oligonucleotide KI467 for AAV titering primer: GCCTATAGTGAGTCGTATTAAG	This paper	Ikegami lab ID: KI467
Recombinant DNA		
pAAV:cTNT::Luciferase	Addgene	Addgene plasmid #69915
pAAV:cTNT::GFP-GS linker	This paper	Ikegami lab ID: bAE263_1
pAAV:cTNT::GFP-icGAS	This paper	Ikegami lab ID: bAE299_1
pAAV:cTNT::cGAS	This paper	Ikegami lab ID: bAE299_3
pLJM1-EGFP	Addgene	Addgene plasmid #19319
Software and algorithms		
ImageJ	N/A	https://imagej.net/ij/
NIS Elements	Nikon Instruments Inc.	RRID:SCR_014329
Odyssey CLx Imaging System	LI-COR Biosciences	https://www.licor.com/bio/odyssey-dlx/
CellProfiler	Broad Institute	https://cellprofiler.org/
Other		
Bio-Rad CFX96 Real-Time PCR Detection System	Bio-Rad	RRID:SCR_018064
Yokogawa CSU-W1 Sora spinning disk confocal microscope	Nikon Instruments Inc.	https://www.microscope.healthcare.nikon.com/products/confocal-microscopes/csu-series/csu-w1
Nikon A1R laser-scanning confocal microscope	Nikon Instruments Inc.	RRID:SCR_020317

FIGURE 1

A Cardiomyocyte-specific Lmna deletion in adult mice

WT: *Lmna^{+/+};Myh6-MerCreMer*
Lmna^{CKO}: *Lmna^{F/F};Myh6-MerCreMer*

Tam 400 µg/day x 4 days

6-8 weeks of age

weeks post Tam

B Abundance of Lamin A/C
Isolated cardiomyocyte at 2 weeks post Tam

Lamin A/GAPDH
Lamin C/GAPDH

C Lamin A/C IHC in hearts
Time post tamoxifen
1 week   2 weeks   3.5 weeks

WT
Lmna^{CKO}

Lamin A/C   DNA

D Heart morphology
Time post tamoxifen
1 week   2 weeks   3.5 weeks

WT
Lmna^{CKO}

E Ejection faction by echocardiography
Wilcoxon test P: 0.9   0.3   0.03   3x10⁻⁴

Ejection Fraction (%)
Weeks post tamoxifen

F Survival analysis
WT (n=48)
Lmna^{CKO} (n=42)
Log-rank test p < 0.0001
Survival probability
Days post tamoxifen

G Transcriptome of left ventricles
Day 7 post Tam   Day 11 post Tam   Day 14 post Tam
Down: 2   Up: 1   Down: 1   Up: 0   Down: 351   Up: 1,752
$-Log_{10}(P\ adj)$
$Log_2[Lmna^{CKO}]/[WT]$

H GO terms for differentially expressed genes at Day 14 post Tam
Up regulated genes
positive reg of cytokine production
reg of cell activation
inflammatory response
neg reg of immune system process
cell activation
reg of immune effector process
immune effector process
leukocyte migration
reg of TNF superfam cytokine product
activation of immune response
$-Log_{10}$(P-value)

Down regulated genes
pos reg of glucose metabolic process
striated muscle adaptation
reg of monoatomic ion transport
monoatom ion transmem transporter act
nuclear steroid receptor act
skeletal muscle adaptation
sulfur compound metabolic process
reg of triglyceride metabolic process
hydro−lyase activity
glucose transmembrane transport
% genes

I Immune-related gene expression at Day 14 post Tam
Cgas   Cxcl10   Ifit1   Ifit3   Ifnb1   Irf3   Irf7   Sting1
$Log_{10}(1 + TPM)$
WT   *Lmna^{CKO}*   *P<0.05, LFC > 1

J Immune cell staining in hearts
CD45   CD68   Merge
WT
Lmna^{CKO}
2 weeks post tamoxifen

K Trichrome staining in hearts
Time post tamoxifen
1 week   2 weeks   3.5 weeks
WT
Lmna^{CKO}

L Lamin A/C reduction in CM causes DCM in adult mice
Lamin A/C reduction in cardiomyocyte → Inflammation-related transcription (2 wk Tam) → Heart functional decline (3 wk Tam)

1124

FIGURE 2

A Lamin A/C immunohistochemistry of heart

B Electronmicroscopy of heart

C Lamin A/C IF in isolated cardiomyocyte

D PCM IF in isolated cardiomyocyte

E Localization of NLS-tdTomato in isolated cardiomyocyte

F Quantification of tdTomato signal

G AAV-mediated gene delivery

H Localization of GFP-icGAS in isolated cardiomyocytes

I GFP-icGAS quantification

J NE rupture frequency

K γ-H2AX quantification

L TUNEL quantification

M NE rupture precedes transcriptional alteration

1125

FIGURE 3

A *cGAS/STING protein expression*

B *GO terms enriched in upregulated genes*

C *AAV-mediated gene delivery*

D *Localization of cGAS in isolated cardiomyocyte*

E *cGAS-STING-related genes*

F *Transcriptome of Cgas⁻/⁻ LmnaCKO hearts*

G *cGAS-STING-related gene expression in Cgas⁻/⁻ LmnaCKO hearts*

H *Ventricular ejection fraction*

I *Trichrome staining in hearts*

J *Survival analysis of Sting⁻/⁻ LmnaCKO mice*

K *NE rupture does not activate cGAS-STING in cardiomyocytes*

FIGURE 4

SUPPLEMENTARY FIGURE S1

A *Cardiomyocyte-restricted Cre expression*

2 weeks post Tam

$Lmna^{+/+;Myh6MCM}+R26^{CAG-LSL-NLS-tdTomato}$

PCM1+DNA · +NLS-tdTomato · +COX4

B *Cre-LoxP mediated inducible Lmna deletion in cardiomyocytes*

Lmna floxed — LoxP — Exon 2 — LoxP
Lmna recombined — Cre

WT: $Lmna^{+/+;Myh6MCM}$
$Lmna^{CKO}$: $Lmna^{F/F;Myh6MCM}$

Cardiomyocyte at 2 weeks post Tam

WT · $Lmna^{CKO}$

650bp / 500bp / 400bp / 200bp / 100bp

598bp (Floxed)
470bp (WT)
153bp (Recombined)

97 82 86
Recombination (%)

C *Lamin A/C immunoblot*

Cardiomyocyte isolated at 2 weeks post Tam

WT · $Lmna^{CKO}$

kDa
75 —
50 —
37 —

Lamin A (74 kDa)
Lamin C (65 kDa)

75 —
50 —
37 —

GAPDH (36 kDa)

Coomassie (post transfer gel)

D *Left ventricular wall thickness by echocardiography*

WT · $Lmna^{CKO}$

Septal wall, systolic (LVAW;s)
P=0.5   0.3   0.4   7x10⁻⁴

Septal wall, diastolic (LVAW;d) (mm)
P=0.7   0.3   0.6   0.8

Free wall, systolic (LVPW;s)
P=0.9   0.6   0.4   4x10⁻⁴

Free wall, diastolic (LVPW;d)
P=0.8   0.3   0.6   0.3

Wall thickness (mm)

Weeks post Tamoxifen   1   2   3   3.5

E *Immune cell quantification in hearts*

WT · $Lmna^{CKO}$

CD45⁺ · CD68⁺ · CD45⁺CD68⁺ · DAPI

Cell count (1000x) per 1.67 mm² area

Wilcoxon test   P=0.008   P=0.016   P=0.008   P=1.0

2 Weeks post Tamoxifen

1128

A *Additional images for EM in hearts*

WT   Day 14 post Tam   *Lmna^{CKO}*

B *Additional images for NLS-tdTomato signals in Lmna^{CKO} CMs*

2 weeks post Tam

WT + *R26^{CAG-LSL-NLS-tdTomato}*   *Lmna^{CKO}* + *R26^{CAG-LSL-NLS-tdTomato}*

PCM1

NLS-tdTomato

Merge

+Phalloidin

C *Localization of cytoplasmic DNA sensor GFP-icGAS in isolated cardiomyocytes*

WT + AAV-GFP-icGAS   2 weeks post Tam

GFP-icGAS   PCM1+DNA   Merge   +Phalloidin

***Lmna^{CKO}*+ AAV-GFP-icGAS**   2 weeks post Tam

GFP-icGAS   PCM1+DNA   Merge   +Phalloidin

D *PCM1 IF in CMs at Day 11 post Tam*

WT   Day 11 post Tam   *Lmna^{CKO}*

PCM1

+DNA

+Phalloidin

E *Number of nuclei with DNA protrusion from PCM1-lost nuclear tips per cell*

		No. nuclei with DNA protrusion from PCM1-lost nuclear tips		
		0	1	≥2
Day 11 post Tam	WT	99% (163)	1% (1)	0% (0)
	Lmna^{CKO}	69% (121)	14% (25)	17% (30)
Day 14 post Tam	WT	100% (163)	0% (0)	0% (0)
	Lmna^{CKO}	46% (78)	25% (42)	29% (49)

% Cells (count)

F *γ-H2AX IF in heart*

WT   *Lmna^{CKO}*

γ-H2AX DNA
Desmin

2 wk post Tam

3.5 wk post Tam

G *TUNEL assay in heart section*

2 weeks post Tam

WT   *Lmna^{CKO}*

TUNEL DNA
Actin

TUNEL

DNaseI + TUNEL

SUPPLEMENTARY FIGURE S3

A *cGAS immunoblot*

B *STING immunoblot*

C *SLAM-ITseq principle*

D *4sU labeling in WT hearts*

E *1,020 bulk RNA−seq upregulated genes detected by SLAM-ITseq*

F *GO terms for differentially expressed genes at Day 14 post Tam*

G *cGAS-STING-related gene expression*

H *Trichrome staining in hearts*

I *Survival analysis of Cgas⁻/⁻ Lmnaᶜᴷᴼ mice*

SUPPLEMENTARY FIGURE S4

A Marker expression in scRNA-seq clusters

B Sample composition within scRNA-seq clusters

C GO terms enriched in downregulated genes

D Select GO term enrichment in upregulated genes

E Cell death related gene expression

F DNA damage repair-related genes