1 Title

Effective mechanical potential of cell–cell interaction in tissues harboring cavity and in

cell sheet toward morphogenesis

5 Short title

Effective potential of cell–cell interaction in 2D and 3D

8 Authors

Hiroshi Koyama[1,2,*], Hisashi Okumura[2,3,4], Tetsuhisa Otani[2,5], Atsushi M. Ito[6], Kazuyuki

Nakamura[7,8], Kagayaki Kato[2,9], and Toshihiko Fujimori[1,2]

[1]Division of Embryology, National Institute for Basic Biology, 5-1 Higashiyama,

Myodaiji, Okazaki, Aichi 444-8787, Japan

[2]SOKENDAI (The Graduate University for Advanced Studies), 5-1 Higashiyama,

Myodaiji, Okazaki, Aichi 444-8787, Japan

[3]Biomolecular Dynamics Simulation Group, Exploratory Research Center on Life and

Living Systems (ExCELLS), National Institutes of Natural Sciences, 5-1 Higashiyama,

Myodaiji, Okazaki, Aichi 444-8787, Japan

[4]Institute for Molecular Science, National Institutes of Natural Sciences, 5-1 Higashiyama,

Myodaiji, Okazaki, Aichi 444-8787, Japan

[5]Division of Cell Structure, National Institute for Physiological Sciences, 5-1

Higashiyama, Myodaiji, Okazaki, Aichi 444-8787, Japan

[6]National Institute for Fusion Science, National Institutes of Natural Sciences, 322-6

Oroshi-cho, Toki, Gifu 509-5292, Japan

[7]School of Interdisciplinary Mathematical Sciences, Meiji University, 4-21-1 Nakano,

Nakano-ku, Tokyo 164-8525, Japan

[8]JST, PRESTO, 4-1-8 Honcho, Kawaguchi, Saitama 332-0012, Japan

[9]Optics and Imaging Facility, Trans-Scale Biology Center, National Institute for Basic

Biology, 38 Nishigonaka, Myodaiji, Okazaki, Aichi 444-8585, Japan

*Correspondence: Hiroshi Koyama

Division of Embryology, National Institute for Basic Biology, 5-1

Higashiyama, Myodaiji, Okazaki, Aichi 444-8787, Japan

Phone: +81 564 59 5862,

hkoyama@nibb.ac.jp.

Conflicts of interest

39   The authors declare no competing financial interests.

Keywords

effective potential of cell–cell interaction, morphological diversity, mechanics, emergent

property of multicellular system, mouse blastocyst, MDCK cyst

This file includes:

Main text

Figure 1 to 8

Abstract

Measurement of mechanical forces of cell–cell interactions is important for studying the

emergence of diverse three-dimensional morphologies of multicellular organisms. We

previously reported an image-based statistical method for inferring the effective

mechanical potentials of pairwise cell–cell interactions, where cell tracking data obtained

by live imaging was fitted by a theoretical model. However, because real tissues harbor

various non-cellular components such as expanding cavities, whether the image-based

method is applicable to such tissue remains elusive. Here we evaluated the applicability

of the method in cavity-harboring tissues. First, we used synthetic data generated by

simulations under cavities with pregiven cell–cell interaction potentials, to which our

inference method was applied. The inferred effective potentials showed a profile modified

from the pregiven potentials, but subsequent analyses revealed that the pregiven

potentials can be reconstructed by substituting the effect of the expanding cavities which

emerged as an additional repulsive component around long-distance of cell–cell

interactions. Interestingly, simulations by using the modified effective potentials

reproduced the cavity-harboring structures, implying that the effect of expanding cavities

was approximately incorporated into the effective pairwise potentials. Then, we applied

our inference method to the mouse blastocyst and cysts of Madin-Darby canine kidney

(MDCK) cells, and found that the inferred effective potentials can reproduce the cavity-

harboring structures. A similar modification of pairwise potentials was also detected in

two-dimensional cell sheets, by which curved sheets such as tubes and cups were

simulated. We conclude that our inference method is applicable to tissues harboring

cavities and cell sheets, and the resultant effective potentials are useful to simulate the

morphologies.

Introduction

In multicellular living systems, various three-dimensional morphologies are

emerged, which is thought to be primarily dependent on the mechanical properties of the

constituent cells (Bazellières et al., 2015; Fletcher et al., 2014; Guillot and Lecuit, 2013;

Maître et al., 2012; Steinberg, 2007). Measurement of cellular mechanical parameters is

inevitable for understanding morphogenetic events. On the other hand, morphogenetic

events are also influenced by non-cellular structures such as cavities, extracellular matrix,

etc (Style et al., 2014; Wang et al., 2018). Therefore, it is challenging to identify all

components contributing to morphogenetic events, and then, to measure their mechanical

forces especially under three-dimensional situations.

Image-based methods for inferring mechanical forces in tissues have been

developed, where mathematical models are assumed which are fitted to microscopic

images (Brodland et al., 2010; Ishihara and Sugimura, 2012; Kondo et al., 2018; Koyama

et al., 2023, 2012; Merkel and Manning, 2017; Miyoshi et al., 2006). These methods

include inference of cell–cell junction tensions in two-dimensional epithelia, cell surface

tensions or bending rigidities, mechanical potential of pairwise cell–cell interactions, etc.

Basically, these methods do not assume non-cellular structures in their mathematical

models, and, it is unknown whether these methods are applicable to tissues with non-

cellular components.

We previously reported a method for inferring mechanical potential of pairwise

cell–cell interactions (Koyama et al., 2023). The pairwise potentials are composed of

attractive and repulsive forces of cell–cell interactions. The attractive forces are at least

partially derived from cell–cell adhesion forces mediated by cell adhesion molecules (Fig.

1A-B), whereas the repulsive forces are from excluded volume effect of cells (Drasdo et

al., 1995; Forgacs and Newman, 2005; Guillot and Lecuit, 2013; Maître et al., 2012;

Merkel and Manning, 2017; Steinberg, 2007; Yong et al., 2015). The summation of these

forces would be dependent on cell–cell distance (Fig. 1C), by which the pairwise potential

energy is calculated. We previously showed that, by applying our method to the mouse

and *C. elegans* early embryos which did not still harbor clear cavities or thick extracellular

matrix, we obtained mechanical potential of pairwise interactions (Koyama et al., 2023).

The profiles of the pairwise potentials were quantitatively different among the tissues, by

which the morphologic features of the corresponding tissues were reproduced.

In the present study, we evaluated whether our inference method is applicable to

cavity-harboring tissues such as the mouse blastocysts (Fig. 1D). First, we theoretically

analyzed the effect of liquid cavities by using synthetic data which were generated by

simulations under pregiven cell–cell forces and cavities. In the case where the cavities are

expanding, the inferred effective pairwise forces of cell–cell interactions became

significantly different from the pregiven forces, where an additional repulsive component

was detected. But we showed that the repulsive component was derived from the

expanding cavities, and that the inferred effective pairwise forces were the summation of

both the pregiven forces and the effect of the expanding cavities. Therefore, simulations

using inferred pairwise forces successfully reproduced the cavity-harboring structures.

Application of our method to the mouse blastocyst and the cysts of MDCK cells also

resulted in the same conclusion. Finally, we systematically analyzed what kind of profiles

of the effective pairwise potentials are critical for forming cavity-harboring structures,

which were presented as phase diagrams, and expanded the analyses to curved sheets.

 Results

Inference of effective pairwise potential in synthetic data with cavity

In our inference method, a cell-particle model is assumed where

attractive/repulsive forces of cell–cell interaction is considered (Fig. 1B-C), and the

mathematical model is fitted to nuclear tracking data (i.e., *xyz*-coordinates of each nucleus

with the time series) as described previously (Koyama et al., 2023). Briefly, the fitting

procedure is based on a minimization of the differences between the *xyz*-coordinates of

particles in the mathematical model and in the nuclear tracking data, with an additional

cost function for cell–cell distance-dependent weight. The fitting parameters are the force

values for each pair of two interacting cells for each time frame, and then, by binned

averaging the inferred forces along cell–cell distances, a distance–force curve is

calculated. In the model to be fitted, we do not explicitly consider non-cellular structures

exerting forces on cells. It can be argued that, if tissues harbor non-cellular structures, the

model should assume the non-cellular structures. But, due to high degrees of freedom in

such models, the fitting problems would become difficult to be solved in general. We

intended to analyze whether valuable information can be obtained when our inference

method is applied to tissues with cavities (Fig. 1D). To this aim, we first generated

simulation data considering both cell–cell interaction forces and cavities, to which we

applied our inference method.

In our simulations based on particulate cells, the cell–cell interaction forces were

set to be provided by the Lenard-Jones (LJ) potential (i.e., the pregiven forces) in a similar

manner to our previous work (Koyama et al., 2023), and expanding cavities was also

provided. In real tissues, cavities are generated by secreting liquid from cells. The cell

layers surrounding the cavities are sealed by the tight junctions, and thus the pressure of

the cavities push the cells outward (Fig. 1D) (Dumortier et al., 2019; Moriwaki et al.,

2007; Wang et al., 2018). In simulations, we implemented expanding spherical cavities

as temporally-evolved spatial constraints; the particulate cells cannot penetrate into the

cavities (Fig. S1A-B and movieS1), leading to that the particles move on the surfaces of

the cavities.

Fig. 2A-I shows the snapshot of the simulations where the spherical cavity was

expanding from the radius = 9.6 to 12.6μm. Fig. 2A-ii shows inferred effective distance–

force (DF) curves under three different conditions of the expanding cavities; the inferred

DF curves were not consistent with the DF curves from the LJ potential (Fig. 2A-ii,

compare with the black broken line), where repulsive components were strongly detected

especially at distant regions, which we call distant energy barrier (DEB). The strength of

the repulsive components seemed to be positively related to the expansion rates. In

contrast, under the conditions of non-expanding cavities, the inferred DF curves were

consistent with the LJ potential (Fig. S1C).

We then examined whether this modification is meaningless or is a result of a

correct reflection of the expanding cavities. By performing simulations under conditions

with the expanding cavities but without the LJ potential (i.e., without cell–cell interaction

forces) (Fig. 2B-i), we solely extracted the influence of the expanding cavities. Fig. 2B-

ii shows that the inferred effective DF curves were absolutely composed of repulsive

components. Then, we checked the raw data where all data points were plotted on the

distance–force space; in Fig. 2C-i, "From B", a heat map of the data points is shown.

Force values calculated by binned averaging are also shown in yellow, which correspond

to the DF curve in Fig. 2B-ii (red line). Interestingly, all the data points were closely

located around the DF curve ("From B"), implying that the DF curve is a good

approximation of the expanding cavities. Probably, due to the effect of the expanding

cavities which causes the increases in distances among the particles, pairwise forces of

particle–particle interactions became apparently repulsive.

Finally, by using the above information, we examined whether the correct DF

curve (i.e., LJ potential) can be reconstructed. We subtracted the DF curves in Fig. 2B

from those in Fig. 2A (Fig. 2C-i, "From A – From B"), and the resultant DF curves were

well consistent with the LJ potential under any expansion rates of the cavities (Fig. 2C-

ii). These results suggest that the effects of the expanding cavities can be addible to

effective DF curves.

We interpret the above results as described in Fig. 2D. A cell receives forces from

cell–cell interactions ("LJ" in Fig. 2D) and from the pressure of the cavity, where the

vector of the pressure is perpendicular to the surface of the cavity (Fig. 2D, top panel).

The repulsive forces detected as cell–cell interactions in Fig. 2A are described in Fig. 2D,

middle panel (light blue vectors), and the vector of the net force (a red vector) for a cell

is perpendicular to the surface of the cavity, if the cells are almost uniformly distributed.

Therefore, the pressure from the cavity can be equivalent to the net force from the DEBs

(Fig. 2D, bottom panel), leading to the successful reconstruction of the LJ potential.

We conclude that the expanding cavities can be approximated as effective

pairwise potentials. In general, effective pairwise forces/potentials are the summation of

different components as explained in Fig. 1C. External factors are one of the components

to be summed up in molecular and colloidal sciences: e.g., by considering hydration of

particles by solvents, effective pairwise potentials are modified so as to contain energy

barriers around distant regions (Israelachvili, 2011; Pettitt and Rossky, 1986), and such

effective potentials have been used to simulate behaviors of the systems. We will show

the simulation outcome from the above inferred effective DF curve later.

Effective forces in mouse blastocyst contained DEB

To investigate whether a repulsive component appeared in effective forces

inferred from the mouse blastocyst (Fig. 3A, illustration). An inner cavity is formed

through liquid secretion from the cells, and the embryos expand while maintaining their

spherical shape (Dumortier et al., 2019; Moriwaki et al., 2007; Wang et al., 2018) (Fig.

3A, bright field and cross-section). Trophectoderm (TE) cells form a single-cell-layered

structure at the outermost surface of the embryo, whereas the cells of the inner cell mass

(ICM) were tightly assembled, forming a cluster (Fig. 3A, illustration). We performed

cell tracking (Fig. 3A, tracking) and inferred effective forces.

Fig. 3B shows the inferred effective DF curve from the blastocyst. The DF curve

contained the repulsive component around distant regions (Fig. 3B, DEB). For

comparison, an inferred DF curve from an embryo before forming a cavity is shown,

which did not contain DEB (Fig. 3B, "compaction" meaning the embryo at the

compaction stage). In addition, Fig. 3C shows the distance–potential (DP) curves which

can be calculated from the DF curves, and the DEB appeared as a long slope.

To further examine whether the DEB was derived from the cavity, we inhibited

the cavitation by using an inhibitor for Na$^+$/H$^+$ exchanger (S3226) (Kawagishi et al.,

2004). Fig. 3D shows the inhibitor-treated embryos; the experimental procedures are

shown in Fig. S2. Fig. 3D shows the inferred effective DP curves, where DEB was

diminished, indicating that the cavity yielded the DEBs.

Effective forces in MDCK cyst contained DEB

We also applied our method to other tissues harboring cavities. MDCK cells can

form cysts containing a cavity when cultured under suspension conditions (Fig. 4A)

(Wang et al., 1990). Fig. 4B and 4C show the effective DF and DP curves, where a DEB

was detected.

Effective potentials reproduced cavity-harboring structures

Effective DF/DP curves are the summation of the multiple components including

the excluded volume effect, adhesive forces, and external factors, and are useful to study

behaviors of systems in molecular and colloidal sciences. We evaluated whether inferred

effective DF/DP curves from the cavity-harboring systems can reproduce their

morphologies. The effective DF curves inferred from the mouse blastocyst, the MDCK

cyst, and the artificial tissue harboring a cavity (Fig. 2-4) were used for simulations, and

the systems were relaxed to find stable states. When initial configurations of particles

were set to be cavity-harboring structures (Fig. 5A), the structures were maintained (Fig.

5B-D, left panels). To examine whether the DEBs contribute to the formation of the

cavities, we artificially eliminated the DEBs from the DF curves, and performed

simulations. We found that cavities were not maintained but aggregates — filled with the

particles — were formed (Fig. 5B-C, "DF curve w/o DEB"). For comparison, we used

the Lenard-Jones potential, and showed that the cavity-harboring structure was not

maintained but became an aggregate (Fig. 5E).

Modeling of distance–force curves with two parameters

To comprehensively understand the effect of the differences of the profiles in

the DF curves on morphogenesis, we modeled the DF curves as a simple mathematical

equation. To analyze the contribution of DEBs, we selected the following equation (Fig.

6A, left panel, blue line):

$$F(D) = \varepsilon(D - D_0)^{-N} \cos\left\{ \frac{2\pi(D - D_0)}{\delta} \right\}, \text{ for } D > D_0 \qquad \text{(Eq. 1)}$$

Here, F is the force, D is the distance. N is the exponent of D, affecting the decay of the

DF curves along D (X-axis) (Fig. 6A, left panel, green broken line). To model DEB, we

introduced a cosine function whose wavelength is affected by δ; the combination of the

above exponential function (Fig. 6A, left panel, green broken line) with the cosine

function (Fig. 6A, left panel, purple broken line) resulted in a function bearing DEB (Fig.

6A, left panel, red line). D_0 can translate the DF curves along D, (Fig. 6A, left panel, red

vs. blue lines). This translation simultaneously modifies the relative positions of DEBs to

the mean diameters (Fig. 6A, left panel, where D_1, mean diameter; D_2, distance at DEB;

$D_3 = D_2/ D_1$, the relative positions of DEB). N also affects the strengths of DEBs (Fig.

6A, right panel, F_{rep}/F_{att} is the relative strength of DEB).

All parameters and variables related to lengths/distances were normalized by

$(\delta/4)$, leading to generation of dimensionless parameters and variables such as $D_0^* = D_0$

$/(\delta/4)$. In addition, ε does not contribute to morphologies under stable states. Therefore,

we can present simulation outcomes in two-dimensional space (D_0^* and N) in a manner

similar to a phase diagram (Fig. 6B) as shown in the next section. Although there can be

other equations modeling DF curves, the advantage of this equation is that it is composed

of just two parameters (D_0^* and N).

Appropriate position and strength of DEB were essential for cavity-harboring structures

First, we examined what kind of DF curves can maintain a cavity-harboring

structures. In simulations of multicellular systems, the morphologies are determined by

both DF curves and initial configurations, because energetic local minimum states (i.e.

metastable state) as well as global minimum states are meaningful. A cavity-harboring

structure was set as the initial configuration (Fig. 6B, "Initial configuration"), and

simulations were performed until the systems reached steady states including metastable

states.

Fig. 6B shows a phase diagram with representative images of the simulations,

on which two regions of cavity-harboring structures were identified ("cavity" and "cavity

with extra particles"). The "cavity" class absolutely depended on the DEBs, because DF

curves whose DEBs were eliminated did not form the "cavity" as shown in Fig. 6C ("w/o

DEB"). We confirmed that the cavity was not broken by slight positional fluctuations

(Movie S2), indicating that the structures were stable. In Fig. 6B, the "cavity" class was

located around smaller N values (less than ~2.0), otherwise the "aggregate" class was

formed, suggesting that the strengths of the DEBs, defined in Fig. 6A (right panel), may

be effective. Furthermore, larger D_0 values were required for the "cavity" class, otherwise

the "multiple clusters/disorganized" class was formed, suggesting that the relative

positions of DEBs ($D_3 = D_2/ D_1$ in Fig. 6A, left panel) should be close to the mean

diameters. In addition, the "cups" and "tubes" classes were found between the "cavity"

and the "multiple clusters/disorganized" classes in Fig. 6B but not in Fig. 6C, indicating

that the DEBs also contribute to the formation of these structures. Under a different

number of particles, a similar trend was observed (Fig. S4). The maintenance of tubes

was confirmed to be dependent on DEBs (Fig. S5).

Profiles of DF curves for describing 2D sheet

To further investigate the capability of DF curves, we next applied a two-

dimensional (2D) sheet as the initial configuration which corresponds to a monolayered

cell sheet (Fig. 7A, "Initial configuration"). Fig. 7A shows a phase diagram with

representative simulation images. The 2D sheets were maintained (the "sheet" class),

which was largely dependent on the presence of DEBs (Fig. 6A, w/ DEBs vs. 6B, w/o

DEBs). We confirmed that the 2D sheets were not broken by slight positional fluctuations

(Movie S2), indicating that the structures were stable. The region of the "sheet" class was

located around larger values of D_0, which was a similar trend to the case of the cavity in

Fig. 6. In addition, the cavity-harboring structures were newly generated even in the

absence of the DEBs (Fig. 7B). By temporally changing N, D_0, and existence of DEB, we

presented possible morphological transitions among the classes in Fig. 6 and 7 (Fig. S6

and movie S3).

Profiles of DF curves for describing spherical/non-spherical aggregates

In Fig. 7, we found that both the "spherical" and "non-spherical aggregate"

classes were generated w/ and w/o DEBs. Smaller values of N and/or D_0 led to the

spherical ones (Fig. 7B). Under smaller values of D_0, because the width (W_1 ; see the

bottom row in Fig. 7) of the attractive force regions in DF curves became relatively long

compared to the mean diameter (D_1, in Fig. 7), the resultant increased attractive forces

may cause the spherical aggregates. Under smaller values of N, the potential energies (P_1,

in Fig. 7) derived from the attractive forces are increased, possibly leading to the spherical

aggregates.

Effective forces in 2D sheets included DEB

As shown in Fig. 7, 2D sheets were maintained under the DF curves including

DEB. Similarly, the cups and tubes in Fig. 6 were also maintained under the DF curves

including DEBs; these structures can be interpreted as a curved sheet. Is the relationship

between these structures and DEB an artifact or are there any reasonable explanations?

To investigate this issue, we used synthetic 2D sheets generated by simulations

(Fig. 8A-i). The 2D simulations were performed under the Lenard-Jones (LJ) potential as

the pregiven force, and the DF curves were inferred. As shown in Fig. 8A-i, the inferred

DF curve was similar to the LJ potential, but the curve contained a clear DEB. This was

contrast to the inferred DF curve from the simulation of a 3D aggregate, where DEB was

rarely detected (Fig. 8A-ii).

We hypothesized that the radial distribution functions (RDF) between 2D and

3D situations were different, because RDFs are related to the profile of distance–potential

curves in atomic/molecular/colloidal sciences (Israelachvili, 2011). RDFs mean the

density of particles at a certain distance from a particle of interest (Fig. 8B). The density

may be reflected as the effective potential energy at the distance; high density corresponds

to low potential energy (Fig. 8B). In other words, by using RDFs, effective pairwise

potentials are calculated if the systems are under equilibrium conditions.

We calculated RDFs in 2D and 3D simulations performed under the LJ potential

(Fig. 8C). In these simulations, fluctuations of the forces of cell–cell interactions were

assumed with the persistency of the fluctuations (the details are described in

Supplementary information). Under the 2D situations, clear peaks were found where the

particle density was high. Importantly, between the peaks, the density was very low (Fig.

8C-i, black arrow). By contrast, under the 3D simulations, relatively unclear peaks were

detected, and the regions with low particle densities almost disappeared (Fig. 8C-ii).

Additionally, even under the 2D simulations, the longer persistency caused disappearance

of the regions with low particle densities, because the long persistency promotes the

movements of the particles. These results suggest that 2D sheets spontaneously yield

regions with low particle density in RDF, probably due to the geometric reasons, and that

this feature can be reflected to the effective pairwise potentials as DEB.

Discussion

In this study, by inferring effective forces of cell–cell interactions in tissues

harboring expanding cavities, we showed that the effect of the cavities was addible to the

effective forces of cell–cell interactions as a repulsive component (DEB). The resultant

distance–force curves reproduced the tissue structures harboring cavities in simulations.

A similar repulsive component was also detected in two-dimension (2D) cell sheets,

which may be derived from geometric reasons of 2D systems. Finally, by assuming

distance–force curves with the repulsive components, tissue structures of 2D sheets, cups,

and tubes were reproduced in simulations. These results suggest that external components

and geometric constraints can be correctly added to the effective forces of cell–cell

interactions. This framework considering the repulsive components can be useful to

simulate and study morphogenesis of 2D sheets and their derivatives (i.e., cavity-

harboring structures, cups, and tubes as curved sheets) (Belmonte et al., 2016; Eiraku et

al., 2011; Honda et al., 2008b; Koyama et al., 2016; Montesano et al., 1991; Rivron et al.,

2018; Serra et al., 2019; Solnica-krezel and Sepich, 2012; Wang et al., 1990).

Influence of external factors on effective potentials

Our results suggest that expanding cavities and geometric features of 2D sheet

can be incorporated into the effective pairwise potentials. In atomic/molecular/colloidal

sciences, it is known that some external factors are addible to effective pairwise potentials.

For instance, particles such as ions and colloids are hydrated by water molecules as

solvents, resulting in that the effective pairwise potentials of the particles contain

repulsive components at distant regions, and the resultant potentials explain the behaviors

of the systems (Israelachvili, 2011). External pressures can modify RDF by increasing

density of the particles, leading to the modification of the effective pairwise potentials. In

multicellular systems, it is unknown what kind of external factors are addible to the

effective pairwise potentials. We speculate that, if external factors uniformly exert forces

on cells, the effect may be correctly incorporated into the effective potentials.

Comparison of other inference methods with other models

To understand morphogenetic events, simulations based on mechanical

parameters are essential. Models such as vertex, Cellular Potts, and phase field models

are often utilized, especially for epithelial cells (Belmonte et al., 2016; Camley et al.,

2014; Fletcher et al., 2014; Honda et al., 2008a; Nonomura, 2012). These models

successfully recapitulated various morphogenetic events, including the formation of the

mouse blastocyst (Honda et al., 2008a). However, it is still challenging to measure cellular

and non-cellular parameters in three-dimensional systems with high spatiotemporal

resolution. Image-based force inference methods are important techniques (Ishihara and

Sugimura, 2012; Tambe et al., 2011). However, in the case that tissues harbor non-cellular

factors, it becomes difficult to infer due to increased degrees of freedom. Our idea can

reduce the degrees of freedom, and may be adapted to other image-based inference

methods when non-cellular factors exist. Our goal is to develop a framework for

explaining diverse morphogenesis in a minimal model. Approximations of some external

factors as effective potentials of cell–cell interaction serve to reduce degrees of freedom.

Materials and methods

Mouse embryos

Embryos were obtained after mating homozygous R26-H2B-EGFP knock-in male

mice which constitutively express EGFP (enhanced green fluorescent protein)-fused H2B

(histone2B proteins) (Kurotaki et al., 2007) and ICR female mice (Japan SLC). The

embryos were cultured in EmbryoMax KSOM +AA with D-Glucose (Millipore, USA)

covered with mineral oil on a glass bottom dish (35-mm; 27-mm ϕ, Matsunami, Japan) at

37°C, and subjected to microscopic imaging. Four distinct embryos were analyzed.

In the experiments using S3226 (Fig. S9), the E2.5 embryos were transferred to the

medium with the drug. After 15 hours culture, some embryos failed to form cavities (Fig.

S9, #1 and #2), while others formed smaller cavities (Fig. S9, #3) than that under no-drug

conditions (Fig. S9, No drugs). Live imaging was performed, and then a rescue process

was carried out, resulting in that the embryos formed, at least, smaller cavities compared

with that under no-drug conditions. The concentration of S3226 was 50μM.

Animal care and experiments were conducted in accordance with National Institutes

of Natural Sciences (NINS), the Guidelines of Animal Experimentation. The animal

experiments were approved by The Institutional Animal Care and Use Committee of

NINS (ref. #17A030, #18A026, #19A021, #20A014, and #21A039).

MDCK cysts

PiggyBac transgenic MDCK cells which constitutively express H2B-EGFP were

constructed as follows. To obtain a DNA fragment bearing H2B-EGFP, pcDNA3-H2B-

EGFP (Abe et al., 2011) was digested with *HindIII*, blunted, and then digested with *NotI*.

pEBMulti-Puro (Wako, Japan) was digested with *BamHI*, blunted, and then digested with

NotI. Then, the DNA fragment bearing H2B-EGFP was inserted into the pEBMulti-Puro

backbone. From this plasmid, a DNA fragment harboring H2B-EGFP was excised again

by *XhoI* and *NotI* digestions, and was then inserted between the *XhoI* and *NotI* sites of

pPB-CAG-BstXI-IP (a gift from Dr. Hitoshi Niwa)(Nakatake et al., 2013). The resultant

plasmid, pPB-CAG-IP-H2B-EGFP, and the PBase-expressing vector pCAGPBase (Guo

et al., 2009) were co-transfected into MDCK cells using Lipofectamine LTX (Invitrogen),

and the transformants were selected in the presence of puromycin (1 μg/mL). The

transformants were expected to express H2B-EGFP under control of the constitutive CAG

promoter.

To generate and culture cysts of MDCK cells harboring H2B-EGFP, MDCK-

conditioned DMEM (Dulbecco's Modified Eagle's Medium containing 10% fetal bovine

serum with $NaHCO_3$, Sigma) was used. The conditioned medium was obtained as

follows: MDCK cells were cultured in 10 mL DMEM on a 95-mm dish for 4 days at 37°C

to reach ~90% confluence, 10 mL fresh DMEM was added, and the cells were cultured

for an additional day. The supernatant was obtained, and filtered through a 0.22-μm

Millipore filter (Millex-GV SLGV033RS, Millipore, USA), yielding conditioned

medium.

The cysts were generated and cultured as follows: MDCK cell suspension in 80%

MDCK-conditioned DMEM and 20% DMEM was placed on glass-bottom dishes coated

with 0.4% agarose (SeaKem GTG agarose, BMA). The cells were cultured for 4–5 days

at 37°C, and then subjected to microscopic imaging. Four distinct cysts were analyzed.

MDCK cells were described previously (Otani et al., 2019), and were originally

provided from Masayuki Murata (University of Tokyo). The identity of the cell line

(derivation from canine) was confirmed by genomic sequencing of targeted loci. The

mycoplasma was removed, and the removal was confirmed by Minerva biolabs Venor

GeM OneStep Mycoplasma detection Kit for Conventional PCR.

Microscopic live imaging

Fluorescence images were acquired on a Nikon A1 laser scanning confocal

microscope (Nikon, Japan) equipped with a 20× objective (PlanApo; Dry; NA=0.75,

Nikon, Japan). A stage-top CO_2 incubation system was used (INUG2-TIZ, Tokai Hit,

Japan) on the inverted microscope. H2B-EGFP was excited using a 488-nm laser.

For imaging of mouse embryos, 50-100 frames were acquired at 3-min intervals in

30–40 z-slices separated by 2.5 µm. For imaging of MDCK cysts, 50-100 frames were

acquired at 10-min intervals in 20–30 z-slices separated by 2.0 µm.

The confocal images obtained from the mouse embryos and MDCK cells were

subjected to the procedure for nuclear detection and tracking. Using the Imaris software

(Oxford instruments/Bitplane, UK), the nuclei were automatically detected, followed by

manual corrections. The nuclear tracking was also performed automatically followed by

manual corrections. Through the manual corrections, the accuracies of the nuclear

detection and tracking became essentially 100%, to the best of our judgment.

Inference method and simulation procedures

The inference method is as we previously reported (Koyama et al., 2023). Simulation

procedures are also as previously reported (Koyama et al., 2023), except that cavities are

considered. Details of the modeling of the cavities are described in Supplementary

information.

Acknowledgement

We thank Drs. Hitoshi Niwa and Yayoi Toyooka for providing the piggy-bac plasmids

and technical suggestions. This work was supported by following grants: Japan Ministry

of Education, Culture, Sports, Science and Technology Grant-in-Aid for Scientific

Research on Innovative Areas "Cross-talk between moving cells and microenvironment

as a basis of emerging order" for H.K., the National Institutes of Natural Sciences (NINS)

program for cross-disciplinary science study for H.K., a Japan Society for the Promotion

of Science (JSPS) Grant-in-Aid for Young Scientists (B) for H.K. (17K15131), for

Scientific Research (C) for T.O. (18K06234), and for Young Scientists (B) for T.O.

(16K18544), a MEXT/JSPS Grant-in-Aid for Scientific Research on Innovative Areas for

T.O. (17H05627), the Inamori Foundation for T.O., and the Takeda Science Foundation

for T.O.

References

Abe, T., Kiyonari, H., Shioi, G., Inoue, K.I., Nakao, K., Aizawa, S., Fujimori, T., 2011.

Establishment of conditional reporter mouse lines at ROSA26 locus for live cell

imaging. Genesis 49, 579–590. https://doi.org/10.1002/dvg.20753

Bazellières, E., Conte, V., Elosegui-artola, A., Serra-picamal, X., Bintanel-morcillo, M.,

Roca-cusachs, P., Muñoz, J.J., Sales-pardo, M., Guimerà, R., Trepat, X., 2015.

Control of cell–cell forces and collective cell dynamics by the intercellular

adhesome. Nat. Cell Biol. 17, 409–420. https://doi.org/10.1038/ncb3135

Belmonte, J.M., Clendenon, S.G., Oliveira, G.M., Swat, M.H., Greene, E. V.,

Jeyaraman, S., Glazier, J.A., Bacallao, R.L., 2016. Virtual-tissue computer

simulations define the roles of cell adhesion and proliferation in the onset of

kidney cystic disease. Mol. Biol. Cell 27, 3673–3685.

https://doi.org/10.1091/mbc.e16-01-0059

Brodland, G.W., Conte, V., Cranston, P.G., Veldhuis, J., Narasimhan, S., Hutson, M.S.,

Jacinto, A., Ulrich, F., Baum, B., Miodownik, M., 2010. Video force microscopy

reveals the mechanics of ventral furrow invagination in Drosophila. Proc. Natl.

Acad. Sci. U. S. A. 107, 22111–22116. https://doi.org/10.1073/pnas.1006591107

Camley, B.A., Zhang, Y., Zhao, Y., Li, B., Ben-Jacob, E., Levine, H., Rappel, W.J.,

2014. Polarity mechanisms such as contact inhibition of locomotion regulate

persistent rotational motion of mammalian cells on micropatterns. Proc. Natl.

Acad. Sci. U. S. A. 111, 14770–14775. https://doi.org/10.1073/pnas.1414498111

Drasdo, D., Kree, R., McCaskill, J.S., 1995. Monte Carlo approach to tissue-cell

populations. Phys. Rev. E 52, 6635–6657.

Dumortier, J.G., Verge-Serandour, M. Le, Tortorelli, A.F., Mielke, A., Plater, L. de,

Turlier, H., Maître, J.-L., 2019. Hydraulic fracturing and active coarsening position

the lumen of the mouse blastocyst. Science (80-.). 365, 465–468.

Eiraku, M., Takata, N., Ishibashi, H., Kawada, M., Sakakura, E., Okuda, S., Sekiguchi,

K., Adachi, T., Sasai, Y., 2011. Self-organizing optic-cup morphogenesis in three-

dimensional culture. Nature 472, 51–6. https://doi.org/10.1038/nature09941

Fletcher, A.G., Osterfield, M., Baker, R.E., Shvartsman, S.Y., 2014. Vertex models of

epithelial morphogenesis. Biophys. J. 106, 2291–2304.

https://doi.org/10.1016/j.bpj.2013.11.4498

Forgacs, G., Newman, S.A., 2005. Biological Physics of the Developing Embryo.

Cambridge University Press.

Guillot, C., Lecuit, T., 2013. Mechanics of epithelial tissue homeostasis and

morphogenesis. Science (80-.). 340, 1185–9.

https://doi.org/10.1126/science.1235249

Guo, G., Yang, J., Nichols, J., Hall, J.S., Eyres, I., Mansfield, W., Smith, A., 2009. Klf4

reverts developmentally programmed restriction of ground state pluripotency.

Development 136, 1063–1069. https://doi.org/10.1242/dev.030957

Honda, H., Motosugi, N., Nagai, T., Tanemura, M., Hiiragi, T., 2008a. Computer

simulation of emerging asymmetry in the mouse blastocyst. Development 135,

1407–1414. https://doi.org/10.1242/dev.014555

Honda, H., Nagai, T., Tanemura, M., 2008b. Two different mechanisms of planar cell

intercalation leading to tissue elongation. Dev. Dyn. 237, 1826–1836.

https://doi.org/10.1002/dvdy.21609

Ishihara, S., Sugimura, K., 2012. Bayesian inference of force dynamics during

morphogenesis. J. Theor. Biol. 313, 201–211.

https://doi.org/10.1016/j.jtbi.2012.08.017

Israelachvili, J.N., 2011. Intermolecular and Surface Forces, 3rd ed. ed. Elsevier Inc.

Kawagishi, R., Tahara, M., Sawada, K., Morishige, K., Sakata, M., Tasaka, K., Murata,

Y., 2004. Na+/H+ exchanger-3 is involved in mouse blastocyst formation. J. Exp.

Zool. Part A Comp. Exp. Biol. 301, 767–775. https://doi.org/10.1002/jez.a.90

Kondo, Y., Aoki, K., Ishii, S., 2018. Inverse tissue mechanics of cell monolayer

expansion. PLoS Comput. Biol. 1–15.

Koyama, H., Okumura, H., Ito, A.M., Nakamura, K., Otani, T., Kato, K., Fujimori, T.,

2023. Effective mechanical potential of cell–cell interaction explains three-

dimensional morphologies during early embryogenesis. PLoS Comput. Biol. 19,

e1011306. https://doi.org/10.1371/journal.pcbi.1011306

Koyama, H., Shi, D., Suzuki, M., Ueno, N., Uemura, T., Fujimori, T., 2016. Mechanical

Regulation of Three-Dimensional Epithelial Fold Pattern Formation in the Mouse

Oviduct. Biophys. J. 111, 650–665. https://doi.org/10.1016/j.bpj.2016.06.032

Koyama, H., Umeda, T., Nakamura, K., Higuchi, T., Kimura, A., 2012. A high-

resolution shape fitting and simulation demonstrated equatorial cell surface

softening during cytokinesis and its promotive role in cytokinesis. PLoS One 7,

e31607. https://doi.org/10.1371/journal.pone.0031607

Kurotaki, Y., Hatta, K., Nakao, K., Nabeshima, Y.-I., Fujimori, T., 2007. Blastocyst

Axis Is Specified Independently of Early Cell Lineage But Aligns with the ZP

Shape. Science 316, 719–23. https://doi.org/10.1126/science.1138591

Maître, J.-L., Berthoumieux, H., Krens, S.F.G., Salbreux, G., Jülicher, F., Paluch, E.,

Heisenberg, C.P., 2012. Adhesion Functions in Cell Sorting by Mechanically

Coupling the Cortices of Adhering Cells. Science (80-.). 338, 253–256.

https://doi.org/10.1126/science.1225399

Merkel, M., Manning, M.L., 2017. Using cell deformation and motion to predict forces

and collective behavior in morphogenesis. Semin. Cell Dev. Biol. 67, 161–169.

https://doi.org/10.1016/j.semcdb.2016.07.029

Miyoshi, H., Satoh, S.K., Yamada, E., Hamaguchi, Y., 2006. Temporal change in local

forces and total force all over the surface of the sea urchin egg during cytokinesis.

Cell Motil. Cytoskeleton 63, 208–221. https://doi.org/10.1002/cm.20118

Montesano, R., Schaller, G., Orci, L., 1991. Induction of epithelial tubular

morphogenesis in vitro by fibroblast-derived soluble factors. Cell 66, 697–711.

Moriwaki, K., Tsukita, S., Furuse, M., 2007. Tight junctions containing claudin 4 and 6

are essential for blastocyst formation in preimplantation mouse embryos. Dev Biol

312, 509–22.

Nakatake, Y., Fujii, S., Masui, S., Sugimoto, T., Torikai-Nishikawa, S., Adachi, K.,

Niwa, H., 2013. Kinetics of drug selection systems in mouse embryonic stem cells.

BMC Biotechnol. 13.

Nonomura, M., 2012. Study on multicellular systems using a phase field model. PLoS

One 7, 1–9. https://doi.org/10.1371/journal.pone.0033501

Otani, T., Nguyen, T.P., Tokuda, S., Sugihara, K., Sugawara, T., Furuse, K., Miura, T.,

Ebnet, K., Furuse, M., 2019. Claudins and JAM-A coordinately regulate tight

junction formation and epithelial polarity. J. Cell Biol.

https://doi.org/10.1083/jcb.201812157

Pettitt, B.M., Rossky, P.J., 1986. Alkali halides in water: Ion–solvent correlations and

ion–ion potentials of mean force at infinite dilution. J. Chem. Phys. 84, 5836–

5844. https://doi.org/10.1063/1.449894

Rivron, N.C., Frias-Aldeguer, J., Vrij, E.J., Boisset, J.C., Korving, J., Vivié, J.,

Truckenmüller, R.K., Van Oudenaarden, A., Van Blitterswijk, C.A., Geijsen, N.,

2018. Blastocyst-like structures generated solely from stem cells. Nature 557, 106–

111. https://doi.org/10.1038/s41586-018-0051-0

Serra, D., Mayr, U., Boni, A., Lukonin, I., Rempfler, M., Meylan, L.C., Stadler, M.B.,

Strnad, P., Papasaikas, P., Vischi, D., Waldt, A., Roma, G., Liberali, P., 2019. Self-

organization and symmetry breaking in intestinal organoid development. Nature

569, 66–72.

Solnica-krezel, L., Sepich, D.S., 2012. Gastrulation: Making and Shaping Germ Layers.

Annu. Rev. Cell Dev. Biol. 28, 687–717. https://doi.org/10.1146/annurev-cellbio-

092910-154043

Steinberg, M.S., 2007. Differential adhesion in morphogenesis : a modern view. Curr.

Opin. Genet. Dev. 17, 1281–286. https://doi.org/10.1016/j.gde.2007.05.002

Style, R.W., Boltyanskiy, R., German, G.K., Hyland, C., Macminn, C.W., Mertz, A.F.,

Wilen, L.A., Xu, Y., Dufresne, E.R., 2014. Traction force microscopy in physics

and biology. Soft Matter 10, 4047–4055. https://doi.org/10.1039/c4sm00264d

Tambe, D.T., Hardin, C.C., Angelini, T.E., Rajendran, K., Park, C.Y., Serra-Picamal,

X., Zhou, E.H., Zaman, M.H., Butler, J.P., Weitz, D.A., Fredberg, J.J., Trepat, X.,

2011. Collective cell guidance by cooperative intercellular forces. Nat. Mater. 10,

469–75. https://doi.org/10.1038/nmat3025

Wang, A.Z., Ojakian, G.K., Nelson, W.J., 1990. Steps in the morphogenesis of a

polarized epithelium. J. Cell Sci. 95, 137–152.

Wang, X., Zhang, Z., Tao, H., Liu, J., Hopyan, S., Sun, Y., 2018. Characterizing Inner

Pressure and Stiffness of Trophoblast and Inner Cell Mass of Blastocysts. Biophys.

J. 115, 2443–2450. https://doi.org/10.1016/j.bpj.2018.11.008

Yong, J., Moeller, J., Hart, K.C., Ramallo, D., Vogel, V., Dunn, A.R., Nelson, W.J.,

Pruitt, B.L., 2015. Spatial distribution of cell–cell and cell–ECM adhesions

regulates force balance while maintaining E-cadherin molecular tension in cell

pairs. Mol. Biol. Cell 26, 2456–2465. https://doi.org/10.1091/mbc.E14-12-1618

Figures

Figure 1: Overview of strategy for inferring the effective potential of cell–cell interactions

A. Microscopic forces contributing to cell–cell interactions are exemplified.

B. The microscopic forces are approximated as attractive/repulsive forces where cells are

described as particles with the mean size.

C. Both the repulsive and attractive forces are provided as cell–cell distance-dependent

functions, and the summation (black broken line) is the distance–force curve of the two

particles. The relationship between the curve and the mean diameters of the cells is shown.

D. A cavity-harboring tissue is illustrated with liquid flux into the cavity.

Figure 2: Influence of external constraints to effective forces of cell–cell interaction

Simulations were performed on the basis of a distance–force (DF) curve obtained from

the Lenard-Jones potential (LJ).

A. Expanding cavities are assumed in the particle model. i) An example of simulations

with the change in the radius of the cavity. The particles are of the mean size of cells. ii)

Inferred effective DF curves with the distant energy barriers (DEBs). The pregiven DF

curve (LJ) is shown in a broken black line. The numbers of the particles are the same

among the different cavities.

B. Expanding cavities are assumed whereas cell–cell interaction forces are not introduced.

i) An example of simulations. ii) Inferred effective DF curves.

C. Reconstruction of cell–cell interaction forces under expanding cavities. i) All the data

points in A and B under the radius of the cavity = 9.7 -> 14.2μm are plotted on distance–

force graph as a heat map, and the binned averages were calculated (yellow) which

correspond to the DF curves in A and B. ii) DF curves obtained by subtracting the DF

curves in B from those in A: i.e., A – B as exemplified in C-i. The definition of the heat

map is described in Supplementary information

D. The subtraction performed in C is schematically explained in the relationship among

A, B, and C. See the main text.

Related figure: Figure S1 (simulation procedures and inferred DF curves under non-

expanding cavities).

Related movie: Movie S1 (simulation)

Figure 3: Inference of the effective force of cell–cell interaction in mouse blastocysts

A. Blastocyst stages of mouse embryo are illustrated, and their confocal microscopic

images are shown: bright field, maximum intensity projection (MIP) and cross-section of

fluorescence of H2B-EGFP. Snapshots of nuclear tracking are also shown; blue spheres

indicate the detected nuclei. ICM cells are located around the bottom region of the image,

as indicated by *. Scale bars = 15μm.

B. Inferred effective DF curve in the mouse blastocyst (ref line). DEB is indicated. For

comparison, an inferred effective DF curve from the compaction stage embryo before

forming cavities is shown which we previously reported (green line) (ref).

C. Inferred effective DP curves in the mouse blastocysts. Four embryos were examined.

An enlarged view around longer distance is also shown in the inset.

D. Inferred effective DP curves in the mouse embryos where the cavitation was inhibited.

S3226 whose target is Na+/H+ exchanger inhibits the cavitation of the blastocyst

(Kawagishi et al., 2004). In the left panel, the microscopic images are shown. In the right

panel, the DF curves from three embryos are shown.

Related figure: Figure S2 (experimental procedures of S3226 treatment)

Figure 4: Inference of effective force of cell–cell interaction in cysts formed by MDCK

cells

A. MDCK cysts formed under suspension conditions are illustrated, and confocal

microscopic images are shown: bright field, maximum intensity projection (MIP) and

cross-section of fluorescence of H2B-EGFP. Scale bars = 10μm.

B and C. Inferred effective DF and DP curves. In C, an enlarged view is also provided. A

DEB is indicated.

Related figures: Figure S3 (DP curves from other cysts).

Figure 5: Outcomes of simulations based on inferred distance–force curves with DEBs

A. Initial configurations of particles are exemplified, that harbor a cavity. A cross-section

is also shown in the right panel.

B-E. Simulations results under the effective DF curves derived from the mouse blastocyst

(B), and the MDCK cyst (C), and the synthetic tissue harboring an expanding cavity (D)

in Fig. 2 are shown. E is the simulation result under the LJ potential for comparison. In

addition to the left panels ("Full DF curves"), the DF curves in which the DEBs were

artificially eliminated were used in the right panel ("DF curves w/o DEB); in D and E,

analyzes for the right panel were not performed ("ND"). For each simulation result, cross-

sections are also shown, and in the case of the mouse blastocyst, two parallel cross-

sections (shown in blue and green) are merged.

Figure 6: Modeling of distance–force curves and diagrams of simulation outcomes;

cavity-harboring structures

A. Modeling of DF curves using four parameters: N, D_0, δ, and ε. In the left panel, a DF

curve without the cosine term is always >0 (repulsive) ($F = \varepsilon D^{-N}$, green broken line).

Upon introduction of the cosine (purple broken line), the curves acquire regions with <0

(attractive) and a DEB (red line). Upon introduction of D_0, the curves are translated along

the distance (blue line). As shown in the blue line, the DEB ranges from $3\delta/4 + D_0$ to

$5\delta/4 + D_0$ (D_2). In the right panel, N affects the gradient of the curves and the height of

DEBs, and thus, the relative strength of DEB is modified.

B. Outcomes of simulations under various values of N and D^* in the presence of DEB.

From the equation defined in A, DF curves from $D = 0$ to D at the end of DEBs (w/ DEB)

were used as exemplified in the graph of DF curve ("w/ DEB"). The initial configuration

is shown in the left bottom. On the diagram, some conditions were designated by symbols

(circle, square, and triangle), and their simulation results are visualized on the bottom row

(blue spheres) with the profiles of the DF curves. Cross-sections are visualized for some

simulation results. The diameters of the blue spheres are equivalent to the mean diameter.

For "cup" [orange rings, mouth of the cup; white dots, particles located on the ring;

crosses, the particles located neither on the ring nor at the center of the ring]. For "tube"

[orange crosses, open ends of the tube]. For "cavity with extra particles" [arrows, extra

particles where multiple- but not single-layered]. D_1, mean diameter; D_2, distance of

DEBs. The definition of each phase is explained in Supplementary information. Gray

crosses, data points of all simulations for depicting the lines separating each phase.

C. Simulation outcomes in the absence of DEBs. DF curves from $D = 0$ to D before

starting DEB (w/o DEB) were used as exemplified in the graph of DF curve ("w/o DEB").

Related figure: Figure S4 (phase diagrams under a different number of particles), S5

(maintenance of tubes was examined under DF curves with DEBs).

Related movie: Movie S2 (the cavity-harboring structure was stably maintained under

fluctuations)

Figure 7: Diagrams of simulation outcomes; 2D sheets, spherical/non-spherical

aggregates

A. Outcomes of simulations starting from a two-dimensional (2D) sheet with slight

positional fluctuations. The results are shown in a similar manner to Fig. 6B, except that

the diameters of the blue spheres are equivalent to 0.4× the mean diameter. D_1, mean

diameter; W_1, width of attractive force regions; P_1, potential energy derived from

attractive forces.

B. Simulation outcomes without DEBs.

Related figure: Figure S6 (possible morphological transitions)

Related movie: Movie S2 (2D sheets were stably maintained under fluctuations), and

Movie S3 (possible morphological transitions).

Figure 8: Radial distribution functions (RDF) in 2D and 3D situations

A. Comparison of inferred DF curves in 2D (i) and 3D (ii) situations. The DF curves are

shown with snapshots of simulations and with the pregiven potentials (LJ). DEB is

indicated. The condition of the simulations is [sampling interval = 3 min, SD value of

force fluctuation = 1000%, and persistency of force fluctuation = 1 min].

B. The relationship between RDF and effective pairwise potentials is described. An RDF

for the central particle (dark blue) is illustrated. Densities of particles are shown as the

function of distances from the central particle. Higher density at a distance means an

energetically favorable state, and therefore, the potential is expected to be lower at the

distance.

C. RDFs in 2D and 3D situations were calculated in simulations under the LJ potential.

The snapshots of the simulations and the RDFs are shown. In these simulations,

fluctuations of the forces given by the LJ potential were introduced with the persistency.

Under longer persistency, the particles tend to easily change their positions. The condition

is [SD value of force fluctuation = 300%, and persistency of force fluctuation = 1 min].

Fig.1 Koyama et al.

Fig.2 Koyama et al.

A Blastocyst

Illustration Bright field Histone-EGFP(nuclei) Tracking
TE   MIP   Cross-section
cavity
ICM

B effective distance-force (DF) curve

Effective force of cell-cell interaction [A.U.]
repulsive / attractive

DEB

Blastocyst
Compaction

cell-cell distance [μm]

C effective distance-potential (DP) curve

Effective potential of cell-cell interaction [A.U.]

DEB

cell-cell distance [μm]

D cavitation-inhibited by S3226

Bright field Cross-section

Effective potential of cell-cell interaction

cell-cell distance [μm]

Fig. 3 Koyama et al.

A

Illustration Bright field Histone-EGFP(nuclei)
                            Cross-section

B effective distance-force (DF) curve

C effective distance-potential (DP) curve

Fig.4 Koyama et al.

Fig.5 Koyama et al.

Fig.6 Koyama et al.

Fig.7 Koyama et al.

A Distance-force curves in 2D sheet vs. 3D aggregate

i. 2D sheet

Effective force of cell-cell interaction [A.U.]
repulsive / attractive
- - - LJ (given)
—— inferred
cell-cell distance [μm]

ii. 3D aggregate

Effective force of cell-cell interaction [A.U.]
repulsive / attractive
- - - LJ (given)
—— inferred
cell-cell distance [μm]

B Relationship between pairwise potential and radial distribution function

Radial distribution function

Density ($\rho(R)$)

high density
→ energetically favorable
→ low potential energy

distance (R)

$$\rho(R) = \rho_0 \left\{ \exp(-U(R)/k_B T) \right\}$$

Effecitve potential of cell-cell interaction

Effective potential ($U(R)$)

C Radial distribution functions in 2D sheet vs. 3D aggregate

i. 2D sheet

Density [number/μm²]
cell-cell distance [μm]

ii. 3D aggregate

Density [number/μm³]
cell-cell distance [μm]

persistency of fluctuation
—— 0 min
—— 3
—— 10
—— 30
- - - 50

Fig.8 Koyama et al.