Novel Gurmarin-like Peptides from *Gymnema sylvestre* and their Interactions with

the Sweet Taste Receptor T1R2/T1R3

Halim Maaroufi

Institut de biologie intégrative et des systèmes (IBIS), Université Laval, Quebec City,
Quebec, Canada. halim.maaroufi@ibis.ulaval.ca

Abstract (228 words)

Gymnema sylvestre (GS) is a traditional medicinal plant known for its hypoglycemic and

hypolipidemic effects. Gurmarin (hereafter Gur-1) is the only known active peptide in

GS. Gur-1 has a suppressive sweet taste effect in rodents but no or only a very weak

effect in humans. Here, eight gurmarin-like peptides (Gur-2 to Gur-9) and their isoforms

are reported in the GS transcriptome. The molecular mechanism of sweet taste

suppression by Gur-1 is still largely unknown. Therefore, the complete architecture of

human and mouse sweet taste receptor T1R2/T1R3 and its interaction with Gur-1 to Gur-

9 were predicted by AlphaFold-Multimer (AF-M) and validated. Only Gur-1 and Gur-2

interact with the T1R2/T1R3 receptor. Indeed, Gur-1 and Gur-2 bind to the region of the

cysteine-rich domain (CRD) and the transmembrane domain (TMD) of the mouse T1R2

subunit. In contrast, only Gur-2 binds to the TMD of the human T1R2 subunit. This result

suggests that Gur-2 may have a suppressive sweet taste effect in humans. Furthermore,

AF-M predicted that Gα-gustducin, a protein involved in sweet taste transduction,

interacts with the intracellular domain of the T1R2 subunit. These results highlight an

unexpected diversity of gurmarin-like peptides in GS and provide the complete predicted

architecture of the human and mouse sweet taste receptor with the putative binding sites

of Gur-1, Gur-2 and Gα-gustducin. Moreover, gurmarin-like peptides may serve as

promising drug scaffolds for the development of antidiabetic molecules.

Introduction

Gurmarin (in reference to "Gurmar" that is mean "sugar destroyer" in Hindi) is the only disulfide-rich peptide of 35 amino acids purified from the leaves of the Indian plant *Gymnema sylvestre* (Imoto et al., 1991; Kamei et al., 1992). Gurmarin (hereafter Gur-1) contains three disulfide bridges forming a cystine knot fold (knottin fold) that confers high resistance to proteolytic, thermal, and chemical degradation (Pallaghy et al., 1994). Gur-1 suppresses the sweet taste responses to sucrose, glucose, glycine, and saccharin in rats (Imoto et al., 1991; Yoshie et al., 1994; Miyasaka and Imoto, 1995) and mice (Ninomiya and Imoto, 1995). However, Gur-1 showed no or only a very weak suppressive sweet taste effect in humans (Imoto et al., 1991). To suppress the sweet taste response, Gur-1 is thought to interact with T1R2/T1R3 receptor in rodents (Shigemura et al., 2008; Ohkuri et al., 2009; Yasumatsu et al., 2009). The T1R2/T1R3 heterodimer, which belongs to class C G-protein-coupled receptor (GPCR), also known as seven-transmembrane (7TM) receptor, is a membrane sweet taste receptor (STR) in eukaryotes (Nelson et al., 2001). STR is expressed in taste buds. T1R2 and T1R3 monomer are also expressed in the gastrointestinal tract (Bezençon et al., 2007), pancreas (Nakagawa et al., 2009; Kyriazis et al., 2012), brain (Ren et al., 2009; Kohno et al., 2016), and testis (Mosinger et al., 2013). T1R2 and T1R3 monomer consist of an N-terminal extracellular ligand-binding (LB) domain (also called Venus flytrap module (VFTM)), a cysteine-rich domain (CRD), the seven-helical transmembrane domain (TMD), and a C-terminal intracellular domain (ICD).

A systematic review and meta-analysis have shown that *G. sylvestre* supplementation has a beneficial effect on glycemic and lipid control in patients with type-2 diabetes mellitus (Devangan et al., 2021). Recently, it has been found that Gur-1 reduces significantly sucrose intake in rodents (Rayo-Morales et al., 2023). Unfortunately, Gur-1 has no or a very weak suppressive sweet taste effect on humans. Therefore, the search for gurmarin-like peptide(s) that may be effective in humans will be helpful in the development of antidiabetic molecules. In this study, the search for the gurmarin-like peptides in the *G. sylvestre* transcriptome reveals a rich arsenal of gurmarin-like peptides. Furthermore, the predicted architecture of the human and mouse sweet taste receptor T1R2/T1R3 and their interaction with Gur-1 and Gur-2 suggest that the sweet taste suppression in rodents by

Gur-1 is probably due to its interaction with the region of the CRD near the TMD of the

mouse T1R2 subunit. Gurmarin-like peptides could be interesting drug scaffolds which

may help to reduce the craving for sugar and thus help to regulate short-term blood

glucose levels.

Materials and methods

Search for Gur-1-like peptides in *G. sylvestre*

TblastN algorithm (http://www.ncbi.nlm.nih.gov/blast) with Gur-1 (UniProt ID:

A0A3S9JKY1 (full sequence) and P25810 (mature sequence)) as query is used to search

in *de novo* transcriptome (RNA-seq) sequences of: (i) leaf, flower and developing fruits

of *G. sylvestre* genotype DGS-22 deposited in NCBI Short Read Archive (SRA) (SRA

accession ID: SRR5965323, SRR5965320, SRR5965321) (Kalariya et al., 2019), (ii)

leaves prior to flowering of *G. sylvestre* genotype DGS-3 (SRA accession ID:

SRR5965322) (Kalariya et al., 2018), and (iii) plant samples (SRA accession ID:

SRR7876667) and leaf tissues (SRA accession ID: SRR3664033) of *G. sylvestre*

(Ayachit et al., 2019).

Chemical properties of gurmarin-like peptides

The chemical structures and properties of gurmarin-like peptides were investigated *in*

silico using PepDraw (http://www.tulane.edu/~biochem/WW/PepDraw/index.html) as

described in Maaroufi and collaborators (Maaroufi et al., 2021).

Phylogeny

Multiple sequences alignment are conducted by Mafft (Katoh and Toh, 2008) and

phylogenetic tree is constructed by PhyML (Guindon et al., 2010).

Residue numbering

The amino acid residues in Gur-1 to Gur-9, mouse and human T1R2 and T1R3 are

numbered without the signal peptide.

Structure prediction with AlphaFold2 and AlphaFold-Multimer

The AlphaFold2 (AF2 v2.2.2) and AlphaFold-Multimer (AF-M v2.2.2) source code was installed locally on a Linux workstation with NVIDIA Quadro RTX6000 GPU.

The structure of Gur-1 has already been determined (PDB ID: 5OLL), while Gur-2 to Gur-9 peptides were predicted using AF2. While AF-M was used to predict the likely complex formation with different combinations of Gur-1 to Gur-9 peptides, T1R2 and T1R3 from mouse (UniProt ID: Q925I4 and Q925D8) and human (UniProt ID: Q8TE23 and Q7RTX0) (Jumper et al., 2021; Evans et al., 2022). AF-M is also used to study the interactions of Gur-1, Gur-2 with mouse T1R2, T1R3 and the Gα-gustducin (GNAT3) (UniProt ID: Q3V3I2).

To test the accuracy of AF-M, the brazzein peptide (UniProt ID: P56552) that has been studied experimentally for its interactions with human T1R2 and T1R3 has been used as input in AF-M with human T1R2 and T1R3.

AF2 and AF-M generate five models that are ranked based on the model confidence pLDDT score (The mean predicted Local Distance Difference Test) for AF2 and the combination of pTM+ipTM scores (pTM for monomers and ipTM for the monomers interface). Thus, pTM reports on the accuracy of prediction within each protein subunit and ipTM on the accuracy of the complex. If AF-M is unsure about an interface between the monomers, those are not docked. Each of the five predicted structure was examined. The structure figures were generated with PyMOL (http://www.pymol.org).

Molecular dynamics simulations

In order to evaluate conformational changes of Gur-1 (PDB ID: 5OLL) and Gur-2 (3D predicted by AF2), molecular dynamics (MD) simulations were performed in GROMACS (2021.4) using the OPLS-AA/L all-atom force field (Kaminski et al., 2001) as described in (Maaroufi et al., 2021), except that Gur-1 and Gur-2 MD simulations were performed for 100 ns and in triplicate.

Structural similarity search

DALI (http://ekhidna2.biocenter.helsinki.fi/dali/), Foldseek (https://search.foldseek.com/search) and the PDB website software

(https://www.rcsb.org/search/advanced/structure) were used to search for structural

similarities of Gur-1 to Gur-9, T1R2 and T1R3 in the PDB and AlphaFold databases

(Van Kempen et al., 2023).

Membrane topology prediction

Membrane T1R2 and T1R3 topology prediction was conducted by MembraneFold

(https://ku.biolib.com/MembraneFold/) (Gutierrez et al., 2022).

Detection of T1R2/T1R3 tunnels/channels

CAVER software (https://loschmidt.chemi.muni.cz/caverweb/) was used for the detection

and analysis of tunnels and channels in T1R2/T1R3 structures.

Binding affinity

The PRODIGY web server (https://wenmr.science.uu.nl/prodigy/) was used to calculate

the binding free energies (ΔG) of interacting protein-protein and peptide-protein

complexes (Xue et al., 2016).

Amino acids interactions in protein-protein complex

PyMOL and ContPro (http://procarb.org/contpro/index.html are used to determine the

bonds between protein-protein and peptide-protein complexes.

Results and discussion

G. sylvestre **transcriptome is rich in gurmarin-like peptides**

Gurmarin (hereafter Gur-1) is the only active peptide extracted from the leaves of

Gymnema sylvestre (Imoto et al., 1991; Kamei et al., 1992). In the aim to find homologs

of Gur-1 (UniProt ID: P25810), a TblastN was performed using the transcriptome (RNA-

seq) sequences of leaves, leaves prior to flowering, flowers and developing fruits of *G.

sylvestre* deposited in the NCBI Short Read Archive (SRA) (Kalariya et al., 2018, 2019;

Ayachit et al., 2019). From this search, 57 gurmarin-like peptides were found. Multiple

sequence alignment (MSA) of these peptides revealed nine groups, named Gur-I to Gur-

IX (Fig. S1). Gur-I group is rich in isoforms of Gur-1. Interestingly, Trp29 whose

mutation to alanine completely abolishes the sweet-suppressing properties of Gur-1, is

conserved in isoforms of Gur-I and Gur-II groups (Sigoillot et al., 2018). It has been

found that Gur-1 has no or only a very weak suppressive sweet taste effect in humans

(Imoto et al., 1991; Miyasaka and Imoto, 1995). Therefore, this arsenal of isoforms might

have a more potent sweet taste inhibitory effect in humans. It is noteworthy that some of

these isoforms have mutations in Cys10 or Cys17 and Cys10/Cys17. Gur-1 has three

disulfide bridges (Cys3-Cys18, Cys10-Cys17 and Cys23-Cys33). It has been shown that

mutagenesis of these six cysteines alters the function of Gur-1. Indeed, mutant Gur-1

analogs with only two native disulfide bonds suppressed the response to sucrose, but not

those to glucose, fructose, saccharin, or glycine in rats (Ota et al., 1998b).

For the continuation of this work, one representative sequence from each group of

gurmarin-like peptides was selected and named Gur-1 (from group I) to Gur-9 (from

group IX) (Fig. 1). The analysis of Gur-1 to Gur-9 showed that they have different

physicochemical properties (Table S1). Gur-2 is more similar to Gur-1 (48.57% of

identity and 65.71% of similarity in amino acid residues). Like Gur-1, Gur-2 has a

hydrophobic and hydrophilic cluster on the surface of the molecule (Ota et al., 1998a).

Gur-2 is more hydrophobic than Gur-1 (Table S1). For Gur-3 to Gur-9, BlastP (expect

threshold of 0.05) against the NCBI database showed no matches, indicating that they are

novel knottin (cystine knot fold) peptides.

```
               1         10           20          30
Gur-1_P25810         QQCVKKDELCIPYYL.DCCEPLECKKVNW.WDHKCIG.......
Gur-2_SRR3664033.37000769.1   ELCQEKDEPCVPNFM.ECCEPYKCRSVSW.WELKCIS.......
Gur-3_SRR7876667.11909172.1   AKCLKKGQKCNPPYT.PCCRPYRCVQTS..RPYKCGI.......
Gur-4_SRR5965322.12953624.2   SDCLPEGALCG..LGIECCPPFHCSEPM..RPFRCRR.......
Gur-5_SRR5965322.2264852.1    PYCLPEGAFCGHQLDIPCCPPFHCSEPM..RPFMCER.......
Gur-6_SRR5965322.10344500.1   LNCRQLLELCGPGVS.DCCPPLECLRTI...ENRCQGYGD....
Gur-7_SRR5965322.12830937.2   LICLPEGALCGGSLG.ECCPPLLCIRTL...DIRCRKLGYV...
Gur-8_SRR5965322.13661248.2   LNCLSIGETCGPGVG.YCCFPLICSVGTEHVGFKCVYVATKSKA
Gur-9_SRR5965322.3024837.1    LSCVEEGNDCEPIEN.ECCPSLECIGSY..YGFKCEHPV.....
consensus>70         ..C......C.......CC.p..C..........C.........
```

1.

Fig. 1. Multiple sequence alignment and clustering of Gur-1 (UniProt ID: P25810) to Gur-9 peptides. Each peptide has six conserved cysteines that are predicted to form a knottin three disulfide bonds. SRA accession ID is indicated with each sequence.

Prediction of the structure of gurmarin-like peptides

The structure of Gur-1 is known (PDB ID: 5OLL). AlphaFold 2 (AF2) is used to predict

the structures of Gur-2 to Gur-9. AF2 produces five ranked models. Each of the five

predicted structures was examined. AF2 predicted structures of all gurmarin-like peptides

in the except of Gur-6. Of note, in the predicted structure of Gur-5 (pLDDT score of

56.77 is bad) the three disulfide bridges are different from those in Gur-1. Therefore,

ESMFold software (based on large language models) that uses only the amino acid

sequence to predict the structure is used for Gur-5 and Gur-6 (Lin et al., 2023). The

results of the structure alignments of Gur-1 to Gur-9 using DALI (Holm, 2020) are

shown in Table S2. The structural similarity dendrogram of Gur-1 to Gur-9 shows that

the structures of Gur-2 to Gur-5 are close to Gur-1, whereas Gur-6/Gur-8 and Gur-7/Gur-

9 are distant from Gur-1 (Fig. S2).

Prediction of the structure of the T1R2/T1R3 heterodimer

Subunits T1R2 and T1R3 of sweet taste receptor consist of a signal peptide, an N-terminal extracellular domain, the seven-helical transmembrane domain (TMD), and a C-terminal intracellular domain (ICD). There is no experimental structure of the monomers or the complete sweet taste receptor T1R2/T1R3. Only the extracellular ligand-binding domains (LBDs, also named Venus flytrap module (VFTM)) of the medaka fish T1R2/T1R3 heterodimer have been determined (PDB ID: 5X2M). Therefore, AF2 was used to predict the structure of the human T1R2 and T1R3 (hT1R2 and hT1R3) and the mouse T1R2 and T1R3 (mT1R2 and mT1R3) monomers (Fig. 2), and AlphaFold-Multimer (AF-M) to predict the structure of the hT1R2/T1R3 and mT1R2/T1R3 heterodimer (Fig. 4). AF-M generates five ranked models. Each of the five predicted structures was examined. The pLDDT (the mean predicted Local Distance Difference Test) score of mT1R2, mT1R3, hT1R2 and hT1R3 monomers are 84.41, 86.97, 85.81 and 87.1, respectively, indicating a high 3D models quality. While the best 3D model of mT1R2/T1R3 and hT1R2/T1R3 has a pTM+ipTM (overall predicted TM (pTM) + predicted interface (ipTM)) score of 0.68. When the ipTM score is greater than 0.6, it indicates that the protein interface is well predicted (Evans et al., 2022). The twilight zone for a good confidence of the predicted interface is found between 0.5 and 0.6. Recently, at CASP15, the accuracy of AF-M has been demonstrated in modeling the quaternary structure of proteins in comparison to bioinformatics methods used prior to AF software (Ozden et al., 2023).

Fig. 2. The membrane topology of the predicted structure of the mouse T1R2 (mT1R2)
by AF2. mT1R2 is formed with an N-terminal Venus flytrap module (VFTM), a cysteine-
rich domain (CRD), a seven-helical transmembrane domain (TMD) and an intracellular
domain (ICD). The accuracy of the predicted model is visualized by different colors
(pLDDT ≥ 90, 90>pLDDT≥70, 70>pLDDT≥50, 50>pLDDT). Amino acid residues of
mT1R2 are numbered without the signal peptide.

Analysis of the hT1R2/T1R3 complex revealed that amino acid residues of hT1R2

involved in interactions with hT1R3 according to Park et collaborators (Park et al., 2019)

are effectively located at the interface between hT1R2 and hT1R3 in the model predicted

by AF-M (Fig. 3). In addition, an ionic bond between Arg217 of hT1R2 and Glu217 of

hT1R3 in the predicted model is consistent with the study of Cui and coworkers who, by

mutating hT1R2 Arg217 to alanine suggested that Arg217 is involved in the

hT1R2/T1R3 interaction (Cui et al., 2006). The predicted binding affinity of mT1R2 and

mT1R3 in the heterodimer (ΔG ~ -23.1 kcal/mol) is similar to that of hT1R2 and hT1R3

(ΔG ~ -21.0 kcal/mol).

The structure similarity search (https://www.rcsb.org/search/advanced/structure) in PDB

database revealed that the predicted structure of the hT1R2 and mT1R3 monomers are

structurally similar to the human calcium-sensing receptor (CaSR) (PDB ID: 7M3G_A)

and human metabotropic glutamate receptor 2 (mGlu2) (PDB ID: 8JD3_A). Like T1R2

and T1R3, CaSR and mGlu2 are a class C G protein-coupled receptors (GPCRs).

Pairwise structure alignment (https://www.rcsb.org/alignment) using the jFATCAT

(flexible) option, aligned T1R2 and T1R3 monomers to CaSR and mGlu2 with an RMSD

of 3.1-3.21 Å at 723-765 amino acid residues, and 2.71-3.76 Å at 700-740 amino acid

residues, respectively. Whereas T1R2 and T1R3 are aligned to the LBDs of the medaka

fish T1R2/T1R3 heterodimer (PDB ID: 5X2M) with an RMSD of 1.89-2.18 Å on 427-

440 amino acid residues.

Fig. 3. Predicted structures of human T1R2/T1R3 (hT1R2/T1R3) and mouse T1R2/T1R3
(mT1R2/T1R3) heterodimer. **A** and **B**, hT1R2/T1R3 (green/cyan) with amino acids
(yellow sticks) of hT1R2 implicated in the interactions with hT1R3 according to Park et
coworkers (Park et al., 2019). **C**, superposition of hT1R2/T1R3 (green/cyan) and
mT1R2/T1R3 (blue/orange). The structure of mT1R2/T1R3 aligns on hT1R2/T1R3 with
an RMSD of 1.84 Å.

Prediction of interactions between gurmarin-like peptides and T1R2 and T1R3

Brazzein and hT1R2/T1R3 interactions as control of AF-M accuracy

It has been demonstrated that AF-M is capable of predicting peptide-protein complex with high accuracy (Chang and Perez, 2023; Teufel et al., 2023). This accuracy sometimes depends on type of proteins. Therefore, the published experimental results of the binding of Brazzein (Brz), a sweet-tasting peptide of 54 amino acids stabilized by four disulfide bonds, with hT1R2 and hT1R3 were used to test the accuracy of the AF-M software for predicting the peptide-protein interactions.

AF-M showed that Brz interacts with the region Ile516-Glu525 in the CRD of hT1R3 subunit of T1R2/T1R3 heterodimer (iptm+ptm = 0.6168) (Fig. S3). This region is involved in the responsiveness to Brz. Indeed, the functional mutagenesis analysis revealed that Ala517 and Phe520, located in the region Ile516-Glu525, jointly determine the different sweet taste sensitivity between human (taster) and mouse (non-taster) toward Brz (Jiang et al., 2004a). In the predicted hT1R2/T1R3 structure, Ala517 and Phe520 interact with two adjacent aromatic residues Tyr39 and Phe38 of Brz, respectively. Interestingly, also two adjacent aromatic residues Trp28 and Trp29 in Gur-1 are important for the sweet-suppressing properties of Gur-1 (Sigoillot et al., 2018). In addition, as for Trp29 in Gur-1 (its mutation to alanine abolishes the suppressive effect of Gur-1), the second aromatic residue Tyr39 is more critical for the Brz activity (Jiang et al., 2004; Sigoillot et al., 2018). Moreover, Ala19 of Brz binds Arg490 of hT1R3. The site-directed mutagenesis of Ala19 has demonstrated its importance for the sweetness effect of Brz (Ghanavatian et al., 2016). Therefore, these experimental results support the accuracy of AF-M for predicting peptide-protein interactions of type peptide-sweet taste receptor.

Interactions of gurmarin-like peptides and T1R2 and T1R3

In situ studies suggest that Gur-1 sensitivity is related to the coupling of T1R2/T1R3 to Gα-gustducin (Shigemura et al., 2008; Ohkuri et al., 2009; Yasumatsu et al., 2009). In addition, the recombinant Gur-1 has been shown to be responsible for the complete blockade of saccharin responses in rat T1R2/T1R3-expressing cells (Sigoillot et al.,

2012a). Therefore, in the aim to understanding the molecular mechanism of sweet taste

suppression by Gur-1 and gurmarin-like peptides, AF-M is used to predict the

interactions of Gur-1 to Gur-9 with hT1R2/T1R3 and mT1R2/T1R3. AF-M is used with

these combinations of human and mouse amino acid sequences (Gur, mT1R2), (Gur,

mT1R3), (Gur, mT1R2, mT1R3), (Gur, hT1R2), (Gur, hT1R3), and (Gur, hT1R2,

hT1R3) (Table 2). The interactions of Gur-1 and gurmarin-like peptides with T1R2 and

T1R3 monomers are studied in the case where, in the taste cell, Gur-1 and Gur-2 bind

T1R2 and/or T1R3 before the dimerization of the sweet taste receptor.

The results showed that only Gur-1 and Gur-2 interact with the T1R2/T1R3 receptor. No

interactions of Gur-3 to Gur-9 are obtained with monomers nor T1R2/T1R3 heterodimer.

This suggests that Gur-3 to Gur-9 may have function other than sweet taste suppression.

AF-M produces five ranked models. In the case of T1R2/T1R3-Gur-1 or T1R2/T1R3-

Gur-2 complexes, two models with different binding sites for Gur-1 or Gur-2 were

obtained with a high pTM+ipTM score. Indeed, AF-M predicted that Gur-1 and Gur-2

interact with the extracellular and intracellular (less probable because Gur-1 and Gur-2

must cross the taste cell membrane) domains of T1R2/T1R3. Previous studies have

hypothesized that Gur-1 may interact with the extracellular region (VFTM and/or CRD)

of mT1R2/T1R3 but not with the intracellular region (Sigoillot et al., 2012b, 2018).

In the extracellular region, Gur-1 and Gur-2 bind to the CRD near the TMD of mT1R2

subunit of mT1R2/T1R3 with binding affinity of -7.1 and -7.7 Kcal/mol, respectively

(Fig. 4 and Fig. S5). Whereas in hT1R2/T1R3 only Trp29 of Gur-2 forms a cation-pi

bond with Arg706 of TMD of hT1R2 subunit with binding affinity of -5.1 Kcal/mol

(Table 1, Tables S3 to S6). Interestingly, Gur-1 do not interact with hT1R2/T1R3. This is

consistent with works that showed that Gur-1 has no or only weak suppressive effect in

human (Imoto et al., 1991; Ninomiya and Imoto, 1995).

Gur-1 binds through its hydrophobic cluster (Tyr13-Tyr14 and Trp28-Trp29) to the

region of CRD near TMD (Fig. 4D). It has been reported that tryptophan residues and

their position in peptides play an important role in the interaction with lipid membranes

(Rydberg et al., 2014). In addition, Obi and Natesan revealed the role of membrane lipids

in the binding affinity, the stability of ligands at the binding site in the G-protein-coupled

receptor (Obi and Natesan, 2022). They suggest considering the ligand-lipids interactions

for calculating the binding affinity. Considering lipids, Gur-1 may have more affinity for

mT1R2/T1R3 than its predicted binding affinity (Table 1). Therefore, the probable

interaction of Gur-1 with lipids may explain why the effect of Gur-1 is not immediate,

Table 1. Predicted interactions by AF-M of Gur-1, Gur-2, T1R2 and T1R3

Input for AF-M	Binding site of Gur-1 and Gur-2	ΔG (kcal mol-1)	pTM+ipTM
mT1R2, Gur-1	ICD	-8.1	0.660
mT1R3, Gur-1	CRD near TMD	-7.3	0.468
mT1R2, mT1R3, Gur-1	ICD mT1R2, CRD near TMD mT1R2	-8.4, -7.1	0.603
mT1R2, Gur-2	ICD	-9.1	0.593
mT1R3, Gur-2	CRD near TMD	-9.7	0.450
mT1R2, mT1R3, Gur-2	ICD mT1R2, CRD near TMD mT1R2	-8.3, -7.7	0.657
hT1R2, Gur-1	ICD	-8.8	0.665
hT1R3, Gur-1	CRD near TMD	-6.9	0.459
hT1R2, hT1R3, Gur-1	ICD hT1R2	-10.2	0.664
hT1R2, Gur-2	ICD	-8.9	0.584
hT1R3, Gur-2	CRD near TMD	-6.4	0.468
hT1R2, hT1R3, Gur-2	ICD hT1R2, CRD near TMD hT1R2	-8.1, -5.1	0.642

AF-M: AlphaFold-Multimer
Mouse T1R2 (mT1R2); human T1R2 (hT1R2)
VFTM: Venus flytrap module; CRD: cysteine-rich domain; TMD: trans-membrane domain; ICD:
intracellular domain

requiring 5 min to achieve maximal suppression of sweet responses (Ninomiya et al.,

1999). It may also explain the long-lasting suppressive effect of Gur-1 (>2-3 h) (Imoto et

al., 1991) and the ineffectiveness of the rinsing procedures (Sigoillot et al., 2018), in

addition to the fact that the suppressed sucrose response after Gur-1 recovered to some

extent after rinsing the tongue with 15 mM β-CD (β-cyclodextrin) for 10 min (Ninomiya

et al., 1998). It has been suggested that β-CD interacts with tyrosine and tryptophan

residues of Gur-1 hydrophobic cluster that facilitates the dissociation of the

mT1R2/T1R3-Gur-1 complex (Imoto et al., 2001). Indeed, in this work Gur-1 and Gur-2

bind to T1R2/T1R3 through their hydrophobic cluster Tyr13-Tyr14/Trp28-Trp29 and

Phe14/Trp28-Trp29, respectively.

In the cytoplasmic region (less probable because Gur-1 must cross the taste cell

membrane), Gur-1 and Gur-2 bind to the intracellular domain (ICD) of hT1R2 and

mT1R2 monomers as well as the T1R2 subunit of the T1R2/T1R3 heterodimer (Fig. 4A

and C) with binding affinities between -8.1 and -10.2 kcal/mol (Table 1). The

electrostatic surface potential showed that the ICD forms a crevice that accommodates

Gur-1 and Gur-2 (Fig. S4). The interaction with the ICD of T1R2 requires that Gur-1 and

Gur-2 cross the taste cell membrane. Otherwise, the results showing that Gur-1 and Gur-2

interact with the ICD have no biological value. As hypotheses, Gur-1 and Gur-2 could

have a propriety (cell-penetrating peptides) to cross the membrane (Gori et al., 2023)

and/or Gur-1 can enter the cell via the predicted channel between T1R2 and T1R3

subunits (Fig. S6).

Finally, Gur-1 and Gur-2 bind to the CRD near TMD in the human and mouse T1R3

monomers but not in T1R2/T1R3 heterodimer (Table 1 and Fig. 4B).

The amino acids of Gur-1 and Gur-2 involved in interactions with T1R2 and T1R3 are

mainly Tyr13-Tyr14 and Trp28-Trp29 for Gur-1 and Phe14 and Trp28-Trp29 for Gur-2

(Tables S3-S6). Interestingly, hydrophobic amino acids Tyr14, Trp28 and Trp29 are

showed involved in the sweet-suppressing properties of Gur-1 (Ota et al., 1998a; Sigoillot

et al., 2018).

Fig. 4. Interactions of Gur-1 with mouse T1R2, T1R3 monomers and T1R2/T1R3
heterodimer. (**A**) Gur-1 (pink) binds in the intracellular domain (ICD) of the T1R2
monomer (green) with Y13, Y14, W28 and W29 amino acid residues. (**B**) Gur-1 binds in
the cysteine-rich domain (CRD) near the transmembrane domain (TMD) of the T1R3
monomer (cyan). (**C**) and (**D**) Gur-1 binds in the ICD and the CRD near TMD of the
T1R2 subunit of T1R2/T1R3 heterodimer (green/cyan), respectively. For details of bonds
between amino acid residues, see Table S3.

Interaction of Gur-2 with chimeric T1R2/T1R3 heterodimer

Gur-2 binds to the CRD-TMD region of mT1R2 subunit of mT1R2/hT1R3 with an

affinity of -7.7 Kcal/mol (Fig. 5A). Whereas only Trp29 of Gur-2 interacts with Arg706

in the TMD of hT1R2 subunit with an affinity of -5.1 Kcal/mol (Fig. 5B). In the aim to

empower the binding of Gur-2 to the hT1R2 subunit the region between amino acids

Thr686 and Gln702 of hT1R2 is replaced by the corresponding region of mT1R2 (Fig.

5E). This replacement improves the binding affinity (-6.8 Kcal/mol) of Gur-2 to chimeric

mouse hT1R2/hT1R3 comparatively to hT1R2/hT1R3 (-5.1 Kcal/mol) (Fig. 5C).

Fig. 5. Interactions of Gur-2 with the mouse, human and chimeric T1R2/T1R3 heterodimer. (**A**) Large interaction interface (yellow) of Gur-2 (salmon) and the mouse T1R2 subunit (mT1R2). (**B**) Only the residue W29 of Gur-2 (grey) interacts with R706 of the human T1R2 subunit (hT1R2). (**C**) Large interaction interface (yellow) of Gur-2 (blue) and the chimeric T1R2/hT1R3 (The region between Thr686 and Gln702 of hT1R2 is replaced by the corresponding region of mT1R2 (**E**)). (**D**) superposition of mT1R2/T1R3-Gur-2 (Gur-2 in salmon), hT1R2/T1R3-Gur-2 (grey) and chimeric T1R2/T1R3-Gur-2 (blue). T1R2 subunit (green) and T1R3 subunit (cyan). (**E**) Pairwise alignment of the region where Gur-2 interacts with mT1R2. Amino acids in mT1R2 that interact with Gur-2 are underlined with green stars and head arrows. The latter represents the two amino acids different in mT1R2 and hT1R2.

Interactions of Gur-1 and Gur-2 with homodimers of T1R2 and T1R3

The use of AF-M with two sequences of mT1R2 and one sequence of Gur-1 or Gur-2

showed only the formation of mT1R2/T1R2 homodimers, but not mT1R2-Gur-1 or

mT1R2-Gur-2 complexes. The same result was obtained with two sequences of mT1R3

and one sequence of Gur-1 or Gur-2. It has been suggested that the putative

homodimerization of T1R3 is not functional (Shimizu et al., 2014). Recently, Belloir and

coworkers found that the overexpressed hT1R2 is predominantly a dimer (Belloir et al.,

2021). AF-M results indicate that Gur-1 and Gur-2 only interact with the T1R2/T1R3

heterodimer. This is consistent with the requirement of heterodimerization of T1R2 and

T1R3 for sweet taste perception.

***In silico* competition assays between Gur-1 and Gur-2**

To compare the binding affinity of Gur-1 and Gur-2 to T1R2, T1R3 and T1R2/T1R3, *in*

silico competition assays were conducted (Chang and Perez, 2023; Banerjee et al., 2023).

To remember that interactions of Gur-1 and Gur-2 with the ICD are considered less

probable because Gur-1 and Gur-2 must cross the taste cell membrane.

For T1R2 as monomer or in T1R2/T1R3 heterodimer, Gur-1 binds or clashes with the

ICD while Gur-2 binds to the CRD near TMD region (Table S7). Whereas for T1R3

monomer, Gur-1 binds or clashes with the TMD near ICD while Gur-2 binds to the CRD

near TMD region. Therefore, comparatively to the results obtained with single gurmarin

peptide (Table 1), Gur-2 has more affinity than Gur-1 for the CRD near TMD region of

T1R2 or T1R3. That would be interesting to use Gur-1 and Gur-2 (and their isoforms)

together in the experiments to know if they may have a synergistic effect on the

suppression of sweet taste in humans.

Effect of Gur-1 or Gur-2 binding on the conformational changes

The pairwise structural alignment was performed to know whether the binding of Gur-1

or Gur-2 to the T1R2/T1R3 heterodimer changes the conformation of the complex. For

mice, structural alignment of the mT1R2/T1R3 with the mT1R2/T1R3-Gur-1 or

mT1R2/T1R3-Gur-2 complex showed an RMSD of 3.02 Å and 3.35 Å over 1624 amino

acid residues, respectively. Whereas hT1R2/T1R3 with hT1R2/T1R3-Gur-2 complex is

3.10 Å over 1600 residues. These results show that the binding of Gur-1 or Gur-2 to the

T1R2/T1R3 induce an important conformational change that may destabilize the active

conformation of T1R2/T1R3 and thus prevent the transmission of the sweet taste signal

from taste cell surface to intracellular signaling pathways (Perez-Aguilar et al., 2019).

Molecular dynamics simulation

Only Gur-1 and Gur-2 among the gurmarin-like peptides interact with the sweet taste

receptor. They share 51% of identity in amino acids and are structurally aligned with an

RMSD of 0.94 Å on all amino acid residues. Gur-1 and Gur-2 show some differences in

binding affinities for T1R2 and T1R3 monomers and the T1R2/T1R3 heterodimer, in

addition that only Gur-2 interacts with hT1R2/T1R3 (Table 1). In the aim to explain these

differences, molecular dynamics (MD) simulations with Gur-1 and Gur-2 for 100 ns were

conducted in GROMACS to compare their stability and conformational changes. As

shown in Fig. 6B, Gur-2 has less conformational flexibility than Gur-1. The most flexible

part of Gur-1 and Gur-2 is the N-terminal extremity. Interestingly, the two other more

flexible regions are those containing Tyr13-Tyr14 and Trp28-Trp29 amino acids of Gur-

1. This result is consistent with that of Sigoillot and coworkers who demonstrated the

flexibility of the Tyr13-Asp16 amino acid cluster and the Trp28-Trp29 residues. Tyr13-

Tyr14 and Trp28-Trp29 are important for the suppressive effect of Gur-1 (Sigoillot et al.,

2018), and in this study they are found the principally amino acids interacting with T1R2

and T1R3. Of note, the difference in the flexibility between Gur-1 and Gur-2 is more

pronounced in the region containing Trp28-Trp29. To characterize the dynamic changes

of the whole peptide, the time evolution of the radius of gyration (Rg) and solvent-

accessible surface area (SASA) of Gur-1 and Gur-2 are displayed (Fig. 6C and D). Rg

and SASA do not show a great difference between Gur-1 and Gur-2. As mentioned above

there is a difference in hydrophobicity, Gur-2 is more hydrophobic than Gur-1 (Table

S1). This difference in hydrophobicity, among others, may explain why Gur-2 interacts

with human T1R2/T1R3.

Fig. 6. Molecular dynamics simulations (100 ns) of Gur-1 (black curve) and Gur-2 (green curve). **A**, Time evolution of Root mean square deviation (RMSD) with respect to initial structure. **B**, root mean square fluctuations (RMSF) of peptides residues. The position of amino acid residues in alignment is identical to this in RMSF plot. **C** and **D** radius of gyration (Rg) and solvent-accessible surface area (SASA), respectively.

Interaction of T1R2/T1R3 and Gα-gustducin

On the tongue, the T1R2/T1R3 receptors are coupled with Gα-gustducin for the

transmission of the sweet taste signal from taste cell surface to intracellular signaling

pathways (Lindemann, 2001; Yasumatsu et al., 2009). Gα-gustducin has been shown to

be a key molecule involved in the transduction pathway for Gur-1 sensitive sweet

responses (Shigemura et al., 2008; Ohkuri et al., 2009; Yasumatsu et al., 2009).

AF-M predicted that mouse Gα-gustducin (mGα-gustducin) interacts only with the ICD

of the mT1R2 subunit in the mT1R2/T1R3 complex (Fig. 7). This result is in agreement

with the results of Sainz and coworkers who suggested that the T1R2 subunit in sweet

taste receptor is responsible for the activation of the coupled G protein (Sainz et al.,

2007), and of Liu and collaborators who showed that heterodimeric metabotropic

glutamate receptors (mGlus) transduce signals with one specific subunit that activates G

protein (Liu et al., 2017).

Fig. 7. Interaction of Gα-gustducin with the mouse T1R2/T1R3. Gα-gustducin (pink)
binds to the intracellular domain (ICD) of T1R2 (green) subunit of T1R2/T1R3
heterodimer (green/cyan). Amino acid residues of T1R2 and T1R3 are numbered without
the signal peptide.

In the binding interface, the C-terminus Cys351, Leu353 and Phe354 residues of mGα-

gustducin interact with Thr820, Ile816 and Met658 residues of the mT1R2 subunit of the

mT1R2/T1R3 complex, respectively. This result is consistent with the work of Hara and

collaborators who showed that the last C-terminus 37 amino acid residues of Gα-

gustducin are important for its activity in yeast (Hara et al., 2012). Additionally, a point

mutation Gly352Pro of Gα-gustducin impaired its ability to activate taste GPCRs (Ruiz-

Avila et al., 2001). It is probable that the Gly352Pro mutation destabilizes the region

containing residues Cys351, Leu353 and Phe354, preventing Gα-gustducin to interact

with the receptor.

Conclusion

This study sheds light on the discovery of novel gurmarin-like peptides in *G. sylvestre*, in

addition to the prediction and the validation of the complete architecture of sweet taste

receiver (STR) and the putative binding sites of Gur-1, Gur-2 and Gα-gustducin. It also

validates previous hypotheses that suggested that hydrophobic residues Tyr13-Tyr14 and

Trp28-Trp29 are involved in the STR-Gur-1 interaction. The putative binding sites will

allow to perform the site-directed mutagenesis of specific amino acids in T1R2 and T1R3

to confirm the validity of AF-M predictions. Interestingly, the putative binding site of

Gur-2 in the human STR pave the way for improving the binding affinity of Gur-2 by

site-directed mutagenesis, and therefore generating peptides with a potent suppressive

sweet taste effect in humans. Moreover, the comparison of the binding of Gur-1, Gur-2

and Brz to mT1R2/T1R3 and hT1R2/T1R3 revealed that in Gur-1 and Gur-2 as in Brz,

two adjacent aromatic amino acids are important for the interactions with the CRD of

T1R2 and T1R3 subunits. It is also interesting to use Gur-1 and Gur-2 (and their

isoforms) together in the future experiments to know if they may have a synergistic effect

on the suppression of the sweet taste in humans. Finally, gurmarin-like peptides could be

interesting drug scaffolds which may help to reduce the craving for sugar and thus help to

regulate short-term blood glucose levels.

References

Ayachit, G., Shaikh, I., Sharma, P., Jani, B., Shukla, L., Sharma, P., Bhairappanavar, S.B., Joshi, C.,
and Das, J. 2019. De novo transcriptome of Gymnema sylvestre identified putative lncRNA and
genes regulating terpenoid biosynthesis pathway. Sci Rep. 9:14876.

Banerjee, S., Varga, J.K., Kumar, M., Zoltsman, G., Isupov, M.N., Rosenzweig, R., Schueler-
Furman, O., and Wiener, R. 2023. Structural study of UFL1-UFC1 interaction uncovers the
importance of UFL1 N-terminal helix for ufmylation. bioRxiv.

Belloir, C., Brulé, M., Tornier, L., Neiers, F., and Briand, L. 2021. Biophysical and functional
characterization of the human TAS1R2 sweet taste receptor overexpressed in a HEK293S
inducible cell line. Sci Rep. 11:22238.

Bezençon, C., Coutre, J. le, and Damak, S. 2007. Taste-signaling proteins are coexpressed in
solitary intestinal epithelial cells. Chem Senses. 32:41–49.

Chang, L., and Perez, A. 2022. AlphaFold encodes the principles to identify high affinity peptide
binders. bioRxiv.

Chang, L., and Perez, A. 2023. Ranking Peptide Binders by Affinity with AlphaFold**. Angew
Chem Int Ed. 62:e202213362.

Cui, M., Jiang, P., Maillet, E., Max, M., Margolskee, R.F., and Osman, R. 2006. The heterodimeric
sweet taste receptor has multiple potential ligand binding sites. Curr Pharm Des. 12:4591–4600.

Devangan, S., Varghese, B., Johny, E., Gurram, S., and Adela, R. 2021. The effect of Gymnema
sylvestre supplementation on glycemic control in type 2 diabetes patients: A systematic review
and meta-analysis. Phytother Res PTR. 35:6802–6812.

Evans, R., O'Neill, M., Pritzel, A., Antropova, N., Senior, A., Green, T., Žídek, A., Bates, R.,
Blackwell, S., Yim, J., et al. 2022. Protein complex prediction with AlphaFold-Multimer. bioRxiv.

Ghanavatian, P., Khalifeh, K., and Jafarian, V. 2016. Structural features and activity of Brazzein
and its mutants upon substitution of a surfaced exposed alanine. Biochimie. 131:20–28.

Gori, A., Lodigiani, G., Colombarolli, S.G., Bergamaschi, G., and Vitali, A. 2023. Cell Penetrating
Peptides: Classification, Mechanisms, Methods of Study, and Applications. ChemMedChem.
18:e202300236.

Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. 2010. New
algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
performance of PhyML 3.0. Syst Biol. 59:307–321.

Gutierrez, S., Tyczynski, W.G., Boomsma, W., Teufel, F., and Winther, O. 2022. MembraneFold:
Visualising transmembrane protein structure and topology. 2022.12.06.518085.

Hara, K., Inada, Y., Ono, T., Kuroda, K., Yasuda-Kamatani, Y., Ishiguro, M., Tanaka, T., Misaka, T.,
Abe, K., and Ueda, M. 2012. Chimeric Yeast G-Protein α Subunit Harboring a 37-Residue C-
Terminal Gustducin-Specific Sequence Is Functional in *Saccharomyces cerevisiae*. Biosci
Biotechnol Biochem. 76:512–516.

Holm, L. 2020. Using Dali for Protein Structure Comparison. Methods Mol Biol Clifton NJ.
2112:29–42.

Imoto, T., Miyasaka, A., Ishima, R., and Akasaka, K. 1991. A novel peptide isolated from the
leaves of Gymnema sylvestre--I. Characterization and its suppressive effect on the neural
responses to sweet taste stimuli in the rat. Comp Biochem Physiol A. 100:309–314.

Imoto, T., Sasamoto, K., and Ninomiya, Y. 2001. Beta-cyclodextrin inhibits the sweet taste
suppressing activity of gurmarin by the formation of an inclusion complex with aromatic
residues in gurmarin. Can J Physiol Pharmacol. 79:836–840.

Jiang, P., Ji, Q., Liu, Z., Snyder, L.A., Benard, L.M.J., Margolskee, R.F., and Max, M. 2004a. The
cysteine-rich region of T1R3 determines responses to intensely sweet proteins. J Biol Chem.
279:45068–45075.

Jiang, P., Ji, Q., Liu, Z., Snyder, L.A., Benard, L.M.J., Margolskee, R.F., and Max, M. 2004b. The
cysteine-rich region of T1R3 determines responses to intensely sweet proteins. J Biol Chem.
279:45068–45075.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K.,
Bates, R., Žídek, A., Potapenko, A., et al. 2021. Highly accurate protein structure prediction with
AlphaFold. Nature. 596:583–589.

Kalariya, K.A., Gajbhiye, N., Minipara, D., Meena, R.P., Kumar, S., Saha, A., Trivedi, A., and
Manivel, P. 2019. Deep sequencing-based de novo transcriptome analysis reveals biosynthesis of
gymnemic acid in Gymnema sylvestre (Retz.) Schult. Ecol Genet Genomics. 13:100047.

Kalariya, K.A., Minipara, D.B., and Manivel, P. 2018. De novo transcriptome analysis deciphered
polyoxypregnane glycoside biosynthesis pathway in Gymnema sylvestre. 3 Biotech. 8:381.

Kamei, K., Takano, R., Miyasaka, A., Imoto, T., and Hara, S. 1992. Amino acid sequence of sweet-
taste-suppressing peptide (gurmarin) from the leaves of Gymnema sylvestre. J Biochem (Tokyo).
111:109–112.

Kaminski, G.A., Friesner, R.A., Tirado-Rives, J., and Jorgensen, W.L. 2001. Evaluation and
Reparametrization of the OPLS-AA Force Field for Proteins via Comparison with Accurate
Quantum Chemical Calculations on Peptides. J Phys Chem B. 105:6474–6487.

Katoh, K., and Toh, H. 2008. Recent developments in the MAFFT multiple sequence alignment
program. Brief Bioinform. 9:286–298.

Kohno, D., Koike, M., Ninomiya, Y., Kojima, I., Kitamura, T., and Yada, T. 2016. Sweet Taste
Receptor Serves to Activate Glucose- and Leptin-Responsive Neurons in the Hypothalamic
Arcuate Nucleus and Participates in Glucose Responsiveness. Front Neurosci. 10:502.

Kyriazis, G.A., Soundarapandian, M.M., and Tyrberg, B. 2012. Sweet taste receptor signaling in
beta cells mediates fructose-induced potentiation of glucose-stimulated insulin secretion. Proc
Natl Acad Sci U S A. 109:E524-532.

Lin, Z., Akin, H., Rao, R., Hie, B., Zhu, Z., Lu, W., Smetanin, N., Verkuil, R., Kabeli, O., Shmueli, Y.,
et al. 2023. Evolutionary-scale prediction of atomic-level protein structure with a language
model. Science. 379:1123–1130.

Lindemann, B. 2001. Receptors and transduction in taste. Nature. 413:219–225.

Liu, J., Zhang, Z., Moreno-Delgado, D., Dalton, J.A., Rovira, X., Trapero, A., Goudet, C., Llebaria,
A., Giraldo, J., Yuan, Q., et al. 2017. Allosteric control of an asymmetric transduction in a G
protein-coupled receptor heterodimer. eLife. 6:e26985.

Maaroufi, H., Potvin, M., Cusson, M., and Levesque, R.C. 2021. Novel antimicrobial anionic
cecropins from the spruce budworm feature a poly-L-aspartic acid C-terminus. Proteins.
89:1205–1215.

Miyasaka, A., and Imoto, T. 1995. Electrophysiological characterization of the inhibitory effect of
a novel peptide gurmarin on the sweet taste response in rats. Brain Res. 676:63–68.

Mosinger, B., Redding, K.M., Parker, M.R., Yevshayeva, V., Yee, K.K., Dyomina, K., Li, Y., and
Margolskee, R.F. 2013. Genetic loss or pharmacological blockade of testes-expressed taste
genes causes male sterility. Proc Natl Acad Sci U S A. 110:12319–12324.

Nakagawa, Y., Nagasawa, M., Yamada, S., Hara, A., Mogami, H., Nikolaev, V.O., Lohse, M.J.,
Shigemura, N., Ninomiya, Y., and Kojima, I. 2009. Sweet taste receptor expressed in pancreatic
beta-cells activates the calcium and cyclic AMP signaling systems and stimulates insulin
secretion. PloS One. 4:e5106.

Nelson, G., Hoon, M.A., Chandrashekar, J., Zhang, Y., Ryba, N.J., and Zuker, C.S. 2001.
Mammalian sweet taste receptors. Cell. 106:381–390.

Ninomiya, Y., and Imoto, T. 1995. Gurmarin inhibition of sweet taste responses in mice. Am J
Physiol. 268:R1019-1025.

Ninomiya, Y., Imoto, T., and Sugimura, T. 1999. Sweet Taste Responses of Mouse Chorda
Tympani Neurons: Existence of Gurmarin-Sensitive and -Insensitive Receptor Components. J
Neurophysiol. 81:3087–3091.

Ninomiya, Y., Inoue, M., and Imoto, T. 1998. Reduction of the Suppressive Effects of Gurmarin
on Sweet Taste Responses by Addition of -Cyclodextrin. Chem Senses. 23:303–307.

Obi, P., and Natesan, S. 2022. Membrane Lipids Are an Integral Part of Transmembrane
Allosteric Sites in GPCRs: A Case Study of Cannabinoid CB1 Receptor Bound to a Negative
Allosteric Modulator, ORG27569, and Analogs. J Med Chem. 65:12240–12255.

Ohkuri, T., Yasumatsu, K., Horio, N., Jyotaki, M., Margolskee, R.F., and Ninomiya, Y. 2009.
Multiple sweet receptors and transduction pathways revealed in knockout mice by temperature
dependence and gurmarin sensitivity. Am J Physiol Regul Integr Comp Physiol. 296:R960-971.

Ota, M., Shimizu, Y., Tonosaki, K., and Ariyoshi, Y. 1998a. Role of hydrophobic amino acids in
gurmarin, a sweetness-suppressing polypeptide. Biopolymers. 45:231–238.

Ota, M., Shimizu, Y., Tonosaki, K., and Ariyoshi, Y. 1998b. Synthesis, characterization, and
sweetness-suppressing activities of gurmarin analogues missing one disulfide bond.
Biopolymers. 46:65–73.

Ozden, B., Kryshtafovych, A., and Karaca, E. 2023. The Impact of AI-Based Modeling on the
Accuracy of Protein Assembly Prediction: Insights from CASP15. Bioinformatics.

Pallaghy, P.K., Nielsen, K.J., Craik, D.J., and Norton, R.S. 1994. A common structural motif
incorporating a cystine knot and a triple-stranded beta-sheet in toxic and inhibitory
polypeptides. Protein Sci Publ Protein Soc. 3:1833–1839.

Park, J., Selvam, B., Sanematsu, K., Shigemura, N., Shukla, D., and Procko, E. 2019. Structural
architecture of a dimeric class C GPCR based on co-trafficking of sweet taste receptor subunits. J
Biol Chem. 294:4759–4774.

Perez-Aguilar, J.M., Kang, S.-G., Zhang, L., and Zhou, R. 2019. Modeling and Structural
Characterization of the Sweet Taste Receptor Heterodimer. ACS Chem Neurosci. 10:4579–4592.

Rayo-Morales, R., Segura-Carretero, A., Poirier, N., Briand, L., and Garcia-Burgos, D. 2023.
Sweet-inhibiting effects of gurmarin on intake during repeated acute and long-term sugar
exposure: A behavioural analysis using an animal model. J Funct Foods. 108:105743.

Ren, X., Zhou, L., Terwilliger, R., Newton, S.S., and Araujo, I.E. de. 2009. Sweet taste signaling
functions as a hypothalamic glucose sensor. Front Integr Neurosci. 3:12.

Ruiz-Avila, L., Wong, G.T., Damak, S., and Margolskee, R.F. 2001. Dominant loss of
responsiveness to sweet and bitter compounds caused by a single mutation in alpha -gustducin.
Proc Natl Acad Sci U S A. 98:8868–8873.

Rydberg, H.A., Kunze, A., Carlsson, N., Altgärde, N., Svedhem, S., and Nordén, B. 2014. Peptide-
membrane interactions of arginine-tryptophan peptides probed using quartz crystal
microbalance with dissipation monitoring. Eur Biophys J EBJ. 43:241–253.

Sainz, E., Cavenagh, M.M., LopezJimenez, N.D., Gutierrez, J.C., Battey, J.F., Northup, J.K., and
Sullivan, S.L. 2007. The G-protein coupling properties of the human sweet and amino acid taste
receptors. Dev Neurobiol. 67:948–959.

Shigemura, N., Nakao, K., Yasuo, T., Murata, Y., Yasumatsu, K., Nakashima, A., Katsukawa, H.,
Sako, N., and Ninomiya, Y. 2008. Gurmarin sensitivity of sweet taste responses is associated with
co-expression patterns of T1r2, T1r3, and gustducin. Biochem Biophys Res Commun. 367:356–
363.

Shimizu, M., Goto, M., Kawai, T., Yamashita, A., and Kusakabe, Y. 2014. Distinct human and
mouse membrane trafficking systems for sweet taste receptors T1r2 and T1r3. PloS One.
9:e100425.

Sigoillot, M., Brockhoff, A., Lescop, E., Poirier, N., Meyerhof, W., and Briand, L. 2012a.
Optimization of the production of gurmarin, a sweet-taste-suppressing protein, secreted by the
methylotrophic yeast Pichia pastoris. Appl Microbiol Biotechnol. 96:1253–1263.

Sigoillot, M., Brockhoff, A., Meyerhof, W., and Briand, L. 2012b. Sweet-taste-suppressing
compounds: current knowledge and perspectives of application. Appl Microbiol Biotechnol.
96:619–630.

Sigoillot, M., Brockhoff, A., Neiers, F., Poirier, N., Belloir, C., Legrand, P., Charron, C., Roblin, P.,
Meyerhof, W., and Briand, L. 2018. The Crystal Structure of Gurmarin, a Sweet Taste-
Suppressing Protein: Identification of the Amino Acid Residues Essential for Inhibition. Chem
Senses. 43:635–643.

Teufel, F., Refsgaard, J.C., Kasimova, M.A., Deibler, K., Madsen, C.T., Stahlhut, C., Grønborg, M.,
Winther, O., and Madsen, D. 2023. Deorphanizing Peptides Using Structure Prediction. J Chem
Inf Model. 63:2651–2655.

Van Kempen, M., Kim, S.S., Tumescheit, C., Mirdita, M., Lee, J., Gilchrist, C.L.M., Söding, J., and
Steinegger, M. 2023. Fast and accurate protein structure search with Foldseek. Nat Biotechnol.

Xue, L.C., Rodrigues, J.P., Kastritis, P.L., Bonvin, A.M., and Vangone, A. 2016. PRODIGY: a web
server for predicting the binding affinity of protein-protein complexes. Bioinforma Oxf Engl.
32:3676–3678.

Yasumatsu, K., Ohkuri, T., Sanematsu, K., Shigemura, N., Katsukawa, H., Sako, N., and Ninomiya,
Y. 2009. Genetically-increased taste cell population with G(alpha)-gustducin-coupled sweet
receptors is associated with increase of gurmarin-sensitive taste nerve fibers in mice. BMC
Neurosci. 10:152.

Yoshie, S., Miyasaka, A., and Imoto, T. 1994. Histological localization of the sweet taste receptor
in rat taste buds by the use of gurmarin, a sweet taste-suppressing peptide. Arch Histol Cytol.
57:531–534.

Acknowledgments

679 I thank members of the Centre of Bioinformatics at the Institut de Biologie Intégrative et
680 des Systèmes (IBIS), Université Laval for their assistance.

681   **Supplementary Material**

683   **Table S1.** Physico-chemical characteristics of Gur-1 and gurmarin-like peptides

	Mature peptide	Length	Net charge	pI	Hydrophobicity (Kcal/mol) [**]
Gur-1	QQCVKKDELCIPYYLDCCEPLECKKVNWWDHKCIG	35	-1	5.44[*]	+37.23
Gur-2	ELCQEKDEPCVPNFMECCEPYKCRSVSWWELKCIS	35	-3	4.18	+35.02
Gur-3	AKCLKKGQKCNPPYTPCCRPYRCVQTSRPYKCGI	34	+8	9.76	+29.10
Gur-4	SDCLPEGALCGLGIECCPPFHCSEPMRPFRCRR	33	0	6.76	+28.49
Gur-5	PYCLPEGAFCGHQLDIPCCPPFHCSEPMRPFMCER	35	-1	5.20	+24.80
Gur-6	LNCRQLLELCGPGVSDCCPPLECLRTIENRCQGYGD	36	-2	4.19	+30.56
Gur-7	LICLPEGALCGGSLGECCPPLLCIRTLDIRCRKLGYV	37	+1	7.60	+19.76
Gur-8	LNCLSIGETCGPGVGYCCFPLICSVGTEHVGFKCVYVATKSKA	43	+1	7.61	+23.80
Gur-9	LSCVEEGNDCEPIENECCPSLECIGSYYGFKCEHPV	36	-7	3.59	+40.12

684   [*]The experimental pI is 4.5 (Imoto et al. 1991).
685   [**]more the number in the column is small more the peptide is hydrophobic (Gur-2 is more hydrophobic than Gur-1).

687   **Table S2**. Pairwise structure alignment of Gur-2 to Gur-9 with Gur-1

Peptides	Z-score	RMSD (A)	Length alignment	Residues number	% Identity
Gur-1	8.8	0.0	35	35	100
Gur-2	7.2	0.9	35	35	51
Gur-3	6.0	1.4	34	34	35
Gur-8	5.1	2.1	35	43	31
Gur-6	5.0	2.0	33	36	42
Gur-9	5.0	2.1	34	36	32
Gur-4	5.0	2.0	33	33	24
Gur-5	4.4	2.8	34	35	21
Gur-7	3.9	2.3	33	37	30

689 **Table S3**. Interactions (≤ 3.5 Å) of Gur-1 amino acid residues with mouse T1R2, T1R3
690 monomers and T1R2/T1R3 heterodimer.

Gur-1	ICD of T1R2	CRD/TMD of T1R3	ICD of T1R2/T1R3	CRD/TMD of T1R2/T1R3
Gln2	Met821, Ser824	Thr522, Cys523, Pro525	Gln817, Thr820, Met821	
Leu9			Tyr654, Met658	
Ile11	Tyr654		Val643, Tyr654, Met658	
Pro12				
Tyr13	Arg582, Tyr741	Pro700	Pro579, Arg582, Ser583, Tyr741, Phe812, Ile816	Arg539, Thr693, Pro695, Pro698
Tyr14	Ser583, Arg636, Ile640, Tyr741, Ile816	Cys526, Asn527, Gln528, Arg542	Arg636, Ile640, Val643, Phe644, Tyr741, Ile816	Gly691, Arg692, Thr693, Ile701
Leu15			Ile816	
Asp16	Asn813	Lys510	Asn813	
Trp28	His661		His661, Tyr664	Arg710, Ser766
Trp29	Phe656, Trp657, Met658, Arg659, Tyr660, His661, Gly662, Pro663		Met588, Gln639, Trp657, Met658, Arg659, His661, Pro663, Tyr664	Asn708, Arg710, Asn711, Leu714
Asp30	Met658		Met658, Arg659	

691 CRD: cysteine-rich domain; ICD: intracellular domain.
692 Amino acid residues of T1R2 and T1R3 are numbered without the signal peptide.
693

695 **Table S4**. Interactions (≤ 3.5 Å) of Gur-2 amino acid residues with mouse T1R2, T1R3
696 monomers and T1R2/T1R3 heterodimer.

Gur2	ICD of T1R2	CRD/TMD of T1R3	ICD of T1R2/T1R3	CRD/TMD of T1R2/T1R3
Leu2	Gln817			
Val11		Ser697		
Asn13	Pro579, Arg582, Tyr741, Phe812, Tyr741, Phe812	Pro700	Gln817, Thr820, Met821	Arg539, Thr693
Phe14	Phe644, Tyr741, Ile816		Arg648, Thr820	Gly691, Thr693
Met15		Arg542, Trp696, Ser697, Leu699		
Glu16	Ser809, Asn813			
Trp28	His661, Gly662, Pro663, Tyr664		Arg582, Ser583, Arg636, Tyr741	
Trp29	Phe656, Trp657, Met658, Arg659, Tyr660, His661, Gly662, Pro663	Leu767, Ala768, Asn769	Arg636, Gln639, Met658	Asn708, Asn711
Glu30	Met658	Trp712	Met658	

697 CRD: cysteine-rich domain; ICD: intracellular domain.
698 Amino acid residues of T1R2 and T1R3 are numbered without the signal peptide.
699

700 **Table S5**. Interactions (≤ 3.5 Å) of Gur-1 amino acid residues with human T1R2, T1R3
701 monomers and T1R2/T1R3 heterodimer.

Gur-1	ICD of T1R2	CRD/TMD of T1R3	ICD of T1R2/T1R3	CRD/TMD of T1R2/T1R3
Leu9	Tyr650, Ser651, Val654		Tyr650, Val654	No interactions
Ile11	Ala639, Tyr650	Asp690, His692, Met693	Ala639, Tyr650	
Pro12		His692, Met693		
Tyr13	Pro575, Arg578, Ser579, Tyr737, Pro805, Phe808, Ile812	His692, Met693	Pro575, Arg578, Ser579, Asn736, Phe808, Ile812	
Tyr14	Ser579, Arg632, Ile636, Ala639, Phe640, Tyr737, Ile812	Gln523, Arg537, Trp691, His692, Leu694, Pro695	Arg632, Ile636, Phe640, Pro734, Thr735, Asn736, Glu739, Ile812	
Leu15	Ala639, Phe640, Ala643, Ile812		Ile812	
Asp16	Asn809, Ile812	His692	Asn809, Ile812	
Trp28	Tyr660	Trp707, Ala763	Gln657, Tyr660	
Trp29	Val654, Gln657, Pro659, Tyr660	Arg703, Thr704, Trp707	Val631, Gln635, Trp653, Val654, Gln657, Gly658, Pro659	
Asp30	Val654, Arg655, Tyr660		Ser651, Val654, Arg655	

702 CRD: cysteine-rich domain; ICD: intracellular domain.
703 Amino acid residues of T1R2 and T1R3 are numbered without the signal peptide.
704

706 **Table S6**. Interactions (≤ 3.5 Å) of Gur-2 amino acid residues with human T1R2, T1R3
707 monomers and T1R2/T1R3 heterodimer.

Gur2	ICD of T1R2	CRD/TMD of T1R3	ICD of T1R2/T1R3	CRD/TMD of T1R2/T1R3
Pro9			Tyr650, Val654	
Val11	Ala639, Ala643, Tyr650	Met693	Ala639, Phe640, Ile812, Thr816	
Pro12	Ile812, Thr816		Ile812, Gln813, Thr816	
Asn13	Gln813, Thr816	His692	Gln813, Thr816, Met817	
Phe14	Ala643, Ser644	Asp690, Trp691, His692, Met693,	Ser644, Thr816, Arg819	
Met15	Pro647	His692	Ala643, Pro647	
Trp28	Arg578, Ser579, Ala580, Gly581, Met584, Arg632, Gln635	Ala763	Arg578, Arg579, Ala580, Gly581, Arg631, Gln635, Tyr737, Glu739	
Trp29	Met584, Gln635, Tyr650, Pro659, Tyr660	Trp707	Arg631, Gln635, Ala639, Trp653, Pro659	Arg706
Glu30	Val654		Val654	

708 CRD: cysteine-rich domain; ICD: intracellular domain.
709 Amino acid residues of T1R2 and T1R3 are numbered without the signal peptide.
710

717 **Table S7**. *In silico* assay competitions between Gur-1, Gur-2, T1R2, T1R3 monomers
718 and T1R2/T1R3 heterodimer.

Input for AF-M	Binding site of Gur-1 and Gur-2	ΔG (kcal mol-1)	pTM+ipTM
mT1R2, Gur-1, Gur-2	Gur-1 in ICD, Gur-2 in CRD near TMD	-10.4, -7.8	0.521
mT1R3, Gur-1, Gur-2	Gur-1 clashes in TMD near ICD, Gur-2 in CRD near TMD	N.D, -7.3	0.376
mT1R2, mT1R3, Gur-1, Gur-2	Gur-1 clashes in ICD of mT1R2, Gur-2 in CRD near TMD of mT1R2	N.D, -6.4	0.570
hT1R2, Gur-1, Gur-2	Gur-1 in ICD, Gur-2 in CRD near TMD	-9.4, -7.2	0.391
hT1R3, Gur-1, Gur-2	Gur-1 in TMD near ICD, Gur-2 in CRD near TMD	-11.0, -6.4	0.453
hT1R2, hT1R3, Gur-1, Gur-2	Gur-1 clashes in the ICD of T1R2, Gur-2 in CRD near TMD of T1R2	N.D, -5.7	0.595

719 AF-M: AlphaFold-Multimer
720 Mouse T1R2 (mT1R2); human T1R2 (hT1R2)
721 VFTM: Venus flytrap module; CRD: cysteine-rich domain; TMD: trans-membrane domain; ICD:
722 intracellular domain
723 N.D: not determined.

Fig. S1. Multiple sequence alignment of *G. sylvestre* gurmarin-like peptides. The aligned
sequences are separated in nine groups, named Gur-I to Gur-IX. Gur-I group that
contains Gur-1 (UniProt ID: A0A3S9JKY1) and its isoforms is the large group. Trp28
and Trp29 that are important for the suppressive sweet taste effect is conserved in Gur-I
and Gur-II groups. SRA accession ID is indicated with each sequence.

Fig. S2. Structural similarity dendrogram of Gur-1 to Gur-9. Labels are linked to
structural summaries. The dendrogram is derived by average linkage clustering of the
structural similarity matrix (Dali Z-scores, see table S2).

Fig. S3. Interactions of Brazzein (Brz) with human T1R2/T1R3 heterodimer. Brz (pink)
binds in cysteine-rich domain (CRD, region I516-E525 in sticks) of hT1R3 (cyan) with
A19 and Y54 amino acid residues that are important for sweet taste effect of Brz. Amino
acid residues of T1R3 are numbered without the signal peptide.

Fig. S4. The interaction of Gur-1 with the mouse sweet taste receptor T1R2/T1R3 (m
T1R2/T1R3). (**A**) the interaction of Gur-1 (pink) with the intracellular domain (ICD) of
mT1R2. (**B**) zoom in the region of the interaction of Gur-1 with the ICD of mT1R2. (**C**)
electrostatic potential of the ICD of mT1R2. The negatively, positively and hydrophobic
amino acid residues are in red, blue and white, respectively.

Fig. S5. Interactions of Gur-2 with the mouse T1R2, T1R3 monomers and T1R2/T1R3
heterodimer. (**A**) and (**B**) Gur-2 (salmon) binds in the ICD of the T1R2 monomer (green)
and in the CRD near TMD of the T1R3 monomer (cyan) with F14, W28 and W29 amino
acid residues, respectively. (**C**) and (**D**) Gur-2 binds in the ICD and the CRD near TMD
of T1R2 subunit of T1R2/T1R3 heterodimer, respectively. For details of bonds between
amino acid residues, see Table S4.

768
769

770
771
Fig. S6. The prediction of tunnels and channels in mouse T1R2/T1R3 structures. (A)
electrostatic potential of Gur-1. Important amino acids Y13-14 and W28-29 for sweet
suppressive effect are present in hydrophobic region. (B) electrostatic potential of
mT1R2/T1R3. The negatively, positively and hydrophobic amino acid residues are in red,
blue and white, respectively. (C) superposition of mT1R2/T1R3-Gur-1 complex and
electrostatic potential of mT1R2/T1R3. Gur-1 binds in the channel entrance. (D)
predicted tunnels and channels in mT1R2-R3 (white spheres). (E) superposition of
mT1R2/T1R3-Gur-1 complex and representation of tunnels and channels of mT1R2/T1R3.
Gur-1 binds in the channel entrance. T1R2, T1R3 and Gur-1 are in green, cyan and pink,
respectively.