Individual differences in wellbeing are supported by separable sets of co-active self- and visual-attention-related brain networks

Yumeng Ma[1,2,3,*] & Jeremy I Skipper[2]

[1] Department of Neuroscience, Physiology and Pharmacology, University College London, London, UK

[2] Experimental Psychology, University College London, London, UK

[3] Department of Psychology, Emory University, Atlanta, GA, USA

[*] Corresponding author: yumeng.ma@emory.edu

Abstract

How does the brain support 'wellbeing'? Because it is a multidimensional construct, it is likely the product of multiple co-active brain networks that vary across individuals. This is perhaps why prior neuroimaging studies have found inconsistent anatomical associations with wellbeing. Furthermore, these used 'laboratory-style' or 'resting-state' methods not amenable to finding manifold networks. To address these issues, we had participants watch a full-length romantic comedy-drama film during functional magnetic resonance imaging. We hypothesised that individual differences in wellbeing measured before scanning would be correlated with individual differences in brain networks associated with 'embodied' and 'narrative' self-related processing. Indeed, searchlight spatial inter-participant representational similarity and subsequent analyses revealed seven sets of co-activated networks associated with individual differences in wellbeing. Two were 'embodied self' related, including brain regions associated with autonomic and affective processing. Three sets were 'narrative self' related, involving speech, language, and autobiographical memory related regions. Finally, two sets of visual-attention related networks emerged. These results suggest that the neurobiology of wellbeing in the real-world is supported by diverse but functionally definable and separable sets of networks. This has implications for psychotherapy where individualised interventions might target, e.g., neuroplasticity in language-related narrative over embodied self or visual-attentional related processes.

Keywords. Brain; Embodied Self; fMRI; Individual Differences; Inter-Participant Correlation; Narrative Self; Neurobiology of Wellbeing; Representational Similarity Analysis; Self; Wellbeing

Significance Statement

Humans deeply care about their own and others wellbeing. Yet, the role the brain plays in generating it remains unestablished, perhaps because wellbeing means different things to different people. Here, we asked whether individual differences in wellbeing are associated with differences in brain networks supporting 'embodied' and 'narrative' selves. That is, how people feel in their bodies and their emotions might contribute more to wellbeing in some individuals. In others, the stories people tell about themselves might be a more important determinant. Supporting this, we found that different self-ratings of wellbeing are associated with different embodied and narrative self-related networks. We also found differences in networks associated with visual-attention, suggesting that what people attend to in the visual world also contributes to wellbeing. These results have implications for how mental health professionals and other individuals might intervene to improve wellbeing by differentially focusing on embodied-self, narrative-self, or visual-attentional related processes.

Introduction

'Wellbeing' is a multidimensional construct, without a single definition (1, 2). It is often viewed from the hedonic or 'subjective wellbeing' perspective, characterised as 'a person feeling and thinking' their 'life is desirable regardless of how others see it.' (3, 4). This view contains both subjective emotional/affective (i.e., feeling) and evaluative/cognitive (i.e., thinking) components (3, 5). It is sometimes contrasted with a eudaimonic or 'psychological wellbeing' perspective that is more associated with meaning, purpose, and self-realisation components of human activity (6–8). Though these general characterizations grossly capture unique components of wellbeing, they are also highly interrelated (9, 10).

Reflecting the multidimensional nature of wellbeing, there are more than 100 self-report measures that are used to evaluate about 200 different dimensions (2, 11). These measures reflect different perspectives, like those above, sometimes with no overlap on any dimension measured (11–14). Nonetheless, they capture something important as they have variously been used to show that wellbeing is associated with better physical health and decreased mortality (15–19), higher chances of recovery from serious mental illnesses like schizophrenia or bipolar disorder (20, 21), increased productivity and higher job performance (22–24), and improved social relationships (25).

As might be expected from a multidimensional construct, these self-report measures correlate wellbeing with a large array of individual differences. To give a non-exhaustive overview, wellbeing is associated with individual differences in hearing (26), interoception (27, 28), visual attention (29), emotional intelligence (30, 31), up-regulating positive emotions (32), following feelings (33), cognitive emotional regulation strategies (34, 35), facets of mindfulness (36, 37), rumination (38), finding meaning (39), narratives about self-growth (40), the abstractness versus concreteness of self-construal (41), degree of past, present, and future thinking (42), nostalgia (43), authenticity (44), 'mental toughness' (45), personality (46), ethical orientation (47), and social participation and outgoingness (48, 49).

Neurobiology of Wellbeing

Despite its importance, our understanding of the neurobiology of wellbeing is limited. Results have been highly variable, with arguably 'no consistent associations' between specific brain regions and wellbeing across studies (50–52). Reviews of the 20 to 60 neuroimaging studies related to wellbeing show that activation spans much of the brain, though not reproducibly (50–55). We suggest that there are a few interrelated reasons for this. The first is implicit in the above, i.e., the multidimensional nature of wellbeing and the individual differences mirroring this dimensionality. This implies that there will be multiple underlying, co-active, and interacting brain networks supporting the manifestation of wellbeing

distributed throughout the whole brain. Furthermore, these networks would be expected to be different or weighted differently depending on individual differences.

Yet, the paradigms used in neuroimaging studies of wellbeing have likely ensured variability of results across studies while not being able to find these different networks or weightings within any one study. Specifically, about half of existing studies can be characterised as 'laboratory-style', involving reductionist paradigms with relatively unnatural but well controlled stimuli and tasks (51). Stimuli include affective pictures, music, playing cards, robot arms, sentences, and video games, among others. Tasks involve emotional reactivity, facial recognition, gender discrimination, imagination, monetary rewards, repetition detection, etc.. The remaining studies use a 'resting-state' paradigm with no controlled stimuli or tasks (51). Participants might be trying to stay awake, fixating cross-hairs, trying not to think, visualising, and/or silently talking to themselves (56, 57). This heterogeneity of stimuli and tasks in laboratory-style and behaviours in resting-state studies likely produces anatomically variable results across studies.

Furthermore, neither paradigm as employed is amenable to finding the many regions and networks likely supporting wellbeing. Laboratory-style studies usually examined only affective perception and/or experienced affect, in exclusion to many other possible dimensions like the evaluative aspects of wellbeing (51). Thus, the subtractive logic often employed could only possibly reveal 'blobs' related to those few dimensions, while obscuring associated networks (due to the region focused methods) and completely hiding networks associated with dimensions not under study. When network analyses were employed in either paradigm, only a few networks could possibly be discovered because only one or a small number of seed regions of interest were used in nearly all studies (51).

Wellbeing and Self

What organising principles might be used to understand the whole-brain distribution of networks we propose to support individual differences in wellbeing? We and others have suggested that wellbeing is self-related and can be mapped onto self-related brain processes (58–60). Indeed, self-report measures of wellbeing include self-related dimensions either implicitly (because all require self-related judgements) or explicitly, e.g., self-acceptance, self-care, self-confidence, self-control, self-discovery, self-efficacy, self-esteem, self-realisation, self-regard, and self-satisfaction (11). Self-related processing encompasses large portions of the brain but might be grossly divided into those supporting 'embodied' and 'narrative' self-related processes (61–65), though this is not to say these are also not intimately interrelated (66, 67).

The 'embodied self' refers to aspects of the self that are not or at least less obviously based on semantic representations and focus on the self in the present moment, like interoceptive awareness, the experience of body ownership, and emotions (68, 69). We suggest that this primarily links embodied self-related processes and supporting brain networks to individual differences in the emotional/affective (i.e., feeling) component of 'subjective wellbeing'. For example, individuals better at interoception are better at emotional regulation and this is particularly reliant on the insula (70–75). Therefore it can be expected that the insula predicts higher wellbeing in those people. Indeed, many of the regions involved in the embodied self (68), nested in emotional processing more generally (76, 77), are also implicated in neuroimaging studies of wellbeing. In addition to the insula, these include the amygdala, anterior cingulate cortex, basal ganglia, dorso- and medial prefrontal cortices, and thalamus (50–55).

In contrast to the embodied self, the 'narrative self' has more obvious semantic content that extends the present to past autobiographical memories and imagined futures (63). Though it is almost never acknowledged, these narratives are typically constructed and manipulated with language (60). We suggest that this links narrative self-related processes and associated brain networks to individual differences in the evaluative/cognitive (i.e., thinking) component of 'subjective wellbeing' and the meaning and purpose components of 'psychological wellbeing'. Therefore, it can be expected that language and autobiographical memory related regions will predict individual differences in wellbeing more in some people. Indeed, of all regions, the superior temporal gyrus (associated with language comprehension) and precuneus (associated with autobiographical memory) are often the most predictive of mental health and wellbeing (for a review, see)(60). These regions also regularly appear in neuroimaging studies of wellbeing (50–55).

Study Overview

In summary, existing work on the neurobiology of wellbeing has yielded variable results, likely because 'wellbeing' is multidimensional and subject to individual differences. Furthermore, the neuroimaging methods used have likely contributed to this variability and are not conducive to discovering the multiple co-activated real-world networks that likely underlie these multiple dimensions. We proposed a general self-related framework for organising expectations about which of these co-activated networks might be associated with individual differences in wellbeing. Specifically, embodied self-related networks will be larger determinants of wellbeing in some while narrative self-related networks will be more important in other individuals.

To be able to test this framework, we developed methods capable of capturing individual differences in a construct that relies on multiple co-active real-world brain networks. This required a stimulus that engages a full range of networks in a manner that is similar to everyday life. We accomplished this by asking 20 participants to complete a wellbeing questionnaire and then watch the romantic comedy-drama film '*500 Days of Summer*' during fMRI (78). Next, we developed an analysis pipeline that could find corresponding individual differences in wellbeing and brain activity, while grouping this activity into networks and sets of co-active networks (Figure 1).

First, 'searchlight spatial inter-participant representational similarity analysis' (SSIP-RSA)(79–81) determines how variations in brain activity patterns across participants in a searchlight at each time point in the film relates to variations in wellbeing as measured by the NIH Toolbox (82)(Figure 1a and 1b). Second, independent component analysis (ICA) decomposes the resulting timeseries into 100 spatial independent components (ICs), i.e., networks, and associated timecourses, while reducing dimensionality (Figure 1c). Third, hierarchical clustering of those timecourses is used to find sets of co-active networks and further reduce dimensionality (Figure 1d, top). Finally, we quantify the functional properties of these sets of networks by spatially correlating them with 10 pre-chosen term-based neuroimaging meta-analyses from a database of thousands of meta-analyses (i.e., Neurosynth)(83).

Figure 1. Analysis pipeline. *a, Searchlight spatial inter-participant representational similarity analysis (SSIP-RSA). (1) For a specific time point (t1), the correlation distance of brain activity patterns inside a searchlight centred at a voxel (v1) with a radius of 6 mm (33 voxels) was calculated between each pair of participants (i, j). (2) The correlation distance of wellbeing score profiles was computed between each pair of participants (i, j). (3) Partial Spearman's rank correlation (ρ) was calculated between participant-by-participant brain and behavioural distance matrices for v1 at t1, controlling for age and gender (not shown). b, The ρ value was calculated for each voxel at each of 5470 time points. c, Independent component (IC) analysis (ICA) was used to decompose the whole-brain timeseries of ρ values into 100 IC brain maps and corresponding time courses. d top, These IC time courses were hierarchically clustered and the resulting dendrogram was cut at the level of eight clades (dotted line). IC maps in each clade were thresholded with an activation probability larger than 0.5 calculated using Gaussian Mixture Model and cluster sizes >50 voxels. Resulting IC maps from the same clade were merged, leaving seven co-activated networks (shown in Figures 3-5). d bottom, Multiple linear regression was also performed on the IC time courses. The latter were treated as the dependent variables to regress against film annotations. Two low-level annotations were also included as controls in the regression model (not shown).*

We hypothesised that some sets of co-active networks representing individual differences in wellbeing would be more associated with terms we selected to correspond to embodied self-related processing. These are the 'autonomic', 'affective', 'emotion', 'interoceptive', and 'self referential' meta-analyses and associated regions (e.g., the insula and anterior cingulate cortex). Another set of networks was hypothesised to be associated with narrative self-related terms, including 'autobiographical', 'language', 'memory retrieval', 'speech perception', and 'speech production' meta-analyses and associated regions (e.g., the precuneus and superior temporal gyrus; see Decoding, Materials and Methods for justification of terms). To corroborate results, we also provide the top 10 correlations for resulting networks and sets of co-active networks with all neuroimaging meta-analyses in the database. Finally, we conducted an exploratory analysis to determine if individual differences in embodied and narrative self-related networks dynamically scale with socio-affective film context by regressing annotations against IC timecourses (Figure 1d, bottom).

Results

We hypothesised that individual differences in wellbeing would be correlated with individual differences in whole brain sets of networks associated with embodied and narrative self-related processing. Wellbeing scores from the wellbeing subdomain of the emotional battery included in the NIH Toolbox were well distributed (Figure 2)(84). SSIP-RSA using these scores followed by ICA and hierarchical clustering (see Figure 1a-d) resulted in eight sets of co-activated networks or clades (where a clade is a distinct group of ICs that share a common point of origin on the dendrogram; Figure 1d and 3a). One of these clades was excluded from further analysis because it did not survive cluster size thresholding. Only ICs corresponding to statistically significant individual differences as determined by a Mantel test and corrected for multiple comparisons are included in clades (all qs < 0.01; cluster sizes > 50 voxels). In what follows, we present clades ordered thematically by the functional properties associated with each clade (Figures 3-5, Table 1).

Figure 2. Wellbeing scores. *Distributions of scores across 20 participants on seven items from the wellbeing subdomain of the NIH toolbox. Items comprising this subdomain are from the Positive Affect, General Life Satisfaction, and Meaning and Purpose questionnaires. The x-axis labels are specific item names from these questionnaires. The y-axis shows T-scores, where a 50 is the mean of the US general*

population and 10 units is the standard deviation. The boxplot in the middle of each violin plot is the maximum, third quartile, median (white dot), first quartile, and minimum T-score. Width of the density plot represents frequency.

Networks Sets 1/2

The co-activated networks in the first two clades are likely functionally related in that they are clustered next to each other (Figure 3a, right). That is, they have similar timeseries suggesting that the information being processed in associated brain regions is thematically related. Supporting this, both clades loaded on different networks that were functionally associated with embodied self-related processing terms. For clade one, these most prominently involved medial frontal cortices anterior to and avoiding the precuneus (Figure 3b, left). It included the anterior and midcingulate and dorsal and ventral medial frontal cortices. Other regions were the right insula and right cerebellum. Collectively, this clade was most correlated with the 'affective', 'autonomic', and 'self referential' neuroimaging meta-analyses (all rs $\geq$ 0.10, Figure 3b, right). The second clade consisted predominantly of the bilateral insula and subcortical regions including the pallidum, putamen, thalamus, and ventral tegmental area (Figure 3c, left). These regions were most correlated with the 'affective', 'autonomic', 'interoceptive', and 'speech production' meta-analyses (all rs $\geq$ 0.11, Figure 3c, right). For both clades, results were supported by a more data driven approach, examining the top 10 neuroimaging meta-analytic associations. That is, both clades had functionally overlapping correlations with the 'mood', 'pain(ful)', and 'reward' term-based meta-analyses (Table 1). Clade one was more associated with, e.g., 'emotional' and 'valence' whereas clade two was more associated with 'reinforcement' and 'reward' overall (Table 1).

Figure 3. Network sets 1/2. *Co-activated networks correlated with individual differences in wellbeing and associated with embodied self-related processing. a, The dendrogram from Figure 1 as coloured by clade. b, Representative brain slices showing all independent components (ICs) from clade 1. c, Representative brain slices showing all ICs from clade 2. Activation on slices in a and b is coloured by z-scores, where yellows represent lower values and red represents higher values (≤ 15.00). Regions*

outlined in black show statistically significant individual differences as determined by a Mantel test (false discovery rate corrected q < 0.01, one-tailed). Numbers below slices are MNI coordinates, where x is L/R (left/right), y is A/P (anterior/posterior), and z is I/S (inferior/superior). Polar plots are the results of hypothesis based correlations between co-activated network maps and term-based meta-analyses. Axis are correlation values with the internal area coloured by clade. Terms in red are related to narrative self-related processing, i.e., autobio (autobiographical), lang (language), mem retriev (memory retrieval), spch percep (speech perception), and spch produc (speech production). Terms in blue are related to the embodied self-related processing, i.e., affective, autonomic, emo (emotion), interocep (interoceptive), and self (self referential). The 'visual' and 'attention' terms in black are included for comparison.

Network Sets 3/4/5

The co-activated networks in clades three through five are likely functionally related in that clades three and five are next to each other and all three clades are near each other on the left side of the dendrogram (Figure 3a). Supporting this, clades, three, four, and five all loaded on regions functionally related to our definition of the narrative self-related processing. Specifically, the third clade consisted of the right Heschl's gyrus, right superior temporal gyrus and left supramarginal gyrus (Figure 4a, left). The clade as a whole was correlated with the 'language', 'speech perception', and 'speech production' meta-analyses (all rs ≥ 0.17, Figure 4a, right). The fourth clade comprised the right ventral precentral gyrus (Figure 4b, left) and correlated with the 'speech production' meta-analysis (r = 0.19; Figure 4b, right). The fifth clade contained left posterior medial precuneus regions (Figure 4c, left). These regions were most correlated with 'attention' and 'memory retrieval' meta-analyses (all rs ≥ 0.10, Figure 4c, right). Again, results are supported by terms derived from the more data driven approach. In addition to those mentioned, these included language related terms like 'articulatory', 'linguistic', 'phonological', 'sentence(s)', 'speech perception', 'production', and 'vocal' and memory related terms like 'autobiographical', 'episodic memory', and 'memory retrieval' (Table 1). Collectively, these clades contained the IC that was the most correlated with individual differences across this entire study, located in the left supramarginal gyrus (ρ = 0.53). Furthermore, clades 3, 4, and 5 were the most predictive of individual differences on average (mean ρ = 0.38 versus 0.29 for clades 1/2 and 0.30 for clades 6/7; Table 1).

a   Co-activated networks, clade 3

x = -52 (L)   x = 51 (R)   y = -45 (P)

b   Co-activated networks, clade 4

x = 52 (R)   y = -11 (P)   z = 32 (S)

c   Co-activated networks, clade 5

x = -3 (L)   x = 8 (R)   x = 12 (R)

Figure 4. Network sets 3/4/5. *Co-activated networks correlated with individual differences in wellbeing and associated with narrative self-related processing. a, Representative brain slices and polar plot from clade 3 (a), clade 4 (b), and clade 5 (c). See Figure 3 for details and abbreviations.*

Network Sets 6/7

The final two clades were functionally related but do not necessarily conform to any a priori hypotheses pertaining to self-related processing. Clade six showed sets of networks that included regions in the occipital lobe like the calcarine sulcus and lingual gyrus (Figure 5a, left) whereas clade seven mainly consists of regions in the dorsolateral prefrontal cortex, bilateral intraparietal sulci, and middle temporal cortices (Figure 5b, left). Table 1 shows that both sets of networks are associated with an array of visual and visual attention related meta-analyses. For these reasons, we included 'visual' and 'attention' in the polar plots in Figures 3-5 for purposes of comparison. Clade six is correlated with the 'visual' (r = 0.20) and 'autobiographical' (r = 0.11) and clade seven with the 'attention' (r = 0.42) and 'visual' (r = 0.29) meta-analyses. Other meta-analytic terms included 'eye fields', 'object(s)', and 'visual stream' for clade six and 'attentional', 'visual motion', and 'visuospatial' for clade seven.

Figure 5. Network sets 6/7. *Co-activated networks correlated with individual differences in wellbeing and associated with visual-attention related processing. Representative brain slices and polar plots from clade 6 (a) and clade 7 (b). See Figure 3 for details and abbreviations.*

Socio-Affective Contexts

Finally, we conducted an exploratory regression analysis to determine whether fluctuations in brain networks that correlated with individual differences are related to socio-affective contexts in '*500 Days of Summer*'. Resulting positive and negative associations would suggest the degree to which networks are influenced by context in the film whereas a lack of association might suggest the relative impermeability to context. To examine this, we regressed valence, arousal, social interaction, and people speaking film annotations and two nuisance regressors generated by Masson and Isik (85) against all IC timecourses (see Figure 1d). There were no statistically significant results after correcting for multiple comparisons at $q < 0.01$ (the threshold we applied throughout). At a less conservative though still corrected threshold, there were 11 components that scaled with socio-affective contexts (all $qs < 0.05$; Figure 6).

Specifically, networks with individual differences that varied with valence in the film are the right calcarine sulcus (IC 5; $\beta = -0.085$, $p < 0.0001$, uncorrected) and anterior midcingulate cortex (IC 31; $\beta = -0.050$, $p = 0.001$, uncorrected; Figure 6a). Networks varying with arousal are the left (IC 21, $\beta = -0.065$, $p < 0.0001$, uncorrected) and right (IC 9; $\beta = -0.062$, $p < 0.0001$, uncorrected) intraparietal sulci and left supramarginal gyrus (IC 8; $\beta = 0.061$, $p = 0.0002$, uncorrected; Figure 6b). Networks varying with the presence of social interactions are the right calcarine sulcus (IC 5; $\beta = 0.063$, $p = 0.0004$, uncorrected), posterior cingulate cortex (IC 52; $\beta = 0.049$, $p = 0.0006$, uncorrected), and midcingulate cortex, subgenual anterior cingulate cortex, and temporoparietal junction (IC 81; $\beta = 0.049$, $p = 0.0012$, uncorrected; Figure 6c). Finally individual difference networks varying with the presence of people speaking are the bilateral intraparietal sulci (IC 35; $\beta = -0.050$, $p = 0.0007$, uncorrected), left lingual gyrus (IC 99; $\beta = 0.046$, $p = 0.0012$, uncorrected), and left supramarginal gyrus (IC8; $\beta = -0.056$, $p = 0.0004$, uncorrected; Figure 6d). Thus, a full 55% of ICs (i.e., 6/11) are from clades 6 and 7 associated with visual attention related processing (see Table 1). Furthermore, 100% of regressors (4/4) were associated with an IC from these visual-attention-related clades.

Figure 6. Socio-affective contexts. *Multiple linear regression of independent component (IC) time courses against the film annotations for valence (a), arousal (b), presence of social interaction (c), and presence of people speaking (d). Shown are thresholded IC maps with time courses that have a statistically significant relationship with film annotation timeseries (false discovery rate corrected q < 0.05, two-tailed permutation, all clusters > 50 voxels). Positive and negative βs are shown in orange and blue, respectively. Numbers are MNI coordinates of slices, where x is L/R (left/right), y is A/P (anterior/posterior), and z is I/S (inferior/superior).*

Discussion

Because it is a multidimensional construct and due to the variability of past neuroimaging results, we proposed that wellbeing is supported by a whole brain distribution of co-active networks, subject to individual differences in weighting. We suggested that these might be broadly understood as being organised around two aspects of self-related processing. Sometimes wellbeing hinges on judgements about physical, interoceptive, and emotional states, i.e., embodied self-related processing. At others, wellbeing is associated with judgments around the effectiveness of the typically language based stories we tell ourselves, i.e., narrative self-related processing. We evaluated this framework by assessing wellbeing and then asking participants to watch a romantic comedy-drama film during fMRI. Using SSIP-RSA combined with ICA and hierarchical clustering (Figure 1), we confirmed hypotheses that individual differences in wellbeing (Figure 2) were correlated with individual differences in sets of co-activated networks, divisible into embodied and narrative self-related processing (Figures 3-4, Table

1). We discovered a third set of visual-attention related co-activated networks that also correlated with individual differences in wellbeing (Figure 5, Table 1). Dynamically varying socio-affective context from the film scaled primarily with individual regions from these visual-attention-related networks (Figure 6).

Embodied Self

We defined embodied self-related processing as encompassing interoceptive awareness, body ownership, and emotional processing. Extensive behavioural and neurobiological research links these processes with mental health and wellbeing (68, 71, 86–93). Consistent with this, we found that some individual differences in wellbeing were correlated with individual differences in brain activity in two closely related sets of co-activated networks that were associated with interception (pain and autonomic processes) and affective/emotional processing (Figure 3; Table 1). Both sets included most of the insula bilaterally, regions consistently associated with interoceptive and emotional awareness (70, 72–75, 94, 95). The first set of co-activated networks additionally contained an array of medial frontal regions anterior to the precuneus, like the anterior cingulate cortex (Figure 3b). These regions are collectively involved in a number of specific functions closely associated with emotional processing, particularly in more anterior portions (96, 97). They also play an important role in self-related processing in general and self-related emotional processing in particular (76, 77, 98, 99). In addition to the insula, the second set of co-activated networks included the pallidum, putamen, thalamus, and ventral tegmental area (Figure 3c). These subcortical regions are primary components of reward processing in the brain (100, 101) and have also been linked to self-related processing (102). The close functional relationship between these two sets of co-activated networks is consistent with recent discussions regarding the interrelationship between emotional and reward processing (103, 104).

Narrative Self

We defined narrative self-related processing as being primarily constructed from language, inner speech, and autobiographical memory and associated brain regions (60). Though research into these relationships is limited, what exists links these processes with mental health and wellbeing (60, 105–115). Consistent with this, we found that individual differences in wellbeing were correlated with three sets of co-activated networks that are associated with language, speech perception and production, and autobiographical memory related processing (Figure 4; Table 1). These narrative self-related networks contained the most predictive single brain region and were collectively the most predictive of individual differences in wellbeing in this study.

Regions in these sets of co-activated networks included the superior temporal, supramarginal, and ventral precentral gyrus that are variously involved in language comprehension, speech perception, and speech production, including inner speech (60, 116–118)(Figure 4a and 4b). They also included medial brain regions. Unlike the anterior distribution in embodied self-related networks, these were posterior, including aspects of the posterior cingulate and precuneus (Figure 4c). Research pertaining to these regions suggest that they are associated with autobiographical memory, self-referential processing, and consciousness (119–121). More generally, they are part of the larger 'default mode network' that has been argued to combine 'episodic memory, language, and semantic memory processes to generate an ongoing internal narrative' (122). Consistent with the various suggestions for anterior to posterior medial divisions, perhaps the more anterior medial regions are more associated with self-related emotional processing (97, 119, 123–128). In contrast, we suggest that these more posterior regions are more associated with language/semantic aspects of self-related narrative processing (60, 129, 130). This is supported by the close functional relationship between all three sets of co-activated networks, suggesting that the superior temporal, supramarginal, ventral precentral, and posterior medial regions all work together (see Figure 3a).

We previously proposed a 'HOLISTIC' model to help account for these links, i.e., how the neurobiology of language and the default mode networks support both consciousness and wellbeing (60). Roughly, the model suggests that our language and inner speech based narratives can become entrenched through learning and use within these language related and default mode networks, making them more automatic and less accessible to conscious awareness. Though these entrenched narratives are typically adaptive, they might not always be appropriate or positive and may become less so over time, leading to decreased wellbeing. Therefore, induction of neuroplasticity in these networks could lead to improved mental health and wellbeing. Indeed, changes to language-based narratives are arguably the primary target and mechanism of psychotherapy and perhaps even meditation (131), leading to plasticity in language and default mode related regions like the precuneus that often best predicts mental health and wellbeing outcomes (60).

Visual-Attention

We framed our hypotheses in terms of embodied and narrative self-related processing but unexpectedly found that individual differences in wellbeing are also correlated with individual differences in co-active networks associated with visual-attention related processing (Figure 5, Table 1). One set of these networks included regions associated more with 'lower level' visual processing like primary visual cortex (Figure 5a). The other set of co-active networks included more 'higher level' visual processing regions

like bilateral middle temporal visual motion processing regions (132, 133). It also included much of the 'dorsal attention network' associated with top-down controlled attentional selection (e.g., the bilateral intraparietal sulcus; Figure 5b)(134). We also found that it was predominantly though not exclusively regions from these two sets of visual-attention related networks that scaled with socio-affective context in the film (Figure 6). However, the latter results should be interpreted with caution as no regions survived our initial q < .01 correction for multiple comparisons (though 11 survived at q < .05).

These results suggest that individual differences in wellbeing might arise when people develop unique but stable attentional strategies for sampling that external world. This is consistent with research showing that mood impacts visual attention (29, 135) and sensory attentional biases are associated with mental health and contribute to wellbeing (136, 137). This warrants expanding the initial framework presented here to include three general neurobiological contributions to individual differences in wellbeing. These are the more internally focused embodied and narrative self-related networks and more externally focused and dynamically determined visual-attention-related networks. Perhaps these individual difference networks are more or less driven by the dominant formats that people experience their thoughts in. For example, experience sampling has been used to identify the 'five frequent phenomena' of inner experience (138) and questionnaire measures have identified 'modes' of thinking (139). From these, individuals more prone to 'sensory awareness'/'feeling'[1], 'inner speaking'/'internal verbalization', or 'inner seeing'/'visual imagers' might weight embodied self-related, narrative self-related, or visual-attention related networks more on average, respectively (138, 139).

Separability

Overall, results associate individual differences in wellbeing with separable embodied and narrative self- and visual-attentional related brain networks. Yet, functional descriptions of these networks do overlap somewhat. Most prominently, excluding the narrative self-related clades, 8.49% of the terms associated with the other clades were auditory and language related (i.e., 27 of 318 total terms, with at least three terms in each clade, Table 1). For example, both the embodied self-related processing clades resulted in anterior insula activity associated with, e.g., the terms 'articulatory' (clade one) and 'phonological' (clade two). The anterior insula is associated with both interoceptive processing and speech production, likely because autonomic/interoceptive processes like respiratory control are necessary for speech production and inner speech (60, 140). Thus, though these descriptions still fit within the general framework outlined

---

[1] This might also include the factor 'representational manipulation' from (139), associated with questions like 'It is easy for me to imagine the sensation of licking a brick'.

here, they nonetheless suggest that clades found in this study are likely less functionally separable than they initially appear.

Indeed, it would be nonsensical to argue that the underlying embodied and narrative self-related and visual-attention-related networks operate independently. Rather, we suggest they are generally interactive and integrative (60, 141). More specifically, the HOLISTIC model introduced above formalises this interactivity and suggests a rationale for the 8.49% prevalence of language related terms in putatively non-language related clades (60). Namely, humans encounter and produce about 150,000 words a day. These constitute a primary tool for categorising our internal and external worlds, allowing us to extend our cognitive abilities by virtually manipulating these experiences with language. Neurobiologically, this can be accomplished because words are linked to interoceptive, emotional, and perceptual experiences in memory. Thus, when we experience emotions or colours, the associated labels are typically activated and vice versa. Because of this interdependence, words can impact wellbeing through entrenchment (as described above) but also through conscious manipulation. For example, labelling and reappraisal are primary tools for emotional regulation and are associated with the embodied, language/narrative, and visual-attention related brain regions described above (142–144). It is perhaps because of such interactivity that the narrative self-related co-active networks are the most correlated with individual differences in wellbeing.

Limitations

A limitation of our initial embodied and narrative self-related framework is our assumption that participants are in fact engaging in self-related processing, associating the film with their own experiences both implicitly and possibly explicitly (e.g., through inner speech). Indeed, it is inconceivable how the film could be understood if they did not (145–148). Nonetheless, because we did not explicitly test this assumption, alternative frameworks are possible. For example, results could be simply described in the more general terms associated with the resulting seven sets of co-active networks, i.e., as 'emotion', 'reward', 'speech perception', 'speech production', 'memory', 'vision', and 'attention'. Alternatively, even more specific networks could be functionally described, down to the 100 sampled. Furthermore, the methodological decisions resulting in seven sets of co-active networks, derived from these 100 networks, were relatively arbitrary and perhaps even hundreds more networks might be described. Though this more agnostic and expansive approach is not wrong, it is problematic in that it lacks any overarching explanatory power for understanding the neurobiology of wellbeing as a whole.

Conclusions

In this study, individual differences in wellbeing are shown to be supported by individual differences in three separable sets of co-activated networks associated with the embodied self, narrative self, and sensory-attentional related processing. These co-activated networks encompass much of the human brain, as might be expected from a multidimensional construct like wellbeing. They can be further subdivided into finer networks with more specific functions that predict individual differences in wellbeing. This holds promise for individualised approaches to medicine. For example, networks particularly associated with negative (and positive) wellbeing might be identified within an individual. These can then become the focus of individual or therapeutic interventions to increase wellbeing (e.g., through psychotherapy). This might take the form of inducing neuroplasticity through pharmacological means and/or manipulating interoception or visual attention more in some individuals or language-related narratives in others.

Material and Methods

This research made use of the 'Naturalistic Neuroimaging Database' (78). The NNDb includes data from 86 participants who completed the NIH Toolbox a few weeks before watching one of 10 previously unseen full-length films during fMRI. Here we analysed data from 20 participants who watched the 2009 romantic comedy-drama film '*500 Days of Summer'* (1 hour and 31 minutes).

Participants

Of the 20 participants, 10 identified as female, with an age range of 19-53 years (M = 27.70, SD = 9.9). All were right-handed, native English speakers, with no history of neurological or psychiatric disorders, taking no medication, with no hearing impairments, and with unimpaired or corrected vision. The study was approved by the University College London Ethics Committee and all participants gave written informed consent.

Wellbeing Measure

Participants completed the majority of the NIH Toolbox on an iPad about 2–4 weeks before fMRI scanning. The tests were administered in a sound shielded testing room and participants were provided with headphones. These included an emotional battery with a wellbeing subdomain that is composed of three components: positive affect, life satisfaction, and meaning and purpose (84). Because not all participants answered the same questions, we included the seven items in the wellbeing subdomain with data from all 20 participants. These were two items from the 'Positive Affect CAT Age 18+ v2.0' (PA010 & PA001), three items from the 'General Life Satisfaction CAT Age 18+ v2.0' (PA049m, PA045m, & PA047), and two items from the 'Meaning and Purpose CAT Age 18+ v2.0' (PA075m & PA051). Raw scores were converted to standard T-scores where 50 represents the mean of the US general population (based on the 2010 Census) and 10 units represent one standard deviation. Scores from the seven items were well distributed, permitting further analysis (Figure 2).

f/MRI Preprocessing

MRI data were obtained on a 1.5T Siemens MAGNETOM Avanto with a 32-channel head coil. Anatomical images were acquired using a MPRAGE sequence (TR = 2730 ms, TE = 3.57 ms, 176 sagittal slices, voxel size = $1 \times 1 \times 1$ mm^3). Functional images were acquired using a multiband echo planar imaging (EPI) sequence (TR = 1000 ms, TE = 54.8 ms, flip angle 75°, 40 interleaved slices, voxel size = $3.2 \times 3.2 \times 3.2$ mm^3, multiband factor = 4, no in-plane acceleration). The film '*500 Days of Summer*' had 5470 whole brain volumes/TRs.

Unless otherwise noted, all preprocessing and data visualisation was done with the AFNI software package and specific programs are indicated where appropriate (149)[2]. First, anatomical images were corrected for intensity non-uniformity, deskulled, and nonlinearly aligned to an MNI template (i.e., MNI152_2009_template_SSW.nii.gz; using '@SSwarper'). Next, Freesurfer's 'recon-all' was run with default parameters (version 7.0)(150, 151)[3]. This allowed us to segment anatomical images into ventricle and white matter regions of interest to be used to make nuisance regressors (see next paragraph) and a group grey matter mask. The latter was created by resampling the anatomical images to functional resolution ('3dresample'). From these, we removed the Freesurfer derived ventricle and white matter regions of interest. We then took the union of these from all participants to create a group grey matter mask (using '3dMean').

Preprocessing of the functional data was done using 'afni_proc.py'[4] and timeseries are available as derivatives on OpenNeuro[5]. Blocks were included for despiking ('despike'), slice-timing correction ('tshift'), volume registration to the run with the least number of outliers and warping to MNI space (in one step, 'align', 'tlrc', and 'volreg'), masking ('mask'), scaling to a mean of 100 ('scale'), and regression ('regress'). The latter used '3dDeconvolve' to remove demeaned motion and motion derivatives, a polynomial degree of two, bandpass filtering (0.01 to 1), ventricular activity (first three principle components), and white matter activity from the timeseries on a per run basis in one step. Finally, as additional protection, hand labelled ICA based artefacts were also removed from the resulting timeseries (for details, see)(78). Note that we did not smooth this data because the searchlight analysis uses a voxel radius of six mm.

SSIP-RSA

We examined individual differences using searchlight spatial inter-participant representational similarity analysis (SSIP-RSA)(79–81). This approach measures how matched inter-participant variability in fMRI activity patterns and wellbeing score profiles are in searchlights at each time point in the film (Figure 1a). Specifically, we quantified inter-participant variability by computing a participant-by-participant distance matrix for both brain patterns (Figure 1a, 1) and wellbeing scores (Figure 1a, 2). Each square of the matrix is the correlation distance of the brain pattern vector or item-level wellbeing score vector between a pair of participants (i, j). The equation for the correlation distance is:

---

[2] https://afni.nimh.nih.gov/
[3] http://www.freesurfer.net/
[4] https://afni.nimh.nih.gov/pub/dist/doc/program_help/afni_proc.py.html
[5] https://openneuro.org/datasets/ds002837/versions/2.0.0

$$1 - \frac{(i - \bar{i}) \cdot (j - \bar{j})}{||\, (i - \bar{i}) \,||_2 \, ||\, (j - \bar{j}) \,||_2}$$

Here the brain distance matrix was computed time point by time point, using spatial patterns in searchlight spheres centred at each voxel with a radius of six mm (or 33 voxels). Unlike typical IP-RSA which uses temporal inter-participant correlation to measure brain synchrony of the whole timeseries, spatial inter-participant analysis allows us to calculate inter-participant variabilities dynamically at all 5470 time points in the film. This provides a foundation for subsequent discovery of networks and co-active networks as well as the possibility to understand how film annotations are related to individual differences.

As controls, age and gender differences between participants were also considered in the SSIP-RSA. Absolute difference between ages of two participants was calculated to compute the participant-by-participant age distance matrix. In the participant-by-participant gender distance matrix, same gender is coded as 1 and different genders as 0. We then calculated the partial Spearman's rank correlation (ρ; controlling for age and gender) between the upper triangles of the brain and wellbeing distance matrices (Figure 1a, 3), and wrote the ρ value back to the voxel in the centre of the searchlight. A significantly positive ρ value means that participants with more similar brain activity patterns also had more similar wellbeing score profiles. Therefore, the ρ value can be a proxy to show the extent a voxel is related to individual differences in wellbeing at a specific time point. The code used to conduct the SSIP-RSA and all subsequent analysis is available for inspection on GitHub[6].

ICA

To identify networks and reduce dimensionality, we performed independent component (IC) analysis on the SSIP-RSA results with 100 dimensions using '*melodic*' from the FSL software package (version 5.0)(152)[7]. By enforcing spatial independence between components, the whole-brian ρ timeseries (Figure 1b) was decomposed into ICs with a spatial brain map and its corresponding timecourse (Figure 1c). The spatial map contains loadings of each voxel for the IC and the time course indicates contributions of that IC at each time point. The original whole-brian ρ timeseries is a linear combination of all IC maps and their associated time courses. We arbitrarily chose 100 components (i.e., dimensions) because the automatic estimation method in melodic did not finish in a reasonable amount of time. To address this

---

[6] https://github.com/YumengMa01/WellbeingNetworks
[7] https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/MELODIC

potential limitation, we also conducted the IC analysis with 50 and 150 dimensions and this resulted in similar spatial networks. All 100 ICs are available for inspection in NeuroVault (153)[8].

Hierarchical Clustering

To identify co-active spatially independent networks and further reduce dimensionality, we used hierarchical clustering with the Ward method to cluster ICs with similar time courses together (using the SciPy clustering package with default parameters; Figure 1d)(154). We made the relatively arbitrary decision to cut the dendrogram at the level of eight clades as this visually appears to parcellate the dendrogram by height into more differentiated clades (see Figure 1d and 3a). Furthermore, cutting higher up would have merged a large number of ICs in some instances, e.g., clades 1 and 2 would be merged, involving 51 networks (see Figure 3a). This might have been difficult to interpret as it would involve much of the brain. Cutting lower down would require potentially interpreting a much larger number of clades, reducing explainability in any one theoretical framework like the self-related framework we propose. Thus, cutting at eight allowed a middle ground that permitted interpretability while permitting us to discover alternative hypotheses. All clades are available for inspection in NeuroVault (153)[9].

Mantel Test

To test if brain regions shown in each of the ICs are statistically significantly associated with individual differences in wellbeing (i.e., the ρ values are significantly high), we conducted a Mantel test (155). This is a permutation test used to measure the significance of correlation between two distance matrices. With the wellbeing, age, and gender distance matrices remaining unchanged, rows and columns of the brain distance matrix were shuffled in the same random order. The partial Spearman's rank correlation (ρ) between the shuffled brain and wellbeing distance matrices was recalculated. After repeating this process 10,000 times, a null distribution of the ρ values was generated and a p-value calculated using a one-tailed test.

Instead of a searchlight analysis, the brain distance matrix was computed using the BOLD signal pattern extracted from voxels with significant contributions to each IC at the peak time point in the corresponding IC time course. In order to select voxels significantly contributing to each IC, an appropriate threshold needed to be chosen. The output IC brain maps from FSL 'melodic' contain z-scores. These are calculated by subtracting the mean of loadings in all voxels from each voxel's loading, and then dividing the result by the standard deviation of loadings in all voxels, so that the brain maps have zero mean and unit

---

[8] https://neurovault.org/collections/15197/
[9] https://neurovault.org/collections/15197/

variance. However, the mean and standard deviation from whole-brain voxels are not the same as those for an underlying null distribution, so the z-scores have no statistical meaning. Therefore, traditional p-value based thresholding cannot be used. Because different z-score thresholds can generate brain regions with different cluster sizes and we do not know which captures individual differences in wellbeing the best, here we used multiple thresholds following the method used by Poppe et al. (156), and did Mantel tests on the threshold that generated the highest SSIP-RSA ρ value for activity at the peak time point in the IC's time course under evaluation.

Specifically, after dividing the loading of each voxel by the maximum loading value in the map, we created brain masks including voxels surviving nine different thresholds separately, based on the normalised z-scores (normalised z larger than 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, and 0.9) and cluster sizes larger than 50 voxels to remove ICs containing small clusters of voxels. This resulted in nine masks for each IC, each containing a different number of voxels. We then re-did the SSIP-RSA on brain patterns inside each of the nine masks at the peak time point and selected the threshold that generated the highest ρ value. We then did the Mantel test, generating a p-value for each IC, correcting for the number of ICs tested using false discovery rate with a threshold of q < 0.01. All Mantel test results are available for inspection in NeuroVault (153)[10].

Note that we did hierarchical clustering on all ICs from the ICA. However, we could have done this step after the Mantel tests were performed. To determine if this affects the results, we examined the clades after hierarchical clustering on just the IC time courses associated with the Mantel results. In terms of interpretation (see Decoding), the results were not substantially different. The primary difference is that clades four and five merged (see Network Sets 3/4/5). If anything, this bolsters the anatomical assertions of the language based narrative framework, suggesting a strong relationship between language related and precuneus regions made throughout.

Artefact Identification

Following the ICA, hierarchical clustering, and the Mantel test, we manually examined all results to identify artefacts. ICs 1, 2, 3, 4, 6, and 83 had no activity after limiting to positive values and applying a minimum cluster size threshold of 50 voxels. ICs 1, 2, 3, 4, and 6 were all in a single clade and that clade was therefore dropped from further analyses. ICA readily picks up artefacts and we hand labelled all components that might be artifactual (157). This identified nine likely artifactual components. These were IC 27 (with activity appearing as confetti throughout the brain, activity crossing multiple tissue

---

boundaries and types, and activity in the white matter), 33 (crossing multiple tissue boundaries, white matter), 45 (crossing multiple tissue boundaries, peaks not located in brain tissue, white matter), 48 (peaks not located in brain tissue), 73 (peaks not located in brain tissue), 82 (peaks not located in brain tissue), 84 (confetti, crossing multiple tissue boundaries, peaks not located in brain tissue, white matter), 94 (peaks not located in brain tissue, white matter), and 95 (peaks not located in brain tissue). All nine of these were excluded from further analyses.

We also examined all Mantel test results and found one outcome in which activity appeared as whole brain confetti (encompassing 9222 voxels), with activity crossing multiple tissue boundaries, and with activity in white matter. Reexamining the original IC suggests it might have been an artefact though it was not labelled as such initially because it was a borderline case. As such, IC 92 was also excluded from further analyses. Note that inclusion of all hand labelled artefacts would not significantly change interpretation of the results as determined by reverse inference (see Decoding, next). All artefact ICs and Mantel test results are available for inspection in NeuroVault (153)[11].

Decoding

To help specify the functions of individual networks and collections of co-active networks associated with individual differences in wellbeing obtained in the prior steps, we performed reverse inference using both NiMARE[12] and Neurosynth[13] 'decoding' (83, 158). Specifically, 10 a priori term-based meta-analyses relevant to our embodied and narrative self-related processing hypotheses were collected using the NiMARE Python package (versions 0.0.12)(158). These derive from the Neurosynth dataset (version 0.7, released July, 2018)(83), constructed from 507,891 voxel activation coordinates reported in 14,371 neuroimaging studies where terms appear at high frequency in the titles and abstracts (compared to studies where those terms do not appear). Next, the Pearson correlation was calculated between these 10 meta-analyses and the vectorized co-active networks or clade maps obtained from the hierarchical clustering analysis, using only ICs significant by the Mantel test not deemed artifactual. For the embodied self-related processing, term-based meta-analyses were 'affective', 'autonomic', 'emotion', 'interoceptive', and 'self referential'. For narrative self-related processing, correlated term-based meta-analyses were 'autobiographical', 'language', 'memory retrieval', 'speech perception', and 'speech production'. These are presented in polar plots where the axes are r-values (Figures 3-5) and only correlations $\geq$ 0.10 are interpreted in the text. In addition to these a priori chosen sets, we also post hoc

---

[11] https://neurovault.org/collections/15197/
[12] https://nimare.readthedocs.io/en/stable/index.html
[13] https://neurosynth.org/

included the correlations with the 'visual' and 'attention' term-based meta-analyses in the Figure 2 polar-plots.

The choice of some of these terms are obvious while others need further justification. The term 'self referential' might be considered to be relevant to both embodied and narrative self-related processing. However, none of the 166 studies associated with the 'self referential' meta-analysis uses the term 'narrative'. More generally, the term 'narrative' was not contained in the titles or abstracts of any papers with a high enough frequency to be included in the neurosynth database. For this reason, we included 'self referential' as an embodied term. Next, 'autobiographical' and 'memory retrieval' might also be considered relevant to both the embodied and narrative self-related processing. However, we defined the embodied self as focused in the present moment whereas the narrative self supports thinking about the past and future (via mental time travel; see Introduction). Furthermore, we have proposed a neurobiological model relating networks associated with language and autobiographical memory to narrative processing (60). For these reasons, we included 'autobiographical' and 'memory retrieval' as narrative self-related terms.

To lend support to our a priori choice of terms, we also conducted data-driven reverse inference, making use of all 1,334 term-based meta-analyses available on the web-based version of Neurosynth (which also uses dataset version 0.7)(83) Specifically, we provide the top 10 functional associations for the peak voxel in each IC network for any term with a z-value greater than zero. We also used the web-based version of Neurosynth[14] and Neurovault[15] to 'decode' the top 10 meta-analyses associated with each of the seven sets of co-active networks or clades. In both cases, reported terms are only those related to cognitive and behavioural function, excluding anatomical and clearly methodological terms. Terms like the 'central sulcus', 'precentral gyrus/sulcus' and 'postcentral gyrus' were considered anatomical as were their functional equivalents like 'motor', 'premotor', 'somatosensory' as they are often used interchangeably in literature. Note that discrepancies between results from the a priori set of terms and these more data-driven analyses are accounted for by differences in using the NiMARE and the web-based neurosynth packages.

Regression

We conducted exploratory analysis to determine if socio-affective contexts in the film participants viewed scaled with co-activated networks determined to be correlated with individual differences in wellbeing

---

[14] https://neurosynth.org/decode/
[15] https://neurovault.org/images/add_for_neurosynth

(Figure 1d). This was accomplished by regressing the IC time courses against six film annotations from Masson and Isik (85). These were valence, arousal, presence of social interactions (social), presence of a person speaking (speaking), pixel brightness, and auditory amplitude. They were generated for each three second film segment after excluding the opening and ending credits. Specifically, valence and arousal annotations were derived by four raters based on a 1-5 Likert scale. As the ratings were moderately consistent between raters (valence Spearman $\rho = 0.62$, arousal $\rho = 0.35$), the mean ratings are used as objective features of the film. The social annotation was generated by two raters based on whether there was actions or communication happening between two or more characters (presence of social interaction = 1, absence = 0). The inter-rater consistency was high with a correlation value $r = 0.86$. The speaking annotation was rated by the same two raters (presence of person speaking = 1, absence = 0). Finally, pixel brightness and auditory amplitude were generated and averaged over each three second segment and are considered 'low-level' controls (85).

Before regression, the film annotation timeseries were normalised and upsampled from three to one second to match the temporal resolution of IC time courses. The IC time courses were shifted by four seconds to account for haemodynamic delay and rise to peak. They were then treated as dependent variables in the multiple linear regression model, while independent variables were the four annotations and two controls (i.e., pixel brightness and auditory amplitude). Considering the autocorrelations in both the IC time courses and the timeseries of film annotations, we used the non-parametric permutation to test significance. The normalised timeseries of the annotation to be tested was randomly shuffled, upsampled, and then put into the multiple linear regression model while the timeseries of other annotations remained unchanged. A null distribution of the β values for the annotation tested can then be obtained after shuffling its timeseries 10,000 times. A two-tailed permutation test was performed to obtain p-values. Because 0.01 was too conservative to yield significant results, we used an alpha value of 0.05 after correcting for the number of ICs and annotations tested using a false discovery rate algorithm.

Acknowledgments

We would like to thank the Birkbeck-UCL Centre for Neuroimaging (BUCNI) for supporting the NNDb. We also thank Annette Glotfelty and Daniel Lametti for helpful feedback on the manuscript and The LAB Lab and UNITy Project for discussion. JIS would also like to thank Boo Boo for everything, Daniel from St Albans for helping me with my craft, and the Barone-Shaw family and the sanctuary of 'Two Pines'.

Author Contributions

YM and JIS conceived of the study and wrote the manuscript. JIS did all fMRI preprocessing. YM did all of the analyses and made all figures.

Conflicting Interests

The authors declare that they have no conflicts of interest.

Open Science Statement

This study was not preregistered. A version of this paper is available as a preprint on bioRxiv. The preprocessed data used in analyses is publically available on OpenNeuro (https://openneuro.org/datasets/ds002837/versions/2.0.0). Code used to generate results are available on GitHub (https://github.com/YumengMa01/WellbeingNetworks). Results are available in Neurovault (https://neurovault.org/collections/15197/).

Table

Table 1. Network Information. Each row is the peak z-score voxel from statistically significant false discovery rate (FDR) corrected wellbeing-modulated independent component (IC) networks, as determined by a Mantel test (all qs < 0.01). Rows are colour coded by co-activated networks or clades (see Figure 1 and 3-5). The number of voxels is for the whole IC network (which might consist of multiple other regions). 'Decoding' provides the top 10 term based meta-analysis associations at both the peak Montreal Neurological Institute (MNI) coordinate and whole clade for only cognitive/behavioural related terms (excluding, e.g., anatomical terms). Other abbreviations: A/P = anterior/posterior; I/S = inferior/superior; L/R = left/right; un/corr = un/corrected.

Clade	IC	Region	Voxels	Peak MNI Coordinates			Decoding (N=10)		ρ	Peak time	Z thresh	Uncorr p	FDR corr q
				x (L/R)	y (A/P)	z (I/S)	Coordinate (z > 0)	Clade					
1	29	Perigenual Anterior Cingulate Cortex	241	-2	28	29	Pain, painful, gain, noxious, monetary, mood, nociceptive	Pain, painful, conflict, referential, self referential, ptsd, rehearsal, coordination, motor performance, default	0.2793	5451	0.4	0.0003	0.0019
1	31	Anterior Midcingulate Cortex	100	2	3	44	Pain, finger, movement, autonomic, movements, sensations, execution, tapping, index finger, target detection		0.2920	612	0.6	0.0004	0.0021
1	32	Ventromedial Prefrontal Cortex	90	-2	45	-3	Default mode, default, mood, reward, valence, smoking, money, gambling, monetary, value		0.3107	2719	0.6	0.0022	0.0054
1	43	Right Cerebellum	184	23	-75	-24	Interpersonal		0.2991	2241	0.3	0.0024	0.0056
1	50	Right Cerebellum	86	26	-74	-45	Verbal working		0.3921	3411	0.6	0.0001	0.0009
1	51	Left Cerebellum	74	-35	-68	-49	Cognitive functions, suppressed, working memory, working		0.2675	2788	0.7	0.0004	0.0021
1	54	Posterior Middle Cingulate Cortex	309	-2	-26	49	Foot, confidence, coordination		0.2557	4519	0.3	0.0010	0.0036
1	60	Right Posterior Insula	143	44	-11	4	Pain, auditory, music, discriminative, painful, musical, nociceptive, listening, audiovisual, noxious		0.3845	2445	0.5	0.0001	0.0009
1	64	Dorsal Medial Frontal Cortex	270	2	-5	63	Movements, movement, motor imagery, hand, sensorimotor, motor task, execution, finger, finger movements, imagery		0.2334	2598	0.4	0.0032	0.0066
1	68	Left Cerebellum	273	-20	-77	-46	Stimulation, pain, electrical, sensory, navigation, sensorimotor		0.2934	4314	0.3	0.0009	0.0036
1	78	Right Cerebellum	460	35	-64	-48	Loop		0.2326	2782	0.2	0.0018	0.0046
1	80	Left Superior Frontal Sulcus	320	-26	27	46	Default, default network, autobiographical, default mode, real world, conceptual, episodic, autobiographical memory, memory, episodic memory		0.2167	2479	0.3	0.0044	0.0082
1	86	Dorsal Medial Frontal Cortex	146	2	43	43	Inference, mind tom, arousal, emotional, social, tom		0.2574	5259	0.5	0.0001	0.0009

				x	y	z							
1	87	Right Anterior Insula	827	44	-1	-9	Pain, intensity, aversive, taste, painful, articulatory, multisensory		0.2148	263	0.2	0.0027	0.0060
1	88	Perigenual Anterior Cingulate Cortex	61	8	47	14	Default, beliefs, default mode		0.3378	1000	0.7	0.0001	0.0009
2	42	Right Putamen	1374	23	-8	1	Finger movements, finger tapping, movements, finger, movement, tapping, sensorimotor, handed, sexual, execution	Gain, pain, reward, anticipation, painful, stop, monetary reward, monetary, sexual, mood	0.2700	3852	0.1	0.0012	0.0036
2	55	Thalamus	654	2	-12	14	Monetary reward, monetary, anticipation, retrieved, video clips, self reported		0.3812	391	0.1	0.0011	0.0036
2	57	Left Posterior Insula	346	-44	-7	-6	Experiences, pain, unfamiliar, acoustic		0.2478	51	0.3	0.0001	0.0009
2	63	Ventral Tegmental Area	274	-2	-17	-6	Reward, motivational, rewards, reinforcement learning, behaviours, reinforcement, motivation, addiction, choices, monetary		0.3229	1527	0.3	0.0003	0.0019
2	67	Left Pallidum	97	-17	-1	-2	Movement, movements, execution, coordination, tapping		0.2518	2198	0.6	0.0056	0.0098
2	72	Right Anterior Insula	1061	38	23	3	Task, gain, calculation, mood, pain, choose, german, demands, tasks, orthographic		0.3142	4697	0.2	0.0004	0.0021
2	96	Left Anterior Insula	359	-35	23	3	Gain, phonological, language, orthographic, competing, words, heard, demands, pain		0.2432	1966	0.3	0.0025	0.0057
3	7	Right Heschl's gyrus	96	59	-11	11	Auditory, pitch, sounds, listening, touch, listened, noise, tactile, production, sound	Auditory, listening, speech, sounds, acoustic, sound, pitch, listened, speech perception, audiovisual	0.3709	5111	0.3	0.0014	0.0039
3	8	Left Supramarginal Gyrus	418	-50	-46	25	Mind tom, tom, phonological, social interaction, reading, speech production, language, listening, linguistic, theory mind		0.5245	1203	0.1	0.0001	0.0009
3	12	Right Superior Temporal Gyrus	176	53	-46	16	Theory mind, tom, mental states, mind tom, mind, motion, theory, watching, empathy, empathic		0.3690	2502	0.4	0.0012	0.0036
3	14	Right Superior Temporal Gyrus	281	53	-23	-3	Listening, speech, auditory, language, sentences, comprehension, voice, linguistic, sentence, speech perception		0.3508	1223	0.5	0.0011	0.0036
4	18	Right Precentral Gyrus	459	47	-16	40	Sensorimotor, finger, speech production, vocal, production	Speech production, production, oral, sensorimotor, vocal, naming, articulatory, movement, movements, finger	0.3753	9	0.2	0.0030	0.0063
5	15	Left Precuneus	241	2	-63	54	Spatial, calculation, judgement task, attention network, navigation	Locations, navigation, orienting, spatial attention, default, default mode, spatial, virtual, visual attention, location	0.3413	836	0.3	0.0020	0.0050
5	40	Left Precuneus	619	14	-65	31	Retrieval, episodic, episodic memory, memory, recognition memory, memory retrieval, amnestic, items, autobiographical		0.3578	1641	0.2	0.0011	0.0036

6	22	Right Lingual Gyrus	475	20	-60	2	Implicit	Place, navigation, objects, visual, episodic, encoding, visual stream, episodic memory, object, autobiographical	0.2430	3965	0.2	0.0029	0.0063
6	58	Right Calcarine Sulcus	55	14	-77	14	Sighted, visual, eye, eye fields, frontal eye, ageing, saccades, lexical		0.3463	1572	0.6	0.0007	0.0031
6	59	Right Cerebellum	179	32	-53	-26	Motor performance, deficient, sensorimotor, motor imagery, execution, movement, movements, hand		0.2318	5230	0.5	0.0035	0.0068
6	62	Left Cerebellum	313	-23	-69	-20	Movements, movement, production, sensorimotor, finger tapping, pitch, tapping, rhythm, musicians, execution		0.2367	2653	0.3	0.0034	0.0068
6	76	Left Parahippocampal Gyrus	229	-29	-41	-11	Episodic, navigation, episodic memory, autobiographical, memory, retrieval, encoding, objects, place, construction		0.3027	1975	0.5	0.0035	0.0068
6	85	Posterior Cingulate Cortex	1279	-2	-52	15	Default, default mode, autobiographical, episodic, memory retrieval, semantic memory, recollection, retrieval, default network, dmn		0.2723	3192	0.1	0.0040	0.0076
6	99	Left Lingual Gyrus	287	-11	-54	-5	Picture, navigation, words, semantic, sensorimotor		0.2383	1838	0.3	0.0014	0.0039
7	9	Right Intraparietal Sulcus	314	35	-74	30	Spatial, object, eye fields, virtual, objects, navigation, tasks, frontal eye, visuospatial, rotation	Tasks, working memory, working, task, visuospatial, attention, calculation, visual, spatial, attentional	0.3618	2219	0.3	0.0053	0.0095
7	10	Left Intraparietal Sulcus	55	-32	-77	33	Retrieval, memory, encoded, navigation, retrieved, episodic, object, episodic memory, ageing, visual		0.3372	3422	0.7	0.0001	0.0009
7	16	Left Premotor Cortex	270	-47	-3	41	Working memory, working, speech production, saccades, phonological, production, eye, sensorimotor, rehearsal, language		0.5137	4027	0.2	0.0001	0.0009
7	19	Right Dorsolateral Prefrontal Cortex	127	44	12	32	Mood, task, interference, demands, cognitive control, tasks, stroop, response time, stroop task, ptsd		0.3320	4498	0.7	0.0012	0.0036
7	21	Left Intraparietal Sulcus	281	-35	-56	48	Memory, working memory, working, retrieval, recognition memory, word, calculation, memory retrieval, words, retrieved		0.2815	3645	0.4	0.0023	0.0055
7	26	Bilateral Middle Temporal Visual Motion Area	307	-41	-76	-3	Visual, objects, visual motion, motion, face recognition, object, face, negative neutral, orthographical, stream		0.3630	2096	0.2	0.0015	0.0041
7	28	Right Intraparietal Sulcus	622	38	-53	45	Tasks, working memory, working, task, memory, load, attentional, calculation, maintenance, demand		0.2677	1987	0.2	0.0048	0.0088
7	35	Right Intraparietal Sulcus	594	44	-32	49	Finger, tactile, planning, movements, hand, spatial, finger movements, tasks, sensorimotor, sequential		0.3522	879	0.3	0.0001	0.0009
7	44	Right Intraparietal Sulcus	153	23	-60	54	Eye movements, eye, visual, eye fields, attention network,		0.2708	2384	0.4	0.0008	0.0034

						spatial, movements, objects, attention, visually						
7	47	Right Frontal Eye Field	55	41	2	47	Eye fields, frontal eye, eye, spatial attention, eye movements, action, eye field, orienting, saccades, tasks		0.2371	481	0.7	0.0005	0.0023
7	66	Left Dorsolateral Prefrontal Gyrus	501	-50	10	25	Phonological, language, semantic, orthographic, words, languages, word, linguistic, reading, tasks		0.3355	801	0.2	0.0002	0.0015
7	70	Middle Temporal Visual Motion Area	235	44	-64	11	Motion, video clips, video, action observation, hands, actions, body, eye, eye movements, visual		0.2994	3546	0.5	0.0002	0.0015
7	90	Midcingulate Cortex	1008	-11	-41	48	Self referential, referential, video clips		0.2271	519	0.2	0.0005	0.0023

References

1. T. J. VanderWeele, *et al.*, Current recommendations on the selection of measures for well-being. *Prev. Med.* **133**, 106004 (2020).

2. J. B. Koslouski, *et al.*, Measuring emotional well-being through subjective report: a scoping review of reviews. *BMJ Open* **12**, e062120 (2022).

3. E. Diener, "Introduction—The Science of Well-Being: Reviews and Theoretical Articles by Ed Diener" in *The Science of Well-Being: The Collected Works of Ed Diener*, E. Diener, Ed. (Springer Netherlands, 2009), pp. 1–10.

4. E. Diener, S. Oishi, L. Tay, Advances in subjective well-being research. *Nat Hum Behav* **2**, 253–260 (2018).

5. K. V. Das, *et al.*, Understanding subjective well-being: perspectives from psychology and public health. *Public Health Rev.* **41**, 25 (2020).

6. C. D. Ryff, Eudaimonic well-being, inequality, and health: Recent findings and future directions. *Int Rev Econ* **64**, 159–178 (2017).

7. R. M. Ryan, E. L. Deci, On happiness and human potentials: a review of research on hedonic and eudaimonic well-being. *Annu. Rev. Psychol.* **52**, 141–166 (2001).

8. C. D. Ryff, Psychological well-being revisited: advances in the science and practice of eudaimonia. *Psychother. Psychosom.* **83**, 10–28 (2014).

9. A. S. Waterman, Two conceptions of happiness: Contrasts of personal expressiveness (eudaimonia) and hedonic enjoyment. *J. Pers. Soc. Psychol.* **64**, 678–691 (1993).

10. C. D. Ryff, B. H. Singer, Know thyself and become what you are: A eudaimonic approach to psychological well-being. *J. Happiness Stud.* **9**, 13–39 (2008).

11. M.-J. Linton, P. Dieppe, A. Medina-Lara, Review of 99 self-report measures for assessing well-being in adults: exploring dimensions of well-being and developments over time. *BMJ Open* **6**, e010641 (2016).

12. J. Lindert, P. A. Bain, L. D. Kubzansky, C. Stein, Well-being measurement and the WHO health policy Health 2010: systematic review of measurement scales. *Eur. J. Public Health* **25**, 731–740 (2015).

13. F. Martela, K. M. Sheldon, Clarifying the Concept of Well-Being: Psychological Need Satisfaction as the Common Core Connecting Eudaimonic and Subjective Well-Being. *Rev. Gen. Psychol.* **23**, 458–474 (2019).

14. P. J. Cooke, T. P. Melchert, K. Connor, Measuring Well-Being: A Review of Instruments. *Couns. Psychol.* **44**, 730–757 (2016).

15. J. K. Boehm, L. D. Kubzansky, The heart's content: the association between positive psychological well-being and cardiovascular health. *Psychol. Bull.* **138**, 655–691 (2012).

16. S. D. Pressman, S. Cohen, Does positive affect influence health? *Psychol. Bull.* **131**, 925–971 (2005).

17. L. D. Kubzansky, *et al.*, Positive Psychological Well-Being and Cardiovascular Disease: JACC Health Promotion Series. *J. Am. Coll. Cardiol.* **72**, 1382–1396 (2018).

18. N. Martín-María, *et al.*, The Impact of Subjective Well-being on Mortality: A Meta-Analysis of Longitudinal Studies in the General Population. *Psychosom. Med.* **79**, 565–575 (2017).

19. A. Steptoe, Happiness and Health. *Annu. Rev. Public Health* **40**, 339–359 (2019).

20. C. L. M. Keyes, Promoting and protecting mental health as flourishing: a complementary strategy for improving national mental health. *Am. Psychol.* **62**, 95–108 (2007).

21. S. Chan, S. Weich, Mental well-being and recovery in serious mental illness: associations between mental well-being and functional status in the Health Survey for England 2014. *BJPsych Open* **6**, e66 (2020).

22. O. I. Gutiérrez, J. D. Polo, M. J. Zambrano, D. C. Molina, Meta-analysis and Scientific Mapping of Well-being and Job Performance. *Span. J. Psychol.* **23**, e43 (2020).

23. A. J. Oswald, E. Proto, D. Sgroi, Happiness and Productivity. *J. Labor Econ.* **33**, 789–822 (2015).

24. S. Lyubomirsky, L. King, E. Diener, The benefits of frequent positive affect: does happiness lead to success? *Psychol. Bull.* **131**, 803–855 (2005).

25. K. Otake, S. Shimai, J. Tanaka-Matsumi, K. Otsui, B. L. Fredrickson, HAPPY PEOPLE BECOME HAPPIER THROUGH KINDNESS: A COUNTING KINDNESSES INTERVENTION. *J. Happiness Stud.* **7**, 361–375 (2006).

26. J. A. Holman, B. W. Y. Hornsby, F. H. Bess, G. Naylor, Can listening-related fatigue influence well-being? Examining associations between hearing loss, fatigue, activity levels and well-being. *Int. J. Audiol.* **60**, 47–59 (2021).

27. C. Ventura-Bort, J. Wendt, M. Weymar, The Role of Interoceptive Sensibility and Emotional Conceptualization for the Experience of Emotions. *Front. Psychol.* **12**, 712418 (2021).

28. E. M. Benau, Self-reported interoceptive accuracy and interoceptive attention differentially correspond to measures of visual attention and self-regard. *PeerJ* **11**, e15348 (2023).

29. H. A. Wadlinger, D. M. Isaacowitz, Positive mood broadens visual attention to positive stimuli. *Motiv. Emot.* **30**, 87–99 (2006).

30. E. J. Austin, D. H. Saklofske, V. Egan, Personality, well-being and health correlates of trait emotional intelligence. *Pers. Individ. Dif.* **38**, 547–558 (2005).

31. M. Zeidner, G. Matthews, R. D. Roberts, The Emotional Intelligence, Health, and Well-Being Nexus: What Have We Learned and What Have We Missed? *Appl. Psychol. Health Well Being* **4**, 1–30 (2012).

32. K. M. Livingstone, S. Srivastava, Up-regulating positive emotions in everyday life: Strategies,

individual differences, and associations with positive emotion and well-being. *J. Res. Pers.* **46**, 504–516 (2012).

33. K. Gasper, K. D. Bramesfeld, Should I follow my feelings? How individual differences in following feelings influence affective well-being, experience, and responsiveness. *J. Res. Pers.* **40**, 986–1014 (2006).

34. J. J. Gross, O. P. John, Individual differences in two emotion regulation processes: implications for affect, relationships, and well-being. *J. Pers. Soc. Psychol.* **85**, 348–362 (2003).

35. S. Balzarotti, F. Biassoni, D. Villani, A. Prunas, P. Velotti, Individual Differences in Cognitive Emotion Regulation: Implications for Subjective and Psychological Well-Being. *J. Happiness Stud.* **17**, 125–143 (2016).

36. A. Roemer, A. Sutton, C. Grimm, O. N. Medvedev, Differential Contribution of the Five Facets of Mindfulness to Well-being and Psychological Distress. *Mindfulness* **12**, 693–700 (2021).

37. A. J. Bergin, K. I. Pakenham, The Stress-Buffering Role of Mindfulness in the Relationship Between Perceived Stress and Psychological Adjustment. *Mindfulness* **7**, 928–939 (2016).

38. D. B. Newman, J. B. Nezlek, Private self-consciousness in daily life: Relationships between rumination and reflection and well-being, and meaning in daily life. *Pers. Individ. Dif.* **136**, 184–189 (2019).

39. M. G. Adler, N. S. Fagley, Appreciation: individual differences in finding value and meaning as a unique predictor of subjective well-being. *J. Pers.* **73**, 79–114 (2005).

40. J. P. Lilgendahl, D. P. McAdams, Constructing stories of self-growth: how individual differences in patterns of autobiographical reasoning relate to well-being in midlife. *J. Pers.* **79**, 391–428 (2011).

41. J. A. Updegraff, E. M. Suh, Happiness is a warm abstract thought: Self-construal abstractness and subjective well-being. *J. Posit. Psychol.* **2**, 18–28 (2007).

42. V. J. Fortunato, J. T. Furey, The theory of MindTime: The relationships between Future, Past, and Present thinking and psychological well-being and distress. *Pers. Individ. Dif.* **50**, 20–24 (2011).

43. K. Layous, J. L. Kurtz, Nostalgia: A potential pathway to greater well-being. *Curr Opin Psychol* **49**, 101548 (2023).

44. A. Sutton, Living the good life: A meta-analysis of authenticity, well-being and engagement. *Pers. Individ. Dif.* **153**, 109645 (2020).

45. Y. Lin, J. Mutz, P. J. Clough, K. A. Papageorgiou, Mental Toughness and Individual Differences in Learning, Educational and Work Performance, Psychological Well-being, and Personality: A Systematic Review. *Front. Psychol.* **8**, 1345 (2017).

46. J. Anglim, S. Horwood, L. D. Smillie, R. J. Marrero, J. K. Wood, Predicting psychological and subjective well-being from personality: A meta-analysis. *Psychol. Bull.* **146**, 279–323 (2020).

47. R. A. Giacalone, C. L. Jurkiewicz, M. Promislo, Ethics and Well-Being: The Paradoxical Implications of Individual Differences in Ethical Orientation. *J. Bus. Ethics* **137**, 491–506 (2016).

48. A. M. Brandmaier, N. Ram, G. G. Wagner, D. Gerstorf, Terminal decline in well-being: The role of multi-indicator constellations of physical health and psychosocial correlates. *Dev. Psychol.* **53**, 996–1012 (2017).

49. R. A. Abbott, *et al.*, The relationship between early personality and midlife psychological well-being: evidence from a UK birth cohort study. *Soc. Psychiatry Psychiatr. Epidemiol.* **43**, 679–687 (2008).

50. M. L. King, The neural correlates of well-being: A systematic review of the human neuroimaging and neuropsychological literature. *Cogn. Affect. Behav. Neurosci.* **19**, 779–796 (2019).

51. C. G. Richter, *et al.*, Brain Imaging Studies of Emotional Well-Being: A Scoping Review. *osf.io* (2023).

52. L. P. de Vries, M. P. van de Weijer, M. Bartels, A systematic review of the neural correlates of well-being reveals no consistent associations. *Neurosci. Biobehav. Rev.* **145**, 105036 (2023).

53. R. Alexander, *et al.*, The neuroscience of positive emotions and affect: Implications for cultivating happiness and wellbeing. *Neurosci. Biobehav. Rev.* **121**, 220–249 (2021).

54. A. Suardi, I. Sotgiu, T. Costa, F. Cauda, M. Rusconi, The neural correlates of happiness: A review of PET and fMRI studies using autobiographical recall methods. *Cogn. Affect. Behav. Neurosci.* **16**, 383–392 (2016).

55. J. R. Tanzer, L. Weyandt, Imaging Happiness: Meta Analysis and Review. *J. Happiness Stud.* **21**, 2693–2734 (2020).

56. J. Gonzalez-Castillo, P. A. Bandettini, Task-based dynamic functional connectivity: Recent findings and open questions. *Neuroimage* **180**, 526–533 (2018).

57. R. T. Hurlburt, B. Alderson-Day, C. Fernyhough, S. Kühn, What goes on in the resting-state? A qualitative glimpse into resting-state experience in the scanner. *Front. Psychol.* **6**, 1535 (2015).

58. L. Koban, P. J. Gianaros, H. Kober, T. D. Wager, The self in context: brain systems linking mental and physical health. *Nat. Rev. Neurosci.* **22**, 309–322 (2021).

59. H. Jo, Y.-Y. Ou, C.-C. Kung, The neural substrate of self- and other-concerned wellbeing: An fMRI study. *PLoS One* **14**, e0203974 (2019).

60. J. I. Skipper, A voice without a mouth no more: The neurobiology of language and consciousness. *Neurosci. Biobehav. Rev.* **140**, 104772 (2022).

61. R. Millière, Looking for the Self: Phenomenology, Neurophysiology and Philosophical Significance of Drug-induced Ego Dissolution. *Front. Hum. Neurosci.* **11**, 245 (2017).

62. R. Millière, R. L. Carhart-Harris, L. Roseman, F.-M. Trautwein, A. Berkovich-Ohana, Psychedelics, Meditation, and Self-Consciousness. *Front. Psychol.* **9**, 1475 (2018).

63. I. Gallagher I., Philosophical conceptions of the self: implications for cognitive science. *Trends Cogn. Sci.* **4**, 14–21 (2000).

64. H. F. Araujo, J. Kaplan, H. Damasio, A. Damasio, Neural correlates of different self domains. *Brain Behav.* **5**, e00409 (2015).

65. P. Frewen, *et al.*, Neuroimaging the consciousness of self: Review, and conceptual-methodological framework. *Neurosci. Biobehav. Rev.* **112**, 164–212 (2020).

66. R. Dings, The dynamic and recursive interplay of embodiment and narrative identity. *Philos. Psychol.* **32**, 186–210 (2019).

67. A. Bortolan, Affectivity and the distinction between minimal and narrative self. *Continental Philosophy Review* **53**, 67–84 (2020).

68. A. K. Seth, Interoceptive inference, emotion, and the embodied self. *Trends Cogn. Sci.* **17**, 565–573 (2013).

69. A. R. Damasio, *The Feeling of what Happens: Body and Emotion in the Making of Consciousness*, Reprint edition (Harcourt Brace, 1999).

70. I. R. Kleckner, *et al.*, Evidence for a Large-Scale Brain System Supporting Allostasis and Interoception in Humans. *Nat Hum Behav* **1** (2017).

71. L. F. Barrett, The theory of constructed emotion: an active inference account of interoception and categorization. *Soc. Cogn. Affect. Neurosci.* **12**, 1–23 (2017).

72. H. D. Critchley, S. Wiens, P. Rotshtein, A. Ohman, R. J. Dolan, Neural systems supporting interoceptive awareness. *Nat. Neurosci.* **7**, 189–195 (2004).

73. Y. Terasawa, H. Fukushima, S. Umeda, How does interoceptive awareness interact with the subjective experience of emotion? An fMRI study. *Hum. Brain Mapp.* **34**, 598–612 (2013).

74. J. Zaki, J. I. Davis, K. N. Ochsner, Overlapping activity in anterior insula during interoception and emotional experience. *Neuroimage* **62**, 493–499 (2012).

75. O. Pollatos, K. Gramann, R. Schandry, Neural systems connecting interoceptive awareness and feelings. *Hum. Brain Mapp.* **28**, 9–18 (2007).

76. Y. Tan, R. Yan, Y. Gao, M. Zhang, G. Northoff, Spatial-topographic nestedness of interoceptive regions within the networks of decision making and emotion regulation: Combining ALE meta-analysis and MACM analysis. *Neuroimage* **260**, 119500 (2022).

77. P. Qin, M. Wang, G. Northoff, Linking bodily, environmental and mental states in the self—A three-level model based on a meta-analysis. *Neurosci. Biobehav. Rev.* **115**, 77–95 (2020).

78. S. Aliko, J. Huang, F. Gheorghiu, S. Meliss, J. I. Skipper, A naturalistic neuroimaging database for understanding the brain using ecological stimuli. *Sci Data* **7**, 347 (2020).

79. P.-H. A. Chen, E. Jolly, J. H. Cheong, L. J. Chang, Intersubject representational similarity analysis reveals individual variations in affective experience when watching erotic movies. *Neuroimage* **216**, 116851 (2020).

80. J. M. van Baar, L. J. Chang, A. G. Sanfey, The computational and neural substrates of moral

strategies in social decision-making. *Nat. Commun.* **10**, 1483 (2019).

81. L. J. Chang, *et al.*, Endogenous variation in ventromedial prefrontal cortex state dynamics during naturalistic viewing reflects affective experience. *Sci Adv* **7** (2021).

82. R. C. Gershon, *et al.*, NIH toolbox for assessment of neurological and behavioral function. *Neurology* **80**, S2–6 (2013).

83. T. Yarkoni, R. A. Poldrack, T. E. Nichols, D. C. Van Essen, T. D. Wager, Large-scale automated synthesis of human functional neuroimaging data. *Nat. Methods* **8**, 665–670 (2011).

84. J. M. Salsman, *et al.*, Emotion assessment using the NIH Toolbox. *Neurology* **80**, S76–86 (2013).

85. H. Lee Masson, L. Isik, Functional selectivity for social interaction perception in the human superior temporal sulcus during natural viewing. *Neuroimage* **245**, 118741 (2021).

86. N. M. Khoury, J. Lutz, Z. Schuman-Olivier, Interoception in Psychiatric Disorders: A Review of Randomized, Controlled Trials with Interoception-Based Interventions. *Harv. Rev. Psychiatry* **26**, 250–263 (2018).

87. M. Tsakiris, H. Critchley, Interoception beyond homeostasis: affect, cognition and mental health. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **371** (2016).

88. T. Pinna, D. J. Edwards, A Systematic Review of Associations Between Interoception, Vagal Tone, and Emotional Regulation: Potential Applications for Mental Health, Wellbeing, Psychological Flexibility, and Chronic Conditions. *Front. Psychol.* **11**, 1792 (2020).

89. F. Adolfi, *et al.*, Convergence of interoception, emotion, and social cognition: A twofold fMRI meta-analysis and lesion approach. *Cortex* **88**, 124–142 (2017).

90. J. A. Chalmers, D. S. Quintana, M. J.-A. Abbott, A. H. Kemp, Anxiety Disorders are Associated with Reduced Heart Rate Variability: A Meta-Analysis. *Front. Psychiatry* **5**, 80 (2014).

91. S. S. Khalsa, *et al.*, Interoception and Mental Health: A Roadmap. *Biol Psychiatry Cogn Neurosci Neuroimaging* **3**, 501–513 (2018).

92. T. Fuchs, J. E. Schlimme, Embodiment and psychopathology: a phenomenological perspective. *Curr. Opin. Psychiatry* **22**, 570–575 (2009).

93. A. K. Seth, K. J. Friston, Active interoceptive inference and the emotional brain. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **371** (2016).

94. X. Gu, P. R. Hof, K. J. Friston, J. Fan, Anterior insular cortex and emotional awareness. *J. Comp. Neurol.* **521**, 3371–3388 (2013).

95. A. D. Craig, How do you feel? Interoception: the sense of the physiological condition of the body. *Nat. Rev. Neurosci.* **3**, 655–666 (2002).

96. A. Etkin, T. Egner, R. Kalisch, Emotional processing in anterior cingulate and medial prefrontal cortex. *Trends Cogn. Sci.* **15**, 85–93 (2011).

97. B. A. Vogt, "Chapter 3 - Cingulate cortex in the three limbic subsystems" in *Handbook of Clinical Neurology*, B. A. Vogt, Ed. (Elsevier, 2019), pp. 39–51.

98. M. D. Lieberman, M. A. Straccia, M. L. Meyer, M. Du, K. M. Tan, Social, self, (situational), and affective processes in medial prefrontal cortex (MPFC): Causal, multivariate, and reverse inference evidence. *Neurosci. Biobehav. Rev.* **99**, 311–328 (2019).

99. D. D. Wagner, J. V. Haxby, T. F. Heatherton, The representation of self and person knowledge in the medial prefrontal cortex. *Wiley Interdiscip. Rev. Cogn. Sci.* **3**, 451–470 (2012).

100. S. N. Haber, B. Knutson, The reward circuit: linking primate anatomy and human imaging. *Neuropsychopharmacology* **35**, 4–26 (2010).

101. K. S. Wang, D. V. Smith, M. R. Delgado, Using fMRI to study reward processing in humans: past, present, and future. *J. Neurophysiol.* **115**, 1664–1678 (2016).

102. G. Northoff, D. J. Hayes, Is our self nothing but reward? *Biol. Psychiatry* **69**, 1019–1025 (2011).

103. D. Sander, L. Nummenmaa, Reward and emotion: an affective neuroscience approach. *Current Opinion in Behavioral Sciences* **39**, 161–167 (2021).

104. S. Padmala, N. Sambuco, L. Pessoa, "Chapter 1 - Interactions between reward motivation and emotional processing" in *Progress in Brain Research*, N. Srinivasan, Ed. (Elsevier, 2019), pp. 1–21.

105. D. R. Baerger, D. P. McAdams, Life Story Coherence and its Relation to Psychological Well-Being. *Narrative Inquiry* **9**, 69–96 (1999).

106. S. Bluck, T. Habermas, The life story schema. *Motiv. Emot.* **24**, 121–147 (2000).

107. J. J. Bauer, D. P. McAdams, J. L. Pals, Narrative identity and eudaimonic well-being. *J. Happiness Stud.* **9**, 81–104 (2008).

108. K. C. McLean, *et al.*, The empirical structure of narrative identity: The initial Big Three. *J. Pers. Soc. Psychol.* **119**, 920–944 (2020).

109. K. A. McLaughlin, T. D. Borkovec, N. J. Sibrava, The effects of worry and rumination on affect states and cognitive activity. *Behav. Ther.* **38**, 23–38 (2007).

110. J. Moffatt, K. J. Mitrenga, B. Alderson-Day, P. Moseley, C. Fernyhough, Inner experience differs in rumination and distraction without a change in electromyographical correlates of inner speech. *PLoS One* **15**, e0238920 (2020).

111. E. Kross, *et al.*, Self-talk as a regulatory mechanism: how you do it matters. *J. Pers. Soc. Psychol.* **106**, 304–324 (2014).

112. C. Reichl, J. F. Schneider, F. M. Spinath, Relation of self-talk frequency to loneliness, need to belong, and health in German adults. *Pers. Individ. Dif.* **54**, 241–245 (2013).

113. K. R. H. Treadwell, P. C. Kendall, Self-talk in youth with anxiety disorders: States of mind, content specificity, and treatment outcome. *J. Consult. Clin. Psychol.* **64**, 941–950 (1996).

114. J. M. Adler, J. Lodi-Smith, F. L. Philippe, I. Houle, The Incremental Validity of Narrative Identity in Predicting Well-Being: A Review of the Field and Recommendations for the Future. *Pers. Soc. Psychol. Rev.* **20**, 142–175 (2016).

115. D. P. McAdams, K. C. McLean, Narrative Identity. *Curr. Dir. Psychol. Sci.* **22**, 233–238 (2013).

116. J. I. Skipper, J. T. Devlin, D. R. Lametti, The hearing ear is always found close to the speaking tongue: Review of the role of the motor system in speech perception. *Brain Lang.* **164**, 77–105 (2017).

117. J. I. Skipper, H. C. Nusbaum, S. L. Small, "Lending a helping hand to hearing: Another motor theory of speech perception" in *Action to Language via the Mirror Neuron System*, M. A. Arbib, Ed. (Cambridge University Press, 2006), pp. 250–286.

118. J. I. Skipper, S. L. Small, "fMRI Studies of Language" in *Encyclopedia of Language & Linguistics (Second Edition)*, K. Brown, Ed. (Elsevier, 2006), pp. 496–511.

119. A. E. Cavanna, M. R. Trimble, The precuneus: a review of its functional anatomy and behavioural correlates. *Brain* **129**, 564–583 (2006).

120. B. L. Foster, *et al.*, A tripartite view of the posterior cingulate cortex. *Nat. Rev. Neurosci.* **24**, 173–189 (2023).

121. S. Wang, L. J. Tepfer, A. A. Taren, D. V. Smith, Functional parcellation of the default mode network: a large-scale meta-analysis. *Sci. Rep.* **10**, 16096 (2020).

122. V. Menon, 20 years of the default mode network: A review and synthesis. *Neuron* (2023) https:/doi.org/10.1016/j.neuron.2023.04.023.

123. S. Daviddi, T. Pedale, P. L. St Jacques, D. L. Schacter, V. Santangelo, Common and distinct correlates of construction and elaboration of episodic-autobiographical memory: An ALE meta-analysis. *Cortex* **163**, 123–138 (2023).

124. G. Mazzoni, *et al.*, Brain activation in highly superior autobiographical memory: The role of the precuneus in the autobiographical memory retrieval network. *Cortex* **120**, 588–602 (2019).

125. J. J. Summerfield, D. Hassabis, E. A. Maguire, Cortical midline involvement in autobiographical memory. *Neuroimage* **44**, 1188–1200 (2009).

126. P. Martinelli, M. Sperduti, P. Piolino, Neural substrates of the self-memory system: new insights from a meta-analysis. *Hum. Brain Mapp.* **34**, 1515–1529 (2013).

127. M. K. Johnson, *et al.*, Dissociating medial frontal and posterior cingulate activity during self-reflection. *Soc. Cogn. Affect. Neurosci.* **1**, 56–64 (2006).

128. P. Qin, *et al.*, Dissociation between anterior and posterior cortical regions during self-specificity and familiarity: a combined fMRI-meta-analytic study. *Hum. Brain Mapp.* **33**, 154–164 (2012).

129. G. Rabini, S. Ubaldi, S. L. Fairhall, Combining concepts across categorical domains: a linking role of the precuneus. *Neurobiol Lang (Camb)* **2**, 354–371 (2021).

130.   J. R. Binder, R. H. Desai, W. W. Graves, L. L. Conant, Where Is the Semantic System? A Critical Review and Meta-Analysis of 120 Functional Neuroimaging Studies. *Cereb. Cortex* **19**, 2767–2796 (2009).

131.   L. Fletcher, S. C. Hayes, Relational frame theory, acceptance and commitment therapy, and a functional analytic definition of mindfulness. *J. Ration. Emot. Cogn. Behav. Ther.* **23**, 315–336 (2005).

132.   J. Gao, *et al.*, Functional Segregation of the Middle Temporal Visual Motion Area Revealed With Coactivation-Based Parcellation. *Front. Neurosci.* **14**, 427 (2020).

133.   M.-H. Grosbras, S. Beaton, S. B. Eickhoff, Brain regions involved in human movement perception: a quantitative voxel-based meta-analysis. *Hum. Brain Mapp.* **33**, 431–454 (2012).

134.   S. Vossel, J. J. Geng, G. R. Fink, Dorsal and ventral attention systems: distinct neural circuits but collaborative roles. *Neuroscientist* **20**, 150–159 (2014).

135.   T. W. Schmitz, E. De Rosa, A. K. Anderson, Opposing Influences of Affective State Valence on Visual Cortical Encoding. *Journal of Neuroscience* **29**, 7199–7207 (2009).

136.   B. Grafton, C. MacLeod, Attentional Bias and Well-Being: How the Bias That Feels Best Can Be Bad for Us (2019) https:/doi.org/10.1093/oxfordhb/9780190653200.013.9 (May 24, 2023).

137.   A. Martinelli, J. Grüll, C. Baum, Attention and interpretation cognitive bias change: A systematic review and meta-analysis of bias modification paradigms. *Behav. Res. Ther.* **157**, 104180 (2022).

138.   R. T. Hurlburt, *et al.*, Measuring the Frequency of Inner-Experience Characteristics. *Perspect. Psychol. Sci.*, 1745691621990379 (2021).

139.   H. Roebuck, G. Lupyan, The Internal Representations Questionnaire: Measuring modes of thinking. *Behav. Res. Methods* **52**, 2053–2070 (2020).

140.   A. Oh, E. G. Duerden, E. W. Pang, The role of the insula in speech and language processing. *Brain Lang.* **135**, 96–103 (2014).

141.   L. Pessoa, On the relationship between emotion and cognition. *Nat. Rev. Neurosci.* **9**, 148–158 (2008).

142.   C. Morawetz, *et al.*, Multiple large-scale neural networks underlying emotion regulation. *Neurosci. Biobehav. Rev.* **116**, 382–395 (2020).

143.   I. Messina, S. Bianco, M. Sambin, R. Viviani, Executive and semantic processes in reappraisal of negative stimuli: insights from a meta-analysis of neuroimaging studies. *Front. Psychol.* **6**, 956 (2015).

144.   J. A. Brooks, *et al.*, The role of language in the experience and perception of emotion: a neuroimaging meta-analysis. *Soc. Cogn. Affect. Neurosci.* **12**, 169–183 (2017).

145.   U. Hasson, J. Chen, C. J. Honey, Hierarchical process memory: memory as an integral component of information processing. *Trends Cogn. Sci.* **19**, 304–313 (2015).

146.   V. Gallese, C. Sinigaglia, What is so special about embodied simulation? *Trends Cogn. Sci.* **15**, 512–519 (2011).

147.   G. Hesslow, The current status of the simulation theory of cognition. *Brain Res.* **1428**, 71–79 (2012).

148.   Y. Yeshurun, M. Nguyen, U. Hasson, The default mode network: where the idiosyncratic self meets the shared social world. *Nat. Rev. Neurosci.* (2021) https:/doi.org/10.1038/s41583-020-00420-w.

149.   R. W. Cox, AFNI: software for analysis and visualisation of functional magnetic resonance neuroimages. *Comput. Biomed. Res.* **29**, 162–173 (1996).

150.   B. Fischl, FreeSurfer. *Neuroimage* **62**, 774–781 (2012).

151.   C. Destrieux, B. Fischl, A. Dale, E. Halgren, Automatic parcellation of human cortical gyri and sulci using standard anatomical nomenclature. *Neuroimage* **53**, 1–15 (2010).

152.   C. F. Beckmann, S. M. Smith, Probabilistic independent component analysis for functional magnetic resonance imaging. *IEEE Trans. Med. Imaging* **23**, 137–152 (2004).

153.   K. J. Gorgolewski, *et al.*, NeuroVault.org: a web-based repository for collecting and sharing unthresholded statistical maps of the human brain. *Front. Neuroinform.* **9**, 8 (2015).

154.   P. Virtanen, *et al.*, SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* **17**, 261–272 (2020).

155.   N. Mantel, The detection of disease clustering and a generalised regression approach. *Cancer Res.* **27**, 209–220 (1967).

156.   A. B. Poppe, *et al.*, Toward a neurometric foundation for probabilistic independent component analysis of fMRI data. *Cogn. Affect. Behav. Neurosci.* **13**, 641–659 (2013).

157.   L. Griffanti, *et al.*, Hand classification of fMRI ICA noise components. *Neuroimage* **154**, 188–205 (2017).

158.   T. Salo, *et al.*, NiMARE: Neuroimaging meta-analysis research environment. *NeuroLibre* **1**, 7 (2022).