# Sex- versus apomictic-specific polymorphisms in the 5'UTR of

APOLLO from *Boechera* shift gene expression from somatic to

reproductive tissues in *Arabidopsis*

**Maryam Honari[1], Joanne R. Ashnest [1,2], Timothy F. Sharbel [1,3]**

[1] Global Institute for Food Security, University of Saskatchewan, Saskatoon, SK, Canada

[2] Present address: School of Agriculture, Environmental and Veterinary Sciences, Charles Sturt
University, Wagga Wagga, NSW, Australia

[3] Present address: Department of Plant Sciences, University of Saskatchewan, Saskatoon, SK,
Canada

**\* Correspondence:**
Corresponding Authors: mhonari@lakeheadu.ca; Tim.Sharbel@usask.ca
12

13 **Keywords: Apomixis, APOLLO, *Boechera*, promoter-swap, 5'UTR, reproduction**
14

## Abstract

Among candidate genes underlying the control components of apomixis, APOLLO is known for
its strong linkage to apomeiosis in the genus *Boechera*. The gene has "apo-alleles", which are
characterized by a set of linked apomixis-specific polymorphisms, and "sex-alleles". All
apomictic *Boechera* genotypes are heterozygous for the apo/sex-alleles, while all sexual
genotypes are homozygous for sex-alleles. In this study, native and synthetic APOLLO
promoters were characterized by detecting the expression level of the β-glucuronidase (GUS)
gene in *Arabidopsis*. Comparing various flower developmental stages in transgenic lines
containing different constructs with 2 kb native transgenic lines revealed that changes to the
APOLLO promoter causes shifts in tissue- and developmental-stage specificity of GUS
expression. Importantly, several apomixis-specific polymorphisms in the 5'UTR change the
timing and location of GUS activity from somatic to reproductive tissues. These synthetic data
simulate a plausible evolutionary process whereby apomixis-specific gene activity can be
achieved.

## Introduction

In contrast to sexual reproduction and segregation of traits in his peas, Mendel's research on

*Hieracium* illustrated a distinct kind of inheritance in which segregation was absent, and which

was later identified as the parthenogenetic development of meiotically-unreduced egg cells

(Bicknell et al, 2016; Mittelsten Scheid, 2022; van Dijk and Noel Ellis 2023). As with

*Hieracium*, many plant species reproduce through apomixis, a natural form of asexual reproduction through which seeds are produced with embryos which are genetic clones of the mother plant (Hojsgaard and Hörandl, 2019; Underwood and Mercier, 2022).

While apomixis can arise via different mechanisms in natural plant populations (Hojsgaard and Hörandl, 2019; Underwood and Mercier, 2022), it fundamentally differs from sexual seed formation by the production of the meiotically-unreduced egg-cell (apomeiosis) which develops into an embryo without fertilization (parthenogenesis) in addition to endosperm development with or without fertilization of the central cell (Hojsgaard and Hörandl, 2019). Many apomicts are furthermore facultative (i.e., producing both sexually- and apomictically-derived embryos), and hence there is interplay between the reproductive pathways (Koltunow et al., 2011; Bicknell et al., 2003; Fehrer et al., 2007; Aliyu et al., 2010). Apomicts are often interspecific hybrids, which has led to the hypothesis that deregulation of genes underlying sexual seed formation underlies the genesis of apomixis (Carman, 1997). Furthermore, many apomictic plants and parthenogenetic animals are also polyploid, with implications for changes in gene expression and mutational load tolerance (among other traits) as being favorable for apomixis (Hojsgaard, 2018).

Apomixis has received much attention because of its potential ability to freeze complex hybrid genotypes (Tucker and Koltunow 2009). It does not exist in crop plants, and the introduction of apomixis from wild apomictic relatives into crops (Underwood et al., 2022), or via *de novo* induction would be highly disruptive to modern agriculture (Spillane et al., 2004; Underwood et al., 2022). Recent progress in inducing apomixis properties in lettuce and rice demonstrates the importance of research in this field to establish new tools to improve global food security (Underwood et al., 2022; Vernet et al., 2022).

The genus *Boechera* Á. Löve & D. Löve (Boecher's rock cress; Brassicaceae)(Dobeš et al., 2006) is an ideal model system to study the evolution of apomixis, a major reason being that sexual and apomict plants can be compared genetically at the same ploidy level (Corral et al., 2013). The genus is characterized by diploid sexuals (basic chromosome number $x = 7$; $2n = 2x = 14$), as well as diploid ($2n = 2x = 14$ or $15$) and triploid ($2n = 3x = 21$) apomicts (Kantama et al., 2007; Alexander et al., 2013; Lovell et al., 2013). A highly reticulate pattern of genetic variation on inter- and intraspecific levels is reflective of numerous diploid and triploid

apomictic lineages in geographically and genetically distinct populations (Rushworth et al.,
2011; Rushworth et al., 2018; Kiefer et al., 2009), whereby rare haploid apomictic pollen can
spread of apomixis factors to establishment new diploid apomictic lineages (Mau et al., 2021).

The introduction of apomixis-like reproduction in plants, using gene mutants (d'Erfurth et al.,
2009), or through genetic constructs based upon candidate loci identified in different wild
apomictic species (Underwood et al., 2022) demonstrate that multiple functional pathways are
available to achieve asexuality. From a functional evolutionary perspective, deregulation of
genes underlying sexual seed formation likely involves multiple loci which are simultaneously
co-opted to induce apomixis, with changes to tissue-specificity and timing of gene expression
leading to the switch from sexual to apomictic seed production in natural populations (Schmidt,
2020).

Here, we demonstrate that sex- vs. apomixis-specific polymorphisms in the 5'UTR of the
apomixis candidate APOLLO gene (TGGCCCGTGAAGTTTATTCC)(Corral et al., 2013) can
modify timing and location of gene expression in Arabidopsis. UTR-control of gene expression
has been demonstrated to be significant in other plant species (Srivastava et al., 2018) as well as
humans (Babu et al. 2021). For example, a novel motif in the 5'UTR of Big Root Biomass (BRB)
gene modulates root biomass in sesame, whereby this motif in single and duplicated copies was
responsible for high and low root biomass in two different accessions respectively (Dossa et al.,
2021). Global spatial analysis of Arabidopsis natural variation revealed 5'UTR splicing of LATE
ELONGATED HYPOCOTYL (LHY) in response to temperature (James et al., 2018). In
addition, blue light exposure in Arabidopsis triggers variation in 5'UTR length resulting in
differential transcription start site (TSSs) usage which in turn influences the post-transcriptional
regulatory processes (Georgy, 2018; Yukio et al., 2018).

In *Boechera,* APOLLO is part of the DEDD 3′→5′ exonuclease superfamily (Corral et al., 2013),
and the expression of its orthologue in *A. thaliana* (AT1G74390) has been reported in flowering
stages, mature plant embryos, petal differentiation and expansion stages, plant embryo bilateral
stages, plant embryo cotyledonary stages and plant embryo globular stages
(https://www.arabidopsis.org). Sexual *Boechera* are homozygous for an APOLLO sex-allele,
while apomicts have both apo- and sex-alleles which differ in sequence composition and
structure (Corral et al., 2013). In *Boechera*, apo-alleles of APOLLO are highly expressed in

developing apomictic ovules during the megaspore mother cell stage, while sex-alleles are not

expressed in apomictic ovules (Corral et al., 2013). No APOLLO expression occurs in the ovules

of sexual plants (Corral et al., 2013). Ovule-specific expression of APOLLO in apomictic

*Boechera* is hypothesized to be associated with differences in the 5'UTR region which differs

between apo- and sex-alleles, whereby the apo-allele 5'UTR is truncated and additionally

contains an internal 20 bp insertion in comparison to the sex-allele (Corral et al., 2013). Here, we

have employed a promoter-swap experiment between apo- and sex-allele specific sequences to

investigate the influence of the apomixis-specific polymorphisms on the gene expression of this

exonuclease in Arabidopsis.

## Materials and methods

### *Plant materials and growth conditions*

*Arabidopsis thaliana* (Columbia-0 ecotype) seeds were obtained from the Arabidopsis Biological

Resource Centre (ABRC). Surface sterilised seeds were sown on MS plates containing ½ MS

(Murashige & Skoog) medium (pH adjusted to 5.7 with 1M NaOH) with 0.8% phytoagar,

stratified at 4°C for 48 h and then transferred to a phytochamber at 20°C with a light cycle of 16h

light (white fluorescent bulbs, with a light intensity of 70 μmol m$^{-2}$ s$^{-1}$) and 8h dark. Two-week

old seedlings were transferred to pots containing Sunshine mix 8 and grown in the same

conditions.

### *Construct generation and plant transformation*

Native and promoter swap constructs were created using various combinations of sex- versus

apo-specific polymorphisms identified in Corral et al. (2013; Figure 1, Supplemental Table 1).

All primers used are described in Supplemental tables 2 and 3. All verified construct sequences

have been submitted to Gene bank, as listed in Supplemental table 4.

To build native *Boechera APOLLO* putative promoter constructs (C1 and C2), approximately 2

kb upstream of start codon, including the 5'UTR, was amplified from *B. stricta* (ES 718; sexual)

and *B. divaricarpa* (ES 517; apomictic) genomic DNA using CloneAmp (Thermo Fischer

Scientific) DNA polymerase. Specific primers were designed based on BAC apo- and sex- allele

sequence libraries (Corral et al. 2013). Synthetic promoter splice variants (C3 to C7) were

constructed using a modified "gene Splicing by Overlap Extension" method (SOE) (Horton et

al., 1989). For C3, C4 and C5, modifications to the 5'UTR indel region were created by
modifying the respective primer sequences (Supplemental figure 1). Construct pC5 was
composed of the 2 kb native apo promoter plus the 5′UTR of the apo-allele, but the apomixis-
specific 20 bp insertion sequence was randomized (but ratios of base pairs remained identical to
the wild-type insertion; Figure 1). Resultant PCR products were purified and cloned into a
pENTR™/D-TOPO™ entry vector via directional TOPO® cloning, and incorporated into the
pBGWFS7 GFP:GUS reporter vector (Nakagawa et al., 2007) using Gateway™ LR Clonase™.
All constructs were verified using Sanger sequencing.

Binary vectors were transformed into the *Agrobacterium tumefaciens* strain GV3101 by
electroporation, and eight-week old wild-type *Arabidopsis* Col-0 plants were transformed by
floral dip (Clough and Bent, 1998). Transformed seeds were selected based on the method by
Harrison et al. (2006). Two-week old glufosinate-resistant seedlings were then transferred to pots
containing Sunshine mix 8, and presence of the transgene was confirmed by PCR of genomic
DNA.

## *Histochemical assay of GUS activity*

Tissue-specific promoter activity for all constructs was assessed by β-glucuronidase (GUS)
activity; seedlings and flowers from at least 10 different independent transformation events were
examined per construct. GUS-staining was performed following the protocol of Jefferson (1987).
Developmental staging was based on pistil and stigma maturity according to Smyth et al. (1990)
and Christensen et al. (1997). We defined 5 developmental stages as in Rojeck et al. (2021) for
analyses based upon pistil length, whereby the first stage (pre-meiotic stage) had a pistil length
of 0.7 mm, and for the post-meiotic stages the pistil lengths were 1.0 mm, 1.5 mm, 2.0 mm and
3.0 mm. Among all of these stages the final stage is considered as post-fertilization stage (Rojeck
et al., 2021)

Tissues were cleared following the protocol of Malamy and Benfey (1997). Samples were
mounted in 50% glycerol on glass microscope slides. Among 10 different pre-assessed
transgenic lines, 3 were used for microscopic analysis and photography, and *A. thaliana* var.
Columbia that was not exposed to the clearing material was used as a control. Promoter activity

in the leaves and flowers of *A. thaliana* was measured by a GUS histochemical assay at two- and five-days post-treatment in a minimum of three flowers per developmental stage (Smyth et al., 1990; Rojek et al., 2018). Images were obtained under a Zeiss Lumar.V12 Stereoscope and Olympus BX61 microscope and analyzed using Image J (https://rsb.info.nih.gov/ij/). To compare intensity of GUS positive areas, the linear region of interest (ROI) method was used, whereby a straight line is drawn through cell images to depict the main intensity of the color (Béziat et al., 2017). The original data was normalized by Min-Max method and analyzed using a two-way analysis of variance (two-way ANOVA). Finally, statistical differences were determined by Tukey test.

## Results

### (a) The 2kb native *Boechera* Apo promoter is expressed in the vascular tissues of *Arabidopsis*

Positive GUS activity for root, shoots, and leaves of two-week old seedlings for transgenic lines carrying the construct pC1 (2 kb native apo promoter) was observed in 33 lines of two-week old seedlings, each with 4 replicates (Figure 2). From this pattern (Figure 2a), three of the 33 lines were negative only for roots, shoots, or leaves, whereas 5 of 33 lines were negative only for roots (Supplemental figure 2). In contrast, no GUS activity was observed for 31 lines of two-week old seedlings transgenic for the construct pC2, carrying the 2 kb Sex native promoter (Figure 2b; Supplemental figure 2).

There was positive GUS activity in leaves of six-week old *Arabidopsis* transgenic lines for pC1 (Figure 2c), and no GUS activity in any plants containing pC2 (Supplemental figure 2).

### (b) Removal of a 20 bp apo-specific insertion from the native 2kb apo-allele changes timing and tissue-specificity of expression

Construct pC3 (2kb native apo promoter plus 5'UTR with apo-specific insertion deleted; Figure 1) GUS activity was measured in 8 transgenic lines, over five different developmental stages (total of 40 flower buds). No GUS activity was observed in the pre-meiotic (0.7 mm) and post meiotic (1.0 mm) flower stages (Figure 3a, b, f, g), while positive GUS activity was observed for

stages 3, 4, and 5 in the style and filament (including vascular tissue) with an increasing activity

toward the final stages (Figure 3c, d, e, h, i, j; P≤0.05, Tukey test; Supplemental figure 3).

Dispersed GUS activity in petals was excluded because they did not show a constant pattern for

all samples.  Thus, removal of the 20 bp apo-specific insertion from its native 2kb allele shifts

expression from somatic to both male and female reproductive tissue.

(c) A 20 bp apo-specific insertion does not induce expression in the 2kb native

*Boechera* sex-allele

Construct pC4 carried the 2 kb native pSex promoter, the 5'UTR of the Sex-allele, with the 20 bp

Apo-insertion added. The GUS activity for 7 transgenic pC4 lines, each of which had five

different developmental stages (total of 35 flower buds) was measured, and no GUS activity was

detected (Supplemental figure 4).

(d) Randomizing the 20 bp apo-specific insertion but maintaining size induces

anther and stigma expression of the native apo-promoter

Construct pC5 was composed of the 2 kb native pApo promoter plus the 5'UTR of the apo-allele,

but a 20 bp randomized (i.e. same nucleotide composition but different sequence) insertion was

substituted for the apomixis-specific 20 bp insertion (Figure 1). GUS activity was observed in

the anthers and receptacle for the pre-meiotic stage, as well as some anther-specific GUS activity

for stages 2 (Figure 4a, b, f, g; Supplemental figure 5). Style-specific GUS activity appeared in

stages 3, and increased in intensity over stages 3, 4, and 5 (Figure 4c, d, e, h, i, j). GUS activity

for the flower receptacle and petals were excluded for quantification because they did not show a

constant pattern for all samples.

(e) The apo-allele promoter with substituted sex-allele 5'UTR induces expression
in polllen

Construct pC6 carried the 2 kb pApo promoter, with a swapped native 5'UTR of sex-allele

(Figure 1), and was analyzed in 4 transgenic lines over five different developmental stages (total

of 20 flower buds). Anther-specific GUS activity with varying intensity was observed for all

developmental stages (Figure 5), with the strongest expression in stage 2 (Figure 5b; P≤0.05;

Tukey test; Supplemental figure 6). We hypothesize that pollen activity is induced (Figure 5g to

k), as is evidenced by post-dehiscence loss of activity (Figure 5d to e).

(f) The sex-allele promoter with substituted apo-allele 5'UTR induces a burst of
expression in both pollen and ovules in pre-zygote stages, followed by decreasing
expression in zygotes

Construct pC7 carried the 2 kb sex promoter with a swapped apo-allele 5'UTR (Figure 1), and

was analyzed in 6 transgenic lines over five developmental stages (total of 30 flower buds). GUS

activity with a varied intensity was observed in the transmitting tract, ovules and anthers for the

first two developmental stages (Figure 6a, b), while only ovule expression occurred in stage 3

(Figures 6c). Ovule expression began in stage one (Figure 6) and lasted for three stages (P≤0.05;

Tukey test; Supplemental figure 7). In stage 4 (Figure 6d), expression in both tissues ceased.

## Discussion

A comparison of gene expression in the micro-dissected ovules of apomictic versus sexual

*Boechera* led to the identification of the APOLLO gene. The heterozygous nature and presence

of apomixis-specific polymorphisms were proposed to underlay gene expression differences

between pre-meiotic ovules of apomictic versus sexual *Boechera* (Corral et al., 2013). Here, we

investigated the nature of regulatory differences between native and synthetic APOLLO

promoter GUS  constructs in transformed Arabidopsis to understand functional differences

between conserved  sex- versus apo-allele polymorphisms in the 5′UTR region of the APOLLO

gene. We have made no effort here to understand sequence variation in the 2 kb promoter

regions of apo- and sex-alleles from multiple accessions of *Boechera*, as our focus was the

5′UTR region of the APOLLO gene. A comparison of the native promoter region sequences

(constructs pC1 and pC2; Supplemental table 4) nonetheless demonstrates that these upstream

regions also differ extensively, with potential implications for specific interactions in the

promoter swap constructs which have not been tested here.

These data demonstrate that evolutionarily conserved apomixis-specific polymorphisms in the

5'UTR of the apo-allele in *Boechera* can cause changes to gene expression timing and location in

*A. thaliana.* Importantly, coordinated expression in both male and female reproductive tissue

can be achieved, thus providing a tool which can potentially be applied to reproductive genetics

in Brassica plants.

**The native *Boechera* apo-allele promoter is active in the somatic tissues of *A. thaliana***

Arabidopsis transgenic lines for the 2 kb native Apo promoter demonstrate expression in the

vascular tissues of roots stems and leaves, and expression ceases before the development of

reproductive tissues (Figure 2). In contrast, no expression was observed for the 2 kb sex

promoter in any Arabidopsis transformants.

In an analysis of the 5'UTR regions of APOLLO, Corral et al. (2013) showed that the *Boechera*

20 bp apomixis-specific insertion of the apo-allele shared a *SORLIP1AT* transcription factor

binding site with the orthologous regions in *A. thaliana* and *Brassica rapa*, as well as a LIM1

transcription factor binding site with the orthologous regions in *A. thaliana*, while the sex-allele

did not. Furthermore, the *Boechera* apo-allele transcripts are truncated at the 5'end of the 20 bp

apo-specific insertion, while the sex-allele transcripts are approximately 35 bp longer (Corral et

al., 2013). Finally, the apo-allele is characterized by exclusive expression in the ovules of

apomictic *Boechera* (Corral et al., 2013), whereas in transformed sexual *A. thaliana* it is

expressed in somatic and not reproductive tissues (Figure 2). In this case, we hypothesize that

expression differences between the apo- and sex-alleles result from the different transcription

factor landscape of the *A. thaliana* genome.

**The apomixis-specific insertion: length and sequence important for expression in male and**

**female reproductive tissues**

Compared to the native apo-allele promoter-5'UTR (Figures 1 and 2), removal of the 20 bp

apomixis-specific insertion shifted expression from somatic to male and female reproductive

tissues (Figure 3). pC3 showed GUS activity in filaments (including vascular tissue) and style in

earlier and later stages of flower development respectively (Figure 3), and thus the absence of the

20 bp Apo-specific insertion changes the locality and timing of expression. Deletion of the 20 bp

Apo-insertion thus likely leads to a change in the recognition site for regulating interacting
elements that control expression in specific floral tissues.

Expression of the native *Boechera* apo-allele is not solely dependent on the apomixis-specific
insertion, as insertion of the 20 bp apomixis-specific sequence into the native *Boechera* sex-
allele produced no GUS activity in *A. thaliana* transformants (Supplemental figure 4).
Alternatively, the 2 kb upstream promoter region of the sex-allele may not be long enough to
drive expression.

Interestingly, the apomixis-specific insertion length is also important for expression in
reproductive tissues, as randomization of the insertion sequence (with identical nucleotide
composition) while maintaining length led to strong early expression in anther heads which
stopped post fertilization. Post-fertilization was then characterized by growing strong expression
in the stigma (Figure 4).

**Strong pollen-specific activity is induced by the apo-promoter/Sex-5'UTR construct.**

Transgenic pC6 lines carrying 2 kb of the apo promoter and the 5'UTR of the sex-allele (i.e.,
with no apomixis-specific 20 bp insertion) showed strong GUS staining in anthers in both pre-
and post-meiotic floral buds (Figure 5). Closer examination of the anthers demonstrated that
strong expression occurred in the pollen grains (Figure 5i, j). This expression ceased post-
fertilization (Figure 5e).

**A coordinated window of activity in anthers and ovules during fertilization**

Finally, transgenic pC7 lines carrying the 2 kb native sex promoter, 5'UTR of the apo-allele with
the 20 bp apo-insertion (Figure 6; Supplemental figure 7) showed a window of GUS activity in
both ovules and anthers in earlier stages of flower development which included the fertilization
stage (Figure 6). As the native sex-allele promoter (pC2) showed no activity in Arabidopsis
flowers (Figure 2), these data demonstrate that the apo 5'UTR can drive both male- and female-
specific expression in *A. thaliana*.

Considering the orthologue of the APOLLO gene in *A. thaliana* (AT1G74390), the NAC45/86-
DEPENDENT EXONUCLEASE-DOMAIN PROTEIN 3 (or NEN3), these data suggest that
polymorphisms shared between the *Boechera* apo-5'UTR and that of *A. thaliana* are functionally

involved with coordinating expression of this gene (or gene family) in male and female tissues

during fertilization.

**The apomixis-specific insertion: size and sequence important for expression in male and female reproductive tissues**

Removal of the 20 bp apomixis-specific insertion from the native apo-allele promoter-5'UTR

(Figure 1) shifted expression from somatic (Figure 2) to male and female reproductive tissues

(Figure 3). pC3 showed GUS activity in filaments and stigma in earlier and later stages of flower

development respectively (Figure 3), and thus the absence of the 20 bp apo-specific insertion

changes the locality and timing of expression. Deletion of the 20 bp apo-insertion thus likely

leads to a change in the recognition site for regulating interacting elements that control APOLLO

expression in specific floral tissues.

Expression of the native *Boechera* apo-allele is not solely dependent on the apomixis-specific

insertion, as insertion of the 20 bp apomixis-specific sequence into the native *Boechera* sex-

allele produced no GUS activity in *A. thaliana* transformants (Supplemental figure 4).

Alternatively, the 2 kb upstream promoter region of the sex-allele may not be long enough to

drive expression.

Interestingly, the apomixis-specific insertion length is also important for expression in

reproductive tissues, as randomization of the insertion sequence while maintaining length led to

strong early expression in anther heads which stopped post fertilization. Post-fertilization was

then characterized by growing strong expression in the style (Figure 4).

**Strong pollen-specific activity is induced by the apo-promoter/sex-5'UTR construct.**

Transgenic pC6 lines carrying 2 kb of the Apo promoter and the 5'UTR of the sex-allele (i.e.

with no apomixis-specific 20 bp insertion) showed strong GUS staining in anthers in both pre-

and post-meiotic floral buds (Figure 5). Closer examination of the anthers demonstrated that

strong expression occurred in the pollen grains (Figure 5i,j). This expression ceased post-

fertilization (Figure 5e).

As the native apo-allele promoter drives expression in *Arabidopsis* while the native sex-allele

promoter does not, the data so far demonstrate that the Apo-promoter upstream of the 5'UTR

drives expression in *A. thaliana*, while the 5'UTR underlies male- or female-specific tissue
expression.

**The 20 bp apo-specific insertion may bind a repressor in meiotic (sexual) cells**

The similarities between the 5′UTR sequences of the *Boechera* apo-allele, *A. thaliana* and *B.*
*rapa* suggest that the apo-allele is ancestral compared to the sex-allele, and that deregulation of
APOLLO in apomicts likely underlies its expression in pre-meiotic tissues (Corral et al. 2013).
This is supported by the data presented here, whereby in *A. thaliana*, a sexually reproducing
species, removal or scrambling of the apo-specific 5′UTR insertion and neighboring sequence
induced expression in both male and female reproductive tissue. Together, these data suggest
that the apo-5′UTR encodes a sequence-specific binding site for a transcriptional or translational
repressor, one that may be specifically expressed in sexual meiotic cells.

**Hybridization and differential gene expression between apomictic versus sexual *Boechera***

The elevated heterozygosity in diploid apomictic *Boechera* can be explained by widespread
hybridization associated with the transition from sexuality to asexuality, or it could be a
byproduct of apomixis itself.  In apomictic individuals the overall heterozygosity is expected to
increase over generations due to the lack of recombination and mutation accumulation (Meselson
effect; Hojsgaard and Hörandl, 2015). Clear evidence for the hybrid origin of highly
heterozygous apomicts has been shown (Beck et al., 2012), in addition to mutation accumulation
on a genomic level in apomictic *Boechera* in conserved non-coding sequences (Lovell et al.,
2017).  Finally, rare haploid pollen produced by apomictic *Boechera* can transfer the complete
apomictic trait in a single generation in both intra- and interspecific crosses onto emasculated
female plants (Mau et al., 2021).  Taken together, these data demonstrate that the conserved apo-
allele of *Boechera* is found with different sexual alleles on a broad geographic scale, not to
mention different genomic "contexts" considering the nuclear and cytoplasmic genomes in
independently derived apomictic lineages (Kiefer and Koch, 2012;  Kiefer et al., 2009).

Here we show that conserved polymorphisms in the native *Boechera* apo-allele 5'UTR are
functionally associated with driving gene expression in both male and female tissues during
important reproductive stages (e.g. gametogenesis and fertilization). The interspecific "collision"
of different sexual genomes may produce apomictic *Boechera* with structural changes such as

homeologous chromosome substitutions and aneuploidy (Kantama et al., 2007). Extending

divergence to regulatory genes, apomixis could result from the novel pattern of gene expressions

created by the interaction of divergent transcriptional regulators (Carman, 2001), and is

supported by the data reported in this study.

**Acknowledgements**

356 Experimental work was funded by a grant from the Global Institute of Food Security at the

357 University of Saskatchewan to Timothy F. Sharbel.  The authors thank Marco Pellino, Andres

358 Posso-Terranova, Zahida Irin, Nazmul Hasan, Angie Li,  Pierre-Luc Pradier for help with

359 greenhouse operations and plant care, as well as laboratory assistance.

# References

Alexander, P. J., Windham, M. D., Beck, J. B., Al-Shehbaz, I. A., Allphin, L, Bailey, C. D (2013).
Molecular phylogenetics and taxonomy of the genus *Boechera* and related genera (Brassicaceae:
Boechereae). *Systematic Botany* 38(1), 192–209. doi.org/10.1600/036364413X661917.

Aliyu, O. M., Schranz, M. E. and Sharbel, T. F. (2010). Quantitative variation for apomictic reproduction
in the genus Boechera (Brassicaceae). *American journal of botany* 97(10), 1719–31.
https://doi.org/10.3732/ajb.1000188

Babu, D., Vannelli, S., Fanelli, A., Mellone, S., Baffico, A. M.,  Corrado , L., et al. (2021). Variants in the
5'UTR Reduce SHOX Expression and Contribute to SHOX Haploinsufficiency. *European Journal of*
*Human Genetics* 29(1): 110–21. https://doi.org/10.1038/s41431-020-0676-y.

Beck, J. B., Alexander, P. J., Allphin, L., Al-Shehbaz, I. A., Rushworth, C., Bailey, C. D., et al., (2012).
Does hybridization drive the transition to asexuality in diploid Boechera? *Evolution* 66(4), 985–995.
https://doi.org/10.1111/j.1558-5646.2011.01507.x

Béziat, Ch., Jürgen Kleine-Vehn., and Feraru, E. (2017). Histochemical Staining of β-Glucuronidase and
Its Spatial Quantification BT  - Plant Hormones: Methods and Protocols. In eds. Jürgen Kleine-Vehn and
Michael Sauer. New York, NY: Springer New York, 73–80. https://doi.org/10.1007/978-1-4939-6469-
7_8.

Bicknell, R. A., Lambie, S. C., & Butler, R. C. (2003). Quantification of progeny classes in two
facultatively apomictic accessions of Hieracium. *Hereditas* 138(1), 11–20. https://doi.org/10.1034/j.1601-
5223.2003.01624.x

Bicknell, R., Catanach, A., Hand, M., & Koltunow, A. (2016). Seeds of doubt: Mendel's choice of
Hieracium to study inheritance, a case of right plant, wrong trait. *Theoretical and Applied Genetics*.
*Springer Verlag* 129(12), 2253–2266. https://doi.org/10.1007/s00122-016-2788-x

Brukhin, V., Osadtchiy, J. V., Florez-Rueda, A. M., Smetanin, D., Bakin, E., Nobre, M. S., et al., (2019).
The Boechera genus as a resource for apomixis research. *Frontiers in Plant Science* 10, 392.
https://doi.org/10.3389/fpls.2019.00392

Carman, J. G. (2001). The gene effect: genome collisions and apomixis. The Flowering of Apomixis:
from Mechanisms to Genetic Engineering. 95-110.

Carman, J. G. (1997). Asynchonous expression of duplicate genes in angiosperms may cause apomixis,
bispory, tetraspory, and polyembryony. *Biological Journal of the Linnean Society* 61(1), 51–94.
https://doi.org/10.1111/j.1095-8312.1997.tb01778.x

Christensen, Cory A, Edward J King, John R Jordan, and G N Drews. 1997. Megagametogenesis in
Arabidopsis Wild Type and the Gf Mutant. *Sexual Plant Reproduction* 10(1): 49–64.
https://doi.org/10.1007/s004970050067.

Clough, S. J. and Bent, A. F. (1998). Floral dip: a simplified method for Agrobacterium -mediated
transformation of Arabidopsis thaliana. *The Plant Journal* 16(6), 735–743.
https://doi.org/10.1046/j.1365-313x.1998.00343.x

Corral, J. M., Vogel, H., Aliyu, O. M., Hensel, G., Thiel, T., Kumlehn, J., & Sharbel, T. F. (2013). A
conserved apomixis-specific polymorphism is correlated with exclusive exonuclease expression in
premeiotic ovules of apomictic Boechera species. *Plant Physiology* 163(4), 1660–1672.
https://doi.org/10.1104/pp.113.222430

D., Gregory Brian. (2018). Shedding some blue light on alternative promoter usage in plants. *PNAS*
115(30), 7654–56. https://doi.org/10.1073/pnas.1809312115.

D'Erfurth, I., Jolivet, S., Froger, N., Catrice, O., Novatchkova, M., & Mercier, R. (2009). Turning meiosis
into mitosis. *PLoS Biology* 7(6), e1000124. https://doi.org/10.1371/journal.pbio.1000124

Van Dijk, Peter J, and T H Noel Ellis. 2023. Gregor Mendel and the Theory of Species Multiplication.
*Genetics* 224(2): iyad046. https://doi.org/10.1093/genetics/iyad046.

Dobeš, Ch., Marcus A. K., Sharbel, T. F. (2006). Embryology, karyology, and modes of reproduction in
the North American genus Boechera (Brassicaceae): A compilation of seven decades of research. *Annals*
*of the Missouri Botanical Garden.* 93(3): 517–34. doi:10.3417/0026-
6493(2007)93[517:EKAMOR]2.0.CO;2

Dossa, K., Zhou, R., Li, D., Liu, A., Qin, Lu., Mmadi, Marie A. et al. (2021). A Novel Motif in the 5'-
UTR of an Orphan Gene 'Big Root Biomass' Modulates Root Biomass in Sesame. *Plant Biotechnology*
*Journal* 19(5): 1065–79. https://doi.org/10.1111/pbi.13531.

Fehrer, J., Gemeinholzer, B., Chrtek, J., Bräutigam, S. (2007). Incongruent plastid and nuclear DNA
phylogenies reveal ancient intergeneric hybridization in Pilosella hawkweeds (Hieracium, Cichorieae,
Asteraceae). *Molecular Phylogenetics and Evolution* 42(2), 347–361.
https://doi.org/10.1016/j.ympev.2006.07.004

Hojsgaard, Diego. (2018). Transient Activation of Apomixis in Sexual Neotriploids May Retain
Genomically Altered States and Enhance Polyploid Establishment. *Frontiers in Plant Science*  9.
https://www.frontiersin.org/article/10.3389/fpls.2018.00230.

Hojsgaard, D., Hörandl, E. (2019). The Rise of Apomixis in natural plant populations. *Frontiers in Plant*
*Science* 10. https://www.frontiersin.org/article/10.3389/fpls.2019.00358.

Horton, Robert M., Hunt, Henry D., Ho, Steffan, N., Pullen, Jeffry K., Pease and Larry, R. (1989). Engineering Hybrid Genes without the Use of Restriction Enzymes: Gene Splicing by Overlap Extension. *Gene* 77(1): 61–68. https://www.sciencedirect.com/science/article/pii/0378111989903594.

Horton, Robert M, Zeling Cai, Steffan N Ho, and Larry R Pease. 2013. Gene Splicing by Overlap Extension: Tailor-Made Genes Using the Polymerase Chain Reaction. *BioTechniques* 54(3): 129–33. https://www.future-science.com/doi/abs/10.2144/000114017.

James, A. B., Sullivan, S., Nimmo, H. G. (2018). Global spatial analysis of Arabidopsis natural variants implicates 5′UTR splicing of LATE ELONGATED HYPOCOTYL in responses to temperature. *Plant, Cell & Environment* 41(7), 1524–38. https://doi.org/10.1111/pce.13188.

Kantama, L., Sharbel, T. F., Schranz, M. E., Mitchell-Olds, T., de Vries, S., de Jong, H. (2007). Diploid apomicts of the Boechera holboellii complex display large-scale chromosome substitutions and aberrant chromosomes. *PNAS* 104(35), 14026–14031. https://doi.org/10.1073/pnas.0706647104.

Kiefer, Ch., Dobeš, ch., Sharbel, T.F., Marcus A. Koch. (2009). Phylogeographic structure of the chloroplast DNA gene pool in North American Boechera – a genus and continental-wide perspective. *Molecular Phylogenetics and Evolution* 52(2), 303–11. https://www.sciencedirect.com/science/article/pii/S1055790309000694.

Kiefer, Ch. and Marcus A Koch. 2012. A Continental-Wide Perspective: The Genepool of Nuclear Encoded Ribosomal DNA and Single-Copy Gene Sequences in North American Boechera (Brassicaceae). *PLOS ONE* 7(5): e36491. https://doi.org/10.1371/journal.pone.0036491.

Koltunow, A. M. G., Johnson, S. D., Rodrigues, J. C. M., Okada, T., Hu, Y., Tsuchiya, T., Wilson, S., et al. (2011). Sexual reproduction is the default mode in apomictic Hieracium subgenus Pilosella, in which two dominant loci function to enable apomixis. *Plant Journal* 66(5), 890–902. https://doi.org/10.1111/j.1365-313X.2011.04556.x

Kurihara, Y., Makita, Y., Kawashima, M., Matsui, M. (2018). Transcripts from Downstream Alternative Transcription Start Sites Evade UORF-Mediated Inhibition of Gene Expression in Arabidopsis. *Proceedings of the National Academy of Sciences* 115(30), 7831–36. https://doi.org/10.1073/pnas.1804971115.

Lovell, J. T., Aliyu, O. M., Mau, M., Schranz, M. E., Koch, M., Kiefer, C., et al. (2013). On the origin and evolution of apomixis in Boechera. *Plant reproduction* 26(4), 309–315. https://doi.org/10.1007/s00497-013-0218-7

J Malamy, J.E., Philip N. Benfey P.N. (1997) Organization and cell differentiation in lateral roots of Arabidopsis thaliana. Development 124 (1): 33–44. https://doi.org/10.1242/dev.124.1.33

Mau, M., Liiving, T., Fomenko, L., Goertzen, R., Paczesniak, D., Böttner, L., et al. (2021). The spread of infectious asexuality through haploid pollen. *New Phytologist* 230(2), 804–820. https://doi.org/10.1111/nph.17174

Mittelsten Scheid, O. (2022). Mendelian and Non-Mendelian Genetics in Model Plants. *The Plant Cell* koac070. https://doi.org/10.1093/plcell/koac070.

Rojek, J., Kapusta, M., Kozieradzka-Kiszkurno, M., Majcher, D., Górniak, M., Sliwinska, E., et al. (2018). Establishing the Cell Biology of Apomictic Reproduction in Diploid Boechera Stricta (Brassicaceae). *Annals of Botany* 122(4): 513–39. https://doi.org/10.1093/aob/mcy114.

Rojek, J., Tucker, M. R., Pinto, S. C., Rychłowski, M., Lichocka, M., Soukupova, H., et al. (2021). Rab-

dependent vesicular traffic affects female gametophyte development in Arabidopsis. *Journal of*
*Experimental Botany* 72, 320–340. https://doi.org/10.1093/jxb/eraa430.

Rushworth, C. A., Song, Bao-Hua,. LEE, Cheng-Ruei and Mitchell-Olds., Th. (2011). Boechera, a Model
System for Ecological Genomics. *Molecular Ecology* 20(23): 4843–57. https://doi.org/10.1111/j.1365-
294X.2011.05340.x.

Rushworth, C. A., Windham, M. D., Keith, R. A., Mitchell-Olds. T. (2018). Ecological differentiation
facilitates fine-scale coexistence of sexual and asexual Boechera. *American Journal of Botany* 105(12),
2051–64. https://doi.org/10.1002/ajb2.1201.

Schmidt, A. (2020). Controlling Apomixis: shared features and distinct characteristics of gene regulation.
*Genes* 11(3), 329. https://pubmed.ncbi.nlm.nih.gov/32245021.

Smyth, D. R., Bowman, J. L., Meyerowitz. E. M. (1990). Early flower development in Arabidopsis. *The*
*Plant cell* 2(8), 755–67. https://pubmed.ncbi.nlm.nih.gov/2152125.

Spillane, Ch., Curtis, M. D., Grossniklaus, U. (2004). Apomixis technology development—Virgin Births
in Farmers' Fields? *Nature Biotechnology* 22(6), 687–91. https://doi.org/10.1038/nbt976.

Srivastava, A. K., Lu, Y., Zinta, G., Lang, Zh., Zhu, J-K. (2018). UTR-Dependent control of gene
expression in pPlants. *Trends in plant science* 23(3), 248–59. https://pubmed.ncbi.nlm.nih.gov/29223924.

Tucker, M. R., Koltunow, A. M. G. (2009). Sexual and Asexual (Apomictic) Seed Development in
Flowering Plants: Molecular, Morphological and Evolutionary Relationships. *Functional Plant Biology*
36(6), 490–504.

Underwood, Ch. J. et al. (2022). A PARTHENOGENESIS Allele from Apomictic Dandelion Can Induce
Egg Cell Division without Fertilization in Lettuce." *Nature Genetics* 54(1), 84–93.
https://doi.org/10.1038/s41588-021-00984-y.

Underwood, Ch. J., Vijverberg, K., Rigola, D., Okamoto, Sh., Oplaat, C., Op den Camp, R. H. M., et al.
(2022). Engineering Apomixis: Clonal Seeds Approaching the Fields. *Annual Review of Plant Biology*.
https://doi.org/10.1146/annurev-arplant-102720-013958.

Vernet, A., Meynard, D., Lian, Q., Mieulet, D., Gibert, O., Bissah, M., et al. (2022). High-Frequency
Synthetic Apomixis in Hybrid Rice. *Nature Communications* 13(1), 7963.
https://doi.org/10.1038/s41467-022-35679-3.

FIGURE 1 Construct pC1, pC2, pC3, pC4, pC5, pC6, and pC7. Constructs pC3 to pC7 created
using SOE (Horton et al. 1989). For pC1, primer pairs of pAPOLLO AF-2 kb and pAPOLLO 1R
were used to amplify 1,945 bp of DNA containing 71 bp of the 5'UTR of the Apo promoter
containing the 20 bp Apo-specific insertion (yellow). For pC2, primer pairs of pAPOLLO SF -2
kb and pAPOLLO 1R were used to amplify 1,970 bp DNA containing 104 bp of the 5'UTR
(gray). For pC3, primer pairs used to amplify 1,880 bp of the Apo promoter are shown in green,
and primer pairs used to amplify 44 bp of the 5'UTR (gray) by deleting the 20 bp Apo-specific
insertion (yellow) are shown in dark-blue. For pC4, primer pairs used to amplify 1,917 bp of the
Sex promoter are shown in green, and primer pairs used to amplify 52 bp of Sex promoter by
incorporating the 20 bp Apo-specific insertion (yellow) in dark-blue, with the 5'UTR of the Sex
allele in gray. For pC5 primer pairs used to amplify 1,880 bp of the Apo promoter are shown in
green, and primer pairs used to amplify 44 bp of the Apo promoter by incorporating a
randomized 20 bp insertion (yellow) in dark-blue, with the 5'UTR of the Apo allele in gray. For
pC6, primer pairs used to amplify 1,870 bp of the Apo-promoter are shown in green, and primer
pairs used to amplify 104 bp of the Sex promoter in dark-blue, with the 5' UTR of the Sex allele
in gray, and the 20 bp Apo-specific insertion (yellow) deleted. For pC7, primer pairs used to
amplify 1,865 bp of the Sex promoter are shown in green, and primer pairs used to amplify 71 bp
5'UTR of the Apo promoter carrying the Apo-specific insertion (yellow) is in dark-blue, with the
5'UTR of the Apo allele in gray color.

FIGURE 2 Expression patterns of Boechera native promoter GUS constructs in Arabidopsis. (A)
GUS distribution of 2 kb native pApo promoter (pC1) in two-week old plant. (B) No GUS
activity observed in 2 kb native pSex promoter (pC2). (A,B) Scale bar=0.5 mm (C) Positive
GUS activity in leaf of six-week old Arabidopsis transgenic for pC1. Scale bar=2.0 mm (Zeiss
Lumar.V12 Stereoscope).

FIGURE 3 Expression patterns of synthetic promoter construct (pC3: GUS) in different
developmental stages of Arabidopsis flowers. (A) 0.7 mm flower. (B) 1.0 mm flower. (C) 1.5
mm flower with GUS activity in filament. (D) 2.0 mm flower with GUS activity in style and
filament. (E) 3.0 mm flower with GUS activity in style and filament. (A-E) Scale bar=1.0 mm
(Zeiss Lumar.V12 Stereoscope), (F) 0.7 mm flower. (G) 1.0 mm flower. (H) 1.5 mm flower
with GUS activity in style and filament. (I) 2.0 mm flower with GUS activity in style and
filament. (J) 3.0 mm flower with GUS activity in style and filament, (F-J) Scale bar=100 µm
(Olympus BX61 microscope).
527
FIGURE 4 Expression patterns of spliced promoter construct (pC5: GUS) in different
developmental stages of Arabidopsis flowers. (A) 0.7 mm flower with GUS activity in anther
and receptacle. (B) 1.0 mm flower with GUS activity in anther and receptacle. (C) 1.5 mm
flower with GUS activity in style and receptacle. (D) 2.0 mm flower with GUS activity in style
and receptacle. (E) 3.0 mm flower with GUS activity in style and receptacle. (A-E) Scale bar=1.0
mm (Zeiss Lumar.V12 Stereoscope). (F) 0.7 mm flower with GUS activity in anthers. (G) 1.0
mm flower with GUS activity in anthers. (H) 1.5 mm flower with GUS activity in style. (I) 2.0
mm flower with GUS activity in style. (J) 3.0 mm flower with GUS activity in style. (G-J) Scale
bar=50 µm

FIGURE 5 Expression patterns of spliced promoter construct (pC6: GUS) in different
developmental stages of Arabidopsis flowers. (A) 0.7 mm flower with GUS activity in anthers,

(B) 1.0 mm flower with GUS activity in anthers, (C) 1.7 mm flower with GUS activity in anthers
(D) 2.0 mm flower with GUS activity in anthers, (E) 3.0 mm flower, (A-E) Scale bar=1.0 mm
(Zeiss Lumar.V12 Stereoscope), (F) Arabidopsis inflorescence, scale bar=2.0 mm, (G) 0.7 mm
flower with GUS activity in anthers, (H) 1.0 mm flower with GUS activity in anthers, (I) 1.5 mm
flower with GUS activity in anthers, (J) 2.0 mm flower with GUS activity in anthers, (K) 3.0 mm
flower, (G-K) Scale bar=50 µm (Olympus BX61 microscope). Pollen-specific GUS expression
demonstrated by black arrows (I, J, K).

FIGURE 6 Expression patterns of promoter construct (pC7: GUS) in different developmental
stages of Arabidopsis flowers. (A) 0.7 mm flower with GUS activity in transmitting tract, ovules
and anthers, (B) 1.0 mm flower with GUS activity in transmitting tract and ovules, (C) 1.5 mm
flower with GUS activity in and transmitting tract and ovules, (D) 2.0 mm flower, (A-D) Scale
bar=1.0 mm (E) The Arabidopsis inflorescence, Scale bar=2.0 mm (Zeiss Lumar.V12
Stereoscope). (F) 1.0 mm flower showing expression in ovules (black arrows) Scale bar=50 µm
(Olympus BX61 microscope).
553