1
2

High quality genome assembly of the brown hare (*Lepus europaeus*) with chromosome-level scaffolding

Craig Michell[1,2], Joanna Collins[3], Pia K. Laine[4], Zsófia Fekete[1,5], Riikka
Tapanainen[1], Jonathan M. D. Wood[3], Steffi Goffart[1] and Jaakko L. O.
Pohjoismäki*[1]

[1]Department of Environmental and Biological Sciences, University of Eastern Finland – Joensuu, Finland
[2]Red Sea Research Center, Biological and Environmental Science and Engineering Division, King Abdullah
University of Science and Technology, – Thuwal, Kingdom of Saudi Arabia
[3]Tree of Life, Wellcome Sanger Institute, Hinxton, Cambridge, UK.
[4]Insitute of Biotechnology, University of Helsinki, Finland.
[5]Hungarian University of Agriculture and Life Sciences, Institute of Genetics and Biotechnology, Gödöllő,
Hungary

18
19   *Corresponding author
20   Correspondence: jaakko.pohjoismaki@uef.fi
21

Abstract

24   We present here a high-quality genome assembly of the brown hare (*Lepus europaeus* Pallas),
25   based on a fibroblast cell line of a male specimen from Liperi, Eastern Finland. This brown hare
26   genome represents the first Finnish contribution to the European Reference Genome Atlas pilot
27   effort to generate reference genomes for European biodiversity.

The genome was assembled using 25X PacBio HiFi sequencing data and scaffolded utilizing a Hi-C chromosome structure capture approach. After manual curation, the assembled genome length was 2,930,972,003 bp with N50 scaffold of 125.8 Mb. 93.16% of the assembly could be assigned to 25 identified chromosomes (23 autosomes plus X and Y), matching the published karyotype. The chromosomes were numbered according to size. The genome has a high degree of completeness based on the BUSCO score (mammalia_odb10 database), Complete: 96.1% [Single copy: 93.1%, Duplicated: 3.0%], Fragmented 0.8%, and Missing 2.9%. The mitochondrial genome of the cell line was sequenced and assembled separately. The final annotated genome has 30,833 genes of which 21,467 code for a polypeptide.

The brown hare genome is particularly interesting as this species readily hybridizes with the mountain hare (*Lepus timidus* L.) at the species contact zone in northern Eurasia, producing fertile offspring and resulting in gene flow between the two species. In addition to providing a useful comparison for population studies, the genome can offer insight into the chromosomal evolution among Glires in general and Lagomorpha in particular. The chromosomal assembly of the genome also demonstrates that the cell line has not acquired karyotypic changes during culture.

Keywords: Lagomorpha, reference genome, chromosome structure, phylogeny, fibroblast cell line

Introduction

The brown hare (*Lepus europaeus* Pallas), also known as the European hare, is a widespread species in the western parts of Eurasia (Bock, 2020). Besides its native range, the brown hare has been introduced to numerous regions, including the British Isles, the Falkland Islands, Canada, South America, Australia and New Zealand (Jaksic, 2023; Petrovan, 2013). In many places the brown hare is regarded as an invasive species and a threat to the local ecosystems (Stott, 2003) or native species (Reid, 2011).

The brown hare is a steppe-adapted species living in open grasslands and avoiding forested regions. Its colonization history in Europe is complex. The species has had glacial refugia during the Pleistocene in the Italian peninsula, the Balkans and Asia Minor (Fickel et al., 2008), with an interesting pre-glacial diversity hotspot in the archipelago of Greece (Minoudi et al., 2018). After the ice age, the species has been expanding its range both naturally as well as facilitated by human-caused changes in the landscape, especially through the expansion of agricultural lands and pastures. As mentioned above, the brown hare has been also frequently introduced by humans to new areas from antiquity to the present day (Petrovan, 2013), resulting in potential mixing of ancestral populations. This is true also for Finland, where the species has arrived naturally from the southeast through the Karelian isthmus in the late 19th century (Ognev, 1940; Siivonen, 1972; Wadén, 1910) but also through local introductions to the southwestern parts of the country in the 1910s (Rikala, 1925; Salonius, 1924; Suomalainen, 1922; Wadén, 1910).

Curiously, Linné was unaware of the existence of the brown hare, despite having described the mountain hare (*Lepus timidus* L.) and the South African Cape hare (*Lepus capensis* L.). In fact, the brown hare is not native to Linné's home country Sweden but was introduced there only a century after his death (Thulin, 2003). Curiously, the species author Peter Simon Pallas never

published a formal description of the brown hare. Instead, the species was authority has been attributed to him because of inclusion of the name *Lepus europaeus* into his identification table of hare species of the world (Pallas, 1778: 30) (Figure 1A). Consequently, there is also no information of the type locality or any other information of the species in the original publication. The type locality of the brown hare has been retrospectively assigned – without too convincing arguments – to Burgundy (France) or Poland (Holden, 2005). Identifying Poland as the type locality was done by the Russian zoologist Sergey Ognev, and is based on Pallas mentioning in the 1778 text the hybrid forms of *Lepus variabilis* (synonym of *L. timidus*) and *L. europaeus* from Poland and Lithuania, as well as citing later works of Pallas for the species distribution (Ognev, 1940). In fact, Ognev (1940: 141) boldly states that "we consider southwestern Poland as a typical terrain for the nominal subspecies; there are hares with the characteristics of the basic form, expressed with full clarity" (translation from Russian).

While being widespread in the temperate regions of Western and Central Europe to the Caspian steppes in historical times, the brown hare began a northward expansion in the 19th century. In the Fennoscandian region, the species reached the St Petersburg area by 1820s (Ognev, 1940), arriving in Finland at the turn of the century (Salonius, 1924; Siivonen, 1972; Thulin, 2003; Wadén, 1910). By 1930, the species had become established throughout southern Finland, with the range expansion stagnating southwest of the N64°-N62° -line until the 1990s. Quite likely benefitting from the ongoing anthropogenic climate change, brown hares range expansion has intensified during the last three decades, currently reaching the polar circle (Levänen et al., 2018a) (Figure 1B), as well as higher altitudes in the Alps (Schai-Braun et al., 2023). In Finland, the expansion seems to be limited with 150 days of annual snow cover (Levänen et al., 2018a). The range expansion together with the past introductions have brought the brown hare increasingly in contact with the mountain hare, especially at the northern edges of the species distribution, constituting a threat especially to the more temperate climate adapted populations and subspecies of the mountain hare (Reid, 2011; Reid and Montgomery, 2007; Thulin, 2003; Thulin et al., 2021). Besides competing, brown hares also hybridize with mountain hares, producing fertile offspring and resulting in gene flow between the species. This gene flow is biased towards the brown hare (Ferreira et al., 2021; Levänen et al., 2018a; Levänen et al., 2018b; Thulin and Tegelström, 2002), which obtains genetic variation from mountain hares, some of which might have adaptive significance (Pohjoismäki et al., 2021). In contrast to the brown hare's expansion towards north, many of the Central European brown hare populations are contracting in range and numbers, likely driven by changes in agricultural practices and land use (Schai-Braun et al., 2013; Smith et al., 2005).

Overall, the brown hare's features as an invasive species outside Europe and expansive species in the north, its hybridization tendency and complex population ancestries make the species highly interesting not only to study population genetics and adaptation mechanisms, but also for the understanding the genomic makeup of species boundaries (Gaertner et al., 2022). These studies would be greatly facilitated by the availability of high quality, chromosomally scaffolded reference genomes (Blaxter et al., 2022).

The last decade has witnessed a boom in genome sequencing technologies, leading to many sequencing initiatives, such as the Earth BioGenome Project (EBP), European Reference Genome Atlas (ERGA) and the Darwin Tree of Life (DToL), aiming to produce high quality reference genomes for a variety of organisms (Blaxter et al., 2022). This has been enabled by the

technological development of high-throughput sequencing of long molecules. The read length N50's > 19 kbp, allows a high contiguity compared to the genome assemblies released in the early 2010's that were based on short read sequencing technologies. Additionally, Hi-C sequencing, a method measuring the contact frequency between all pairs of loci in the genome, has allowed producing pretext maps of chromosomes (Ghurye et al., 2019), enabling chromosome level assemblies with a precision that has been previously achieved only with model organisms (Lawniczak et al., 2022).

In the presented work, we apply the PacBio HiFi long-range sequencing technology with Hi-C sequencing to produce a high-quality reference genome of the brown hare (*Lepus europaeus*). While there is no previous genome assembly for the species, a mountain hare assembly (GCA_009760805) exists, representing the Irish subspecies *L. t. hibernicus* and is based on a female specimen, thus lacking the Y-chromosome. This assembly is also a so-called "pseudoreference", assembled to chromosomes using the rabbit (*Oryctolagus cuniculus* L.) reference genome (Marques et al., 2020). The original rabbit reference genome had been established using whole-genome shotgun sequencing of females representing the inbred Thorbecke New Zealand White line and was quite fragmented (Carneiro et al., 2014), updated only recently with long-read sequencing data (Bai et al., 2021). Notably, the diploid chromosome number in domestic rabbits is 2n = 44 (Korstanje et al., 1999), whereas it is 48 in brown hare and mountain hare (Gustavsson, 1972).

To obtain a reference genome conforming to current standards, we used a fibroblast cell line (LE1) of a male specimen of brown hare (*Lepus europaeus*) from Liperi, Eastern Finland (Gaertner et al., 2022), as a source of high-molecular weight (HMW) DNA and fresh RNA. Compared to solid tissues, fibroblasts are optimal for Hi-C, as the method was originally developed for cells growing in a monolayer (Lieberman-Aiden et al., 2009). Now, over 240 years after the first mention of the name *Lepus europaeus* in the literature (Figure 1A), we have expanded the original seven-word description to encompass 2,930,972,003 genomic letters. These letters have been assembled in a highly complete (BUSCO mammalia_odb10 database, complete: 96.1%), and continuous (N50 scaffold: 125.8 Mb) manner with 93.16% of the sequence being placed in the identified 23 autosomes, and X and Y sex chromosomes. The presented genome assembly not only contains the identity of the brown hare as a species, but also provides a gateway to detailed knowledge of its biology and evolutionary history. Besides being the first Finnish contribution to the European Reference Genome Atlas (Mazzoni et al., 2023), the brown hare reference genome represents a continuing effort to map and understand our planets biodiversity.

Methods

Sampling and confirming of the species identity
A young male brown hare was sampled in October 2018 in Kuorinka, Liperi, Finland (62.6207 N, 29.4478 E, Figure 1B). In Finland, brown hares are highly dependent on similar man-made environments (Levänen et al., 2018a; Levänen et al., 2019), and also the collection location is agricultural area with a mosaic of fields, shrubs and mixed forest with a strong brown hare population and only occasional stray mountain hares. Geographically, the sampled population is part of the same distribution continuum as with species proposed type locality, through Russian

Karelia and the Baltic states to Poland. This is important, as is recommendable that the species
reference genome would represent or be closely related to the type locality population
(Lawniczak et al., 2022).

The sampling had minimal impact on the local brown hare population and no impact on
the habitat. As brown hares are legal game animals in Finland and the hunting followed the
regional hunting seasons and legislation (Metsästyslaki [Hunting law] 1993/615/5§), the sampling
adheres to the ARRIVE guidelines and no ethical assessment for the sampling was required. The
sampling also did not involve International Trade in Endangered Species of Wild Fauna and Flora
(CITES) or other export of specimens, as defined by the Convention on Biological Diversity (CBD).

The species identity was confirmed at site based on the morphological features
distinguishing the brown hare from the mountain hare, the only other hare species in northern
Europe. Further analysis of the coding part of the LE1 genome and mitochondrial DNA
haplotyping showed minimal ancestral admixture with mountain hares (Gaertner et al., 2022). In
fact, unlike many other brown hares in the region, showing adaptive introgression of mountain
hare specific UCP1 alleles (Pohjoismäki et al., 2021), the LE1 specimen is homozygous for the
common ancestral brown hare allele, UCP02.

Generation and vouchering of the cell line
The fibroblast cell line (LE1) was isolated from the specimen as described earlier (Gaertner et al.,
2022) and is deposited as cryopreserved living cells under voucher number ZFMK-TIS-69747 into
the biobank of the Stiftung Leibniz-Institut zur Analyse des Biodiversitätswandels, Zoological
Research Museum Alexander König (ZFMK), Bonn, Germany. Additional identifiers for the sample
are mLepEur2 and ERGA specimen ID ERGA_FI_3610_002 (COPO portal, https://copo-
project.org).

HMW DNA extraction and PacBio HiFi sequencing
A salting out-method was used to extract high molecular weight DNA from cells grown to
confluency on a 10 cm cell culture dish. Briefly, the cells were detached from the dish using
trypsin, followed by centrifugation and two washes with PBS. The salting out-method then
followed the 10X Genomics "Salting Out Method for DNA extraction from cells" protocol (10x-
Genomics, 2017). The high molecular weight DNA was quantified using a Qubit 3.0 (Thermo Fisher
Scientific™, Malaysia), followed by qualification with an 0.8 % Agarose gel. The DNA was sent for
the library preparation and sequencing at the DNA Sequencing and Genomics Laboratory,
Institute of Biotechnology, University of Helsinki on the PacBio Sequel II. In brief, the DNA was
quantified using Qubit and DNA fragment length was measured using Fragment Analyzer (HS large
fragment kit). DNA was then sheared with Megaruptor 3 (Diagenode) (45/200/30) to produce
DNA of length 24 kb. Buffer was then replaced with PacBio's Elution buffer using AMPure beads.
Repair and A-tailing, adapter ligation, nuclease treatment and cleanup with SMRTbell cleanup
beads was then performed with the SMRTbell prep kit 3.0. Larger fragments (greater than 10 000
bp) were then purified with the BluePippin (Sage Science) (0.75% DF Marker S1 high pass 6-10kb
v3) and DNA was further purified with PacBio AMPure beads and treated with DNA damage repair
mix (PacBio) (37°C for 60 mins) and then purified with AMPure clean-up beads and eluted to 11
ul PacBio's Elution buffer. Libraries were sequenced on the PacBio Sequel II at a concentration of
90 pM using PacBio's instructions (provided by the SMRTlink software). Sequence data from two

flow cells was produced. Genomescope2 was used to visualize the HiFi sequencing results
(Ranallo-Benavidez et al., 2020).
204
Mitochondrial DNA (mtDNA) sequencing
The mitochondrial genomes of our hare cell lines (Gaertner et al., 2022) have been sequenced
for the purposes of another study using approximately 2 kb overlapping PCR-amplified fragments
of mtDNA. The primers used to amplify the mountain hare mtDNA were as follows:
Le93F: TTGTTTTGTAGCAAGTTTACACATGC
Le184R: GCTTAATACCTGCTCCTCTTGATCTA
Le1580F: TTAAACCCATAGTTGGCCTAAAAGC
Le1635R: TTGAGCTTTAACGCTTTCTTAATTGA
Le3045F: AGGCGTATTATTTATCCTAGCAACCT
Le3175R: CCTCATAAGAAATGGTCTGTGCGA
Le3921F: CCCCCTAATCTTTTCCATCATCCTAT
Le4482R: TCATCCTATATGGGCAATTGAGGAAT
Le4689F: AGGCTTTATTCCAAAGTGAATTATTATTCA
Le5417R: AGGCTCCAAATAAAAGGTAGAGAGTT
Le6696F: ATACCGTCTCATCAATAGGCTCCTTC
Le6756R: ATAAAGATTATTACTATTACAGCGGTTAGA
Le8603F: AGCCTATATCTACATGATAATACTTAATGA
Le8698R: CGGATAAGGCCCCGGTAAGTGG
Le10552F: TTGAAGCAACACTAATCCCTACACTA
Le10613R: TCGTTCTGTTTGATTACCTCATCGT
Le11301F: ACCATTAACCTTCTAGGAGAGCTTCT
Le11807R: AGGATAATGATTGAGACGGCTATTGA
Le12407F: GTCTAATCCTAGCTGCTACAGGTAAG
Le12791R: GAGCATAAAAAGAGTATAGCTTTGAA
Le14204F: ATTGTTAACCACTCTCTAATCGACCT
Le14514R: CCAATGTTTCAGGTTTCTAGGTAAGT
Lt16056F: TGGGGTATGCTTGGACTCAAC
Le16119R: TCGTCTACAATAAGTGCACCGG
233
In total, 12 separate reactions were prepared to cover the mitochondria genome:
1.   Lt16056F + Le184R: 1871 bp
2.   Le93F + Le1635R: 1543 bp
3.   Le1580F + Le3175R: 1596 bp
4.   Le3045F + Le4482R: 1438 bp
5.   Le3921F + Le5417R: 1497 bp
6.   Le4689F + Le6756R: 2068 bp
7.   Le6696F + Le8698R: 2003 bp
8.   Le8603F + Le10613R: 2011 bp
9.   Le10552F + Le11807R: 1256 bp
10.   Le11301F + Le12791R: 1491 bp
11.   Le12407F + Le14514R: 2108 bp

12.   Le14204F + Le16119R: 1916 bp

(Expected fragment size based on the published *Lepus europaeus* mtDNA sequence from Sweden

[NC_004028.1]).

The fragments were amplified from total DNA preparations using a PCR program with a 1 min

94 °C denaturing step, followed by 35 cycles of 94 °C for 15 s, 56 °C for 15 s and 72 °C for 2 min

and a final 3 min elongation step at 72 °C. The obtained products were gel purified using the

GeneJET gel extraction kit (Thermo Fisher Scientific™) and sent for sequencing using Illumina

MiniSeq at the Genome Center of Eastern Finland.

The sequence of the non-coding region-containing PCR fragment (Lt16056F + Lt184R) was further

validated by Sanger sequencing, applying also the following additional sequencing primer:

Le101F: TATAAATTCCTGCCAAACCCCAAAAA

Figure 1. The brown hare. (A) An excerpt from the identification table by Pallas (1778: 30), which

has been assigned as the species description for *Lepus europaeus*. "Tips of the ears longer than

the head, tail black above." (translation from Latin). (B) The geographic location of the LE1 sample

used in this study. The grey-shaded area shows the approximate current distribution of the

brown hare in Finland, based on the data in the Finnish Biodiversity Information Facility (FinBIF,

https://laji.fi/en). Individual records from the north are from settlemens or towns. The arctic

circle (black line) runs along the 66.6 ° parallel. Background photo shows a typical Finnish male

brown hare. Photo taken in Joensuu, 20 km East from the LE1 sampling site by Dr. Mervi
Kunnasranta.
270
Hi-C library preparation
Hi-C sequencing libraries were prepared following the protocol of (Belaghzal et al., 2017) with the
following changes: 1.) To prepare a diverse sequencing library, we performed the Hi-C protocol in
triplicates, 2.) Size fractionation was performed using the NEBNext Ultra II FS DNA module (New
England Biolabs), 3.) Library preparation was performed using the NEBNext Ultra II library
preparation kit for Illumina (New England Biolabs) and 4.) Library enrichment was performed
using triplicate PCR reactions with six cycles of PCR. The PCR reactions were then purified using
Ampure XP beads (Beckman Coulter Life Sciences) at a ratio of 0.9X. The final clean libraries were
quantified using Qbit, followed by agarose gel electrophoresis to confirm the fragment size. The
sequencing was performed on a single lane of the Illumina NovaSeq 6000 using the SP flowcell
with paired-end chemistry 2 x 150bp.
282
Genome assembly
HiFiasm version 0.18.7 (Cheng et al., 2021) was used to assemble the 25X (rq >0.99) PacBio HiFi
reads using the recommended arguments *-l3 –h1 and –h2 –primary* to integrate the Hi-C read
data and produce a primary assembly. We then continued with the scaffolding of the primary
assembly and annotation. To process the HiC data, we first mapped the Hi-C data to the primary
genome assembly using BWA-mem version 0.7.17 (Li and Durbin, 2009) with the arguments *-SP*.
The mapped reads were then parsed and filtered using pairtools version 1.0.2 (Open2C et al.,
2023). To parse the aligned Hi-C reads, we used the options *–no-flip –dd-columns mapq –walks-*
policy mask. The parsed pair data was then sorted and deduplicated using default arguments with
pairtools. Finally, we selected unique-unique and unique-rescued pairs and split these into the
pairs file and bam file for input in YaHS version 1.1 (Zhou et al., 2023). YAHS was run using the
default parameters with the primary contig assembly and the filtered Hi-C bam file. HiFiasm
operates as a haplotype-aware assembler, counting and assembling unique k-mers. Utilizing k-
mer coverage information, the assembler identifies regions in the genome with expected
heterozygous coverage and subsequently amalgamates them into a unified sequence. To refine
the assembly further, Hi-C data is incorporated to elucidate the phasing of variants and ascertain
connected genomic regions. Consequently, the assembler's output provides a primary assembly
along with an alternative assembly, effectively separating the haplotypes by their distinct alleles
within the genomic content (please see https://lh3.github.io/2021/04/17/concepts-in-phased-
assemblies). Contiguity and general genome statistics were calculated using QUAST version 5.2.0
(Mikheenko et al., 2018). We assessed the completeness of the genome by calculating the
number of complete single copy orthologs with BUSCO version 5.1.2 (Manni et al., 2021), using
the mammalia_odb10 database as well as the more lineage specific glires_odb10 database.
306
Genome annotation
Repeat annotation of the genome was performed with EDTA version 2.1.0 (Ou et al., 2019), a *de*
novo repeat identification pipeline. Using the repeat library produced by EDTA, we masked the
scaffolded genome using RepeatMasker version 4.1.1 (Smit et al., 2013-2015). RNA-seq data from
the same cell line (SRA accession number: SRR18740842) as well as RNA-seq data from other *L.*

timidus libraries (SRA accession number: SRR10020054, SRR10020055, SRR10020060,
SRR10491719, SRR18740839, SRR18740840, and SRR18740841) was collected from the sequence
read archive (SRA). The previously produced RNA-seq data (Gaertner et al., 2022) was trimmed
using fastp version 0.23.2 (Chen et al., 2018) and mapped against the masked genome using
HISAT2 version 2.1.0 (Kim et al., 2019) with the default parameters. Furthermore, we included
the protein sequences of the assembled transcripts from this cell line as further evidence for the
genome annotation. These lines of gene evidence were included in the annotation using BRAKER3
version 3.02 (Bruna et al., 2021). The original annotation has been further augmented by the NCBI
Eukaryotic Genome Annotation Pipeline (https://www.ncbi.nlm.nih.gov/genome/annotation_
euk/process/). Telomeric sequences [AACCCT]n were identified using a Telomere Identification
toolkit (tidk) version 0.2.31. The telomeric sequence copy number was then calculated in
windows of 200kb for visualization using Circos (Krzywinski et al., 2009).
324
Mitochondrial DNA assembly and annotation
Mitochondrial DNA was assembled from the PCR-amplified and Illumina sequenced mtDNA using
the MitoZ pipeline (Meng et al., 2019). After comparing the results of the pipeline's outputs with
different kmer options, we selected the best assembly as final. Run options used in the final
assembly were --clade Chordata –fastq_read_length 150, --requiring_taxa Chordata --
genetic_code 2 --kmers_megahit 21 29 39 59 79 99 119 141. The tools invoked by the pipeline
included fastp (Chen et al., 2018) for cleaning the raw data, MEGAHIT (Li et al., 2015) for assembly,
after which sequences were filtered to ensure the correct taxa by HMMER (Wheeler and Eddy,
2013) and further filtered for completeness of protein-coding genes. Annotation steps were done
using TBLASTN (Gertz et al., 2006), GeneWise (Birney et al., 2004) and MiTFi (Juhling et al., 2012).
Final manual curation, the annotation of the non-coding region (NCR) as well as the illustration
of the mitochondrial genome was done using Geneious® 10.2.6 (Biomatters. Available from
https://www.geneious.com). The functional loci on the NCR were identified based on the
similarity with the human (NC_012920) and mouse (FJ374652) NCR sequences.
339
340 **Manual curation**
341 The assembled and annotated genome was manually curated to further improve its quality based
342 on Hi-C contact maps as described in (Howe et al., 2021), allowing to break or join erroneus
343 scaffold assemblies and removing duplicated contigs.
344
345 **Comparison with previous assembly**
346 We performed a comparison of our scaffolded assembly to the current *L. timidus* genome
347 assembly (NCBI Accession number: GCA_009760805). Mapping to the genome was performed
348 using minimap2 version 2.21 (Li, 2018) with the arguments *-asm5*. A dot plot of the alignment
349 was created using the R script pafCoordsDotPlotly.R (https://github.com/tpoorten/dotPlotly).
350

Results

352
The genome assemblies can be accessed via BioSample accession SAMN33984520 as well as
BioProject accession PRJNA1009711 for the primary assembly and PRJNA1009710 for the

alternative assembly. The Final annotated assembly can be found under NCBI Reference
Sequence assembly accession GCF_033115175.1, which can be viewed also using the NCBI
Genome Data Viewer
(https://www.ncbi.nlm.nih.gov/genome/gdv/browser/genome/?id=GCF_033115175.1).
359

Genome assembly

The expected haploid genome size of the *L. europaeus* is similar to *L. timidus*, which has been
estimated to be 3.25 pg (3,178,500,000 bp) (Vinogradov, 1998), containing 23 autosomal
chromosomes and two sex chromosomes (Gustavsson, 1972). PacBio HiFi sequencing with two
flow cells resulted in 25 X coverage of the expected genome size. The sequence N50 of the HiFi
data was 19.97 kb. When k-mer size of 21 is used, the resulting genome size from the PacBio HiFi
data is 2.4 Gbp (Figure 2A). The Illumina sequencing of the Hi-C data produced 494,285,585 paired
reads, representing about 51 X coverage of the genome, with a duplication rate of 17 %. Assembly
with HiFiasm yielded a contig assembly of 2.96 Gbp made up of 671 contigs with a contig N50 of
58 Mbp. The longest contig was 149 Mbp. Using the uniquely mapped Hi-C data, we were able to
scaffold the contigs and fix misassembled contigs (Figure 2B). Although the genome size (2.96
Gbp) and the largest scaffold (149 Mbp) were similar to the contig assembly, the scaffold N50 was
increased to 124 Mbp (N=10) after the Hi-C scaffolding. The obtained scaffold N90 was 21 Mbp
(N=25). The BUSCO scores of the *L. europaeus* assembly suggest it to be near-complete, with the
following results (%- of the total genome): Complete: 96.1% [Single copy: 93.1 %, Duplicated: 3.0
%], Fragmented 0.8 %, and Missing 2.9 % based on the mammalia_odb10 database.
To further improve the assembly, manual curation was performed, resulting in 76 scaffold
breaks, 81 joins and removal of 62 erroneously duplicated contigs (Table 1). These interventions
led to an increase in scaffold N50 of 1.69 % from 123.7 Mb to 125.8 Mb, and a reduction in scaffold
count of 9.21 % from 716 to 650. Of the finalized assembly, 93.16 % could be assigned to 25
identified chromosomes (23 autosomes plus X and Y), matching the known karyotype.
Chromosomes were numbered according to size (Table 2). The curated genome length was
2,930,972,003 bp.

Figure 2. A) Genomescope2 profile of the PacBio HiFi data. B) YaHS pretext map of Hi-C scaffolding after manual curation. C) Dot plot comparison with the mountain hare pseudoreference (GCA_009760805) assembly with rabbit chromosomal assignments. Each grey horizontal line represents a contig end. Names of the non-chromosomal contigs have been omitted for clarity. Rabbit chromosomes corresponding two hare chromosomes as well as the X chromosome with an inversion are are highlighted.

392 **Table 1.** The brown hare reference genome chromosome assignment and assembly statistics

Chromosome assignment	
Chr length	2,730,543,680 bp
Autosomes	23
X	1
Y	1
Unlocalised	19

Statistics for primary assembly after manual curation	
# scaffolds	650
Total scaffold length	2,930,972,003
Average scaffold length	4,509,187.70
Scaffold N50	125,776,599
Scaffold auN	117,151,857.99
Scaffold L50	10
Largest scaffold	184,151,585
Smallest scaffold	1000
# contigs	1068
Total contig length	2,930,889,045
Average contig length	2,744,278.13
Contig N50	43,409,466
Contig auN	47,572,960.95
Contig L50	21
Largest contig	141,329,000
Smallest contig	1000
# gaps in scaffolds	418
Total gap length in scaffolds	82,958
Average gap length in scaffolds	198.46
Gap N50 in scaffolds	200
Gap auN in scaffolds	199.70
Gap L50 in scaffolds	208
Largest gap in scaffolds	200
Smallest gap in scaffolds	31
Base composition	
A	821,587,687
C	645,897,412
G	644,604,632
T	818,799,314

GC content	44.03 %
# soft-masked bases	0
# segments	1,068
Total segment length	2,930,889,045
Average segment length	2,744,278.13
# gaps	418
# paths	650

394 **Table 2.** Chromosome assemblies of the brown hare reference genome with their corresponding
395 GenBank access numbers.

Chromosome	GenBank	Size (bp)	GC content (%)
1	CM065551.1	184,151,585	42.0
2	CM065552.1	176,467,698	42.0
3	CM065553.1	169,342,607	42.0
4	CM065554.1	166,606,208	43.0
5	CM065555.1	159,335,674	46.0
6	CM065556.1	132,443,831	42.0
7	CM065557.1	132,293,278	44.0
8	CM065558.1	130,587,543	41.0
9	CM065559.1	125,776,599	44.5
10	CM065560.1	125,704,418	46.0
11	CM065561.1	117,559,561	44.0
12	CM065562.1	105,667,338	44.0
13	CM065563.1	100,925,866	44.0
14	CM065564.1	99,301,477	44.0
15	CM065565.1	95,895,056	42.0
16	CM065566.1	91,637,918	42.5
17	CM065567.1	84,898,765	45.5
18	CM065568.1	77,563,730	48.5
19	CM065569.1	72,700,603	47.5
20	CM065570.1	69,331,453	45.5
21	CM065571.1	60,609,710	47.5
22	CM065572.1	44,025,767	47.0
23	CM065573.1	30,716,498	53.0
X	CM065574.1	135,183,192	40.5
Y	CM065575.1	33,560,193	41.0
MT	OR915849.1	17,540	38.6

Genome annotation

EDTA produced a curated custom repeat library of 7,045 repetitive elements. Interestingly, the proportion of the genome masked as repetitive elements was higher than expected at 46.8 % of the genome. In the mountain hare pseudoreference genome assembly (GCA_009760805), only 28 % of the genome was masked, and similarly the *k*-mer based estimate of the repetitive elements was 26.7 %. To annotate the genome, we mapped ~1.6 billion RNA-seq reads from previously published RNA-seq libraries, including this cell line. The average mapping rate of all libraries was 89.95 %, which we consider a good indicator of the quality of this genome. Using BRAKER3, we were able to annotate 19,906 gene models in the genome. We then ran BUSCO on the predicted genes and the results show a good level of completeness: Complete: 81.1 % [Single copy: 63.3 %, Duplicate: 17.8 %], Fragmented: 1.1 %, Missing: 17.8 %, Total: 9226 when compared to the mammalia_odb10 database. The final annotation (NCBI RefSeq GCF_033115175.1-RS_2024_01) contains 30,833 genes of which 21,467 code for a polypeptide. Telomeric sequences on both ends of the chromosome were detected on 13 chromosomes (Figure 3A). Furthermore, telomeric sequences were found in high copy number throughout the chromosome length, a feature previously noted using FISH in mountain hares (Forsyth et al., 2005).

Comparison to the mountain hare assembly

Minimap2 was able to align 97.3 % of the contig sequences from the mountain hare pseudoreference genome assembly (GCA_009760805) to this genome assembly (Figure 2C). As pointed out earlier, the mountain hare genome assembly has been scaffolded against the rabbit reference genome (Giska et al., 2022), which has a different chromosome count (Beklemisheva et al., 2011) and explains the few notable inversions and structural differences between the two genomes (Figure 2C). For example, the chr7 and chr 12 assemblies of the brown hare have a good degree of synteny with the mountain hare pseudoreference (= rabbit) chr1, with the chr12 being in inverted orientation. Similarly, the brown hare chr13 and chr16 assemblies have a good synteny and orientation with the rabbit chr2.

Mitochondrial genome

While it is possible to recover the entire mitochondrial genome from the HiFi sequencing data (Uliano-Silva et al., 2023), we had already previously sequenced the mitochondrial genomes of our hare cell lines for the purposes of another study. The assembly of the Illumina data provided a 16,836 bp mtDNA sequence, which was complete and circular, with a 142 bp non-repetitive overlap between the 3´ and 5´ ends of the original assembly removed from the final version. All the expected vertebrate mitochondrial genes, 37 in total, were found (Figure 3B).

Interestingly, the primary assembly of the Illumina data was notably smaller than expected based on the original PCR amplicons, with a notable difference between the expected and obtained non-coding region (NCR) length. Sanger sequencing of the PCR-fragment containing the NCR (Lt16056F + Le184R) revealed that it contained a sequence run consisting of six head-to-tail repeats of a 140 bp element between OriH and tRNA-Phe, explaining the incorrect assembly of the NCR sequence from the short-read Illumina data. With the addition of the repeat elements, the final mitochondrial genome size of our specimen is 17,540 bp.

Based on the sequence similarity with human and mouse mitochondrial genomes, we were able to tentatively identify the termination associated sequence (TAS, 2 nt difference to

human), required for the generation of the 7S DNA and thus the displacement loop (D-loop)
structure on the NCR (Crews et al., 1979). Interestingly, assuming that the 7S DNA is roughly the
same size as in other mammals (600-650 nt), its 5'-end would map to the beginning of another
highly repetitive 240 bp region, likely to have structural significance. This is a notably longer
sequence run than the tRNA-like sequence reported at the replication origin of *L. timidus* mtDNA
(Melo-Ferreira et al., 2014). We have preliminary assigned this structure to correspond to the
heavy-strand origin of replication (OriH), supported by the fact that a similar, tRNA sequence-
derived hairpin structure is required for the priming of the origin of the light-strand replication
(OriL) (Fuste et al., 2010). Experimental validation of the proposed OriH is required.
450
451

452
453 **Figure 3.** A reference genome for the brown hare. A) A size map of 23 autosomes as well as X and
454 Y sex chromosomes of *Lepus europaeus*. Telomeric sequences and their relative lengths are
455 indicated on the inner circle. B) A schematic illustration of the mountain hare mitochondrial
456 genome. Genes encoded by the H-strand are placed inside and the L-strand genes outside of the
457 circle. NCR = non-coding region.

Discussion

Given the challenges of assembling a large mammalian genome, we have produced a high-quality
genome assembly for the brown hare. The genome assembly has a high degree of contiguity
(Contig N50 43 Mbp) and completeness (BUSCO complete 96.1 %). Interestingly, the genome
contains a high number of repetitive elements, a fact that warrants further investigation to
elucidate their identity. While vast majority of the sequences seem to come from retroelements
or endoretroviruses, we cannot fully exclude that some multicopy host genes could be masked.
A detailed analysis of the repeat element libraries from this and another vertebrate genome
assembly releases might be warranted.

The genome assembly is in chromosome scale, and all 23 autosomes as well as X and Y
chromosomes could be scaffolded. Compared with the closely related mountain hare
pseudoreference assembly, the brown hare genome assembly is much more contiguous (Table 1)
and inclusion of telomeric repeats on both ends of several chromosomes (Figure 3A) further
demonstrates the telomere-to-telomere continuity of these assemblies. The presented assembly
also resolves the expected chromosome structure for the species (Figure 2B, C), which is
informative regarding the karyotype difference between hares (n = 24) and rabbits (n = 22).
Comparison of the brown hare assembly with the mountain hare pseudoreference chromosomes
shows that the rabbit chromosome 1 corresponds to the brown hare chromosomes 12 and 17,
with the chromosomal arm corresponding to the former is in an inversed orientation. Similarly,
the rabbit chromosome 2 corresponds to the hare chromosomes 13 and 16, both in the same
orientation. It should also be noted that as the hare chromosomes were numbered according to
their size, the numbering in general does not match those of the rabbit chromosomes (Figure 2C)

The obtained 17,540 bp long mitochondrial genome assembly also provides some
interesting aspects of mtDNA sequence variation within mammalian species. The size difference
to human (16,569 bp) and mouse (16,298 bp) mitochondrial genomes is caused by a rather long
non-coding region (2,102 bp vs. 1,123 bp and 879 bp in humans and mice, respectively), longer
rRNA and tRNA genes, as well as additional non-coding nucleotides between genes. The
mitochondrial genome of our specimen is slightly smaller than the 17,734 bp previous NCBI
Reference Sequence for brown hare (NC_004028.1) from Sweden. The size difference is mostly
due to a copy number difference in tandem repeat elements within the non-coding region, with
seven in the NC_004028.1 and six in our specimen. Indels elsewhere in the NCR explain the rest
of the difference.

The main technical challenge for highly continuous, good quality reference genome
assemblies is the availability of relatively large quantities of intact HMW DNA for HiFi sequencing.
This typically requires snap freezing of tissue samples and assuring an intact cold chain of below
-80 °C for their sending, handling, and long-term storage (Januszczak and Holroyd, 2023). While
development in sequencing technologies can reduce the required DNA amounts, obtaining high
sequencing coverage over a large vertebrate genome is easier when material is plentiful. Similarly,
RNA is very sensitive to degradation in post-mortem tissue samples, also requiring immediate
preparation or snap-freezing to be suitable for RNA-seq. Although transcript data is not obligatory,
it is highly useful for the purpose of genome annotation. The requirement of large samples of
fresh tissue for -omics purposes complicates the sampling of most vertebrates species. For
example, in Finland all land vertebrates are protected by law (animal protection law 247/1996,
hunting law 615/1993), with very few exceptions, such as game animals and pest species.
Capturing and lethal sampling is especially problematic for endangered species. Not only will the
sampling require extensive permits, but it is also difficult to justify and conduct ethically. Here,
the generation of primary fibroblast cell lines can offer a solution. Although the source specimen
for the cell line utilized in this study was hunted, fibroblasts can be isolated from relatively small
skin biopsies, such as ear clippings (Seluanov et al., 2010), with minimally invasive sampling and
harm to the individual. These cells can be expanded in tissue culture and, when immortalized,
provide a highly scalable source of fresh DNA and RNA. Furthermore, living cells can be stored as
frozen stocks in biobanks for decades (Fazekas et al., 2017). Tissue sampling can be done directly
into the growth medium in ambient temperature, allowing several days for the transport to the

laboratory without a need for dry ice or N_2-cryocontainers, which greatly simplifies collection and
logistics. The same applies to the isolated cells. As an example, we sent living cell suspensions
from Joensuu, Finland to the ZFMK biobank in Bonn, Germany, over regular airmail in ambient
temperature. The receiving laboratory can then amplify the cells by culturing and cryopreserve
them in sufficient replicate stocks. It is worth to point out that the obtained genome assembly
scaffolded to 23 autosomes, X and Y chromosomes, as expected for the species, shows no
evidence of karyotypic drift caused by cell culture, dispelling possible concerns for the use of cell
lines for such work (Gaertner et al., 2022).
The brown hare is an iconic European mammal, locally familiar to a wider public of nature-
goers, farmers and hunters, while being at the same time an invasive species on many other
continents. It has populations with complex ancestral makeup due to isolated Pleistocene refugia,
frequent introductions and cross-species hybridization, making it an exceptionally interesting
mammal for genetic studies. Furthermore, depending on the local environmental context, brown
hares can be both an endangered as well as a rapidly expanding species, prompting opportunities
for both conservation as well as invasion genetics studies. A high-quality reference genome
assembly will allow us also to peer deeper into genome structure and chromosome evolution,
which will be particularly interesting from the viewpoint of genetic introgression (Juric et al.,
2016) and maintenance of species boundaries. Any issues of genetic compatibility between
species should leave their fingerprint in the genomes (Skov et al., 2023), providing interesting
insight into the speciation mechanisms (Wolf et al., 2010). Having a high-quality genome for both
species has important implications for the fields of evolutionary and conservation biology. Due to
the high contiguity of the genome assembly, it will be possible to perform accurate analyses of
linkage disequilibrium as well as the identification of runs of homozygosity (ROH). Accurate ROH
analysis is useful for identifying and understanding the impact of inbreeding burden on the
genome of increasingly threatened and isolated species, as quantitative traits can be associated
with ROH burden (Ceballos et al., 2018). Furthermore, this information might provide important
insights into the hybridization of the brown hare with the mountain hare, as well as help to
pinpoint genomic regions that might be helping the species to adapt to local environmental
conditions during the ongoing range expansion (Pohjoismäki et al., 2021). Other *Lepus* reference
genome assemblies available soon will help to complete this picture. Together, this genome
assembly and the mountain hare genome assembly will provide a solid base for future studies.
543

Acknowledgements

We acknowledge the DNA Sequencing and Genomics Laboratory, Institute of Biotechnology,
University of Helsinki for the PacBio and Hi-C sequencing, with special thanks to Dr Petri Auvinen
and Dr Martyn James for helpful discussions and personal interest towards the project. Dr Seppo
Helisalmi and Dr Joose Raivo from the Gene diagnostics lab at Genome Center of Eastern Finland
are thanked for providing the mtDNA sequencing service. We are grateful to the European
Reference Genome Atlas (ERGA) for the opportunity to include our mountain hare genome into
their pilot project. Dr Ann McCartney is thanked for her kind help throughout the process and Mr
Jukka Pusa for his contribution in obtaining the brown hare specimen. We thank Ms Anita

Data, scripts, code, and supplementary information availability

The genome assembly is available from the NCBI database under BioProject ID PRJNA950335. The cell line is available through the corresponding author as well as from the ZFMK biobank.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Funding

This study belongs to the xHARES consortium funded by the R'Life initiative of the Academy of Finland, grant no. 329264.

References

10x-Genomics, 2017. Salting out method for DNA extraction from cells. Sample Preparation Demostrated Protocol. CG00016. Rev A., 1-5.

Bai, Y., Lin, W., Xu, J., Song, J., Yang, D., Chen, Y.E., Li, L., Li, Y., Wang, Z., Zhang, J., 2021. Improving the genome assembly of rabbits with long-read sequencing. Genomics 113, 3216-3223.

Beklemisheva, V.R., Romanenko, S.A., Biltueva, L.S., Trifonov, V.A., Vorobieva, N.V., Serdukova, N.A., Rubtsova, N.V., Brandler, O.V., O'Brien, P.C., Yang, F., Stanyon, R., Ferguson-Smith, M.A., Graphodatsky, A.S., 2011. Reconstruction of karyotype evolution in core Glires. I. The genome homology revealed by comparative chromosome painting. Chromosome Res 19, 549-565.

Belaghzal, H., Dekker, J., Gibcus, J.H., 2017. Hi-C 2.0: An optimized Hi-C procedure for high-resolution genome-wide mapping of chromosome conformation. Methods 123, 56-65.

Birney, E., Clamp, M., Durbin, R., 2004. GeneWise and Genomewise. Genome Res 14, 988-995.

Blaxter, M., Archibald, J.M., Childers, A.K., Coddington, J.A., Crandall, K.A., Di Palma, F., Durbin, R., Edwards, S.V., Graves, J.A.M., Hackett, K.J., Hall, N., Jarvis, E.D., Johnson, R.N., Karlsson, E.K., Kress, W.J., Kuraku, S., Lawniczak, M.K.N., Lindblad-Toh, K., Lopez, J.V., Moran, N.A., Robinson, G.E., Ryder, O.A., Shapiro, B., Soltis, P.S., Warnow, T., Zhang, G.J., Lewin, H.A., 2022. Why sequence all eukaryotes? P Natl Acad Sci USA 119.

Bock, A., 2020. Lepus europaeus (Lagomorpha: Leporidae). Mammalian Species 52, 125-142.

Bruna, T., Hoff, K.J., Lomsadze, A., Stanke, M., Borodovsky, M., 2021. BRAKER2: automatic
eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein
database. NAR Genom Bioinform 3, lqaa108.
Carneiro, M., Rubin, C.J., Di Palma, F., Albert, F.W., Alfoldi, J., Martinez Barrio, A., Pielberg, G.,
Rafati, N., Sayyab, S., Turner-Maier, J., Younis, S., Afonso, S., Aken, B., Alves, J.M., Barrell, D.,
Bolet, G., Boucher, S., Burbano, H.A., Campos, R., Chang, J.L., Duranthon, V., Fontanesi, L.,
Garreau, H., Heiman, D., Johnson, J., Mage, R.G., Peng, Z., Queney, G., Rogel-Gaillard, C.,
Ruffier, M., Searle, S., Villafuerte, R., Xiong, A., Young, S., Forsberg-Nilsson, K., Good, J.M.,
Lander, E.S., Ferrand, N., Lindblad-Toh, K., Andersson, L., 2014. Rabbit genome analysis
reveals a polygenic basis for phenotypic change during domestication. Science 345, 1074-
1079.
Ceballos, F.C., Joshi, P.K., Clark, D.W., Ramsay, M., Wilson, J.F., 2018. Runs of homozygosity:
windows into population history and trait architecture. Nat Rev Genet 19, 220-234.
Chen, S.F., Zhou, Y.Q., Chen, Y.R., Gu, J., 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 34, 884-890.
Cheng, H., Concepcion, G.T., Feng, X., Zhang, H., Li, H., 2021. Haplotype-resolved de novo
assembly using phased assembly graphs with hifiasm. Nat Methods 18, 170-175.
Crews, S., Ojala, D., Posakony, J., Nishiguchi, J., Attardi, G., 1979. Nucleotide sequence of a region
of human mitochondrial DNA containing the precisely identified origin of replication.
Nature 277, 192-198.
Fazekas, J., Grunt, T.W., Jensen-Jarolim, E., Singer, J., 2017. Long term storage in liquid nitrogen
leads to only minor phenotypic and gene expression changes in the mammary carcinoma
model cell line BT474. Oncotarget 8, 35076-35087.
Ferreira, M.S., Jones, M.R., Callahan, C.M., Farelo, L., Tolesa, Z., Suchentrunk, F., Boursot, P., Mills,
L.S., Alves, P.C., Good, J.M., Melo-Ferreira, J., 2021. The Legacy of Recurrent Introgression
during the Radiation of Hares. Syst Biol 70, 593-607.
Fickel, J., Hauffe, H.C., Pecchioli, E., Soriguer, R., Vapa, L., Pitra, C., 2008. Cladogenesis of the
European brown hare (Lepus europaeus Pallas, 1778). Eur J Wildlife Res 54, 495-510.
Forsyth, N.R., Elder, F.F., Shay, J.W., Wright, W.E., 2005. Lagomorphs (rabbits, pikas and hares) do
not use telomere-directed replicative aging in vitro. Mech Ageing Dev 126, 685-691.
Fuste, J.M., Wanrooij, S., Jemt, E., Granycome, C.E., Cluett, T.J., Shi, Y., Atanassova, N., Holt, I.J.,
Gustafsson, C.M., Falkenberg, M., 2010. Mitochondrial RNA polymerase is needed for
activation of the origin of light-strand DNA replication. Mol Cell 37, 67-78.
Gaertner, K., Michell, C., Tapanainen, R., Goffart, S., Saari, S., Soininmaki, M., Dufour, E.,
Pohjoismaki, J.L.O., 2022. Molecular phenotyping uncovers differences in basic
housekeeping functions among closely related species of hares (Lepus spp., Lagomorpha:
Leporidae). Mol Ecol.
Gertz, E.M., Yu, Y.K., Agarwala, R., Schaffer, A.A., Altschul, S.F., 2006. Composition-based statistics
and translated nucleotide searches: improving the TBLASTN module of BLAST. BMC Biol 4,
41.
Ghurye, J., Rhie, A., Walenz, B.P., Schmitt, A., Selvaraj, S., Pop, M., Phillippy, A.M., Koren, S., 2019.
Integrating Hi-C links with assembly graphs for chromosome-scale assembly. PLoS Comput
Biol 15, e1007273.

Giska, I., Pimenta, J., Farelo, L., Boursot, P., Hacklander, K., Jenny, H., Reid, N., Montgomery, W.I.,
Prodohl, P.A., Alves, P.C., Melo-Ferreira, J., 2022. The evolutionary pathways for local
adaptation in mountain hares. Mol Ecol 31, 1487-1503.

Gustavsson, I., 1972. Mitotic and meiotic chromosomes of the variable hare (Lepus timidus L.),
the common hare (Lepus europaeus Pall.) and their hybrids. Hereditas 67, 27-34.

Holden, M.E., 2005. Family Gliridae, in: Wilson, D.E., Reeder, D.M. (Eds.), Mammal Species of the
World: A Taxonomic and Geographic Reference. Johns Hopkins University Press, Baltimore,
USA, pp. 819-841.

Howe, K., Chow, W., Collins, J., Pelan, S., Pointon, D.-L., Sims, Y., Torrance, J., Tracey, A., Wood, J.,
2021. Significantly improving the quality of genome assemblies through curation.
GigaScience 10.

Jaksic, F.M., 2023. Historical ecology and current status of the European hare Lepus europaeus in
South America: a new species in new countries. Stud Neotrop Fauna E.

Januszczak, I., Holroyd, N., 2023. DToL Tissue and Blood Sampling Standard Operating Procedure:
Chordata: Vertebrata (general). protocols.io
dx.doi.org/10.17504/protocols.io.n92ldmjool5b/v1.

Juhling, F., Putz, J., Bernt, M., Donath, A., Middendorf, M., Florentz, C., Stadler, P.F., 2012.
Improved systematic tRNA gene annotation allows new insights into the evolution of
mitochondrial tRNA structures and into the mechanisms of mitochondrial genome
rearrangements. Nucleic Acids Res 40, 2833-2845.

Juric, I., Aeschbacher, S., Coop, G., 2016. The Strength of Selection against Neanderthal
Introgression. PLoS Genet 12, e1006340.

Kim, D., Paggi, J.M., Park, C., Bennett, C., Salzberg, S.L., 2019. Graph-based genome alignment
and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol 37, 907-915.

Korstanje, R., O'Brien, P.C., Yang, F., Rens, W., Bosma, A.A., van Lith, H.A., van Zutphen, L.F.,
Ferguson-Smith, M.A., 1999. Complete homology maps of the rabbit (Oryctolagus
cuniculus) and human by reciprocal chromosome painting. Cytogenet Cell Genet 86, 317-
322.

Krzywinski, M., Schein, J., Birol, I., Connors, J., Gascoyne, R., Horsman, D., Jones, S.J., Marra, M.A.,
2009. Circos: an information aesthetic for comparative genomics. Genome Res 19, 1639-
1645.

Lawniczak, M.K.N., Durbin, R., Flicek, P., Lindblad-Toh, K., Wei, X., Archibald, J.M., Baker, W.J.,
Belov, K., Blaxter, M.L., Marques Bonet, T., Childers, A.K., Coddington, J.A., Crandall, K.A.,
Crawford, A.J., Davey, R.P., Di Palma, F., Fang, Q., Haerty, W., Hall, N., Hoff, K.J., Howe, K.,
Jarvis, E.D., Johnson, W.E., Johnson, R.N., Kersey, P.J., Liu, X., Lopez, J.V., Myers, E.W.,
Pettersson, O.V., Phillippy, A.M., Poelchau, M.F., Pruitt, K.D., Rhie, A., Castilla-Rubio, J.C.,
Sahu, S.K., Salmon, N.A., Soltis, P.S., Swarbreck, D., Thibaud-Nissen, F., Wang, S., Wegrzyn,
J.L., Zhang, G., Zhang, H., Lewin, H.A., Richards, S., 2022. Standards recommendations for
the Earth BioGenome Project. Proc Natl Acad Sci U S A 119.

Levänen, R., Kunnasranta, M., Pohjoismäki, J., 2018a. Mitochondrial DNA introgression at the
northern edge of the brown hare (Lepus europaeus) range. Ann Zool Fenn 55, 15-24.

Levänen, R., Pohjoismäki, J., Kunnasranta, M., 2019. Home ranges of semi-urban brown hares
(Lepus europaeus) and mountain hares (Lepus timidus) at northern latitudes. Ann Zool Fenn
56, 107–120.

Levänen, R., Thulin, C.G., Spong, G., Pohjoismäki, J.L.O., 2018b. Widespread introgression of
mountain hare genes into Fennoscandian brown hare populations. Plos One 13.

Li, D., Liu, C.M., Luo, R., Sadakane, K., Lam, T.W., 2015. MEGAHIT: an ultra-fast single-node solution
for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics
31, 1674-1676.

Li, H., 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34, 3094-
3100.

Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics 25, 1754-1760.

Lieberman-Aiden, E., van Berkum, N.L., Williams, L., Imakaev, M., Ragoczy, T., Telling, A., Amit, I.,
Lajoie, B.R., Sabo, P.J., Dorschner, M.O., Sandstrom, R., Bernstein, B., Bender, M.A.,
Groudine, M., Gnirke, A., Stamatoyannopoulos, J., Mirny, L.A., Lander, E.S., Dekker, J., 2009.
Comprehensive mapping of long-range interactions reveals folding principles of the human
genome. Science 326, 289-293.

Manni, M., Berkeley, M.R., Seppey, M., Zdobnov, E.M., 2021. BUSCO: Assessing Genomic Data
Quality and Beyond. Curr Protoc 1, e323.

Marques, J.P., Seixas, F.A., Farelo, L., Callahan, C.M., Good, J.M., Montgomery, W.I., Reid, N., Alves,
P.C., Boursot, P., Melo-Ferreira, J., 2020. An Annotated Draft Genome of the Mountain Hare
(Lepus timidus). Genome Biol Evol 12, 3656-3662.

Mazzoni, C.J., Ciofi, C., Waterhouse, R.M., 2023. Biodiversity: an atlas of European reference
genomes. Nature 619, 252.

Melo-Ferreira, J., Vilela, J., Fonseca, M.M., da Fonseca, R.R., Boursot, P., Alves, P.C., 2014. The
elusive nature of adaptive mitochondrial DNA evolution of an arctic lineage prone to
frequent introgression. Genome Biol Evol 6, 886-896.

Meng, G., Li, Y., Yang, C., Liu, S., 2019. MitoZ: a toolkit for animal mitochondrial genome assembly,
annotation and visualization. Nucleic Acids Res 47, e63.

Mikheenko, A., Prjibelski, A., Saveliev, V., Antipov, D., Gurevich, A., 2018. Versatile genome
assembly evaluation with QUAST-LG. Bioinformatics 34, i142-i150.

Minoudi, S., Papapetridis, I., Karaiskou, N., Chatzinikos, E., Triantaphyllidis, C., Abatzopoulos, T.J.,
Triantafyllidis, A., 2018. Genetic analyses of brown hare (Lepus europaeus) support limited
migration and translocation of Greek populations. Plos One 13, e0206327.

Ognev, S.I., 1940. 157. Lepus (Eulagus) europeaus Pall. (1778) заяц-русак, Mammals of the USSR
and adjacent countries: rodents (In Russian). Akademiya Nauk SSSR, Moscow, pp. 112-161.

Open2C, Abdennur, N., Fudenberg, G., Flyamer, I.M., Galitsyna, A.A., Goloborodko, A., Imakaev,
M., Venev, S.V., 2023. Pairtools: from sequencing data to chromosome contacts. bioRxiv.

Ou, S., Su, W., Liao, Y., Chougule, K., Agda, J.R.A., Hellinga, A.J., Lugo, C.S.B., Elliott, T.A., Ware, D.,
Peterson, T., Jiang, N., Hirsch, C.N., Hufford, M.B., 2019. Benchmarking transposable
element annotation methods for creation of a streamlined, comprehensive pipeline.
Genome Biol 20, 275.

Petrovan, S., 2013. Lepus europaeus (European hare), CABI Compendium.

Pohjoismäki, J.L.O., Michell, C., Levänen, R., Smith, S., 2021. Hybridization with mountain hares
increases the functional allelic repertoire in brown hares. Sci Rep 11, 15771.

Ranallo-Benavidez, T.R., Jaron, K.S., Schatz, M.C., 2020. GenomeScope 2.0 and Smudgeplot for
reference-free profiling of polyploid genomes. Nat Commun 11, 1432.

Reid, N., 2011. European hare (Lepus europaeus) invasion ecology: implication for the conservation of the endemic Irish hare (Lepus timidus hibernicus). Biol Invasions 13, 559-569.

Reid, N., Montgomery, W.I., 2007. Is naturalisation of the brown hare in Ireland a threat to the endemic Irish hare? Biol Environ 107b, 129-138.

Rikala, K., 1925. Ajojahdilla [flush hunting of hares]. Metsästys ja Kalastus 2, 44-48.

Salonius, A.H., 1924. Rusakkojäniksen leviämisalue Suomessa [Distribution of the brown hare in Finland]. In Finnish. Metsästys ja Kalastus, 165-169.

Schai-Braun, S.C., Schwienbacher, S., Smith, S., Hackländer, K., 2023. Coexistence of European hares and Alpine mountain hares in the Alps: what drives the occurrence and frequency of their hybrids? Journal of Zoology 320, 214-225.

Schai-Braun, S.C., Weber, D., Hacklander, K., 2013. Spring and autumn habitat preferences of active European hares (Lepus europaeus) in an agricultural area with low hare density. Eur J Wildlife Res 59, 387-397.

Seluanov, A., Vaidya, A., Gorbunova, V., 2010. Establishing primary adult fibroblast cultures from rodents. J Vis Exp.

Siivonen, L., 1972. Rusakko. Kustannusosakeyhtiö Otava, Keuruu, Finland.

Skov, L., Coll Macia, M., Lucotte, E.A., Cavassim, M.I.A., Castellano, D., Schierup, M.H., Munch, K., 2023. Extraordinary selection on the human X chromosome associated with archaic admixture. Cell Genom 3, 100274.

Smit, A.F.A., Hubley, R., Green, P., 2013-2015. RepeatMasker Open-4.0. http://www.repeatmasker.org.

Smith, R.K., Jennings, N.V., Harris, S., 2005. A quantitative analysis of the abundance and demography of European hares Lepus europaeus in relation to habitat type, intensity of agriculture and climate. Mammal Rev 35, 1-24.

Stott, P., 2003. Use of space by sympatric European hares (Lepus europaeus) and European rabbits (Oryctolagus cuniculus) in Australia. Mamm Biol 68, 317-327.

Suomalainen, E.W., 1922. Rusakkojäniksen leveneminen Suomessa [Distribution of the brown hare in Finland]. In Finnish. Metsästys ja Kalastus 6, 165-169.

Thulin, C.-G., 2003. The distribution of mountain hares (Lepus timidus, L. 1758) in Europe: A challenge from brown hares (L. europaeus, Pall. 1778)? Mammal Rev 33, 29-42.

Thulin, C.-G., Tegelström, H., 2002. Biased geographical distribution of mitochondrial DNA that passed the species barrier from mountain hares to brown hares (genus Lepus): an effect of genetic incompatibility and mating behaviour? J Zool 258, 299-306.

Thulin, C.G., Winiger, A., Tallian, A.G., Kindberg, J., 2021. Hunting harvest data in Sweden indicate precipitous decline in the native mountain hare subspecies Lepus timidus sylvaticus (heath hare). J Nat Conserv 64.

Uliano-Silva, M., Ferreira, J., Krasheninnikova, K., Darwin Tree of Life, C., Formenti, G., Abueg, L., Torrance, J., Myers, E.W., Durbin, R., Blaxter, M., McCarthy, S.A., 2023. MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity reads. BMC Bioinformatics 24, 288.

Vinogradov, A.E., 1998. Genome size and GC-percent in vertebrates as determined by flow cytometry: The triangular relationship. Cytometry 31, 100-109.

Wadén, D.J., 1910. Rusakkojäniksen (Lepus europaeus) nykyisestä leviämisestä Suomessa [The
current distribution of the brown hare in Finland]. In Finnish. Suomen metsästyslehti 8, 204-
207.
Wheeler, T.J., Eddy, S.R., 2013. nhmmer: DNA homology search with profile HMMs. Bioinformatics
29, 2487-2489.
Wolf, J.B., Lindell, J., Backstrom, N., 2010. Speciation genetics: current status and evolving
approaches. Philos Trans R Soc Lond B Biol Sci 365, 1717-1733.
Zhou, C., McCarthy, S.A., Durbin, R., 2023. YaHS: yet another Hi-C scaffolding tool. Bioinformatics
39.
774