1
2

# High quality genome assembly of the brown hare (*Lepus europaeus*) with chromosome-level scaffolding

Craig Michell[1,2], Joanna Collins[3], Pia K. Laine[4], Zsófia Fekete[1,5], Riikka Tapanainen[1], Jonathan M. D. Wood[3], Steffi Goffart[1] and Jaakko L. O. Pohjoismäki*[1]

[1]Department of Environmental and Biological Sciences, University of Eastern Finland – Joensuu, Finland
[2]Red Sea Research Center, Biological and Environmental Science and Engineering Division, King Abdullah University of Science and Technology,  – Thuwal, Kingdom of Saudi Arabia
[3]Tree of Life, Wellcome Sanger Institute, Hinxton, Cambridge, UK.
[4]Insitute of Biotechnology, University of Helsinki, Finland.
[5]Hungarian University of Agriculture and Life Sciences, Institute of Genetics and Biotechnology, Gödöllő, Hungary

*Corresponding author
Correspondence: jaakko.pohjoismaki@uef.fi

## Abstract

We present here a high-quality genome assembly of the brown hare (*Lepus europaeus* Pallas), based on a fibroblast cell line of a male specimen from Liperi, Eastern Finland. This brown hare genome represents the first Finnish contribution to the European Reference Genome Atlas pilot effort to generate reference genomes for European biodiversity.

The genome was assembled using 25X PacBio HiFi sequencing data and scaffolded utilizing a Hi-C chromosome structure capture approach. After manual curation, the assembled genome length was 2,930,972,003 bp with N50 scaffold of 125.8 Mb. 93.16% of the assembly could be assigned to 25 identified chromosomes (23 autosomes plus X and Y), matching the published karyotype. The chromosomes were numbered according to size. The genome has a high degree of completeness based on the BUSCO score (mammalia_odb10 database), Complete: 96.1% [Single copy: 93.1%, Duplicated: 3.0%], Fragmented 0.8%, and Missing 2.9%. The mitochondrial genome of the cell line was sequenced and assembled separately.

The brown hare genome is particularly interesting as this species readily hybridizes with the mountain hare (*Lepus timidus* L.) at the species contact zone in northern Eurasia, producing

fertile offspring and resulting in gene flow between the two species. In addition to providing a useful comparison for population studies, the genome can offer insight into the chromosomal evolution among Glires in general and Lagomorpha in particular. The chromosomal assembly of the genome also demonstrates that the cell line has not acquired karyotypic changes during culture.

**Keywords:** Lagomorpha, reference genome, chromosome structure, phylogeny, fibroblast cell line

## Introduction

The brown hare (*Lepus europaeus* Pallas), also known as the European hare, is a widespread species in the western parts of Eurasia (Bock, 2020). Besides its native range, the brown hare has been introduced to numerous regions, including the British Isles, the Falkland Islands, Canada, South America, Australia and New Zealand (Jaksic, 2023; Petrovan, 2013). In many places the brown hare is regarded as an invasive species and a threat to the local ecosystems (Stott, 2003) or native species (Reid, 2011).

The brown hare is a steppe-adapted species living in open grasslands and avoiding forested regions. Its colonization history in Europe is complex. The species has had glacial refugia during the Pleistocene in the Italian peninsula, the Balkans and Asia Minor (Fickel *et al*, 2008), with an interesting pre-glacial diversity hotspot in the archipelago of Greece (Minoudi *et al*, 2018). After the ice age, the species has been expanding its range both naturally as well as facilitated by human-caused changes in the landscape, especially through the expansion of agricultural lands and pastures. As mentioned above, the brown hare has been also frequently introduced by humans to new areas from antiquity to the present day (Petrovan, 2013), resulting in potential mixing of ancestral populations. This is true also for Finland, where the species has arrived naturally from the southeast through the Karelian isthmus in the late 19th century (Ognev, 1940; Siivonen, 1972) but also through local introductions to the southwestern parts of the country in the 1910s (Rikala, 1925).

Curiously, Linné was unaware of the existence of the brown hare, despite having described the mountain hare (*Lepus timidus* L.) and the South African Cape hare (*Lepus capensis* L.). In fact, the brown hare is not native to Linné's home country Sweden but was introduced there only a century after his death (Thulin, 2003). Curiously, the species author Peter Simon Pallas never published a formal description of the brown hare. Instead, the species was authority has been attributed to him because of inclusion of the name *Lepus europaeus* into his identification table of hare species of the world (Pallas, 1778: 30) (Figure 1A). Consequently, there is also no information of the type locality or any other information of the species in the original publication. The type locality of the brown hare has been retrospectively assigned – without too convincing arguments – to Burgundy (France) or Poland (Holden, 2005). Identifying Poland as the type locality was done by the Russian zoologist Sergey Ognev, and is based on Pallas mentioning in the 1778 text the hybrid forms of *Lepus variabilis* (synonym of *L. timidus*) and *L. europaeus* from Poland and Lithuania, as well as citing later works of Pallas for the species distribution (Ognev, 1940). In fact, Ognev (1940: 141) boldly states that "we consider southwestern Poland as a typical

terrain for the nominal subspecies; there are hares with the characteristics of the basic form,
expressed with full clarity" (translation from Russian).

While being widespread in the temperate regions of Western and Central Europe to the
Caspian steppes in historical times, the brown hare began a northward expansion in the 19$^{th}$
century. In the Fennoscandian region, the species reached the St Petersburg area by 1820s
(Ognev, 1940), arriving in Finland at the turn of the century (Siivonen, 1972; Thulin, 2003). By
1930, the species had become established throughout southern Finland, with the range
expansion stagnating southwest of the N64°-N62° -line until the 1990s. Quite likely benefitting
from the ongoing anthropogenic climate change, brown hares range expansion has intensified
during the last three decades, currently reaching the polar circle (Levanen *et al*, 2018a) (Figure
1B), as well as higher altitudes in the Alps (Schai-Braun *et al*, 2023). In Finland, the expansion
seem to be limited with 150 days of annual snow cover (Levanen *et al.*, 2018a). The range
expansion together with the past introductions have brought the brown hare increasingly in
contact with the mountain hare, especially at the northern edges of the species distribution,
constituting a threat especially to the more temperate climate adapted populations and
subspecies of the mountain hare (Reid, 2011; Reid & Montgomery, 2007; Thulin, 2003; Thulin *et*
*al*, 2021). Besides competing, brown hares also hybridize with mountain hares, producing fertile
offspring and resulting in gene flow between the species. This gene flow is biased towards the
brown hare (Ferreira *et al*, 2021; Levanen *et al.*, 2018a; Levanen *et al*, 2018b; Thulin & Tegelström,
2002), which obtains genetic variation from mountain hares, some of which might have adaptive
significance (Pohjoismaki *et al*, 2021). In contrast to the brown hare's expansion towards north,
many of the Central European brown hare populations are contracting in range and numbers,
likely driven by changes in agricultural practices and land use (Schai-Braun *et al*, 2013; Smith *et*
*al*, 2005).

Overall, the brown hare's features as an invasive species outside Europe and expansive
species in the north, its hybridization tendency and complex population ancestries make the
species highly interesting not only to study population genetics and adaptation mechanisms, but
also for the understanding the genomic makeup of species boundaries (Gaertner *et al*, 2022).
These studies would be greatly facilitated by the availability of high quality, chromosomally
scaffolded reference genomes (Blaxter *et al*, 2022).

The last decade has witnessed a boom in genome sequencing technologies, leading to
many sequencing initiatives, such as the Earth BioGenome Project (EBP), European Reference
Genome Atlas (ERGA) and the Darwin Tree of Life (DToL), aiming to produce high quality reference
genomes for a variety of organisms (Blaxter *et al.*, 2022). This has been enabled by the
technological development of high-throughput sequencing of long molecules. The read length
N50's > 19 kbp, allows a high contiguity compared to the genome assemblies released in the early
2010's that were based on short read sequencing technologies. Additionally, Hi-C sequencing, a
method measuring the contact frequency between all pairs of loci in the genome, has allowed
producing pretext maps of chromosomes (Ghurye *et al*, 2019), enabling chromosome level
assemblies with a precision that has been previously achieved only with model organisms
(Lawniczak *et al*, 2022).

In the presented work, we apply the PacBio HiFi long-range sequencing technology with
Hi-C sequencing to produce a high-quality reference genome of the brown hare (*Lepus*
*europaeus*). While there is no existing genome assembly for the species, a mountain hare

assembly (GCA_009760805) exists, representing the Irish subspecies *L. t. hibernicus* and is based on a female specimen, thus lacking the Y-chromosome. This assembly is also a so-called "pseudoreference", assembled to chromosomes using the rabbit (*Oryctolagus cuniculus* L.) reference genome (Marques *et al*, 2020). The original rabbit reference genome had been established using whole-genome shotgun sequencing of females representing the inbred Thorbecke New Zealand White line and was quite fragmented (Carneiro *et al*, 2014), updated only recently with long-read sequencing data (Bai *et al*, 2021). Notably, the diploid chromosome number in domestic rabbits is 2n = 44 (Korstanje *et al*, 1999), whereas it is 48 in brown hare and mountain hare (Gustavsson, 1972).

To obtain a reference genome conforming to current standards, we used a fibroblast cell line (LE1) of a male specimen of brown hare (*Lepus europaeus*) from Liperi, Eastern Finland (Gaertner *et al.*, 2022), as a source of high-molecular weight (HMW) DNA and fresh RNA. Compared to solid tissues, fibroblasts are optimal for Hi-C, as the method was originally developed for cells growing in a monolayer (Lieberman-Aiden *et al*, 2009). Now, over 240 years after the first mention of the name *Lepus europaeus* in the literature (Figure 1A), we have expanded the original seven-word description to encompass 2,930,972,003 genomic letters. These letters have been assembled in a highly complete (BUSCO mammalia_odb10 database, complete: 96.1%), and continuous (N50 scaffold: 125.8 Mb) manner with 93.16% of the sequence being placed in the identified 23 autosomes, and X and Y sex chromosomes. The presented genome assembly not only contains the identity of the brown hare as a species, but also provides a gateway to detailed knowledge of its biology and evolutionary history. Besides being the first Finnish contribution to the European Reference Genome Atlas (Mazzoni *et al*, 2023), the brown hare reference genome represents a continuing effort to map and understand our planets biodiversity.

# Methods

### Sampling and confirming of the species identity

A young male brown hare was hunted in October 2018 in Kuorinka, Liperi, Finland (62.6207 N, 29.4478 E, Figure 1B). In Finland, brown hares are highly dependent on similar man-made environments (Levanen *et al.*, 2018a; Levänen *et al*, 2019), and also the collection location is agricultural area with a mosaic of fields, shrubs and mixed forest with a strong brown hare population and only occasional stray mountain hares. Geographically, the sampled population is part of the same distribution continuum as with species proposed type locality, through Russian Karelia and the Baltic states to Poland. This is important, as is recommendable that the species reference genome would represent or be closely related to the type locality population (Lawniczak *et al.*, 2022).

The sampling had minimal impact on the local brown hare population and no impact on the habitat. As brown hares are legal game animals in Finland and the hunting followed the regional hunting seasons and legislation (Metsästyslaki [Hunting law] 1993/615/5§), the sampling adheres to the ARRIVE guidelines and no ethical assessment for the sampling was required. The sampling also did not involve International Trade in Endangered Species of Wild Fauna and Flora (CITES) or other export of specimens, as defined by the Convention on Biological Diversity (CBD).

The species identity was confirmed at site based on the morphological features
distinguishing the brown hare from the mountain hare, the only other hare species in northern
Europe. Further analysis of the coding part of the LE1 genome and mitochondrial DNA
haplotyping showed minimal ancestral admixture with mountain hares (Gaertner *et al.*, 2022). In
fact, unlike many other brown hares in the region, showing adaptive introgression of mountain
hare specific UCP1 alleles (Pohjoismaki *et al.*, 2021), the LE1 specimen is homozygous for the
common ancestral brown hare allele, UCP02.

**Generation and vouchering of the cell line**
The fibroblast cell line (LE1) was isolated from the specimen as described earlier (Gaertner
*et al.*, 2022) and is deposited as cryopreserved living cells under voucher number ZFMK-TIS-69747
into the biobank of the Stiftung Leibniz-Institut zur Analyse des Biodiversitätswandels, Zoological
Research Museum Alexander König (ZFMK), Bonn, Germany. Additional identifiers for the sample
are mLepEur2 and ERGA specimen ID ERGA_FI_3610_002 (COPO portal, https://copo-
project.org).

**HMW DNA extraction and PacBio HiFi sequencing**
A salting out-method was used to extract high molecular weight DNA from cells grown to
confluency on a 10 cm cell culture dish. Briefly, the cells were detached from the dish using
trypsin, followed by centrifugation and two washes with PBS. The salting out-method then
followed the 10X Genomics "Salting Out Method for DNA extraction from cells" protocol (10x-
Genomics, 2017). The high molecular weight DNA was quantified using a Qubit 3.0 (Thermo Fisher
Scientific™, Malaysia), followed by qualification with an 0.8 % Agarose gel. The DNA was sent for
the library preparation and sequencing at the DNA Sequencing and Genomics Laboratory,
Institute of Biotechnology, University of Helsinki on the PacBio Sequel II. In brief, the DNA was
quantified using Qubit and DNA fragment length was measured using Fragment Analyzer (HS large
fragment kit). DNA was then sheared with Megaruptor 3 (Diagenode) (45/200/30) to produce
DNA of length 24 kb. Buffer was then replaced with PacBio's Elution buffer using AMPure beads.
Repair and A-tailing, adapter ligation, nuclease treatment and cleanup with SMRTbell cleanup
beads was then performed with the SMRTbell prep kit 3.0. Larger fragments (greater than 10 000
bp) were then purified with the BluePippin (Sage Science) (0.75% DF Marker S1 high pass 6-10kb
v3) and DNA was further purified with PacBio AMPure beads and treated with DNA damage repair
mix (PacBio) (37oC for 60 mins) and then purified with AMPure clean-up beads and eluted to 11
ul PacBio's Elution buffer. Libraries were sequenced on the PacBio Sequel II at a concentration of
90 pM using PacBio's instructions (provided by the SMRTlink software). Sequence data from two
flow cells was produced. Genomescope2 was used to visualize the HiFi sequencing results
(Ranallo-Benavidez *et al*, 2020).

**Mitochondrial DNA (mtDNA) sequencing**
The mitochondrial genomes of our hare cell lines (Gaertner *et al.*, 2022) have been
sequenced for the purposes of another study using approximately 2 kb overlapping PCR-amplified
fragments of mtDNA. The primers used to amplify the mountain hare mtDNA were as follows:
Le93F: TTGTTTTGTAGCAAGTTTACACATGC
Le184R: GCTTAATACCTGCTCCTCTTGATCTA

Le1580F: TTAAACCCATAGTTGGCCTAAAAGC
Le1635R: TTGAGCTTTAACGCTTTCTTAATTGA
Le3045F: AGGCGTATTATTTATCCTAGCAACCT
Le3175R: CCTCATAAGAAATGGTCTGTGCGA
Le3921F: CCCCCTAATCTTTTCCATCATCCTAT
Le4482R: TCATCCTATATGGGCAATTGAGGAAT
Le4689F: AGGCTTTATTCCAAAGTGAATTATTATTCA
Le5417R: AGGCTCCAAATAAAAGGTAGAGAGTT
Le6696F: ATACCGTCTCATCAATAGGCTCCTTC
Le6756R: ATAAAGATTATTACTATTACAGCGGTTAGA
Le8603F: AGCCTATATCTACATGATAATACTTAATGA
Le8698R: CGGATAAGGCCCCGGTAAGTGG
Le10552F: TTGAAGCAACACTAATCCCTACACTA
Le10613R: TCGTTCTGTTTGATTACCTCATCGT
Le11301F: ACCATTAACCTTCTAGGAGAGCTTCT
Le11807R: AGGATAATGATTGAGACGGCTATTGA
Le12407F: GTCTAATCCTAGCTGCTACAGGTAAG
Le12791R: GAGCATAAAAAGAGTATAGCTTTGAA
Le14204F: ATTGTTAACCACTCTCTAATCGACCT
Le14514R: CCAATGTTTCAGGTTTCTAGGTAAGT
Lt16056F: TGGGGTATGCTTGGACTCAAC
Le16119R: TCGTCTACAATAAGTGCACCGG
232
In total, 12 separate reactions were prepared to cover the mitochondria genome:
1.    Lt16056F + Le184R: 1871 bp
2.    Le93F + Le1635R: 1543 bp
3.    Le1580F + Le3175R: 1596 bp
4.    Le3045F + Le4482R: 1438 bp
5.    Le3921F + Le5417R: 1497 bp
6.    Le4689F + Le6756R: 2068 bp
7.    Le6696F + Le8698R: 2003 bp
8.    Le8603F + Le10613R: 2011 bp
9.    Le10552F + Le11807R: 1256 bp
10.    Le11301F + Le12791R: 1491 bp
11.    Le12407F + Le14514R: 2108 bp
12.    Le14204F + Le16119R: 1916 bp
(Expected fragment size based on the published *Lepus europaeus* mtDNA sequence from Sweden
[NC_004028.1]).
248
The fragments were amplified from total DNA preparations using a PCR program with a 1 min
94 °C denaturing step, followed by 35 cycles of 94 °C for 15 s, 56 °C for 15 s and 72 °C for 2 min
and a final 3 min elongation step at 72 °C. The obtained products were gel purified using the
GeneJET gel extraction kit (Thermo Fisher Scientific™) and sent for sequencing using Illumina
MiniSeq at the Genome Center of Eastern Finland.

The sequence of the non-coding region-containing PCR fragment (Lt16056F + Lt184R) was further

validated by Sanger sequencing, applying also the following additional sequencing primer:

Le101F: TATAAATTCCTGCCAAACCCCAAAAA

**Figure 1**. The brown hare. (A) An excerpt from the identification table by Pallas (1778: 30), which

has been recognized as the species description for *Lepus europaeus*. "Tips of the ears longer than

the head, tail black above." (translation from Latin). (B) The geographic location of the LE1 sample

used in this study. The grey-shaded area shows the approximate current distribution of the

brown hare in Finland, based on the data in the Finnish Biodiversity Information Facility (FinBIF,

https://laji.fi/en). Individual records from the north are from settlemens or towns. The arctic

circle (black line) runs along the 66.6 ° parallel. Background photo shows a typical Finnish male

brown hare. Photo taken in Joensuu, 20 km East from the LE1 sampling site by Dr. Mervi

Kunnasranta.

**Hi-C library preparation**

Hi-C sequencing libraries were prepared following the protocol of (Belaghzal *et al*, 2017)

with the following changes: 1.) To prepare a diverse sequencing library, we performed the Hi-C

protocol in triplicates, 2.) Size fractionation was performed using the NEBNext Ultra II FS DNA

module (New England Biolabs), 3.) Library preparation was performed using the NEBNext Ultra II

library preparation kit for Illumina (New England Biolabs) and 4.) Library enrichment was

performed using triplicate PCR reactions with six cycles of PCR. The PCR reactions were then
purified using Ampure XP beads (Beckman Coulter Life Sciences) at a ratio of 0.9X. The final clean
libraries were quantified using Qbit, followed by agarose gel electrophoresis to confirm the
fragment size. The sequencing was performed on a single lane of the Illumina NovaSeq 6000 using
the SP flowcell with paired-end chemistry 2 x 150bp.
281
**Genome assembly**
HiFiasm version 0.18.7 (Cheng *et al*, 2021) was used to assemble the 25X (rq >0.99) PacBio
HiFi reads using the recommended arguments *-l3 –h1 and –h2 –primary* to integrate the Hi-C read
data and produce a primary assembly. We then continued with the scaffolding of the primary
assembly and annotation. To process the HiC data, we first mapped the Hi-C data to the primary
genome assembly using BWA-mem version 0.7.17 (Li & Durbin, 2009) with the arguments *-SP*.
The mapped reads were then parsed and filtered using pairtools version 1.0.2 (Open2C *et al*,
2023). To parse the aligned Hi-C reads, we used the options *–no-flip –dd-columns mapq –walks-*
*policy mask*. The parsed pair data was then sorted and deduplicated using default arguments with
pairtools. Finally, we selected unique-unique and unique-rescued pairs and split these into the
pairs file and bam file for input in YaHS version 1.1 (Zhou *et al*, 2023). YAHS was run using the
default parameters with the primary contig assembly and the filtered Hi-C bam file. HiFiasm
operates as a haplotype-aware assembler, counting and assembling unique k-mers. Utilizing k-
mer coverage information, the assembler identifies regions in the genome with expected
heterozygous coverage and subsequently amalgamates them into a unified sequence. To refine
the assembly further, Hi-C data is incorporated to elucidate the phasing of variants and ascertain
connected genomic regions. Consequently, the assembler's output provides a primary assembly
along with an alternative assembly, effectively separating the haplotypes by their distinct alleles
within the genomic content (please see https://lh3.github.io/2021/04/17/concepts-in-phased-
assemblies). Contiguity and general genome statistics were calculated using QUAST version 5.2.0
(Mikheenko *et al*, 2018). We assessed the completeness of the genome by calculating the number
of complete single copy orthologs with BUSCO version 5.1.2 (Manni *et al*, 2021), using the
mammalia_odb10 database as well as the more lineage specific glires_odb10 database.
305
**Genome annotation**
Repeat annotation of the genome was performed with EDTA version 2.1.0 (Ou *et al*, 2019),
a *de novo* repeat identification pipeline. Using the repeat library produced by EDTA, we masked
the scaffolded genome using RepeatMasker version 4.1.1 (Smit *et al*, 2013-2015). RNA-seq data
from the same cell line (SRA accession number: SRR18740842) as well as RNA-seq data from other
*L. timidus* libraries (SRA accession number: SRR10020054, SRR10020055, SRR10020060,
SRR10491719, SRR18740839, SRR18740840, and SRR18740841) was collected from the sequence
read archive (SRA). The previously produced RNA-seq data (Gaertner *et al.*, 2022) was trimmed
using fastp version 0.23.2 (Chen *et al*, 2018) and mapped against the masked genome using
HISAT2 version 2.1.0 (Kim *et al*, 2019) with the default parameters. Furthermore, we included the
protein sequences of the assembled transcripts from this cell line as further evidence for the
genome annotation. These lines of gene evidence were included in the annotation using BRAKER3
version 3.02 (Bruna *et al*, 2021). Telomeric sequences [AACCCT]n were identified using a Telomere

Identification toolkit (tidk) version 0.2.31. The telomeric sequence copy number was then calculated in windows of 200kb for visualization using Circos (Krzywinski *et al*, 2009).

**Mitochondrial DNA assembly and annotation**

Mitochondrial DNA was assembled from the PCR-amplified and Illumina sequenced mtDNA using the MitoZ pipeline (Meng *et al*, 2019). After comparing the results of the pipeline's outputs with different kmer options, we selected the best assembly as final. Run options used in the final assembly were --clade Chordata –fastq_read_length 150, --requiring_taxa Chordata --genetic_code 2 --kmers_megahit 21 29 39 59 79 99 119 141. The tools invoked by the pipeline included fastp (Chen *et al.*, 2018) for cleaning the raw data, MEGAHIT (Li *et al*, 2015) for assembly, after which sequences were filtered to ensure the correct taxa by HMMER (Wheeler & Eddy, 2013) and further filtered for completeness of protein-coding genes. Annotation steps were done using TBLASTN (Gertz *et al*, 2006), GeneWise (Birney *et al*, 2004) and MiTFi (Juhling *et al*, 2012). Final manual curation, the annotation of the non-coding region (NCR) as well as the illustration of the mitochondrial genome was done using Geneious® 10.2.6 (Biomatters. Available from https://www.geneious.com). The functional loci on the NCR were identified based on the similarity with the human (NC_012920) and mouse (FJ374652) NCR sequences.

**Manual curation**

The assembled and annotated genome was manually curated to further improve its quality based on Hi-C contact maps as described in (Howe *et al*, 2021), allowing to break or join erroneus scaffold assemblies and removing duplicated contigs.

**Comparison with previous assembly**

We performed a comparison of our scaffolded assembly to the current *L. timidus* genome assembly (NCBI Accession number: GCA_009760805). Mapping to the genome was performed using minimap2 version 2.21 (Li, 2018) with the arguments *-asm5*. A dot plot of the alignment was created using the R script pafCoordsDotPlotly.R (https://github.com/tpoorten/dotPlotly).

# Results

The genome assemblies can be accessed via BioSample accession SAMN33984520 as well as BioProject accession PRJNA1009711 for the primary assembly and PRJNA1009710 for the alternative assembly.

**Genome assembly**

The expected haploid genome size of the *L. europaeus* is similar to *L. timidus*, which has been estimated to be 3.25 pg (3,178,500,000 bp) (Vinogradov, 1998), containing 23 autosomal chromosomes and two sex chromosomes (Gustavsson, 1972). PacBio HiFi sequencing with two flow cells resulted in 25 X coverage of the expected genome size. The sequence N50 of the HiFi data was 19.97 kb. When k-mer size of 21 is used, the resulting genome size from the PacBio HiFi data is 2.4 Gbp (Figure 2A). The Illumina sequencing of the Hi-C data produced 494,285,585 paired reads, representing about 51 X coverage of the genome, with a duplication rate of 17 %. Assembly

with HiFiasm yielded a contig assembly of 2.96 Gbp made up of 671 contigs with a contig N50 of
58 Mbp. The longest contig was 149 Mbp. Using the uniquely mapped Hi-C data, we were able to
scaffold the contigs and fix misassembled contigs (Figure 2B). Although the genome size (2.96
Gbp) and the largest scaffold (149 Mbp) were similar to the contig assembly, the scaffold N50 was
increased to 124 Mbp (N=10) after the Hi-C scaffolding. The obtained scaffold N90 was 21 Mbp
(N=25). The BUSCO scores of the *L. europaeus* assembly suggest it to be near-complete, with the
following results (%- of the total genome): Complete: 96.1% [Single copy: 93.1 %, Duplicated: 3.0
%], Fragmented 0.8 %, and Missing 2.9 % based on the mammalia_odb10 database.
To further improve the assembly, manual curation was performed, resulting in 76 scaffold
breaks, 81 joins and removal of 62 erroneously duplicated contigs (Table 1).  These interventions
led to an increase in scaffold N50 of 1.69 % from 123.7 Mb to 125.8 Mb, and a reduction in scaffold
count of 9.21 % from 716 to 650. Of the finalized assembly, 93.16 % could be assigned to 25
identified chromosomes (23 autosomes plus X and Y), matching the known karyotype.
Chromosomes were numbered according to size (Table 2). The curated genome length was
2,930,972,003 bp.

**Figure 2.** A) Genomescope2 profile of the PacBio HiFi data. B) YaHS pretext map of Hi-C scaffolding after manual curation. C) Dot plot comparison with the mountain hare pseudoreference (GCA_009760805) assembly with rabbit chromosomal assignments. Each grey horizontal line represents a contig end. Names of the non-chromosomal contigs have been omitted for clarity. Rabbit chromosomes corresponding two hare chromosomes as well as the X chromosome with an inversion are are highlighted.

386 **Table 1.** The brown hare reference genome chromosome assignment and assembly statistics

| Chromosome assignment | |
|---|---|
| **Chr length** | 2,730,543,680 bp |
| **Autosomes** | 23 |
| X | 1 |
| Y | 1 |
| Unlocalised | 19 |

| Statistics for primary assembly after manual curation | |
|---|---|
| **# scaffolds** | 650 |
| Total scaffold length | 2,930,972,003 |
| **Average scaffold length** | 4,509,187.70 |
| **Scaffold N50** | 125,776,599 |
| **Scaffold auN** | 117,151,857.99 |
| **Scaffold L50** | 10 |
| Largest scaffold | 184,151,585 |
| **Smallest scaffold** | 1000 |
| # contigs | 1068 |
| Total contig length | 2,930,889,045 |
| **Average contig length** | 2,744,278.13 |
| **Contig N50** | 43,409,466 |
| **Contig auN** | 47,572,960.95 |
| **Contig L50** | 21 |
| Largest contig | 141,329,000 |
| **Smallest contig** | 1000 |
| # gaps in scaffolds | 418 |
| Total gap length in scaffolds | 82,958 |
| Average gap length in scaffolds | 198.46 |
| **Gap N50 in scaffolds** | 200 |
| Gap auN in scaffolds | 199.70 |
| **Gap L50 in scaffolds** | 208 |
| Largest gap in scaffolds | 200 |
| Smallest gap in scaffolds | 31 |
| Base composition | |
| **A** | 821,587,687 |
| **C** | 645,897,412 |
| **G** | 644,604,632 |
| **T** | 818,799,314 |

| | |
|---|---|
| **GC content** | 44.03 % |
| **# soft-masked bases** | 0 |
| # segments | 1,068 |
| Total segment length | 2,930,889,045 |
| **Average segment length** | 2,744,278.13 |
| **# gaps** | 418 |
| # paths | 650 |
| | |

388 **Table 2.** Chromosome assemblies of the brown hare reference genome with their corresponding
389 GenBank access numbers.

| Chromosome | GenBank | Size (bp) | GC content (%) |
|---|---|---|---|
| 1 | CM065551.1 | 184,151,585 | 42.0 |
| 2 | CM065552.1 | 176,467,698 | 42.0 |
| 3 | CM065553.1 | 169,342,607 | 42.0 |
| 4 | CM065554.1 | 166,606,208 | 43.0 |
| 5 | CM065555.1 | 159,335,674 | 46.0 |
| 6 | CM065556.1 | 132,443,831 | 42.0 |
| 7 | CM065557.1 | 132,293,278 | 44.0 |
| 8 | CM065558.1 | 130,587,543 | 41.0 |
| 9 | CM065559.1 | 125,776,599 | 44.5 |
| 10 | CM065560.1 | 125,704,418 | 46.0 |
| 11 | CM065561.1 | 117,559,561 | 44.0 |
| 12 | CM065562.1 | 105,667,338 | 44.0 |
| 13 | CM065563.1 | 100,925,866 | 44.0 |
| 14 | CM065564.1 | 99,301,477 | 44.0 |
| 15 | CM065565.1 | 95,895,056 | 42.0 |
| 16 | CM065566.1 | 91,637,918 | 42.5 |
| 17 | CM065567.1 | 84,898,765 | 45.5 |
| 18 | CM065568.1 | 77,563,730 | 48.5 |
| 19 | CM065569.1 | 72,700,603 | 47.5 |
| 20 | CM065570.1 | 69,331,453 | 45.5 |
| 21 | CM065571.1 | 60,609,710 | 47.5 |
| 22 | CM065572.1 | 44,025,767 | 47.0 |
| 23 | CM065573.1 | 30,716,498 | 53.0 |
| X | CM065574.1 | 135,183,192 | 40.5 |
| Y | CM065575.1 | 33,560,193 | 41.0 |
| MT | OR915849.1 | 17,540 | 38.6 |

**Genome annotation**

EDTA produced a curated custom repeat library of 7,045 repetitive elements. Interestingly, the proportion of the genome masked as repetitive elements was higher than expected at 46.8 % of the genome. In the mountain hare pseudoreference genome assembly (GCA_009760805), only 28 % of the genome was masked, and similarly the *k*-mer based estimate of the repetitive elements was 26.7 %. To annotate the genome, we mapped ~1.6 billion RNA-seq reads from previously published RNA-seq libraries, including this cell line. The average mapping rate of all libraries was 89.95 %, which we consider a good indicator of the quality of this genome. Using BRAKER3, we were able to annotate 19,906 gene models in the genome. We then ran BUSCO on the predicted genes and the results show a good level of completeness: Complete: 81.1 % [Single copy: 63.3 %, Duplicate: 17.8 %], Fragmented: 1.1 %, Missing: 17.8 %, Total: 9226 when compared to the mammalia_odb10 database. Telomeric sequences on both ends of the chromosome were detected on 13 chromosomes (Figure 3A). Furthermore, telomeric sequences were found in high copy number throughout the chromosome length, a feature previously noted using FISH in mountain hares (Forsyth *et al*, 2005).

**Comparison to the mountain hare assembly**

Minimap2 was able to align 97.3 % of the contig sequences from the mountain hare pseudoreference genome assembly (GCA_009760805) to this genome assembly (Figure 2C). As pointed out earlier, the mountain hare genome assembly has been scaffolded against the rabbit reference genome (Giska *et al*, 2022), which has a different chromosome count (Beklemisheva *et al*, 2011) and explains the few notable inversions and structural differences between the two genomes (Figure 2C). For example, the chr7 and chr 12 assemblies of the brown hare have a good degree of synteny with the mountain hare pseudoreference (= rabbit) chr1, with the chr12 being in inverted orientation. Similarly, the brown hare chr13 and chr16 assemblies have a good synteny and orientation with the rabbit chr2.

**Mitochondrial genome**

While it is possible to recover the entire mitochondrial genome from the HiFi sequencing data (Uliano-Silva *et al*, 2023), we had already previously sequenced the mitochondrial genomes of our hare cell lines for the purposes of another study. The assembly of the Illumina data provided a 16,836 bp mtDNA sequence, which was complete and circular, with a 142 bp non-repetitive overlap between the 3´ and 5´ ends of the original assembly removed from the final version. All the expected vertebrate mitochondrial genes, 37 in total, were found (Figure 3B).

Interestingly, the primary assembly of the Illumina data was notably smaller than expected based on the original PCR amplicons, with a notable difference between the expected and obtained non-coding region (NCR) length. Sanger sequencing of the PCR-fragment containing the NCR (Lt16056F + Le184R) revealed that it contained a sequence run consisting of six head-to-tail repeats of a 140 bp element between OriH and tRNA-Phe, explaining the incorrect assembly of the NCR sequence from the short-read Illumina data. With the addition of the repeat elements, the final mitochondrial genome size of our specimen is 17,540 bp.

Based on the sequence similarity with human and mouse mitochondrial genomes, we were able to tentatively identify the termination associated sequence (TAS, 2 nt difference to human), required for the generation of the 7S DNA and thus the displacement loop (D-loop)

structure on the NCR (Crews *et al*, 1979). Interestingly, assuming that the 7S DNA is roughly the same size as in other mammals (600-650 nt), its 5'-end would map to the beginning of another highly repetitive 240 bp region, likely to have structural significance. This is a notably longer sequence run than the tRNA-like sequence reported at the replication origin of *L. timidus* mtDNA (Melo-Ferreira *et al*, 2014). We have preliminary assigned this structure to correspond to the heavy-strand origin of replication (OriH), supported by the fact that a similar, tRNA sequence-derived hairpin structure is required for the priming of the origin of the light-strand replication (OriL) (Fuste *et al*, 2010). Experimental validation of the proposed OriH is required.

**Figure 3.** A reference genome for the brown hare. A) A size map of 23 autosomes as well as X and Y sex chromosomes of *Lepus europaeus*. Telomeric sequences and their relative lengths are indicated on the inner circle. B) A schematic illustration of the mountain hare mitochondrial genome. Genes encoded by the H-strand are placed inside and the L-strand genes outside of the circle. NCR = non-coding region.

## Discussion

Given the challenges of assembling a large mammalian genome, we have produced a high-quality genome assembly for the brown hare. The genome assembly has a high degree of contiguity (Contig N50 43 Mbp) and completeness (BUSCO complete 96.1 %). Interestingly, the genome contains a high number of repetitive elements, a fact that warrants further investigation to elucidate their identity. While vast majority of the sequences seem to come from retroelements or endoretroviruses, we cannot fully exclude that some multicopy host genes could be masked. A detailed analysis of the repeat element libraries from this and another vertebrate genome assembly releases might be warranted.

The genome assembly is in chromosome scale, and all 23 autosomes as well as X and Y chromosomes could be scaffolded. Compared with the closely related mountain hare

pseudoreference assembly, the brown hare genome assembly is much more contiguous (Table 1)
and inclusion of telomeric repeats on both ends of several chromosomes (Figure 3A) further
demonstrates the telomere-to-telomere continuity of these assemblies. The presented assembly
also resolves the expected chromosome structure for the species (Figure 2B, C), which is
informative regarding the karyotype difference between hares (n = 24) and rabbits (n = 22).
Comparison of the brown hare assembly with the mountain hare pseudoreference chromosomes
shows that the rabbit chromosome 1 corresponds to the brown hare chromosomes 12 and 17,
with the chromosomal arm corresponding to the former is in an inversed orientation. Similarly,
the rabbit chromosome 2 corresponds to the hare chromosomes 13 and 16, both in the same
orientation. It should also be noted that as the hare chromosomes were numbered according to
their size, the numbering in general does not match those of the rabbit chromosomes (Figure 2C)
The obtained 17,540 bp long mitochondrial genome assembly also provides some
interesting aspects of mtDNA sequence variation within mammalian species. The size difference
to human (16,569 bp) and mouse (16,298 bp) mitochondrial genomes is caused by a rather long
non-coding region (2,102 bp vs. 1,123 bp and 879 bp in humans and mice, respectively), longer
rRNA and tRNA genes, as well as additional non-coding nucleotides between genes. The
mitochondrial genome of our specimen is slightly smaller than the 17,734 bp previous NCBI
Reference Sequence for brown hare (NC_004028.1) from Sweden. The size difference is mostly
due to a copy number difference in tandem repeat elements within the non-coding region, with
seven in the NC_004028.1 and six in our specimen. Indels elsewhere in the NCR explain the rest
of the difference.
The main technical challenge for highly continuous, good quality reference genome
assemblies is the availability of relatively large quantities of intact HMW DNA for HiFi sequencing.
This typically requires snap freezing of tissue samples and assuring an intact cold chain of below
-80 °C for their sending, handling, and long-term storage (Januszczak & Holroyd, 2023). While
development in sequencing technologies can reduce the required DNA amounts, obtaining high
sequencing coverage over a large vertebrate genome is easier when material is plentiful. Similarly,
RNA is very sensitive to degradation in post-mortem tissue samples, also requiring immediate
preparation or snap-freezing to be suitable for RNA-seq. Although transcript data is not obligatory,
it is highly useful for the purpose of genome annotation. The requirement of large samples of
fresh tissue for -omics purposes complicates the sampling of most vertebrates species. For
example, in Finland all land vertebrates are protected by law (animal protection law 247/1996,
hunting law 615/1993), with very few exceptions, such as game animals and pest species.
Capturing and lethal sampling is especially problematic for endangered species. Not only will the
sampling require extensive permits, but it is also difficult to justify and conduct ethically. Here,
the generation of primary fibroblast cell lines can offer a solution. Although the source specimen
for the cell line utilized in this study was hunted, fibroblasts can be isolated from relatively small
skin biopsies, such as ear clippings (Seluanov *et al*, 2010), with minimally invasive sampling and
harm to the individual. These cells can be expanded in tissue culture and, when immortalized,
provide a highly scalable source of fresh DNA and RNA. Furthermore, living cells can be stored as
frozen stocks in biobanks for decades (Fazekas *et al*, 2017). Tissue sampling can be done directly
into the growth medium in ambient temperature, allowing several days for the transport to the
laboratory without a need for dry ice or $N_2$-cryocontainers, which greatly simplifies collection and
logistics. The same applies to the isolated cells. As an example, we sent living cell suspensions

from Joensuu, Finland to the ZFMK biobank in Bonn, Germany, over regular airmail in ambient
temperature. The receiving laboratory can then amplify the cells by culturing and cryopreserve
them in sufficient replicate stocks. It is worth to point out that the obtained genome assembly
scaffolded to 23 autosomes, X and Y chromosomes, as expected for the species, shows no
evidence of karyotypic drift caused by cell culture, dispelling possible concerns for the use of cell
lines for such work (Gaertner *et al.*, 2022).

The brown hare is an iconic European mammal, locally familiar to a wider public of nature-
goers, farmers and hunters, while being at the same time an invasive species on many other
continents. It has populations with complex ancestral makeup due to isolated Pleistocene refugia,
frequent introductions and cross-species hybridization, making it an exceptionally interesting
mammal for genetic studies. Furthermore, depending on the local environmental context, brown
hares can be both an endangered as well as a rapidly expanding species, prompting opportunities
for both conservation as well as invasion genetics studies. A high-quality reference genome
assembly will allow us also to peer deeper into genome structure and chromosome evolution,
which will be particularly interesting from the viewpoint of genetic introgression (Juric *et al*, 2016)
and maintenance of species boundaries. Any issues of genetic compatibility between species
should leave their fingerprint in the genomes (Skov *et al*, 2023), providing interesting insight into
the speciation mechanisms (Wolf *et al*, 2010). Having a high-quality genome for both species has
important implications for the fields of evolutionary and conservation biology. Due to the high
contiguity of the genome assembly, it will be possible to perform accurate analyses of linkage
disequilibrium as well as the identification of runs of homozygosity (ROH). Accurate ROH analysis
is useful for identifying and understanding the impact of inbreeding burden on the genome of
increasingly threatened and isolated species, as quantitative traits can be associated with ROH
burden (Ceballos *et al*, 2018). Furthermore, this information might provide important insights
into the hybridization of the brown hare with the mountain hare, as well as help to pinpoint
genomic regions that might be helping the species to adapt to local environmental conditions
during the ongoing range expansion (Pohjoismaki *et al.*, 2021). Other *Lepus* reference genome
assemblies available soon will help to complete this picture. Together, this genome assembly and
the mountain hare genome assembly will provide a solid base for future studies.

## Acknowledgements
We acknowledge the DNA Sequencing and Genomics Laboratory, Institute of Biotechnology,
University of Helsinki for the PacBio and Hi-C sequencing, with special thanks to Dr Petri Auvinen
and Dr Martyn James for helpful discussions and personal interest towards the project. Dr Seppo
Helisalmi and Dr Joose Raivo from the Gene diagnostics lab at Genome Center of Eastern Finland
are thanked for providing the mtDNA sequencing service. We are grateful to the European
Reference Genome Atlas (ERGA) for the opportunity to include our mountain hare genome into
their pilot project. Dr Ann McCartney is thanked for her kind help throughout the process and Mr
Jukka Pusa for his contribution in obtaining the brown hare specimen. We thank Ms Anita
Kervinen for her valuable laboratory assistance and Dr. Mervi Kunnasranta for kindly allowing to
use her hare photo in the publication.

## Data, scripts, code, and supplementary information availability

The genome assembly is available from the NCBI database under BioProject ID PRJNA950335. The cell line is available through the corresponding author as well as from the ZFMK biobank.

## Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

## Funding

This study belongs to the xHARES consortium funded by the R'Life initiative of the Academy of Finland, grant no. 329264.

## References

10x-Genomics (2017) Salting out method for DNA extraction from cells. *Sample Preparation Demostrated Protocol* CG00016. Rev A.: 1-5

Bai Y, Lin W, Xu J, Song J, Yang D, Chen YE, Li L, Li Y, Wang Z, Zhang J (2021) Improving the genome assembly of rabbits with long-read sequencing. *Genomics* 113: 3216-3223

Beklemisheva VR, Romanenko SA, Biltueva LS, Trifonov VA, Vorobieva NV, Serdukova NA, Rubtsova NV, Brandler OV, O'Brien PC, Yang F *et al* (2011) Reconstruction of karyotype evolution in core Glires. I. The genome homology revealed by comparative chromosome painting. *Chromosome Res* 19: 549-565

Belaghzal H, Dekker J, Gibcus JH (2017) Hi-C 2.0: An optimized Hi-C procedure for high-resolution genome-wide mapping of chromosome conformation. *Methods* 123: 56-65

Birney E, Clamp M, Durbin R (2004) GeneWise and Genomewise. *Genome Res* 14: 988-995

Blaxter M, Archibald JM, Childers AK, Coddington JA, Crandall KA, Di Palma F, Durbin R, Edwards SV, Graves JAM, Hackett KJ *et al* (2022) Why sequence all eukaryotes? *P Natl Acad Sci USA* 119

Bock A (2020) Lepus europaeus (Lagomorpha: Leporidae). *Mammalian Species* 52: 125-142

Bruna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M (2021) BRAKER2: automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database. *NAR Genom Bioinform* 3: lqaa108

Carneiro M, Rubin CJ, Di Palma F, Albert FW, Alfoldi J, Martinez Barrio A, Pielberg G, Rafati N, Sayyab S, Turner-Maier J *et al* (2014) Rabbit genome analysis reveals a polygenic basis for phenotypic change during domestication. *Science* 345: 1074-1079

Ceballos FC, Joshi PK, Clark DW, Ramsay M, Wilson JF (2018) Runs of homozygosity: windows into population history and trait architecture. *Nat Rev Genet* 19: 220-234

Chen SF, Zhou YQ, Chen YR, Gu J (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* 34: 884-890

Cheng H, Concepcion GT, Feng X, Zhang H, Li H (2021) Haplotype-resolved de novo assembly using
phased assembly graphs with hifiasm. *Nat Methods* 18: 170-175
Crews S, Ojala D, Posakony J, Nishiguchi J, Attardi G (1979) Nucleotide sequence of a region of
human mitochondrial DNA containing the precisely identified origin of replication. *Nature*
277: 192-198
Fazekas J, Grunt TW, Jensen-Jarolim E, Singer J (2017) Long term storage in liquid nitrogen leads
to only minor phenotypic and gene expression changes in the mammary carcinoma model
cell line BT474. *Oncotarget* 8: 35076-35087
Ferreira MS, Jones MR, Callahan CM, Farelo L, Tolesa Z, Suchentrunk F, Boursot P, Mills LS, Alves
PC, Good JM *et al* (2021) The Legacy of Recurrent Introgression during the Radiation of
Hares. *Syst Biol* 70: 593-607
Fickel J, Hauffe HC, Pecchioli E, Soriguer R, Vapa L, Pitra C (2008) Cladogenesis of the European
brown hare (Lepus europaeus Pallas, 1778). *Eur J Wildlife Res* 54: 495-510
Forsyth NR, Elder FF, Shay JW, Wright WE (2005) Lagomorphs (rabbits, pikas and hares) do not
use telomere-directed replicative aging in vitro. *Mech Ageing Dev* 126: 685-691
Fuste JM, Wanrooij S, Jemt E, Granycome CE, Cluett TJ, Shi Y, Atanassova N, Holt IJ, Gustafsson
CM, Falkenberg M (2010) Mitochondrial RNA polymerase is needed for activation of the
origin of light-strand DNA replication. *Mol Cell* 37: 67-78
Gaertner K, Michell C, Tapanainen R, Goffart S, Saari S, Soininmaki M, Dufour E, Pohjoismaki JLO
(2022) Molecular phenotyping uncovers differences in basic housekeeping functions among
closely related species of hares (Lepus spp., Lagomorpha: Leporidae). *Mol Ecol*
Gertz EM, Yu YK, Agarwala R, Schaffer AA, Altschul SF (2006) Composition-based statistics and
translated nucleotide searches: improving the TBLASTN module of BLAST. *BMC Biol* 4: 41
Ghurye J, Rhie A, Walenz BP, Schmitt A, Selvaraj S, Pop M, Phillippy AM, Koren S (2019) Integrating
Hi-C links with assembly graphs for chromosome-scale assembly. *PLoS Comput Biol* 15:
e1007273
Giska I, Pimenta J, Farelo L, Boursot P, Hacklander K, Jenny H, Reid N, Montgomery WI, Prodohl
PA, Alves PC *et al* (2022) The evolutionary pathways for local adaptation in mountain hares.
*Mol Ecol* 31: 1487-1503
Gustavsson I (1972) Mitotic and meiotic chromosomes of the variable hare (Lepus timidus L.), the
common hare (Lepus europaeus Pall.) and their hybrids. *Hereditas* 67: 27-34
Holden ME (2005) Family Gliridae. In: *Mammal Species of the World: A Taxonomic and Geographic*
*Reference*, Wilson D.E., Reeder D.M. (eds.) pp. 819-841. Johns Hopkins University Press:
Baltimore, USA
Howe K, Chow W, Collins J, Pelan S, Pointon D-L, Sims Y, Torrance J, Tracey A, Wood J (2021)
Significantly improving the quality of genome assemblies through curation. *GigaScience* 10
Jaksic FM (2023) Historical ecology and current status of the European hare
in South America: a new species in new countries. *Stud Neotrop Fauna E*
Januszczak I, Holroyd N (2023) DToL Tissue and Blood Sampling Standard Operating Procedure:
Chordata:          Vertebrata          (general).          *protocolsio*
dx.doi.org/10.17504/protocols.io.n92ldmjool5b/v1
Juhling F, Putz J, Bernt M, Donath A, Middendorf M, Florentz C, Stadler PF (2012) Improved
systematic tRNA gene annotation allows new insights into the evolution of mitochondrial

tRNA structures and into the mechanisms of mitochondrial genome rearrangements.
*Nucleic Acids Res* 40: 2833-2845

Juric I, Aeschbacher S, Coop G (2016) The Strength of Selection against Neanderthal Introgression.
*PLoS Genet* 12: e1006340

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL (2019) Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 37: 907-915

Korstanje R, O'Brien PC, Yang F, Rens W, Bosma AA, van Lith HA, van Zutphen LF, Ferguson-Smith
MA (1999) Complete homology maps of the rabbit (Oryctolagus cuniculus) and human by
reciprocal chromosome painting. *Cytogenet Cell Genet* 86: 317-322

Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra MA (2009)
Circos: an information aesthetic for comparative genomics. *Genome Res* 19: 1639-1645

Lawniczak MKN, Durbin R, Flicek P, Lindblad-Toh K, Wei X, Archibald JM, Baker WJ, Belov K, Blaxter
ML, Marques Bonet T *et al* (2022) Standards recommendations for the Earth BioGenome
Project. *Proc Natl Acad Sci U S A* 119

Levanen R, Kunnasranta M, Pohjoismaki J (2018a) Mitochondrial DNA introgression at the
northern edge of the brown hare (Lepus europaeus) range. *Ann Zool Fenn* 55: 15-24

Levänen R, Pohjoismäki J, Kunnasranta M (2019) Home ranges of semi-urban brown hares (Lepus
europaeus) and mountain hares (Lepus timidus) at northern latitudes. *Ann Zool Fenn* 56:
107–120

Levanen R, Thulin CG, Spong G, Pohjoismaki JLO (2018b) Widespread introgression of mountain
hare genes into Fennoscandian brown hare populations. *Plos One* 13

Li D, Liu CM, Luo R, Sadakane K, Lam TW (2015) MEGAHIT: an ultra-fast single-node solution for
large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* 31:
1674-1676

Li H (2018) Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34: 3094-
3100

Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform.
*Bioinformatics* 25: 1754-1760

Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling A, Amit I, Lajoie BR,
Sabo PJ, Dorschner MO *et al* (2009) Comprehensive mapping of long-range interactions
reveals folding principles of the human genome. *Science* 326: 289-293

Manni M, Berkeley MR, Seppey M, Zdobnov EM (2021) BUSCO: Assessing Genomic Data Quality
and Beyond. *Curr Protoc* 1: e323

Marques JP, Seixas FA, Farelo L, Callahan CM, Good JM, Montgomery WI, Reid N, Alves PC, Boursot
P, Melo-Ferreira J (2020) An Annotated Draft Genome of the Mountain Hare (Lepus timidus).
*Genome Biol Evol* 12: 3656-3662

Mazzoni CJ, Ciofi C, Waterhouse RM (2023) Biodiversity: an atlas of European reference genomes.
*Nature* 619: 252

Melo-Ferreira J, Vilela J, Fonseca MM, da Fonseca RR, Boursot P, Alves PC (2014) The elusive
nature of adaptive mitochondrial DNA evolution of an arctic lineage prone to frequent
introgression. *Genome Biol Evol* 6: 886-896

Meng G, Li Y, Yang C, Liu S (2019) MitoZ: a toolkit for animal mitochondrial genome assembly,
annotation and visualization. *Nucleic Acids Res* 47: e63

Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A (2018) Versatile genome assembly
evaluation with QUAST-LG. *Bioinformatics* 34: i142-i150

Minoudi S, Papapetridis I, Karaiskou N, Chatzinikos E, Triantaphyllidis C, Abatzopoulos TJ,
Triantafyllidis A (2018) Genetic analyses of brown hare (Lepus europaeus) support limited
migration and translocation of Greek populations. *Plos One* 13: e0206327

Ognev SI (1940) 157. Lepus (Eulagus) europeaus Pall. (1778) заяц-русак. In: *Mammals of the USSR*
*and adjacent countries: rodents (In Russian)*, pp. 112-161. Akademiya Nauk SSSR: Moscow

Open2C, Abdennur N, Fudenberg G, Flyamer IM, Galitsyna AA, Goloborodko A, Imakaev M, Venev
SV (2023) Pairtools: from sequencing data to chromosome contacts. *bioRxiv*

Ou S, Su W, Liao Y, Chougule K, Agda JRA, Hellinga AJ, Lugo CSB, Elliott TA, Ware D, Peterson T *et*
*al* (2019) Benchmarking transposable element annotation methods for creation of a
streamlined, comprehensive pipeline. *Genome Biol* 20: 275

Petrovan S Lepus europaeus (European hare) (2013) [DATASET]

Pohjoismaki JLO, Michell C, Levanen R, Smith S (2021) Hybridization with mountain hares
increases the functional allelic repertoire in brown hares. *Sci Rep* 11: 15771

Ranallo-Benavidez TR, Jaron KS, Schatz MC (2020) GenomeScope 2.0 and Smudgeplot for
reference-free profiling of polyploid genomes. *Nat Commun* 11: 1432

Reid N (2011) European hare (Lepus europaeus) invasion ecology: implication for the
conservation of the endemic Irish hare (Lepus timidus hibernicus). *Biol Invasions* 13: 559-
569

Reid N, Montgomery WI (2007) Is naturalisation of the brown hare in Ireland a threat to the
endemic Irish hare? *Biol Environ* 107b: 129-138

Rikala K (1925) Ajojahdilla [flush hunting of hares]. *Metsästys ja Kalastus* 2: 44-48

Schai-Braun SC, Schwienbacher S, Smith S, Hackländer K (2023) Coexistence of European hares
and Alpine mountain hares in the Alps: what drives the occurrence and frequency of their
hybrids? *Journal of Zoology* 320: 214-225

Schai-Braun SC, Weber D, Hacklander K (2013) Spring and autumn habitat preferences of active
European hares (Lepus europaeus) in an agricultural area with low hare density. *Eur J*
*Wildlife Res* 59: 387-397

Seluanov A, Vaidya A, Gorbunova V (2010) Establishing primary adult fibroblast cultures from
rodents. *J Vis Exp*

Siivonen L (1972) *Rusakko*. Kustannusosakeyhtiö Otava, Keuruu, Finland

Skov L, Coll Macia M, Lucotte EA, Cavassim MIA, Castellano D, Schierup MH, Munch K (2023)
Extraordinary selection on the human X chromosome associated with archaic admixture.
*Cell Genom* 3: 100274

Smit AFA, Hubley R, Green P, 2013-2015. RepeatMasker Open-4.0. http://www.repeatmasker.org.

Smith RK, Jennings NV, Harris S (2005) A quantitative analysis of the abundance and demography
of European hares Lepus europaeus in relation to habitat type, intensity of agriculture and
climate. *Mammal Rev* 35: 1-24

Stott P (2003) Use of space by sympatric European hares (Lepus europaeus) and European rabbits
(Oryctolagus cuniculus) in Australia. *Mamm Biol* 68: 317-327

Thulin C-G (2003) The distribution of mountain hares (Lepus timidus, L. 1758) in Europe: A
challenge from brown hares (L. europaeus, Pall. 1778)? *Mammal Rev* 33: 29-42

Thulin C-G, Tegelström H (2002) Biased geographical distribution of mitochondrial DNA that
passed the species barrier from mountain hares to brown hares (genus Lepus): an effect of
genetic incompatibility and mating behaviour? *J Zool* 258: 299-306
Thulin CG, Winiger A, Tallian AG, Kindberg J (2021) Hunting harvest data in Sweden indicate
precipitous decline in the native mountain hare subspecies Lepus timidus sylvaticus (heath
hare). *J Nat Conserv* 64
Uliano-Silva M, Ferreira J, Krasheninnikova K, Darwin Tree of Life C, Formenti G, Abueg L, Torrance
J, Myers EW, Durbin R, Blaxter M *et al* (2023) MitoHiFi: a python pipeline for mitochondrial
genome assembly from PacBio high fidelity reads. *BMC Bioinformatics* 24: 288
Vinogradov AE (1998) Genome size and GC-percent in vertebrates as determined by flow
cytometry: The triangular relationship. *Cytometry* 31: 100-109
Wheeler TJ, Eddy SR (2013) nhmmer: DNA homology search with profile HMMs. *Bioinformatics*
29: 2487-2489
Wolf JB, Lindell J, Backstrom N (2010) Speciation genetics: current status and evolving
approaches. *Philos Trans R Soc Lond B Biol Sci* 365: 1717-1733
Zhou C, McCarthy SA, Durbin R (2023) YaHS: yet another Hi-C scaffolding tool. *Bioinformatics* 39
732