Phylogeny and disease links of a widespread and ancient gut phage lineage

Patrick A. de Jonge[1,2,3], Bert-Jan H. van den Born[1,2,3], Aeilko H. Zwinderman[4], Max

Nieuwdorp[1], Bas E. Dutilh[5,6†], Hilde Herrema[1,2,3†]*

1: Department of Internal and Experimental Vascular Medicine; Amsterdam UMC; Location

AMC; University of Amsterdam; Amsterdam; the Netherlands.

2: Amsterdam Gastroenterology, Endocrinology & Metabolism; Endocrinology, metabolism &

nutrition; Amsterdam UMC; Amsterdam; the Netherlands.

3: Amsterdam Cardiovascular Sciences; Diabetes & Metabolism; Amsterdam UMC;

Amsterdam; the Netherlands.

4: Department of Clinical Epidemiology; Biostatistics and Bioinformatics; Amsterdam UMC;

Location AMC; University of Amsterdam; Amsterdam; the Netherlands.

5: Theoretical Biology and Bioinformatics; Science for Life; Utrecht University; Utrecht; the

Netherlands.

6: Institute of Biodiversity; Faculty of Biological Sciences; Cluster of Excellence Balance of

the Microverse; Friedrich-Schiller-University Jena; Jena; Germany.

[†]: These authors contributed equally.

*: Correspondence: h.j.herrema@amsterdamumc.nl

Abstract

Viruses are a core component of the human microbiome, impacting health and disease

through interactions with gut bacteria[1] and the immune system[2]. Most viruses in the human

microbiome are bacteriophages, which exclusively infect bacteria. Individual gut

bacteriophages can affect bacterial bile acid deconjugation[3], and can alter their infection

strategy based on dietary content[4]. Up to recently, most studies of the gut virome have

focused on low taxonomic scales (e.g., viral operational taxonomic units), hampering

population-level analyses. We previously identified the expansive and widespread

Candidatus *Heliusviridae* bacteriophage family in a cohort with inhabitants of Amsterdam,

the Netherlands. Here, we study their biodiversity and evolution in a wide variety of human

populations. With a more detailed phylogeny based on sequences from six viral genome

databases, we now propose the Candidatus order *Heliusvirales* to accommodate these

viruses. We identify *Ca. Heliusvirales* viruses in 80% of 5,441 individuals across 39 studies,

and also in thirteen out of sixteen analyzed metagenomes from ancient humans that lived in

Europe and North America between 1,000 and 5,000 years ago. We show that a large *Ca.*

Heliusvirales lineage has diversified starting with human dispersion across the globe some

50,000 years ago. Ancient peoples and modern hunter-gatherers further have distinct *Ca.*

Heliusvirales populations that are characterized by lower richness than modern urbanized

people. Within urbanized people, those suffering from type 1 and type 2 diabetes, as well as

inflammatory bowel disease, have higher *Ca. Heliusvirales* richness than healthy controls.

We thus conclude that these ancient core members of the human gut virome have thrived

with increasingly westernized lifestyles of the human population.

<u>Introduction</u>

Bacteriophages are the second most common biological entities in the human gut

microbiome, after their bacterial targets[5]. As obligate parasites of bacteria, phages play a

key role in regulating the structure and function of the gut microbiome in health and

disease[6]. For example, phage community alterations have been linked to diseases such as

inflammatory bowel disease[1,7], metabolic syndrome[8], and malnutrition[9]. Individual human gut

phages have further been found to mediate diet-induced bacterial lysis[4], bacterial bile acid

metabolism[3], and gut colonization[10]. But while phages are evidently crucial to the functioning

of the gut microbiome, their study at higher taxonomic levels was long restricted to

morphology- rather than phylogeny-based taxonomy.

This knowledge gap is partly due to the absence of a universal viral marker gene akin

to ribosomal RNA in cellular organisms. Bacteriophage taxonomy was originally based on

nucleic acid composition (with the majority being dsDNA viruses) and viral particle

morphology[11], with the bulk of known bacteriophage diversity being classified as

Siphoviridae, Myoviridae, or *Podoviridae.* These families, as well as the order *Caudovirales,*

proved to be so genomically diverse that they were recently abolished and reclassified in the

class *Caudoviricetes*[12,13]. A striking example of the discovery of higher taxonomic lineage

was the highly prevalent and abundant human gut phage crAssphage[14] and its relatives[15] in

metagenomics data. These viruses are now accepted as the phylogenetics-based

Crassvirales order by the International Committee for the Taxonomy of Viruses (ICTV). But

while the description of the *Crassvirales* is a step forward in the taxonomy of the human gut

virome, many other important lineages remain unclassified.

We recently identified the putative phage family *Ca. Heliusviridae* in a study of gut

virome perturbations associated with metabolic syndrome, a collection of risk factors for

cardiometabolic disease[8]. The expansive family was detected in over 90% of almost 200

study participants and may thus be part of a persistent core of human gut bacteriophage

lineages[16,17]. But as our previous study focused on a single cohort, it remains unknown how

widespread these phages are in the general population, as well as in other environments.

Furthermore, their relation to human health beyond metabolic syndrome is yet to be

elucidated. Their further adoption in virome studies necessitates a robust taxonomic

classification based on comprehensive phylogenetic analysis of sequences from diverse

sources[18–21].

Here, we present a comprehensive study of *Ca. Heliusviridae* proteomic and

phylogenetic diversity. Comparisons of member phages derived from several large

metagenomic databases show both the distinctness of this family from known ICTV

taxonomic lineages, and their phylogenetic structure. We further provide evidence that

increasing ecological richness of this phage family in the gut is linked to both urbanized

lifestyles and several (cardiometabolic) diseases.

<u>Results</u>

Helius-phages are ubiquitous in bacteriophage databases

We previously reported an expansive and widespread clade of human gut phages,

tentatively christened as the family *Ca. Heliusviridae*, after the HELIUS (Healthy Life In an

Urban Setting) cohort that we studied[8]. These phages shared nine marker genes: four

structural proteins (terminase large subunit or TerL, portal protein, major capsid protein, and

a head maturation protease), three replication-related ones (DNA polymerase I, helicase,

and nuclease), and two proteins of unknown function (**Figure 1a**)[8]. To study these phages

more extensively, we analyzed 1,314,988 contigs of over 25 kbp from seven phage

databases with profile hidden Markov models (HMMs) of all nine marker genes (**Figure 1b**).

Of the 111,534 contigs with at least one homolog to a marker gene, 2,322 were deemed

complete by checkv[22]. Notably, the 1,007 complete genomes with hits to all marker genes

distinctly had bit-scores of over 600 in homology searches against the terminase large

subunit (TerL) marker gene, while this threshold was seldomly passed by genomes carrying

fewer marker genes (**Figure 1c**). The TerL, which encodes an ATP-driven molecular motor

involved in genome translocation into the viral capsid, is often highly conserved among

Caudoviricetes phages and consequently frequently used as marker gene[13,15]. We thus

selected the 1,253 complete viral genomes with bit-scores of over 600 against the TerL

HMM for further study. Phage-specific gene annotation of all 1,253 genomes with pharokka[23]

refined the function of several marker genes: the DNA polymerase I was revealed to instead

be an exonuclease, while the ones of unknown function encode a DNA polymerase and

ssDNA-binding protein (**Figure 1a**). We next determined which genes were most common

among these phages by forming clusters of homologous proteins (i.e., protein clusters). The

only universal protein cluster was the TerL, and the next eight most common protein clusters

represented the other marker genes (**Figure 1d**, **Supplementary Table 1**). Genes with hits

against the HMMs of the major capsid protein, head maturation protease, helicase, and DNA

polymerase genes were spread out over multiple protein clusters, indicating some

heterogeneity in their sequences. Overall, all nine marker genes were found in 1000-1253 of

the genomes (**Supplementary Table 2**). Thus, the nine originally identified marker genes

are extremely common, though not universal, among *Ca. Heliusviridae.*

Helius-phages are taxonomically and ecologically distinct

We next sought to establish more detailed phylogenies of these phages. We previously

divided the *Ca. Heliusviridae* into three groups: *alpha, beta,* and *gamma*. In an approximate

maximum-likelihood tree of TerL genes, we again discerned three large well-supported

clades (Shimodaira–Hasegawa-like approximate likelihood ratio test[24] ≥80 and ultrafast

bootstrap approximation[25] ≥95, **Figure 2a**), but these did not completely align with the

previous putative groups. The largest clade contained all but one of the partial genomes

from the previously proposed *gamma*-group, but groups *alpha* and *beta* were spread out

across the other two clades. This reflects the two main clades in group *beta* that were

evident in the earlier analysis[8], while their different architecture in the TerL-based tree

versus the earlier concatenated tree likely reflects horizontal gene transfer of at least some

of the core genes among these groups. Furthermore, large subclades in all three clades

contained no representative of the earlier groups, and neither did a number of smaller clades

across the tree which were not part of the three main clades. In the face of this expanded

biodiversity, we set out to draw more detailed taxonomic boundaries.

The TerL tree also made it evident that these phages are better classified at the

order-level. The reasoning behind this is as follows: the three largest clades were divisible

into smaller clades that at minimum shared 5-57% of their protein content, too low for them

to be considered different viral genera (**Figure 2b, Supplementary Figure 1**)[26]. In line with

the viral taxonomic hierarchy[27], we classify these clades as subfamilies. The higher-level

clades are then necessarily families, and the entirety of the tree an order. Consequently, we

renamed the Ca. *Heliusviridae* as the Candidatus viral order *Heliusvirales.* Because Helios

was the ancient Greek solar deity, we named families after various solar deities. The three

largest families are the *Ca. Utuviridae* (Sumerian), *Ca. Aineviridae* (Irish), and *Ca.*

Hathorviridae (Egyptian). Several smaller well-supported and high-level clades were also

placed at the family rank due to their very low (<15%) shared protein content

(**Supplementary Figure 1)**. The subfamilies within the three largest families, and three small

high-level clades classified as subfamilies, all shared at least 30% of their protein content

with each other (**Figure 2b, Supplementary Figure 1)**. All subfamilies were named after

words for sun in languages from around the world (**Figure 2a, Supplementary Table 3**).

In order for a group of phages to be deemed a defined lineage, they need to be a

monophyletic group that is distinct from established taxonomy[26]. To establish this, we

performed proteomic-based analyses on the 1,237 complete genomes versus the national

center for biotechnological information (NCBI) reference sequence database[28]. In a genomic

protein-sharing network, our genomes formed a cluster separate from RefSeq phages

(**Supplementary Figure 2a**). This was confirmed in a phage proteomic tree, were they also

formed a separate monophyletic clade distinct from ICTV-recognized and RefSeq

sequences (**Supplementary Figure 2b**). Together, these results show that the *Ca.*

Heliusviriales are a cohesive lineage that is unrelated to any of the viral families that are

currently accepted by the ICTV.

A genomic network of the *Ca. Heliusvirales* showed that several subfamilies formed

distinct clusters, notably the second- and third-largest subfamilies: the *Ca. Mataharivirinae*

(*Ca. Hathorviridae*) and *Ca. Mehrvirinae* (*Ca. Aineviridae)* (**Figure 2c**). Their distinctness

was reflected by the environments in which they were found and the bacteria that they infect

(**Figure 3a and b**), as well as their divergent GC content when compared to other members

of their family (**Supplementary Figure 3**). Although animal-, terrestrial-, and aquatic-derived

phages dominate some *Ca. Heliusvirales* lineages*,* the *Lachnospiraceae* phages in the *Ca.*

Mataharivirinae were enriched in human gut-derived phages (hypergeometric test, *p* =

2.8×10^{-21}). Within this family, these phages seem to have a broad host range, as on average

they had CRISPR-spacer hits (0-1 mismatch) against 3.18±2.75 genera (median 2), with the

most common host predictions being *Acetatifactor sp900066365, Agathobacter faecis,* and

Blautia_A wexlerae_A (**Supplementary Figure 4**). The *Ca. Mehrvirinae* were similarly

enriched in phages from the human oral cavity (hypergeometric test, *p* = 1.5×10^{-17}). This

environmental preference was also reflected in their most common hosts: the facultatively
anaerobic *Streptococcaceae*. These two subfamilies are thus specialized for distinct
environments. Similarly, the *Ca. Nitavirinae* formed a separate cluster in the genomic
network, reflecting their hosts in the *Actinobacteriota* rather than the *Firmicutes* infected by
other *Ca. Heliusvirales*. So too did the distinctness of the *Ca. Zonviridae* reflect their hosts in
the *Negativicutes*. Unlike the recently defined *Crassvirales*, which all infect within the
Bacteroidetes phylum[29], the *Ca. Heliusvirales* as a collective thus have a very broad host
range across multiple phyla.

The majority of complete *Ca. Heliusvirales* genomes (56.9%) carried identifiable
genes involved with integration and excision, meaning they were temperate phages. This
also differed among the subfamilies, with notably higher amounts among the *Ca. Ra-* and
Mataharivirinae in the *Ca. Hathorviridae* (96.7% and 82% of genomes), as well as the *Ca.*
Mehrvirinae (85% of genomes, **Supplementary Figure 5**). Considerably lower fractions of
temperate phages were present among the subfamilies of the *Ca. Utuviridae* (overall
47.6%). It is thus possible that besides contrasting environments and host species, the
different *Ca. Heliusvirales* lineages also differ in lifestyles.

Putative auxiliary metabolic genes were very common among *Ca. Heliusvirales*,
being present in 78.1% of genomes (**Figure 3c**), though we found no clear family- or
subfamily specific auxiliary metabolic genes. Many were part of cysteine and methionine
metabolism, with the most common genes being DNA (cytosine-5)-methyltransferase 1
(K00558, present in 67% of genomes) and S-adenosylmethionine synthetase (K00789,
present in 58.9% of genomes). Both of these genes likely aid the phage in defending against
bacterial restriction-modification defenses[30,31]. After cysteine and methionine metabolism,
sulfur metabolism-associated genes were most commonly found. This is in line with previous
findings that bacteriophages can use sulfur metabolism to reprogram bacterial metabolism
for additional energy production during phage particle formation[32].

Helius-phages have been associated with humans since ancient times

Across our datasets, we found *Ca. Heliusvirales* TerL genes from every continent (**Figure**

4a), indicating that they are widespread. Only two countries had human gut-associated

samples without any detected *Ca. Heliusvirales*: El Salvador and Kazakhstan. Indeed,

analysis of 7,166 human gut metagenomes of 5,441 individuals from 38 studies[33] found *Ca.*

Heliusvirales TerL genes in 4,491 individuals (80%, **Figure 4b**). Outside the gut, prevalence

of *Ca. Heliusvirales* was markedly lower: we detected their presence in 18/54 skin samples,

17/61 vaginal samples, and 115/312 oral samples. While the *Ca. Heliusvirales* thus occur in

various body sites, they are especially prevalent in the gut. This is in line with the finding that

their most common hosts are in the *Firmicutes* (e.g., the *Lachnospiraceae*), which is one of

the main bacterial phyla in the human gut.

We hypothesized that high *Ca. Heliusvirales* prevalence reflected its ancient

association to the human lineage. To explore this, we re-analyzed whole metagenome

shotgun sequencing datasets from paleofeces in Austria[34] and North America[35], and

intestinal biopsies of Ötzi the iceman[36]. Among the North-American samples, we found 36

Ca. Heliusvirales TerL genes across all eight of the studied samples (**Figure 4a**). These

samples were from three different locations across the USA and Mexico, and were dated

between 10-920 CE. We furthermore identified four *Ca. Heliusvirales* TerL genes in small-

and large-intestinal samples of Ötzi the iceman, who lived around 3,200 BCE[37]. In the

Austrian samples, we however found only four TerL sequences in one of the four Austrian

samples, which was dated at between 652-544 BCE. Since all other samples from both

North America and Ötzi the iceman did contain *Ca. Heliusvirales*, their absence from 3/4

ancient Austrian samples may reflect higher levels of degradation or fragmentation of these

samples rather than their true absence among these individuals.

The presence of *Ca. Heliusvirales* in multiple pre-colonial North Americans imply that they

were part of the human gut microbiome before human migration to the Americas (about

15,000 years ago[38]). We confirmed this hypothesis with a time-measured phylogenetic tree

of the *Ca. Hathorviridae*, which we chose for this analysis because it contains the large and

highly human-gut-associated *Ca. Mataharivirinae*. A tree with ancient North-American

sequences and a selection of modern sequences that reflect *Ca. Hathorviridae* tree topology

(**Supplementary Figure 6**) showed that these phages began diversifying about 65,000

years ago, with a 95% highest posterior density (HPD) interval of 44,000-87,000 years ago

(**Figure 4c**). This is notably around the time that modern humans started to disperse

worldwide from Africa (40,000-60,000 years ago)[39]. Further diversification of the *Ca.*

Hathorviridae into its two largest subfamilies (*Ca. Mataharivirinae* and *Ca. Ravirinae*)

occurred about 34,000 years ago (95% HDP 24,000-45,000 years ago), indicating that these

phages adapted to specific human populations. Indeed, a concatenated tree of all universal

genes in the complete *Ca. Mataharivirinae*, showed notable clustering of sequences from

China (**Supplementary Figure 7**).

Ancient gut microbiomes have similar diversity to those of modern populations

consuming non-urbanized diets[35], which are more diverse than those consuming urbanized

diets[40,41]. By extension, this is likely true for the gut virome as well. To further explore the

relation between ancient and modern *Ca. Heliusvirales*, we compared them to samples

derived from urbanized and non-urbanized populations. To minimize interstudy batch-

effects, we selected two studies that included both hunter-gatherers (from Tanzania or Peru)

and urbanized people (from Italy or the USA)[40,41]. Within both studies, we observed a

significantly higher *Ca. Heliusvirales* richness among urbanized people (Wilcoxon signed-

rank test, Benjamini-Hochberg-adjusted *p*=0.026 and 0.01, **Supplementary Figure 8a**).

Combining the values of modern urbanized and hunter-gatherer populations with ancient

ones showed significantly higher *Ca. Heliusvirales* richness in urbanized people than both

hunter-gatherers (linear mixed-effect model, Benjamini-Hochberg-adjusted *p*=4.5x10^{-5}) and

ancients (linear mixed-effect model, Benjamini-Hochberg-adjusted *p*=1.2x10^{-5}), but not

between hunter-gatherers and ancients (linear mixed-effect model, Benjamini-Hochberg-

adjusted *p*=0.076, **Figure 4d**). This interestingly is the reverse of overall gut bacterial

richness, which decreases with urbanization[42]. Abundance, measured as the fraction of

reads that mapped to *Ca. Heliusvirales* TerL genes, was not significantly different between

the three populations (**Supplementary Figure 8b**). *Ca. Heliusvirales* thus seem to diversify

in tandem with urbanization.

We next analyzed *Ca. Heliusvirales* β-diversity among urbanized, hunter-gatherer,

and ancient samples with a phylogenetic tree of TerL genes from these samples and those

from complete genomes as depicted in Figure 2a. Non-metric multidimensional scales

(NMDS) on unweighted UniFrac distances intriguingly showed that all three populations

(urban, hunter-getherers, ancients) had *Ca. Heliusvirales* populations that were distinct from

each other permanova *q*=0.001, **Figure 4e**). Among ancient samples, the *Ca. Xiheviridae*,

Ca. Mithraviridae, Ca. Tilevirinae, and *Ca. Ghrianvirinae* were all more prevalent than in both

hunter-gatherers and urban populations (Fisher's exact test, Benjamini-Hochberg-adjusted *p*

< 0.05, **Figure 4f**). Meanwhile, the *Ca. Sontsevirinae* and *Ca. Dzuwavirinae* were more

prevalent, and the *Ca. Kuarahyvirinae* and *Ca. Nayviriane* were less prevalent among

modern urban populations. With the exception of the *Ca. Ghrianvirinae*, which target

Bacteroidacaeae, these significantly altered lineages notably did not seem to infect aberrant

hosts than the other subfamilies. These results are, however, at least partially caused by the

greater sequencing depth of the ancient samples. In fact, taxonomic richness expressed as

the number of observed *Heliusvirales* subfamilies per billion sequenced bases, showed the

same pattern as overall richness (**Supplementary figure 8c**). Yet despite a higher

sequencing depth, ancient samples were notably mostly lacking the *Ca. Mataharivirinae*,

which is the largest of the subfamilies. This intriguingly suggests that this subfamily in

particular is associated with modern and urbanized human gut microbiomes. Our results

could reflect that *Ca. Heliusvirales* hosts in the *Lachnospiraceae* thrive in the urbanized gut

microbiome, or, since the samples we analyzed are not virome-specific, that these phages

have increased rates of lysogenation.

Helius-phage richness is associated with disease

Our previous research focused on gut virome alterations in metabolic syndrome (MetS). To

study a wider range of illnesses, we now identified *Ca. Heliusvirales* sequences in

assemblies of fourteen studies of diverse human-derived samples[33] by searching for the

TerL marker gene. Within-study analyses of richness alterations identified significantly

altered *Ca. Heliusvirales* richness in four out of twelve illnesses (Wilcoxon signed-rank test

Benjamini-Hochberg adjusted p < 0.05): type 1 diabetes (T1D), type 2 diabetes (T2D),

inflammatory bowel disease (IBD), and liver cirrhosis (**Figure 5a).** In the former three,

richness was elevated when compared to healthy controls, while in liver cirrhosis it was

decreased. These disease linkages are interesting in light of the *Lachnospiraceae* hosts of

many *Ca. Heliusvirales*, because this bacterial family includes species involved in short-

chained fatty acid production (e.g., *Roseburia intestinalis, Anaerobutyricum hallii,*

Coprococcus eutactus, and *Blautia wexlerae* **Supplementary Figure 4**)[43], which are often

associated with the healthy gut. Indeed, increased *Lachnospiraceae* abundance in T1D-[44]

and T2D-patients[45] has been reported, though a study of in vitro incubations in a relatively

small cohort size (n=26) linked IBD to decreased *Lachnospiraceae* abundance[46]. For the

former two diseases, this could explain the greater richness and prevalence in *Ca.*

Heliusvirales phages among such patients. The alterations in the *Ca. Heliusvirales* order are

notably counter to ASV-level analysis of the entire gut virome, which was found to be

unchanged in T1D[47], T2D[48], and IBD[1]. This underscores the value of analyzing higher

taxonomic ranks in the context of the gut virome.

Besides differences in *Ca. Heliusvirales* richness, the populations of these phages

also differed in each of the four studies (unweighted UniFrac, PERMANOVA Benjamini-

Hochberg adjusted *p* < 0.05, **Figure 5b**). These population-wide differences between

healthy controls and people with T1D, T2D, IBD, and cirrhosis expressed themselves in

distinct patterns of prevalence at the subfamily level (**Figure 5c**). The *Ca. Uternevirinae* in

particular were significantly more prevalent among T1D patients than healthy controls, but

more prevalent in controls than IBD and cirrhosis patients (Fisher's exact test, Benjamini-

Hochberg adjusted *p* < 0.05). Additionally, the *Ca. Nayvirinae* were more prevalent in

controls in the IBD cohort, but less so in the T2D cohort, while the *Ca. Oorunvirinae* were

more prevalent in controls in cirrhosis and less so in T2D. These subfamilies are all part of

the *Ca. Utuviridae*, consistent with our previous linkages of viruses in this family to metabolic

syndrome[8]. Thus, while some of these results could be explained by technical differences

(e.g., different sample sizes between controls and T1D patients), the findings across five

different cohorts linking members of the *Ca. Utuviridae* to different cardiometabolic diseases

robustly show they are associated with a wide range of disease states. The *Ca. Nayvirinae*

were notably less likely temperate than other *Ca. Heliusvirales* (31% temperate phage

versus 56.9% overall, **Supplementary Figure 5**), as where the *Ca. Atenviridae*. Their

association with healthy over IBD patients is thus in line with previous findings that this

disease is correlated with a loss of lytic phages[1]. The *Ca. Nayvirinae* were also depleted in

urbanized populations compared to hunter-gatherer and ancient ones (**Figure 4f**), perhaps

indicating that they are the viral component of VANISH (volatile and/or associated negatively

with industrialized societies of humans) gut microbiome taxa[49]. Conversely, its positive

associations with T1D (non-significant) and T2D (significant) points toward a role in disease,

underscoring the complex interrelations between the human gut virome and health.

Discussion

Elucidating higher-order structures among gut phages is essential for gaining a deeper

understanding of the viromes in the human gut and beyond. This process is underway with

the re-evaluation of viral taxonomy along genomic lines[26]. Recently, the prevalent

Crassvirales phage order has undergone the arduous process of classification[29,50], and here

we add the *Ca. Heliusvirales* order as a second genome-based classification of phages

widespread in the human gut. For this, we used a combination of marker genes and

proteomics-based approaches, though our phylogeny of the TerL gene was the main driver

for taxonomic classification. As this gene is well-conserved among the *Duplodnaviria*, to

which dsDNA bacteriophages belong[51], we considered the phylogeny built with this gene is

the most reliable method for defining *Ca. Heliusvirales* taxonomy. Our usage of the TerL

gene to define taxonomy is akin to the *Crassvirales*[12], but unlike the concatenated phylogeny

of marker genes used to define the *Herelleviridae*[13]. Naturally, the absence of any other

universal marker gene beside the TerL gene among the *Ca. Heliusvirales* invalidates a

concatenated phylogeny approach. This absence of universal marker genes also indicates

that relatively large amounts of horizontal gene transfer events occur within the *Ca.*

Heliusvirales, meaning that they may be interesting targets to study evolutionary dynamics of

human gut viruses.

The *Ca. Heliusvirales* order is highly diverse in environmental origin, geographic

distribution, and bacterial hosts. While we found *Ca. Heliusvirales* phages in terrestrial,

marine, and gut microbiomes of both cattle and wild animals, the majority of genomes

derived from human gut samples. While at face-value this may be assigned to

anthropocentric sampling bias, the IMGVR database included in our analysis contains about

six times more aquatic and terrestrial phages than human-derived ones[52]. Furthermore, non-

animal-associated *Ca. Heliusvirales* phages are largely consigned to specific families,

indicating environmental differentiation.

Ca. Heliusvirales phages associated with the human gut were identified in countries

from every permanently inhabited continent, and in about 80% of the population. This

percentage could well be an underestimation, as shallower sequencing depths may lead to

false negatives. In either case, this order is clearly a core component of the human gut

virome. This has been the case since the deep past of modern humans, as evidenced by the

presence of these phages in palaeofaeces and our accompanying evolutionary studies. We

dated the diversification of the *Ca. Hathorviridae* family roughly to the dispersal of human

populations out of Africa 60,000 years ago[39]. This dating is notably more recent than the

previously studied diversification of the human-associated bacterial species

Methanobrevibacter smithii[35] and *M.oralis*[53] at respectively 85,000 and 126,000 years ago.

During their rapid evolution, these phages perhaps diversified in response to microbiome

alterations influenced by changing human geography and diet, both main drivers of human

microbiome composition[54,55]. Earlier studies showed co-linearity of *Crassvirales* between

humans and great apes[56], and estimated a diversification of these phages in the past several

centuries[57]. This estimate of much more recent diversification was, however, noted for its

uncertainty and focused only on the *crAssphage* species rather than the family-level analysis

presented here.

We also found that higher *Ca. Heliusvirales* richness is associated with increased

urbanization. This is remarkable, as microbiome diversity as a whole is lower with increasing

urbanization[40,41]. The opposite observations in *Ca. Heliusvirales* could reflect the success of

their bacterial hosts to replicate in urbanized microbiomes: indeed, Hadza hunter-gatherers

from Tanzania and Matses hunter-gatherers from Peru have lower levels of

Lachnospiraceae bacteria such as *Blautia* and *Ruminococcus* than people in urbanized

environments[41,58].

While the *Ca. Heliusvirales* as a whole are near universally present in the human gut,

no single viral genome in this putative order is. The linkages to lifestyle and disease

demonstrated here are thus only evident when studying higher viral taxonomic levels. In line

with the recent reassessment of viral taxonomy along genomic lines[12], this study thus shows

how future definition of additional viral orders and classes can provide key insights into the

ecology and disease-associations of the gut virome.

<u>References</u>

1. Clooney, A. G. *et al.* Whole-Virome Analysis Sheds Light on Viral Dark Matter in Inflammatory

Bowel Disease. *Cell Host Microbe* **26**, 764-778.e5 (2019).

2. Fluckiger, A. *et al.* Cross-reactivity between tumor MHC class I–restricted antigens and an

enterococcal bacteriophage. *Science* **369**, 936–942 (2020).

3. Campbell, D. E. *et al.* Infection with Bacteroides Phage BV01 Alters the Host Transcriptome and

Bile Acid Metabolism in a Common Human Gut Microbe. *Cell Rep.* **32**, 108142 (2020).

4. Oh, J.-H. *et al.* Dietary Fructose and Microbiota-Derived Short-Chain Fatty Acids Promote

Bacteriophage Production in the Gut Symbiont Lactobacillus reuteri. *Cell Host Microbe* **25**, 273-

284.e6 (2019).

5. Shkoporov, A. N. & Hill, C. Bacteriophages of the Human Gut: The "Known Unknown" of the

Microbiome. *Cell Host Microbe* **25**, 195–209 (2019).

6. Cao, Z. *et al.* The gut virome: A new microbiome component in health and disease. *eBioMedicine*

81, 104113 (2022).

7. Norman, J. M. *et al.* Disease-Specific Alterations in the Enteric Virome in Inflammatory Bowel

Disease. *Cell* **160**, 447–460 (2015).

8. de Jonge, P. A. *et al.* Gut virome profiling identifies a widespread bacteriophage family associated

with metabolic syndrome. *Nat. Commun.* **13**, 3594 (2022).

9. Reyes, A. *et al.* Gut DNA viromes of Malawian twins discordant for severe acute malnutrition.

Proc. Natl. Acad. Sci. **112**, 11941–11946 (2015).

10.   Brown, E. M. *et al.* Gut microbiome ADP-ribosyltransferases are widespread phage-encoded

fitness factors. *Cell Host Microbe* **29**, 1351-1365.e11 (2021).

11.   Simmonds, P. *et al.* Four principles to establish a universal virus taxonomy. *PLOS Biol.* **21**,

e3001922 (2023).

12.   Turner, D. *et al.* Abolishment of morphology-based taxa and change to binomial species

names: 2022 taxonomy update of the ICTV bacterial viruses subcommittee. *Arch. Virol.* **168**, 74

(2023).

13.   Barylski, J. *et al.* Analysis of Spounaviruses as a Case Study for the Overdue Reclassification

of Tailed Phages. *Syst. Biol.* **69**, 110–123 (2020).

14.   Dutilh, B. E. *et al.* A highly abundant bacteriophage discovered in the unknown sequences of

human faecal metagenomes. *Nat. Commun.* **5**, 4498 (2014).

15.   Koonin, E. V. & Yutin, N. The crAss-like Phage Group: How Metagenomics Reshaped the

Human Virome. *Trends Microbiol.* **28**, 349–359 (2020).

16.   Shkoporov, A. N. *et al.* The Human Gut Virome Is Highly Diverse, Stable, and Individual

Specific. *Cell Host Microbe* **26**, 527-541.e5 (2019).

414 17. Manrique, P. *et al.* Healthy human gut phageome. *Proc. Natl. Acad. Sci.* **113**, 10400–10405

415 (2016).

416 18. Camarillo-Guerrero, L. F., Almeida, A., Rangel-Pineros, G., Finn, R. D. & Lawley, T. D. Massive

417 expansion of human gut bacteriophage diversity. *Cell* **184**, 1098-1109.e9 (2021).

418 19. Gregory, A. C. *et al.* The Gut Virome Database Reveals Age-Dependent Patterns of Virome

419 Diversity in the Human Gut. *Cell Host Microbe* **28**, 724-740.e8 (2020).

420 20. Tisza, M. J. & Buck, C. B. A catalog of tens of thousands of viruses from human metagenomes

421 reveals hidden associations with chronic diseases. *Proc. Natl. Acad. Sci.* **118**, e2023202118 (2021).

422 21. Roux, S. *et al.* IMG/VR v3: an integrated ecological and evolutionary framework for

423 interrogating genomes of uncultivated viruses. *Nucleic Acids Res.* **49**, D764–D775 (2021).

424 22. Nayfach, S. *et al.* CheckV assesses the quality and completeness of metagenome-assembled

425 viral genomes. *Nat. Biotechnol.* **39**, 578–585 (2021).

426 23. Bouras, G. *et al.* Pharokka: a fast scalable bacteriophage annotation tool. *Bioinformatics* **39**,

427 btac776 (2023).

428 24. Guindon, S. *et al.* New Algorithms and Methods to Estimate Maximum-Likelihood

429 Phylogenies: Assessing the Performance of PhyML 3.0. *Syst. Biol.* **59**, 307–321 (2010).

430 25. Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2: Improving

431 the Ultrafast Bootstrap Approximation. *Mol. Biol. Evol.* **35**, 518–522 (2018).

432 26. Turner, D., Kropinski, A. M. & Adriaenssens, E. M. A Roadmap for Genome-Based Phage

433 Taxonomy. *Viruses* **13**, 506 (2021).

434 27. Gorbalenya, A. E. *et al.* The new scope of virus taxonomy: partitioning the virosphere into 15

435 hierarchical ranks. *Nat. Microbiol.* **5**, 668–674 (2020).

436 28. O'Leary, N. A. *et al.* Reference sequence (RefSeq) database at NCBI: current status,

437 taxonomic expansion, and functional annotation. *Nucleic Acids Res.* **44**, D733–D745 (2016).

29.   Smith, L., Goldobina, E., Govi, B. & Shkoporov, A. N. Bacteriophages of the Order

Crassvirales: What Do We Currently Know about This Keystone Component of the Human Gut

Virome? *Biomolecules* **13**, 584 (2023).

30.   Samson, J. E., Magadán, A. H., Sabri, M. & Moineau, S. Revenge of the phages: defeating

bacterial defences. *Nat. Rev. Microbiol.* **11**, 675–687 (2013).

31.   Guo, X. *et al.* Structure and mechanism of a phage-encoded SAM lyase revises catalytic

function of enzyme family. *eLife* **10**, e61818.

32.   Howard-Varona, C. *et al.* Phage-specific metabolic reprogramming of virocells. *ISME J.* **14**,

881–895 (2020).

33.   Pasolli, E. *et al.* Extensive Unexplored Human Microbiome Diversity Revealed by Over

150,000 Genomes from Metagenomes Spanning Age, Geography, and Lifestyle. *Cell* **176**, 649-

662.e20 (2019).

34.   Maixner, F. *et al.* Hallstatt miners consumed blue cheese and beer during the Iron Age and

retained a non-Westernized gut microbiome until the Baroque period. *Curr. Biol.* **31**, 5149-

5162.e6 (2021).

35.   Wibowo, M. C. *et al.* Reconstruction of ancient microbial genomes from the human gut.

Nature **594**, 234–239 (2021).

36.   Maixner, F. *et al.* The 5300-year-old Helicobacter pylori genome of the Iceman. *Science* **351**,

162–165 (2016).

37.   Bonani, G., Ivy, S. D., Hajdas, I., Niklaus, T. R. & Suter, M. Ams 14C Age Determinations of

Tissue, Bone and Grass Samples from the Ötztal Ice Man. *Radiocarbon* **36**, 247–250 (1994).

38.   Willerslev, E. & Meltzer, D. J. Peopling of the Americas as inferred from ancient genomics.

Nature **594**, 356–364 (2021).

39.   Bergström, A., Stringer, C., Hajdinjak, M., Scerri, E. M. L. & Skoglund, P. Origins of modern

human ancestry. *Nature* **590**, 229–237 (2021).

40.   Rampelli, S. *et al.* Metagenome Sequencing of the Hadza Hunter-Gatherer Gut Microbiota.

Curr. Biol. CB **25**, 1682–1693 (2015).

41.   Obregon-Tito, A. J. *et al.* Subsistence strategies in traditional societies distinguish gut

microbiomes. *Nat. Commun.* **6**, 6505 (2015).

42.   Sonnenburg, E. D. & Sonnenburg, J. L. The ancestral and industrialized gut microbiota and

implications for human health. *Nat. Rev. Microbiol.* **17**, 383–390 (2019).

43.   Vacca, M. *et al.* The Controversial Role of Human Gut Lachnospiraceae. *Microorganisms* **8**,

573 (2020).

44.   Kostic, A. D. *et al.* The Dynamics of the Human Infant Gut Microbiome in Development and

in Progression toward Type 1 Diabetes. *Cell Host Microbe* **17**, 260–273 (2015).

45.   Qin, J. *et al.* A metagenome-wide association study of gut microbiota in type 2 diabetes.

Nature **490**, 55–60 (2012).

46.   Sasaki, K. *et al.* Construction of a Model Culture System of Human Colonic Microbiota to

Detect Decreased Lachnospiraceae Abundance and Butyrogenesis in the Feces of Ulcerative

Colitis Patients. *Biotechnol. J.* **14**, 1800555 (2019).

47.   Zhao, G. *et al.* Intestinal virome changes precede autoimmunity in type I diabetes-

susceptible children. *Proc. Natl. Acad. Sci.* **114**, E6166–E6175 (2017).

48.   Ma, Y., You, X., Mai, G., Tokuyasu, T. & Liu, C. A human gut phage catalog correlates the gut

phageome with type 2 diabetes. *Microbiome* **6**, 24 (2018).

49.   Fragiadakis, G. K. *et al.* Links between environment, diet, and the hunter-gatherer

microbiome. *Gut Microbes* **10**, 216–227 (2019).

50.   Yutin, N. *et al.* Discovery of an expansive bacteriophage family that includes the most

abundant viruses from the human gut. *Nat. Microbiol.* **3**, 38–46 (2018).

51.   Krupovic, M., Dolja, V. V. & Koonin, E. V. The LUCA and its complex virome. *Nat. Rev.*

Microbiol. **18**, 661–670 (2020).

52.   Camargo, A. P. *et al.* IMG/VR v4: an expanded database of uncultivated virus genomes

within a framework of extensive functional, taxonomic, and ecological metadata. *Nucleic Acids*

Res. **51**, D733–D743 (2023).

53.   Weyrich, L. S. *et al.* Neanderthal behaviour, diet, and disease inferred from ancient DNA in

dental calculus. *Nature* **544**, 357–361 (2017).

54.   He, Y. *et al.* Regional variation limits applications of healthy gut microbiome reference

ranges and disease models. *Nat. Med.* **24**, 1532–1535 (2018).

55.   Zmora, N., Suez, J. & Elinav, E. You are what you eat: diet, health and the gut microbiota.

Nat. Rev. Gastroenterol. Hepatol. **16**, 35–56 (2019).

56.   Edwards, R. A. *et al.* Global phylogeography and ancient evolution of the widespread human

gut virus crAssphage. *Nat. Microbiol.* **4**, 1727–1736 (2019).

57.   Honap, T. P. *et al.* Biogeographic study of human gut-associated crAssphage suggests

impacts from industrialization and recent expansion. *PLOS ONE* **15**, e0226930 (2020).

58.   Schnorr, S. L. *et al.* Gut microbiome of the Hadza hunter-gatherers. *Nat. Commun.* **5**, 3654

(2014).

59.   Snijder, M. B. *et al.* Cohort profile: the Healthy Life in an Urban Setting (HELIUS) study in

Amsterdam, The Netherlands. *BMJ Open* **7**, e017873 (2017).

60.   Nayfach, S. *et al.* Metagenomic compendium of 189,680 DNA viruses from the human gut

microbiome. *Nat. Microbiol.* **6**, 960–970 (2021).

61.   Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site

identification. *BMC Bioinformatics* **11**, 119 (2010).

62.   Eddy, S. R. Accelerated Profile HMM Searches. *PLOS Comput. Biol.* **7**, e1002195 (2011).

63.   Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for the

analysis of massive data sets. *Nat. Biotechnol.* **35**, 1026–1028 (2017).

64.   Kieft, K., Zhou, Z. & Anantharaman, K. VIBRANT: automated recovery, annotation and

curation of microbial viruses, and evaluation of viral community function from genomic

sequences. *Microbiome* **8**, 90 (2020).

65.   Nishimura, Y. *et al.* ViPTree: the viral proteomic tree server. *Bioinformatics* **33**, 2379–2380

(2017).

66.   Bin Jang, H. *et al.* Taxonomic assignment of uncultivated prokaryotic virus genomes is

enabled by gene-sharing networks. *Nat. Biotechnol.* **37**, 632–639 (2019).

67.   Shannon, P. *et al.* Cytoscape: A Software Environment for Integrated Models of Biomolecular

Interaction Networks. *Genome Res.* **13**, 2498–2504 (2003).

68.   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree

display and annotation. *Nucleic Acids Res.* **49**, W293–W296 (2021).

69.   Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973 (2009).

70.   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference

in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

71.   Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.

ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**, 587–

589 (2017).

72.   Almeida, A. *et al.* A unified catalog of 204,938 reference genomes from the human gut

microbiome. *Nat. Biotechnol.* **39**, 105–114 (2021).

73.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics **34**, i884–i890 (2018).

74.   Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P. A. metaSPAdes: a new versatile

metagenomic assembler. *Genome Res.* **27**, 824–834 (2017).

75.   Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25 (2009).

76.   Borry, M., Hübner, A., Rohrlach, A. B. & Warinner, C. PyDamage: automated ancient damage

identification and estimation for contigs in ancient DNA de novo assembly. *PeerJ* **9**, e11845

(2021).

77.   Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

78.   Bouckaert, R. *et al.* BEAST 2.5: An advanced software platform for Bayesian evolutionary

analysis. *PLOS Comput. Biol.* **15**, e1006650 (2019).

79.   Bouckaert, R. R. OBAMA: OBAMA for Bayesian amino-acid model averaging. *PeerJ* **8**, e9460

(2020).

<u>Methods</u>

Data

All bioinformatic tools were used with default settings, unless explicitly stated otherwise.

Statistical analyses were done in R v4.2.1. All p-values were adjusted for multiple testing

with the Benjamini-Hochberg method.

The primary dataset of *Heliusviridae* marker genes were derived from 298 contigs

that were used in our earlier study[8], which studied the gut virome in the general population-

based multi-ethnic Healthy Life in an Urban Setting (HELIUS) cohort[59]. Contig datasets that

were used to identify *Heliusviridae* consisted of 63,481 genomes from this previous study.

This included all viral contigs before removing redundant ones. Further data sources

consisted of four recently constructed databases of gut bacteriophages: the gut phage

database (GPD, 142,809 genomes)[18], cenote-taker 2 human virome database (CHVD,

93,860 genomes)[20], gut virome database (GVD, 33,242 genomes)[19], and metagenomic gut

virus catalogue (MGV, 189,680 genomes)[60]. Further contigs were derived from fourth

version of the integrated microbial genomes viral dataset (IMG/VR, 15,722,824 genomes)[21]

and bacterial viruses from the NCBI viral reference sequence dataset (RefSeq, 4,220

genomes) release 216. For studies of urbanization- and disease-links, metagenome

assemblies were obtained from Pasolli *et al*[33]. of 7,166 human gut metagenomes of 5,441

individuals from 38 studies.

Identification of putative Heliusviridae sequences

Since previous findings showed *Heliusviridae* genomes to be around 50,000-100,000 bp

long, we first selected contigs of at least 25,000 bp without any ambiguous bases from all

datasets. This resulted in an effective search space of 1,314,988 contigs. Open reading

frames (ORFs) were then predicted using prodigal v2.6.3[61] (option --meta).

For each of the nine *Heliusviridae* marker genes, a profile hidden Markov model

(HMM) was constructed from protein sequences derived from our previous study using

hmmbuild v3.3[62]. The resulting profile HMMs were used to search against predicted ORFs

using hmmsearch v3.3[62]. For phylogenetic studies, we selected contigs with an hmmsearch

hit (bit-score ≥600) against the *Heliusviridae* terminase large subunit (TerL). Duplicates were

removed with dedupe from BBTools v38.84 (option minidentity=100), resulting in a non-

redundant dataset of 6,359 contigs. These were then analyzed for completeness with checkv

v1.0.1[22] (using database version 1.4), and we selected contigs that had 100% completeness,

no warnings, and were not recognized as prophages. These genomes were fully annotated

with pharokka v1.2.1[23] (options -m, -g prodigal), and their ORFs were grouped in protein

clusters using mmseqs2 v14.7e284[63]. GC contents were derived from pharokka output, and

temperate phages were defined as those with at least one ORF with a predicted function in

the "integration and excision" category. Potential auxiliary metabolic genes encoded by the

genomes were identified with VIBRANT v1.2.1[64].

Viral sequence clustering and proteomic analysis

Separation between our phages and those in the NCBI refseq database was analyzed with

both ViPTreeGen v1.1.3[65] and vContact2 v0.11.3[66]. The vContact2 network was visualized

with Cytoscape v3.9.1[67], while the ViPTreeGen tree was visualized with the interactive tree

of life (iTOL) webtool[68]. To analyze the grouping of genomes among the high quality contigs,

we used them to constructed a second vContact2 network.

Phylogenetic tree construction

Before phylogenetic tree construction, incomplete protein sequences were removed by

selecting only sequences that were longer than the 25th percentile minus 1.5 times the inter-

quantile ranges of all lengths. As outgroup, the homologous TerL sequence from

Staphylococcus phage LH1 (NC055041) was added. Marker genes were aligned using

MAFFT v4.753 (options --maxiterate, --localpair), after which positions with more than 90%

gaps were trimmed with trimal v1.4.rev15[69] (option -gt 0.1). Finally, a tree was built using

IQTree v2.2.0.3[70], using model finder[71] and 1000 iterations of the ultrafast bootstrap

approximation[25] and SH-like approximate likelihood ratio test. IQTree performed ten

separate tree iterations and selected the one with the best log likelihood score (options -B

1000 -alrt 1000 --runs 10). The tree was visualized with the interactive tree of life (iTOL)

webtool[68].

Heliusvirales hosts

We determined the hosts infected by the *Heliusvirales* with two complementary methods,

adapted from previous work[60]. Firstly, we identified phage-host links by matching CRISPR

spacers. For this, we downloaded 1,846,441 CRISPR spacers identified[60] among the

Bacteria and Archaea in the Unified Human Gastrointestinal Genomes (UHGG) dataset[72].

We then created a subset of 1,742,522 spacers with lengths between 30 and 40 bp and

without undefined bases. These we used for a blastn search against *Heliusvirales* genomes

(options -task blastn-short). Hits were finally filtered for those with 0-2 mismatches across

the entire length. Our second host-linking approach focused on identifying phages as

prophages. We downloaded all 286,997 bacterial and archaeal genomes from the Unified

Human Gastrointestinal Genomes (UHGG) dataset[72]. Viral genomes were then used for a

blastn search against these genomes, after which hits with over 96% identity and a length of

over 1000 bp were selected. Prophage and CRISPR-spacer hits were combined and the

most common phage-host link was determined for each viral genome. If one bacterial

lineage was responsible for 50+% of all phage-host linkage hits, it was considered as the

host of that phage.

Analysis of ancient samples

For analysis of ancient metagenomes, reads were downloaded from two separate

studies[34,35]. These were trimmed with fastp v.0.23.2[73] (options --detect_adapter_for_pe) and

error-corrected with tadpole from the bbmap package v38.90 (https://jgi.doe.gov/data-and-

tools/bbtools, options mode=correct, ecc=t, prefilter=2). Error corrected reads were

assembled with metaSPAdes v3.15.5[74]. The ancient origins of assembled contigs were

determined by mapping reads to contigs of ≥1,000 bp with bowtie2 v2.4.2[75], and determining

C-to-T degradation at the 5' end of sequencing reads with PyDamage v0.70[76]. Contigs with a

predicted accuracy of over 0.5 were considered of ancient origins.

Heliusvirales prevalence and relation to urbanization

To determine *Heliusvirales* prevalence, we firstly gathered geographic and environmental

metadata from all genomes used to build the main phylogenetic tree. Second, contigs

obtained from Pasolli et al[33] were analyzed for the presence of *Heliusvirales* TerL genes. To

do this, ORFs were predicted with Prodigal v2.6.3[61] (option –meta), and used for an

hmmsearch with the profile hmm of the TerL constructed from sequences identified in our

previous study. Hits with bit-scores of over 600 were selected. The same approach was

used to identify *Heliusvirales* TerL genes among ancient populations. Enrichment of phages

from certain environments among selected subfamilies was calculated with the phyper

function from base R.

To analyze the relation of *Heliusvirales* richness to urbanization, richness was

calculated by dividing the number of identified TerL genes by the number of reads in billions.

Significance testing of richness differences after combining studies was done with linear

mixed effects models where the study was entered as a random effect, using the lme

function from the nlme R package v3.1-162 and Anova function from the rstatix R package

v0.7.2. The same approach was used to calculate significant richness differences among

pooled disease-related cohorts.

Assignment of TerL sequences from ancient- and disease-related samples was done

by constructing a phylogenetic tree with TerL genes from all high-quality *Heliusvirales*

genomes that had been used in the main phylogeny. For the ancient, hunter-gatherers, and

urban populations, a single such tree was built, while for the disease-related datasets one

tree was constructed per project. Trees were constructed with IQTree v2.2.0.3 as described

above, and pairwise cophenetic distances were calculated from them with the ape R

package v5.7-1. Each sequence was then assigned to the same taxonomy as its closest

neighbor of which the taxonomy was known.

For the ancient, hunter-gatherers, and urban populations, taxonomic assignment at

the family level was used to calculate UniFrac distances, from which a non-metric

multidimensional scaling (NMDS) was constructed. Permutated ANOVA significance levels

were calculated with the adonis2 function in the vegan R package v2.6-4. To determine

significant differences in subfamily prevalence between populations with different lifestyles,

Fisher exact tests were calculated using the fisher.test function from base R.

Timed tree construction

To reconstruct an evolutionary timed tree of the *Ca. Hathorviridae*, we first constructed a

regular tree on members of this family. For this, we selected the TerL genes of all *Ca.*

Hathorviridae from both the complete genomes and the ancient samples. This was used to

build a phylogenetic tree with IQTree as described above. From this tree, we selected

thirteen complete genomes and all three ancient genome fragments that combined reflected

the architecture of the *Ca. Hathorviridae.* All ORFs from these genomes were then clustered

with mmseqs2 v14.7e284[63], after which the six protein clusters that were universally present

were selected. This number was relatively low because the ancient sequences were genome

fragments of 15,076, 19,664, and 67,881 bp long. The six universal PCs were a TerL, head

maturation protease, major head protein, portal protein, a hypothetical protein, and

terminase small subunit. For each protein, an alignment was made with MAFFT v4.753

(options --maxiterate, --localpair)[77], from which a timed tree was inferred using BEAST

v2.7.3[78]. BEAST was run with the following models: Birth death model, Coalescent constant

population, Coalescent exponential population, Coalescent bayesian skyline, Coalescent

extended bayesian skyline, each until the estimated sample size was above 200. The

coalescent exponential population was the best-fitting model. BEAST was run with OBAMA

v1.1.1[79] for amino-acid model averaging.

Acknowledgements

P.A.d.J. was supported by a Talent Development grant of the Amsterdam Gastroenterology,

Endocrinology, and Metabolism research school at the Amsterdam UMC. MN was supported

by a personal ZONMW-VICI grant 2020 (09150182010020). HH was supported by an

Aspasia premium (015.017.050) and Dutch Diabetes Research Foundation (2019.82.004).

The HELIUS study is conducted by the Amsterdam University Medical Centers, location

AMC and the Public Health Service of Amsterdam. Both organizations provided core support

for HELIUS. The HELIUS study is also funded by the Dutch Heart Foundation, the

Netherlands Organization for Health Research and Development (ZonMw), the European

Union (FP-7), and the European Fund for the Integration of non-EU immigrants (EIF). We

are most grateful to the participants of the HELIUS study and the management team,

research nurses, interviewers, research assistants and other staff who have taken part in

gathering the data of this study.

Author contributions

P.A.d.J. conducted data analysis and wrote the manuscript; B.J.v.d.B., A.H.Z., and M.N. assisted with experimental design and data interpretation; P.A.d.J., B.E.D., and H.H. designed the study. All authors read and provided input on the manuscript.

Competing interests

M.N. owns stock in, consults for, and has intellectual property rights in Caelus Health. He consults for Kaleido. None of these are directly relevant to the current paper. The remaining authors declare no competing interests.

Figure legends

Figure 1: Presence of near-universal marker genes is linked to TerL homology. **a:** Annotation of NatCom_15117, a complete genome with inverse terminal repeats. The nine marker genes are denoted above the genome line, with re-annotated functions in parentheses. Other annotated genes are below the genome line. **b:** Overview of the procedure by which genomes were selected from seven phage databases. **c:** density plots of bit-scores from a search with the TerL HMM, separated by the total amount of marker genes found in the genome. Dots are individual complete genomes. **d:** prevalence of the most common protein clusters. Red denotes marker genes, grey others.

Figure 2: the *Ca. Heliusvirales* are a large order of gut phages with distinct subgroups. **a:** a rooted approximate maximum-likelihood tree of *Ca. Heliusvirales* TerL genes. The outgroup

at the extreme left-hand side of the tree is *Triavirus LH1* (NC_055041). **b:** chart depicting the

pairwise percent of shared genes among the three main families (top) and their subfamilies

(bottom). The n reflects the number of genomes in the lineage. For an overview of all

families and subfamilies, see **Supplementary Figure 1**. Box plots show the median (middle

line), 25th, and 75th percentile (box), with the 25th percentile minus and the 75th percentile

plus 1.5 times the interquartile range (whiskers), and outliers (single points). **c:** protein-

sharing network of all sequences in **a**. Node edge color denotes family, while fill denote

subfamilies, both according to the same colors as use in **a**.

Figure 3: high *Ca. Heliusvirales* diversity in environmental origin and host species. **a:**

environmental origins of the *Ca. Heliusvirales* subfamilies. **b:** bacterial host assignment of

Ca. Heliusvirales subfamilies. **c:** pathway assignment of putative auxiliary metabolic genes

(AMGs) and morons encoded by the *Ca. Heliusvirales*. For brevity, the 25 pathways that

were most commonly found were selected.

Figure 4: the *Ca. Heliusvirales* are widespread and ancient. **a:** map showing in which

countries *Ca. Heliusvirales* TerL-containing sequences were found. Data pertains to the

seven bacteriophage databases described in the methods. Red dots show the locations of

ancient gut microbiomes that were analyzed for *Ca. Heliusvirales* TerL presence, with their

dates, gene-, and sample-counts. **b:** prevalence of *Ca. Heliusvirales* TerL genes among

individuals of 39 gut metagenome studies. The red bar is the prevalence among the totality

of all studies. **c:** tip-dated phylogenetic tree of *Ca. Hathorviridae* sequences from modern

and ancient samples. Bold sequences are from ancient samples. Bar plots denote 95%

confidence, values denote mean and 95% confidence intervals **d:** boxplots of the number of

Ca. Heliusvirales TerL genes found per person, per billion sequenced bases, with jittered

points separated by study. Significance is according to nested linear mixed-effect models

where the study was used as a random effect. Box plots show the median (middle line),

25th, and 75th percentile (box), with the 25th percentile minus and the 75th percentile plus

1.5 times the interquartile range (whiskers). Comparisons of richness in urbanized vs hunter-

gatherer populations within each study can be found in Supplementary Figure 2. **e:** non-

metric multidimensional scaling (NMDS) on unweighted UniFrac distances, colored by

lifestyle. **f:** Prevalence of *Ca. Heliusvirales* subfamilies in ancient peoples, hunter-gatherers,

and urbanized peoples, with significance according to Fisher exact tests for subfamilies with

significant values. All p-values were adjusted for multiple testing with the Benjamini-

Hochberg procedure, and are denoted as follows * ≤ 0.05, ** ≤0.01, *** ≤ 0.001, **** ≤

0.0001. Lineages without any representatives among the samples were not displayed.

Figure 5: disease associations of *Ca. Heliusvirales* phages in gut microbiomes. **a:** four

studies in which a significant difference was found in the number of *Ca. Heliusvirales* TerL

genes per person per billion sequenced bases. Significance is according to Wilcoxon signed-

rank tests. Box plots show the median (middle line), 25th, and 75th percentile (box), with the

25th percentile minus and the 75th percentile plus 1.5 times the interquartile range

(whiskers). **b:** non-metric multidimensional scaling (NMDS) on unweighted UniFrac

distances, colored by disease status. **c:** the same tree as in **Figure 1a**, but with all

subfamilies collapsed. Boxes show the difference in prevalence between controls and

people with disease. All p-values were adjusted for multiple testing with the Benjamini-

Hochberg procedure.

Supplementary Figure 1: shared gene content for all families (top) and subfamilies (bottom).

Supplementary Figure 2: **a**: vContact2 genome network of the *Ca. Heliusvirales* and all viral

sequences in the NCBI RefSeq database. **b:** VipTree proteome tree of the same sequences

in **a**.

Supplementary Figure 3: GC-content of the *Ca. Heliusvirales* by subfamily.

Supplementary Figure 4: the most common host predictions in the six largest subfamilies.

Supplementary Figure 5: the percentage of temperate phages across the subfamilies, as

determined by the presence of integration and excision genes in pharokka annotations. The

dashed red line is the average for all *Ca. Heliusvirales*

Supplementary Figure 6: An approximate maximum-likelihood tree of *Ca. Hathorviridae* TerL

genes, with color coding conform genome origin. Dots show branches with SH-like

approximate likelihood ratio test ≥80 and ultrafast bootstrap approximation ≥95.

Supplementary Figure 7: An approximate maximum-likelihood tree of *Ca. Matahariviridae*

TerL genes, with color coding conform genome origin. Dots show branches with SH-like

approximate likelihood ratio test ≥80 and ultrafast bootstrap approximation ≥95.

Supplementary Figure 8: **a**: comparisons of *Ca. Heliusvirales* richness between urban and

hunter-gatherer populations in two studies. Q-values are according to Wilcoxon signed-rank

tests. **b:** comparison of *Ca. Heliusvirales* TerL abundance between lifestyles. **c:** taxonomic

Ca. Heliusvirales richness comparison between ancient, hunter-gatherer, and urban

populations. Q-values according to linear mixed-effect models with the study as a random

effect. All p-values are adjusted for multiple testing with the Benjamini-Hochberg method.

a

DNA, RNA and nucleotide metabolism
moron, auxiliary metabolic gene and host takeover
head and packaging
tail
others/unknown

5 kb

DUF2815
(ssDNA binding)
DNA pol.
(exonucl.)
DUF2800
(DNA pol.)
helicase
nuclease
(endonucl.)
TerL
portal
protease
major capsid

transcr. regulator
transcr. activator
toxin
endonucl.
partition nucl.
GGCT
toxin
tail sheath
tail prot.
tail prot.
toxin
tape measure
tail prot.
tail prot.
baseplate
tail prot.
endolysin

b

IMG/VR CHVD GPD GVD MGV RefSeq deJonge

1,314,988 contigs ≥25 kbp

111,534 contigs ≥1 marker gene

2,322 complete genomes

1,253 ≥600 bit TerL

c

marker genes in genome

TerL bitscore

d

protein cluster

PC_1 — TerL
PC_2 — portal protein
PC_3 — major capsid protein
PC_4 — helicase
PC_5 — exonuclease
PC_6 — ssDNA binding protein
PC_7 — endonuclease
PC_8 — DNA polymerase
PC_9 — head maturation protease
PC_10 — holin
PC_11 — DNA methyltransferase
PC_12 — head-tail adaptor Ad1
PC_13 — integrase
PC_14 — hypothetical protein
PC_56 — head maturation protease
PC_70 — DNA polymerase
PC_122 — helicase
PC_172 — major capsid protein
PC_230 — major capsid protein
PC_3709 — major capsid protein

genomes

a

family
- *Aineviridae*
- *Atenviridae*
- *Auroraviridae*
- *Gunanaviridae*
- *Hathorviridae*
- *Hunahpuviridae*
- *Hyperionviridae*
- *Intiviridae*
- *Mithraviridae*
- *Shapeshviridae*
- *Utuviridae*
- *Xiheviridae*
- unclassified

subfamily
- *Antuvirinae*
- *Aurinkovirinae*
- *Dzuwavirinae*
- *Ghrianvirinae*
- *Gunesvirinae*
- *Haevirinae*
- *Haulvirinae*
- *Iavirinae*
- *Ilangavirinae*
- *Kuarahyvirinae*
- *Mataharivirinae*
- *Mehrvirinae*
- *Nayvirinae*
- *Nitavirinae*
- *Oorunvirinae*
- *Quyashvirinae*
- *Ravirinae*
- *Shamsvirinae*
- *Sikinikvirinae*
- *Sontsevirinae*
- *Taiyangvirinae*
- *Tilevirinae*
- *Uternevirinae*
- *Zonvirinae*

de Jonge 2023
- Alpha
- Beta
- Gamma

• SH-aLRT ≥80 & UFboot ≥95

de Jonge 2023 subfamily family

b

family

- *Utuviridae* (n=593)
- *Hathorviridae* (n=162)
- *Aineviridae* (n=287)

subfamily

- *Tilevirinae* (n=22)
- *Taiyangvirinae* (n=64)
- *Shamsvirinae* (n=17)
- *Quyashvirinae* (n=14)
- *Panonpoevirinae* (n=8)
- *Oorunvirinae* (n=61)
- *Nitavirinae* (n=41)
- *Nayvirinae* (n=29)
- *Gunesvirinae* (n=36)
- *Ghrianvirinae* (n=31)
- *Gevirinae* (n=10)
- *Dzuwavirinae* (n=42)
- *Aurinkovirinae* (n=126)
- *Arawvirinae* (n=48)
- *Antuvirinae* (n=33)
- *Ravirinae* (n=30)
- *Mataharivirinae* (n=128)
- *Zonvirinae* (n=66)
- *Mehrvirinae* (n=107)
- *Ilangavirinae* (n=10)
- *Haulvirinae* (n=88)

homologous proteins (%)

0   25   50   75   100

c

Mataharivirinae

Mehrvirinae

Nitavirinae

a

Arid West cave
595–635 CE
12 TerL in 3/3 samples

Hallstatt
1040 BCE
2 TerL in 1/4 samples

Boomerang shelter
10–60 CE
10 TerL in 2/2 samples

Ötzi
3230 BCE
4 TerL in 4/4 samples

Zape
725–920 CE
14 TerL in 3/3 samples

Ca. Heliusvirales
TerL found in:
- only human
- only non-human
- both human and non-human
- none detected/no data

b

Study (top to bottom):
DavidLA-2015, SmitsSA-2017, VatanenT-2016, HanniganGD-2017, OlmMR-2017, RampelliS-2015, LomanNJ-2013, AsnicarF-2017, FerrettiP-2018, QinN-2014, VincentC-2016, LiJ-2017, ChengpingW-2017, LiuW-2016, QinJ-2012, SchirmerM-2016, NielsenHB-2014, ZeeviD-2015, total average, LoombaR-2017, HMP-2012, LeChatelierE-2013, BackhedF-2015, KosticAD-2015, Obregon-TitoAJ-2015, BritoIL-2016, YuJ-2015, ZellerG-2014, KarlssonFH-2013, IjazUZ-2017, CosteaPI-2017, LiJ-2014, FengQ-2015, VogtmannE-2016, XieH-2016, RaymondF-2016, LouisS-2016, LiSS-2016, Bengtsson-PalmeJ-2015

individuals with *Ca. Heliusvirales*

c

SRR12557711_NODE_2296
NatCom_59710
NatCom_24345
IMGVR_15533314
CHVD_93067
IMGVR_15563536   *Ca. Mataharivirinae*
SRR12557733_NODE_95
IMGVR_6929766
IMGVR_15498387

IMGVR_15670726
IMGVR_15355857
NatCom_26447   *Ca. Ravirinae*
SRR12557705_NODE_1404
IMGVR_516903
IMGVR_4510279
IMGVR_15430980

29.4 [39.3–20.4]
33.6 [45.1–23.5]
64.9 [86.8–44.5]
15.9 [21.5–11.0]
58.5 [79.6–39.5]

time (thousands of years)

d

TerL/1B bases

0.076
1.3×10⁻⁵
4.5×10⁻⁵

ancient   h.-g.   urban

e

NMDS2 vs NMDS1

- ancient
- hunter-gatherer
- urban

PERMANOVA	R²	q
urban vs. h.-g.	0.087	0.001
urban vs. ancient	0.121	0.001
ancient vs. h.-g.	0.061	0.001

f

present: yes / no

ancient | hunter-gatherer | urban | a./h.-g. | a./u. | u./h.-g.

Aineviridae: Zonvirinae *, Mehrvirinae, Haulvirinae
Atenviridae: Atenvirinae
Hathorviridae: Ravirinae, Mataharivirinae, Hyperionvirinae
Mithraviridae * *
Sontsevirinae, Sikinikvirinae, Haevirinae **
Utuviridae: Uternevirinae *, Tilevirinae *** **, Taiyangvirinae * **, Quyashvirinae, Oorunvirinae, Nayvirinae * *, Kuarahyvirinae *** **, Gunesvirinae **, Ghrianvirinae *** ** *, Dzuwavirinae, Aurinkovirinae, Antuvirinae * **
Xiheviridae ** **

a

Heliusvirales

b

Tree scale: 0.1

Ca. Heliusvirales
ICTV taxa
(other) RefSeq genomes

NatCom 59754
IMGVR 15341856
CHVD 19045
IMGVR 15348144
IMGVR 15609269
IMGVR 15498387
IMGVR 9172387
IMGVR 15373252
IMGVR 6929766
CHVD 14936
IMGVR 15563536
IMGVR 15605860
CHVD 31162
IMGVR 15580039
IMGVR 15595382
IMGVR 15509441
IMGVR 14221724
IMGVR 14243793
CHVD 53930
IMGVR 15364386
IMGVR 14193823
CHVD 14685
NatCom 31361
CHVD 93067
IMGVR 1427301
CHVD 1518
IMGVR 14246410
IMGVR 14197755
IMGVR 14217481
IMGVR 15603243
IMGVR 15343016
IMGVR 14290941
IMGVR 14211640
CHVD 2900
IMGVR 15695648
IMGVR 14323260
CHVD 20962
IMGVR 15587674
IMGVR 15621192
CHVD 36908
IMGVR 14220269
IMGVR 14150610
IMGVR 14311113
IMGVR 15414627
IMGVR 15408941
IMGVR 15698826
IMGVR 15486627
IMGVR 15381572
IMGVR 15593580
IMGVR 15453176
CHVD 34414
IMGVR 15388483
IMGVR 15663995
CHVD 36340
IMGVR 15683585
IMGVR 15563120
NatCom 29765
IMGVR 15672362
IMGVR 15601699
NatCom 43249
IMGVR 15532807
IMGVR 15527989
IMGVR 9124972
IMGVR 15672172
IMGVR 15660478
IMGVR 15701077
IMGVR 15620626
IMGVR 15575799
IMGVR 15576548
NatCom 40364
IMGVR 15414642
NatCom 6238
CHVD 3628
CHVD 1848
CHVD 35621
CHVD 6839
CHVD 21130
IMGVR 15592815
IMGVR 15601958
IMGVR 15694224
IMGVR 15590078
CHVD 1926
IMGVR 15615779
CHVD 36939
IMGVR 15358895
IMGVR 2863652
CHVD 33808
NatCom 13861
IMGVR 15389782
CHVD 1682
CHVD 34914
IMGVR 3101438
IMGVR 15634752
CHVD 14102
NatCom 10489
IMGVR 15638643
IMGVR 15570843
NatCom 59739
CHVD 21456
CHVD 2868
CHVD 41559
NatCom 59710
CHVD 2215
CHVD 19302
CHVD 10992
NatCom 24345
IMGVR 15627977
IMGVR 15641467
CHVD 30457
IMGVR 15623193
IMGVR 15639336
IMGVR 15671427
IMGVR 15528566
IMGVR 15625545
CHVD 15722
CHVD 14001
CHVD 16261
IMGVR 15533314
IMGVR 15378578
IMGVR 15671396
IMGVR 15644778
NatCom 22372
IMGVR 15635960
IMGVR 15603481
CHVD 13426
IMGVR 15494557
CHVD 21473

0.1

continent

Africa
Asia
Europe
North America
Unknown origin

b