Altered reactivity to threatening stimuli in *Drosophila* models of Parkinson's disease, revealed by a trial-

based assay

Márton Kajtor[1], Viktor A. Billes[2,3], Bálint Király[1,4], Hannah Stabb[1], Katalin Sviatkó[1,5], Eszter Ujvári[1], Diána

Balázsfi[1], Sophie E. Seidenbecher[6], Duda Kvitsiani[7], Tibor Vellai[2,3], Balázs Hangya[1,*]

1 Lendület Laboratory of Systems Neuroscience, Institute of Experimental Medicine, H-1083 Budapest,

Hungary

2 Department of Genetics, Eötvös Loránd University, H-1117 Budapest, Hungary

3 ELKH-ELTE Genetic Research Group, H-1117 Budapest, Hungary

4 Department of Biological Physics, Eötvös Loránd University, H-1117 Budapest, Hungary

5 János Szentágothai Doctoral School of Neurosciences, Semmelweis University, H-1085 Budapest,

Hungary

6 University of Technology Nuremberg, Nuremberg, Germany

7 Department of Molecular Biology and Genetics, Danish Research Institute of Translational Neuroscience,

Aarhus University, Aarhus, Denmark

* Correspondence: Balázs Hangya, hangya.balazs@koki.hu

Abstract

The fruit fly *Drosophila melanogaster* emerges as an affordable, genetically tractable model of behavior

and brain diseases. However, despite the surprising level of evolutionary conservation from flies to

humans, significant genetic and circuit-level differences hinder the interpretability of fruit fly models for

human disease. Therefore, to facilitate fly-to-human translation with more direct behavior-level

comparison, we surveyed the rarely-exploited, rich behavioral repertoire of fruit flies with genetic

alterations relevant to Parkinson's disease (PD). Flies displayed variable behaviors, including freezing,

slowing and running, in response to predator-mimicking passing shadows used as threatening stimuli in a

single-animal trial-based assay. We found that the expression of human mutant Parkin in flies resulted in

reduced walking speed and decreased reactivity to passing shadows. Flies with dopamine receptor

mutations showed similar alterations, consistent with the motor and cognitive deficits typical in humans

with PD. However, *Drosophila* overexpressing the human form of α-synuclein manifested in only

moderate phenotypical alterations, suggesting that other fruit fly models may be favored in PD research.

We also found age-dependent trends in behavioral choice across the fly lifespan, while dopamine receptor

mutant flies maintained their decreased general reactivity throughout all age groups. Our data

demonstrate that single-trial behavioral analysis can reveal subtle behavioral changes in mutant flies that

can be used to further our understanding of disease pathomechanisms and help gauge the validity of

genetic *Drosophila* models of neurodegeneration.

Introduction

Parkinson's disease (PD) affects over 6 million people worldwide, representing the second most common

neurodegenerative disease. PD is slowly progressing and typically leads to years of aggravating disability,

thereby placing a huge burden on families, health care systems and the society, measured in hundreds of

billions of dollars annually (Olesen et al., 2012; Obeso et al., 2017; Przedborski, 2017; Bloem et al., 2021).

Therefore, a hitherto elusive disease-modifying therapy is of prominent priority, envisioned to be fueled

by research into basic mechanisms of the disease.

Patients with PD display a diverse set of motor and non-motor symptoms with an underlying progressive

loss of midbrain dopaminergic neurons (Schapira, 2009; Schapira et al., 2017; Bloem et al., 2021),

mechanisms of which are widely studied in animal models of the disease. Primate models offer the

advantage of more direct translatability but are restricted to hard-to-handle neurotoxin approaches and

do not promise fast progress. Unlike the genetic mouse models of Alzheimer's disease, most of the

successful rodent models of PD are also toxin-based, making it difficult to exploit and further our

understanding of the genetic bases of the disease (Cannon and Greenamyre, 2010; Bové and Perier, 2012;

Breger and Fuzzati Armentero, 2019). This left a niche for the genetically tractable, affordable fruit flies

as genetic models of PD (Feany and Bender, 2000; Guo, 2012; Hewitt and Whitworth, 2017), building on

the homologies between the vertebrate basal ganglia and the fruit fly central complex (Strausfeld and

Hirth, 2013). However, substantial genetic, anatomical, physiological, and behavioral discrepancies

between insects and mammals call for better validation methods of fruit fly models for human disease.

To address this, here we used a single-animal trial-based behavioral assay that facilitates fine-grained

assessment of phenotypical behavioral changes in fruit flies with mutations relevant to understanding

human PD.

Genes linked to familial forms of PD may serve as an ideal basis for genetic disease models. Notably, mutations in the *PARK2* gene, which encodes the Parkin protein involved in maintaining mitochondrial integrity, are associated with autosomal-recessive forms of PD (Guo, 2012). Parkin loss-of-function mutant flies were found to have advanced mitochondrial aging, structural mitochondrial damage and a consequential selective loss of dopaminergic neurons (Cackovic et al., 2018), leading to motor deficits assessed by a climbing assay (Chambers et al., 2013; Cackovic et al., 2018), as well as non-motor PD phenotypes including memory deficits (Julienne et al., 2017). Mutations in the *SNCA* gene of α-Synuclein (α-Syn) are also associated with familiar forms of PD (Polymeropoulos et al., 1997) and α-Syn has been shown to accumulate in Lewy-bodies and Lewy-neurites (Spillantini et al., 1997). α-Syn proteins have been found in the pre-synaptic terminals in humans and mice (Kahle et al., 2000) and are thought to be involved in dopamine (DA) synthesis under physiological conditions by reducing tyrosine hydroxylase activity (Perez et al., 2002). Expression of human α-Syn in flies has been proposed as a genetic model of PD, showing age-dependent loss of dopaminergic neurons and locomotor dysfunction in a climbing assay (Feany and Bender, 2000; Haywood and Staveley, 2006). However, other studies found normal locomotion and dopaminergic cell counts in these flies, casting doubts on the validity of this model (Pesah et al., 2005; Nagoshi, 2018).

Despite the observed homology between mammalian and fruit fly DA systems in motor control and the establishment of *Drosophila* PD models based on human genetic information derived from familial PD patients, the role of *Drosophila* DA receptors in locomotor control is not well characterized. Nevertheless, it has been demonstrated that the D1-like DA receptor mediates ethanol-induced locomotion in the ellipsoid body (Kong et al., 2010). Furthermore, *Dop1R1* has been shown to be involved in turning behavior for goal-directed locomotion (Kottler et al., 2019) and startle-induced negative geotaxis (Sun et al., 2018). To shed light on the possible roles of DA receptors in threat-induced motor behaviors, we tested the behavioral responses of three dopamine receptor (*Dop1R1*, *Dop1R2* and *DopEcR*) insertion mutant lines to predator-mimicking passing shadows and compared them to established fruit fly PD models with partially known locomotor deficits.

We found that flies expressing the R275W mutant allele of human Parkin ('Parkin flies') showed slower average locomotion speed, which, in contrast, was not characteristic of flies expressing the A53T mutant allele of the human *SNCA* gene ('α-Syn flies'). Parkin flies also showed less frequent freezing and generally less behavioral reactivity to passing shadows compared to controls, whereas α-Syn flies showed increased durations of stopping after the stimuli. Dopamine receptor mutant flies showed reduced speed and less

behavioral reactivity similar to Parkin flies. *Dop1R1* mutant flies exhibited more pronounced behavioral

alterations than the other two receptor mutants in most of the parameters tested. These data

demonstrate that mutations in DA receptor genes lead to specific patterns of behavioral deficits in

Drosophila; hence, these dopamine receptor paralogs may have different functions in behavioral control.

The modest phenotype of A53T α-Syn compared to Parkin flies suggests that the latter should be favored

as a genetic model of human PD, at least with respect to the motor deficits examined in this study. We

further propose that single-trial analyses such as those we present here help us gain a better

understanding of the behavioral changes in fruit fly models of PD and are strong tools for validating

Drosophila models of human diseases.

Results

A single-animal trial-based assay to test behavioral responses to predator-mimicking passing shadows

We designed a behavioral apparatus to examine the responses of individual flies to predator-mimicking

passing shadows. To do this, we designed a transparent plexiglass arena (Fig. 1a), featuring 13 tunnels (53

mm x 5 mm) for simultaneous tracking of 13 individual flies. The height of the tunnels (1 mm) was designed

to allow free walking in two dimensions but prevent jumps and flight. To simulate predatory threat, we

created passing 'shadows' (Fig. 1a, right; see Methods) with a sliding red screen presented on a 10.1-inch

display placed on the top of the arena. A high frame-rate camera was placed under the arena (Fig. 1a,

left), allowing us to simultaneously record the movement of the animals and the shadows. All 13 tunnels

housed a single fly in each session, enabling the collection of single-animal data from 13 flies in parallel.

The locomotion of individual flies was tracked by custom software developed using the Bonsai visual

programming environment (Fig. S1, see Methods). We recorded 40-minute-long sessions, consisting of 40

trials of 2-second-long shadow presentations separated by pseudorandom inter-trial-intervals to prevent

the animals from learning temporal expectations of the shadow presentations (Fig. 1b).

Fruit flies exhibit a rich behavioral repertoire upon threatening stimuli, including freezing (or stopping)

and various escape behaviors such as jumping, slow or fast take-off and running, modulated by walking

speed at the time of the threatening stimulus (Zacarias et al., 2018). To study these behaviors, we

calculated the speed and acceleration of fruit flies based on the tracked x-y position of their center of

mass (Fig. 2a) and aligned these signals to the presentations of the shadow stimuli. To identify separate

response types to the threating stimuli, we applied hierarchical clustering on these stimulus-aligned speed

traces (see Methods). This analysis consistently revealed three main stereotypical behavioral responses across flies in addition to the trials where no significant response was evoked (Fig. 2b). Since jumping and flying was not possible in the arena, fruit flies were restricted to choose among freezing, slowing (also observed in (Zacarias et al., 2018)) and running. While this clustering approach was sufficient to reveal response types qualitatively, cluster boundaries were sensitive to genotype-specific differences across groups of flies (Király and Hangya, 2022). Therefore, for rigorous group comparisons, we determined exact definitions of each response type based on fly speed and acceleration (Fig. 2c). Reactions were considered robust if the absolute value of the acceleration reached 200 mm/s^2; otherwise, the trial was classified as a 'no reaction' trial. We considered the reaction as a 'stop' if the animal was moving before shadow presentation and its speed decreased to zero in the first second relative to the shadow presentation. If fly speed decreased to a non-zero value, the trial was defined as 'slow down', and if the fly accelerated after the shadow presentation, the trial was classified as 'speed up' (see also Methods).

Parkinson's model and dopamine receptor mutant fruit flies showed reduced walking speed and decreased reactivity to threatening stimuli

To generate PD fly models, we expressed the mutant human Parkin (275W) and α-Syn (A53T) coding transgenes (*UAS-Parkin-275W* and *UAS-α-Syn-A53T*, respectively), applying the UAS-Gal4 system (Duffy, 2002). Transgenes were driven by *Ddc-Gal4* inducing the expression of the correspondent human mutant proteins in dopaminergic and serotoninergic fly neurons. Parkin and α-Syn flies were compared to control animals from the same genetic background without mutant transgenes (*Ddc-Gal4* animals were crossed with *iso w^1118^*, the examined F1 generation flies are referred to as *iso w^1118^* Fig. 3) or mutants overexpressing GFP (Fig. S2). The same level of eye pigmentation and vision of the compared genotypes was achieved by the prior replacement of the *w** mutant first chromosome of the applied *Ddc-Gal4* stock for that of the wild-type . The dopamine receptor mutant groups were compared to their parental strains without the mutations (*y^1^ w^67c23^* served as control for *Dop1R1* and *Dop1R2* and *w^1118^* for *DopEcR*; Fig. 3).

To test whether mutant flies showed differences in overall locomotion independent of the threatening stimuli, we analyzed average fly speed in the 200 ms time windows before stimulus presentation. We found that Parkin flies showed reduced mean speed compared to controls which expressed the Gal4 driver alone (24.96% reduction, p = 6.55 × 10^{-6}, Mann-Whitney U-test; Fig. 3a, top), while α-Syn flies did not show significant reduction (p = 0.799, Mann–Whitney U-test). We observed similar changes in dopamine

receptor mutant flies, where the *Dop1R1* and *DopEcR* defective lines showed a robust mean speed decrease compared to the control groups (*Dop1R1*, 19.68% decrease compared to $y^1 w^{67c2}$, p = 0.0016; *DopEcR*, 32.97% decrease compared to w^{1118}, p = 0.0034; Fig. 3a, bottom), while the *Dop1R2* mutant flies only showed a non-significant speed decrease (21.73% compared to $y^1 w^{67c23}$, p = 0.1009, Mann–Whitney U-test).

Next, we tested whether mutant flies showed a difference in their reaction to threatening stimuli. We found that PD model flies showed line-specific alterations in their freezing behavior. Parkin flies froze after stimuli less frequently, stopping in 37.72% of trials compared to the 43.48% observed in the *iso w^{1118}* animals (p = 0.0043, Mann Whitney U-test; Fig. 3b, top). However, in the stop trials, they showed normal duration of pauses in locomotion (p = 0.8018 compared to *iso w^{1118}*, Mann Whitney U-test; Fig. 3c, top). In contrast, α-Syn flies showed an unchanged stopping frequency when compared to controls (p = 0.0768 compared to *iso w^{1118}*, Mann-Whitney U-test), but their stop durations showed a large increase (116.98% increase compared to *iso w^{1118}*, p = 2.18×10^{-11}, Mann-Whitney U-test). Interestingly, in contrast to PD model flies, dopamine receptor mutant flies did not show significant differences in their freezing behavior relative to controls (stop proportion: *Dop1R1*, p = 0.0631 compared to $y^1 w^{67c23}$; *Dop1R2*, p = 0.1838 compared to $y^1 w^{67c23}$; *DopEcR*, p = 0.1445 compared to w^{1118}; Mann-Whitney U-test; Fig. 3b, bottom; stop duration: *Dop1R1*, p = 0.893 compared to $y^1 w^{67c23}$; *Dop1R2*, p = 0.154 compared to $y^1 w^{67c23}$; *DopEcR*, p = 0.3576 compared to w^{1118}; Mann-Whitney U-test; Fig. 3c, bottom).

α-Syn flies showed a reduced aptitude to increase their speed, or 'run', upon encountering threatening stimuli (26.67% decrease compared to *iso w^{1118}*, p = 8.4×10^{-6}, Mann-Whitney U-test; Fig. 3d, top). Similar results were found in *Dop1R1* (25% decrease compared to $y^1 w^{67c23}$, p = 0.0122; Mann-Whitney U-test; Fig. 3d, bottom) and *Dop1R2* (41.67% decrease compared to $y^1 w^{67c23}$, p = 0.0024, Mann-Whitney U-test), but not in *DopEcR* mutant flies (compared to w^{1118}, p = 0.6598, Mann-Whitney U-test). Frequency of slowing, i.e., reducing their speed without freezing in a full stop, was moderately decreased in Parkin (14.38% decrease compared to *iso w^{1118}*, p = 0.0723, Mann-Whitney U-test; Fig. 3e, top) and *DopEcR* mutant flies (15.28% decrease compared to w^{1118}, p = 0.020, Mann-Whitney U-test; Fig. 3e, bottom).

Overall, all mutations tested resulted in decreased reactivity to threatening stimuli, confirmed by a significant increase in the proportion of trials where no reactions were detected (Fig. 3f). This effect was significant for Parkin flies (30% increase compared to *iso w^{1118}*, p = 0.0043, Mann-Whitney U-test), as well as for *Dop1R1* (50% increase compared to $y^1 w^{67c23}$, p = 0.0173, Mann-Whitney U-test) and *DopEcR* mutant flies (50% increase compared to w^{1118}, p = 0.0167, Mann-Whitney U-test), but not for α-Syn (α-Syn vs. *iso*

w^{1118}, p = 0.151, Mann-Whitney U-test) and *Dop1R2* mutant flies (*Dop1R2* vs. *y^1 w^{67c23}*, p = 0.1903, Mann-

Whitney U-test).

Reaction to threatening stimuli depends on fly walking speed

It has been shown that fruit flies may exhibit a different choice of escape behavior based on their

momentary speed when encountering the threat (Zacarias et al., 2018). Therefore, we tested whether

mutations in genes relevant to PD caused a change in this speed - behavioral response relationship. We

calculated the probability of stopping, slowing, speeding up and no reaction as a function of speed for all

mutants, as well as the ratio of each response type conditioned on walking speed at the time of shadow

presentations (Fig. 4a-b). These analyses confirmed that 'running' and 'no reaction' was most likely at

slow (< 5 mm/s) or zero walking speeds, while slowing down was more frequent as the walking speed

increased. Freezing was most frequently observed in the 5-13 mm/s speed range. We did not observe

significant difference between groups in their speed - behavioral response relationship for any of the

response types (stops, p = 0.4226; slow down, p = 0.6025; speed up, p = 0.9074; no reaction p = 0.4692;

two-way ANOVA genotype × speed interaction).

Since the distribution of the expression of various escape behaviors depended on walking speed, which

was also different across the mutant lines tested (Fig. 3), we asked whether the difference in baseline

speed could explain the observed differences in behavioral reactions. Therefore, we performed

simulations where reaction probabilities were based on the baseline walking speed distribution of each

mutant line to test whether behavioral responses could be predicted based on speed alone (Fig. S3). In

most of the groups we found a significant difference between the predicted and the measured response

type distributions, suggesting a role for other behavioral differences among the mutant lines beyond the

baseline speed differences (Parkin, p = 0.0487; α-Syn, p < 0.000001; *Dop1R1*, p = 0.332; *Dop1R2*, p =

0.006157; *DopEcR*, p = 0.003901, chi-square test).

Changes in escape behavior from the first to the fourth week of life

The age of *Drosophilae* may have a significant influence on their responses to threatening stimuli. To test

this, we examined control and dopamine receptor mutant flies (*w^{1118}*, *y^1 w^{67c23}*, *Dop1R1*, *Dop1R2*, *DopEcR*)

in 5 age groups: from 1-day-old to 4-week-old (Fig. 5; Fig. S4). The animals were kept under conditions

that accelerated their aging (29°C and 70% humidity). Our data showed that the behavior of 1-day-old

flies was significantly different from all other age groups. One-day-old flies stopped significantly more

often, and the frequency of stopping gradually decreased with age in all groups (two-way ANOVA, age, p

= 1.87 x 10^{-20}, f = 26.11; genotype, p = 1.3 x 10^{-29}, f = 38.42; genotype × age, p = 3.18 x 10^{-6}, f = 3.54; 1-

day-old vs. 1-week-old, p = 1.33 x 10^{-5}; 1-week-old vs. 2-week-old, p = 0.017, 2-week-old vs. 3-week-old,

p = 0.63; 3-week-old vs. 4-week-old, p = 0.098, Tukey's test for post hoc analysis). In contrast, the

probability of slowing down showed a substantial increase from 1-day-old to 1-week-old animals, and then

remained unchanged until the 4th week (two-way ANOVA, age, p = 9.11 x 10^{-15}, f = 18.72; genotype, p =

1.4 x 10^{-21}, f = 27.59; genotype × age, p = 0.0028, f = 2.291; 1-day-old vs. 1-week-old, 1-week-old vs. 2-

week-old, p = 2.56 x 10^{-8}; p = 1; 2-week-old vs. 3-week-old, p = 0.13; 3-week-old vs. 4-week-old, p = 0.093;

Tukey's test for post hoc analysis). The probability of speeding up showed a similar trend, except that the

4-week-old animals showed a significant decrease compared to 3-week-old flies (two-way ANOVA, age, p

= 2.48 x 10^{-15}, f = 19.44; genotype, p = 2.28 x 10^{-44}, f = 59.48; genotype × age, p = 1.1 x 10^{-7}, f = 4.12; 1-day-

old vs 1-week-old, p = 9.92 × 10^{-9}; 1-week-old vs. 2-week-old, p = 0.79; 2-week-old vs. 3-week-old, p =

0.75; 3-week-old vs. 4-week-old, p = 0.024; Tukey's test for post hoc analysis). Thus, different types of

escape reactions showed similar trends across the fly lifespan, while dopamine receptor mutant flies

maintained their decreased general reactivity throughout all age groups.

Discussion

We tested *Drosophila* mutant strains relevant for Parkinson's disease in a single-trial single-animal

behavioral assay. Our tests revealed strain-specific behavioral alterations in flies' reactions to predator-

mimicking passing shadows, serving as proof of principle demonstration of the viability of single-trial

approaches in *Drosophila*, a method widely used in mammalian studies. Specifically, we found reduced

walking speed, decreased freezing frequency and decreased overall reactivity in Parkin flies. In contrast,

α-Syn flies merely showed an increased freezing duration without a concomitant change in freezing

frequency, suggesting that Parkin flies better recapitulate some behavioral features of human PD

progression. *Dop1R* mutant flies resembled Parkin flies in their decreased walking speed and decreased

reactivity to the threatening stimuli. The distribution of the behavioral response fruit flies chose to

execute depended on their speed at the time of stimuli, and this speed - behavioral response relationship

was robust across the tested genotypes. Nevertheless, differences in walking speed alone did not explain

the strain-specific behavioral alterations. Age-dependence of the behavioral choice was also found to be

stereotypic across the tested genotypes.

Recent studies increasingly recognize the importance of investigating subtle changes in fly behavior to

better understand the manifestations of locomotor and other disorders (Geissmann et al., 2017; Zacarias

et al., 2018; Seidenbecher et al., 2020). Along these lines, Aggarwal and colleagues introduced an

automated climbing assay for fruit flies and revealed subtle motor deficits in different mutant strains

(Aggarwal et al., 2019). We expanded the scope of these studies by examining a diverse behavioral

repertoire of *Drosophila* in response to threatening stimuli. The diversity of freeze vs. flee responses of

fruit flies, including different durations of stopping as well as running, were accurately described by

Zacarias et al. (Zacarias et al., 2018), where they also identified DNp09 neurons as important controllers

of running and freezing responses. We built on this behavioral characterization for our definitions of fly

reactions to predator-mimicking stimuli.

Mutations in the *PARK2* gene lead to impaired ubiquitination and a consequential mitochondrial

dysfunction (Guo, 2010, 2012), and early-onset PD in human patients. This led to the widespread use of

Parkin flies as a genetic model pf PD (Guo, 2010; Chambers et al., 2013; Aggarwal et al., 2019). Of note,

flies that lack Parkin display a significant degeneration of dopaminergic neurons (Whitworth et al., 2005).

We found similarities between Parkin and *Dop1R* mutant flies in their altered responses to predator-

mimicking passing shadows: reduced walking speed and decreased overall reactivity, suggesting that the

lack of dopamine action through *Dop1R* may be one of the common pathways underlying motor deficits.

This is in line with a recent study demonstrating impaired startle-induced geotaxis and locomotor

reactivity in *Dop1R* mutant flies (Sun et al., 2018), similar to what had been shown for Parkin flies

(Aggarwal et al., 2019), and a demonstration of dopaminergic control over walking speed (Marquis and

Wilson, 2022). In contrast, flies expressing human α-syn showed moderate changes in motor behavior

except for a marked prolongation of freezing duration upon threating stimuli. This is in accordance with

studies suggesting that α-syn misexpression is not fully penetrant under some conditions (Pesah et al.,

2005; Nagoshi, 2018), and calls for the use of other genetic *Drosophila* models, including Parkin flies, in

studying the pathophysics of human PD.

The dopamine/ecdysteroid receptor *DopEcR* is a G-protein-coupled dual receptor for dopamine and the

steroid hormone ecdysone. It has been proposed that the *DopEcR* may serve as an integrative hub for

dopamine-mediated actions and stress responses in fruit flies (Petruccelli et al., 2020). We found that

DopEcR mutant flies showed decreased mean walking speed, decreased probability of slowing down and

decreased overall behavioral reactivity in response to predator-mimicking passing shadows. This pattern of alterations was largely similar to those observed in Parkin and *Dop1R* mutant flies, suggesting that the *DopEcR* may convey similar dopamine-mediated functions as *DopR1* at least in those motor aspects tested in the present study. A recent work demonstrated that serotonin also modulates both walking speed and startle response in flies, suggesting a complex neuromodulatory control over *Drosophila* motor behavior (Howard et al., 2019).

In humans, degeneration of midbrain dopaminergic neurons has been found as a direct cause of PD symptoms and DA supplementation with its precursor levodopa has been one of the most successful therapeutic approaches to date. The fruit fly and mammalian DA systems show a number of homologies which are thought to serve as a good basis for *Drosophila* PD models (Feany and Bender, 2000; Hewitt and Whitworth, 2017). Specifically, a deep-running evolutional conservation has been revealed between the arthropod central complex and the vertebrate basal ganglia, where GABAergic and dopaminergic neurons play key roles in motor control in both phylae (Strausfeld and Hirth, 2013). Clusters of fly dopaminergic neurons that project to the ellipsoid body, the fan-shaped body and the lateral accessory lobes are thought to share homologies with the striatum-projecting dopaminergic neurons of the substantia nigra in mammals. Both the fruit fly central complex and vertebrate basal ganglia mediate a range of functions from motor control and sensorimotor integration to action selection and decision making to motivation (Strauss, 2002; Claridge-Chang et al., 2009; Kong et al., 2010) and learning (McCurdy et al., 2021; Zolin et al., 2021; Fisher et al., 2022; Qiao et al., 2022; Taisz and Jefferis, 2022; Yamada et al., 2023). It has been found that fruit flies lacking the DA-synthetizing tyrosine hydroxylase enzyme in the central nervous system show reduced activity and locomotor deficit that worsen with age, and also exhibit marked impairments in associative learning based on both appetitive and aversive reinforcement (Riemensperger et al., 2011). This is in line with mammalian studies emphasizing the role of the dopaminergic nigrostriatal pathway in controlling goal-directed action (Eban-Rothschild et al., 2016), reinforcement learning (Schultz et al., 1997; Lak et al., 2014), and more recently, conveying aversive information (Menegas et al., 2015, 2018). Our findings are consistent with the apparent homologies across fruit fly and mammalian dopaminergic systems and suggest that a more in-depth investigation of specific behavioral changes and underlying dopamine-related dysfunctions may yield translatable results. We also observed age-related changes in motor behavior in response to threatening stimuli, particularly in 4-week-old flies, which may parallel the age-dependent worsening of PD symptoms in humans.

Acknowledgements

We thank the Anne Von Phillipsborn lab, DANDRITE, Aarhus University for the kind gift of the DA receptor mutant lines. Most of the *Drosophila* stocks were obtained from the Bloomington Drosophila Stock Center (NIH P40OD018537). We thank the FENS-Kavli Network of Excellence for fruitful discussions. B.H. was supported by the Eotvos Lorand Research Network. T.V. was supported by the grants OTKA (Hungarian Scientific Research Fund) K132439 and VEKOP (VEKOP-2.3.2-16-2017-00014), and by the ELKH/MTA-ELTE Genetics Research Group (01062).

Data and code availability

All analysis functions are available at https://github.com/hangyabalazs/Drosophila-behavior-analysis. All data are available at https://figshare.com/s/40539ee040a269bf05e5.

Methods

Animals

We used 2-weeks-old *Drosophila melanogaster* (both males and females) raised at 24°C and 60% humidity in a natural light-dark cycle for experiments presented in Figures 1-4. For age-group comparisons (Fig.5.), fruit flies aged 1, 7, 14, 21 and 28 days were used. These flies were raised at 29°C and 70% humidity to accelerate aging.

Two sets of mutant groups were used for behavioral comparison. In the first set of animals we used *w*; UAS-Parkin-R275W* (created in the laboratory of Kah-Leong Lim, Neurodegeneration Research Laboratory, National Neuroscience Institute, Singapore)(Wang et al., 2007; Kovács et al., 2017) and *w*; UAS-alpha-Syn-A53T* (BDSC 8148) from the Eotvos Lorand University, with y^1, v^1; *UAS-GFP* (BDSC 35786) and *+; P{Ddc-GAL4.L}4.36* (modified version of BDSC 7009, the original first chromosome w^{1118} has been replaced to w^+ in this study) as controls. For the second set of animals we used *Dop1R1*, *Dop1R2* and *DopEcR* mutant flies kindly donated by the Anne Von Phillipsborn lab, DANDRITE, Aarhus University and y^1 *w*; Mi{MIC}Dop1R1*^{MI04437} (BDSC 43773), y^1 *w*; Mi{MIC}Dop1R2*^{MI08664} (BDSC 51098), and w^{1118}; *PBac{PB}DopEcR*^{c02142}/*TM6B, Tb*^{1} (BDSC 10847) with w^{1118} (BDSC 5905) y^1 and y^1 w^{67c23} (BDSC 6599) as controls from the Bloomington stock centre. Table 1. shows the number of recorded flies for the age group comparisons, and Table 2. shows the number of recorded flies for comparing the mutant groups.

	Dop1R1	*Dop1R2*	*DopEcR*	w^{1118}	$y^1\ w^{67c23}$
1-day-old	35	36	36	35	41
1-week-old	36	54	36	36	36
2-week-old	36	36	36	36	30
3-week-old	35	35	29	36	36
4-week-old	30	36	34	30	30

328   Table 1. Number of recorded flies for the age group comparisons.

2-week-old	No. of flies
Dop1R1	45
Dop1R2	47
DopEcR	35
w^{1118}	46
$y^1,\ w^{67c23}$	40
Parkin	78
α-Syn	69
iso w^{1118}	92
+GFP	65

Table 2. Number of recorded flies for comparing the mutant groups.

Arena

We designed an arena made of four layers of plexiglass and metal that formed 13 tunnels (Fig. 1a). Each

tunnel was 52mm x 5mm × 1mm, in which individual flies were able to move freely in two dimensions but

could not fly. The bottom layer was a standard transparent plexiglass layer. The second layer contained

the tunnel walls made of metal to prevent horizontal spread of light. The third layer was the top of the

tunnels, made of transparent plexiglass, which could slide above the tunnel. It also contained a small hole

on the side, where the flies could be inserted into the tunnel through a pipette. The fourth layer was a

metal cover designed with cut-outs that matched the shape and position of the tunnels to prevent flies in

the neighboring tunnels from detecting the shadow before it reached their position.

Movement tracking

Flies were tracked by using a FLIR Camera (FLIR Blackfly S USB3 FLIR Systems, Wilsonville, OR, US) placed

under the arena (Fig. 1a). The frame rate was set to 100 frames/second to allow the detection of position

with high temporal resolution. The recorded images were processed by Bonsai (2.5.2) tracking software

(Lopes et al., 2015), using customized Bonsai code. The code extracts each tunnel areas separately from

the image and detects flies by selecting the biggest and darkest pixel object in the tunnel area in real time.

Fly position coordinates were calculated from the centroid. The software stored x-y position coordinates

along with their timestamps in csv files, which were later processed in Matlab R2016a (Mathworks, Natick,

MA, US).

Predator-mimicking shadow stimuli

A 10.1-inch screen (HannStar HSD101PWW1-A00,) was placed on the top of the arena. The screen

presented a yellow background. We implemented 'shadow' stimuli in red color, as these stimuli were

suggested to be perceived by fruit flies as dark shadows and also enabled continuous tracking of the

animals (Sharkey et al., 2020). The red color screen represented a large enough contrast change to evoke

escape behaviors of flies, but also provided sufficient brightness to enable continuous motion tracking. A

passing shadow stimulus was chosen, as it was affecting all tunnels simultaneously. To mimic an

approaching shadow, a red screen slid in from the side, stayed on for 2 seconds, then slid out. The shadow

stimuli were separated by pseudorandomized inter-trial intervals with an approximately exponential

distribution with a mean of 52.2 seconds, preventing flies from anticipating the next shadow stimulus.

Identification of behavioral response types to the threatening stimuli

Animal speed was calculated as a function of time in a -200 ms to 1000 ms window around each shadow

stimulus in 10 ms time bins. These speed functions were normalized by subtracting the average speed in

the 200 ms window before the shadow presentation to reveal stimulus induced absolute instantaneous

speed changes. Principal component analysis (PCA) was used to reduce dimensionality of the normalized

speed functions in the 1 s interval following shadow presentation in a way that the variance between trials

is maximally preserved in the low-dimensional representation. Agglomerative hierarchical clustering was

performed in the space spanned by the first three principal components to identify trials with distinct

characteristic response types.

Behavior detection

After identifying the four most frequently observed animal responses to shadow presentations, we

algorithmically defined them based on the speed and acceleration thresholds. First, we examined the

speed of the animal in the 200 ms time window before the shadow (v_{pre}); stopping/freezing or slowing

down was only possible if the fly had been moving, defined by $v_{pre} > 0$ mm/s. Second, we examined the

acceleration (a_{post}) in the 1 s interval following the shadow using a 100 ms moving average window to

characterize the flies' response. Reactions were considered robust if the absolute value of the acceleration

reached 200 mm/s^2. A trial was characterized as a 'stop' trial, if the average v_{pre} was above 0 mm/s, the

fly's first reaction was deceleration, and 0 mm/s speed was reached before accelerating again. If the first

reaction was deceleration but 0 mm/s speed was not reached, the trial was classified as a 'slow down'
trial. In 'speed up' trials, the first response following the shadow was acceleration. If the absolute value
of a_{post} did not reach the 200 mm/s^2 threshold during the 1 s interval following the shadow, the trial was
labeled as a 'no reaction' trial.

Data analysis and statistics

Data analysis was performed in Matlab R2016a (Mathworks, Natick, MA, US) using custom-written code.
Statistical comparison between groups of flies was performed by two-sided Mann-Whitney U-test. Exact
p values are reported in the Results section. The effect of movement speed on the distribution of
behavioral response types was tested using Monte Carlo simulation (Fig.S3). First, we calculated the
probability of each response type at different speed values from the grand average of all trails of every
animal. Then we simulated behavior of virtual flies by drawing random velocity values from the velocity
distribution of each genotype and then randomly selecting a reaction based on the reaction probabilities
associated with the drawn velocity. Finally, we calculated reaction probabilities for the virtual flies and
compared it with real data from animals of the same genotype. Differences were statistically tested by
Chi-squared test.

Figure Legends

Fig. 1. Behavioral apparatus and experimental design.

a, Top left, schematic of the experimental setup. Top right, schematic of the fly arena. Bottom left,
photograph of the experimental setup. Bottom right, a photograph of the fly arena. **b,** Timeline of a
session showing the shadow presentations in red over the yellow background.

Fig. 2. Behavioral characterization of the escape behavior repertoire of individual fruit flies.

a, Schematic for calculating speed and acceleration based on tracked position coordinates. **b,** Four
characteristic escape behaviors categorized by PCA for an example session of a control fly (w1118; from
left; stop, slow down, speed up, no reaction). Top, color-coded heatmaps indicating the walking speed of
the fly (blue, low speed; yellow, high speed), aligned to shadow presentations (orange line). Bottom,
average moving speed triggered on the shadow presentations (orange line). **c,** Threshold-based
classification of behavioral responses (from left, stop, slow down, speed up, no reaction). Top, single-trial
example for each response type. Time intervals for calculating the average speed before the shadow
presentation (orange) as well as the speed and acceleration after the shadow presentation (green) are

marked. Threshold values for each response type are displayed above the graphs. Bottom, average walking speed across the trials from all session of w^{1118} flies, sorted by the type of behavioral response. Line and errorshade show mean ± SE.

Fig. 3. Parkinson's model and dopamine receptor mutant fruit flies showed reduced walking speed and decreased reactivity to threatening stimuli.

a, Distribution of the mean speed measured in the time window [-0.2, 0] seconds relative to shadow presentation for the different mutant groups. Top, Parkin flies showed reduced mean speed compared to controls (Parkin vs. *iso w^{1118}*; p = 6.55 × 10^{-6}, Mann–Whitney U-test). Bottom, *Dop1R1* and *DopEcR* mutant flies showed reduced mean speed compared to controls (*$y^1 w^{67c23}$* and *w^{1118}* respectively; p = 0.0016, p = 0.0034; Mann-Whitney U-test). **b,** Distribution of the proportion of stop trials in different mutant groups. Parkin flies showed a reduced tendency for stopping compared to *iso w^{1118}* (p = 0.0043, Mann Whitney U test). **c,** Distribution of stop duration in different mutant groups. α-Syn flies showed increased stop durations compared to *iso w^{1118}* (p = 2.18 × 10^{-11}, Mann-Whitney U-test). **d,** Distribution of the proportion of speed up trials in different mutant groups. Top, α-Syn flies showed a reduced tendency to speed up compared to their controls (p = 8.4 × 10^{-6}, Mann-Whitney U-test). Bottom, *Dop1R1* and *Dop1R2* mutant flies also showed a significantly reduced tendency to speed up compared to their controls (p = 0.0122 and p = 0.0024, respectively; Mann-Whitney U-test). **e,** Distribution of the proportion of slow down trials in different mutant groups. Bottom, *DopEcR* showed a 15.04% decrease compared to w^{1118} (p = 0.020, Mann-Whitney U-test). **f,** Distribution of the proportion of 'no reaction' trials in different mutant groups. Top, Parkin mutants showed reduced reactivity compared to *iso w^{1118}* controls (p = 0.0043, Mann-Whitney U-test). Bottom, *Dop1R1* and *DopEcR* mutants also showed reduced reactivity compared to *$y^1 w^{67c23}$* and *w^{1118}* controls, respectively (p = 0.0173 and p = 0.0167, respectively; Mann-Whitney U-test). Box-whisker plots show median, interquartile range and non-outlier range. *, p < 0.05; **, p < 0.01; ***, p < 0.001. Exact genotypes: *iso w^{1118}*: *+; +; Ddc-Gal4/+*. Parkin: *+; +; Ddc-Gal4/UAS-Parkin-R275W*. α-Syn: *+; +; Ddc-Gal4/UAS-α-Syn-A53T*. *$y^1 w^{67c23}$*: *$y^1 w^{67c23}$*. *Dop1R*: *$y^1 w^*$; Mi{MIC}Dop1R1^{MI04437}*. *Dop1R2*: *$y^1 w^*$; Mi{MIC}Dop1R2^{MI08664}*. *DopEcR*: *w^{1118}; PBac{PB}DopEcR^{c02142}/TM6B, Tb^1*.

Fig. 4. Reaction to threatening stimuli depends on fly walking speed.

a, Probability of a given response type as a function of average fly speed before the shadow presentation (200 ms pre-stimulus time window). From left, stop, slow down, speed up and no reaction trials are quantified. Top, Parkin and α-Syn flies. Bottom, *Dop1R1*, *Dop1R2* and *DopEcR* mutant flies. **b,** Proportion

of a given response type as a function of average fly speed before the shadow presentation. Top, Parkin and α-Syn flies. Bottom, *Dop1R1*, *Dop1R2* and *DopEcR* mutant flies. Exact genotypes: *iso w^{1118}: +; +; Ddc-Gal4/+.* Parkin: *+; +; Ddc-Gal4/UAS-Parkin-R275W.* α-Syn: *+; +; Ddc-Gal4/UAS-α-Syn-A53T. y^1 w^{67c23}: y^1 w^{67c23}. Dop1R: y^1 w*; Mi{MIC}Dop1R1^{MI04437}. Dop1R2: y^1 w*; Mi{MIC}Dop1R2^{MI08664}. DopEcR: w^{1118}; PBac{PB}DopEcR^{c02142}/TM6B, Tb^1.*

Fig. 5. Changes in escape behavior from the first to the fourth week of life

Proportion of responses as a function of age for different groups of mutants and controls. From left to right, stop, slow down, speed up and no reaction trials are quantified. Lines and errorshades show mean and standard error. Exact genotypes: *iso w^{1118}: +; +; Ddc-Gal4/+.* Parkin: *+; +; Ddc-Gal4/UAS-Parkin-R275W.* α-Syn: *+; +; Ddc-Gal4/UAS-α-Syn-A53T. y^1 w^{67c23}: y^1 w^{67c23}. Dop1R: y^1 w*; Mi{MIC}Dop1R1^{MI04437}. Dop1R2: y^1 w*; Mi{MIC}Dop1R2^{MI08664}. DopEcR: w^{1118}; PBac{PB}DopEcR^{c02142}/TM6B, Tb^1.*

Supplementary Figure Legends

Fig. S1. Picture of the arena from the camera view

The 13 tunnels of the arena from the camera view with the tracking lines showing that the animals can move in 2 dimnesions. The tracking line color indicates time, with dark blue and red marking the beginning and the end of the tracking, respectively.

Fig. S2. Parkin and α-Syn fly group escape reaction compared to GFP mutant flies

a, Distribution of the mean speed measured in the 0.2 second before shadow presentation for Parkin, α-Syn and +GFP mutant groups. Parkin and α-Syn flies showed reduced mean speed compared to +GFP controls (Parkin vs. +GFP, p = 1.58 x 10^{-11}; α-Syn vs. +GFP, p = 2.85 x 10^{-4}; Mann–Whitney U-test). **b,** Distribution of the proportion of stop trials in different mutant groups. Parkin flies showed reduced tendency for stopping compared to the +GFP group (Parkin vs. +GFP, p = 7.1 x 10^{-5}; α-Syn vs. +GFP, p = 0.67; Mann–Whitney U-test). **c,** Distribution of stop duration in different mutant groups. α-Syn flies showed increased stop durations compared to GFP (Parkin vs. +GFP, p = 0.17; α-Syn vs. +GFP, p = 3.98 x 10^{-10}; Mann–Whitney U-test). **d,** Distribution of the proportion of speed up trials in different mutant groups. Both Parkin and α-Syn flies showed a reduced tendency to speed up compared to +GFP controls (Parkin vs. +GFP, p = 8.3 x 10^{-5}; α-Syn vs. +GFP, p = 1.96 x 10^{-12}; Mann–Whitney U-test). **e**, Distribution of the proportion of slow down trials in different mutant groups. Parkin flies showed a reduced tendency to slow down compared to +GFP controls (Parkin vs. +GFP, p = 0.044; α-Syn vs. +GFP, p = 0.465; Mann–Whitney U-test). **f**, Distribution of the proportion of 'no reaction' trials in different mutant groups. Both

Parkin and α-Syn flies showed reduced reactivity compared to +GFP controls (Parkin vs. +GFP, p = 9.4 x 10^{-13}; α-Syn vs. +GFP, p = 1,3 x 10^{-8}; Mann-Whitney U-test). *, p < 0.05; **, p < 0.01; ***, p < 0.001. Exact genotypes: *iso w^{1118}: +; +; Ddc-Gal4/+*. Parkin: *+; +; Ddc-Gal4/UAS-Parkin-R275W*. α-Syn: *+; +; Ddc-Gal4/UAS-α-Syn-A53T*.

Fig. S3. Differences in walking speed do not explain different escape behaviors

Real and simulated proportion of response types are shown for the mutant groups tested. For the simulations, random velocity values were drawn from the velocity distribution of each genotype and then a response type was randomly selected based on the response type distributions associated with the drawn velocity. Bar plots show the proportion of response types color-coded. For each mutant group, the right bar shows the measured proportion of escape responses, while the left bar shows the simulated distribution. We found significant differences between the real and simulated distributions for the following groups: Parkin (p = 0.0487), α-Syn (p < 0.000001), *Dop1R2* (p = 0.006157) and *DopEcR* (p = 0.003901). We found no significant differences for the *Dop1R1* group (p = 0.332; Chi-square test for all the statistics). *, p < 0.05; **, p < 0.01; ***, p < 0.001. Exact genotypes: *iso w^{1118}: +; +; Ddc-Gal4/+.* Parkin: *+; +; Ddc-Gal4/UAS-Parkin-R275W*. α-Syn: *+; +; Ddc-Gal4/UAS-α-Syn-A53T*. *y^1 w^{67c23}: y^1 w^{67c23}*. *Dop1R: y^1 w*; Mi{MIC}Dop1R1^{MI04437}*. *Dop1R2: y^1 w*; Mi{MIC}Dop1R2^{MI08664}*. *DopEcR: w^{1118}; PBac{PB}DopEcR^{c02142}/TM6B, Tb^1*.

Fig. S4. Changes in escape behavior from the first to the fourth week of life

a, Average walking speed as a function of age for different groups of controls (left) and mutants (right). Lines and errorshades show mean and standard error. Mean speed showed significant differenceis among age groups (two-way ANOVA, age, f = 41.38, p = 9.39 x 10^{-32}; genotype, f = 42.19, p = 2.46 x 10^{-32}; genotype × age, f = 4.36, p = 2.667 x 10^{-8};Tukey's post hoc test: 1 day old vs. 1 week old, p = 9.92 x 10^{-9}; 1 week old vs. 2 weeks old, p = 7.34 x 10^{-7}; 2 weeks old vs. 3 weeks old, p = 0.085; 3 weeks old vs. 4 weeks old, p = 0.0043). **b,** Stop duration as a function of age for different groups of controls (left) and mutants (right). We found that age did not effect the duration of stops (two-way ANOVA, age, f = 0.85, p = 0.49; genotype, f = 0.91, p = 0.46; genotype × age, f = 2.53, p = 0.0008; Tukey's post hoc test: 1 day old vs. 1 week old, p = 0.98; 1 week old vs. 2 weeks old, p = 0.992; 2 weeks old vs. 3 weeks old, p = 0.578; 3 weeks old vs. 4 weeks old, p = 0.9759). Lines and errorshades show mean and standard error.

References

Aggarwal A, Reichert H, VijayRaghavan K (2019) A locomotor assay reveals deficits in heterozygous
Parkinson's disease model and proprioceptive mutants in adult Drosophila. Proc Natl Acad Sci U S A
116:24830–24839.

Bloem BR, Okun MS, Klein C (2021) Parkinson's disease. Lancet 397:2284–2303.

Bové J, Perier C (2012) Neurotoxin-based models of Parkinson's disease. Neuroscience 211:51–76.

Breger LS, Fuzzati Armentero MT (2019) Genetically engineered animal models of Parkinson's disease:
From worm to rodent. Eur J Neurosci 49:533–560.

Cackovic J, Gutierrez-Luke S, Call GB, Juba A, O'Brien S, Jun CH, Buhlman LM (2018) Vulnerable parkin
loss-of-function Drosophila dopaminergic neurons have advanced mitochondrial aging,
mitochondrial network loss and transiently reduced autophagosome recruitment. Front Cell
Neurosci 12:1–14.

Cannon JR, Greenamyre JT (2010) Neurotoxic in vivo models of Parkinson's disease. In: Progress in Brain
Research, pp 17–33. Elsevier B.V.

Chambers RP, Call GB, Meyer D, Smith J, Techau JA, Pearman K, Buhlman LM (2013) Nicotine increases
lifespan and rescues olfactory and motor deficits in a Drosophila model of Parkinson's disease.
Behav Brain Res 253:95–102.

Claridge-Chang A, Roorda RD, Vrontou E, Sjulson L, Li H, Hirsh J, Miesenböck G (2009) Writing Memories
with Light-Addressable Reinforcement Circuitry. Cell 139:405–415.

Duffy JB (2002) GAL4 system indrosophila: A fly geneticist's swiss army knife. genesis 34:1–15.

Eban-Rothschild A, Rothschild G, Giardino WJ, Jones JR, De Lecea L (2016) VTA dopaminergic neurons
regulate ethologically relevant sleep-wake behaviors. Nat Neurosci 19:1356–1366.

Feany MB, Bender WW (2000) A Drosophila model of Parkinson's disease. Nature 404:394–398.

Fisher YE, Marquis M, D'Alessandro I, Wilson RI (2022) Dopamine promotes head direction plasticity
during orienting movements. Nature 612:316–322.

Geissmann Q, Garcia Rodriguez L, Beckwith EJ, French AS, Jamasb AR, Gilestro GF (2017) Ethoscopes: An
open platform for high-throughput ethomics. PLOS Biol 15:e2003026.

Guo M (2010) What have we learned from Drosophila models of Parkinson's disease? In: Progress in

Brain Research, pp 2–16. Elsevier B.V.

Guo M (2012) Drosophila as a model to study mitochondrial dysfunction in Parkinson's disease. Cold Spring Harb Perspect Med 2:1–18.

Haywood AFM, Staveley BE (2006) Mutant α-synuclein-induced degeneration is reduced by parkin in a fly model of Parkinson's disease. Genome 49:505–510.

Hewitt VL, Whitworth AJ (2017) Mechanisms of Parkinson's Disease. In: Current Topics in Developmental Biology, 1st ed., pp 173–200. Elsevier Inc.

Howard CE, Chen C-L, Tabachnik T, Hormigo R, Ramdya P, Mann RS (2019) Serotonergic Modulation of Walking in Drosophila. Curr Biol 29:4218-4230.e8.

Julienne H, Buhl E, Leslie DS, Hodge JJL (2017) Drosophila PINK1 and parkin loss-of-function mutants display a range of non-motor Parkinson's disease phenotypes. Neurobiol Dis 104:15–23.

Kahle PJ, Neumann M, Ozmen L, Müller V, Jacobsen H, Schindzielorz A, Okochi M, Leimer U, Van Der Putten H, Probst A, Kremmer E, Kretzschmar HA, Haass C (2000) Subcellular localization of wild-type and Parkinson's disease-associated mutant α-synuclein in human and transgenic mouse brain. J Neurosci 20:6365–6373.

Király B, Hangya B (2022) Navigating the Statistical Minefield of Model Selection and Clustering in Neuroscience. eNeuro 9.

Kong EC, Woo K, Li H, Lebestky T, Mayer N, Sniffen MR, Heberlein U, Bainton RJ, Hirsh J, Wolf FW (2010) A pair of dopamine neurons target the D1-like dopamine receptor dopr in the central complex to promote ethanol-stimulated locomotion in drosophila. PLoS One 5.

Kottler B, Faville R, Bridi JC, Hirth F (2019) Inverse Control of Turning Behavior by Dopamine D1 Receptor Signaling in Columnar and Ring Neurons of the Central Complex in Drosophila. Curr Biol 29:567-577.e6.

Kovács T et al. (2017) The small molecule AUTEN-99 (autophagy enhancer-99) prevents the progression of neurodegenerative symptoms. Sci Rep 7:42014.

Lak A, Stauffer WR, Schultz W (2014) Dopamine prediction error responses integrate subjective value from different reward dimensions. Proc Natl Acad Sci U S A 111:2343–2348.

Lopes G, Bonacchi N, Frazão J, Neto JP, Atallah B V., Soares S, Moreira L, Matias S, Itskov PM, Correia PA,

Medina RE, Calcaterra L, Dreosti E, Paton JJ, Kampff AR (2015) Bonsai: an event-based framework

for processing and controlling data streams. Front Neuroinform 9:7.

Marquis M, Wilson RI (2022) Locomotor and olfactory responses in dopamine neurons of the Drosophila

superior-lateral brain. Curr Biol 32:5406-5414.e5.

McCurdy LY, Sareen P, Davoudian PA, Nitabach MN (2021) Dopaminergic mechanism underlying reward-

encoding of punishment omission during reversal learning in Drosophila. Nat Commun 12:1115.

Menegas W, Akiti K, Amo R, Uchida N, Watabe-Uchida M (2018) Dopamine neurons projecting to the

posterior striatum reinforce avoidance of threatening stimuli. Nat Neurosci 21:1421–1430.

Menegas W, Bergan JF, Ogawa SK, Isogai Y, Venkataraju KU, Osten P, Uchida N, Watabe-Uchida M

(2015) Dopamine neurons projecting to the posterior striatum form an anatomically distinct

subclass. Elife 4:1–30.

Nagoshi E (2018) Drosophila Models of Sporadic Parkinson's Disease. Int J Mol Sci 19:3343.

Obeso JA et al. (2017) Past, present, and future of Parkinson's disease: A special essay on the 200th

Anniversary of the Shaking Palsy. Mov Disord 32:1264–1310.

Olesen J, Gustavsson a., Svensson M, Wittchen HU, Jönsson B (2012) The economic cost of brain

disorders in Europe. Eur J Neurol 19:155–162.

Perez RG, Waymire JC, Lin E, Liu JJ, Guo F, Zigmond MJ (2002) A Role for α-Synuclein in the Regulation of

Dopamine Biosynthesis. J Neurosci 22:3090–3099.

Pesah Y, Burgess H, Middlebrooks B, Ronningen K, Prosser J, Tirunagaru V, Zysk J, Mardon G (2005)

Whole-mount analysis reveals normal numbers of dopaminergic neurons following misexpression

of?-Synuclein inDrosophila. genesis 41:154–159.

Petruccelli E, Lark A, Mrkvicka JA, Kitamoto T (2020) Significance of DopEcR, a G-protein coupled

dopamine/ecdysteroid receptor, in physiological and behavioral response to stressors. J

Neurogenet 34:55–68.

Polymeropoulos MH et al. (1997) Mutation in the α-synuclein gene identified in families with Parkinson's

disease. Science (80-) 276:2045–2047.

Przedborski S (2017) The two-century journey of Parkinson disease research. Nat Rev Neurosci 18:251–259.

Qiao J, Yang S, Geng H, Yung W-H, Ke Y (2022) Input-timing-dependent plasticity at incoming synapses of the mushroom body facilitates olfactory learning in Drosophila. Curr Biol 32:4869-4880.e4.

Riemensperger T, Isabel G, Coulom H, Neuser K, Seugnet L, Kume K, Iché-Torres M, Cassar M, Strauss R, Preat T, Hirsh J, Birman S (2011) Behavioral consequences of dopamine deficiency in the Drosophila central nervous system. Proc Natl Acad Sci U S A 108:834–839.

Schapira AHV (2009) Neurobiology and treatment of Parkinson's disease. Trends Pharmacol Sci 30:41–47.

Schapira AHV, Chaudhuri KR, Jenner P (2017) Non-motor features of Parkinson disease. Nat Rev Neurosci 18:435–450.

Schultz W, Dayan P, Montague PR (1997) A neural substrate of prediction and reward. Science 275:1593–1599.

Seidenbecher SE, Sanders JI, von Philipsborn AC, Kvitsiani D (2020) Reward foraging task and model-based analysis reveal how fruit flies learn value of available options Skoulakis EMC, ed. PLoS One 15:e0239616.

Sharkey CR, Blanco J, Leibowitz MM, Pinto-Benito D, Wardill TJ (2020) The spectral sensitivity of Drosophila photoreceptors. Sci Rep.

Spillantini MG, Schmidt ML, Lee VM-Y, Trojanowski JQ, Jakes R, Goedert M (1997) α-Synuclein in Lewy bodies. Nature 388:839–840.

Strausfeld NJ, Hirth F (2013) Deep homology of arthropod central complex and vertebrate basal ganglia. Science (80-) 340:157–161.

Strauss R (2002) The central complex and the genetic dissection of locomotor behaviour. Curr Opin Neurobiol 12:633–638.

Sun J, Xu AQ, Giraud J, Poppinga H, Riemensperger T, Fiala A, Birman S (2018) Neural control of startle-induced locomotion by the mushroom bodies and associated neurons in drosophila. Front Syst Neurosci 12:1–18.

Taisz I, Jefferis GSXE (2022) Speed of learning depends on turning. Nature 612:216–217.

Wang C, Lu R, Ouyang X, Ho MWL, Chia W, Yu F, Lim K-L (2007) Drosophila Overexpressing Parkin R275W
Mutant Exhibits Dopaminergic Neuron Degeneration and Mitochondrial Abnormalities. J Neurosci
27:8563–8570.

Whitworth AJ, Theodore DA, Greene JC, Beneš H, Wes PD, Pallanck LJ (2005) Increased glutathione S -
transferase activity rescues dopaminergic neuron loss in a Drosophila model of Parkinson's disease.
Proc Natl Acad Sci 102:8024–8029.

Yamada D, Bushey D, Li F, Hibbard KL, Sammons M, Funke J, Litwin-Kumar A, Hige T, Aso Y (2023)
Hierarchical architecture of dopaminergic circuits enables second-order conditioning in Drosophila.
Elife 12:1–30.

Zacarias R, Namiki S, Card GM, Vasconcelos ML, Moita MA (2018) Speed dependent descending control
of freezing behavior in Drosophila melanogaster. Nat Commun 9:1–11.

Zolin A, Cohn R, Pang R, Siliciano AF, Fairhall AL, Ruta V (2021) Context-dependent representations of
movement in Drosophila dopaminergic reinforcement pathways. Nat Neurosci 24:1555–1566.

    Figures

    Figure 1

622 **Figure 2**

a

Tracking software

x, y

y

$$d = \sqrt{\Delta x^2 + \Delta y^2}$$

d Δy

Δx

x

$v = \frac{\Delta d}{\Delta t}$

v shadow

$$a = \frac{\Delta v}{\Delta t}$$

t

b

Stop	Slow down	Speed up	No reaction

Trial #

250   25   35   2500

v (mm/s): 25 20 15 10 5 0

v (mm/s) 30 / 20 / 10 / 0

-1 0 1   t (s) (all panels)

c

Stop   **Slow down**   **Speed up**   **No reaction**

$\overline{v}_{pre} > 0, \min(v_{post}) = 0$ $a_{post} < -200$ mm/s²

$\overline{v}_{pre} > 0, \min(v_{post}) > 0$ $a_{post} < -200$ mm/s²

$a_{post} > 200$ mm/s²

$|a_{post}| < 200$ mm/s²

v (mm/s) 20 / 10 / 0

v (mm/s) 10 / 5 / 0

-1 0 1   t (s) (all panels)

Figure 3

Figure 4

Figure 5

Supplementary Figures

635   **Figure S1**

     Figure S2

Figure S3

Figure S4