Cooperative progression of colitis and leukemia modulated by clonal hematopoiesis via PTX3/IL-1β pro-inflammatory signaling

Hang He[1,2], Yuchen Wen[1,2], Hanzhi Yu[1,2], Jingjing Liu[1,2], Qingran Huo[1,2], Wenyan Jin[1,2], Zhiqin Wang[1,2], Guohui Du[3], Jun Du[4], Huaquan Wang[5], Zhigang Zhao[6], Zhigang Cai[1,2,4,5,*]

[1]The Province and Ministry Co-sponsored Collaborative Innovation Center for Medical Epigenetics; Department of Pharmacology, School of Basic Medical Science, Tianjin Medical University, Tianjin, China.
[2]National Key Laboratory of Blood Science, Tianjin, China.
[3]Department of Research and Development, Beijing SeekGene BioSciences Co. Ltd, Beijing, China.
[4]Department of Rheumatology and Immunology, Tianjin Medical University Tianjin General Hospital, Tianjin, China.
[5]Department of Hematology, Tianjin Medical University Tianjin General Hospital, Tianjin, China.
[6]Department of Medical Oncology, Tianjin First Central Hospital, School of Medicine, Nankai University, Tianjin, China.

*Correspondence:
Zhigang Cai, Professor, Ph.D.
Tianjin Medical University
No.22, Qixiangtai Road, Tianjin, China 200037
E-mail: US36zcai@tmu.edu.cn
Phone: (+86)18622633722

Abstract

Clonal hematopoiesis (CH) is suggested as an important risk factor for all-cause mortality and multiple chronic diseases including hematological neoplasms, cardiovascular diseases, and potentially a spectrum of autoimmune or immune-deficiency diseases. Mutations in *TET2* are one of the first identified, most important and prevalent genetic drivers of CH. However, cooperative factors and mechanisms underlying *TET2*-deficiency related CH (TedCH) remain largely unknown. Controversially, it is recently suggested that certain diseases take place prior to TedCH and promote TedCH on the contrary, indicating diseases in non-hematopoietic organs may act as environmental non-genetic drivers of CH. To clarify relationships between immune-dysfunctional diseases and CH, here we tested impact of various challenges on TedCH. We found that expedited TedCH depended on establishment of inflammatory environment. Primary or chimeric *Tet2*-mutant mice spontaneously developed co-symptoms reminiscent of human chronic colitis and myeloid leukemia, which was exacerbated by feeding with DSS, an experimental inducer of ulcerative colitis. Single cell RNA-seq (scRNA-seq) analysis reveals in depth the damage of colon and bone marrow in the *Tet2*-mutant mice in physiological condition or fed with DSS, along with increase of dysbacteriosis indicated by gut microbiome analysis. Results from colon scRNA-seq from both mouse and human highlight important roles of PTX3/IL-1β pro-inflammatory signaling in promoting colitis or leukemia. Finally, TedCH trajectory and inflammation in colon and bone marrow was ameliorated by treatment of IL-1R1 inhibitor Anakinra. Our study suggests that PTX3/IL-1β signaling and clonal hematopoiesis cooperates and plays important roles in gut-bone marrow axis and related diseases including colitis and leukemia.

Key words

Clonal hematopoiesis, colitis, leukemia, inflammation, *TET2, PTX3*, *IL-1β*, Anakinra

Highlights

1. Certain environmental factors such as Dextran Sulfate Sodium (DSS), an experimental inducer of ulcerative colitis, promotes *Tet2*-difficiency related clonal hematopoiesis (TedCH)

2. Colitis and leukemia are spontaneously and simultaneously developed in *Tet2*-defficient primary or chimeric mice, along with increased pathogenic gut microbiomes, indicating aberrant gut-bone marrow axis in the mutant mice.

3. Single cell RNA-seq analysis reveals enhanced PTX3, a soluble pattern recognition molecule (PRM) and IL-1β pro-inflammatory signaling in colitis and leukemia.

4. *In vivo* function of the PTX3/IL-1β pro-inflammatory signaling in TedCH is also indicated in human colitis and validated experimental settings.

Running Title

Colitis and Leukemia Co-modulated by PTX3/IL-1β Signaling and Clonal Hematopoiesis

Introduction

Clonal hematopoiesis (CH) is a phenomenon whereby hematopoietic stem cells (HSCs) carry genetic mutations with advantageous growth potential, resulting in aberrant expansion of certain immature and mature hematopoietic cell population over time (Genovese et al., 2014; Steensma et al., 2015). Similar phenomenon, somatic mutation-driven clonal expansion of certain cells, takes place in non-hematopoietic organs such as skin, esophagus, and brain (Jaiswal and Ebert, 2019). Recent studies suggest that CH is associated with an increased risk of developing hematological malignancies including acute myeloid leukemia (AML) and myelodysplastic syndrome (MDS). Scoring CH is able to stratify high-risk CH-carriers (Weeks et al., 2022). Interestingly, CH is also associated with non-hematological diseases including all-cause mortality, cardiovascular disease, chronic obstructive pulmonary disease, and gout (metabolic arthritis, an auto-immune disease in bone juncture) (Agrawal et al., 2022; Jaiswal and Libby, 2020; Jaiswal et al., 2017; Miller et al., 2022). Accordingly, World Health Organization recently recognizes that CH is a precursor lesion of myeloid neoplasms (Khoury et al., 2022). Therefore, characterizing drivers and consequences of CH will not only assist in investigating clonal expansion of somatic cells in non- hematopoietic organs but also help in understanding the etiology of numerous chronic diseases many years prior to their onset (Weeks et al., 2022).

Loss-of-function heterozygous mutations in the epigenetic regulator *TET2* (Ten-Eleven Translocation methyl-cytosine dioxygenase-2) were among the first discovered and one of the most prevalent and important drivers of CH. Unsurprisingly now, mutations in *TET2* are also the top prevalent single-gene mutations in MDS, an age-related hematological disease with abnormal hematopoietic stem and progenitor cells (HSPCs) (Delhommeau et al., 2009; Nazha et al., 2021). *Tet2*-deficient primary or chimeric mouse models display a competitive advantage over *Tet2*-deficient HSPCs, skewing myeloid differentiation, and a tendency to transform into full-blown leukemia when co-operative mutations and environmental factors involved in (e.g., $Flt3^{ITD/+}$ or $NRas^{G12D}$) (Moran-Crusio et al., 2011) (Cai et al., 2020a; Chu et al., 2012). Loss of *TET2* results in increase of DNA methylation at various genomic sites, such as promoters, enhancers, and CpG islands (Rasmussen et al., 2015). Recently we and others reported that *Tet2*-deficient HSPCs exhibit significantly increased expression of the inflammatory cytokine interleukin-6 (IL-6) and show stronger anti-apoptotic and self-renewal abilities after exposure to acute stimulus such as IL-6 or lipopolysaccharides (LPS), leading to the development of an age-dependent chronic myelomonocytic leukemia (CMML)-like disease in the mutant mice over time (Cai et al., 2018; Zhang et al., 2015). Microbial signals are a major source of inflammatory inducer and interestingly *Tet2* mutant mice with antibiotic treatment or raised in germ-free condition developed minimal CMML-like phenotypes (Meisel et al., 2018). In line with these studies, a most recent study based on *in vivo* models rather than generating chimeric *Tet2*-mutant models demonstrated that interleukin-1 (IL-1α and IL-1β), is able to act as an external factor in driving *Tet2*-deficiency related clonal hematopoiesis (TedCH) over age; genetic loss of their receptor *IL1-R1* mitigated *Tet2*-deficiency related abnormalities (Burns et al., 2022; Caiado et al., 2023). In summary, results from several independent studies strongly suggest that *Tet2*-deficient HSPCs manifest TedCH phenomenon and cooperate with intrinsic and extrinsic factors to develop stronger symptoms such as full-blown leukemia AML.

However, through mathematical analysis and experimental validation, a recent

study of TedCH suggests that hypercholesterolemia-induced atherosclerosis took place prior to clonal hematopoiesis and may act as a driver of *Tet2*-deficient HSCs proliferation on the contrary. This study suggests that certain diseases or environmental conditions, for example, atherosclerosis drives the onset and trajectory of TedCH (Heyde et al., 2021). Atherosclerosis is a complicated and chronic disease where inflammation dysregulation explains in part (if not all) the etiology: mononuclear cells from the blood, including white blood cells, are recruited to the vessel wall in response to tissue damage (Libby et al., 2002). To further demonstrate that "diseases" was able to take prior to TedCH and drive TedCH, the authors presented an additional experimental model, a sleep fragmentation model, and demonstrated that sleep disturbance drives TedCH (Heyde et al., 2021). Nonetheless, clarifying the drivers and consequences of TedCH is the most critical challenge in the area. Controlling the trajectory of TedCH will also assist in developing strategies to mitigate CH-related diseases.

In previous studies, we have identified LPS, an inducer of innate-immunity related TLR4 signaling, and mutations in *Flt3*, *Ins2* or lncRNA *Morrbid* cooperates with *Tet2* deficiency to exacerbate or mitigate TedCH or leukemic phenotypes(Cai et al., 2020a; Cai et al., 2018; Cai et al., 2021; Cai et al., 2020b). To extensively explore the potential drivers of CH, we further assessed the impact of another five different environmental factors on TedCH and confirmed that accelerated TedCH depends on the establishment of inflammatory environment. Based on the results of single cell RNA-seq in the organs of colon and bone marrow (mouse and human both included) and microbe sequencing in gut, we propose that PTX3/IL-1β signaling pathway, a tight complement/pro-inflammation cascade, play important roles in gut-bone marrow axis which in turn regulates clonal hematopoiesis. We suggest that the PTX3/IL-1β signaling should be appreciated to mitigate TedCH trajectory and related diseases including colitis and leukemia.

Results
DSS but not STZ, 5-FU, or additional irradiation promotes TedCH
Competitive bone marrow transplantation (cBMT) assays were developed for analyzing TedCH in the chimeric mice and four different treatments were chosen for testing environmental impacts on TedCH: DSS, STZ, 5-FU, and additional irradiation (**Figure 1A**). 5-FU and additional irradiation are well-known direct regulators of hematopoiesis while DSS induces colitis and systematic inflammation and STZ induces hypoglycemia. TedCH trajectory was monitored every 3 weeks by analyzing CD45.1, CD45.2 and CD45.1/CD45.2 chimerism in the peripheral blood (PB) of the chimeric mice. As shown in **Figure 1B** and **C**, expedited TedCH were observed upon DSS feeding compared to vehicle (Veh) feeding. Treatments with STZ or additional irradiation appear to have no effect on TedCH while 5-FU has significant inhibition on TedCH (**Figure 1D** to **F; Supplemental Figure 1A** to **C**). As DSS induces inflammation and colitis, we conclude that inflammation and colitis positively cooperate TedCH trajectory in the cBMT experimental setting.

DSS exacerbates colitis burden and aberrant dysbacteriosis in *Tet2*$^{+/-}$ primary mice
Since DSS expedited TedCH in the chimeric mice, we then asked in detail what are the in-middle biological and pathological processes and consequences during the

DSS challenge. We chose *Tet2*-deficient primary mice for further study since they are more stable and readily monitored in a large cohort without irradiation involvement (irradiation itself may damage gut histology). A previous report suggests that acute DSS challenge exacerbate inflammatory bowel disease (IBD) in *Tet2*-deficient primary (Zhang et al., 2015). We turned to ask about the impact of chronic colitis by three cycles of low-dose DSS feeding (**Figure 2A**). Consistently, even low-dose of DSS also induced exacerbated colitis in *Tet2*-deficient primary mice (*Tet2*$^{+/-}$_DSS vs. WT_DSS, **Figure 2B-E**). Upon DSS challenge, increased diseases score (based on colon H&E staining) were observed and confirmed by expression of gut barrier markers *Muc2* and *Occludin* (**Figure 2F** and **G**; and **Supplemental Figure 2A**). Interestingly, *Tet2*-deficient primary mice appear to develop colitis spontaneously as shown by FITC-D staining in the blood serum and H&E staining (*Tet2*$^{+/-}$_Veh vs. WT_Veh, **Figure 2H**; and **Supplemental Figure 2A**).

To ask to what extent *Tet2*-deficient primary mice manifest aberrant colon immunity, we performed 16S RNA sequence and compared the microbial composition between WT and *Tet2*-deficient primary mice (*Tet2*$^{+/-}$_Veh vs. WT_Veh), along with the comparison between WT and *Tet2*-deficient primary mice fed with DSS (*Tet2*$^{+/-}$_DSS vs. WT_DSS). Results of the microbial composition data calculated by principal component analysis (PCA) suggest the four groups distinguish each other very well (**Figure 2I**). Although the Simpson index of operational taxonomic units (OUT, measuring microbial community composition) level fails to distinguish the four groups (a higher Simpson index indicates lower community diversity, **Figure 2J**), the Chaos index of OUT level significantly distinguishes microbial communities of *Tet2*$^{+/-}$_Veh mice from that of WT_Veh mice (a higher Chaos index indicates higher community richness, **Figure 2K**). Upon DSS challenge, the Chaos index of OUT level in *Tet2*$^{+/-}$_DSS and WT_DSS were decreased compared to that in *Tet2*$^{+/-}$_Veh and WT_Veh, suggesting certain bacteria were dominant expanded. Further microbial analysis suggests that: (1) at phylum level, *Tet2*_DSS mice exhibited most increments in *Proteobacteria* and *Actinobacteria* (both are pathogenic and toxic to the animals, **Figure 2L**); (2) at genus level, different microbiome compositions are revealed for the comparison between *Tet2*$^{+/-}$_Veh and WT_Veh (**Figure 2M**). Detailed comparison between *Tet2*$^{+/-}$_Veh and WT_Veh or between *Tet2*$^{+/-}$_DSS and WT_DSS reveal that aberrant changes in microbial composition including probiotic microbials (circled by blue rectangles) or pathogenic microbials (circled by pink rectangles) (**Figure 2N** and **O; and Supplemental Figure 2B** and **C**). Analysis combining hematological parameters and microbial composition further confirm that the pathogenic microbial *Proteobacteria* are involved in alteration of Hemoglobin level and neutrophil percentage (**Supplemental Figure 2D** and **E**). Taken together, results from colon pathology and microbial 16S RNA sequencing suggest that without DSS feeding, *Tet2*-deficient primary mice have already manifest subtle colitis and aberrant microbial composition, especially increase of pathogenic microbials; upon DSS challenge, *Tet2*-deficient primary mice manifest exacerbated colitis and growth of pathogenic microbial in the colon tissue.

DSS induces aberrant hematopoiesis and skewed myelopoiesis, which are consistently revealed by flow cytometry and scRNA-seq analysis

To directly clarify the DSS effect on hematopoiesis, cells from blood, spleen and bone marrow were subjected to hematological cell counts, histology, and flow cytometry analysis. Consistently with our previous report that *Tet2*-defficient primary

mice manifest skewed myelopoiesis with or without LPS treatment (Cai et al., 2018), we observed similar trends in the *Tet2*-defficient primary mice compared to wild-type controls (**Supplemental Figure 3A** to **H**). As flow cytometry relies only on limited molecular markers but scRNA-seq transcriptomic assays can offer much more comprehensive landscape of various HSPCs, Lin⁻ bone marrow cells from the 4 groups of animals were sorted for scRNA-seq analysis. After quality controls, around 5000 to 9000 cells

from each group were included in the further annotation analysis (**Figure 3A**). A total of 11 main populations including hematopoietic stem cells (HSC) and granucyte-monocyte progenitors (GMP-1, GMP-2, and GMP-3) are annotated and have relative different developmental hierarchies, indicated by the relative position in the plots of Uniform Manifold Approximation and Projection (UMAP). Cells that are very close to mature neutrophils, monocytes and B cells are also annotated and marked as neutrophil progenitors (NP), monocyte progenitors (MP) and progenitor of B cells (Pro_B). Each group has all of these 11 cellular populations and their compositions are quantified at percentage portion (**Figure 3B**). The expression of the annotation markers supports the developmental identify of the cells (**Figure 3C** and **D**; and **Supplemental Figure 4A**). Interestingly, in the one-by-one comparison, markers for HSCs (*Ifitm1*, *Egr*1 and *Nr4a1*) are upregulated in the comparison between *Tet2*⁺/⁻_Veh vs. WT_Veh, or between *Tet2*⁺/⁻_DSS vs. *Tet2*⁺/⁻_Veh, or between WT_DSS vs. WT_Veh, or between *Tet2*⁺/⁻_DSS vs. WT_DSS (HSCs genes are underlined in **Figure 3D** and **E**; and **Supplemental Figure 4B**). Analysis of differentially expressed genes (DEGs) reveals that stemness-related genes including *Egr1* are upregulated in the pools of HSC and GMP-1, as indicated in the scRNA-seq dataset in the comparison between *Tet2*⁺/⁻_Veh vs. WT_Veh or between *Tet2*⁺/⁻_DSS vs. WT_DSS (**Supplemental Figure 4C**). The heatmap of the top 50 signaling pathways in the pool of HSC and GMP-1 further indicate that inflammatory signaling pathways such as IL6-Jak-Stat3 signaling, inflammatory response, interferon alpha and grammar response are upregulated in the *Tet2*⁺/⁻_Veh compared to WT_Veh; similar trends are observed in the *Tet2*⁺/⁻_DSS compared to WT_DSS (**Supplemental Figure 4D** to **F**). Interestingly, we also observed that higher cell-cycle pathways take place in the HSC and GMP-1 of *Tet2*⁺/⁻_Veh compared to WT_Veh, or in the HSC and GMP-1 of *Tet2*⁺/⁻_DSS compared to WT_DSS (**Supplemental Figure 4D** to **F**). When numerous candidate marker genes that are critical for HSCs are considered, we found that the expression level of *Cd34* and *Il1r1* are increased (notice the max level of each panels of the four groups) and the their expression "region" are expanded (notice the circle in dashed-line) (**Figure 3G**), indicating pro-inflammatory IL1b signaling is involved in the HSCs of Tet2mut_Veh and the DSS-treated mice, compared to that of WT_Veh controls. RNA velocity analysis also suggest that the hematopoietic developmental trajectory is altered in Tet2_Veh or DSS treated mice compared to WT_Veh (**Figure 3H** and **Supplemental Figure 4G**) (Bergen et al., 2020; La Manno et al., 2018; Li et al., 2023). Taken together, scRNA-seq results revealed similar skewed myelopoiesis as shown by flow cytometry and also revealed alterations specific to the pools of HSC and GMP in the *Tet2*⁺/⁻_Veh or *Tet2*⁺/⁻_DSS mice compared to the respective controls.

scRNA-seq analysis of colon tissue highlights that cell-to-cell talk between *Ptx3*⁺ fibroblasts and *Il-1β*⁺ monocytes in promoting inflammation

Colon tissue from the 4 group of mice were also used for scRNA-seq analysis since this approach can provide a full-map of cell-to-cell talk at the single-cell resolution. After quality controls, around 3000 to 6000 cells from each group were

included in the further annotation analysis (**Figure 4A**). A total of 5 main populations including epithelial cells, stromal cells, and immune cells (myeloid cells, B cells, and T cells) are annotated in the UMAP plot for the first step (**Figure 4A**). Each group has these five main cell populations. However, *Tet2*$^{+/-}$_Veh, WT_DSS and *Tet2*$^{+/-}$_DSS appear to have a higher percentage of stromal cells compared to the WT_Veh control (**Figure 4B** and **C**). Expression density of the respective markers for the five main cell population is shown in **Figure 4D**. The overall enriched signaling pathway analysis suggests that inflammatory signaling pathway is upregulated in the myeloid cells of *Tet2*$^{+/-}$_Veh, WT_DSS and *Tet2*$^{+/-}$_DSS compared to the WT_Veh control (green rectangle, **Figure4E;** and **Supplemental Figure 5A**). Importantly, cell-cycle related pathway is also upregulated in the myeloid cells of *Tet2*$^{+/-}$_Veh compared to WT_Veh control (pink rectangle, **Figure4E**). Interestingly, we also observed upregulated epithelia-mesenchymal-transition pathway (EMT) in the stromal cells of *Tet2*$^{+/-}$_Veh, WT_DSS and *Tet2*$^{+/-}$_DSS compared to the WT_Veh control (blue rectangle, **Figure4E**; and **Supplemental Figure 5C**). Overall, the pathway enrichment analysis in each single cell population suggests that myeloid cells and stromal cells grossly have the most alterations in the comparison between *Tet2*$^{+/-}$_Veh vs. WT_Veh or between WT_DSS vs. WT_Veh. We then filtered out myeloid cells and stromal for further cell-to-cell talk analysis.

First, stromal cells and myeloid cells from the colon UMAP pool were subject to further clustering to distinguish sub-populations. In total, 8 sub-populations of cells in the stromal cells and 7 sub-populations of cells in the myeloid cells are annotated according to the typical marker gene expression (**Figure 4F;** and **Supplemental Figure 5B**). Expression of *Ptx3* and *Il-1β* are plotted in the UMAP space of colon tissue, stromal cells, or myeloid cells as indicated (**Figure 4G**). To our surprise, expression of *Ptx3* appears to be specific to stromal cells in the colon while expression of *Il-1β* appears to be specific to stromal cells in the colon in the scRNA-seq dataset.

Five sub-populations from the stromal cells and six sub-populations from the myeloid cells had the most alterations in the percentage and therefore were used for calculating cell-to-cell talk strength (**Figure 4H**). Indeed, *Cx3crl1*$^{+}$-myeloid cells appear to have stronger interaction with *Ptx3*$^{+}$-stromal cells in *Tet2*$^{+/-}$_Veh compared to WT_Veh (left two panels, **Figure 4H**). Myofibroblasts also appear to strongly interact with many myeloid cells in *Tet2*$^{+/-}$_Veh compared to WT_Veh (left two panels, **Figure 4H**). Upon DSS feeding, interaction between *Ptx3*$^{+}$-stromal cells and several myeloid cells are dramatically activated (right two panels, **Figure 4H**).

To dissect the detailed molecular events in the *Il-1β*$^{+}$-myeloid cells and *Ptx3*$^{+}$-stromal cells, five of myeloid sub-populations were used for NicheNet assay (Browaeys et al., 2020). As shown in **Figure 4I** to **L**, IL-1β/IL1-R1 signaling is actively involved in the cell-cell interaction, associated with expression of many downstream genes encoding complement protein proteins or inflammatory ligands (*C1ra*, *C1s1*, *C3* and *Cxcl12*). Surprisingly, *Il1r1* along with *C1ra*, *C1s1*, *C3* and *Cxcl12* all are dominantly expressed in the stromal cells of colon, compared with other four main cell population (**Supplemental Figure 5D**). Taken together, our detailed cell-cell talk analysis based on the colon scRNA-seq dataset from the 4 groups of mice strongly suggest that PTX3/IL-1β axis recruits inflammatory and complement cascade to module colon immunity in *Tet2*-deficient mice fed with DSS or raised in normal condition.

Aberrant expression of *PTX3* and *IL-1β* are indicated clinical samples with

leukemia or colitis.

To guarantee that PTX3/IL-1β signaling involves in human colitis or leukemia, we searched publicly available datasets of bulk RNA-seq or scRNA-seq and compared their expression between disease group and healthy controls. Data shown in **Figure 5A** to **D** suggest that increased expression of *PTX3* is detected in the myeloid leukemia group (datasets with AML or MDS patients were analyzed). Interestingly, mutations in *TET2* appear to have higher expression of *PTX3* also (**Figure 5A** and **D**). Importantly, higher expression of *PTX3* has a worse prognosis in the TCGA-LAML cohort while expression of *IL-1β* failed to stratify the cohort (**Figure 5A** and **D**). Additionally, analysis of colitis samples also suggests that higher expression of *PTX3* and *IL-1β* took place in some cohorts of colitis samples compared to the controls (**Figure 5E** to **G**). Taken together, these results warrant that the PTX3/IL-1β bundle indeed involves in human colitis or leukemia.

As we surprisingly observed that *Ptx3* and *Il-1r1* are dominantly (if not specifically) expressed stromal cells of colon tissue in mice (**Figure 4G** and **M;** and **Supplemental Figure 7A** and **B**), we want to characterize if this pattern is conserved in human. We downloaded the publicly available scRNA-seq datasets of colitis and re-analyzed the cells (Smillie et al., 2019). Results from **Figure 5H** to **M**, along with **Supplemental Figure 6,** suggest that the percentage of *PTX3*[+]-fibroblasts indeed increased according to the grade of colitis. Although partial myeloid cells express *PTX3* themselves, *PTX3* is also dominantly expressed in stromal cells. Taken together, results from human clinical samples suggest that PTX3/IL-1β signaling may have important function in colitis and leukemia.

Pharmacological blockage of IL-1β signaling by Anakinra inhibits PTX3 expression and inflammation in colon and mitigates TedCH in blood

PTX3 is a soluble pattern recognition molecule (PRM) and has been reported to directly involves in IL-1β signaling (Bonavita et al., 2015). We postulate that the PTX3/IL-1β signaling pathway is present at the protein level in the colon of *Tet2*[+/-] mice with chronic inflammation. To validate this hypothesis, we performed western blot analysis to examine the expression of Il-1β and Ptx3 (**Supplemental Figure 7**C and **D**). Additionally, immunohistochemical (IHC) staining revealed high levels of Ptx3 expression in inflammatory cells infiltrating the colonic tissues of *Tet2*[+/-] mice (**Supplemental Figure 7E** and **F**).

As inhibitors specific to PTX3/Ptx3 activity are not available, we turn to use IL1R1 specific inhibitor Anakinra to assess the function of Ptx3 in the TedCH, colitis and aberrant hematopoiesis. cBMT assays for TedCH were conducted along with DSS and/or Anakinra treatment (**Figure 6A**). Consistent with its anti-inflammation function, application of Anakinra mitigate TedCH expansion in the mice feed with Vehicle or DSS (**Figure 6B;** and **Supplemental Figure 7G**). Aberrant hematopoiesis also appears to be partially rescued in the TedCH_Anakinra mice compared to TedCH_PBS mice (**Figure 6B;** and **Supplemental Figure 7H** and **K**). Importantly, application of Anakinra reduced the expression of both Ptx3 and Il-1β (**Figure 6D** and **E**). The reduced expression of Ptx3 upon Anakinra is also validated by immunohistochemistry (IHC) and pathologic index is largely ameliorated upon Anakinra treatment (**Figure 6F** to **I**). Taken together, results from Anakinra treatment suggest that inhibition of Il-1β signaling modulated Ptx3 expression and inflammation severity, along with TedCH trajectory and aberrant hematopoiesis.

Discussion

Since clonal hematopoiesis was for the first time recognized as a new risk medical entity to broadly predict diseases and even all-cause mortality around a decade ago, numerous genetic drivers or passengers of clonal hematopoiesis have been characterized, along with a recent report indicating their trajectories were able to be mathematically computed in human samples (Abelson et al., 2018; Fabre et al., 2022; Genovese et al., 2014; Weeks et al., 2022; Weinstock et al., 2023).To understand the mechanisms underlying CH, it was proposed that different genetic drivers of CH may favor corresponding environmental factors to strengthen or amplify the onset and consequence of CH (Caiado et al., 2021). For example, mutations in *TP53* and *PPM1D,* two top-10 prevalent drivers of CH, were selected when irradiation was applied in the therapy or after exposure to cisplatin and doxorubicin treatment (Wong et al., 2015) (Hsu et al., 2018). Our previous studies have shown that mutations in *TET2*, the first identified and one of the top-3 driver of CH, favors inflammatory condition on the contrary; blocking inflammation by genetic loss of *Morrbid* or administration of anti-inflammation drugs E3330 or SHP099 mitigate TedCH in experimental models (Cai et al., 2020a; Cai et al., 2018; Cai et al., 2021; Cai et al., 2020b). However, whether and how other environmental factors (positive, negative, or neutral to TedCH), intra-cell or intra-organ or host-pathogen communications drives TedCH remain incompletely understood.

To reach these aims, we first assessed the role of five different environmental factors in TedCH (4 factors are included in this study; and another one based on $Tet2^{+/-}$ HSPCs mixed with genetically stable inflammatory HSPCs will be reported elsewhere) and prioritized the impact of DSS, an inducer of infection with toxic pathogens and colitis in gut. We further characterized the impact of DSS on the *Tet2*-deficient chimeric or primary mice by multi-omic tools including single cell RNA sequencing or gut microbe sequencing. Although administration of the other three challenges, irradiation, 5-FU (depletion of immune cells and hematopoietic cells), or STZ (toxic damager of pancreases and inducer of high blood glucose) failed to accelerate TedCH, chronic colitis induced by three cycles of feeding with DSS or mixing with IL-1β-secreting genetically stable inflammatory HSPCs expedite the trajectory of TedCH. These results suggest that: 1) only certain specific environmental factors play a positive role to cooperate with TedCH in experimental models; 2) these "positive" factors may have a direct link to inflammation or innate immunity; 3) an intra-organ communication, gut-bone marrow axis, which could be stimulated by host-pathogen axis or just by dysregulated auto-immunity, play a role in TedCH and leukemogenesis.

These observations from us are consistent with a previous study by *Meisel* et al. who directly studied the pathogen-host interaction in *Tet2*-mutation mediated leukemic diseases. In the study, *Meisel* et al. systematically studied why *Tet2*-mutant primary mice did not develop MPN or CMML phenotypes with 100% penetration and proposed that microbial signals cooperate to induce pre-leukemic MPN phenotype in the mice (host-pathogen axis in TedCH-related diseases). Using *Tet2*-mutant mice raised in germ-free condition, the authors further solidly validate their conclusions (Meisel et al., 2018; Zeng et al., 2019). However, most of the experimental mice and our human are exposed to microbial environment during life. Where the initial inflammation take place (gut, bone marrow or even other organ?) and how it triggers CH and changes its trajectory remain to be elucidated.

Considering the importance of inflammation in *Tet2*-mutation related diseases, in previous studies we integrated multiplayers and proposed a working model for *Tet2*-

mutation related diseases: TedCH recruits a feed-forward loop and inflammation acts
as an amplifier in the loop for diseases progression including chronic leukemia and
acute leukemia (Cai et al., 2018; Cai et al., 2021). Given that DSS promotes TedCH in
chimeric mice, we then tested if DSS could promote *Tet2*-mediated CMML to
transform into full-blown AML. During the short-period of colitis, we indeed notices
exacerbated phenotypes in *Tet2*mutant_DSS mice in both gut permeability (worsen
colitis) and bone marrow hematopoiesis (malignant transformation to MDS/MPN-
like). Unlike that we observed full-blown AML in aged *Tet2$^{+/-}$;Flt3$^{ITD/ITD}$* or
Tet2$^{+/-}$;Ins2$^{+/-}$, we failed to observe full-blown AML in the DSS-challenged mice.
These results suggest that inflammation as an amplifier to induce full-blown AML
may depend on adequate time.
Furthermore, single-cell RNA-seq (scRNA-seq) today becomes a very powerful
tool to dissect the etiology of complicated diseases at single cell level and molecular
level. Both scRNA-seq and bulk RNA-seq were applied in this study to delineate the
widespread transcriptional changes in bone marrow and gut (note that only lineage-
negative bone marrow cells were sorted for scRNA-seq analysis to prioritize
alterations in hematopoiesis). Consistent with previous studies, functional enrichment
analysis of DEGs in HSC and CMP showed that stem cell self-renewal and immune
responses related genes were upregulated in *Tet2*-deficient mice treated with DSS.
These observations were unlikely to be gained years ago since minimal HSC and
CMP cells could be sorted for RNA-seq and the sorted cells may not represent their
physiological conditions. Indeed, analysis of scRNA-seq in gut of *Tet2*-mutant_veh
and *Tet2*-mutant_DSS mice helps us to reveale a previously unappreciated alteration
in *Ptx3$^+$*-fibroblast of gut and the inferred PTX3/IL-1β signaling [note that mice were
raised in specific pathogen-free (SPF) condition with or without feeding with DSS].
These results make us conclude that *Tet2*-mutant mice appear to develop colitis and
leukemia simultaneously, very subtle if not strong, reminiscent of human symptoms.
Consistently, the involvement of PTX3/IL-1β inflammatory signaling is also
confirmed when we re-analyzed human colitis scRNA datasets.
Recent studies report that patients with inflammatory bowel disease (IBD) are
associated with clonal hematopoiesis and mutations in *DNMT3A* and *PPM1D* are
prevalent although mutations in *TET2* is not dominant in the cohort on the contrary
(Zhang et al., 2019). The colitis patients harbor mutant hematopoietic clones and
often present with clinical symptoms of MDS/AML (Cumbo et al., 2022). Moreover,
intestinal barrier dysfunction is also reported at diagnosis in patients with MDS and
AML (Khan et al., 2021). Along with these clinical observations, we conclude that
Tet2-mutant mice manifest co-symptoms of colitis and leukemia and that a molecular
link, PTX3/IL-1β signaling, may modulate both diseases in the mice.
Changes in the gut microbiome are considered a key regulator of the conversion of
normal hematopoiesis to diseased hematopoiesis, such as anemia and neutropenia
caused by IBD or long-term antibiotic use (Deshmukh et al., 2014). A study has
shown that an imbalance in the microbiota induced by antibiotic exposure or sterile
conditions can lead to dysregulation of hematopoiesis (Yan et al., 2018). Our data
from 16S rRNA gene sequencing indicated that *Tet2$^{+/-}$* mice had disturbed gut
microbiota with significantly reduced probiotic abundance prior to the addition of
inflammatory factors. For example, chronically infected *Tet2$^{+/-}$* mice exhibited
increased *Proteobacteria* and *Actinobacteria*, two toxic gut pathogens to mammals.
We also observed that there was a positive correlation between these microbiotas and
PB hematological parameters (**Supplemental Figure 2E**). Previous clinical studies
have shown that patients undergoing hematopoietic stem cell transplantation (HSCT)

have reduced gut microbiota diversity and increased *Proteobacteria*, which correlates with the risk of patients developing bloodstream infections during HSCT (Taur et al., 2012). These results suggest a potential role of microbial signaling in exacerbating the disease burden of IBD in $Tet2^{+/-}$ mice as well as hematologic malignancy transformation. Functional analysis in the appropriate facility with adequate biological protection, for example pathogen infection or fecal microbial transplantation, will offer direct evidence with regards the interaction between specific-pathogens and TedCH or *Tet2*-mutant mice (Zeng et al., 2023).

Consistent with previous studies (Burns et al., 2022; Caiado et al., 2023), in the context of DSS challenged-TedCH, our data suggest that anakinra eliminates the $Tet2^{+/-}$ clones in the clonal expansion in neutrophil and HSPC compartments, inhibits gut inflammation, and restores the adaptive hematopoietic injury induced by chronic inflammation. However, it does not fully restore the established clonal dominance in the HSPC pool, which may be partially addressed by increasing the treatment dose and duration. Dissecting the role in a cell-type specific deletion will assist in further understanding the cellular communications, i.e., immune cells with epithelia cells and offer a new knowledge for understanding gut-bone marrow axis and co-symptoms of colitis and leukemia. Based on our pharmacological data using the IL-1R inhibitor Anakinra, it is feasible to speculate that: 1) cocktails of antibiotics administration would also be able to slowdown TedCH, which should be tested in future studies; 2) blocking the IL-1β/PTX3 axis may represent a promising therapeutic target for preventing and treating infection-related clonal hematopoiesis and associated diseases. Since specific PTX3 inhibitors, except PTX3 antibodies, are still not available in research-grade or clinical-grade, developing PTX3 inhibitors or combining them with methylation drugs may represent potential therapeutic approaches for targeting *TET2* mutant hematopoiesis in the future.

In summary, the present study demonstrates that gut-bone marrow axis plays an important role in the co-symptoms of colitis and leukemia in the *Tet2*-mutant chimeric and primary mice. Dysbacteriosis, chronic inflammation and clonal hematopoiesis are all assembled to promote the progression the co-symptoms over age. Our single-cell transcriptomic analysis highlights and appreciates the role of a population of gut $Ptx3^{+}$-fibroblasts in driving the entire diseases. Mitigation of inflammation by IL-1R1 inhibitor Anakinra can downregulate Ptx3, repair gut barrier and slowdown TedCH. The study therefore demonstrates that PTX3/IL-1β signaling-associated clonal hematopoiesis plays important roles in gut-bone marrow axis and related chronic diseases including colitis and leukemia.

Materials and Methods
Materials and Methods including mice breeding, experimental procedures, mouse disease scoring and histology, computational analysis of single-cell RNA-seq and gut microbiome sequencing, collection, and analysis of clinical RNA-seq datasets, and statistical analysis are provided in detail in the **Supplemental File-1**. Computational environments with Linux and R language for analyzing scRNA-seq data and gut microbiome sequencing data have been described in our previous study (He et al., 2022). Custom scripts for analyzing and visualizing the omics results are available upon reasonable request.

Conflict of Interest
The authors declare that the research was conducted in the absence of any commercial

549   or financial relationships that could be construed as no potential conflict of interest.
550

Author Contributions
551
ZC and HH conceived and designed the study, wrote the computational scripts for
sequencing analysis, visualized the results, drafted, and wrote the manuscript. HH
performed most of the experiments and analysis. YW, HY, QH, WJ, ZW and JD
assisted the experimental verification or validations from other database/cohorts and
edited the manuscript. HW and ZZ contributed critical reagents and analyzed the data.
All authors contributed to the editing and revision of the manuscript.
558

Funding
559
560   This work was supported in part by grants from the Tianjin Medical University Talent
561   Program and from National Science Foundation of China to ZC (No.82170173).
562

Acknowledgments
563
We thank members of Cai laboratory and colleagues of Tianjin Medical University for
their technical and administration support, critics and helpful suggestions improve the
manuscript. We would also thank Drs. Mingjiang Xu, Tao Cheng, Hui Cheng,
Zhiqiang Liu for sharing us reagents.

Figure Legends:
Figure 1. Assessing impacts of various treatments on TedCH trajectory.
(A) Competitive bone marrow transplantation (cBMT) assays were used for searching
cooperative environmental factors of TedCH. See Methods for detailed cBMT
experimental procedures. Treatments were applied once TedCH is set up at the 4~6
weeks post cBMT. Results of four different treatments (DSS, STZ, 5-FU and
additional irradiation) were included in this study. As shown in the Figure 1A, DSS
expedited TedCH. Expedited TedCH was also observed when $Tet2^{+/-}$ HSPCs were
mixed with genetically stable inflammatory HSPCs and the detail results will be
reported in another study (Wen and Cai $et\ al$., manuscript in preparation, 2023).
cBMT, competitive bone marrow transplantation; PB, peripheral blood.
(B and C) Representative flow profiles of PB from the chimeric mice feed with DSS
or normal water (vehicle, veh) (B) and the quantified trajectories of TedCH in the
chimeric mice (C).
(D-F) Quantified trajectories of TedCH in the chimeric mice treated with STZ (D), or
5-FU (E), or additional irradiation (F) as indicated.
Data are shown as means ± SEMs. Number of biological repeats (animals): n = 5~7.
Experiment of DSS treatment were repeated twice. Experiments of STZ or 5-FU or
additional irradiation treatment were performed once.*, $p < 0.05$; **, $p < 0.01$; ***, p
< 0.001; ****, $p < 0.0001$.
See also **Supplemental Figure 1** for representative flow profiles of PB from the
chimeric mice with or without STZ, 5-FU and additional irradiation treatment.
590

Figure 2. $Tet2$-deficient mice manifest exacerbated colitis and aberrant
microbiota composition when fed with DSS or vehicle.
(A) Experimental scheme for induction of chronic colitis by DSS. Three cycles of
DSS feeding were performed during the 50-day experimental procedure. Four groups
of animals were included in the entire procedure: WT_Veh, $Tet2^{+/-}$_Veh, WT_DSS,
$Tet2^{+/-}$_DSS.
(B and C) Body weights and disease scores (disease activity index, DAI) were
monitored daily and plotted during the induction.

(D-F) *Tet2*-deficient mice manifest exacerbated colitis based on colon pathology.
Colons from the 4 groups of mice were photographed (D) and quantified based on
colon length (E). Damage scores of colons were monitored and plotted based on H&E
staining (F).
(G) Expression of two classic gut barrier markers *Muc2* and *Occludin* in colon were
quantified by qRT-PCR.
(H) Damage of gut barrier were quantified by FITC-dextran staining in serum of PB.
(I-N) Microbiota (16S rRNA) sequencing analysis of gut in mice feed with DSS or
vehicle. The PCA plot of the 4 groups of microbiota samples suggests obvious
separation in microbial composition (I). α-diversity of the microbial community based
on observed operational taxonomic units were quantified by Simpson or Chao index
(J and K). Average relative abundance of prevalent microbiota at the phylum level or
at the genus level among the four groups (L and M). Detailed comparison of relative
taxon abundance between the $Tet2^{+/-}$_Veh vs. WT_Veh or between $Tet2^{+/-}$_DSS vs.
WT_DSS are also shown in (N) and (O).
PCA, principal component analysis.
Data are shown as means ± SEMs in A to H. Number of biological repeats (animals):
n = 3~7. Experiments of DSS treatment on primary mice were repeated three times. *,
p < 0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001.
See also **Supplemental Figure 2** for representative photography of H&E staining of
colons and additional microbiota analysis.
620
Figure 3. Single-cell RNA-seq analysis of HSPCs from *Tet2*-deficient mice when
feed with DSS or vehicle.
(A) UMAP plot showing 11 main populations in Lin⁻ BM cells. A total of 26,341
high-quality cells with an average of about 3000 genes per cell were included in the
UMAP plot. After quality control during the dataset analysis, the group of WT_Veh
has 7267 cells; *Tet2mut_Veh* has 4793 cells. WT_DSS has 5270 cells; *Tet2mut_DSS*
has 9011cells. 1, HSC: hematopoietic stem cell; 2, CLP: common lymphoid
progenitor; 3, GMP-1: granulocyte-macrophage progenitor-1, which may be close to
CMP (common myeloid progenitor) in flow cytometry; 4, GMP-3: granulocyte-
macrophage progenitor; 5, MMEP: mast and megakaryocyte-erythroid progenitor; 6,
GMP-2: granulocyte-macrophage progenitor-2, which may be close to monocyte-
dendrite progenitor; 7, MP: monocyte progenitor; 8, MkP: megakaryocyte progenitor;
9, NP: neutrophil progenitors; 10, ErP: erythroid progenitor; 11, Pro_B: progenitor of
B cell.
(B) Stacking bar plot showing the portion of the 11 main annotated populations in
each Lin⁻ BM cell sample.
(C and D) Representative annotation markers of the 11 main cell populations in the
scRNA-seq analysis of Lin⁻ BM cell. In C, expression of *Flt3*, *Cebpe*, *Il6ra* and *Spi1*
are highlighted in WT_Veh dataset. In D, each cell population has 4 typical markers in
the dot plots.
(E) Top 20 DEGs (10 up-regulated and 10 down-regulated) are showed in each
comparison when average expression of genes is calculated in each sample.
(F and G) Expression level and region of *Cd34* and *Il1r1* in the 4 group of scRNA-seq
datasets.
(H) Hematopoietic cell trajectory inferred by RNA velocity with 24 selected critical
genes, which are with known function in hematopoiesis. Notice the abnormal
trajectory in NP and MP region, compared to WT_Veh.
DEG, differential expressed gene. UMAP, uniform manifold approximation and

projection.
See also **Supplemental Figure 3 and 4** for flow cytometry results and additional
analysis of scRNA-seq Lin⁻ BM dataset.
652
Figure 4. Single-cell RNA-seq analysis of colon tissue from *Tet2*-deficient mice
when feed with DSS or vehicle.
(A) UMAP plot showing 5 main populations in colon tissues. After lysis of colon
tissue, all cells from the colon tissue were subjected for scRNA-seq. A total of 18,092
high-quality cells with an average of about 3000 genes per cell were included in the
UMAP plot. After quality control during the dataset analysis, the group of *WT_Veh*
has 5589 cells; *Tet2mut_Veh* has 5649 cells; *WT_DSS* has 3382 cells; *Tet2mut_DSS*
has 4372 cells. The 5 main populations include epithelial cells (red), stromal cells
(yellow) and three major types of immune cells: myeloid cells (green), B cells (blue)
and T cells (dark blue).
(B) Four individual UMAP plots are separately showed according to the sample
source.
(C) Stacking bar plot showing the portion of the 5 main annotated populations in each
colon sample.
(D) Expression of representative annotation markers for the 5 main cell populations in
the UMAP plot of colons. *Dcn* for stromal cells; *Krt*8 for epithelial cells; *Lyz2* for
myeloid cells; *Igha* for B cells; and *Trbc2* for T cells.
(E) Heatmap of dysregulated biological pathways in the 5 populations of colon tissues
from 4 groups of samples. A full version of the heatmap is showed in the
Supplemental Figure 5. Of note, Il6-Jak-Stat3 signaling, inflammatory response and
interferon alpha response are upregulated in *Tet2mut_Veh, WT_DSS, and*
Tet2mut_DSS in myeloid cell, compared to *WT_Veh* (circled by green rectangle).
Additionally, proliferation-related pathways appear to be upregulated in myeloid cells
of *Tet2mut_Veh* compared to *WT_Veh;* proliferation-related pathways appears to
upregulated in T cells of *WT_DSS* and *Tet2mut_DSS* compared to *WT_Veh* (circled by
pink rectangle). Interestingly, the epithelial mesenchymal transitional (EMT) pathway
appears to be dysregulated in *Tet2mut_Veh, WT_DSS,* and *Tet2mut_DSS* in myeloid
cell, compared to *WT_Veh* (circled by blue rectangle).
(F) Stromal cells and myeloid cells are subjected for further clustering by UMAP as
they have most significantly upregulated biological pathways as indicated in (E). Top
panel, UMAP plot of stromal cells and stacking plot showing potion of the sub-
population; bottom panel, UMAP plot of myeloid cells and stacking plot showing
potion of the sub-population. As indicated, 10 subpopulations and 7 subpopulations
are annotated in stromal cells and myeloid cells respectively.
(G) Expression of *Ptx3* and *Il-1β* in the colon tissues (left panel) or in stromal cells
(up-right panel) or in myeloid cells (bottom-right panel), respectively.
(H) Heatmap of cell-to-cell talk strength between subpopulations of stromal cells and
myeloid cells.
(I-L) Detailed molecular events of the cell-to-cell talk between myeloid cells and
stromal cells. Of note, *Ptx3*⁺ fibroblasts and *Il-1β*⁺ monocytes appear to be most
involved. *Il-1β*⁺ monocytes express numerous ligands including *Il-1β* (I and J). *Il-1β*
binds *Il-1r1* in *Ptx3*⁺ fibroblasts and stimulate a couple of downstream genes (K and
L).
(M) Expression of *Il-1β* receptor *Il1r1* and relevant genes encoding complement
components or inflammation regulators (*C1ra, C1s1, C3* and *Cxcl12*) were plotted on
UMAP of colon tissue (top panel) or on the UMAP plot of stromal cells (bottom
14

699 panel).
700 See also **Supplemental Figure 5** for additional analysis of scRNA-seq colon dataset.
701

Figure 5. Aberrant PTX3/IL-1β signaling is indicated in human leukemia or
colitis through analysis of clinical samples.
(A) Expression of *PTX3* and *IL-1β* in unsorted AML patients (left panel) or in AML
patients carrying *TET2* mutation (right panel). Bulk RNA-seq datasets of bone
marrow were included in the analysis.
(B-C) Prognosis value of marker genes *PTX3* and *IL-1β* in the TCGA-LAML cohort.
(D) Expression of *PTX3* and *IL-1β* in unsorted MDS patients (left panel, bone marrow
samples) or in MDS patients carrying *TET2* mutation (right panel, bone marrow
samples).
(E-G) Expression of *PTX3* and *IL-1β* in patients with ulcerative colitis and healthy
controls, based on three independent cohorts. Bulk RNA-seq datasets of colon tissues
were included in E and F; Bulk RNA-seq dataset of blood were included in G.
(H-L) Aberrant expression of *PTX3* and *IL-1β* was also detected using scRNA-seq
dataset of human colitis samples. The dataset is downloaded from the website Broad
Data Use Oversight System (https://duos.broadinstitute.org) and re-analyzed. The
cohorts include two groups of human colon samples: colons from healthy controls
(n=10) and colons from ulcerative colitis patients (n=7). Patient samples from both
non-inflamed region and inflamed region were n=7 respectively. Inflammation scores
suggesting the three group has significant difference overall (H). UMAP plots of the
human colon scRNA-seq dataset also show five main population as similar as Figure
4A: epithelia cells, stromal cells and three immune cells (myeloid cells, B cells and T
cells) (I). Stromal cells were highlighted by the dotted lines (I). Stromal cells were
further annotated by UMAP plots and $PTX3^+$ fibroblasts were screened out by dotted
lines (J). Of note, stacking plots indicate percentage of $PTX3^+$ fibroblasts were
increased according the inflammation grade of the colitis (K). Expression of *PTX3* is
shown in the UMAP plot of colon tissue (L) and three groups of stromal cells (M) as
indicated.
See also **Supplemental Figure 6** for additional analysis of human colitis scRNA-seq
dataset.
731

Figure 6. Pharmacological blockage of Il-1β signaling by Anakinra inhibits Ptx3
expression and inflammation in colon and mitigates TedCH in blood.
(A) Schematic representation of Anakinra treatment on the DSS-induced expedited
TedCH.
(B) Quantification of PB chimerism in TedCH mice treated with DSS and/or
Anakinra.
(C) Quantification of HSPCs including LSK, HSC, MPP2, MPP3/4, CMP and GMP
in 4 groups of TedCH mice.
(D-E) The proteins level of Il-1β and Ptx3 were measured in colon tissues and
quantified.
(F-G) Representative H&E staining of colon tissue from each group and diseases
score (damages in colon tissue) were quantified accordingly. Scale bar is 500 μm.
(H-I) Representative immunohistochemical (IHC) staining of Ptx3 in colon (H) and
semi-quantitative analysis of immunohistochemistry staining of Ptx3 based on
integrated optical density (IOD) (I). scale bar is 100 μm.
Data are shown as means ± SEMs in A-H. Number of biological repeats (animal
mouse): n = 5~7. Experiments of DSS treatment combining with Anakina on TedCH

mice were repeated twice. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.
See also **Supplemental Figure 7** for additional analysis of Ptx3 expression in TedCH
colons.
752
753
Figure 7. Schematic illustration for the working model about clonal
hematopoiesis cooperating with inflammation and dysbacteriosis in gut-bone
marrow axis to promote colitis and leukemia.
We propose that gut-bone marrow axis plays important role in the co-symptoms of
colitis and leukemia in the *Tet2*-mutant chimeric and primary mice. Dysbacteriosis,
chronic inflammation and clonal hematopoiesis are all assembled to promote the
progression the co-symptoms over age. See the main text for further discussion.
761
762
References

Abelson, S., Collord, G., Ng, S.W.K., Weissbrod, O., Mendelson Cohen, N., Niemeyer, E., Barda, N.,
Zuzarte, P.C., Heisler, L., Sundaravadanam, Y.*, et al.* (2018). Prediction of acute myeloid leukaemia risk
in healthy individuals. Nature *559*, 400-404.
Agrawal, M., Niroula, A., Cunin, P., McConkey, M., Shkolnik, V., Kim, P.G., Wong, W.J., Weeks, L.D.,
Lin, A.E., Miller, P.G.*, et al.* (2022). TET2-mutant clonal hematopoiesis and risk of gout. Blood *140*,
1094-1103.
Bergen, V., Lange, M., Peidli, S., Wolf, F.A., and Theis, F.J. (2020). Generalizing RNA velocity to
transient cell states through dynamical modeling. Nature Biotechnology *38*, 1408-1414.
Bonavita, E., Gentile, S., Rubino, M., Maina, V., Papait, R., Kunderfranco, P., Greco, C., Feruglio, F.,
Molgora, M., Laface, I.*, et al.* (2015). PTX3 is an extrinsic oncosuppressor regulating complement-
dependent inflammation in cancer. Cell *160*, 700-714.
Browaeys, R., Saelens, W., and Saeys, Y. (2020). NicheNet: modeling intercellular communication by
linking ligands to target genes. Nat Methods *17*, 159-162.
Burns, S.S., Kumar, R., Pasupuleti, S.K., So, K., Zhang, C., and Kapur, R. (2022). Il-1r1 drives
leukemogenesis induced by Tet2 loss. Leukemia *36*, 2531-2534.
Cai, Z., Aguilera, F., Ramdas, B., Daulatabad, S.V., Srivastava, R., Kotzin, J.J., Carroll, M., Wertheim,
G., Williams, A., Janga, S.C.*, et al.* (2020a). Targeting Bim via a lncRNA Morrbid Regulates the Survival
of Preleukemic and Leukemic Cells. Cell Rep *31*, 107816.
Cai, Z., Kotzin, J.J., Ramdas, B., Chen, S., Nelanuthala, S., Palam, L.R., Pandey, R., Mali, R.S., Liu, Y.,
Kelley, M.R.*, et al.* (2018). Inhibition of Inflammatory Signaling in Tet2 Mutant Preleukemic Cells
Mitigates Stress-Induced Abnormalities and Clonal Hematopoiesis. Cell Stem Cell *23*, 833-849.e835.
Cai, Z., Lu, X., Zhang, C., Nelanuthala, S., Aguilera, F., Hadley, A., Ramdas, B., Fang, F., Nephew, K.,
Kotzin, J.J.*, et al.* (2021). Hyperglycemia cooperates with Tet2 heterozygosity to induce leukemia driven
by proinflammatory cytokine-induced lncRNA Morrbid. J Clin Invest *131*.
Cai, Z., Zhang, C., Kotzin, J.J., Williams, A., Henao-Mejia, J., and Kapur, R. (2020b). Role of lncRNA
Morrbid in PTPN11(Shp2)E76K-driven juvenile myelomonocytic leukemia. Blood Adv *4*, 3246-3251.
Caiado, F., Kovtonyuk, L.V., Gonullu, N.G., Fullin, J., Boettcher, S., and Manz, M.G. (2023). Aging
drives Tet2+/- clonal hematopoiesis via IL-1 signaling. Blood *141*, 886-903.
Caiado, F., Pietras, E.M., and Manz, M.G. (2021). Inflammation as a regulator of hematopoietic stem
cell function in disease, aging, and clonal selection. J Exp Med *218*.
Chu, S.H., Heiser, D., Li, L., Kaplan, I., Collector, M., Huso, D., Sharkis, S.J., Civin, C., and Small, D.

(2012). FLT3-ITD knockin impairs hematopoietic stem cell quiescence/homeostasis, leading to
myeloproliferative neoplasm. Cell Stem Cell *11*, 346-358.

Cumbo, C., Tarantini, F., Zagaria, A., Anelli, L., Minervini, C.F., Coccaro, N., Tota, G., Impera, L.,
Parciante, E., Conserva, M.R.*, et al.* (2022). Clonal Hematopoiesis at the Crossroads of Inflammatory
Bowel Diseases and Hematological Malignancies: A Biological Link? Front Oncol *12*, 873896.

Delhommeau, F., Dupont, S., Della Valle, V., James, C., Trannoy, S., Massé, A., Kosmider, O., Le
Couedic, J.P., Robert, F., Alberdi, A.*, et al.* (2009). Mutation in TET2 in myeloid cancers. N Engl J Med
360, 2289-2301.

Deshmukh, H.S., Liu, Y., Menkiti, O.R., Mei, J., Dai, N., O'Leary, C.E., Oliver, P.M., Kolls, J.K., Weiser,
J.N., and Worthen, G.S. (2014). The microbiota regulates neutrophil homeostasis and host resistance to
Escherichia coli K1 sepsis in neonatal mice. Nat Med *20*, 524-530.

Fabre, M.A., de Almeida, J.G., Fiorillo, E., Mitchell, E., Damaskou, A., Rak, J., Orrù, V., Marongiu, M.,
Chapman, M.S., Vijayabaskar, M.S.*, et al.* (2022). The longitudinal dynamics and natural history of
clonal haematopoiesis. Nature *606*, 335-342.

Genovese, G., Kähler, A.K., Handsaker, R.E., Lindberg, J., Rose, S.A., Bakhoum, S.F., Chambert, K.,
Mick, E., Neale, B.M., Fromer, M.*, et al.* (2014). Clonal hematopoiesis and blood-cancer risk inferred
from blood DNA sequence. N Engl J Med *371*, 2477-2487.

He, H., Wang, Z., Yu, H., Zhang, G., Wen, Y., and Cai, Z. (2022). Prioritizing risk genes as novel
stratification biomarkers for acute monocytic leukemia by integrative analysis. Discov Oncol *13*, 55.

Heyde, A., Rohde, D., McAlpine, C.S., Zhang, S., Hoyer, F.F., Gerold, J.M., Cheek, D., Iwamoto, Y.,
Schloss, M.J., Vandoorne, K.*, et al.* (2021). Increased stem cell proliferation in atherosclerosis accelerates
clonal hematopoiesis. Cell *184*, 1348-1361.e1322.

Hsu, J.I., Dayaram, T., Tovy, A., De Braekeleer, E., Jeong, M., Wang, F., Zhang, J., Heffernan, T.P., Gera,
S., Kovacs, J.J.*, et al.* (2018). PPM1D Mutations Drive Clonal Hematopoiesis in Response to Cytotoxic
Chemotherapy. Cell Stem Cell *23*, 700-713.e706.

Jaiswal, S., and Ebert, B.L. (2019). Clonal hematopoiesis in human aging and disease. Science *366*.

Jaiswal, S., and Libby, P. (2020). Clonal haematopoiesis: connecting ageing and inflammation in
cardiovascular disease. Nat Rev Cardiol *17*, 137-144.

Jaiswal, S., Natarajan, P., Silver, A.J., Gibson, C.J., Bick, A.G., Shvartz, E., McConkey, M., Gupta, N.,
Gabriel, S., Ardissino, D.*, et al.* (2017). Clonal Hematopoiesis and Risk of Atherosclerotic
Cardiovascular Disease. N Engl J Med *377*, 111-121.

Khan, N., Patel, D., Trivedi, C., Kavani, H., Pernes, T., Medvedeva, E., Lewis, J., Xie, D., and Yang, Y.X.
(2021). Incidence of Acute Myeloid Leukemia and Myelodysplastic Syndrome in Patients With
Inflammatory Bowel Disease and the Impact of Thiopurines on Their Risk. Am J Gastroenterol *116*, 741-
747.

Khoury, J.D., Solary, E., Abla, O., Akkari, Y., Alaggio, R., Apperley, J.F., Bejar, R., Berti, E., Busque, L.,
Chan, J.K.C.*, et al.* (2022). The 5th edition of the World Health Organization Classification of
Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. Leukemia *36*, 1703-1719.

La Manno, G., Soldatov, R., Zeisel, A., Braun, E., Hochgerner, H., Petukhov, V., Lidschreiber, K., Kastriti,
M.E., Lönnerberg, P., Furlan, A.*, et al.* (2018). RNA velocity of single cells. Nature *560*, 494-498.

Li, S., Zhang, P., Chen, W., Ye, L., Brannan, K.W., Le, N.-T., Abe, J.-i., Cooke, J.P., and Wang, G. (2023).
A relay velocity model infers cell-dependent RNA velocity. Nature Biotechnology.

Libby, P., Ridker, P.M., and Maseri, A. (2002). Inflammation and atherosclerosis. Circulation *105*, 1135-
1143.

Meisel, M., Hinterleitner, R., Pacis, A., Chen, L., Earley, Z.M., Mayassi, T., Pierre, J.F., Ernest, J.D.,
Galipeau, H.J., Thuille, N., *et al.* (2018). Microbial signals drive pre-leukaemic myeloproliferation in a
Tet2-deficient host. Nature *557*, 580-584.

Miller, P.G., Qiao, D., Rojas-Quintero, J., Honigberg, M.C., Sperling, A.S., Gibson, C.J., Bick, A.G.,
Niroula, A., McConkey, M.E., Sandoval, B., *et al.* (2022). Association of clonal hematopoiesis with
chronic obstructive pulmonary disease. Blood *139*, 357-368.

Moran-Crusio, K., Reavie, L., Shih, A., Abdel-Wahab, O., Ndiaye-Lobry, D., Lobry, C., Figueroa, M.E.,
Vasanthakumar, A., Patel, J., Zhao, X., *et al.* (2011). Tet2 loss leads to increased hematopoietic stem cell
self-renewal and myeloid transformation. Cancer Cell *20*, 11-24.

Nazha, A., Komrokji, R., Meggendorfer, M., Jia, X., Radakovich, N., Shreve, J., Hilton, C.B., Nagata,
Y., Hamilton, B.K., Mukherjee, S., *et al.* (2021). Personalized Prediction Model to Risk Stratify Patients
With Myelodysplastic Syndromes. J Clin Oncol *39*, 3737-3746.

Rasmussen, K.D., Jia, G., Johansen, J.V., Pedersen, M.T., Rapin, N., Bagger, F.O., Porse, B.T., Bernard,
O.A., Christensen, J., and Helin, K. (2015). Loss of TET2 in hematopoietic cells leads to DNA
hypermethylation of active enhancers and induction of leukemogenesis. Genes Dev *29*, 910-922.

Smillie, C.S., Biton, M., Ordovas-Montanes, J., Sullivan, K.M., Burgin, G., Graham, D.B., Herbst, R.H.,
Rogel, N., Slyper, M., Waldman, J., *et al.* (2019). Intra- and Inter-cellular Rewiring of the Human Colon
during Ulcerative Colitis. Cell *178*, 714-730.e722.

Steensma, D.P., Bejar, R., Jaiswal, S., Lindsley, R.C., Sekeres, M.A., Hasserjian, R.P., and Ebert, B.L.
(2015). Clonal hematopoiesis of indeterminate potential and its distinction from myelodysplastic
syndromes. Blood *126*, 9-16.

Taur, Y., Xavier, J.B., Lipuma, L., Ubeda, C., Goldberg, J., Gobourne, A., Lee, Y.J., Dubin, K.A., Socci,
N.D., Viale, A., *et al.* (2012). Intestinal domination and the risk of bacteremia in patients undergoing
allogeneic hematopoietic stem cell transplantation. Clin Infect Dis *55*, 905-914.

Weeks, L.D., Niroula, A., Neuberg, D.S., Wong, W.J., Lindsley, R.C., Luskin, M.R., Berliner, N., Stone,
R.M., DeAngelo, D.J., Soiffer, R.J., *et al.* (2022). Prediction of Risk for Myeloid Malignancy in Clonal
Hematopoiesis. Blood *140*, 2229-2231.

Weinstock, J.S., Gopakumar, J., Burugula, B.B., Uddin, M.M., Jahn, N., Belk, J.A., Bouzid, H., Daniel,
B., Miao, Z., Ly, N., *et al.* (2023). Aberrant activation of TCL1A promotes stem cell expansion in clonal
haematopoiesis. Nature *616*, 755-763.

Wong, T.N., Ramsingh, G., Young, A.L., Miller, C.A., Touma, W., Welch, J.S., Lamprecht, T.L., Shen,
D., Hundal, J., Fulton, R.S., *et al.* (2015). Role of TP53 mutations in the origin and evolution of therapy-
related acute myeloid leukaemia. Nature *518*, 552-555.

Yan, H., Baldridge, M.T., and King, K.Y. (2018). Hematopoiesis and the bacterial microbiome. Blood
132, 559-564.

Zeng, H., He, H., Guo, L., Li, J., Lee, M., Han, W., Guzman, A.G., Zang, S., Zhou, Y., Zhang, X., *et al.*
(2019). Antibiotic treatment ameliorates Ten-eleven translocation 2 (TET2) loss-of-function associated
hematological malignancies. Cancer Lett *467*, 1-8.

Zeng, X., Li, X., Li, X., Wei, C., Shi, C., Hu, K., Kong, D., Luo, Q., Xu, Y., Shan, W., *et al.* (2023). Fecal
microbiota transplantation from young mice rejuvenates aged hematopoietic stem cells by suppressing
inflammation. Blood *141*, 1691-1707.

Zhang, C.R.C., Nix, D., Gregory, M., Ciorba, M.A., Ostrander, E.L., Newberry, R.D., Spencer, D.H., and
Challen, G.A. (2019). Inflammatory cytokines promote clonal hematopoiesis with specific mutations in
ulcerative colitis patients. Exp Hematol *80*, 36-41.e33.

Zhang, Q., Zhao, K., Shen, Q., Han, Y., Gu, Y., Li, X., Zhao, D., Liu, Y., Wang, C., Zhang, X*., et al.*
(2015). Tet2 is required to resolve inflammation by recruiting Hdac2 to specifically repress IL-6. Nature
525, 389-393.
886

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6

Figure 7.

Supplemental Figure 1

Supplementary Figure 2

Supplemental Figure 3

Supplemental Figure 4

A expression plots for *Gata1*, *Pf4*, *Gata2*, *Pax5*, *Il1b*, *Il6*

B *Tet2*mut_Veh vs WT_Veh and *Tet2*mut_DSS vs WT_DSS (average logFC vs Cell types)

C Violin plots of Egr1, Socs3, Junb, Lyz2, S100a8, S100a9 expression in HSC

D GO analysis of upregulated of DEGs of GMP_1 in *Tet2*mut_DSS and GO analysis of upregulated of DEGs of HSC in *Tet2*mut_DSS

Supplemental Figure 5

Supplementary Figure 6

Supplementary Figure 7

A *Ptx3* expression in Colon from scRNA-seq data

B *Ptx3* expression in stromal cells from colon

C

D

E

WT-Veh *Tet2*⁺/⁻-Veh WT-DSS *Tet2*⁺/⁻-DSS

F

G 1:5 mix of BMMC before transplantation

H

I

J

K