Linking genomic prediction for muscle fat content in

Atlantic salmon to underlying changes in lipid metabolism

regulation

Thomas N. Harvey[1*], Hanne Dvergedal[1], Lars Grønvold[1], Yang Jin[1], Jørgen Ødegård[1,2], Sven

Arild Korsvoll[2], Tim Knutsen[2], Torgeir R. Hvidsten[3], Simen R. Sandve[1]

[1]Department of Animal and Aquacultural Sciences, Faculty of Biosciences, Norwegian

University of Life Sciences, P. O. Box 5003, NO-1433 Ås, Norway

[2]AquaGen AS, P. O. Box 1240, NO-7462 Trondheim, Norway

[3]Faculty of Chemistry, Biotechnology and Food Sciences, Norwegian University of Life

Sciences, NO-1432 Ås, Norway

[*]Corresponding author

E-mail addresses:

TH: thomas.n.harvey@nmbu.no

Abstract

Muscle fat content is an important production trait in Atlantic salmon (*Salmo salar*) because it influences the flavor, texture, and nutritional properties of the fillet. Genomic selection can be applied to alter muscle fat content, however how such selection changes the underlying molecular physiology of these animals is unknown. Here, we examine the link between genomic prediction and underlying molecular physiology by correlating genomic breeding values for fat content to liver gene expression in 184 fish. We found that Salmon with higher genomic breeding values had higher expression of genes in lipid metabolism pathways. This included key lipid metabolism genes *hmgcrab*, *fasn-b*, *fads2d5*, and *fads2d6*, and lipid transporters *fatp2f*, *fabp7b*, and *apobc*. We also found several regulators of lipid metabolism with negative correlation to genomic breeding vales, including *pparg-b*, *fxr-a*, and *fxr-b*. A quantitative trait loci analysis for variation in gene expression levels (eQTLs) for 167 trait associated genes found that 71 genes had at least one eQTL, and that most were trans eQTLs. Closer examination revealed distinct eQTL clustering on chromosomes 3 and 6, indicating the presence of putative common regulator in these regions. Taken together, these results suggest that increased fat content in high genomic breeding value salmon is associated with elevated lipid synthesis, elevated lipid transport, and reduced glycerolipid breakdown; and that this is at least partly achieved by selection on genetic variants that impact the function of top-level transcription factors involved in liver metabolism. Our study sheds light on how genomic selection alters lipid content in Atlantic salmon, and the results could be used to prioritize SNPs to improve the efficiency of genomic selection in the future.

Keywords: Atlantic salmon, genomic selection, lipid metabolism, fat content, gene expression, eQTL

1   Background

Intramuscular fat content is an important quality parameter in most production animals because it influences the texture, flavor, and nutritional properties of the meat. The latter is especially important in Atlantic salmon as it is considered an excellent source of healthy omega-3 fatty acids in human diets and higher fat equates to higher levels of omega-3s reaching the end consumer (Tocher, 2015). Lipid content is a highly polygenic trait, with many genes explaining a small fraction of the total genetic variation (Pena et al., 2016). In Atlantic salmon, lipid content has been reported to have a heritability of 0.18 to 0.2 (Kristjánsson et al., 2020; Tsai et al., 2015), so breeding for fat content in Atlantic salmon is possible.

Breeding for complex traits such as growth rate, disease resistance, and filet properties has been accelerated in recent years with the widespread adoption of genomic selection. This breeding strategy takes advantage of genome wide single nucleotide polymorphism (SNP) data to calculate genomic breeding values (GBV) for each individual based on the genotype and phenotype of their parents, enabling rational selection of breeding pairs (Meuwissen et al., 2001). Genomic selection in salmon has already been successfully applied; with breeding programs improving lice resistance and filet color faster and more reliably than traditional breeding programs (Ødegård et al., 2014). Genomic selection for fat content has been successfully applied to rainbow trout (Hu et al., 2020), however such methods have yet to be applied to fillet fat in Atlantic salmon.

Application of genomic selection to production traits traditionally does not consider the biology of the target trait, and rather weighs all SNPs equally to predict genomic effects (Meuwissen et al., 2001). However, biological knowledge about the trait of interest can be used to improve the power and accuracy of genomic predictions (de las Heras-Saldana et al., 2020; MacLeod et al., 2016). For this reason, it is useful to understand the underlying changes in molecular

physiology occurring during genomic selection to prioritize SNPs for future rounds of genomic

selection. In Atlantic salmon, lipid homeostasis is achieved by balancing dietary intake from

the gut, *de-novo* synthesis in the body, and excretion through gut or biliary systems, which

involves the coordinated action of thousands of genes. Liver plays a central role in lipid

metabolism, controlling the flow of dietary lipids between different parts of the body through

absorption and secretion of different lipoproteins to the circulatory system (Vance and Vance,

2008). Liver is therefore a logical place to analyze the effect of genomic selection for fat

content on the molecular physiology of the fish.

In this study, we calculate GBVs for filet fat content in 184 fish based on genotype and

phenotype data from a training set of 2487 fish. We sequenced the liver transcriptomes of all

184 fish and associate gene expression with GBV. Finally, we took the most significant GBV

associated genes (padj $< 10^{-4}$) together with a manually curated subset of known lipid related

GBV associated genes and identify eQTLs for each (Figure 1a). Our aim is to improve our

understanding of the genetic basis for differences in fat content by interrogating the link

between GBVs and gene regulation in the liver.

2   Methods

2.1   Fish and housing

A family experiment with Atlantic salmon was carried out at the fish laboratory, Norwegian

University of Life Sciences (NMBU), Aas, Norway. The family experiment is explained in

detail by Dvergedal *et al.* (Dvergedal et al., 2019). Broodstock from AquaGen's breeding

population (22 males and 23 females) were used to generate 23 families.

From the eyed egg stage until the start of the experiment, all families were communally reared

in a single tank until the start of the experiment. When the fish reached 5-10 g, they were pit-

tagged with a 2 x 12 mm unique glass tag (RFID Solutions, Hafrsfjord, Norway), and a fin-clip

was collected for DNA-extraction and genotyping of a total of 2,300 fish. Fin clips (20 mg)

were incubated in lysis buffer and treated with proteinase K (20 μg/ml) at 56 ℃ overnight. The

following day, DNA was isolated from the lysate using the sbeadex livestock kit (LGC

Genomics) according to the manufacturer's protocol (Thermo Fisher Scientific) at Biobank AS

(Hamar, Norway). The DNA concentration was measured using a Nanodrop 8000 (Thermo

Fisher Scientific). All fish were genotyped using AquaGen's custom Axiom®SNP (single-

nucleotide polymorphism) genotyping array from Thermo Fisher Scientific (former Affymetrix)

(San Diego, CA, USA). This SNP-chip contains 56,177 SNPs which were originally identified

based on Illumina HiSeq reads (10-15x coverage) from 29 individuals from AquaGen's

breeding population. Genotyping was done at CIGENE (Aas, Norway). Genotypes were called

from the raw data using the Axiom Power Tools software from Affymetrix. Individuals having

a Dish-QC score below 0.82, and/or a call-rate below 0.97 were deleted from further analyses.

A priori to the 12-day test, the parentage of each fish was established using genomic

relationship likelihood for parentage assignment (Grashei et al., 2018), and families were

allocated to tanks, 50 fish per tank, and 2 tanks per family. Except for nine tanks in which the

number of fish varied between 42 and 54, due to some mortality before the start of the

experiment or an increased number due to a counting mistake. The total number of fish was

2,281 and families were fed a fishmeal-based diet, as described in Dvergedal *et al.* (Dvergedal

et al., 2019). The phenotypic data were registered individually for relative growth, as described

by Dvergedal *et al.* (Dvergedal et al., 2019).

2.2   Estimating genomic breeding values (GBV) for fat content

Estimated genomic breeding values (GBV) for fillet fat content were predicted using a training

data set consisting of 2487 genotyped (50-70k SNP chip) and phenotyped (fillet fat content)

fish from two slaughter tests performed on the parental year-class 2014 (766 fish of the parental

generation of the fish in the current study) and 2017 (1721 fish of the same generation as in the

current study). Phenotypes for fat content were obtained using Norwegian quality cuts (NQC)

from each fish that were subsequently frozen. The cut was thawed, skin and central bone

removed, and homogenized for fat measurement on a NIR XDS machine. Fat content was

predicted using a proprietary NQC model owned by Cargill. Average fat (standard deviation)

was 12.96% (1.22%) and 17.41% (2.84%) for, respectively, year-classes 2014 and 2017.

Additionally, 59 samples were analyzed by an independent lab (Eurofins) for validation. A

linear genomic animal model (GBLUP) was used to obtain EBV:

121
$$\mathbf{y_{fat}} = \mathbf{X\mu} + \mathbf{Zg} + \mathbf{e}$$

Where $\mathbf{y_{fat}}$ is a vector of fat content phenotypes (standardized within each year-class), $\mathbf{\mu}$ is a

vector of the two year-class intercepts, $\mathbf{g} \sim N(\mathbf{0}, \mathbf{G}\sigma_g^2)$ is a vector of polygenic effects (including

both 2014 and 2017 year-class fish in the current study), $\mathbf{e}$ is a vector of random residuals, σ_g^2

is the additive genetic variance, $\mathbf{G} = \rho\mathbf{MM}'$ is the genomic relationship matrix, $\mathbf{M}$ is a centered

genotype matrix (one row per individual and one column per locus), $\rho = \frac{1}{2\sum p_i(1-p_i)}$ and p_i is

the allele frequency at locus i. The estimated genomic heritability (across the two year-classes)

for fillet fat content was rather moderate (0.23 ± 0.03). SNP marker effects estimates were

obtained as: $\hat{\mathbf{m}} = \rho^{-1}\mathbf{M}'\mathbf{G}^{-1}\hat{\mathbf{g}}$ (Legarra et al., 2018). Using the estimated marker effects GBVs

for the fish in the current study were predicted as:

131
$$\hat{\mathbf{g}}_{sample} = M_{sample}\hat{\mathbf{m}}$$

Where $\hat{\mathbf{g}}_{sample}$ is a vector of GBVs (on the standardized scale) for fat content of fish in the

current study and M_{sample} is a (centered) genotype matrix of the same fish.

2.3   RNA extraction and transcriptomic sequencing

Four individual fish from each family were used for RNA isolation. RNA was extracted from

liver of each individual fish using RNeasy Plus Universal Kit (Qiagen, Hilden, Germany),

according to the manufacturer's instructions. The concentration of RNA was determined by a

Nanodrop 8000 (Thermo Fisher Scientific, Waltham, USA), and RNA integrity was examined

by using a 2100 Bioanalyzer (Agilent Technologies, Santa Clara, USA). All RNA samples had

RNA integrity (RIN) values higher than 8. Sequencing libraries were generated using TruSeq

Stranded mRNA Library Prep Kit (Illumina, San Diego, USA) according to the manufacturer's

protocol. Libraries were delivered to Norwegian Sequencing Centre (Oslo, Norway) where all

184 samples were merged into one flow cell and sequenced using 100bp single-end mRNA

sequencing (RNA-seq) on Illumina Hiseq 2500 (Illumina, San Diego, CA, USA).

Raw fastq file of reads sequences are publicly available on ArrayExpress under accession

number E-MTAB-8305. Gene expression was quantified using the Salmon quasi-mapper

version 0.13.1 (Patro et al., 2017) against the Atlantic salmon transcriptome (ICSASG_v2).

2.4   Trait association analysis

Trait associated genes (TAGs) were detected by correlating gene expression to GBV using

edgeR (Robinson et al., 2009). Genes with low read counts, i.e. less than 0.5 count per million

(CPM) in 50% of the samples, were removed and GBV was scaled (mean = 0 and standard

deviation =1) prior to analysis. The linear regression model was

$$Y = \beta_0 + \beta_1 x + \epsilon$$

where Y is gene expression and x is scaled GBV value. Genes were classified as TAGs if the

slope (β_1) was significant, i.e. $\beta_1 \neq 0$ with false discover rate adjusted p value < 0.05. We also

ran the analysis with family as a fixed effect factor in the model, however this only slightly

changed the number of TAGs and did not influence the conclusion, so we use the simpler model

in the analysis. KEGG enrichment analysis of the TAGs with positive and negative slope was

performed with the kegga function in the limma R package (Ritchie et al., 2015). Translations

between human readable gene names and NCBI RefSeq annotation gene ID's can be found in

File S1.

2.5   Genome-wide association analysis

To associate variation in TAGs with host genetics, a genome-wide association study was done

using TAGs with an adjusted p-value <0.0001 ($n = 121$) and genes known to be associated with

fat metabolism in the liver ($n = 46$) as response variables. The analysis was carried out by a

linear mixed-model algorithm implemented in a genome-wide complex trait analysis (GCTA)

(Yang et al., 2011). The leave one chromosome out option (--mlm-loco) was used, meaning

that the chromosome harboring the SNP tested for was left out when building the genetic

relationship matrix (GRM). The linear mixed model can be written as:

$$Y_i = a + bx + g_i^- + \varepsilon_i,$$

where Y_i is one of the TAGs of individual i, a is the intercept, b is the fixed regression of the

candidate SNP to be tested for association, x is the SNP genotype indicator variable coded as

0, 1, or 2, g_i^- i is the random polygenic effect for individual $i \sim N(0, G\sigma_g^2)$ where **G** is

the GRM and σ_g^2 is the variance component for the polygenic effect, and ε_i is the random

residual. In this algorithm, σ_g^2 is re-estimated each time a chromosome is left out from the

calculation of the GRM. The dataset was filtered, and individuals with < 10% missing

genotypes were kept ($n = 2279$). Further, it was required that SNPs should have minor allele

frequency (MAF) $\geq$ 1% and a call rate > 90%. After filtering, 54,200 SNPs could be included

in the analysis. The level of significance for SNP was evaluated with a built-in likelihood-ratio

test, and the threshold value for genome-wide significance was calculated using the Bonferroni

correction (0.05/ 54200 = 9.23 × 10−7), corresponding to a -log10 *p*-value of 6.03.

2.6　　Allele distribution of top SNPs for TAGs associated with lipid metabolism in liver

For TAGs with a significant genome-wide eQTL association to lipid metabolism in liver, the

allele distribution for the top SNP (the SNP with the highest -log10 *p*-value) was examined

with PLINK 2.00 (Chang et al., 2015) using the --recode option, which creates a new file after

applying sample/variant filters and the --extract option to create a file with the top SNP of

interest.

2.7　　Co-expression network analysis

We assembled an independent RNA-seq expression data set comprising 112 liver samples

spanning different diets and life stages in fresh water (Gillard et al., 2018). Raw RNA-Seq data

can be found in the European Nucleotide Archive (ENA) under project accession no.

PRJEB24480.　　Read　　counts　　for　　Atlantic　　salmon　　genes　　(NCBI:

GCF_000233375.1_ICSASG_v2) were estimated using Salmon (Patro et al., 2017). Code and

data available at https://gitlab.com/garethgillard/megaLiverRNA. 27,786 genes with at least 10

mapped reads in more than 10 samples were retained for further analysis. Read counts were

normalized using the varianceStabilizingTransformation-function from the R package DESeq2

(Love et al., 2014).

For co-expression network inference, we used the Weighted Gene Co-expression Network

Analysis (WGCNA) R package (Langfelder and Horvath, 2008) and the function

blockwiseModules with the bicor correlation measure and parameters *power = 3 (scale free*

topology fit with an R2 of 0.8), maxBlockSize = 30000, networkType = "signed", TOMType =

"signed", corType = "bicor", maxPOutliers = 0.05, replaceMissingAdjacencies = TRUE,

pamStage = F, deepSplit = 1, minModuleSize = 5, minKMEtoStay = 0.2, minCoreKME = 0.2,

minCoreKMESize = 2, reassignThreshold = 0 and mergeCutHeight = 0.2. A hard correlation

threshold of 0.5 was used to visualize the network in Cytoscape (https://cytoscape.org).

3 Results

3.1 Correlations between liver gene expression levels and GBV

We first calculated GBV for the 184 fish used in this study based on a training set of 2487 fish

(Figure 1a and b). RNA sequencing and trait association analysis revealed a total of 710 TAGs

(padj < 0.05) positively correlated to GBV and 653 TAGs negatively correlated to GBV (Figure

1c, File S2). By comparing the number of TAGs to total genes within each KEGG pathway,

we found that the positive TAGs were enriched (p < 0.05) in 45 KEGG pathways (Figure 1d,

File S3). Many of the most significantly enriched KEGG pathways were related to lipid

metabolism including "*PPAR signaling*", "*fatty acid degradation", "fatty acid biosynthesis*",

"*biosynthesis of unsaturated fatty acids*" and "*glycerolipid metabolism*" (Figure 1d). In

addition, pathways related to amino acid metabolism and energy metabolism were enriched for

positively correlated TAGs (Figure 1d). Only eight pathways were found enriched (p < 0.05)

for the negatively correlated TAGs. This included the lipid metabolism pathway *"primary bile*

acid synthesis" which produces bile acids from cholesterol that aid in lipid solubilization in the

intestine.

Figure 1: Correlation of GBV to gene expression. A) Flow chart of our analysis. B) Distribution of GBVs in the 184 salmon used for RNAseq analysis. C) Slope distribution among positively and negatively correlated TAGs. D) KEGG enrichment analysis of positively and negatively correlated TAGs. Only significantly enriched pathways (p < 0.05) are shown.

3.2 Regulation of lipid metabolism genes

To further dissect the link between lipid metabolism and GBV for fat content, we performed an in-depth analyses of a manually annotated set of genes involved in lipid metabolism in salmon (Gillard et al., 2018). We found that the TAG list was clearly enriched for lipid genes (fisher's p-value 2.2e-16, odds ratio 4.92) with 45 (6.3%) lipid genes found in the 710 TAGs

positively correlated to GBV, and 10 lipid genes (1.5%) found in the 653 TAGs negatively

correlated with GBV (Figure 2a). Intriguingly, the GBV-associated lipid genes covered genes

involved in many enzymatic steps in the cholesterol biosynthesis pathway including the

enzyme controlling the rate limiting step of cholesterol biosynthesis, *hmgcrab*. Other lipid

metabolism enzyme-encoding genes associated to fat content GBV included fatty acid synthase

(*fasn-b),* all three polyunsaturated fatty acid desaturases *(fads2d5 fads2d6a, and fads2d6b)*

(figure 2e), and several lipid and fatty acid transporters (*fatp2f*, *fabp7b*, *ldlrab-a*, and *apobc)*

(Figure 2c).

We also found several lipid regulatory genes positively correlated with GBV (Figure 2),

including all three isoforms of SREPB1 (*srebp1b, srebp1c, srebp1d)* (Figure 2f), known

regulators of lipid biosynthesis (Shimano and Sato, 2017). Lipid metabolism genes negatively

correlated to GBV included the important lipid oxidation regulator *pparg-b* (figure 2b) as well

as both copies of farnesoid x receptor (*fxr-a* and *fxr-b*), which plays a key role in hepatic

triglyceride homeostasis and is involved in suppression of lipogenesis (Jiao et al., 2015). The

lipid gene that was most negatively associated with GBV was monoacylglycerol lipase (*mgll*)

which is involved glycerolipid breakdown.

Figure 2: Correlation between GBV and lipid metabolism gene expression. A) Volcano plot of the regression results between gene expression and GBV. Genes involved in lipid metabolism are coloured and important lipid metabolism genes are labelled. Size corresponds to mean log2TPM values. The dashed red line indicates the padj < 0.05 cutoff. Correlation of GBV and gene expression are shown on the right for *pparg-b* (B), *fatp2f* (C), and *mgll* (D), *fads2d6a* (E), and *srebp1d* (F).

3.3   eQTL analyses highlights several trans-acting loci impacting many genes

To explore the genetic architecture of gene expression differences between fish with high and low GBV, we used linear regressions to identify genetic variants associated gene expression levels (File S4). In total 167 genes associated with GBV fat were examined, which included the top 121 significant TAGs (padj < 0.0001), as well as 46 significant TAGs from lipid pathways (padj < 0.05) (Figure 2a). Genes that were not annotated on a chromosome (i.e belonged to a smaller unplaced scaffold) were discarded from this analysis.

In total, 71 genes had at least one significant eQTL (genome wide significance level, $p < 10^{-6}$), with a mean number of 1.4 eQTLs per gene (Figure 3a). Dissecting the positions of eQTL

signals relative to these genes showed that 21 genes had a significant association in cis (no more than 10 Mbp from the gene). Considering only "top associations" for each gene reveals a clear tendency for eQTL-associations to variants on other chromosomes (i.e. trans associations) (Figure 3b). Seven of the genes with significant eQTL's were known lipid metabolism genes; two of which are monoglyceride lipase (*mgll*) and fatty acid desaturase 2-like (*fads2d6b*) (Figure 3c-f).

Figure 3: GWAS of top GBV TAGs and selected lipid metabolism genes. A) Number of significant (p < 10^{-6}) eQTLs per gene. B) Number of genes with the top eQTL in cis (< 10 MB from gene), trans (different chromosome), or trans (same chromosome). C) Manhattan plot of SNPs associated with *fads2d6b* gene expression. Significance cutoffs are indicated by red (strict) and blue (relaxed) lines. D) Genotype distribution of the top SNP for *fads2d6b*. E and F) Same as C and D for the gene *mgll*.

Since the genes tested for eQTLs were associated with differences in one quite specific

molecular trait (i.e. lipid metabolism), it is plausible that a few top regulators of key lipid-

metabolism pathways could impact the expression of many of the genes in our study. The high

numbers of trans eQTL signals (Figure 3b) supports this idea. Hence, to test if any

chromosomal regions showed sign of harboring such major regulators, we analyzed the

distribution of top trans-eQTL associations across chromosomes. Using a relaxed p-value

cutoff for significant associations ($p < 10^{-4}$), two chromosomes (3 and 6) were clearly enriched

for top trans-eQTLs relative to the total number of SNPs on each chromosome (Figure 4a). It

is worth noting, that the enrichment signal on chromosome 6 dropped rapidly as we increased

p-value cutoff stringency and was diminished at $p < 10^{-6}$.

Next, we performed an in-depth analysis of the eQTL signals on chromosome 3 and 6, with

specific focus on lipid-metabolism genes. We hypothesized that if regulators of lipid-

metabolism pathways were located on these chromosomes we would find clusters of trans-

eQTLs for genes that are co-regulated in liver. We therefore first used a large gene expression

dataset of 112 liver samples to estimate co-expression modules (i.e. genes whereby their

expression correlates across different samples) (Figure 4b). Genes within these co-expression

modules are predicted to share transcriptional regulators. We then associated genes with trans-

eQTLs on chromsomes 6 and 3 to specific co-expression modules. Even though the trans-eQTL

enrichment on chromosome 3 contained many more highly significant associations compared

to chromosome 6 (Figure S1), there was no obvious clustering of trans-eQTL signals among

co-expressed genes on this chromosome. However, the central region (35-50 Mbp) of

chromosome 6 displayed a clear clustering of trans-eQTL signals to co-expressed genes.

Although these genes belonged to three different co-expression modules, these three modules

were virtually identical in the co-expression network (Figure 4b). Interestingly, this genomic

region harbors two copies of srebp-2-like genes, known to regulate various aspects of lipid-

metabolism in vertebrates, which also belong to one of these three co-expression modules.

Figure 4: Trans eQTL and gene co-expression. A) Relationship between total number of

SNPs for each chromosome and the number of genes having top trans-eQTL signals on each

chromosome. B) Co-expression network from liver RNAseq data highlighting three co-

expression modules which contain genes with top trans eQTL signals ($p < 10^{-4}$) on chromosome

6. C) Circos plot showing the top trans-eQTL links between trait associated genes and

chromosome 6. Colors reflect the co-expression module that the trait associated genes belong

to. We have indicated the position of a major lipid metabolism transcription factor (srebp2) on

chromosome 6.

4   Discussion

Our results clearly demonstrate that genomic selection for high muscle fat content in salmon

drives changes in expression of genes involved in lipid metabolism in the liver. Variation in

muscle lipid content among fish could be due to differences in 1) uptake of lipids from the diet,

2) *de-novo* lipogenesis in the body, 3) lipid transport and deposition between different part of

the body, 4) lipid degradation by beta-oxidation and efflux through bile synthesis, or 5) a

combination. Our results have shown that GBV was positively associated with gene expression

in lipid transport (*fatp2f*, *fabp7b*), *de novo* lipogenesis (*fas*), fatty acid desaturation (*fads2d5*,

fads2d6a, *fads2d6b*, and *fads2d6c*), cholesterol biosynthesis (*hmgcrab*), and negatively

associated with genes in glycerolipid breakdown (*mgll*), beta-oxidation (*pparg-b*) and bile acid

synthesis (*fxr*) in liver. This suggests that higher deposition of lipids in high GBV salmon was

most likely due to a combination of reduced glycerolipid breakdown, elevated lipid synthesis,

and elevated lipid transport. Both endogenous and exogeneous lipids in liver are packaged into

very-low-density lipoproteins (VLDL) which are secreted into the vasculature. Lipids in VLDL

are taken up by the peripheral tissues such as muscle, and the leftover lipoproteins (low-density

lipoproteins, LDL) are taken back by liver through the LDL receptor (LDLR) (Wang, 2007).

Most of the lipoproteins including VLDL and LDL contain a copy of apolipoprotein B (APOB),

an essential component for its structure (Elovson et al., 1988). Our study has found GBV to be

positively correlated to *ldlrab-a*, and *apobc*. This could suggest high GBV was associated with

increased amount and turnover of lipoproteins in the bloodstream. Additionally, we have also

identified two *fxr* genes which were negatively correelated to GBV. Since fxr is a key regulator

and positively correlated to bile salt synthesis in fish (Wen et al., 2021), this suggests that high

GBV fish has decreased lipid excretion through bile salt production pathway. Under this

scenario, a higher proportion of newly synthesized saturated and monounsaturated fatty acid,

and diet derived polyunsaturated fatty acid containing triacylglycerol are exported as

lipoproteins from the liver and deposited in peripheral tissues including muscle where the fat

phenotype was measured.

We find expression of key genes in lipid biosynthesis pathways, including *fas*, *fads2d5*,

fads2d6a, *fads2d6b*, and *fads2d6c* were positively correlated to GBV. This is likely due to

higher expression of sterol regulatory element-binding protein 1 (SREBP1), which is the key

positive regulator of fatty acid *de-novo* synthesis and LC-PUFA synthesis in salmon (Carmona-

Antoñanzas et al., 2014; Minghetti et al., 2011). Other genes of LC-PUFA synthesis, the *elovl2*

and *elovl5* genes, were not correlated to GBV. This could be because these genes are not

controlled by SREBP1 transcription factor in salmon (Datsomor et al., 2019). A similar study

using high and low muscle fat lines of rainbow trout also found higher expression of lipogenic

genes in high fat lines and hypothesized that this was due to a more active target of rapamycin

(TOR) signalling pathway (Skiba-Cassy et al., 2009). Fat and lean rainbow trout lines displayed

a similar ratio of phosphorylated and native TOR, however fat lines had significantly higher

levels of TOR protein. We did not find TOR mRNA to be associated with GBV of muscle lipid

levels in Atlantic salmon, however TOR protein abundance could be regulated at the

posttranslational level. In addition, we found that ATP-citrate lyase (*acyl*, geneID:106589258)

gene expression was positively associated to GBV, which agrees with the study in rainbow

trout (Skiba-Cassy et al., 2009). ACYL acts as a metabolic switch linking glucose and lipid

metabolism that diverts citrate from the TCA cycle into lipogenesis by converting it to acetyl-

CoA and oxaloacetate in the cytosol. This enables elevated lipogenesis by increasing the

available pool of acetyl-CoA to be converted to fatty acids by FAS.

Our eQTL analysis revealed an unexpectedly high number of trans-eQTLs on chromosome 6

associated to co-regulated genes that were TAGs in our analysis. Moreover, these chromosome

trans-eQTL associations were mostly originating from a smaller region around 50 Mbp,

pointing to a potential common transcription factor. Two of the genes in this region are paralogs

of SREBP2-like, a known regulator of lipid metabolism. Srebp2 is a key transcriptional

regulator controlling cholesterol metabolism in fish and mammals (Carmona-Antoñanzas et al.,

2014; Madison, 2016), and our study has found many positively correlated TAGs involved in

de-novo cholesterol synthesis including *hmgcrab*. Additionally, positively correlated *acyl*

suggests an increased acetyl-CoA pool. Although we did not find *srebp2* to be associated with

GBV in our analysis, SREBP is known to be highly regulated post-transcriptionally through

interactions with SREBP cleavage-activating protein (SCAP). SCAP forms a complex with

SREBP and facilitates the cleavage of SREBPs by site-1 protease, thereby releasing active

NH2-terminal fragments from the ER membrane to nucleus, activating gene expression

(Nohturfft et al., 1998). Therefore, this lack of association could be explained by variation in

srebp2 protein structure resulting in increasing SCAP or Site-1 protease binding activity

without influencing *srebp2* expression itself. Alternatively, it may be that the trans-eQTL

signal cluster on chromosome 6 is driven by another, so far unknown regulator of lipid

metabolism on chromosome 6.

Although lipid synthesis and transport in the liver contributes to lipid content in the muscle,

our results only tell part of the story. Since there is considerable variation in fat deposition and

turnover in salmon (Dvergedal et al., 2020; Dvergedal et al., 2019) variation in the regulation

of lipid metabolism in the muscle must also be a large contributor to the high fat phenotype.

High muscle fat has previously been associated with downregulation of genes related to lipid

catabolism and upregulation of genes associated to glycogenolysis (Horn et al., 2019), which

may signal a transition in how fish utilize energy stores. Additionally, differences in hormonal

signalling between the brain, adipose, and muscle tissues could contribute to the high fat

phenotype. To further improve our understanding of what makes a salmon fat, future studies

need to address these aspects of the salmon molecular physiology.

5 Conclusion

We demonstrate that genomic selection using estimated breeding values for fat content drives

changes in lipid metabolism in Atlantic salmon. Fish with high GBV for muscle fat content

had overall higher gene expression in lipid biosynthesis and transport pathways and lower

expression of genes involved in glycerolipid breakdown. This is important validation for

genomic selection as a strategy to improve lipid content and the results could be used to

prioritize SNPs in future estimates of breeding values.

Ethics approval and consent to participate

The experiment used phenotypic data, which were collected from a family experiment with Atlantic salmon carried out at the fish laboratory, Norwegian University of Life Sciences (NMBU), Aas, Norway, following the laws and regulations for experiments on live animals in EU (Directive 2010/637EU) and Norway (FOR-2015-06-18-761). The experiment was approved by the Norwegian Food Safety Authority (FOTS ID 11676).

Data availability

The genotypic data are owned by AquaGen AS, used under license for this study, and not publicly available. Phenotypic data can be made available on request. All RNAseq data is publicly available on ArrayExpress under accession number E-MTAB-8305.

Declaration of competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding

This study was supported by The Norwegian University of Life Sciences, AquaGen AS, Foods of Norway (a Centre for Research-based Innovation in the Research Council of Norway; grant no. 237841/O30), and GenoSysFat (the Research Council of Norway Havbruk; grant no. 244164/E40).

Acknowledgements

We thank Bjørn Reidar Hansen, Harald Støkken and Bjørn Frode Eriksen for help and assistance at the fish laboratory, and Ricardo Tavares Benicio, Ragnhild Ånestad, Milena Bjelanovic, Jon Øvrum Hansen, Mathabela Nelson, Shomorin Oluwaseun George and Ingrid

415   Marie Håkenåsen for their help during feed production, experiment, and sampling. A thank

416   goes to all that contributed during sampling of experimental material.

References

Carmona-Antoñanzas, G., Tocher, D.R., Martinez-Rubio, L., Leaver, M.J., 2014. Conservation
of lipid metabolic gene transcriptional regulatory networks in fish and mammals. Gene 534(1),
1-9. https://doi.org/https://doi.org/10.1016/j.gene.2013.10.040

Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M., Lee, J.J., 2015. Second-
generation PLINK: rising to the challenge of larger and richer datasets. GigaScience 4(1).
https://doi.org/10.1186/s13742-015-0047-8

Datsomor, A.K., Zic, N., Li, K., Olsen, R.E., Jin, Y., Vik, J.O., Edvardsen, R.B., Grammes, F.,
Wargelius, A., Winge, P., 2019. CRISPR/Cas9-mediated ablation of elovl2 in Atlantic salmon
(Salmo salar L.) inhibits elongation of polyunsaturated fatty acids and induces Srebp-1 and
target genes. Scientific Reports 9(1). https://doi.org/10.1038/s41598-019-43862-8

de las Heras-Saldana, S., Lopez, B.I., Moghaddar, N., Park, W., Park, J.-e., Chung, K.Y., Lim,
D., Lee, S.H., Shin, D., van der Werf, J.H.J., 2020. Use of gene expression and whole-genome
sequence information to improve the accuracy of genomic prediction for carcass traits in
Hanwoo cattle. Genet. Sel. Evol. 52(1), 54. https://doi.org/10.1186/s12711-020-00574-2

Dvergedal, H., Harvey, T.N., Jin, Y., Ødegård, J., Grønvold, L., Sandve, S.R., Våge, D.I., Moen,
T., Klemetsdal, G., 2020. Genomic regions and signaling pathways associated with indicator
traits for feed efficiency in juvenile Atlantic salmon (Salmo salar). Genet. Sel. Evol. 52(1), 66.
https://doi.org/10.1186/s12711-020-00587-x

Dvergedal, H., Ødegård, J., Øverland, M., Mydland, L.T., Klemetsdal, G., 2019. Selection for
feed efficiency in Atlantic salmon using individual indicator traits based on stable isotope
profiling. Genet. Sel. Evol. 51(1), 13. https://doi.org/10.1186/s12711-019-0455-9

Elovson, J., Chatterton, J.E., Bell, G.T., Schumaker, V.N., Reuben, M.A., Puppione, D.L.,
Reeve, J.R., Young, N.L., 1988. Plasma very low density lipoproteins contain a single molecule
of apolipoprotein B. J. Lipid Res. 29(11), 1461-1473.
https://doi.org/https://doi.org/10.1016/S0022-2275(20)38425-X

Gillard, G., Harvey, T.N., Gjuvsland, A., Jin, Y., Thomassen, M., Lien, S., Leaver, M.,
Torgersen, J.S., Hvidsten, T.R., Vik, J.O., Sandve, S.R., 2018. Life-stage-associated
remodelling of lipid metabolism regulation in Atlantic salmon. Mol. Ecol. 27(5), 1200-1213.
https://doi.org/10.1111/mec.14533

Grashei, K.E., Ødegård, J., Meuwissen, T.H.E., 2018. Using genomic relationship likelihood
for parentage assignment. Genet. Sel. Evol. 50(1), 26. https://doi.org/10.1186/s12711-018-
0397-7

Horn, S.S., Sonesson, A.K., Krasnov, A., Moghadam, H., Hillestad, B., Meuwissen, T.H.E.,
Ruyter, B., 2019. Individual differences in EPA and DHA content of Atlantic salmon are
associated with gene expression of key metabolic processes. Scientific Reports 9(1), 3889.

https://doi.org/10.1038/s41598-019-40391-2

Hu, G., Gu, W., Cheng, L., Wang, B., 2020. Genomic prediction and variance component
estimation for carcass fat content in rainbow trout using SNP markers. J. World Aquacult. Soc.
51(2), 501-511. https://doi.org/https://doi.org/10.1111/jwas.12677

Jiao, Y., Lu, Y., Li, X.-y., 2015. Farnesoid X receptor: a master regulator of hepatic triglyceride
and glucose homeostasis. Acta Pharmacologica Sinica 36(1), 44-50.
https://doi.org/10.1038/aps.2014.116

Kristjánsson, Ó.H., Gjerde, B., Ødegård, J., Lillehammer, M., 2020. Quantitative Genetics of
Growth Rate and Filet Quality Traits in Atlantic Salmon Inferred From a Longitudinal Bayesian
Model for the Left-Censored Gaussian Trait Growth Rate. Frontiers in Genetics 11.
https://doi.org/https://doi.org/10.3389/fgene.2020.573265

Langfelder, P., Horvath, S., 2008. WGCNA: an R package for weighted correlation network
analysis. BMC Bioinformatics 9(1), 559. https://doi.org/10.1186/1471-2105-9-559

Legarra, A., Ricard, A., Varona, L., 2018. GWAS by GBLUP: Single and Multimarker
EMMAX and Bayes Factors, with an Example in Detection of a Major Gene for Horse Gait.
G3 Genes|Genomes|Genetics 8(7), 2301-2308. https://doi.org/10.1534/g3.118.200336

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biology 15(12), 550. https://doi.org/10.1186/s13059-
014-0550-8

MacLeod, I.M., Bowman, P.J., Vander Jagt, C.J., Haile-Mariam, M., Kemper, K.E.,
Chamberlain, A.J., Schrooten, C., Hayes, B.J., Goddard, M.E., 2016. Exploiting biological
priors and sequence variants enhances QTL discovery and genomic prediction of complex traits.
BMC Genomics 17(1), 144. https://doi.org/10.1186/s12864-016-2443-6

Madison, B.B., 2016. Srebp2: A master regulator of sterol and fatty acid synthesis1. J. Lipid
Res. 57(3), 333-335. https://doi.org/https://doi.org/10.1194/jlr.C066712

Meuwissen, T.H.E., Hayes, B.J., Goddard, M.E., 2001. Prediction of Total Genetic Value Using
Genome-Wide Dense Marker Maps. Genetics 157(4), 1819-1829.
https://doi.org/10.1093/genetics/157.4.1819

Minghetti, M., Leaver, M.J., Tocher, D.R., 2011. Transcriptional control mechanisms of genes
of lipid and fatty acid metabolism in the Atlantic salmon (Salmo salar L.) established cell line,
SHK-1. Biochim. Biophys. Acta 1811(3), 194-202.
https://doi.org/https://doi.org/10.1016/j.bbalip.2010.12.008

Nohturfft, A., Brown, M.S., Goldstein, J.L., 1998. Sterols regulate processing of carbohydrate
chains of wild-type SREBP cleavage-activating protein (SCAP), but not sterol-resistant
mutants Y298C or D443N. Proceedings of the National Academy of Sciences 95(22), 12848-
12853. https://doi.org/10.1073/pnas.95.22.12848

Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., Kingsford, C., 2017. Salmon provides fast and
bias-aware quantification of transcript expression. Nat. Methods 14(4), 417-419.
https://doi.org/10.1038/nmeth.4197

Pena, R.N., Ros-Freixedes, R., Tor, M., Estany, J., 2016. Genetic Marker Discovery in Complex
Traits: A Field Example on Fat Content and Composition in Pigs. Int J Mol Sci 17(12).
https://doi.org/10.3390/ijms17122100

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., Smyth, G.K., 2015. limma
powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic
Acids Res. 43(7), e47-e47. https://doi.org/10.1093/nar/gkv007

Robinson, M.D., McCarthy, D.J., Smyth, G.K., 2009. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26(1), 139-140.
https://doi.org/10.1093/bioinformatics/btp616

Shimano, H., Sato, R., 2017. SREBP-regulated lipid metabolism: convergent physiology —
divergent pathophysiology. Nature Reviews Endocrinology 13(12), 710-730.
https://doi.org/10.1038/nrendo.2017.91

Skiba-Cassy, S., Lansard, M., Panserat, S., Médale, F., 2009. Rainbow trout genetically
selected for greater muscle fat content display increased activation of liver TOR signaling and
lipogenic gene expression. American Journal of Physiology-Regulatory, Integrative and
Comparative Physiology 297(5), R1421-R1429. https://doi.org/10.1152/ajpregu.00312.2009

Tocher, D.R., 2015. Omega-3 long-chain polyunsaturated fatty acids and aquaculture in
perspective. Aquaculture 449, 94-107. https://doi.org/10.1016/j.aquaculture.2015.01.010

Tsai, H.Y., Hamilton, A., Guy, D.R., Tinch, A.E., Bishop, S.C., Houston, R.D., 2015. The
genetic architecture of growth and fillet traits in farmed Atlantic salmon (Salmo salar). BMC
Genet. 16, 51. https://doi.org/10.1186/s12863-015-0215-y

Vance, D.E., Vance, J.E., 2008. CHAPTER 8 - Phospholipid biosynthesis in eukaryotes, in:
Vance, D.E., Vance, J.E. (Eds.), Biochemistry of Lipids, Lipoproteins and Membranes (Fifth
Edition). Elsevier, San Diego, pp. 213-244.

Wang, D.Q.H., 2007. Regulation of Intestinal Cholesterol Absorption. Annu. Rev. Physiol.
69(1), 221-248. https://doi.org/10.1146/annurev.physiol.69.031905.160725

Wen, J., Mercado, G.P., Volland, A., Doden, H.L., Lickwar, C.R., Crooks, T., Kakiyama, G.,
Kelly, C., Cocchiaro, J.L., Ridlon, J.M., Rawls, J.F., 2021. Fxr signaling and microbial
metabolism of bile salts in the zebrafish intestine. Science Advances 7(30), eabg1371.
https://doi.org/10.1126/sciadv.abg1371

Yang, J., Lee, S.H., Goddard, M.E., Visscher, P.M., 2011. GCTA: A Tool for Genome-wide
Complex Trait Analysis. The American Journal of Human Genetics 88(1), 76-82.
https://doi.org/https://doi.org/10.1016/j.ajhg.2010.11.011

Ødegård, J., Moen, T., Santi, N., Korsvoll, S.A., Kjøglum, S., Meuwissen, T.H.E., 2014.
Genomic prediction in an admixed population of Atlantic salmon (Salmo salar). Frontiers in
Genetics 5. https://doi.org/10.3389/fgene.2014.00402
528

Supplementary data

Figure S1: Trans eQTL links on chromosome 3. Circos plot showing the top trans-eQTL

links between trait associated genes and chromosome 3. Colors reflect the co-expression

module that the trait associated genes belong to from Figure 4b.

File S1: Human readable gene name to NCBI id translations. List of human readable gene

names of lipid genes used in our analysis and their associated NCBI refseq identifiers. Column

1 - NCBI gene ID, column 2 – human readable gene name, column 3 – gene description.

File S2: Trait associated genes for fat content in Atlantic salmon. List of all salmon genes

significantly associated to GBV. Column 1 – Gene ID, column 2 – log2 fold change, column 3

– adjusted p-value, column 4 – gene name, column 5 – gene product.

File S3: Enriched KEGG pathways for GBV associated genes. List of all significantly ($p <$

0.05) enriched KEGG pathways among GBV associated genes. Column 1 – pathway name,

column 2 – pathway ID, column 3 – type of GBV correlation for genes in pathway, column 4

– number genes in pathway, column 5 – number of TAGs in pathway, column 6 – enrichment

p-value.

File S4: eQTL results of selected TAGs. List of SNPs significantly associated to TAG

expression ($p < 10^{-6}$). Column 1 – NCBI gene ID of TAG, column 2 – Chromosome containing

SNP, column 3 – SNP identifier, column 4 – Physical position of SNP, column 5 – Reference

allele, column 6 – Second allele, column 7 – Frequency of the reference allele, column 8 – SNP

effect, column 9 – Standard error, column 10 – p-value.