Title:

RdDM mutants reduce histone methylation rather than DNA methylation at the paramutated

maize *b1* enhancer

Authors:

Iris Hövel[1], Rechien Bader[1], Marieke Louwers[1,2], Max Haring[1,3], Kevin Peek[1], Jonathan I.

Gent[4] and Maike Stam[1]

Author Affiliations:

[1] Swammerdam Institute for Life Sciences, Universiteit van Amsterdam, P.O. box 1210, 1090

GE Amsterdam, The Netherlands.

[2] argenx BV, Industriepark Zwijnaarde 7, 9052 Zwijnaarde (Ghent), Belgium

[3] Universiteit van Amsterdam, P.O. box 19185, 1000 GD Amsterdam, The Netherlands.

[4] Department of Plant Biology, University of Georgia, Athens, Georgia 30602, USA.

Corresponding Author:

m.e.stam@uva.nl (e-mail), +31-20-5257655 (phone)

The author responsible for distribution of materials integral to the findings presented in this

article in accordance with the policy described in the Instructions for Authors

(https://academic.oup.com/plphys/pages/general-instructions) is: Maike Stam.

Short Running head:

Silencing release without DNA methylation loss

Abstract

Paramutation is the transfer of mitotically and meiotically heritable silencing information between two alleles. With paramutation at the maize *booster1* (*b1*) locus, the low expressed *B'* epiallele heritably changes the high expressed *B-I* epiallele into *B'* with 100% frequency. This requires specific tandem repeats and multiple components of the RNA-directed DNA methylation (RdDM) pathway, including the RNA-dependent RNA polymerase (RdRP, *mediator of paramutation1*, MOP1), the second-largest subunit of RNA polymerase IV and V (NRP(D/E)2a, *mediator of paramutation2,* MOP2), and the largest subunit of RNA Polymerase IV (NRPD1, *mediator of paramutation3*, MOP3). Mutations in *mop* genes prevent paramutation and release silencing at the *B'* epiallele. In this study we investigated the effect of mutations in *mop1*, *mop2* and *mop3* on chromatin structure and DNA methylation at the *B'* epiallele, and especially the regulatory hepta-repeat 100 kb upstream of the *b1* gene.

We show that mutations in *mop1* and *mop3* result in decreased repressive histone modifications H3K9me2 and H3K27me2 at the hepta-repeat. Associated with this decrease are partial activation of the hepta-repeat enhancer function, formation of a multi-loop structure, and elevated *b1* expression. In *mop2* mutants, which do not show elevated *b1* expression, H3K9me2 and H3K27me2 and a single-loop structure like in wild type *B'* are retained. Surprisingly, high DNA methylation levels at the *B'* hepta-repeat remains in all three mutants. Our results raise the possibility of MOP factors mediating RNA-directed histone methylation rather than RNA-directed DNA methylation at the *b1* locus.

Introduction

Paramutation involves meiotically heritable epigenetic changes in gene expression caused by *trans*-communication between homologous alleles (Hövel et al., 2015; Hollick, 2017; Springer and McGinnis, 2015; Ronsseray, 2015). Paramutation has been studied in plants, such as maize and tomato, but also in animals, such as mice, Drosophila and *C. elegans*. In maize, paramutation has been best studied at genes encoding transcriptional regulators of the flavonoid pigmentation pathway, *r1 (red1)*, *b1 (booster1)*, *pl1 (plant color1)*, and *p1 (pericarp color1)*. Most alleles of these genes do not participate in paramutation; they are neutral to paramutation; only specific alleles do. The ability to participate in paramutation is often dependent on repeated sequences (for reviews see Hövel et al., 2015; Hollick, 2017). The alleles participating in paramutation at the *b1* locus are *B′* and *B-I*. *B′* and *B-I* have the same DNA sequence; they are epialleles (Stam et al., 2002a). *B′* is the paramutated, inactive state of *b1* (light pigmented plant), and *B-I* the paramutable, active state of *b1* (dark pigmented plant). In a *B′/B-I* heterozygote, the low expressed *B'* epiallele is paramutagenic; it changes the high expressed *B-I* epiallele into *B'* with 100% efficiency, reducing the *B-I* transcription rate 10-20 fold (Patterson et al., 1993). The change of *B-I* into *B'* is mitotically and meiotically heritable and requires multiple tandem copies of part of an 853-nt sequence located ~100 kb upstream of the *b1* transcription start site (TSS) (Stam et al., 2002a; Stam et al., 2002b; Belele et al., 2013). *B-I* and *B'* carry seven copies of the 853-nt sequence (hepta-repeat), while *b1* alleles insensitive (neutral) to paramutation carry one copy of the repeat unit. Our previous study implicated DNA hypermethylation at the hepta-repeat in both the maintenance and establishment of silencing at the *b1* hepta-repeat, while the presence of Histone H3 lysine K27 dimethylation (H3K27me2) at the hepta-repeat and *b1* coding region was associated with the maintenance of the silenced *B'* state (Haring et al., 2010). Besides their role in paramutation, the 853-nt tandem repeats carry transcriptional enhancer sequences that are required for tissue-specific enhancement of *b1* expression (Belele et al., 2013; Louwers et al., 2009a). In *B-I*, tissue-specific activation of *b1* expression in husk tissue is associated with H3 acetylation (at K9 and K14) and nucleosome depletion at the hepta-repeat (Haring et al., 2010). High *b1* expression levels are in addition associated with chromosomal interactions between the hepta-repeat, TSS and other regulatory regions at the *b1* locus, ~15, ~47 and ~107 kb upstream of the TSS (Louwers et al., 2009a). In *B'*, chromosomal interactions with the TSS, but not with the ~15, ~47 and ~107 kb regions were observed.

To this date, several genes, called *mediator of paramutation* (*mop*) and *required to*

maintain repression (rmr), have been identified to be required to establish paramutation at

multiple maize loci (*b1, pl1, r1, p1*) (Dorweiler et al., 2000; Hollick et al., 2005; Sidorenko et

al., 2009; Sidorenko and Chandler, 2008; Sloan et al., 2014; Stonaker et al., 2009). In mutants

of these genes, the paramutagenic alleles can no longer silence the expression of paramutable

alleles. The *mop1* and *rmr* genes identified to play such role in maize share homology with

genes involved in the RNA-directed DNA methylation (RdDM) pathway in *Arabidopsis*

thaliana. This pathway induces *de novo* DNA methylation (5mC) in all sequence contexts

(CG, CHG and CHH, where H is an A, T, or C) at transposable elements and selected genes

(Matzke and Mosher, 2014; Rymen et al., 2020; Zemach et al., 2013). In the RdDM pathway,

plant-specific RNA Polymerase IV (Pol IV) transcripts are converted into double-stranded

RNA by RNA-DEPENDENT RNA POLYMERASE 2 (RDR2), and processed into 24-nt

small interfering RNAs (siRNAs) by DICER-LIKE 3 (DCL3). One strand of the siRNAs is

incorporated into a complex containing ARGONAUTE 4 (AGO4), targeting the resulting

complex to complementary transcripts generated by RNA Polymerase V (Pol V).

Subsequently, DOMAINS REARRANGED METHYLTRANSFERASE2 (DRM2) is

recruited, mediating *de novo* DNA methylation of cytosines in all sequence contexts (CG,

CHG and CHH; H = A, T or C). This in turn leads to recruitment of histone modifying

enzymes and chromomethyltransferases such as CROMOMETHYLASE 3 (CMT3), which

methylates DNA in the CHG context. It also leads to recruitment of the replication-coupled

DNA methyltransferase MET1, which methylates in the CG context.

107   The first gene identified to be required for the establishment of paramutation was

mop1, encoding a putative ortholog of RDR2 in Arabidopsis (Alleman et al., 2006; Dorweiler

et al., 2000; Haag et al., 2014; Woodhouse et al., 2006a). Subsequently *mop2/rmr7* and

mop3/rmr6 were identified (Erhard et al., 2009; Haag et al., 2014; Hollick et al., 2005;

Sidorenko et al., 2009; Sloan et al., 2014; Stonaker et al., 2009); *mop2/rmr7* encodes the

maize ortholog of NRP(D/E)2a, the second largest subunit of Pol IV (NRPD1) and Pol V

(NRPE1), while *mop3/rmr6*, encodes NRPD1, the largest subunit of Pol IV. Mutations in

mop1, *mop2/rmr7* and *rmr6* have been shown to cause a severe decrease in the global 24-nt

siRNA levels, which is in line with their expected function in the RdDM pathway (Erhard et

al., 2009; Jia et al., 2009; Nobuta et al., 2008; Sidorenko et al., 2009; Stonaker et al., 2009). In

addition, for *mop1-1* and *mop2-1* mutants, diminished siRNAs levels at the *b1* hepta-repeat

were shown (Arteaga-Vazquez et al., 2010; Sidorenko et al., 2009). The nature of the

contribution of RdDM is, however, still unclear. The *b1* repeats are transcribed and yield

siRNAs not only from *B'*, but also *B-I* and a neutral *b1* allele, suggesting that presence of

siRNAs is not sufficient for paramutation (Alleman et al., 2006; Arteaga-Vazquez et al.,
2010; Sidorenko et al., 2009).

In line with a role for RdDM, *mop1*, *mop2* and *mop3* mutations are shown to induce a
reduction in DNA methylation at specific transposons, especially at DNA sequence regions
with high levels of DNA methylation in a CHH context, also called mCHH islands (Gent et
al., 2014, 2013; Li et al., 2015, 2014). This reduction in DNA methylation at mCHH islands
was indicated to occur in all cytosine contexts and was strongest in *mop3* and weakest in
mop2 mutants. Decreased DNA methylation levels were associated with increased expression
of several transposons (Jia et al., 2009; Li et al., 2015; Woodhouse et al., 2006b).

The *mop1*, *mop2/rmr7* and *mop3/rmr6* gene products are not only necessary for
paramutation, they are also reported to be required to maintain the silent state of *B'* and *Pl'*,
the paramutagenic *b1* and *pl1* loci (Dorweiler et al., 2000; Hollick et al., 2005; Sidorenko et
al., 2009; Sloan et al., 2014; Stonaker et al., 2009). In these mutants, increased expression
and/or plant pigmentation was observed. Upon crossing out the mutations, depending on the
gene and the mutation studied, the epialleles either lose expression and become
paramutagenic again or retain expression and are indistinguishable from their non-
paramutated states. The first is true for *B'; B'* does not revert to *B-I* in any of the three mutants
(Dorweiler et al., 2000) Similarly, *Pl'* does not revert to the active *Pl-Rh* state in *rmr7* mutants
(Stonaker et al., 2009). Thus, the epigenetic memory of *B'* and *Pl'* persists in mutants in spite
of increased epiallele expression. In *rmr6* mutants, however, the epigenetic memory of *Pl'* can
be erased: *Pl'* can stably revert to *Pl-Rh* in *rmr6* mutants (Hollick et al., 2005).

To get a better understanding of the role of the *mop* genes in the maintenance of *B'*
silencing, more specifically their role in the maintenance of DNA methylation and repressive
histone modifications, we examined the effect of *mop1*, *mop2* and *mop3* mutants on
chromatin structure and DNA methylation at the *B'* epiallele, and especially the regulatory
hepta-repeat 100 kb upstream of the *b1* coding sequence. Surprisingly, our results show that
none of the three *mop* genes is required to maintain high DNA methylation levels at the *B'*
hepta-repeat. High DNA methylation levels are maintained in all three mutants. In *mop2*
mutants, the *B'* epiallele in addition remains low expressed and retains high levels of the
repressive histone modifications H3K9me2 and H3K27me2. The elevated *b1* expression
levels in *mop1* and *mop3* mutants are, however, associated with significant reductions in
H3K9me2 and H3K27me2 at the hepta-repeat. We propose that this decrease in repressive
histone modifications allows the activation of the hepta-repeat enhancer function despite the
high DNA methylation levels. Our results indicate that MOP1 and MOP3 play a role in

maintaining H3K9me2 and H3K27me2 independent of a role in establishing DNA
methylation.

Results

159
Approach

To disentangle the effect of the *mop1-1, mop2-1, mop2-2* and *mop3-1* mutations (Alleman et
al., 2006; Dorweiler et al., 2000; Li et al., 2015; Sidorenko et al., 2009; Sloan et al., 2014) on
the epigenetic marks and chromatin features that define the mitotically and meiotically
heritable *B'* state from the effect on tissue-specific activation of *B'* expression, where
relevant, experiments were performed on two types of tissues: seedling and husk tissue. In
seedling tissue, comprising 1-month old seedlings with the exposed leaf blades and roots
removed, *B'* as well as *B-I* are very low expressed (Haring et al., 2010). In husk tissue (the
leaves surrounding the ear) expression of *b1* is transcriptionally activated, resulting in high *B-*
I and low *B'* expression. *B'* in a wild-type background was compared to *B'* in *mop* mutant
backgrounds. Plants carrying the *B-I* allele served as a positive control for an active *b1*
epiallele. The *mop1-1, mop2-2* and *mop3-1* mutations act recessive in preventing
paramutation between *B'* and *B-I*, and in their effect on pigmentation (Dorweiler et al., 2000;
Sidorenko et al., 2009; Sloan et al., 2014), therefore for these mutants only homozygous
mutant tissues were used in our analyses. The *mop2-1* mutation acts dominant in preventing
paramutation, and was reported to act recessive for enhancing pigmentation (Sidorenko et al.,
2009). Tissues of both heterozygous *Mop2/mop2-1,* and homozygous *mop2-1* mutants were
used in our analyses. In the text individuals homozygous for *B'* and a *mop* mutation are
referred to as *B' mop1-1, B' mop2-1, B' mop2-2,* and *B' mop3-1. Mop2 B'/mop2-1 B'*
individuals are referred to as *B' Mop2/mop2-1*. Wild type *B'* and *B-I* are referred to as *B'* and
B-I.

***B'* expression levels are increased in *mop*1 and *mop*3, but not *mop*2 mutants**
To monitor the *b1* gene expression levels in the *mop* mutants in our greenhouse conditions,
we performed RNA blot analysis using seedling and husk tissue from the mutants as well as
B' and *B-I* wild type individuals as comparison.
In seedling tissue of *B' mop1-1, B' Mop2/mop2-1, B' mop2-1*, *B' mop3-1* and *B-I* no
expression of *b1* could be detected (triplicate samples; data not shown). As reported
previously, in wild-type husk tissue, in which the *b1* gene is transcriptionally activated, the *B'*

allele was lowly expressed, while *B-I* transcript levels were increased ~20-fold compared to

the *B'* levels (Figure 1) (Louwers et al., 2009a). In husk tissue from *B' mop1-1* and *B' mop3-*

1, the *b1* expression levels were ~3 fold and ~5-fold increased relative to that in *B'* plants,

respectively (Figure 1). The latter upregulation is in line with an 8-fold increase of *B'*

transcription in husk tissue observed in an *rmr6-1* mutant (Erhard et al., 2009; Hollick et al.,

2005). Both *B' mop1-1* and *B' mop3-1* adult plants display as intensely dark purple tissues as

B-I wildtype plants, indicating that the *b1* RNA expression levels in all three genotypes are

above a threshold needed to activate the maize pigmentation pathway. In *B' Mop2/mop2-1*

husk tissue, expression of *B'* remained low, while the expression in *B' mop2-1* and *B' mop2-2*

was slightly higher, in line with (slightly) higher pigment levels (Figure 1 and Supplemental

Figure S1**)**. In conclusion, although all three mutations prevent paramutation between *B'* and

B-I, the RNA blot data shows they have different effects on the *B'* expression level.

Regulatory sequences at the *b1* locus are activated in *B' mop1-1* and *B' mop3-1*

To test whether changes in *b1* transcript level in *B' mop1-1*, *B' mop2* and *B' mop3-1* plants

are mediated by the activation of the previously identified distant regulatory sequences at the

b1 locus (Belele et al., 2013; Louwers et al., 2009a; Stam et al., 2002a), we performed

chromatin immunoprecipitation (ChIP) with an anti-histone H3K9K14 acetylation (H3ac)

antibody on chromatin isolated from husk, tissue in which the *b1* gene is transcriptionally

activated. The precipitate was analyzed by quantitative PCR (qPCR) and normalized to *actin1*

values (Haring et al., 2007). The targeted genomic regions are shown in Figure 2A, and

Supplemental Figure S2; the primers used are listed in Supplemental Table S1. Compared to

the levels in *B'* wild-type, elevated levels of H3ac were clearly observed in husk tissue of *B'*

mop1-1 and *B' mop3-1* at the *b1* locus (Figure 2B; Supplemental Figure S3 and Supplemental

Table S2), whereby the H3ac levels at the hepta-repeat where generally higher in *B' mop3-1*

than *B' mop1-1* tissue. This is in line with the higher *B'* expression levels in *mop3-1*

compared to *mop1-1*. The H3ac levels in both genotypes were, however, generally still

significantly lower than in the highly transcribed *B-I* genotype, indicating that the enhancer

sequences at the *b1* hepta-repeat are intermediately activated in *mop1-1* and *mop3-1* plants. In

line with low *B'* transcript levels, in heterozygous and homozygous *mop2-1* mutants, *B'*

showed only very low H3ac levels at the hepta-repeat and other regions at the *b1* locus,

similar as seen for *B'* in a wild-type background (Figure 2B; Supplemental Figure S3 and

Supplemental Table S2). In conclusion, the higher *b1* transcript levels in *B' mop1-1* and *B'*

mop3-1 husk tissue are associated with higher histone acetylation levels at the hepta-repeat,

indicating activation of the distant regulatory sequences

at the *b1* locus.

Activated *b1* regulatory sequences are associated with a multi-loop structure

The low expression level of the *B'* epiallele in husk tissue has been associated with the

formation of a single loop between the *b1* TSS and hepta-repeat 100 kb upstream, whereas

transcriptional activation of the high expressed *B-I* epiallele has been associated with the

formation of a multi-loop structure between the TSS and regulatory regions ~15, ~47, ~100

and ~107 kb upstream (Louwers et al., 2009a). To examine if the activation of the regulatory

sequences at the *b1* locus in the *mop* mutants is associated with the formation of a multi-loop

structure, chromosome conformation capture (3C) was applied, using husk tissue from *B-I*, *B'*

(Louwers et al., 2009a) and *B' mop1-1*, *B' Mop2/mop2-1* and *B' mop3-1* plants, and *Bgl*II as

restriction enzyme. 3C allows the identification of physical interactions between selected

genomic regions. The TSS (fragment 1), hepta-repeat (fragment X) and a fragment ~47 kb

upstream (fragment VII) were used as a viewpoint (bait) (Figure 3, Supplemental Figure S2).

The *S-adenosyl methionine decarboxylase* (*Sam*) locus was used as an unrelated, internal

control for data normalization (Louwers et al., 2009a).

Our data show that in *mop1-1* and *mop3-1* husk tissue, the *B'* epiallele shows a *B-I*-

like multi-loop conformation, whereas in *Mop2/mop2-1* tissue *B'* shows a *B'*-like single-loop

conformation (Figure 3A-C). When using fragment I (*b1* TSS) as viewpoint, in all samples

the hepta-repeat (fragment X) showed high interaction frequencies (Figure 3A), whereby the

B' epiallele showed the lowest interaction frequencies in a wild-type and *Mop2/mop2-1*

background, and the highest in a *B' mop3-1* and *B-I* background. In addition, fragment I also

interacted with fragments XII (-107 kb), VII (-47 kb), and VI (-40 kb) and IV (-15 kb),

whereby high interaction frequencies with fragments VII, VI and IV where only observed in

B' mop1-1, *B' mop3-1* and *B-I* tissue, consistent with a multi-loop structure. When using

fragment X and VII as viewpoints (Figure 3, B and C), the high interaction frequencies

between the TSS, hepta-repeat and regions ~47, and ~15 kb upstream were confirmed for *B'*

mop1-1 and *B' mop3-1*. The interactions involving the region ~107 kb upstream were

confirmed for all genotypes, with *B'* in a wild type and *Mop2/mop2-1* background showing

the lowest interaction frequencies.

In summary, our 3C data show that the transcriptional enhancement of *B'* expression

in the *mop1-1* and *mop3-1* mutant background is associated with the formation of a *B-I*-like

multi-loop structure, whereas the low *B'* expression level in the *Mop2/mop2-1* background is

associated with a *B'*-like single-loop structure as observed.

_B'_ hepta-repeat DNA methylation level is retained in all _mop_ mutants

At RdDM-target loci, the *mop1, mop2* and *mop3* genes have been indicated to play a major

role in maintaining CHH methylation (mCHH), and a minor role in maintaining CG and CHG

methylation (mCG, mCHG) (Gent et al., 2014; Li et al., 2015, 2014). In these studies, a

decrease in mCG and mCHG seemed most prominent in a *mop3* and least prominent in a

mop2 mutant. Previous experiments showed that the silenced *B'* allele is DNA

hypermethylated at the hepta-repeat junction regions compared to *B-I* (Haring et al., 2010).

To examine if transcriptional activation of *B'* in *mop* mutants is associated with a decrease in

DNA methylation, DNA blotting and bisulfite sequencing experiments were performed.

DNA blot analyses were performed as described in Haring *et al.* (2010), using

genomic DNA from leaf and husk tissue, isolated from 4 to 27 individuals per genotype at

different stages of development. DNA was isolated and digested with *Eco*RI or *Bam*HI, which

released a discrete fragment containing the entire hepta-repeat, and eight different methylation

sensitive restriction enzymes (Figure 4), size fractionated, blotted and hybridized with a

repeat probe (for representative examples, and the sequence and sites examined see

Supplemental Figure S2, S4 and S5). To check for complete digestion, probes recognizing

unmethylated sequence regions within the *b1* locus were used. To approximate the DNA

methylation levels, relative band intensities of all detected restriction fragments were

computationally compared to all theoretical possible combinations of fragment intensities.

Compared to *B'* tissue, in *B' mop1-1* and *B' mop3-1* tissue we observed only a slight

decrease in DNA methylation level at the *B'* repeat junction regions, with the most decrease

observed in *B' mop3-1* tissues (Figure 4, Supplemental Figure S4 and S6). The decrease

occurred in several, but not all plants, and was observed at specific restriction sites, in a

stochastic manner; a decrease at a particular site was not necessarily associated with a

decrease at another site in the same sample. In *B' Mop2/mop2-1, mop2-1* as well as *mop2-2*

samples, no decrease in DNA methylation was observed at the hepta-repeat (Figure 4,

Supplemental Figure S4 and S6).

In addition to the slight decrease in DNA methylation, in most *B' mop1-1* and all *B'*

mop2 samples we detected hypermethylation at the repeat junctions (Figure 4 and

Supplemental Figure S4 and S6; *Hha*I and *Hae*II). In the *mop3-1* mutant, however, a

stochastic decrease in DNA methylation was observed at the site hypermethylated in the
mop1-1 and *mop2* mutants.

We expected that during the development of *B' mop1-1* plants, as previously
hypothesized (Ohtsu et al., 2007), a progressive decrease in DNA methylation levels would
be observed due to cell division. To test this hypothesis, DNA blot analysis was performed on
DNA from six to eight leaves collected during the development of six homozygous *B' mop1-*
1 plants (Supplemental Figure S7). In most plants monitored (5 of 6), we did detect a slight
decrease in DNA methylation, but in general, the maximum decrease in DNA methylation
was already observed in the second leaf stage, suggesting that this reduction in DNA
methylation may already have occurred in very early developmental stages such as embryos,
in which *Mop1* appears higher expressed compared to seedlings and leaves (Sekhon et al.,
2011; Woodhouse et al., 2006b).

DNA methylation at the *B'* repeat junction regions in *mop1-1*, *Mop2/mop2-1* and
mop3-1, and *B'* and *B-I* wild type was also monitored by targeted bisulfite sequencing, which
provides single base-pair resolution data on cytosine methylation. DNA was obtained from
leaf 4 tissue of V4 stage plants, and *Fie2* was used as a control for complete conversion
(Gutiérrez-Marcos et al., 2006). As observed with DNA blot analysis, the *B'* repeat junction
region was hypermethylated compared to that of *B-I*, both in wild-type and *mop* mutant leaf
tissue (Figure 4B, Supplemental Figure S8). Bisulfite sequencing confirmed the lack of
significant DNA demethylation in the *mop* mutants. In fact, there was no detectable decrease
in DNA methylation. This difference from the DNA blot analysis method could be due to the
limited number of samples that can be easily examined by targeted bisulfite sequencing. Also
note that bisulfite sequencing represents individual molecules in individual cells while DNA
blot analysis represents large population of cells. In addition, there may be variation between
the seven repeats, and DNA blot analyses provide the overall picture. Intriguingly, in contrast
to the expectations (Li et al., 2014), the vast majority of DNA methylation observed at the *B'*
allele was in the CG and CHG context, and the low level in the CHH context (3%) was
similar in *B-I* and *B' mop* samples (Figure 4B and C, Supplemental Figure S8). These
findings are in contrast to reported observations that the RdDM pathway, in which the MOP
proteins play a role, mediates *de novo* DNA methylation in all sequence contexts, including
CHH (Gent et al., 2014; Li et al., 2015, 2014; Matzke and Mosher, 2014).

Together, our data indicate that i) activation of the enhancer function of the hepta-
repeat occurs in the presence of high DNA methylation levels, ii) the MOP proteins plays a

minor role in maintaining DNA methylation at the *b1* hepta-repeat, and iii) the *B'* repeat

junction regions show very low mCHH levels.

Genome-wide variation in mCHH levels exists at RdDM loci

Our finding that the *B'* repeat junction region, which is likely to be an RdDM locus, has little

mCHH, prompted us to explore whether similar loci, with MOP1-dependent 24-nt siRNAs,

but little mCHH, are present elsewhere in the genome. To do this, we used published small

RNA and DNA methylation datasets from wild-type and *mop1-1* mutant developing ear (Gent

et al., 2014). First, we compared the abundance of perfect-matching, uniquely-mapping 24-nt

siRNAs across the maize B73 genome to identify loci with abundant siRNAs in wild-type and

at least ten-fold reduced coverage in the *mop1-1* mutant. We then categorized all identified

loci based on mCHH levels in developing ear, whereby loci with mCHH values of less

than 0.02 (mC/total C) were categorized as low mCHH *mop1* loci, and greater than or equal

to 0.02 were categorized as high mCHH *mop1* loci. For increased stringency, we used 50-bp

non-overlapping intervals across the genome, and only included loci where at least two

adjacent loci were both low or both high mCHH. All adjacent 50-bp loci fitting the same

category were then merged, producing 69 low mCHH *mop1* loci and 3107 high mCHH *mop1*

loci, all of which were at least 100-bp in length (Supplemental Figure S9). Note that these

numbers are certainly smaller than the actual numbers of such loci in the genome. For this

analysis, our aim was to determine whether the *b1* enhancer is an anomaly, not to rigorously

identify and categorize all *mop1* loci in the genome. Excluding multi-mapping siRNAs is the

largest limiting factor of this analysis, but was necessary for robust interpretation of the data.

Based on these numbers, approximately 2% of *mop1* loci have low mCHH levels, like

observed for the *B'* repeat junction region. While both the low and high mCHH *mop1* loci had

high abundance of 24-nt siRNAs, the siRNAs at high mCHH loci were on average about two-

fold more abundant than at low mCHH loci. Both sets were dominated by 24nt siRNAs,

though other lengths were also present and were also similarly more abundant at high mCHH

loci (Supplemental Figure S9).

In whole-genome studies like this, selecting for the tails of a distribution (in this case

low mCHH) will always yield some number of false positives. To evaluate whether the loci

we identified have reproducibly low mCHH, we used data from a different tissue (developing

second leaf), produced using a different method (enzymatic conversion rather than bisulfite

conversion) (Hufford et al., 2021). In spite of a general reduction in mCHH in leaf compared

to ear, the general patterns were highly reproducible, not just for mCHH, but also mCG and

mCHG (**Figure 5**). In particular, the mean mCHH in leaf was 0.02 in low mCHH loci, but

0.10 in high mCHH loci, and both mCG and mCHG showed high, heterochromatin-like levels

in both sets of loci (Gent et al., 2014). These data also indicate that the majority of low

mCHH loci with MOP1-dependent siRNAs in ear also have low mCHH in leaf.

Transcriptional activation is not associated with a decrease in nucleosome occupancy at

the *B'* repeats

Nucleosome occupancy plays an important role in gene regulation (Jiang and Pugh, 2009;

Struhl and Segal, 2013), with nucleosome depletion at regulatory regions being associated

with gene expression. To test whether the increased *B'* expression level in *mop1-1* and *mop3-*

1 plants (Figure 1) is associated with decreased nucleosome occupancy, ChIP-qPCR

experiments were performed using an anti-histone H3 antibody (H3core) that recognizes

canonical H3 and H3 variants. As observed before, the hepta-repeats display a low and high

nucleosome occupancy in *B-I* and *B'* husk, respectively (Figure 6, Supplemental Figure S10)

(Haring et al., 2010). This is consistent with the hepta-repeat acting as a transcriptional

enhancer in *B-I*, but not *B'* plants. In husk tissue of *B' mop1-1* plants, however, compared to

B' wildtype plants no significant differences in nucleosome occupancy level were observed at

the hepta-repeat; they were significantly different from those in *B-I* plants, but not *B'* plants

(Figure 6, Supplemental Figure S10, Supplemental Table S2). This indicates that in *B' mop1-*

1, the increased *B'* RNA level is not facilitated by nucleosome displacement. The histone H3

levels at the hepta-repeat in *B' mop3-1* plants were not significant different from both *B'* and

B-I plants, indicating they were at an intermediate level and the elevated *B'* RNA levels in *B'*

mop3-1 plants could be slightly facilitated by nucleosome displacement. *Mop2/mop2-1* plants

showed similar results at the *B'* hepta-repeat as *mop1-1* plants, while *mop2-1* homozygous

plants showed H3core levels that were significantly different from both *B'* and *B-I* wildtype.

They were higher than in *B'*.

Transcriptional activation is associated with reduced H3K9me2 and H3K27me2 levels

The silencing of transposons and other repeats is associated with DNA methylation,

H3K9me2 and H3K27me2 (Bernatavichute et al., 2008; Gent et al., 2014; Liu et al., 2021;

Roudier et al., 2011; West et al., 2014). In maize, H3K9me2 and H3K27me2 are both

enriched in heterochromatin, and they are enriched at low and moderately expressed genes,

respectively (relative to highly expressed genes) (Gent et al., 2014). Our previous study

indicated that the *B'* gene body is marked by H3K27me2 in both seedling and husk tissue,

while the *B'* hepta-repeat carries H3K9me2 and H3K27me2 only in husk (Haring et al., 2010). To examine if activation of the *B'* epiallele in *mop* mutant plants is associated with reduced H3K9me2 and H3K27me2 levels, ChIP-qPCR experiments were performed on seedling and husk tissue, using monoclonal antibodies with a much better signal-to-noise ratio than the antibodies used in our previous study (Figure 7, see Methods) (Haring et al., 2010). The *B'* and *B-I* epialleles are transcriptionally activated in husk, but not in seedling tissue. A high-copy *Copia* sequence was used to normalize the data (Haring et al., 2010, 2007).

Unlike our previous results, H3K9me2 was significantly enriched at the hepta-repeat in *B'* seedling as well as husk tissue, suggesting that the *B'* hepta-repeat is marked by H3K9me2 in a tissue-independent manner (Figure 7A, B, Supplemental Figure S11A, B). The *b1* gene itself in the *B'* epiallelic state did not show significant H3K9me2 levels, nor in the *B-I* epiallele. In line with H2K27me2 being associated with repressed sequences, H3K27me2 levels were significant enriched at the hepta-repeat, gene body and other sequences in *B'* and *B-I* seedlings, and *B'* husk tissue, tissue in which the *b1* gene is not activated (Figure 7C, D, Supplemental Figure S11C, D, Supplementary Table S2). In *B-I* husk tissue, where *b1* transcript levels are high (Figure 1), only background levels of H3K27me2 were detected across the *b1* locus.

In seedling tissue, in which the *b1* gene is transcriptionally inactive, the H3K9me2 levels were significantly lower at the *B'* hepta-repeat in *mop3-1* plants than in wild type *B'* plants (Figure 7A, C, Supplemental Table S2). The same is true for the H3K27me2 levels for both *mop1-1* and *mop3-1* seedlings. In husk tissue, both the H3K9me2 and H3K27me2 levels were significantly lower in *mop1-1* and *mop3-1* plants compared to wild type *B'* plants (Figure 7B, D and Supplemental Figure S11B, D, Supplementary Table S2). The significant decrease in H3K9me2 (*mop3-1*) and H3K27me2 (*mop1-1* and *mop3-1*) levels at the hepta-repeat is already observed in seedling tissue and therefore independent of transcriptional activation of the *b1* gene, suggesting this decrease is a true effect of these *mop* mutations. At the gene body (UTR, Ex3), only the H3K27me2 levels in *B'* *mop3-1* husk tissue were significantly different from those in wild-type *B'* plants, in line with the relatively high expression level in *B' mop3-1* plants. The increased *b1* expression levels in *B' mop1-1* husk tissue occurs without a significant decrease of H3K27me2 levels at the gene body.

In *Mop2/mop2-1*, *mop2-1* husk tissue, similar levels of H3K9me2 and H3K27me2 were observed at the *B'* hepta-repeat as in wild type *B'* tissue (Figure 7B, D and Supplemental Figure S11B, D, Supplementary Table S2). H3K27me2 levels at the *b1* coding region were also not significantly different. These observations were confirmed using the independent

mop2-2 mutant, showing that the results are not specific for a dominant mutation like *mop2-1;*

repressive chromatin marks at the *B'* epiallele are also retained in the *mop2-2* mutant.

In conclusion, in *mop1-1* and *mop3-1* mutants the repressive H3K9me2 and/or

H3K27me2 marks are reduced at the *B'* hepta-repeat. For H3K9me2, this reduction is tissue-

independent for *mop3-1*, while for H3K27me2 this is true for both mutants. We propose that

these lower levels allow transcriptional activation of the *B'* epiallele in husk tissue. The

retention of these marks in *mop2* mutants correlates with the lack of a significant increase in

B' transcript levels in these mutants.

431 **Discussion**

In this study we show that the RNA-dependent RNA polymerase (RdRP, MOP1) and the

largest subunit of RNA polymerase IV (NRPD1, MOP3), proteins in the RNA-directed DNA

methylation (RdDM) pathway, are involved in maintaining high levels of the repressive

histone marks H3K9me2 and H3K27me2 at the *b1* regulatory hepta-repeat, but not in

maintaining high DNA methylation levels. In the *mop1* and *mop3* mutants strong and

reproducible reductions in the levels of the repressive H3K9me2 and H3K27me2 histone

marks are detected at the regulatory hepta-repeat enhancer, but only weak and stochastic

reductions in DNA methylation. We also show that *mop1* and *mop3* mutants, which prevent

paramutation at multiple maize loci including *b1*, allow partial activation of the regulatory

hepta-repeat enhancer at the *B'* epiallele in the presence of high DNA methylation and

nucleosome occupancy levels.

Role of MOP proteins in maintenance of DNA methylation

Mutations in *Mop1* (RdRP), *Mop2* (NRP(D/E)2a), and *Mop3/Rmr6* (NRPD1) prevent

paramutation and significantly reduce overall 24-nt siRNAs levels (Arteaga-Vazquez et al.,

2010; Erhard et al., 2009; Nobuta et al., 2008; Sidorenko et al., 2009; Stonaker et al., 2009).

The *mop1-1* and *mop2-1* mutants have in addition been shown to significantly reduce the

levels of hepta-repeat-derived 24-nt siRNAs (Arteaga-Vazquez et al., 2010; Sidorenko et al.,

2009). We, however, did not observe significant reductions in mCG and mCHG at the repeat

junction region in any of the mutants examined. Earlier studies in maize did report reductions

in DNA methylation at so-called mCHH islands, whereby decreases were more severe in

mop3-1 than *mop1-1* plants and more in mCHG than in mCG (Gent et al., 2014; Li et al.,

2014). In addition, studies in Arabidopsis, showed decreased mCHG and mCG levels in *rdr2*,

and *nrpd1* mutants (Tang et al., 2016; Zemach et al., 2013). We only observed stochastic

decreases in DNA methylation with our DNA blot analyses in *mop1-1* and *mop3-1* mutants,
and to a larger extent in *mop3-1* than *mop1-1* plants. Not all plants, however, showed such
decrease at the restriction sites monitored, therefore, the activation of the hepta-repeat
enhancer in these mutants is only weakly associated with non-reproducible decreases in DNA
methylation.

Role for RdDM components in maintenance of repressive histone modifications.
The activation of the enhancer at the *B'* hepta-repeat in the *mop1-1* and *mop3-1* mutants is
associated with significant reductions in the repressive H3K9me2 and/or H3K27me2 marks.
Given the well-established roles for siRNAs in directing DNA methylation, the reductions in
these histone modifications would normally be explained as a downstream consequence of
loss of DNA methylation. In principle, the MOP siRNA machinery could also directly target
H3K9me2 or H3K27me2 through unidentified protein-protein interactions between argonaute
proteins and histone methyltransferases. Such a scenario is supported by the existence of
RNA-directed histone modification without RNA-directed DNA methylation in diverse fungi
and animals, including *Drosophila melanogaster*, *Caenorhabditis elegans* and
Schizosaccharomyces pombe (Castel and Martienssen, 2013; Gu et al., 2012; Pal-Bhadra et
al., 2004; Verdel et al., 2004). In addition, a few studies have implicated siRNAs in directing
H3K9me2 in Arabidopsis (Jackel et al., 2016; Parent et al., 2021), although effects on
H3K9me2 and DNA methylation could not be unequivocally disentangled.

Interestingly, it has been shown that tandem repeats are efficiently and stably silenced
by H3K9 methylation through RNA-mediated mechanisms (Grewal, 2023). Small RNAs
derived from only one or a limited number of repeats can target all the repeats at a given locus
and mediate a high density of H3K9 methylation, which in turn recruits histone deacetylases
(HDACs) that stabilize heterochromatic regions (Grewal, 2023; Zofall et al., 2022). Similarly,
small RNAs may be required to target high H3K9me2 levels at the hepta-repeat, which in turn
recruits HDACs stabilizing the silencing.

Resolving the hierarchy of histone methylation and DNA methylation is complicated
by indirect effects of developmental gene regulation. In husk, where the hepta-repeat serves
as a weak enhancer for the *b1* gene in the mutants, we expect that some of the reduction in
histone methylation is due to the presence of transcription factors and associated chromatin
modifiers. In seedling tissue, however, in which *b1* is transcriptionally repressed, the
reduction should be independent of transcriptional activation of *b1*. Thus, the stronger

reduction of H3K27me2 over H3K9me2 in *B' mop1-1* seedling could suggest a more direct

effect of the MOP siRNA pathway on H3K27me2.

A clear link between H3K27me2 and RdDM in plants has not been shown before.

H3K27me2 is, as H3K27me3, intimately associated with transcriptional silencing through

Polycomb group (PcG) proteins (Conway et al., 2015). While H3K27me3 is mainly

associated with repressed genes, H3K27me2 is localized more ubiquitously throughout the

genome. Data from animals indicate that, unlike H3K27me3, H3K27me2 is not associated

with stable binding of Polycomb Repressive Complex 2 (PRC2); it is indicated to act as a

protective layer against unspecific transcriptional activation (Ferrari et al., 2014; Lee et al.,

2015). This includes preventing the activation of enhancer sequences in inappropriate cell

types. In agreement with these findings, in Arabidopsis and maize, H3K27me2 is enriched at

silenced promoters, genes and repeats (Gent et al., 2014; Locatelli et al., 2009; Mathieu et al.,

2005; Park et al., 2012; Roudier et al., 2011), and in wheat at euchromatic transposons at

distal chromosome ends (Liu et al., 2021). In Arabidopsis, H3K27me2 enrichment seems

independent of the presence of H3K9me2 or DNA methylation (Mathieu et al., 2005), and in

wheat mutually exclusive with H3K9me2 (Liu et al., 2021), suggesting that H3K27me2 and

RdDM are independent of each other. In line with this idea, in maize, high H3K27me2 levels

are observed at both RdDM and non-RdDM loci (Gent et al., 2014). In maize

heterochromatin, however, H3K27me2 largely overlaps with H3K9me2 and DNA

methylation, except that H3K27me2 is found at more loci (Gent et al., 2014). Links between

small-RNA mediated transcriptional silencing and H3K27 methylation have been observed in

protozoa. In *Tetrahymena thermophila*, small-RNA dependent recruitment of H3K27me3 to

developmentally transcriptionally silenced loci has been observed (Liu et al., 2007). A strong

relation between H3K27me2 enrichment and small RNA-mediated transcriptional silencing

has been observed in the protozoan *Entamoeba histolytica* (Foda and Singh, 2015). Here we

show that in *B' mop1-1* and *mop3-1* plants, the H3K27me2 levels at the hepta-repeat are

significantly lower than in *B'* wild type plants, suggesting that in maize MOP factors can

mediate RNA-directed histone methylation.

Our observation of *mop* mutants resulting in reduced H3K27me2 and H3K9me2 levels

does not resolve the relationship and potential hierarchy between these two histone

modifications. It is also possible that the MOP siRNA pathway recruits more than one histone

methyltransferase. In this scenario, both H3K27 and H3K9 could be direct targets, whereby

the respective histone methyltransferases may be recruited at different levels in different

tissues. Regardless of which histone modifier could be part of the MOP siRNA pathway at the

b1 hepta-repeat, it is likely not an anomaly. Our finding that 2% of *mop1* siRNA loci have

low levels of mCHH is consistent with a common function of 24-nt siRNAs in something

other than directing DNA methylation.

Reinforcing cycle between mCHG and H3K9me2

How could a large role for siRNAs in directing histone modifications go unnoticed? For one

reason, it is difficult to separate DNA methylation from histone modifications, especially

given the massive pleiotropic effects of mutants. At the *b1* hepta-repeat, however, we

observed a tissue-independent decrease in H3K9me2 in *mop3-1*, while mCHG remains high.

mCHG is normally tightly linked to H3K9me2 because chromomethylases (CMTs) bind

H3K9me2 containing nucleosomes and methylate cytosines in the CHG context (Bartee et al.,

2001; Du et al., 2014, 2012; Lindroth et al., 2001; Stroud et al., 2014, 2013). In addition,

H3K9 histone methyltransferases bind to methylated cytosines to methylate H3K9 (Du et al.,

2014; Johnson et al., 2007). In maize, the CMTs ZMET2 and ZMET5 are required for mCHG

(Fu et al., 2018; Gent et al., 2014; Li et al., 2014; Papa et al., 2001); ZMET2 also binds

H3K9me2 (Du et al 2012). At the *B'* hepta-repeat, CHG methylation is, however, not

decreased in *mop1-1* and *mop3-1* mutants, while H3K9me2 is significantly decreased,

suggesting that CHG methylation is not sufficient to recruit H3K9 methyltransferases to

maintain high H3K9me2 levels at the *B'* hepta-repeat. This supports the idea that the MOP1

RdRP and especially MOP3 NRPD1 play a role in recruiting H3K9 methyltransferases. It is

unclear, however, how the high mCHG levels are maintained. Possibly, the reduced levels of

H3K9me2 are still sufficient for ZMET2/5 to maintain high mCHG levels.

Gene activation in presence of repressive marks

Previously, we postulated that low DNA methylation levels and lack of H3K9me2 and

H3K27me2 at the *B-I* hepta-repeat would allow the binding of transcription factors driving

activation of the hepta-repeat enhancer (Haring et al., 2010; Louwers et al., 2009a). We

showed that activation of the *B-I* hepta-repeat enhancer was associated with histone

acetylation and chromosomal interactions between the hepta-repeat, TSS and additional

regulatory sequences at the *b1* locus, ultimately resulting in tissue-specific transcriptional

enhancement of *b1*. The repressive marks at the *B'* hepta-repeat, DNA methylation,

H3K9me2, and H3K27me2 were hypothesized to hamper enhancer activation and block the

formation of a multi-loop structure, resulting in low *b1* expression. Accordingly, the presence

of these repressive epigenetic marks are also negatively correlated with enhancer activity in

other systems (Ferrari et al., 2014; Plank and Dean, 2014; Schmitz et al., 2022; Sekhon et al.,

2012; Xie et al., 2013). Here we show the remarkable finding that, in *mop1-1* and *mop3-1*

mutant plants, the regulatory sequences at the *b1* locus can be activated in the presence of

high levels of DNA methylation and nucleosomes. This activation is illustrated by increased

b1 RNA and H3ac levels, and the formation of a multi-loop structure at the *B'* epiallele (Figs.

1-3). The restoration of the *b1* expression is, however, only to about 15% (*mop1-1*) or 23%

(*mop3-1*) of the *B-I* level (Figure 1). Accordingly, the H3ac levels are (slightly) lower than

observed for *B-I*. We hypothesize that in *B' mop1-1* or *mop3-1* husk tissue, a reduction in

H3K9me2 and H3K27me2 levels at each hepta-repeat is sufficient to allow transcription

factors and other protein factors involved in enhancer function to bind the hepta-repeat, albeit

with a relatively low affinity. This then results in an intermediately active enhancer that can

physically interact with the additional regulatory sequences, ~107, ~47 and ~15 kb upstream

of the TSS; together they mediate the formation of an active transcription complex at the *b1*

promoter, upregulating *b1* transcription. Alternatively, there could be near 100% reduction of

H3K9me2 and H3K27me2 at a limited number of *b1* alleles, activating *b1* expression in only

a fraction of cells. In either case, the high DNA methylation levels at the hepta-repeat

enhancer allow the binding of proteins involved in enhancer activation to some extent.

574   Unlike in the *mop1-1* and *mop3-1* mutants, in both the dominant and recessive *mop2*

mutants there is no significant decrease in H3K9me2 and H3K27me2 at the hepta-repeat

enhancer, and the DNA methylation levels are, if anything, even slightly increased. Retention

of these repressive marks in *mop2* mutants correlates with a lack of enhancer activation,

which is associated with the formation of a single loop structure and no significantly

increased *B'* transcript levels in these mutants. *Mop2/Rmr7* codes for NRP(D/E)2a, the

second largest subunit of Pol IV and Pol V (Sidorenko et al., 2009; Stonaker et al., 2009).

NRP(D/E)2a has two paralogs, NRP(D/E)2b and NRPE2c. NRP(D/E)2b is also part of Pol IV

and V while NRPE2c is only part of Pol V (Haag et al., 2014; Sidorenko et al., 2009).

Nevertheless, neither of these paralogs is able to substitute for NRP(D/E)2a with respect to

paramutation in terms of the establishment of DNA and histone methylation at the *b1* hepta-

repeat. However, one of the paralogs may substitute for NRP(D/E)2a to maintain these marks.

Arabidopsis contains one functional homologue of NRP(D/E)2a, (NRP(D/E)2; (Herr et al.,

2005; Kanno et al., 2005; Onodera et al., 2005; Pontier et al., 2005)). DNA blot analysis on

Arabidopsis mutants of NRP(D/E)2 indicated a slight decrease in DNA methylation at 5S

genes. So far, no effects on histone modifications are reported. In conclusion, our results

indicate that repression of the *B'* epiallele can be maintained independent of MOP2 but does
require MOP1 and MOP3.
592
Effect on chromatin structure by mop mutants
The different *mop* mutants all hamper paramutation and block RdDM, but affect the
chromatin structure differently. In the *mop1-1* and *mop3-1* mutants, repression at the *B'*
hepta-repeat enhancer is released to different extents, but not in *mop2* mutants. We know little
about the effect of these mutants on the binding of the RdDM components that are still
functional. We expect, like observed for Arabidopsis, the MOP1 RdRP to stably bind the
largest subunit of Pol IV (NRPD1/Mop3/Rmr6) (Law et al., 2011; Mishra et al., 2021). The
effect of a lack of a functional MOP1 on the binding of the remaining RdDM components to
chromatin in a *mop1-1* mutant is unknown.
We hypothesize that in a *mop3* mutant, Pol V, but not Pol IV, is still functional,
resulting in transcription of the *B'* hepta-repeat by Pol V in the absence of RdDM. This
transcription may lead to the observed decrease of H3K9me2 and H3K27me2 at the *B'* hepta-
repeat, allowing enhancer activation. In line with this hypothesis, in Arabidopsis the
production of Pol V transcripts is independent of mutations affecting components in siRNA
biogenesis, and *de novo* and maintenance DNA methylation (DICER1-4, RDR2, DRM2,
MET1 and DDM1) (Wierzbicki et al., 2008).
As discussed above, we cannot exclude that in *mop2* mutants, NRP(D/E)2a is replaced
by NRP(D/E)2b or 2c, which may still be able to repress the *B'* hepta-repeat. Alternatively,
both Pol IV and Pol V become non-functional and in absence of their transcription activity,
the *B'* hepta-repeat stays repressed. In Arabidopsis, which has only one NRP(D/E), it has
been seen that in the absence of NRP(D/E)2, the largest subunit of Pol V is destabilized,
while NRPD1 is not (Pontier et al., 2005). If also true in maize, the Pol V protein complex
would no longer bind, while the Pol IV complex may be dysfunctional.
The dissimilar effect of *mop1*, *mop2* and *mop3* mutations on the repressed state of the
B' hepta-repeat is in line with findings for *pl1*. In 2-10% of wild-type plants, *Pl'*
spontaneously reverts to the *Pl-Rh* state (Hollick et al., 1995) and is therefore less stable than
the *B'* state, for which no reversion to *B-I* has been observed. In *rmr6-1* (*mop3*) mutants, *Pl'*
changes to *Pl-Rh* with an increased reversion rate of 36% (Hollick et al., 2005), and also in
mop1-1 mutants a higher reversion rate to *Pl-Rh* was observed than in wild-type plants
(Dorweiler et al., 2000). In *rmr7* (*mop2*) mutants, however, *Pl'* does not revert to *Pl-Rh*,

indicating that the *Pl'* epiallele is more stably repressed in an *nrp(d/e)2a* mutant than in wild-
type plants (Stonaker et al., 2009).

Epigenetic memory

For *mop1-1* and *rmr6* (*mop3*) mutant plants, in which the production of siRNAs is severely
decreased (Arteaga-Vazquez et al., 2010; Erhard et al., 2009; Nobuta et al., 2008; Sidorenko
et al., 2009; Stonaker et al., 2009), it has been shown that the epigenetic memory of the
paramutagenic *B'* state is maintained (Dorweiler et al., 2000; Hollick et al., 2005). Upon
crossing out the *mop* mutations, *B'* can again paramutate *B-I* and is again lowly expressed,
indicating the RdDM pathway is not involved in maintaining the epigenetic memory. We
hypothesize that mCG and mCHG play a crucial role in maintaining the *B'* epigenetic state
through meiosis, as in all *mop* mutants, the *B'* hepta-repeat remains associated with high
mCG and mCHG levels. DNA methylation has been shown to play a central role in the stable
transgenerational inheritance of silent epigenetic states in plants (Fitz-James and Cavalli,
2022; Mathieu et al., 2007). In line with this, Calarco et al. (2012) showed that mCG and
mCHG are largely maintained in the germline. We, however, cannot entirely exclude a role
for H3K9me2 and H3K27me2, even when the *B'* hepta-repeat H3K27me2 levels are
significantly reduced tissue-independently in *mop1-1* and *mop3-1* mutants, and H3K9me2
levels in the *mop3-1* mutant. H3K9 methylation has been indicated to play a role in
transgenerational inheritance in *S. pombe*, *D. melanogaster* and *C. elegans* (Fitz-James and
Cavalli, 2022), and immunocytological data suggests that H3K9me2 may be retained during
meiosis (Underwood et al., 2018). Currently, there is no role reported for H3K27me2 in
transgenerational inheritance in plants.

Concluding remarks
This study showed that, in the presence of high DNA methylation, a reduction in the
repressive H3K9me2 and H3K27me2 marks at the regulatory *b1* hepta-repeat is sufficient to
allow partial activation of the hepta-repeat enhancer function. This indicates that the mere
presence of DNA methylation is not sufficient to entirely block enhancer function and that
high levels of H3K9me2 and H3K27me2 are involved in blocking enhancer function. We also
showed that high mCHG levels can be maintained in the presence of significantly reduced
H3K9me2 levels, suggesting the reduced H3K9me2 levels are still sufficient for ZMET2/5 to
maintain high mCHG levels. Intriguingly, we show that in maize, the RdRP and largest
subunit of RNA Pol IV involved in RdDM also play a role in the maintenance of H3K9me2

and H3K27me2. More experiments are necessary to show if this maintenance role is confined
to a limited number of sequences or occurs genome-wide. Our results also emphasize the
possibility of RNA-directed histone modification in plants, as occurs in diverse fungi and
animals. Future studies will be needed to examine the role of the MOP siRNA pathway in
H3K9me2 and H3K27me2, and if RNA-directed histone modifications play a role in
paramutation.

Methods

Genetic Stocks and Plant Materials

The plants stocks used in this study (*B'/B' Mop1/Mop1* (K55), *B-I/B-I Mop1/Mop1* (W23),
B'/B' Mop1/mop1-1 (K55/W23), *Mop2 B'/mop2-1 B'* (W23/K55), *Mop2 B'/mop2-2 B'*
(W23/K55), *B'/B' Mop3/mop3-1* (W23/K55) were obtained from V.L. Chandler (University
of Arizona, Tucson, AZ) and grown in greenhouse conditions. Homozygous *mop* mutants
were obtained by crossing *B'/B' mop1-1/Mop1* with *B'/B' mop1-1/mop1-1, Mop2 B'/mop2-1*
B' with *mop2-1 B'/mop2-1 B', Mop2 B'/mop2-2 B'* with *mop2-2 B'/mop2-2 B',* and *B'/B'*
Mop3/mop3-1 with *B'/B' mop3-1/mop3-1*, respectively. Homozygous progeny were identified
using PCR genotyping (*mop1-1*) followed by sanger sequencing (*mop2*) and/or scoring
pigment levels (*mop1-1*, *mop3-1*) (Sidorenko et al., 2009; Sloan et al., 2014). Heterozygous
B' Mop2/mop2-1 mutants for experiments were mainly obtained from a cross of *Mop2*
B'/mop2-1 B' with *mop2-1 B'/mop2-1 B'.* In addition, for a few RNA and DNA isolations also
samples from a cross between wild type *B'* and homozygous *mop2-1 B'* plants were used (data
in Supplemental Figure S1, S6; derivation is indicated in the figures). The seedling tissue
used for ChIP experiments comprised of 1-month-old seedlings with the roots and exposed
leaf blades removed. The husk tissues employed for RNA, ChIP and 3C consisted of all
leaves surrounding the maize ear, whereby the tough, outer leaves were discarded (Haring et
al., 2010; Louwers et al., 2009a). Husks were harvested at the time of silk emergence. For
DNA methylation analysis, mainly leaf blade tissue served as input material.

RNA Analysis

RNA blot analysis was performed as described previously (Louwers et al., 2009a; Haring et
al., 2010). RNA was isolated from husk tissue and 10 µg of RNA was size-fractionated by
formaldehyde gel electrophoresis, blotted and hybridized with probes against the *b1* or *Sam*

gene (Louwers et al., 2009a). RNA blots were hybridized with probes recognizing the coding

region of *b1* or *Sam*. Band intensities were quantified using a Storm 840 phosphorimager and

ImageQuant software (GE LifeSciences) and relative transcript levels of *b1* were calculated

by normalization to transcript levels of the *Sam* housekeeping gene.

Chromatin Immunoprecipitation

ChIP-qPCR experiments were performed on husk and seedling tissue as previously described

(Haring et al., 2010, 2007). Antibodies were used against H3K9ac/K14ac (Upstate #06-599),

H3K9me2 (Cell signaling #4658), H3K27me2 (Cell signaling #9728) and histone H3 (Abcam

#1791). For the no-antibody reactions, 20 μl Rabbit serum/ChIP sample (Sigma R9133) was

used. The specificity of the H3K9me2 and H3K27me2 antibodies was validated by slot blots

containing different dilutions of the target peptides (Egelhofer et al., 2011). All ChIP-qPCR

experiments were done using PLAT SYBR QPCR 500 (Invitrogen/Fisher, # 11733046) on an

Applied Biosystems™ 7500 Real-Time PCR System, repeated 2-4 times with chromatin

derived from different plants (see legends Figures 2,6 and 7, and Supplemental Figures S3,

S11, S11). Primers that were used for the qPCR analysis are listed in Supplemental Table S1.

Actin and *copia* served to normalize ChIP-qPCR data.

To test if *B'*, *B-I*, *B' mop1-1, B' Mop2/mop2-1*, *B' mop2-1*, *B' mop2-2* and *B' mop3-1*

behaved similar for the enrichment of the measured histone modifications in either seedling or

husk tissue, two-way Analysis of variance (ANOVA) tests were used. Given the limited

sample number, we assumed all datasets to be normally distributed and having similar

variances. When the ANOVA tests indicated significant differences between the different

(epi)genotypes, a pairwise t-test with a Bonferroni correction for multiple testing was

performed to uncover which individual pairs differ significantly. The output from the

ANOVAs and pairwise t-tests is summarized in Supplemental Table S2.

The *Copia* sequence behaves, on average, as a non-RdDM locus (Parent et al., 2021)

and could therefore be used to normalize ChIP-qPCR data obtained for *B' mop1-1* plants. In

previous experiments, *Copia* sequences were used to normalize ChIP-qPCR data obtained

with antibodies recognizing H3K9me2 and H3K27me2 (Haring et al., 2010, 2007). The

majority of *Copia* elements are depleted of siRNAs (Gent et al., 2013), indicating they are

rarely targeted by RdDM. To validate that the specific type of *Copia* sequence used in our

previous experiments was not targeted by RdDM and could be employed for normalization in

mop mutant backgrounds, the 107-nt sequence was blasted to the maize B73 genome (version

3) using parameters optimized for low similarity (e-value 1e-15) and including low

complexity regions. Of the 163 matches, the 97 that were greater or equal to 95-nt in length,

carried at least part of both primer binding sites, and had sequence coverage in a bisulfite

sequencing experiment (Gent et al., 2014) were selected. These 97 loci were examined for the

presence of CHH methylation, a hallmark of RdDM. Their average CHH methylation level

was 1.66%, which is very close to the 1.34% of CHH methylation at non-RdDM loci and

significantly different from the average CHH methylation level at RdDM loci (14.51%). In

addition, analysis of H3K9me2 data from the same study revealed that these loci had an

H3K9me2 enrichment value of 1.10, which is characteristic of non-RdDM loci (H3K9me2

enrichment of 1.12) rather than RdDM loci (H3K9me2 enrichment of 0.55) (Gent et al.,

2014). This indicated that our *Copia* control sequence behaves, on average, as a non-RdDM

locus and that it could be used to normalize our ChIP-qPCR data obtained from wild-type and

mop mutant plants.

Chromatin Conformation analysis (3C)

3C analyses were performed on *B' mop1-1* husk tissue as described previously (Louwers et

al., 2009a, 2009b). The data for *B' Mop1/Mop1* and *B-I Mop1/Mop1* have been published

before (Louwers et al., 2009a) and are shown for comparison. The experiments on *B' mop1-1*

tissues were conducted at the same time and using the same methods as the experiments on *B'*

Mop1/Mop1 and *B-I Mop1/Mop1* tissues. The primers and TaqMan probes used are listed in

Louwers *et al.* (2009a) In 3C experiments, to correct for primer amplification efficiencies, for

each primer pair the qPCR data was normalized to a random ligation control sample produced

using a BAC clone containing the *b1* locus and an amplified control template for the *Sam*

locus. To control for quantity and quality of the 3C samples, the data were normalized to 3C

values measured for the *Sam* locus.

DNA Methylation Analysis by DNA Blotting

DNA blot analysis was performed as described previously (Haring et al., 2010). Most DNA

samples were derived from leaves collected at different stages of development (Supplemental

Figures S4, S6 and S7). DNA was isolated (Dellaporta et al., 1983) and 5 μg of DNA was

digested with different methylation sensitive restriction enzymes in combination with *Eco*RI

or *Bam*HI according to the manufacturer's specifications. After size fractionation by

electrophoresis the gels were blotted onto membrane (Blotting-Nylon 66 membranes, Sigma-

Aldrich # 15356), followed by UV fixation. The resulting blots were hybridized to a 853-nt

repeat probe. The relative band intensities obtained for the enzymes *Hpa*I, *Pst*I, *Hha*I, *Hae*II

*Alu*I, *Sau*96I, *Bsm*AI were quantified for all (*Hpa*I, *Hha*I, *Hae*II) or representative examples

(*Pst*I, *Alu*I, *Sau*96I, *Bsm*AI), using a Storm 840 phosphorimager and ImageQuant software

(GE LifeSciences). The most probable DNA methylation patterns were identified by the least

squares fit comparing the relative band intensities with theoretical possible intensities derived

from all possible options for DNA methylation patterns (Haring et al., 2010). *Bsm*AI

generated relatively small fragments, including similarly sized 5' and 3' border fragments

complicating the computational analysis. The degrees of DNA methylation indicated (Figure

4A, Supplemental Figure S6, S7) are our best estimates. To test for complete digestion,

probes recognizing unmethylated DNA regions at the *b1* locus were used (probes A, D2 and

21, Figure 1 in Haring et al., 2010).

DNA Methylation Analysis by Bisulfite Sequencing

For bisulfite sequencing, genomic maize DNA was extracted (Dellaporta et al., 1983) from

leaf four of two *B'*, two *B-I* and three *B' mop1-1* plants in a V4 stage (Ritchie et al., 1986),

and 400 ng of DNA was treated with bisulfite using the EZ DNA Methylation-Gold kit

(Zymo Research, D5006). The DNA regions of interest were PCR-amplified (10 min 95°C,

followed by 40 PCR cycles (30 sec 95°C, 30 sec 50°C (repeat) or 52°C (*Fie2*), 30 sec 72°C),

and 5 min at 72°C). The amplification was performed using MethylTaq DNA polymerase

(Diagenode, C09010010), a forward (KL1310; TGGTGTTTAAAAATTYATGTTTTTGTG)

and reverse primer (KL1844; TCCACRARTCATCRTCTCCAAACA) for a 320-nt repeat

junction fragment and a forward (AAGATTTGAGATTYGATTTGAAGTGTG) and reverse

primer (ACTTTCCCCTCCRCCTAATTCTCCTTA) for a 226-nt *Fie2* fragment that served

as a control for complete bisulfite conversion (similar to the -302 to -91 fragment described in

Gutierrez-Marcos et al., 2006). PCR fragments were cloned into the pJET 1.2 vector

(CloneJet PCR Cloning Kit, Thermo Scientific) following the manufacturer's instructions.

Clones carrying inserts of the correct size were identified by PCR. DNA was isolated from

positive clones (GeneJet Plasmid Miniprep Kit, Thermo scientific) and subjected to Sanger

sequencing. To determine the percentage, sequence context and pattern of cytosine

methylation, sequences were analyzed using Kismeth (Gruntman et al., 2008). For each DNA

sample approximately 15 clones were sequenced for *Fie2*, and 30 clones for the repeat

junction region.

Genomic Alignments and Data Analysis of DNA methylation-seq and Small RNA Reads

Small RNA sequences from developing ear were obtained from the Sequence Read Archive,
runs SRR1583941 and SRR1583942 for the *mop1-1* mutant; and SRR1583943 and
SRR1583944 for wild-type (Gent et al., 2014). Both were in a genetic background closely
related to B73. The methylC-seq data from the same study were derived from pure B73
developing ear, runs SRR1583945, SRR1583946, and SRR1583947. The Enzymatic Methyl-
seq (EM-seq) data from developing second leaf of B73 were from the NAM founders
assembly project and available through the European Nucleotide Archive, runs ERR5347654,
ERR5347655, and ERR5347656 (Hufford et al., 2021).

Both methylC-seq and EM-seq reads were processed and mapped with BS-
Seeker2 (Guo et al., 2013) according to published methods (Hufford et al., 2021), except that
no mismatches were allowed (-m 0) and mapping was done in end-to-end mode (--bt2--end-
to-end). Potential PCR duplicates were removed from the ear methylC-seq reads using
the BS-Seeker2 FilterReads tool. Methylation values and coverage for genomic loci were
obtained using the CGmapTools mtr tool for each of the three sequence contexts separately
(Guo et al., 2018). For defining low mCHH and high mCHH *mop1* loci, a minimum effective
coverage of 20 was required, where effective coverage is the sum of the read coverage of each
individual cytosine in each locus. Methylation values were measured as the average of each
site in each locus. Low mCHH *mop1* loci were defined as having less than 0.02 mCHH, and
high mCHH *mop1* loci as having greater than or equal to 0.02 mCHH in the source B73 ear
methylome (Gent et al., 2014). These were initially identified based on 50-bp non-
overlapping intervals, but all adjacent loci that met criteria for low mCHH were merged, and
all adjacent loci that met criteria for high mCHH were also merged. Only merged loci
consisting of at least two adjacent loci were included in the final sets, giving them minimum
lengths of 100 bp. The methylation of these loci was reevaluated using the leaf methylome
(Hufford et al., 2021), but with a constraint that only loci with at least five cytosines of the
specific cytosine context being spanned by reads were included in the analysis.

Small RNA-seq reads were quality filtered, trimmed of adapters, and filtered for
lengths of 20-25 nt using cutadapt (Martin, 2011), parameters -q 20 -a
AGATCGGAAGAGCACACGTCT -e .05 -O 5 --discard-untrimmed -m 20 -M 25. Reads
were aligned to the genome with Bowtie2, –very-sensitive parameters (Langmead and
Salzberg, 2012). Reads that overlapped at least 90% of their lengths with tRNA, 5S RNA,
NOR, or miRNA loci were removed using the BEDTools intersect tool with parameters -v -f
.9 (Quinlan and Hall, 2010). These excluded loci were identified using published methods
(Gent et al., 2022), but applied to the B73 v5 reference genome (Hufford et al., 2021). The

read counts for miRNA loci were used for normalization in downstream steps. The two input
files for wild-type and the two for the *mop1-1* mutant were combined into single files using
the SAMtools merge tool (Li et al., 2009) Uniquely-mapping reads were selected using the
SAMtools view tool with the -q20 parameter. Perfect-matching siRNAs were selected using
grep for the pattern 'NM:i:0'. 24-nt siRNAs were selected from the complete set of 20 to 25-nt
siRNAs using grep for the pattern '24M'. Read coverage over 50-bp intervals of the genome
was obtained using the BEDTools coverage tool. All intervals were identified that had at least
25 overlapping reads and which spanned at least 30 bp of the interval from wild-type. In
addition, they had to have a minimum of ten-fold fewer overlapping reads in the *mop1-1*
mutant than in wild-type (normalized by the number of miRNA reads). These, called *mop1*
loci, were then further filtered using methylation data as to generate low mCHH and high
mCHH *mop1* loci as described above.

837
Gene IDs
mop1: Zm00001eb080370
mop2/rmr7: Zm00001eb068960
mop3/rmr6: Zm00001eb035080

Acknowledgements
We would like to thank Vicki Chandler and Lyudmila Sidorenko for providing seeds. We are
grateful to Nicole Riddle and Sarah Gadel who tested the specificity of the H3K9me2 and
H3K27me2 Cell signaling antibodies with Slot Blots. Shiv Grewal is thanked for advice and
fruitful discussion, Ludek Tikovsky and Harold Lemereis for taking excellent care of the
maize plants, and Christel Rooker for proofreading of the manuscript. I.H. was supported by
the Systems Biology Research Priority fund of the University of Amsterdam. K.P. is
supported by the Topsector Horticulture & Starting materials; J.I.G was supported by a
National Science Foundation grant (2114797). M.S. was supported by the Royal Netherlands
Academy of Arts and Sciences (KNAW).

Author contributions
I.H., M.L., M.H., J.I.G. and M.S. designed the research; R.B., I.H., M.L., M.H and J.I.G.
performed research, analyzed data and made figures; K.P. performed statistical analysis and
generated figures; I.H., J.I.G. and M.S. wrote the manuscript.

Supplemental data

Supplemental Figure S1. *b1* expression levels in maize husk tissues.

Supplemental Figure S2. Schematic representation of the *b1* and *Sam* locus.

Supplemental Figure S3. *B'* hepta-repeat is marked by H3 acetylation in *mop1-1* and *mop3-*
1 mutants.

Supplemental Figure S4. In *mop* mutants slight differences in DNA methylation were
observed at the *B'* repeat junction regions by DNA blot analysis.

Supplemental Figure S5. Consensus double-stranded DNA sequence of the 2nd-6th repeat.

Supplemental Figure S6. Detailed DNA methylation data for *B'* in *mop* mutants using
methylation sensitive restriction enzymes and DNA blot analysis.

Supplemental Figure S7. DNA methylation levels at *B'* hepta-repeat do not progressively
decrease during development of a *B' mop1-1* plant.

Supplemental Figure S8. Targeted bisulfite data for *B'*, *B-I*, *B' mop1-1*, *B' Mop2/mop2-1*
and *B' mop3-1* DNA derived from leaf 4 of V4 stage plants.

Supplemental Figure S9. Identification of low and high mCHH *mop1* loci and their
respective siRNA levels.

Supplemental Figure S10. No decrease in nucleosome occupancy was observed in the *mop*
mutants.

Supplemental Figure S11. Transcriptional activation in *mop1-1* and *mop3-1* is associated
with reduced levels of repressive histone marks.

Supplemental Table S1. Primers used in ChIP-qPCR experiments.

Supplemental Table S2. Statistical analyses of the ChIP-qPCR experiments.

References

Alleman, M., Sidorenko, L., McGinnis, K., Seshadri, V., Dorweiler, J.E., White, J., Sikkink,
K., Chandler, V.L., 2006. An RNA-dependent RNA polymerase is required for
paramutation in maize. Nature 442, 295–298. https://doi.org/10.1038/nature04884
Arteaga-Vazquez, M., Sidorenko, L., Rabanal, F.A., Shrivistava, R., Nobuta, K., Green, P.J.,
Meyers, B.C., Chandler, V.L., 2010. RNA-mediated trans -communication can
establish paramutation at the b1 locus in maize. PNAS 107, 12986–12991.
https://doi.org/10.1073/pnas.1007972107/-
/DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1007972107
Bartee, L., Malagnac, F., Bender, J., 2001. Arabidopsis cmt3 chromomethylase mutations
block non-CG methylation and silencing of an endogenous gene. Genes Dev 15,
1753–1758. https://doi.org/10.1101/gad.905701
Belele, C.L., Sidorenko, L., Stam, M., Bader, R., Arteaga-Vazquez, M.A., Chandler, V.L.,
2013. Specific Tandem Repeats Are Sufficient for Paramutation-Induced Trans-

Generational Silencing. PLoS Genetics 9.
https://doi.org/10.1371/journal.pgen.1003773
Bernatavichute, Y.V., Zhang, X., Cokus, S., Pellegrini, M., Jacobsen, S.E., 2008. Genome-
wide association of histone H3 lysine nine methylation with CHG DNA methylation
in Arabidopsis thaliana. PLoS One 3, e3156.
https://doi.org/10.1371/journal.pone.0003156
Calarco, J.P., Borges, F., Donoghue, M.T.A., Van Ex, F., Jullien, P.E., Lopes, T., Gardner, R.,
Berger, F., Feijó, J.A., Becker, J.D., Martienssen, R.A., 2012. Reprogramming of
DNA Methylation in Pollen Guides Epigenetic Inheritance via Small RNA. Cell 151,
194–205. https://doi.org/10.1016/j.cell.2012.09.001
Castel, S.E., Martienssen, R.A., 2013. RNA interference in the nucleus: roles for small RNAs
in transcription, epigenetics and beyond. Nat Rev Genet 14, 100–112.
https://doi.org/10.1038/nrg3355
Conway, E., Healy, E., Bracken, A.P., 2015. PRC2 mediated H3K27 methylations in cellular
identity and cancer. Curr Opin Cell Biol 37, 42–48.
https://doi.org/10.1016/j.ceb.2015.10.003
Dellaporta, S.L., Wood, J., Hicks, J.B., 1983. A plant DNA minipreparation: Version II. Plant
Mol Biol Rep 1, 19–21. https://doi.org/10.1007/BF02712670
Dorweiler, J.E., Carey, C.C., Kubo, K.M., Hollick, J.B., Kermicle, J.L., Chandler, V.L., 2000.
mediator of paramutation1 is required for establishment and maintenance of
paramutation at multiple maize loci. Plant Cell 12, 2101–2118.
https://doi.org/10.1105/tpc.12.11.2101
Du, J., Johnson, L.M., Groth, M., Feng, S., Hale, C.J., Li, S., Vashisht, A.A., Wohlschlegel,
J.A., Patel, D.J., Jacobsen, S.E., 2014. Mechanism of DNA methylation-directed
histone methylation by KRYPTONITE. Mol Cell 55, 495–504.
https://doi.org/10.1016/j.molcel.2014.06.009
Du, J., Zhong, X., Bernatavichute, Y.V., Stroud, H., Feng, S., Caro, E., Vashisht, A.A.,
Terragni, J., Chin, H.G., Tu, A., Hetzel, J., Wohlschlegel, J.A., Pradhan, S., Patel,
D.J., Jacobsen, S.E., 2012. Dual binding of chromomethylase domains to H3K9me2-
containing nucleosomes directs DNA methylation in plants. Cell 151, 167–180.
https://doi.org/10.1016/j.cell.2012.07.034
Egelhofer, T.A., Minoda, A., Klugman, S., Lee, K., Kolasinska-Zwierz, P., Alekseyenko,
A.A., Cheung, M.-S., Day, D.S., Gadel, S., Gorchakov, A.A., Gu, T., Kharchenko,
P.V., Kuan, S., Latorre, I., Linder-Basso, D., Luu, Y., Ngo, Q., Perry, M.,
Rechtsteiner, A., Riddle, N.C., Schwartz, Y.B., Shanower, G.A., Vielle, A., Ahringer,
J., Elgin, S.C.R., Kuroda, M.I., Pirrotta, V., Ren, B., Strome, S., Park, P.J., Karpen,
G.H., Hawkins, R.D., Lieb, J.D., 2011. An assessment of histone-modification
antibody quality. Nat Struct Mol Biol 18, 91–93. https://doi.org/10.1038/nsmb.1972
Erhard, K.F., Stonaker, J.L., Parkinson, S.E., Al., E., 2009. RNA Polymerase IV Functions in
Paramutation in Zea mays. Science 323, 1201–1205.
Ferrari, K.J., Scelfo, A., Jammula, S., Cuomo, A., Barozzi, I., Stützer, A., Fischle, W.,
Bonaldi, T., Pasini, D., 2014. Polycomb-dependent H3K27me1 and H3K27me2
regulate active transcription and enhancer fidelity. Mol Cell 53, 49–62.
https://doi.org/10.1016/j.molcel.2013.10.030
Fitz-James, M.H., Cavalli, G., 2022. Molecular mechanisms of transgenerational epigenetic
inheritance. Nat Rev Genet 23, 325–341. https://doi.org/10.1038/s41576-021-00438-5
Foda, B.M., Singh, U., 2015. Dimethylated H3K27 Is a Repressive Epigenetic Histone Mark
in the Protist Entamoeba histolytica and Is Significantly Enriched in Genes Silenced
via the RNAi Pathway. Journal of Biological Chemistry 290, 21114–21130.
https://doi.org/10.1074/jbc.M115.647263

Fu, F.-F., Dawe, R.K., Gent, J.I., 2018. Loss of RNA-Directed DNA Methylation in Maize
Chromomethylase and DDM1-Type Nucleosome Remodeler Mutants. The Plant Cell
30, 1617–1627. https://doi.org/10.1105/tpc.18.00053
Gent, Jonathan I, Ellis, N.A., Guo, L., Harkess, A.E., Yao, Y., Zhang, X., Dawe, R.K., 2013.
CHH islands : de novo DNA methylation in near-gene chromatin regulation in maize.
Genome research 23, 628–637. https://doi.org/10.1101/gr.146985.112.as
Gent, J I, Ellis, N.A., Guo, L., Harkess, A.E., Yao, Y., Zhang, X., Dawe, R.K., 2013. CHH
islands: de novo DNA methylation in near-gene chromatin regulation in maize.
Genome Research 23, 628–637. https://doi.org/10.1101/gr.146985.112
Gent, J.I., Higgins, K.M., Swentowsky, K.W., Fu, F.-F., Zeng, Y., Kim, D.W., Dawe, R.K.,
Springer, N.M., Anderson, S.N., 2022. The maize gene maternal derepression of r1
encodes a DNA glycosylase that demethylates DNA and reduces siRNA expression in
the endosperm. Plant Cell 34, 3685–3701. https://doi.org/10.1093/plcell/koac199
Gent, Jonathan I, Madzima, T.F., Bader, R., Kent, M.R., Zhang, X., Stam, M., McGinnis,
K.M., Dawe, R.K., 2014b. Accessible DNA and Relative Depletion of H3K9me2 at
Maize Loci Undergoing RNA-Directed DNA Methylation. The Plant Cell Online 26,
4903–4917. https://doi.org/10.1105/tpc.114.130427
Grewal, S.I.S., 2023. The molecular basis of heterochromatin assembly and epigenetic
inheritance. Mol Cell 83, 1767–1785. https://doi.org/10.1016/j.molcel.2023.04.020
Gruntman, E., Qi, Y., Slotkin, R.K., Roeder, T., Martienssen, R.A., Sachidanandam, R., 2008.
Kismeth: Analyzer of plant methylation states through bisulfite sequencing. BMC
Bioinformatics 9, 371. https://doi.org/10.1186/1471-2105-9-371
Gu, S.G., Pak, J., Guang, S., Maniar, J.M., Kennedy, S., Fire, A., 2012. Amplification of
siRNA in Caenorhabditis elegans generates a transgenerational sequence-targeted
histone H3 lysine 9 methylation footprint. Nat Genet 44, 157–164.
https://doi.org/10.1038/ng.1039
Guo, W., Fiziev, P., Yan, W., Cokus, S., Sun, X., Zhang, M.Q., Chen, P.Y., Pellegrini, M.,
2013. BS-Seeker2: A versatile aligning pipeline for bisulfite sequencing data. BMC
Genomics 14, 1. https://doi.org/10.1186/1471-2164-14-774
Guo, W., Zhu, P., Pellegrini, M., Zhang, M.Q., Wang, X., Ni, Z., 2018. CGmapTools
improves the precision of heterozygous SNV calls and supports allele-specific
methylation detection and visualization in bisulfite-sequencing data. Bioinformatics
34, 381–387. https://doi.org/10.1093/bioinformatics/btx595
Gutiérrez-Marcos, J.F., Costa, L.M., Dal Prà, M., Scholten, S., Kranz, E., Perez, P.,
Dickinson, H.G., 2006. Epigenetic asymmetry of imprinted genes in plant gametes.
Nat Genet 38, 876–878. https://doi.org/10.1038/ng1828
Haag, J.R., Brower-Toland, B., Krieger, E.K., Sidorenko, L., Nicora, C.D., Norbeck, A.D.,
Irsigler, A., LaRue, H., Brzeski, J., McGinnis, K., Ivashuta, S., Pasa-Tolic, L.,
Chandler, V.L., Pikaard, C.S., 2014. Functional diversification of maize RNA
polymerase IV and V subtypes via alternative catalytic subunits. Cell reports 9, 378–
90. https://doi.org/10.1016/j.celrep.2014.08.067
Haring, M., Bader, R., Louwers, M., Schwabe, A., van Driel, R., Stam, M., 2010a. The role of
DNA methylation, nucleosome occupancy and histone modifications in paramutation.
The Plant journal : for cell and molecular biology 63, 366–78.
https://doi.org/10.1111/j.1365-313X.2010.04245.x
Haring, M., Offermann, S., Danker, T., Horst, I., Peterhansel, C., Stam, M., 2007. Chromatin
immunoprecipitation: optimization, quantitative analysis and data normalization. Plant
Methods 3, 11. https://doi.org/10.1186/1746-4811-3-11

Herr, A.J., Jensen, M.B., Dalmay, T., Baulcombe, D.C., 2005. RNA polymerase IV directs
silencing of endogenous DNA. Science 308, 118–120.
https://doi.org/10.1126/science.1106910
Hollick, J.B., 2017. Paramutation and related phenomena in diverse species. Nat Rev Genet
18, 5–23. https://doi.org/10.1038/nrg.2016.115
Hollick, Jay B, Kermicle, J.L., Parkinson, S.E., 2005. Rmr6 maintains meiotic inheritance of
paramutant states in Zea mays. Genetics 171, 725–40.
https://doi.org/10.1534/genetics.105.045260
Hollick, J. B., Kermicle, J.L., Parkinson, S.E., 2005. Rmr6 maintains meiotic inheritance of
paramutant states in Zea mays. Genetics 171, 725–740.
https://doi.org/10.1534/genetics.105.045260
Hollick, J.B., Patterson, G.I., Coe, E.H., Cone, J.K.C., Chandler, V.L., 1995. Allelic
Interactions Heritably Alter the Activity of a Metastable Maize pl Allele. Genetics
141, 709–719.
Hövel, I., Pearson, N.A., Stam, M., 2015. Cis-acting determinants of paramutation. Semin
Cell Dev Biol 44, 22–32. https://doi.org/10.1016/j.semcdb.2015.08.012
Hufford, M.B., Seetharam, A.S., Woodhouse, M.R., Chougule, K.M., Ou, S., Liu, J., Ricci,
W.A., Guo, T., Olson, A., Qiu, Y., Della Coletta, R., Tittes, S., Hudson, A.I., Marand,
A.P., Wei, S., Lu, Z., Wang, B., Tello-Ruiz, M.K., Piri, R.D., Wang, N., Kim, D.W.,
Zeng, Y., O'Connor, C.H., Li, X., Gilbert, A.M., Baggs, E., Krasileva, K.V.,
Portwood, J.L., Cannon, E.K.S., Andorf, C.M., Manchanda, N., Snodgrass, S.J.,
Hufnagel, D.E., Jiang, Q., Pedersen, S., Syring, M.L., Kudrna, D.A., Llaca, V.,
Fengler, K., Schmitz, R.J., Ross-Ibarra, J., Yu, J., Gent, J.I., Hirsch, C.N., Ware, D.,
Dawe, R.K., 2021. De novo assembly, annotation, and comparative analysis of 26
diverse maize genomes. Science 373, 655–662.
https://doi.org/10.1126/science.abg5289
Jackel, J.N., Storer, J.M., Coursey, T., Bisaro, D.M., 2016. Arabidopsis RNA Polymerases IV
and V Are Required To Establish H3K9 Methylation, but Not Cytosine Methylation,
on Geminivirus Chromatin. Journal of Virology 90, 7529–7540.
https://doi.org/10.1128/jvi.00656-16
Jia, Y., Lisch, D.R., Ohtsu, K., Scanlon, M.J., Nettleton, D., Schnable, P.S., 2009. Loss of
RNA-dependent RNA polymerase 2 (RDR2) function causes widespread and
unexpected changes in the expression of transposons, genes, and 24-nt small RNAs.
PLoS Genet 5, e1000737. https://doi.org/10.1371/journal.pgen.1000737
Jiang, C., Pugh, B.F., 2009. Nucleosome positioning and gene regulation: advances through
genomics. Nat Rev Genet 10, 161–172. https://doi.org/10.1038/nrg2522
Johnson, L.M., Bostick, M., Zhang, X., Kraft, E., Henderson, I., Callis, J., Jacobsen, S.E.,
2007. The SRA methyl-cytosine-binding domain links DNA and histone methylation.
Curr Biol 17, 379–384. https://doi.org/10.1016/j.cub.2007.01.009
Kanno, T., Huettel, B., Mette, M.F., Aufsatz, W., Jaligot, E., Daxinger, L., Kreil, D.P.,
Matzke, M., Matzke, A.J.M., 2005. Atypical RNA polymerase subunits required for
RNA-directed DNA methylation. Nat Genet 37, 761–765.
https://doi.org/10.1038/ng1580
Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods
9, 357–359. https://doi.org/10.1038/nmeth.1923
Law, J.A., Vashisht, A.A., Wohlschlegel, J.A., Jacobsen, S.E., 2011. SHH1, a homeodomain
protein required for DNA methylation, as well as RDR2, RDM4, and chromatin
remodeling factors, associate with RNA polymerase IV. PLoS Genet 7, e1002195.
https://doi.org/10.1371/journal.pgen.1002195

Lee, H.-G., Kahn, T.G., Simcox, A., Schwartz, Y.B., Pirrotta, V., 2015. Genome-wide
activities of Polycomb complexes control pervasive transcription. Genome Res. 25,
1170–1181. https://doi.org/10.1101/gr.188920.114
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R., 1000 Genome Project Data Processing Subgroup, 2009. The Sequence
Alignment/Map format and SAMtools. Bioinformatics 25, 2078–2079.
https://doi.org/10.1093/bioinformatics/btp352
Li, Q., Eichten, S.R., Hermanson, P.J., Zaunbrecher, V.M., Song, J., Wendt, J., Rosenbaum,
H., Madzima, T.F., Sloan, A.E., Huang, J., Burgess, D.L., Richmond, T.A., McGinnis,
K.M., Meeley, R.B., Danilevskaya, O.N., Vaughn, M.W., Kaeppler, S.M., Jeddeloh,
J.A., Springer, N.M., 2014. Genetic Perturbation of the Maize Methylome. The Plant
Cell Online 26, 4602–4616. https://doi.org/10.1105/tpc.114.133140
Li, Q., Gent, J.I., Zynda, G., Song, J., Makarevitch, I., Hirsch, C.D., Hirsch, C.N., Dawe,
R.K., Madzima, T.F., McGinnis, K.M., Lisch, D., Schmitz, R.J., Vaughn, M.W.,
Springer, N.M., 2015. RNA-directed DNA methylation enforces boundaries between
heterochromatin and euchromatin in the maize genome. Proceedings of the National
Academy of Sciences 112, 14728–14733. https://doi.org/10.1073/pnas.1514680112
Lindroth, A.M., Cao, X., Jackson, J.P., Zilberman, D., McCallum, C.M., Henikoff, S.,
Jacobsen, S.E., 2001. Requirement of CHROMOMETHYLASE3 for maintenance of
CpXpG methylation. Science 292, 2077–2080.
https://doi.org/10.1126/science.1059745
Liu, Y., Taverna, S.D., Muratore, T.L., Shabanowitz, J., Hunt, D.F., Allis, C.D., 2007. RNAi-
dependent H3K27 methylation is required for heterochromatin formation and DNA
elimination in Tetrahymena. Genes Dev 21, 1530–1545.
https://doi.org/10.1101/gad.1544207
Liu, Y., Yuan, J., Jia, G., Ye, W., Jeffrey Chen, Z., Song, Q., 2021. Histone H3K27
dimethylation landscapes contribute to genome stability and genetic recombination
during wheat polyploidization. The Plant Journal 105, 678–690.
https://doi.org/10.1111/tpj.15063
Locatelli, S., Piatti, P., Motto, M., Rossi, V., 2009. Chromatin and DNA modifications in the
Opaque2-mediated regulation of gene transcription during maize endosperm
development. Plant Cell 21, 1410–1427. https://doi.org/10.1105/tpc.109.067256
Louwers, M., Bader, R., Haring, M., van Driel, R., de Laat, W., Stam, M., 2009a. Tissue- and
expression level-specific chromatin looping at maize b1 epialleles. The Plant Cell 21,
832–42. https://doi.org/10.1105/tpc.108.064329
Louwers, M., Splinter, E., van Driel, R., de Laat, W., Stam, M., 2009b. Studying physical
chromatin interactions in plants using Chromosome Conformation Capture (3C). Nat
Protoc 4, 1216–1229. https://doi.org/10.1038/nprot.2009.113
Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet.journal 17, 10–12. https://doi.org/10.14806/ej.17.1.200
Mathieu, O., Probst, A.V., Paszkowski, J., 2005. Distinct regulation of histone H3
methylation at lysines 27 and 9 by CpG methylation in Arabidopsis. The EMBO
Journal 24, 2783–2791. https://doi.org/10.1038/sj.emboj.7600743
Mathieu, O., Reinders, J., Caikovski, M., Smathajitt, C., Paszkowski, J., 2007.
Transgenerational stability of the Arabidopsis epigenome is coordinated by CG
methylation. Cell 130, 851–862. https://doi.org/10.1016/j.cell.2007.07.007
Matzke, M.A., Mosher, R.A., 2014. RNA-directed DNA methylation: an epigenetic pathway
of increasing complexity. Nat Rev Genet 15, 394–408.
https://doi.org/10.1038/nrg3683

Mishra, V., Singh, J., Wang, F., Zhang, Y., Fukudome, A., Trinidad, J.C., Takagi, Y.,
Pikaard, C.S., 2021. Assembly of a dsRNA synthesizing complex: RNA-
DEPENDENT RNA POLYMERASE 2 contacts the largest subunit of NUCLEAR
RNA POLYMERASE IV. Proc Natl Acad Sci U S A 118, e2019276118.
https://doi.org/10.1073/pnas.2019276118

Nobuta, K., Lu, C., Shrivastava, R., Pillay, M., De Paoli, E., Accerbi, M., Arteaga-Vazquez,
M., Sidorenko, L., Jeong, D.-H., Yen, Y., Green, P.J., Chandler, V.L., Meyers, B.C.,
2008a. Distinct size distribution of endogeneous siRNAs in maize: Evidence from
deep sequencing in the mop1-1 mutant. Proc Natl Acad Sci U S A 105, 14958–14963.
https://doi.org/10.1073/pnas.0808066105

Nobuta, K., Lu, C., Shrivastava, R., Pillay, M., Paoli, E. De, Accerbi, M., Arteaga-vazquez,
M., Sidorenko, L., Jeong, D., Yen, Y., Green, P.J., Chandler, V.L., Meyers, B.C.,
2008b. Distinct size distribution of endogenous siRNAs in maize : Evidence from
deep sequencing in the mop1-1 mutant. PNAS 105, 14958–14963.

Ohtsu, K., Smith, M.B., Emrich, S.J., Borsuk, L.A., Zhou, R., Chen, T., Zhang, X.,
Timmermans, M.C.P., Beck, J., Buckner, B., Janick-Buckner, D., Nettleton, D.,
Scanlon, M.J., Schnable, P.S., 2007. Global gene expression analysis of the shoot
apical meristem of maize (Zea mays L.). Plant J 52, 391–404.
https://doi.org/10.1111/j.1365-313X.2007.03244.x

Onodera, Y., Haag, J.R., Ream, T., Costa Nunes, P., Pontes, O., Pikaard, C.S., 2005. Plant
nuclear RNA polymerase IV mediates siRNA and DNA methylation-dependent
heterochromatin formation. Cell 120, 613–622.
https://doi.org/10.1016/j.cell.2005.02.007

Pal-Bhadra, M., Leibovitch, B.A., Gandhi, S.G., Rao, M., Bhadra, U., Birchler, J.A., Elgin,
S.C.R., 2004. Heterochromatic Silencing and HP1 Localization in Drosophila Are
Dependent on the RNAi Machinery. Science 303, 669–672.
https://doi.org/10.1126/science.1092653

Papa, C.M., Springer, N.M., Muszynski, M.G., Meeley, R., Kaeppler, S.M., 2001. Maize
chromomethylase Zea methyltransferase2 is required for CpNpG methylation. Plant
Cell 13, 1919–1928. https://doi.org/10.1105/tpc.010064

Parent, J.-S., Cahn, J., Herridge, R.P., Grimanelli, D., Martienssen, R.A., 2021. Small RNAs
guide histone methylation in Arabidopsis embryos. Genes Dev. 35, 841–846.
https://doi.org/10.1101/gad.343871.120

Park, S., Oh, S., Nocker, S. van, 2012. Genomic and Gene-Level Distribution of Histone H3
Dimethyl Lysine-27 (H3K27me2) in Arabidopsis. PLOS ONE 7, e52855.
https://doi.org/10.1371/journal.pone.0052855

Patterson, G.I., Thorpe, C.J., Chandler, V.L., 1993. Paramutation, an allelic interaction, is
associated with a stable and heritable reduction of transcription of the maize b
regulatory gene. Genetics 135, 881–894. https://doi.org/10.1093/genetics/135.3.881

Plank, J.L., Dean, A., 2014. Enhancer function: mechanistic and genome-wide insights come
together. Mol Cell 55, 5–14. https://doi.org/10.1016/j.molcel.2014.06.015

Pontier, D., Yahubyan, G., Vega, D., Bulski, A., Saez-Vasquez, J., Hakimi, M.-A., Lerbs-
Mache, S., Colot, V., Lagrange, T., 2005. Reinforcement of silencing at transposons
and highly repeated sequences requires the concerted action of two distinct RNA
polymerases IV in Arabidopsis. Genes Dev 19, 2030–2040.
https://doi.org/10.1101/gad.348405

Quinlan, A.R., Hall, I.M., 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics 26, 841–842. https://doi.org/10.1093/bioinformatics/btq033

Ritchie, S.W., Hanway, J.J., Benson, G.O., 1986. How a corn plant develops, 1986
(Departmental Report).

Ronsseray, S., 2015. Paramutation phenomena in non-vertebrate animals. Semin Cell Dev
Biol 44, 39–46. https://doi.org/10.1016/j.semcdb.2015.08.009
Roudier, F., Ahmed, I., Bérard, C., Sarazin, A., Mary-Huard, T., Cortijo, S., Bouyer, D.,
Caillieux, E., Duvernois-Berthet, E., Al-Shikhley, L., Giraut, L., Després, B.,
Drevensek, S., Barneche, F., Dèrozier, S., Brunaud, V., Aubourg, S., Schnittger, A.,
Bowler, C., Martin-Magniette, M.-L., Robin, S., Caboche, M., Colot, V., 2011.
Integrative epigenomic mapping defines four main chromatin states in Arabidopsis:
Organization of the Arabidopsis epigenome. The EMBO Journal 30, 1928–1938.
https://doi.org/10.1038/emboj.2011.103
Rymen, B., Ferrafiat, L., Blevins, T., 2020. Non-coding RNA polymerases that silence
transposable elements and reprogram gene expression in plants. Transcription 11,
172–191. https://doi.org/10.1080/21541264.2020.1825906
Schmitz, R.J., Grotewold, E., Stam, M., 2022. Cis-regulatory sequences in plants: Their
importance, discovery, and future challenges. Plant Cell 34, 718–741.
https://doi.org/10.1093/plcell/koab281
Sekhon, R.S., Lin, H., Childs, K.L., Hansey, C.N., Buell, C.R., de Leon, N., Kaeppler, S.M.,
2011. Genome-wide atlas of transcription during maize development. Plant J 66, 553–
563. https://doi.org/10.1111/j.1365-313X.2011.04527.x
Sekhon, R.S., Wang, P.-H., Sidorenko, L., Chandler, V.L., Chopra, S., 2012. Maize Unstable
factor for orange1 is required for maintaining silencing associated with paramutation
at the pericarp color1 and booster1 loci. PLoS Genet 8, e1002980.
https://doi.org/10.1371/journal.pgen.1002980
Sidorenko, L., Chandler, V., 2008. RNA-dependent RNA polymerase is required for
enhancer-mediated transcriptional silencing associated with paramutation at the maize
p1 gene. Genetics 180, 1983–1993. https://doi.org/10.1534/genetics.108.095281
Sidorenko, L., Dorweiler, J.E., Cigan, a M., Arteaga-Vazquez, M., Vyas, M., Kermicle, J.,
Jurcin, D., Brzeski, J., Cai, Y., Chandler, V.L., 2009. A dominant mutation in
mediator of paramutation2, one of three second-largest subunits of a plant-specific
RNA polymerase, disrupts multiple siRNA silencing processes. PLoS genetics 5,
e1000725. https://doi.org/10.1371/journal.pgen.1000725
Sloan, Amy E., Sidorenko, L., McGinnis, K.M., 2014. Diverse gene-silencing mechanisms
with distinct requirements for RNA polymerase subunits in Zea mays. Genetics 198,
1031–1042. https://doi.org/10.1534/genetics.114.168518
Springer, N.M., McGinnis, K.M., 2015. Paramutation in evolution, population genetics and
breeding. Semin Cell Dev Biol 44, 33–38.
https://doi.org/10.1016/j.semcdb.2015.08.010
Stam, M., Belele, C., Dorweiler, J.E., Chandler, V.L., 2002a. Differential chromatin structure
within a tandem array 100 kb upstream of the maize b1 locus is associated with
paramutation. Genes and Development 16, 1906–1918.
https://doi.org/10.1101/gad.1006702
Stam, M., Belele, C., Ramakrishna, W., Dorweiler, J.E., Bennetzen, J.L., Chandler, V.L.,
2002b. The regulatory regions required for B' paramutation and expression are located
far upstream of the maize b1 transcribed sequences. Genetics 162, 917–930.
https://doi.org/10.1093/genetics/162.2.917
Stonaker, Jennifer L., Lim, J.P., Erhard, K.F., Hollick, J.B., 2009. Diversity of Pol IV
function is defined by mutations at the maize rmr7 locus. PLoS Genet 5, e1000706.
https://doi.org/10.1371/journal.pgen.1000706
Stroud, H., Do, T., Du, J., Zhong, X., Feng, S., Johnson, L., Patel, D.J., Jacobsen, S.E., 2014.
Non-CG methylation patterns shape the epigenetic landscape in Arabidopsis. Nat
Struct Mol Biol 21, 64–72. https://doi.org/10.1038/nsmb.2735

Stroud, H., Greenberg, M.V.C., Feng, S., Bernatavichute, Y.V., Jacobsen, S.E., 2013. Comprehensive analysis of silencing mutants reveals complex regulation of the Arabidopsis methylome. Cell 152, 352–364. https://doi.org/10.1016/j.cell.2012.10.054

Struhl, K., Segal, E., 2013. Determinants of nucleosome positioning. Nat Struct Mol Biol 20, 267–273. https://doi.org/10.1038/nsmb.2506

Tang, K., Lang, Z., Zhang, H., Zhu, J.-K., 2016. The DNA demethylase ROS1 targets genomic regions with distinct chromatin modifications. Nat Plants 2, 16169. https://doi.org/10.1038/nplants.2016.169

Underwood, C.J., Choi, K., Lambing, C., Zhao, X., Serra, H., Borges, F., Simorowski, J., Ernst, E., Jacob, Y., Henderson, I.R., Martienssen, R.A., 2018. Epigenetic activation of meiotic recombination near Arabidopsis thaliana centromeres via loss of H3K9me2 and non-CG DNA methylation. Genome Res 28, 519–531. https://doi.org/10.1101/gr.227116.117

Verdel, A., Jia, S., Gerber, S., Sugiyama, T., Gygi, S., Grewal, S.I.S., Moazed, D., 2004. RNAi-mediated targeting of heterochromatin by the RITS complex. Science 303, 672–676. https://doi.org/10.1126/science.1093686

West, P.T., Li, Q., Ji, L., Eichten, S.R., Song, J., Vaughn, M.W., Schmitz, R.J., Springer, N.M., 2014. Genomic distribution of H3K9me2 and DNA methylation in a maize genome. PLoS One 9, e105267. https://doi.org/10.1371/journal.pone.0105267

Wierzbicki, A.T., Haag, J.R., Pikaard, C.S., 2008. Noncoding transcription by RNA polymerase Pol IVb/Pol V mediates transcriptional silencing of overlapping and adjacent genes. Cell 135, 635–648. https://doi.org/10.1016/j.cell.2008.09.035

Woodhouse, M.R., Freeling, M., Lisch, D., 2006a. Initiation, establishment, and maintenance of heritable MuDR transposon silencing in maize are mediated by distinct factors. PLoS Biol 4, e339. https://doi.org/10.1371/journal.pbio.0040339

Woodhouse, M.R., Freeling, M., Lisch, D., 2006b. The mop1 (mediator of paramutation1) mutant progressively reactivates one of the two genes encoded by the MuDR transposon in maize. Genetics 172, 579–592. https://doi.org/10.1534/genetics.105.051383

Xie, W., Schultz, M.D., Lister, R., Hou, Z., Rajagopal, N., Ray, P., Whitaker, J.W., Tian, S., Hawkins, R.D., Leung, D., Yang, H., Wang, T., Lee, A.Y., Swanson, S.A., Zhang, J., Zhu, Y., Kim, A., Nery, J.R., Urich, M.A., Kuan, S., Yen, C., Klugman, S., Yu, P., Suknuntha, K., Propson, N.E., Chen, H., Edsall, L.E., Wagner, U., Li, Y., Ye, Z., Kulkarni, A., Xuan, Z., Chung, W.-Y., Chi, N.C., Antosiewicz-Bourget, J.E., Slukvin, I., Stewart, R., Zhang, M.Q., Wang, W., Thomson, J.A., Ecker, J.R., Ren, B., 2013. Epigenomic analysis of multilineage differentiation of human embryonic stem cells. Cell 153, 1134–1148. https://doi.org/10.1016/j.cell.2013.04.022

Zemach, A., Kim, M.Y., Hsieh, P.-H., Coleman-Derr, D., Eshed-Williams, L., Thao, K., Harmer, S.L., Zilberman, D., 2013. The Arabidopsis Nucleosome Remodeler DDM1 Allows DNA Methyltransferases to Access H1-Containing Heterochromatin. Cell 153, 193–205. https://doi.org/10.1016/j.cell.2013.02.033

Zofall, M., Sandhu, R., Holla, S., Wheeler, D., Grewal, S.I.S., 2022. Histone deacetylation primes self-propagation of heterochromatin domains to promote epigenetic inheritance. Nat Struct Mol Biol 29, 898–909. https://doi.org/10.1038/s41594-022-00830-7

Table S1. Primers used in ChIP-qPCR experiments.

Gene	Primer set	Sequence	Position relative to TSS	Primer name
b1	5'R	5' CTC GGG GTC AAA TGG ACG G 3'	107.0 kb upstream Fw	M85
		5' GCG GTC ACA ACC TTT TCA GAT C 3'	106.9 kb upstream Rv	M317
	R1	5' GGT TTG CTG CAT CCT TGA CC 3'	~ 100 kb upstream Fw	M12
		5' CTG CAC AGG CTG CAG CAG 3'	~ 100 kb upstream Rv	M295
	R2	5'-GTG GAC AAA TAG TGC ATT CAC CTC AC-3'	~ 100 kb upstream Fw	M111
		5'-GAA ACT GAA CTA ACC AAT GTG GCT-3'	~ 100 kb upstream Rv	M296
	R3	5'-CAA GAT CCA TTG AAC ATC TTG TCC-3'	~ 100 kb upstream Fw	M297
		5'-CAT GTG TGA GGG TGA TGC TGC G-3'	~ 100 kb upstream Rv	M10
	R4	5'-CTA TGC TCG TGC GAA GGG TGG TC-3'	~ 100 kb upstream Fw	M11
		5'-CAG ATC CCC TAT CTA ATC CCA TGT-3'	~ 100 kb upstream Rv	M298
	R5	5'-GCG GGT GAC CAG TGT TT-3'	~ 100 kb upstream Fw	M668
		5'-GAC CTC TTG GAG CCC TCT AAT TTG-3'	~ 100 kb upstream Rv	M669
	R6	5'-CAA ATT AGA GGG CTC CAA GAG GTC-3'	~ 100 kb upstream Fw	M112
		5'-GTT GTG TAC TGC AGT GTT AGG TAG-3'	~ 100 kb upstream Rv	M15
	e	5' CTG GCG GCA CTA AAA AAC G 3'	45.3 kb upstream Fw	M975
		5' TGT GCC CAC CTT TAT TGT GAG TT 3'	45.2 kb upstream Rv	M976
	h	5' ACC AGT CCA ACC TAG CTT GGA AT 3'	41.9 kb upstream Fw	M611
		5' GTC ATT GCG TGC GGC TAT C 3'	41.8 kb upstream Rv	M334
	l	5' TCC ATC TCA CCT CTC ATT GTA TCT TT 3'	8.6 kb upstream Fw	M984
		5' AAA TCG TGT ATA TGG TCG TTT AAA ACA T 3'	8.5 kb upstream Rv	M985
	5'b	5'-GGT GTG CAC ACC ATT AAT TGA-3'	1.2 kb upstream Fw	M119
		5'-CGA TAT TTT GGT GAA AAC TGT TC-3'	1.0 kb upstream Rv	M102
	UTR	5' CTC TTC TGA TCT TCT TCA CCG TCT C 3'	0.4 kb downstream Fw	M108
		5' ATT CCC GGG CGG CCG CGC CTT ACT AAT CCT TC 3'	0.5 kb downstream Rv	M44
	Ex3	5' ACC AGC TGC TCA TGC AGA GGA 3'	1.3 kb downstream Fw	M120
		5' AGG AAG GCG TAG GTC ATG CAG AT 3'	1.4 kb downstream Rv	M121
Actin		5'-GGT GTG CAC ACC ATT AAT TGA-3'	coding region of Actin	M70
		5'-CAC TTT CTG CTC ATG GTT TAA GG-3'	coding region of Actin	M414
Copia		5'-CGA TGT GAA GAC AGC ATT CCT-3'	coding region of Copia	M688
		5'-CTC AAG TGA CAT CCC ATG TGT-3'	coding region of Copia	M689

 Figure legends

 Figure 1. *b1* expression levels in maize husk tissues. RNA blot analysis of RNA from *B'*, *B-I*

 and *B' mop1-1, B' Mop2/mop2-1, B' mop3-1* husk tissue, using probes recognizing the coding

 region of *b1* (exons 7-9) and *Sam* (see Supplemental Figure S2B). The intensities of the bands

 representing full-length transcripts were quantified. The indicated values represent the

 average plus standard deviation of *b1* transcript levels normalized to *Sam* transcript levels.

 Figure 2. The regulatory hepta-repeat is activated in *mop1-1* and *mop3-1*, but not in *mop2*

 mutants. A) Schematic representation of the *b1* locus including coding region (white box) and

 hepta-repeat enhancer (arrowheads). Regions monitored in ChIP are indicated below in green.

 The fragments (I-XII) examined by 3C (see Figure 3) are indicated as dark grey boxes. B)

 ChIP-qPCR experiments were performed on husk tissue from *B'*, *B' mop1-1*, *B' Mop/mop2-1*,

 B' mop2-1, *B' mop3-1* and *B-I* plants with an antibody recognizing H3ac. ChIP signals were

 normalized to *actin* ChIP signals. Error bars indicate standard error of the mean (SEM) of

 three *(B', B' mop1-1, B' Mop/mop2-1)*, or four (*B' mop2-1*, *B' mop3-I, B-I)* biological

 replicates. See Supplemental Table S2 for summary of ChIP statistics.

 Figure 3. *B'* shows a multi-loop structure in the *mop1-1* and *mop3-1* mutant background, and

 a single loop structure in *Mop2/mop2-1* mutant background. At the top a schematic

 representation of the *b1* locus including the coding region (white box) and hepta-repeat

 enhancer (arrowheads). The *Bgl*II fragments (I-XII) examined by 3C are indicated as dark

 grey boxes. The viewpoints I (TSS; panel A), X (Hepta-repeat; panel B), and VII (~47kb

 upstream; panel C) are indicated by black vertical bars. Data were normalized using

 crosslinking frequencies measured for the *Sam* locus (Supplementary Figure S2). Error bars

 indicate the SEM of four biological replicates each for *B' mop1-1*, *B' Mop2/mop2-1* and *B'*

 mop3-1. Data for *B'* and *B-I* (reproduced from Louwers et al., 2009) are shown in shading for

 comparison. The experiments for *B'*, *B-I*, and the *mop* mutants were done in parallel.

 Figure 4. Differentially methylated repeat junction region retains high levels of symmetric

 DNA methylation in *mop* mutants. A) Summary of the DNA methylation levels data obtained

 by DNA blot analysis. DNA derived from *B'* wild type, *B-I* and *B' mop* mutants was digested

 with the methylation-insensitive *Eco*RI or *Bam*HI and the methylation-sensitive enzymes

 indicated. A zoom of part of the hepta-repeat, spanning the repeat junction region, is shown.

 The consensus repeat methylation profiles of *B'* and *B-I* (Haring et al., 2010) are shown. DNA

methylation levels are specified by colors: 0-12.5%, white; 12.5-37.5%, yellow; 37.5-62.5%,

orange; 62.5-87.5%, red; 87.5-100% methylation, dark red. H, *Hpa*I; P, *Pst*I; E, *Hha*I and

*Hae*II; A, *Alu*I; S, *Sau*96I; U, *Sau*3AI; B, *Bsm*AI; Numbers discriminate individual sites

present more than once every repeat. For details on the quantification of the DNA methylation

levels see Haring *et al.* (2010). B) DNA methylation profiles of the *b1* repeat junction region

at base-pair resolution for *B'* and *B-I*, generated by targeted bisulfite sequencing. This region

includes the R1 and R6 loci assayed by ChIP-qPCR. See Supplementary Fig. S8 for all data

sets. The dot-plots show the DNA methylation for 30 and 31 individual *B'* and *B-I* clones,

respectively. Methylated and unmethylated cytosines are represented by filled and empty

circles, respectively. Red, CG; Blue, CHG; Green, CHH. The restriction sites examined by

DNA blotting are indicated above the panel for comparison. E indicates a site cut by both

*Hha*I and *Hae*II. C) Total fraction of methylated cytosines in the repeat junction region in all

three sequence contexts as measured by bisulfite sequencing. The mean and standard

deviations shown are calculated from three (*B' mop1-1*) or two (all other genotypes)

biological replicates. *M/mop2* indicates *Mop2/mop2-1.*

1295

1296   **Figure 5.** The maize genome contains low-mCHH and high-mCHH RdDM loci. Average

1297   DNA methylation levels for high and low mCHH loci in the developing second leaf of B73.

1298   Values of 5-methylcytosine are relative to the total number of cytosines (mC/total C) in each

1299   sequence context (CG, CHG, or CHH).

1300

Figure 6. No significant decrease in nucleosome occupancy was observed in the *mop*

mutants. ChIP experiments were performed with an antibody recognizing histone H3, using

husk tissue from *B'* (light green), *B' Mop/mop2-1* (green), *B' mop2-1* (black), *B' mop1-1*

(magenta)*, B' mop3-1* (blue), and *B-I* (purple) plants. Data was normalized to *actin* values.

The error bars indicate the standard error of the mean (SEM) of two (*B' mop1-1*), three (*B'*,

B' Mop/mop2-1, *B' mop3-1),* or four (*B' mop2-1*, *B-I*) replicate experiments. For the no-

antibody control signals see Supplemental Figure S10.

1308

Figure 7. Transcriptional activation in *mop1-1* and *mop3-1* is associated with reduced levels

of repressive histone marks. ChIP-qPCR experiments were performed on seedling (A, C) and

husk tissue (B, D) from *B'*, *B' Mop/mop2-1*, *B' mop2-1*, *B' mop1-1*, *B' mop3-1* and *B-I* plants

with antibodies recognizing H3K9me2 (A, B), and H3K27me2 (C, D). ChIP signals were

normalized to *Copia* signals. Error bars indicate the SEM of multiple biological replicates.

For seedlings, three (*B', B' mop1-1*) or two (*B' mop3-1, B-I)* replicates were performed. For

husks (B, D), two (*B' mop2-2*), three (*B', B' mop3-1, B-I*), four (*B' mop1-1, B' Mop/mop2-1*,

B' mop2-1) replicates were performed. For the no-antibody control signals see Supplemental

Figure S11.

1318

Figure 1

Figure 1. *b1* expression levels in maize husk tissues. RNA blot analysis of RNA from *B'*, *B-I* and *B' mop1-1*, *B' Mop2/mop2-1*, *B' mop3-1* husk tissue, using probes recognizing the coding region of *b1* (exons 7-9) and *Sam* (see Supplemental Figure S2B). The intensities of the bands representing full-length transcripts were quantified. The indicated values represent the average plus standard deviation of *b1* transcript levels normalized to *Sam* transcript levels.

Figure 2

Figure 2. The regulatory hepta-repeat is activated in *mop1-1* and *mop3-1*, but not in *mop2* mutants. A) Schematic representation of the *b1* locus including coding region (white box) and hepta-repeat enhancer (arrowheads). Regions monitored in ChIP are indicated below in green. The fragments (I-XII) examined by 3C (see Figure 3) are indicated as dark grey boxes. B) ChIP-qPCR experiments were performed on husk tissue from *B'*, *B' mop1-1*, *B' Mop/mop2-1*, *B' mop2-1*, *B' mop3-1* and *B-I* plants with an antibody recognizing H3ac. ChIP signals were normalized to *actin* ChIP signals. Error bars indicate standard error of the mean (SEM) of three *(B', B' mop1-1, B' Mop/mop2-1)*, or four *(B' mop2-1, B' mop3-I, B-I)* biological replicates. See Supplemental Table S2 for summary of ChIP statistics.

Figure 3

Figure 3. *B'* shows a multi-loop structure in the *mop1-1* and *mop3-1* mutant background, and a single loop structure in *Mop2/mop2-1* mutant background. At the top a schematic representation of the *b1* locus including the coding region (white box) and hepta-repeat enhancer (arrowheads). The *Bgl*II fragments (I-XII) examined by 3C are indicated as dark grey boxes. The viewpoints I (TSS; panel A), X (Hepta-repeat; panel B), and VII (~47kb upstream; panel C) are indicated by black vertical bars. Data were normalized using crosslinking frequencies measured for the *Sam* locus (Supplementary Figure S2). Error bars indicate the SEM of four biological replicates each for *B' mop1-1*, *B' Mop2/mop2-1* and *B' mop3-1*. Data for *B'* and *B-I* (reproduced from Louwers et al., 2009) are shown in shading for comparison. The experiments for *B'*, *B-I*, and the *mop* mutants were done in parallel.

Figure 4

Figure 4. Differentially methylated repeat junction region retains high levels of symmetric DNA methylation in *mop* mutants. A) Summary of the DNA methylation levels data obtained by DNA blot analysis. DNA derived from *B'* wild type, *B-I* and *B' mop* mutants was digested with the methylation-insensitive *Eco*RI or *Bam*HI and the methylation-sensitive enzymes indicated. A zoom of part of the hepta-repeat, spanning the repeat junction region, is shown. The consensus repeat methylation profiles of *B'* and *B-I* (Haring et al., 2010) are shown. DNA methylation levels are specified by colors: 0-12.5%, white; 12.5-37.5%, yellow; 37.5-62.5%, orange; 62.5-87.5%, red; 87.5-100% methylation, dark red. H, *Hpa*I; P, *Pst*I; E, *Hha*I and *Hae*II; A, *Alu*I; S, *Sau*96I; U, *Sau*3AI; B, *Bsm*AI; Numbers discriminate individual sites present more than once every repeat. For details on the quantification of the DNA methylation levels see Haring *et al.* (2010). B) DNA methylation profiles of the *b1* repeat junction region at base-pair resolution for *B'* and *B-I*, generated by targeted bisulfite sequencing. This region includes the R1 and R6 loci assayed by ChIP-qPCR. See Supplementary Fig. S8 for all data sets. The dot-plots show the DNA methylation for 30 and 31 individual *B'* and *B-I* clones, respectively. Methylated and unmethylated cytosines are represented by filled and empty circles, respectively. Red, CG; Blue, CHG; Green, CHH. The restriction sites examined by DNA blotting are indicated above the panel for comparison. E indicates a site cut by both *Hha*I and *Hae*II. C) Total fraction of methylated cytosines in the repeat junction region in all three sequence contexts as measured by bisulfite sequencing. The mean and standard deviations shown are calculated from three (*B' mop1-1*) or two (all other genotypes) biological replicates. *M/mop2* indicates *Mop2/mop2-1*.

Figure 5

Figure 5. The maize genome contains low-mCHH and high-mCHH RdDM loci. Average DNA methylation levels for high and low mCHH loci in the developing second leaf of B73. Values of 5-methylcytosine are relative to the total number of cytosines (mC/total C) in each sequence context (CG, CHG, or CHH).

Figure 6

Figure 6. No significant decrease in nucleosome occupancy was observed in the *mop* mutants. ChIP experiments were performed with an antibody recognizing histone H3, using husk tissue from *B'* (light green), *B' Mop/mop2-1* (green), *B' mop2-1* (black), *B' mop1-1* (magenta), *B' mop3-1* (blue), and *B-I* (purple) plants. Data was normalized to *actin* values. The error bars indicate the standard error of the mean (SEM) of two (*B' mop1-1*), three (*B'*, *B' Mop/mop2-1*, *B' mop3-1*), or four (*B' mop2-1*, *B-I*) replicate experiments. For the no-antibody control signals see Supplemental Figure S10.

Figure 7

Figure 7. Transcriptional activation in *mop1-1* and *mop3-1* is associated with reduced levels of repressive histone marks. ChIP-qPCR experiments were performed on seedling (A, C) and husk tissue (B, D) from *B'*, *B' Mop/mop2-1*, *B' mop2-1*, *B' mop1-1*, *B' mop3-1* and *B-I* plants with antibodies recognizing H3K9me2 (A, B), and H3K27me2 (C, D). ChIP signals were normalized to *Copia* signals. Error bars indicate the SEM of multiple biological replicates. For seedlings, three (*B'*, *B' mop1-1*) or two (*B' mop3-1*, *B-I*) replicates were performed. For husks (B, D), two (*B' mop2-2*), three (*B'*, *B' mop3-1*, *B-I*), or four (*B' mop1-1*, *B' Mop/mop2-1*, *B' mop2-1*) replicates were performed. For the no-antibody control signals see Supplemental Figure S11.

Parsed Citations

Alleman, M., Sidorenko, L., McGinnis, K., Seshadri, V., Dorweiler, J.E., White, J., Sikkink, K., Chandler, V.L., 2006. An RNA-dependent RNA polymerase is required for paramutation in maize. Nature 442, 295–298. https://doi.org/10.1038/nature04884
  Google Scholar: Author Only Title Only Author and Title

Arteaga-Vazquez, M., Sidorenko, L., Rabanal, F.A, Shrivistava, R., Nobuta, K., Green, P.J., Meyers, B.C., Chandler, V.L., 2010. RNA-mediated trans -communication can establish paramutation at the b1 locus in maize. PNAS 107, 12986–12991. https://doi.org/10.1073/pnas.1007972107/-/DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1007972107
  Google Scholar: Author Only Title Only Author and Title

Bartee, L., Malagnac, F., Bender, J., 2001. Arabidopsis cmt3 chromomethylase mutations block non-CG methylation and silencing of an endogenous gene. Genes Dev 15, 1753–1758. https://doi.org/10.1101/gad.905701
  Google Scholar: Author Only Title Only Author and Title

Belele, C.L., Sidorenko, L., Stam, M., Bader, R., Arteaga-Vazquez, M.A, Chandler, V.L., 2013. Specific Tandem Repeats Are Sufficient for Paramutation-Induced Trans-Generational Silencing. PLoS Genetics 9. https://doi.org/10.1371/journal.pgen.1003773
  Google Scholar: Author Only Title Only Author and Title

Bernatavichute, Y.V., Zhang, X., Cokus, S., Pellegrini, M., Jacobsen, S.E., 2008. Genome-wide association of histone H3 lysine nine methylation with CHG DNA methylation in Arabidopsis thaliana. PLoS One 3, e3156. https://doi.org/10.1371/journal.pone.0003156
  Google Scholar: Author Only Title Only Author and Title

Calarco, J.P., Borges, F., Donoghue, M.T.A, Van Ex, F., Jullien, P.E., Lopes, T., Gardner, R., Berger, F., Feijó, J.A, Becker, J.D., Martienssen, R.A., 2012. Reprogramming of DNA Methylation in Pollen Guides Epigenetic Inheritance via Small RNA. Cell 151, 194–205. https://doi.org/10.1016/j.cell.2012.09.001
  Google Scholar: Author Only Title Only Author and Title

Castel, S.E., Martienssen, R.A., 2013. RNA interference in the nucleus: roles for small RNAs in transcription, epigenetics and beyond. Nat Rev Genet 14, 100–112. https://doi.org/10.1038/nrg3355
  Google Scholar: Author Only Title Only Author and Title

Conway, E., Healy, E., Bracken, A.P., 2015. PRC2 mediated H3K27 methylations in cellular identity and cancer. Curr Opin Cell Biol 37, 42–48. https://doi.org/10.1016/j.ceb.2015.10.003
  Google Scholar: Author Only Title Only Author and Title

Dellaporta, S.L., Wood, J., Hicks, J.B., 1983. A plant DNA minipreparation: Version II. Plant Mol Biol Rep 1, 19–21. https://doi.org/10.1007/BF02712670
  Google Scholar: Author Only Title Only Author and Title

Dorweiler, J.E., Carey, C.C., Kubo, K.M., Hollick, J.B., Kermicle, J.L., Chandler, V.L., 2000. mediator of paramutation1 is required for establishment and maintenance of paramutation at multiple maize loci. Plant Cell 12, 2101–2118. https://doi.org/10.1105/tpc.12.11.2101
  Google Scholar: Author Only Title Only Author and Title

Du, J., Johnson, L.M., Groth, M., Feng, S., Hale, C.J., Li, S., Vashisht, A.A, Wohlschlegel, J.A, Patel, D.J., Jacobsen, S.E., 2014. Mechanism of DNA methylation-directed histone methylation by KRYPTONITE. Mol Cell 55, 495–504. https://doi.org/10.1016/j.molcel.2014.06.009
  Google Scholar: Author Only Title Only Author and Title

Du, J., Zhong, X., Bernatavichute, Y.V., Stroud, H., Feng, S., Caro, E., Vashisht, A.A, Terragni, J., Chin, H.G., Tu, A, Hetzel, J., Wohlschlegel, J.A, Pradhan, S., Patel, D.J., Jacobsen, S.E., 2012. Dual binding of chromomethylase domains to H3K9me2-containing nucleosomes directs DNA methylation in plants. Cell 151, 167–180. https://doi.org/10.1016/j.cell.2012.07.034
  Google Scholar: Author Only Title Only Author and Title

Egelhofer, T.A, Minoda, A, Klugman, S., Lee, K., Kolasinska-Zwierz, P., Alekseyenko, A.A, Cheung, M.-S., Day, D.S., Gadel, S., Gorchakov, A.A, Gu, T., Kharchenko, P.V., Kuan, S., Latorre, I., Linder-Basso, D., Luu, Y., Ngo, Q., Perry, M., Rechtsteiner, A, Riddle, N.C., Schwartz, Y.B., Shanower, G.A, Vielle, A, Ahringer, J., Elgin, S.C.R., Kuroda, M.I., Pirrotta, V., Ren, B., Strome, S., Park, P.J., Karpen, G.H., Hawkins, R.D., Lieb, J.D., 2011. An assessment of histone-modification antibody quality. Nat Struct Mol Biol 18, 91–93. https://doi.org/10.1038/nsmb.1972
  Google Scholar: Author Only Title Only Author and Title

Erhard, K.F., Stonaker, J.L., Parkinson, S.E., Al., E., 2009. RNA Polymerase IV Functions in Paramutation in Zea mays. Science 323, 1201–1205.
  Google Scholar: Author Only Title Only Author and Title

Ferrari, K.J., Scelfo, A, Jammula, S., Cuomo, A, Barozzi, I., Stützer, A, Fischle, W., Bonaldi, T., Pasini, D., 2014. Polycomb-dependent H3K27me1 and H3K27me2 regulate active transcription and enhancer fidelity. Mol Cell 53, 49–62. https://doi.org/10.1016/j.molcel.2013.10.030

Google Scholar: Author Only Title Only Author and Title

Fitz-James, M.H., Cavalli, G., 2022. Molecular mechanisms of transgenerational epigenetic inheritance. Nat Rev Genet 23, 325–341. https://doi.org/10.1038/s41576-021-00438-5
Google Scholar: Author Only Title Only Author and Title

Foda, B.M., Singh, U., 2015. Dimethylated H3K27 Is a Repressive Epigenetic Histone Mark in the Protist Entamoeba histolytica and Is Significantly Enriched in Genes Silenced via the RNAi Pathway. Journal of Biological Chemistry 290, 21114–21130. https://doi.org/10.1074/jbc.M115.647263
Google Scholar: Author Only Title Only Author and Title

Fu, F.-F., Dawe, R.K., Gent, J.I., 2018. Loss of RNA-Directed DNA Methylation in Maize Chromomethylase and DDM1-Type Nucleosome Remodeler Mutants. The Plant Cell 30, 1617–1627. https://doi.org/10.1105/tpc.18.00053
Google Scholar: Author Only Title Only Author and Title

Gent, Jonathan I, Ellis, N.A, Guo, L., Harkess, A.E., Yao, Y., Zhang, X., Dawe, R.K., 2013. CHH islands : de novo DNA methylation in near-gene chromatin regulation in maize. Genome research 23, 628–637. https://doi.org/10.1101/gr.146985.112.as
Google Scholar: Author Only Title Only Author and Title

Gent, J I, Ellis, N.A., Guo, L., Harkess, A.E., Yao, Y., Zhang, X., Dawe, R.K., 2013. CHH islands: de novo DNA methylation in near-gene chromatin regulation in maize. Genome Research 23, 628–637. https://doi.org/10.1101/gr.146985.112
Google Scholar: Author Only Title Only Author and Title

Gent, J.I., Higgins, K.M., Swentowsky, K.W., Fu, F.-F., Zeng, Y., Kim, D.W., Dawe, R.K., Springer, N.M., Anderson, S.N., 2022. The maize gene maternal derepression of r1 encodes a DNA glycosylase that demethylates DNA and reduces siRNA expression in the endosperm. Plant Cell 34, 3685–3701. https://doi.org/10.1093/plcell/koac199
Google Scholar: Author Only Title Only Author and Title

Gent, Jonathan I, Madzima, T.F., Bader, R., Kent, M.R., Zhang, X., Stam, M., McGinnis, K.M., Dawe, R.K., 2014b. Accessible DNA and Relative Depletion of H3K9me2 at Maize Loci Undergoing RNA-Directed DNA Methylation. The Plant Cell Online 26, 4903–4917. https://doi.org/10.1105/tpc.114.130427
Google Scholar: Author Only Title Only Author and Title

Grewal, S.I.S., 2023. The molecular basis of heterochromatin assembly and epigenetic inheritance. Mol Cell 83, 1767–1785. https://doi.org/10.1016/j.molcel.2023.04.020
Google Scholar: Author Only Title Only Author and Title

Gruntman, E., Qi, Y., Slotkin, R.K., Roeder, T., Martienssen, R.A., Sachidanandam, R., 2008. Kismeth: Analyzer of plant methylation states through bisulfite sequencing. BMC Bioinformatics 9, 371. https://doi.org/10.1186/1471-2105-9-371
Google Scholar: Author Only Title Only Author and Title

Gu, S.G., Pak, J., Guang, S., Maniar, J.M., Kennedy, S., Fire, A, 2012. Amplification of siRNA in Caenorhabditis elegans generates a transgenerational sequence-targeted histone H3 lysine 9 methylation footprint. Nat Genet 44, 157–164. https://doi.org/10.1038/ng.1039
Google Scholar: Author Only Title Only Author and Title

Guo, W., Fiziev, P., Yan, W., Cokus, S., Sun, X., Zhang, M.Q., Chen, P.Y., Pellegrini, M., 2013. BS-Seeker2: A versatile aligning pipeline for bisulfite sequencing data. BMC Genomics 14, 1. https://doi.org/10.1186/1471-2164-14-774
Google Scholar: Author Only Title Only Author and Title

Guo, W., Zhu, P., Pellegrini, M., Zhang, M.Q., Wang, X., Ni, Z, 2018. CGmapTools improves the precision of heterozygous SNV calls and supports allele-specific methylation detection and visualization in bisulfite-sequencing data. Bioinformatics 34, 381–387. https://doi.org/10.1093/bioinformatics/btx595
Google Scholar: Author Only Title Only Author and Title

Gutiérrez-Marcos, J.F., Costa, L.M., Dal Prà, M., Scholten, S., Kranz, E., Perez, P., Dickinson, H.G., 2006. Epigenetic asymmetry of imprinted genes in plant gametes. Nat Genet 38, 876–878. https://doi.org/10.1038/ng1828
Google Scholar: Author Only Title Only Author and Title

Haag, J.R., Brower-Toland, B., Krieger, E.K., Sidorenko, L., Nicora, C.D., Norbeck, A.D., Irsigler, A, LaRue, H., Brzeski, J., McGinnis, K., Ivashuta, S., Pasa-Tolic, L., Chandler, V.L., Pikaard, C.S., 2014. Functional diversification of maize RNA polymerase IV and V subtypes via alternative catalytic subunits. Cell reports 9, 378–90. https://doi.org/10.1016/j.celrep.2014.08.067
Google Scholar: Author Only Title Only Author and Title

Haring, M., Bader, R., Louwers, M., Schwabe, A, van Driel, R., Stam, M., 2010a. The role of DNA methylation, nucleosome occupancy and histone modifications in paramutation. The Plant journal : for cell and molecular biology 63, 366–78. https://doi.org/10.1111/j.1365-313X.2010.04245.x
Google Scholar: Author Only Title Only Author and Title

Haring, M., Offermann, S., Danker, T., Horst, I., Peterhansel, C., Stam, M., 2007. Chromatin immunoprecipitation: optimization,

quantitative analysis and data normalization. Plant Methods 3, 11. https://doi.org/10.1186/1746-4811-3-11
Google Scholar: Author Only Title Only Author and Title

Herr, A.J., Jensen, M.B., Dalmay, T., Baulcombe, D.C., 2005. RNA polymerase IV directs silencing of endogenous DNA. Science 308, 118–120. https://doi.org/10.1126/science.1106910
Google Scholar: Author Only Title Only Author and Title

Hollick, J.B., 2017. Paramutation and related phenomena in diverse species. Nat Rev Genet 18, 5–23. https://doi.org/10.1038/nrg.2016.115
Google Scholar: Author Only Title Only Author and Title

Hollick, Jay B, Kermicle, J.L., Parkinson, S.E., 2005. Rmr6 maintains meiotic inheritance of paramutant states in Zea mays. Genetics 171, 725–40. https://doi.org/10.1534/genetics.105.045260
Google Scholar: Author Only Title Only Author and Title

Hollick, J. B., Kermicle, J.L., Parkinson, S.E., 2005. Rmr6 maintains meiotic inheritance of paramutant states in Zea mays. Genetics 171, 725–740. https://doi.org/10.1534/genetics.105.045260
Google Scholar: Author Only Title Only Author and Title

Hollick, J.B., Patterson, G.I., Coe, E.H., Cone, J.K.C., Chandler, V.L., 1995. Allelic Interactions Heritably Alter the Activity of a Metastable Maize pl Allele. Genetics 141, 709–719.
Google Scholar: Author Only Title Only Author and Title

Hövel, I., Pearson, N.A., Stam, M., 2015. Cis-acting determinants of paramutation. Semin Cell Dev Biol 44, 22–32. https://doi.org/10.1016/j.semcdb.2015.08.012
Google Scholar: Author Only Title Only Author and Title

Hufford, M.B., Seetharam, A.S., Woodhouse, M.R., Chougule, K.M., Ou, S., Liu, J., Ricci, W.A., Guo, T., Olson, A, Qiu, Y., Della Coletta, R., Tittes, S., Hudson, A.I., Marand, A.P., Wei, S., Lu, Z, Wang, B., Tello-Ruiz, M.K., Piri, R.D., Wang, N., Kim, D.W., Zeng, Y., O'Connor, C.H., Li, X., Gilbert, A.M., Baggs, E., Krasileva, K.V., Portwood, J.L., Cannon, E.K.S., Andorf, C.M., Manchanda, N., Snodgrass, S.J., Hufnagel, D.E., Jiang, Q., Pedersen, S., Syring, M.L., Kudrna, D.A, Llaca, V., Fengler, K., Schmitz, R.J., Ross-Ibarra, J., Yu, J., Gent, J.I., Hirsch, C.N., Ware, D., Dawe, R.K., 2021. De novo assembly, annotation, and comparative analysis of 26 diverse maize genomes. Science 373, 655–662. https://doi.org/10.1126/science.abg5289
Google Scholar: Author Only Title Only Author and Title

Jackel, J.N., Storer, J.M., Coursey, T., Bisaro, D.M., 2016. Arabidopsis RNA Polymerases IV and V Are Required To Establish H3K9 Methylation, but Not Cytosine Methylation, on Geminivirus Chromatin. Journal of Virology 90, 7529–7540. https://doi.org/10.1128/jvi.00656-16
Google Scholar: Author Only Title Only Author and Title

Jia, Y., Lisch, D.R., Ohtsu, K., Scanlon, M.J., Nettleton, D., Schnable, P.S., 2009. Loss of RNA-dependent RNA polymerase 2 (RDR2) function causes widespread and unexpected changes in the expression of transposons, genes, and 24-nt small RNAs. PLoS Genet 5, e1000737. https://doi.org/10.1371/journal.pgen.1000737
Google Scholar: Author Only Title Only Author and Title

Jiang, C., Pugh, B.F., 2009. Nucleosome positioning and gene regulation: advances through genomics. Nat Rev Genet 10, 161–172. https://doi.org/10.1038/nrg2522
Google Scholar: Author Only Title Only Author and Title

Johnson, L.M., Bostick, M., Zhang, X., Kraft, E., Henderson, I., Callis, J., Jacobsen, S.E., 2007. The SRA methyl-cytosine-binding domain links DNA and histone methylation. Curr Biol 17, 379–384. https://doi.org/10.1016/j.cub.2007.01.009
Google Scholar: Author Only Title Only Author and Title

Kanno, T., Huettel, B., Mette, M.F., Aufsatz, W., Jaligot, E., Daxinger, L., Kreil, D.P., Matzke, M., Matzke, A.J.M., 2005. Atypical RNA polymerase subunits required for RNA-directed DNA methylation. Nat Genet 37, 761–765. https://doi.org/10.1038/ng1580
Google Scholar: Author Only Title Only Author and Title

Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9, 357–359. https://doi.org/10.1038/nmeth.1923
Google Scholar: Author Only Title Only Author and Title

Law, J.A, Vashisht, A.A, Wohlschlegel, J.A, Jacobsen, S.E., 2011. SHH1, a homeodomain protein required for DNA methylation, as well as RDR2, RDM4, and chromatin remodeling factors, associate with RNA polymerase IV. PLoS Genet 7, e1002195. https://doi.org/10.1371/journal.pgen.1002195
Google Scholar: Author Only Title Only Author and Title

Lee, H.-G., Kahn, T.G., Simcox, A, Schwartz, Y.B., Pirrotta, V., 2015. Genome-wide activities of Polycomb complexes control pervasive transcription. Genome Res. 25, 1170–1181. https://doi.org/10.1101/gr.188920.114
Google Scholar: Author Only Title Only Author and Title

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., 1000 Genome Project Data Processing Subgroup, 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25, 2078–2079. https://doi.org/10.1093/bioinformatics/btp352
  Google Scholar: Author Only Title Only Author and Title

Li, Q., Eichten, S.R., Hermanson, P.J., Zaunbrecher, V.M., Song, J., Wendt, J., Rosenbaum, H., Madzima, T.F., Sloan, A.E., Huang, J., Burgess, D.L., Richmond, T.A., McGinnis, K.M., Meeley, R.B., Danilevskaya, O.N., Vaughn, M.W., Kaeppler, S.M., Jeddeloh, J.A., Springer, N.M., 2014. Genetic Perturbation of the Maize Methylome. The Plant Cell Online 26, 4602–4616. https://doi.org/10.1105/tpc.114.133140
  Google Scholar: Author Only Title Only Author and Title

Li, Q., Gent, J.I., Zynda, G., Song, J., Makarevitch, I., Hirsch, C.D., Hirsch, C.N., Dawe, R.K., Madzima, T.F., McGinnis, K.M., Lisch, D., Schmitz, R.J., Vaughn, M.W., Springer, N.M., 2015. RNA-directed DNA methylation enforces boundaries between heterochromatin and euchromatin in the maize genome. Proceedings of the National Academy of Sciences 112, 14728–14733. https://doi.org/10.1073/pnas.1514680112
  Google Scholar: Author Only Title Only Author and Title

Lindroth, A.M., Cao, X., Jackson, J.P., Zilberman, D., McCallum, C.M., Henikoff, S., Jacobsen, S.E., 2001. Requirement of CHROMOMETHYLASE3 for maintenance of CpXpG methylation. Science 292, 2077–2080. https://doi.org/10.1126/science.1059745
  Google Scholar: Author Only Title Only Author and Title

Liu, Y., Taverna, S.D., Muratore, T.L., Shabanowitz, J., Hunt, D.F., Allis, C.D., 2007. RNAi-dependent H3K27 methylation is required for heterochromatin formation and DNA elimination in Tetrahymena. Genes Dev 21, 1530–1545. https://doi.org/10.1101/gad.1544207
  Google Scholar: Author Only Title Only Author and Title

Liu, Y., Yuan, J., Jia, G., Ye, W., Jeffrey Chen, Z., Song, Q., 2021. Histone H3K27 dimethylation landscapes contribute to genome stability and genetic recombination during wheat polyploidization. The Plant Journal 105, 678–690. https://doi.org/10.1111/tpj.15063
  Google Scholar: Author Only Title Only Author and Title

Locatelli, S., Piatti, P., Motto, M., Rossi, V., 2009. Chromatin and DNA modifications in the Opaque2-mediated regulation of gene transcription during maize endosperm development. Plant Cell 21, 1410–1427. https://doi.org/10.1105/tpc.109.067256
  Google Scholar: Author Only Title Only Author and Title

Louwers, M., Bader, R., Haring, M., van Driel, R., de Laat, W., Stam, M., 2009a. Tissue- and expression level-specific chromatin looping at maize b1 epialleles. The Plant Cell 21, 832–42. https://doi.org/10.1105/tpc.108.064329
  Google Scholar: Author Only Title Only Author and Title

Louwers, M., Splinter, E., van Driel, R., de Laat, W., Stam, M., 2009b. Studying physical chromatin interactions in plants using Chromosome Conformation Capture (3C). Nat Protoc 4, 1216–1229. https://doi.org/10.1038/nprot.2009.113
  Google Scholar: Author Only Title Only Author and Title

Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal 17, 10–12. https://doi.org/10.14806/ej.17.1.200
  Google Scholar: Author Only Title Only Author and Title

Mathieu, O., Probst, A.V., Paszkowski, J., 2005. Distinct regulation of histone H3 methylation at lysines 27 and 9 by CpG methylation in Arabidopsis. The EMBO Journal 24, 2783–2791. https://doi.org/10.1038/sj.emboj.7600743
  Google Scholar: Author Only Title Only Author and Title

Mathieu, O., Reinders, J., Caikovski, M., Smathajitt, C., Paszkowski, J., 2007. Transgenerational stability of the Arabidopsis epigenome is coordinated by CG methylation. Cell 130, 851–862. https://doi.org/10.1016/j.cell.2007.07.007
  Google Scholar: Author Only Title Only Author and Title

Matzke, M.A, Mosher, R.A, 2014. RNA-directed DNA methylation: an epigenetic pathway of increasing complexity. Nat Rev Genet 15, 394–408. https://doi.org/10.1038/nrg3683
  Google Scholar: Author Only Title Only Author and Title

Mishra, V., Singh, J., Wang, F., Zhang, Y., Fukudome, A, Trinidad, J.C., Takagi, Y., Pikaard, C.S., 2021. Assembly of a dsRNA synthesizing complex: RNA-DEPENDENT RNA POLYMERASE 2 contacts the largest subunit of NUCLEAR RNA POLYMERASE IV. Proc Natl Acad Sci U S A 118, e2019276118. https://doi.org/10.1073/pnas.2019276118
  Google Scholar: Author Only Title Only Author and Title

Nobuta, K., Lu, C., Shrivastava, R., Pillay, M., De Paoli, E., Accerbi, M., Arteaga-Vazquez, M., Sidorenko, L., Jeong, D.-H., Yen, Y., Green, P.J., Chandler, V.L., Meyers, B.C., 2008a. Distinct size distribution of endogeneous siRNAs in maize: Evidence from deep sequencing in the mop1-1 mutant. Proc Natl Acad Sci U S A 105, 14958–14963. https://doi.org/10.1073/pnas.0808066105
  Google Scholar: Author Only Title Only Author and Title

Nobuta, K., Lu, C., Shrivastava, R., Pillay, M., Paoli, E. De, Accerbi, M., Arteaga-vazquez, M., Sidorenko, L., Jeong, D., Yen, Y.,

Green, P.J., Chandler, V.L., Meyers, B.C., 2008b. Distinct size distribution of endogenous siRNAs in maize : Evidence from deep sequencing in the mop1-1 mutant. PNAS 105, 14958–14963.
Google Scholar: Author Only Title Only Author and Title

Ohtsu, K., Smith, M.B., Emrich, S.J., Borsuk, L.A, Zhou, R., Chen, T., Zhang, X., Timmermans, M.C.P., Beck, J., Buckner, B., Janick-Buckner, D., Nettleton, D., Scanlon, M.J., Schnable, P.S., 2007. Global gene expression analysis of the shoot apical meristem of maize (Zea mays L.). Plant J 52, 391–404. https://doi.org/10.1111/j.1365-313X.2007.03244.x
Google Scholar: Author Only Title Only Author and Title

Onodera, Y., Haag, J.R., Ream, T., Costa Nunes, P., Pontes, O., Pikaard, C.S., 2005. Plant nuclear RNA polymerase IV mediates siRNA and DNA methylation-dependent heterochromatin formation. Cell 120, 613–622. https://doi.org/10.1016/j.cell.2005.02.007
Google Scholar: Author Only Title Only Author and Title

Pal-Bhadra, M., Leibovitch, B.A, Gandhi, S.G., Rao, M., Bhadra, U., Birchler, J.A, Elgin, S.C.R., 2004. Heterochromatic Silencing and HP1 Localization in Drosophila Are Dependent on the RNAi Machinery. Science 303, 669–672. https://doi.org/10.1126/science.1092653
Google Scholar: Author Only Title Only Author and Title

Papa, C.M., Springer, N.M., Muszynski, M.G., Meeley, R., Kaeppler, S.M., 2001. Maize chromomethylase Zea methyltransferase2 is required for CpNpG methylation. Plant Cell 13, 1919–1928. https://doi.org/10.1105/tpc.010064
Google Scholar: Author Only Title Only Author and Title

Parent, J.-S., Cahn, J., Herridge, R.P., Grimanelli, D., Martienssen, R.A, 2021. Small RNAs guide histone methylation in Arabidopsis embryos. Genes Dev. 35, 841–846. https://doi.org/10.1101/gad.343871.120
Google Scholar: Author Only Title Only Author and Title

Park, S., Oh, S., Nocker, S. van, 2012. Genomic and Gene-Level Distribution of Histone H3 Dimethyl Lysine-27 (H3K27me2) in Arabidopsis. PLOS ONE 7, e52855. https://doi.org/10.1371/journal.pone.0052855
Google Scholar: Author Only Title Only Author and Title

Patterson, G.I., Thorpe, C.J., Chandler, V.L., 1993. Paramutation, an allelic interaction, is associated with a stable and heritable reduction of transcription of the maize b regulatory gene. Genetics 135, 881–894. https://doi.org/10.1093/genetics/135.3.881
Google Scholar: Author Only Title Only Author and Title

Plank, J.L., Dean, A, 2014. Enhancer function: mechanistic and genome-wide insights come together. Mol Cell 55, 5–14. https://doi.org/10.1016/j.molcel.2014.06.015
Google Scholar: Author Only Title Only Author and Title

Pontier, D., Yahubyan, G., Vega, D., Bulski, A, Saez-Vasquez, J., Hakimi, M.-A, Lerbs-Mache, S., Colot, V., Lagrange, T., 2005. Reinforcement of silencing at transposons and highly repeated sequences requires the concerted action of two distinct RNA polymerases IV in Arabidopsis. Genes Dev 19, 2030–2040. https://doi.org/10.1101/gad.348405
Google Scholar: Author Only Title Only Author and Title

Quinlan, A.R., Hall, I.M., 2010. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics 26, 841–842. https://doi.org/10.1093/bioinformatics/btq033
Google Scholar: Author Only Title Only Author and Title

Ritchie, S.W., Hanway, J.J., Benson, G.O., 1986. How a corn plant develops, 1986 (Departmental Report).

Ronsseray, S., 2015. Paramutation phenomena in non-vertebrate animals. Semin Cell Dev Biol 44, 39–46. https://doi.org/10.1016/j.semcdb.2015.08.009
Google Scholar: Author Only Title Only Author and Title

Roudier, F., Ahmed, I., Bérard, C., Sarazin, A, Mary-Huard, T., Cortijo, S., Bouyer, D., Caillieux, E., Duvernois-Berthet, E., Al-Shikhley, L., Giraut, L., Després, B., Drevensek, S., Barneche, F., Dèrozier, S., Brunaud, V., Aubourg, S., Schnittger, A, Bowler, C., Martin-Magniette, M.-L., Robin, S., Caboche, M., Colot, V., 2011. Integrative epigenomic mapping defines four main chromatin states in Arabidopsis: Organization of the Arabidopsis epigenome. The EMBO Journal 30, 1928–1938. https://doi.org/10.1038/emboj.2011.103
Google Scholar: Author Only Title Only Author and Title

Rymen, B., Ferrafiat, L., Blevins, T., 2020. Non-coding RNA polymerases that silence transposable elements and reprogram gene expression in plants. Transcription 11, 172–191. https://doi.org/10.1080/21541264.2020.1825906
Google Scholar: Author Only Title Only Author and Title

Schmitz, R.J., Grotewold, E., Stam, M., 2022. Cis-regulatory sequences in plants: Their importance, discovery, and future challenges. Plant Cell 34, 718–741. https://doi.org/10.1093/plcell/koab281
Google Scholar: Author Only Title Only Author and Title

Sekhon, R.S., Lin, H., Childs, K.L., Hansey, C.N., Buell, C.R., de Leon, N., Kaeppler, S.M., 2011. Genome-wide atlas of transcription during maize development. Plant J 66, 553–563. https://doi.org/10.1111/j.1365-313X.2011.04527.x

Google Scholar: Author Only Title Only Author and Title

Sekhon, R.S., Wang, P.-H., Sidorenko, L., Chandler, V.L., Chopra, S., 2012. Maize Unstable factor for orange1 is required for maintaining silencing associated with paramutation at the pericarp color1 and booster1 loci. PLoS Genet 8, e1002980. https://doi.org/10.1371/journal.pgen.1002980
Google Scholar: Author Only Title Only Author and Title

Sidorenko, L., Chandler, V., 2008. RNA-dependent RNA polymerase is required for enhancer-mediated transcriptional silencing associated with paramutation at the maize p1 gene. Genetics 180, 1983–1993. https://doi.org/10.1534/genetics.108.095281
Google Scholar: Author Only Title Only Author and Title

Sidorenko, L., Dorweiler, J.E., Cigan, a M., Arteaga-Vazquez, M., Vyas, M., Kermicle, J., Jurcin, D., Brzeski, J., Cai, Y., Chandler, V.L., 2009. A dominant mutation in mediator of paramutation2, one of three second-largest subunits of a plant-specific RNA polymerase, disrupts multiple siRNA silencing processes. PLoS genetics 5, e1000725. https://doi.org/10.1371/journal.pgen.1000725
Google Scholar: Author Only Title Only Author and Title

Sloan, Amy E., Sidorenko, L., McGinnis, K.M., 2014. Diverse gene-silencing mechanisms with distinct requirements for RNA polymerase subunits in Zea mays. Genetics 198, 1031–1042. https://doi.org/10.1534/genetics.114.168518
Google Scholar: Author Only Title Only Author and Title

Springer, N.M., McGinnis, K.M., 2015. Paramutation in evolution, population genetics and breeding. Semin Cell Dev Biol 44, 33–38. https://doi.org/10.1016/j.semcdb.2015.08.010
Google Scholar: Author Only Title Only Author and Title

Stam, M., Belele, C., Dorweiler, J.E., Chandler, V.L., 2002a. Differential chromatin structure within a tandem array 100 kb upstream of the maize b1 locus is associated with paramutation. Genes and Development 16, 1906–1918. https://doi.org/10.1101/gad.1006702
Google Scholar: Author Only Title Only Author and Title

Stam, M., Belele, C., Ramakrishna, W., Dorweiler, J.E., Bennetzen, J.L., Chandler, V.L., 2002b. The regulatory regions required for B' paramutation and expression are located far upstream of the maize b1 transcribed sequences. Genetics 162, 917–930. https://doi.org/10.1093/genetics/162.2.917
Google Scholar: Author Only Title Only Author and Title

Stonaker, Jennifer L., Lim, J.P., Erhard, K.F., Hollick, J.B., 2009. Diversity of Pol IV function is defined by mutations at the maize rmr7 locus. PLoS Genet 5, e1000706. https://doi.org/10.1371/journal.pgen.1000706
Google Scholar: Author Only Title Only Author and Title

Stroud, H., Do, T., Du, J., Zhong, X., Feng, S., Johnson, L., Patel, D.J., Jacobsen, S.E., 2014. Non-CG methylation patterns shape the epigenetic landscape in Arabidopsis. Nat Struct Mol Biol 21, 64–72. https://doi.org/10.1038/nsmb.2735
Google Scholar: Author Only Title Only Author and Title

Stroud, H., Greenberg, M.V.C., Feng, S., Bernatavichute, Y.V., Jacobsen, S.E., 2013. Comprehensive analysis of silencing mutants reveals complex regulation of the Arabidopsis methylome. Cell 152, 352–364. https://doi.org/10.1016/j.cell.2012.10.054
Google Scholar: Author Only Title Only Author and Title

Struhl, K., Segal, E., 2013. Determinants of nucleosome positioning. Nat Struct Mol Biol 20, 267–273. https://doi.org/10.1038/nsmb.2506
Google Scholar: Author Only Title Only Author and Title

Tang, K., Lang, Z., Zhang, H., Zhu, J.-K., 2016. The DNA demethylase ROS1 targets genomic regions with distinct chromatin modifications. Nat Plants 2, 16169. https://doi.org/10.1038/nplants.2016.169
Google Scholar: Author Only Title Only Author and Title

Underwood, C.J., Choi, K., Lambing, C., Zhao, X., Serra, H., Borges, F., Simorowski, J., Ernst, E., Jacob, Y., Henderson, I.R., Martienssen, R.A., 2018. Epigenetic activation of meiotic recombination near Arabidopsis thaliana centromeres via loss of H3K9me2 and non-CG DNA methylation. Genome Res 28, 519–531. https://doi.org/10.1101/gr.227116.117
Google Scholar: Author Only Title Only Author and Title

Verdel, A., Jia, S., Gerber, S., Sugiyama, T., Gygi, S., Grewal, S.I.S., Moazed, D., 2004. RNAi-mediated targeting of heterochromatin by the RITS complex. Science 303, 672–676. https://doi.org/10.1126/science.1093686
Google Scholar: Author Only Title Only Author and Title

West, P.T., Li, Q., Ji, L., Eichten, S.R., Song, J., Vaughn, M.W., Schmitz, R.J., Springer, N.M., 2014. Genomic distribution of H3K9me2 and DNA methylation in a maize genome. PLoS One 9, e105267. https://doi.org/10.1371/journal.pone.0105267
Google Scholar: Author Only Title Only Author and Title

Wierzbicki, A.T., Haag, J.R., Pikaard, C.S., 2008. Noncoding transcription by RNA polymerase Pol IVb/Pol V mediates transcriptional silencing of overlapping and adjacent genes. Cell 135, 635–648. https://doi.org/10.1016/j.cell.2008.09.035

Google Scholar: Author Only Title Only Author and Title

Woodhouse, M.R., Freeling, M., Lisch, D., 2006a. Initiation, establishment, and maintenance of heritable MuDR transposon silencing in maize are mediated by distinct factors. PLoS Biol 4, e339. https://doi.org/10.1371/journal.pbio.0040339
Google Scholar: Author Only Title Only Author and Title

Woodhouse, M.R., Freeling, M., Lisch, D., 2006b. The mop1 (mediator of paramutation1) mutant progressively reactivates one of the two genes encoded by the MuDR transposon in maize. Genetics 172, 579–592. https://doi.org/10.1534/genetics.105.051383
Google Scholar: Author Only Title Only Author and Title

Xie, W., Schultz, M.D., Lister, R., Hou, Z, Rajagopal, N., Ray, P., Whitaker, J.W., Tian, S., Hawkins, R.D., Leung, D., Yang, H., Wang, T., Lee, A.Y., Swanson, S.A, Zhang, J., Zhu, Y., Kim, A, Nery, J.R., Urich, M.A, Kuan, S., Yen, C., Klugman, S., Yu, P., Suknuntha, K., Propson, N.E., Chen, H., Edsall, L.E., Wagner, U., Li, Y., Ye, Z, Kulkarni, A, Xuan, Z, Chung, W.-Y., Chi, N.C., Antosiewicz-Bourget, J.E., Slukvin, I., Stewart, R., Zhang, M.Q., Wang, W., Thomson, J.A, Ecker, J.R., Ren, B., 2013. Epigenomic analysis of multilineage differentiation of human embryonic stem cells. Cell 153, 1134–1148. https://doi.org/10.1016/j.cell.2013.04.022
Google Scholar: Author Only Title Only Author and Title

Zemach, A, Kim, M.Y., Hsieh, P.-H., Coleman-Derr, D., Eshed-Williams, L., Thao, K., Harmer, S.L., Zilberman, D., 2013. The Arabidopsis Nucleosome Remodeler DDM1 Allows DNA Methyltransferases to Access H1-Containing Heterochromatin. Cell 153, 193–205. https://doi.org/10.1016/j.cell.2013.02.033
Google Scholar: Author Only Title Only Author and Title

Zofall, M., Sandhu, R., Holla, S., Wheeler, D., Grewal, S.I.S., 2022. Histone deacetylation primes self-propagation of heterochromatin domains to promote epigenetic inheritance. Nat Struct Mol Biol 29, 898–909. https://doi.org/10.1038/s41594-022-00830-7
Google Scholar: Author Only Title Only Author and Title