Rab7a activation promotes degradation of select tight junction proteins at the blood-brain barrier after ischemic stroke

Azzurra Cottarelli[1,2,*], Sanjid Shahriar[2,3,**], Ahmet Arac[4,**], Michael Glendinning[1], Mary Claire Tuohy[1], Grace Prochilo[1], Jason B. Neal[1,5], Aimee L. Edinger[6], and Dritan Agalliu[1,2,*]

Departments of [1]Neurology and [2]Pathology and Cell Biology, Columbia University Irving Medical Center, New York, NY, 10032, USA

[3]Wyss Institute for Biologically Inspired Engineering, Boston, MA 02115, USA

[4]Department of Neurology, David Geffen School of Medicine, University of California in Los Angeles, Los Angeles, CA, 90095, USA

[5]Baylor Scott and White Health, Dallas, TX, 75226, USA

[6]Departments of Developmental and Cell Biology and Pharmaceutical Sciences, University of California, Irvine, CA 92697, USA

[**] equal contribution

*Corresponding authors:

Azzurra Cottarelli, Ph.D.   Email: ac4023@cumc.columbia.edu

Dritan Agalliu, Ph.D.   Email: da191@cumc.columbia.edu

Lead contact:

Dritan Agalliu, Ph.D.   Email: da191@cumc.columbia.edu

SUMMARY (150 words)

The stability of tight junctions (TJs) between endothelial cells (ECs) is essential to maintain blood-brain barrier (BBB) function in the healthy brain. Following ischemic stroke, TJ strand dismantlement due to protein degradation leads to BBB dysfunction, yet the mechanisms driving this process are poorly understood. Here, we show that endothelial-specific ablation of Rab7a, a small GTPase that regulates endolysosomal protein degradation, reduces stroke-induced TJ strand disassembly resulting in decreased paracellular BBB permeability and improved neuronal outcomes. Two pro-inflammatory cytokines, TNFα and IL1β, but not glucose and oxygen deprivation, induce Rab7a activation via Ccz1 in brain ECs *in vitro*, leading to increased TJ protein degradation and impaired paracellular barrier function. Silencing Rab7a in brain ECs *in vitro* reduces cytokine-driven endothelial barrier dysfunction by suppressing degradation of a key BBB TJ protein, Claudin-5. Thus, Rab7a activation by inflammatory cytokines promotes degradation of select TJ proteins leading to BBB dysfunction after ischemic stroke.

KEYWORDS

Rab7a, tight junctions, blood-brain barrier, ischemic stroke, Claudin-5, inflammation, cytokines

INTRODUCTION

The stability of tight junction (TJs) strands formed between endothelial cells (ECs) lining the blood vessels is a critical structural component of the blood-brain barrier (BBB) that, together with low rates of transcellular transport through sparse transcytosis vesicles, restricts the permeability of blood-derived proteins and immune cells into the healthy central nervous system (CNS) to preserve neuronal function (reviewed in [1]). Disintegration of TJ strands occurs in several neurological disorders associated with BBB damage including ischemic stroke, a complex and devastating neurological condition that is the second leading cause of death and the third leading cause of disability worldwide (reviewed in [2-5]). ECs that survive in the peri-infarct region of the brain, but do not maintain BBB properties, facilitate damage to the adjacent parenchyma and exacerbate the clinical prognosis of stroke (reviewed in [1]). Therefore, targeting the molecular pathways that promote stabilization of BBB TJ strands is a highly attractive therapeutic strategy to improve stroke outcomes. Yet, the molecular mechanisms that drive persistent degradation of TJ proteins and loss of BBB integrity *in vivo* at the acute phase of ischemic stroke remain unclear.

During the reperfusion phase of acute ischemic stroke, there is a biphasic increase in BBB permeability that contributes to vasogenic edema, hemorrhage and increased mortality [6-9]. Dynamic imaging studies using green fluorescent protein (GFP)-tagged TJs have found that junctional strands are continuously remodeled at a slow rate in the healthy CNS ECs[10, 11]. However CNS EC junctional strands become highly dynamic and show an increased frequency and number of structural abnormalities (protrusions or gaps) between 48-58 hours after the transient middle cerebral artery occlusion (t-MCAO), a rodent model for ischemic stroke [10]. This phase coincides with increased expression in several pro-inflammatory cytokines, as well as recruitment of immune

cells into the CNS (reviewed in [12, 13]), that together contribute to stroke pathophysiology and

clinical outcomes [14-16]. These structural BBB TJ abnormalities are not unique to ischemic stroke.

They are also found in primary neuroinflammatory CNS diseases to allow immune cell infiltration

into the CNS (reviewed in [1]) and have been observed by *in vivo* two photon imaging in

experimental autoimmune encephalomyelitis (EAE), a rodent model for human multiple sclerosis

prior and throughout the course of the disease [11]. The temporal coincidence of TJ abnormalities

and inflammation across multiple CNS diseases emphasizes a critical role for pro-inflammatory

mechanisms in driving TJ strand disassembly.

What mechanisms drive BBB TJ strand disassembly after ischemic stroke? Caveolae have

been proposed to promote internalization and degradation of TJ transmembrane proteins such as

Claudin-5 and Occludin in CNS and peripheral ECs [17] as well as epithelial cells [18]. However,

Caveolin-1 (Cav-1) [-/-] mice lacking caveolae [19, 20] show increased paracellular BBB permeability

and infarct size after t-MCAO compared to wild-type mice [10, 21], suggesting that TJ strand

disassembly following brain ischemia is not mediated by a caveolar-dependent mechanism.

Enhanced endocytotic degradation of distinct BBB transmembrane and intracellular TJ proteins in

response to inflammatory cytokines is as a potential mechanism for persistent TJ strand

disassembly after ischemic stroke. EC junctional strand protrusions are inhibited by treatment with

inhibitors of endocytosis [18], and these structures express the early endosomal marker EEA-1 [11],

indicative of a transition into early endosomes. During endocytosis, early endosomes and their

cargoes can either be sorted back to the plasma membrane as recycling endosomes, or directed to

degradation by maturing into late endosomes/multivesicular bodies that ultimately fuse with

lysosomes [22]. Cell membrane trafficking of various cargoes, including transmembrane proteins, is

regulated by small GTPases of the Rab, Arf and Arl families, which cycle between inactive (GDP-bound) and active (GTP-bound) conformational states [23]. The small GTPase Rab7a is critical for cargo sorting into the late endosome, biogenesis of lysosomes and phagocytosis, as well as regulation of substrate degradation, antigen presentation, cell signaling/survival and microbial infection [24, 25].

Although the endolysosomal functions of Rab7 have been extensively studied in mammalian cells, how its activity is regulated is not well understood. In cancer cells, growth factor deprivation increases the fraction of Rab7a associated with membranes and the Rab7a-GTP (active) form, and triggers cell death due to degradation of nutrient transporters essential for survival [26-28]. Conversely, Rab7a inhibition in the setting of growth factor withdrawal promotes cancer cell survival due to recycling of transporters to the cell surface [29, 30]. Regulatory proteins that control Rab7a GTPase activity have been difficult to identify in cells. TBC1D15, has been identified as a putative Rab7a inactivator, since it accelerates GTP hydrolysis by Rab7a *in vitro* and its overexpression disrupts lysosomal morphology and blocks growth factor withdrawal-induced cell death in cancer cells [29]. Ccz1, a guanidine exchange factor (GEF) that acts together with vacuolar fusion protein Mon1, has been shown to promote both Rab7a activation and Rab7a-mediated membrane fusion in yeast [31] and eukaryotic cells [32-35]. Nevertheless, it is unclear whether Mon1/Ccz1 complex activates Rab7a in ECs and the role of Rab7a in EC function has not been fully elucidated. Rab5a and Rab7a regulate VEGFR2 receptor trafficking and the response of ECs to VEGF-A [36] via the Rab effector protein RABEP2 [37]. Recently, the liver kinase B1 (LBK1) has also been identified as a Rab7a effector in ECs that promotes Neuropilin-1 receptor lysosomal trafficking and degradation to inhibit angiogenesis [38]. In lung ECs, Rab4 activation and Rab9

inhibition are critical for vascular permeability through regulation of VE-Cadherin (a transmembrane adherens junction protein) localization to cell junctions [39]. Yet, the role of endolysosomal proteins in regulation of BBB function after ischemic stroke has not been explored.

In this study, we show that endothelial-specific deletion of Rab7a reduces stroke-induced TJ strand disassembly resulting in reduced paracellular BBB permeability and improved neuronal outcomes. TNFα and IL1β, two pro-inflammatory cytokines that are upregulated in the brain after ischemic stroke [13] induce Rab7a activation via Ccz1 in brain ECs *in vitro*, leading to increased TJ protein degradation and impaired paracellular barrier function. In contrast, silencing Rab7a in brain ECs *in vitro* suppresses cytokine-driven barrier dysfunction by preventing degradation of Claudin-5 protein and maintaining its localization at cell-cell junctions. These findings represent a new mechanism by which inflammatory cytokines promote degradation of select TJ proteins via Rab7a activation leading to BBB dysfunction after ischemic stroke.

RESULTS

Rab7a is essential in CNS ECs for triggering structural TJ abnormalities and increased paracellular BBB permeability 48 hours after t-MCAO.

Studies in T cells and cancer cells have found that Rab7a inhibition is critical to prevent degradation of nutrient transporters and support cell survival [27, 29, 30]. Therefore, we hypothesized that Rab7a activation in ECs may be a critical step to promote degradation of BBB TJ transmembrane proteins after ischemic stroke. To test this hypothesis, we generated a Rab7a-inducible EC knockout mouse strain (*iEC-KO - Rab7a$^{fl/fl}$; VEC-Cre$^{ERT2+/-}$; eGFP::Claudin5$^{+/-}$*) to

delete Rab7a in ECs upon administration of 4-OH-tamoxifen (**Figure 1A, top panel**). These mice

also express the eGFP::Claudin5 fusion protein in ECs to visualize TJ strand morphology [10]. We

confirmed the efficiency and specificity of Rab7a deletion in ECs by decreased Rab7a protein

expression in cortical Lectin (BSL)$^+$ blood vessels of *Rab7a iEC-KO* compared to *Rab7a $^{fl/fl}$* mice

(*Rab7a$^{fl/fl}$; eGFP-Claudin5$^{+/-}$*; referred as WT) by immunofluorescence (**Figures 1B-C",**

arrowheads). Rab7a protein was detected equally in other cell types of both WT and *iEC-KO*

cortices (**Figures 1B-C", arrows**). We induced ischemic stroke in these mice by performing t-

MCAO for 45 minutes and analyzed TJ integrity 48 hours after reperfusion (**Figure 1A, bottom**

panel), a timepoint at which BBB dysfunction is dominated mainly by TJ strand disassembly [10].

WT and *iEC-KO* mice displayed a normal BBB TJ morphology by immunofluorescence for

eGFP::Claudin5 and ZO-1 in GLUT1$^+$ vessels of the contralateral cortex (**Figure 1D-E"**). In

contrast, blood vessels in the ipsilateral somatosensory cortex of WT mice had a high fraction of

TJ strands with abnormalities, including gaps [~50%; **Figure 1F-F' (white arrows), H**] and GFP$^+$

membrane protrusions [~35%; **Figure 1F-F" (yellow arrowhead**), **I**]. The fractions of TJ strands

with gaps and protrusions were significantly lower in the ipsilateral cortex of *iEC-KO* (~18% and

20%, respectively) compared to WT mice, but they were not significantly different from those in

the contralateral cortexes of both genotypes (**Figure 1G-G", H, I**).

To determine the effects of Rab7a deletion on total levels of adherens junction (AJ; VE-

Cadherin) and TJ [Claudin-5, Occludin, and ZO-1] proteins in ECs after ischemic stroke, lysates

were collected from the ipsilateral and contralateral cortices of both genotypes at 48 hours after t-

MCAO and assessed via Western blotting (**Figure 2A**). All AJ and TJ proteins were significantly

reduced in the ipsilateral cortex of WT mice at 48 hours after t-MCAO, compared to both the WT

healthy (t-MCAO sham) and WT contralateral cortex (**Figure 2B-E**). In contrast, Claudin-5,

Occludin and ZO-1 protein levels were higher in the ipsilateral cortex of *iEC-KO* mice, and were

not significantly different from those in healthy and contralateral *iEC-KO* cortex (p>0.05; **Figure**

2B-D). Importantly, Claudin-5 protein levels were also significantly higher in the ipsilateral cortex

of *iEC-KO* compared to WT mice (p=0.04), whereas Occludin and ZO-1 protein levels showed

higher, but not significantly different, levels in the *iEC-KO* compared to WT ipsilateral cortex

[Occludin (p=0.18), ZO-1 (p=0.3)] (**Figure 2B-D**). Finally, VE-cadherin was significantly reduced

in the ipsilateral cortex of WT mice at 48 hours after t-MCAO, compared to both the WT healthy

(t-MCAO sham) and WT contralateral cortex, and its levels were not rescued by deletion of Rab7a

in ECs (**Figure 2A, E**). Therefore, Rab7a deletion in ECs rescues selectively Claudin-5, a key

BBB TJ protein [40], but is less effective for other TJ or AJ proteins after ischemic stroke.

To determine the effect of Rab7a EC-specific deletion on changes in paracellular BBB

permeability after ischemic stroke, we injected biocytin-TMR, an 890 Da dye that preferentially

extravasates blood vessels through the paracellular route[10, 11], 48 hours after t-MCAO reperfusion,

and analyzed its distribution in seven distinct bregma regions of the brain. WT mice showed an

intense biocytin-TMR fluorescence signal in the ipsilateral cortex of all analyzed bregma regions

after stroke, with extravascular leakage localized to both the cortex and the striatum in most mice

(**Figure 3A, B**). In contrast, the extravascular tracer leakage was less frequent and less extensive

in the ipsilateral cortex of all analyzed bregma regions from *iEC-KO* brains (**Figure 3C, D**). There

was an overall decrease in the volume of tracer leakage in *iEC-KO* compared to *WT* brains (**Figure**

3E). In addition, the average area of tracer leakage in each bregma region, representing the fraction

of the brain section affected by BBB impairment, and the amount of tracer leakage, measured as

fluorescence intensity of the extravasated dye in the parenchyma, were also significantly reduced

in *iEC-KO* mice (**Figure 3F, G**). Thus, Rab7a EC-specific deletion rescues structural and

functional BBB abnormalities related to TJ disassembly at 48 hours after t-MCAO.

To determine if Rab7a deletion in ECs affects exclusively paracellular BBB permeability

after ischemic stroke, we analyzed the leakage of endogenous serum immunoglobulin G (IgG) as

a readout of transcellular transport across the BBB 48 hours after reperfusion. No significant

differences in volume, area, or fluorescence intensity of serum IgG leakage were observed between

the WT and *iEC-KO* brains (**Figure S1A-G**). Correspondingly, the levels of Cav-1, a protein that

mediates transcellular transport across ECs, were comparable in lysates from the ipsilateral and

contralateral cortex of WT and *iEC-KO* mice 48 hours after t-MCAO (**Figure S1H, I**), confirming

that Rab7a EC-specific deletion did not affect the transcellular BBB leakage after ischemic stroke

[10]. Overall, Rab7a EC-specific deletion rescues significantly the paracellular, but not transcellular,

BBB permeability at the acute phase after ischemic stroke.

Rab7a iEC-KO mice show improved neuronal outcomes 48 hours after t-MCAO.

To determine whether the rescue of paracellular BBB permeability after ischemic stroke in

iEC-KO mice has a neuroprotective effect, we assessed the number of NeuN$^+$ cells, a neuronal

marker, in the ipsilateral and contralateral somatosensory cortexes of both genotypes at seven

distinct bregma regions. NeuN staining appeared missing or fragmented in the ipsilateral

somatosensory cortex of WT mice, but it was markedly rescued in the ipsilateral cortex of *iEC-*

KO mice (**Figure 4A-I'**). Quantification of the NeuN$^+$ / DAPI$^+$ cell ratio showed a significant

reduction in the ipsilateral compared to the contralateral cortex of WT mice (the number of DAPI

cells did not change significantly between these two regions) (**Figure 4J**). However, this phenotype was rescued in *iEC-KO* ipsilateral cortex where the cell ratio of NeuN$^+$ / DAPI was comparable to the contralateral cortex in all analyzed bregma regions (**Figure 4J**). Consistent with a larger number of NeuN$^+$ neurons in the cortex after ischemic stroke by immunofluorescence, *iEC-KO* showed fewer neurological deficits with a lower modified Bederson neurological score (0-5) after a series of behavioral tests (forelimb flexion, circling, resistance to lateral push as described in [41]), compared to WT mice 48 hours after t-MCAO (**Figure 4K**). These data demonstrate that *iEC-KO* have an overall improvement in neuronal outcomes after t-MCAO, likely due to a smaller stroke volume (**Figure 3E**).

To examine if the rescue in structural and functional paracellular BBB impairment observed in *iEC-KO* mice after ischemic stroke reduces local inflammation, we analyzed myeloid cells (activated microglia and infiltrating macrophages) and resting microglia in the stroke core, border and contralateral cortex of both genotypes 48 hours after t-MCAO (**Figure S2A, E**; yellow outlined area and green boxed regions 1-3). Resting microglial cells (Iba1$^+$ CD68$^-$ cells with ramified branches) were found outside the BBB leakage area both at the stroke border and in the contralateral cortex (Figure **S2C-D''', G-H'''**; white arrows). In contrast, myeloid cells (Iba1$^+$ CD68$^+$ cells with ameboid morphology representing both activated microglia and macrophages) were found in the stroke core and border within the BBB leakage area in both *WT* and *iEC-KO* cortexes (**Figure S2B-C''', F-G'''**; white arrowheads within the red border). The ratio of CD68$^+$ Iba1$^+$ over total Iba1$^+$ cells was similar in the cortex of both genotypes at all analyzed bregma regions with no significant differences between the two genotypes (**Figure S2I**). Although somewhat paradoxical, the similar local inflammatory changes found in both WT and *iEC-KO*

cortexes in the acute phase (48 hours) of ischemic stroke could be a consequence of a partial, rather

than a complete, rescue in BBB permeability in mutant mice since Rab7a EC-specific deletion

rescues significantly the paracellular, but not transcellular, BBB permeability at the acute phase of

disease.

Pro-inflammatory cytokines, TNFα and IL1β, but not glucose and oxygen deprivation,

induce Rab7a activation via upregulation of Ccz1 in brain ECs *in vitro*.

Rab7a cycles between inactive (GDP-bound) and active (GTP-bound) states to regulate

various cell biological processes [24, 25]. Rab-interacting lysosomal protein (RILP) binds selectively

to Rab7-GTP and recruits the dynein-dynactin motor complex to facilitate vesicle movement

toward the minus end of microtubules [42]. The N terminus region of RILP is required to recruit

dynein motors, but not for Rab7-GTP binding; therefore an N-terminal truncation of RILP can be

used to assess Rab7 activation in cells [29]. To investigate the molecular mechanisms responsible for

Rab7a activation in ischemic stroke, we cultured primary mouse brain ECs (mBECs) *in vitro* under

conditions of either glucose and oxygen deprivation (OGD), or inflammation, and quantified the

amount of GTP-bound Rab7a by using a GST-RILP fusion protein to pull down selectively the

active protein from the cell lysate (**Figure 5A**). First, we tested whether the ischemic environment

in stroke promotes Rab7a activation by growing mBECs under OGD conditions for 48 hours.

Surprisingly, both the total and GTP-bound Rab7a levels were decreased significantly after 48

hours of OGD treatment (**Figure 5G-I**), suggesting that ischemic conditions do not activate Rab7a

protein. The lower levels of Rab7a in OGD conditions were not associated with increased cell

death (data not shown). Since ischemic stroke induces inflammation[13], we then tested the ability

of several pro-inflammatory cytokines to induce Rab7a activation. Treatment with a combination

of TNFα and IL1β did not increase the total amount of Rab7a, but produced a trend, albeit non-

significant, towards increased GTP-bound Rab7a levels after 24 hours (**Figure 5B-D**). After 48

hours of TNFα and IL1β treatment, Rab7a-GTP levels were significantly increased without a

change in the total Rab7a levels (**Figure 5B, E, F**). Three other pro-inflammatory cytokines that

are upregulated after ischemic stroke, IL-21 [43], CCL2 [44] and IL-17A [45], had no effect on either

total or activated Rab7a levels in mBECs at 48 hours after treatment (**Figure S3**), indicating that

select pro-inflammatory cytokines promote Rab7a activation in primary mBECs.

Next, we examined expression of Ccz1, a GEF that acts together with vacuolar fusion

protein Mon1 to promote both Rab7a activation and Rab7a-mediated membrane fusion in yeast [31]

and eukaryotic cells [32-35]. Ccz1 expression was significantly increased in mBECs 24 hours after

TNFα and IL1β exposure. However, this effect was transient and Ccz1 protein returned to control

levels at 48 hours after cytokine treatment (**Figure 5J, K**). Thus, TNFα and IL1β induce expression

of Ccz1 in mBECs within 24 hours, which likely results in Rab7a activation peaking between 24-

48 hours to promote degradation of TJ proteins.

Rab7a knockdown reduces cytokine-driven EC barrier dysfunction *in vitro* by preventing

degradation of Claudin-5 protein.

To investigate the downstream effect of Rab7a activation on brain EC barrier integrity at

the cell biological level, we generated Rab7a knockdown primary mBECs via transfection of

stealth RNAi duplexes targeting the *Rab7a* mRNA transcript. We verified the efficiency of Rab7a

protein downregulation by immunofluorescence and Western blotting, and found that Rab7a

expression was decreased by 75% in mBECs transfected with the siRNA (siRab7a) compared to

those transfected with a scrambled siRNA (siCTRL). This knockdown efficiency was observed in the presence and absence of TNFα and IL1β (**Figure S4A-F**). We examined whether Rab7a knockdown affected the levels of other proteins in the endolysosomal pathway, such as the early endosomal marker EEA-1 and the lysosomal marker LAMP-1 by Western blotting. We confirmed that EEA-1 and LAMP-1 protein levels were not different in both siCTRL- and siRab7a-transfected mBECs with or without cytokine treatment (**Figure S4E, G, H**). Since Rab7a activity maintains lysosome mass, we quantified the lysosome mass per cell in both siCTRL- and siRab7a-transfected mBECs after immunostaining for the pH sensor Lysotracker. The lysosome mass per cell was similar between siCTRL and siRab7a treatments with or without cytokine treatment (**Figure S4I-M**). Together, these data demonstrate that Rab7 knockdown does not affect other endolysosomal proteins and the lysosome mass in mBECs.

Since Rab7a plays a critical role in autophagy (reviewed in [46]), we examined whether Rab7a knockdown affected expression of the two autophagy proteins Atg5 and p62 under basal and inflammatory conditions in mBECs. Under basal conditions, the levels of Atg5 and p62 proteins were similar in both siCTRL- and siRab7a-transfected mBECs (**Figure S5A-C**). As expected, the Atg5 and p62 protein levels were upregulated under inflammatory conditions in both siCTRL- and siRab7a-transfected mBECs (**Figure S5A-C**). Moreover, p62 levels were significantly higher in siRab7a-transfected mBECs after cytokine treatment, suggesting that Rab7a knockdown induced p62 accumulation in mBECs (**Figure S5A-C**), likely due to reduced autophagy.

To determine if Rab7a knockdown rescues cytokine-mediated increase in mBEC

paracellular permeability we analyzed the transendothelial electrical resistance (TEER) in

siCTRL- and siRab7a-transfected cells after cytokine treatment. In the absence of cytokines,

siRab7a-transfected mBECs show a decreased basal TEER compared to siCTRL-transfected cells

(**Figure 6A-C**). Treatment of siCTRL-transfected mBECs with TNFα and IL1β for 48 hours

significantly reduced the TEER. However, this cytokine mediated disruption was mitigated in

siRab7a-transfected mBECs (**Figure 6A-C**). To investigate which TJ proteins are regulated by

Rab7a in mBECs, we examined expression and subcellular localization of TJ proteins (Claudin-5,

Occludin and ZO-1) at 48 hours after cytokine addition. Cytokine treatment significantly reduced

the levels of Claudin-5 and ZO-1, but not Occludin, proteins in mBECs. Furthermore, transfection

with siRab7a rescued cytokine-induced degradation of Claudin-5, but not ZO-1, protein levels in

mBECs (**Figure 6D-G**).

At the basal level, siRab7a-transfected mBECs showed irregular cell junctions compared

to siCTRL-transfected cells (**Figure 6H-H", J-J"**), a phenotype resembling the lack of ZO-1/ZO-

2 in epithelial cells [47], although ZO-1 protein co-localized with Claudin-5. This observation likely

explains the lower TEER seen with siRab7a-treated mBECs under basal conditions (**Figure 6A**).

Treatment with inflammatory cytokines induced the appearance of gaps and protrusions in

Claudin-5[+] cell junction strands in siCTRL-treated mBECs, and loss of ZO-1 protein from most

cell junctions (**Figure 6I-I"**). Although cytokine-treated Rab7a-deficient mBECs show an

exacerbated jagged cell junction phenotype due to the absence of ZO1, very few gaps and

protrusions in Claudin-5[+] strands were observed after exposure to pro-inflammatory cytokines

(**Figure 6K-K"**). Cytokine treatment, however, downregulated potently ZO-1 protein in both

control and siRab7a-transfected mBECs (**Figure 6I-I", K-K"**), indicating that Rab7a knockdown

has very selective effects on degradation of distinct TJ proteins.

Since Cav-1 protein has been shown to regulate Claudin-5 protein levels in mBECs after

exposure to inflammatory chemokines (e.g. CCL2)[17], we examined its levels and the caveolar-

mediated transcytosis in Rab7a knockdown cells. Cav-1 protein levels were increased in primary

BECs after exposure to TNFα and IL1β independently of Rab7a levels (**Figure S5I-J**). In addition,

we performed an albumin-594 uptake assay to measure Caveolin-1-dependent transcellular

transport, and confirmed that albumin-594 uptake was equally increased in siCTRL- and siRab7a-

transfected mBECs upon treatment with cytokines (**Figure S5D-H**). Therefore, the rescue of

Claudin-5 degradation with pro-inflammatory cytokines following Rab7a knockdown in mBECs,

is not due to changes in Caveolin-1 protein levels and caveolar-mediated transport. In conclusion,

the *in vivo* and *in vitro* data demonstrate that Rab7a is necessary for cytokine-induced degradation

of select BBB TJ proteins (e.g. Claudin-5) and enhanced paracellular permeability in mouse brain

endothelial cells, but is not involved in the increased caveolar transcellular transport in brain ECs.

DISCUSSION

The disassembly of cell junctions due to persistent degradation of adherens and tight junction

proteins has long been considered as the main contributor to increased BBB permeability in

multiple neuropathological conditions, including ischemic stroke. However, the cell biological

mechanisms driving this process are still debated. OGD, perhaps the most obvious trigger, has

been proposed to promote junctional protein degradation through activation of several proteases

[48, 49]. Surprisingly, we find that OGD does not activate Rab7a in primary brain endothelial cells,

suggesting that OGD-induced junctional protein degradation does not require Rab7a activation.

Our findings support previous studies that two pro-inflammatory cytokines, IL-1β and TNF-α,

which are upregulated in stroke tissue between 24-48 hours after reperfusion and coincide with

immune cell infiltration into the CNS (reviewed in [12, 13]), induce internalization and degradation

of BBB TJ proteins [17, 18]. We further demonstrate that IL-1β and TNF-α, but not other pro-

inflammatory cytokines found in the stroke milieu (e.g. CCL2, IL-21, IL-17A), induce Rab7a

activation likely via GEF Ccz1 [31-33], which then drives degradation of selective BBB cell junction

proteins. Other pro-inflammatory cytokines/chemokines that have been implicated in BBB

dysfunction such as IL-17A, IL-21 and CCL2 may promote TJ degradation through Rab7a-

independent mechanisms, since they cannot activate Rab7a in primary brain ECs. Therefore, pro-

inflammatory cytokines likely mediate BBB TJ degradation at the acute phase of ischemic stroke

through both Rab7a-dependent and -independent mechanisms.

Cav-1 has been previously implicated in tight junction transmembrane protein

internalization and degradation [17, 18, 48]. Our data demonstrate that Rab7a-mediated TJ rescue is not

accompanied by changes in Caveolin-1 protein levels or caveolar-mediated transcellular

permeability *in vitro* and *in vivo* (**Figures S1I, S5I**). Moreover, the pro-inflammatory chemokine

CCL2 is unable to induce Rab7a activation in primary brain endothelial cells (**Figure S3D-F**),

although it has been shown to promote Caveolin-1-mediated internalization of Claudin-5 in the

bEnd3.0 cell line [17]. Although we cannot exclude the possibility that Caveolin-1-dependent cargo

sorting is a putative mechanism of junctional protein degradation under some conditions, it is likely

not the predominant mechanism by which inflammation-mediated degradation of cell junction

proteins occurs in the setting of acute ischemic stroke. Thus, the cellular mechanisms of BBB TJ

protein degradation are context dependent and likely differ across distinct neurological diseases.

Overall our findings *in vivo* and *in vitro* suggest a model for how the state of Rab7a

activation in brain ECs regulates recycling and degradation of BBB TJ proteins under healthy and

disease states (**Figure 6L**). We propose that under physiological conditions, similar levels of active

and inactive Rab7a protein levels regulate a constant internalization, recycling and degradation of

transmembrane adherens and tight junction proteins, in order to maintain their steady-state levels

at cell junctions (**Figure 6L, left panel**). When ECs are exposed to a subset of pro-inflammatory

cytokines (e.g. TNFα and IL1β), such as under acute conditions of ischemic stroke, they

upregulate Ccz1 expression, a GEF which promotes Rab7a activation [31-33]. Rab7a hyperactivation

results in an imbalance of the endolysosomal pathway towards degradation rather than recycling,

drives cell junction transmembrane proteins to the lysosomes, and triggers tight junction

dismantlement which further exacerbates paracellular BBB permeability (**Figure 6L, right panel**).

Our study identifies a new molecular mechanism controlling EC junctional integrity in diseased

BBB, and establishes Rab7a as a key regulator of TJ protein turnover via the endolysosomal

pathway in CNS inflammation. These findings have implications for not only ischemic stroke, but

potentially other neurological disorders characterized by inflammation and BBB dysfunction. For

example, in EAE, structural BBB TJ abnormalities precede the onset of disease and persist

throughout the course of EAE to allow immune cell infiltration into the CNS [11], emphasizing a

critical role for pro-inflammatory mechanisms in driving TJ strand disassembly. Based on our

findings in this study, we postulate that early and persistent Rab7a activation due to pro-

inflammatory cytokines Il-1β and TNF-α may likely drive degradation of BBB TJ transmembrane

proteins in EAE leading to early and persistent BBB leakage [11].

The proposed mechanism by which Rab7a activation triggers degradation of junctional

proteins to promote persistent dismantlement of cell junctions implies a potential lack of

selectivity, since the endolysosomal degradation is a fundamental cell biological process.

However, our genetic loss-of-function data suggest that Rab7a elimination both *in vivo* and *in vitro*

rescues both the levels and localization of selective transmembrane cell junction proteins such as

Claudin-5, a key BBB TJ protein. In contrast, Rab7a elimination does not rescue significantly the

levels of Occludin (a transmembrane TJ protein) and ZO-1 (an intracellular TJ component that

anchors transmembrane junction proteins to the actin cytoskeleton). This discrepancy could result

because either low levels of Rab7a may be sufficient to degrade Occludin and ZO-1 protein, or

their degradation could be mediated by a Rab7a-independent mechanism under inflammatory

conditions. Moreover, Rab7a knockdown has no effect on the levels of Caveolin-1 or Slc2a1 (Glut-

1), which are both integral membrane proteins. A possible explanation is that TNFα, which

activates Rab7a, also promotes the formation of actin stress fibers [50] that further induce

disassembly and internalization of junctional proteins [51]. In conclusion, our findings represent a

new mechanism by which inflammatory cytokines promote degradation of select TJ proteins via

Rab7a activation, leading to BBB dysfunction after ischemic stroke. Recently, a small chemical

molecule (CID 1067700) has been identified as a potent inhibitor of Rab7a-GTPase activity with

(K_i) values that range from 12.90-19.70 nM [52]. Future studies will determine whether this

compound can reduce or prevent BBB breakdown and disease outcomes after ischemic stroke.

ACKNOWLEDGEMENTS

We thank Martin Hsu (UNC) and Julian Smith (University of Washington) for initial help with the

project and the generation of *Rab7a-iECKO* mice, respectively; Tyler Cutforth for scientific and

editorial feedback on this project. A.C., S.S., M.G., G.P. and D.A. are supported by grants from

the National Eye Institute (R01EY033994), National Heart Lung and Blood Institute

(R61HL159949), National Institute of Aging (RF1AG078352), National Institute of Neurological

Disorders and Stroke (R21NS130265), the Leducq Foundation (FDNLEDQ 15CVD 02) and in

part by a gift donation from the Ju Foundation. Development of the Rab floxed mouse was

supported by a grant from the National Cancer Institute to ALE (K08 CA100526).

AUTHOR CONTRIBUTIONS

Conceptualization: A.C., D.A.; Methodology: A.C., S.S., A.A., M.G., G.P., J.N., M.C.T., A. L. E.,

D.A.; Formal analysis: A.C., S.S., M.G., M.C.T.; Investigation: A.C., S.S., A.A., M.G., M.C.T.;

Resources: A.L.E, D.A.; Data curation: A.C., S.S., M.G., M.C.T.; Writing - original draft: A.C.,

D.A.; Writing - review & editing: A.C., S.S., A.A., M.C.T., A.L.E, M.C.T., D.A.; Visualization:

A.C.; Supervision: D.A., Project administration: D.A., Funding acquisition: D.A.

DECLARATION OF INTEREST

The authors declare no competing or financial interests relevant to the subject matter in this

contribution.

 FIGURES AND FIGURE LEGENDS

Figure 1. Rab7a is essential for structural abnormalities found in BBB tight junctions 48 hours after t-MCAO.

(**A**) Diagrams showing the breeding strategy to generate the inducible endothelial Rab7a knockout (*Rab7a iEC-KO*) mice and the experimental setup to analyze the paracellular blood-brain barrier (BBB) permeability after t-MCAO. (**B-C"**) Immunofluorescence images for Griffonia

(Bandeiraea) Simplicifolia Lectin I (BSL, green), Rab7a (red) and DAPI (blue) in the cortex of

healthy WT (**B-B"**) and *Rab7a iEC-KO* (**C-C"**) mice. Cortical vessels are delineated by yellow

dotted lines (**B'**, **C'**). Boxed areas are magnified in the orthogonal view images on the right (**B"**,

C"). Arrowheads indicate endothelial Rab7a, arrows indicate non-endothelial Rab7a. There is very

little residual Rab7a protein in ECs of mutant mice. Scale bar: 30 μm. (**D-G"**)

Immunofluorescence images for eGFP-Claudin5 (green), ZO-1 (blue) and BSL (red) in the

contralateral (**D-E"**) and ipsilateral (**F-G"**) cortex of WT and *Rab7a iEC-KO* mice 48 hours after

t-MCAO. White arrows point to gaps, yellow arrowheads show protrusions in BBB tight junction

strands. Scale bar: 25 μm. (**H**, **I**) Quantification of junction gaps (**H**) and protrusions (**I**) in the

contralateral and ipsilateral cortex of *WT* and *Rab7a iEC-KO* mice 48 hours after t-MCAO. Each

dot represents an animal (n=7-8 mice / group). Data are means ± s.e.m. *: $p<0.05$; n.s.: $p>0.05$;

one-way ANOVA with post-hoc Tukey's correction.

Figure 2. Rab7a regulates degradation of selective BBB TJs 48 hours after t-MCAO.

(**A**) Western blots for BBB junctional proteins ZO-1, VE-cadherin, Occludin and Claudin-5 and β-actin (control) of brain lysates collected from either the healthy or contralateral and ipsilateral ischemic cortex of *WT* and *Rab7a iEC-KO* mice 48 hours after t-MCAO. The respective molecular

weights for each protein are shown on the right. (**B-E**) Quantification of Claudin-5, Occludin, ZO-1 and VE-cadherin protein levels in either healthy, contralateral or ipsilateral ischemic brain lysates of *WT* and *Rab7a iEC-KO* mice 48 hours after t-MCAO. Each dot represents an animal (n=4 mice / group). Data are means ± s.e.m and normalized to WT healthy levels. ***: p<0.001; **: p<0.01; *: p<0.05; n.s.: p>0.05; one-way ANOVA with post-hoc Tukey's correction.

Figure 3. *Rab7a iEC-KO* **mice show a partial rescue in the paracellular BBB permeability 48 hours after t-MCAO.**

(**A-D**) Fluorescent micrographs and heatmaps showing biocytin-TMR tracer extravasation in seven distinct brain regions in WT (**A, B**) and *Rab7a iEC-KO* (**C, D**) mice 48 hours after t-MCAO (biocytin-TMR was injected 30 minutes before analysis). Dotted lines outline the border of the brain section and the leakage area. The heatmaps show the fraction of animals displaying BBB leakage in each region represented as a scale (0-100%) of red hues. (**E**) Quantification of biocytin-TMR leakage volume in the brain of WT and *Rab7a iEC-KO* mice 48 hours after t-MCAO. n=8 animals/group. Data are means ± s.e.m. *: p<0.05; Student's t-student. (**F**) Quantification of biocytin-TMR leakage area in each bregma region from WT and *Rab7a iEC-KO* mice 48 hours after t-MCAO. Each dot represents the average of n=8 animals / group. Data are means ± s.e.m. *: p<0.05; one-way ANOVA with post-hoc Tukey's correction. (**G**) Quantification of the intensity of biocytin-TMR leakage in the ipsilateral cortex of each bregma region from WT and *Rab7a iEC-KO* mice 48 hours after t-MCAO. The dotted line represents the average fluorescence intensity in the contralateral cortex. Each dot represents the average of n=8 animals. Data are means ± s.e.m. *: p<0.05; one-way ANOVA with post-hoc Tukey's correction. Please see also **Figure S1**.

Figure 4. *Rab7a iEC-KO* **mice show improved neuronal survival and neurological score 48 hours after t-MCAO.**

(**A-I'**) Immunofluorescence analysis for NeuN (green) and DAPI (blue) in the contralateral and ipsilateral cortex of WT and *Rab7a iEC-KO* mice 48 hours after t-MCAO. (**A, D, G**) Diagrams illustrate the bregma regions and the red boxes outline the cortical area shown in each micrograph in **B-I'**. (**B-I'**) White arrows indicate abnormal and white arrowheads indicate normal NeuN staining in the ipsilateral cortex. Scale bar: 25 μm. (**J**) Quantification of the fraction of viable neurons (NeuN/DAPI ratio) in the contralateral (dotted lines) and ipsilateral (solid lines) cortexes in seven bregma regions of WT and *Rab7a iEC-KO* mice 48 hours after t-MCAO. Each dot represents the average of n=8 animals / group. Data are means ± s.e.m. *: $p<0.05$; one-way

490 ANOVA with post-hoc Tukey's correction. (**K**) Analysis of neurological score in WT and *Rab7a*

491 *iEC-KO* mice 24 and 48 hours after t-MCAO. Each dot represents one animal (n=8 mice / group).

492 Data are medians ± 95% C.I. *: $p<0.05$; **: $p<0.01$; Kruskal-Wallis one-way ANOVA. Please see

493 also **Figure S2**.

Figure 5. Proinflammatory cytokines TNFα and IL1β induce Rab7a activation in mouse brain endothelial cells via upregulation of Ccz1.

(**A**) Schematic diagram of the experimental setup. Primary mouse brain endothelial cells (mBECs) are grown to confluence and switched to low-serum media overnight prior to addition of pro-inflammatory cytokines (TNFα, IL-1β; 10 ng/mL). Protein lysates are collected after 24 or 48 hours of cytokine treatment. Rab7a-GTP is pulled down using a GST-RILP fusion protein that binds glutathione immobilized on sepharose beads. Western blotting is then used to detect both Rab7a and Rab7a-GTP protein levels. (**B**) Western blot of active and total Rab7a proteins in mBECs treated with TNFα and IL1β for either 24 or 48 hours and untreated control cells. GST and β-actin are used to normalize active Rab7a and total Rab7a levels, respectively. The molecular weight of each protein is shown on the right. (**C-F**) Quantification of total (**C, E**) and active (**D, F**) Rab7a levels in mBECs treated with TNFα and IL1β for 24 hours (**C, D**) and 48 hours (**E, F**) and untreated cells. Each dot represents an independent experiment. Data are means ± s.e.m. (**F**) ***: $p<0.005$; Student's t-test. (**G**) Western blot for active and total Rab7a in mBECs grown in oxygen and glucose deprivation (OGD) conditions for 48 hours and untreated control cells. GST and β-actin are used to normalize active Rab7a and total Rab7a levels, respectively. The molecular weight of each protein is shown on the right. (**H, I**) Quantification of total (**H**) and active (**I**) Rab7a levels in mBECs grown in OGD conditions for 48 hours and control cells. Each dot represents an independent experiment. Data are means ± s.e.m.; **: $p<0.01$; *: $p<0.05$; n.s. $p>0.05$ Student's t-test. (**J**) Western blot for Ccz1 in mBECs treated with TNFα and IL1β for 24 and 48 hours and untreated cells. (**K**) Quantification of Ccz1 protein levels in mBECs treated with TNFα and IL1β for 24 and 48 hours and untreated cells. Each dot represents an independent experiment. Data are means ± s.e.m. ****: $p<0.001$; one-way ANOVA with post-hoc Tukey's correction. Please see also **Figure S3**.

Figure 6. Rab7a knockdown reduces cytokine-driven EC barrier dysfunction by preventing degradation of Claudin-5 protein in mouse brain endothelial cells.

(**A**) Transendothelial electrical resistance (TEER) of primary mouse brain endothelial cells (mBECs) transfected with either an siRNA targeting Rab7a (siRab7a), or a scrambled siRNA (siCTRL) in the absence or presence of inflammatory cytokines (TNFα, IL-1β 10 ng/mL). The gray area represents the period of time during which cells were treated with cytokines or vehicle. (**B, C**) Quantification of the area under the curve (A.U.C.) during treatment with cytokines [gray area in (a)] and of the percentage of TEER reduction 48 hours after treatment with cytokines (right panel). Each dot represents an independent experiment. Data are means ± s.e.m. **: $p<0.01$; *: $p<0.05$; n.s. $p>0.05$; one-way ANOVA with post-hoc Tukey's correction (left graph), and Student's t-test (right graph). (**D**) Western blot for junctional proteins ZO-1, Occludin and Claudin-5 in siCTRL- and siRab7a-transfected mBECs with or without cytokines. β-actin serves as control. The molecular weights of the proteins are shown on the right. (**E-G**) Quantification of Claudin-5, Occludin and ZO-1 levels in siCTRL- and siRab7a-transfected with or without cytokines. Each dot represents an independent experiment (n=4). Data are means ± s.e.m.; ***: $p<0.001$; **: $p<0.01$; *: $p<0.05$; n.s. $p>0.05$; one-way ANOVA with post-hoc Tukey's correction. (**H-K"**) Immunofluorescence for Claudin 5 (green) and ZO-1 (red) in mBECs transfected with either siCTRL (**H-I"**) or siRab7a (**J-K"**) in the presence or absence of cytokines. Cytokine treatment induces fewer structural abnormalities in Claudin-5 localization at cell junctions of siRab7a-treated mBECs. (**L**) Working model for the role of Rab7a in tight junction instability after ischemic stroke. In healthy conditions, Rab7a cycles equally between its active and inactive states, balancing the rates of degradation and recycling of internalized membrane proteins including adherens and tight junction proteins. Under pathological conditions, such as ischemic stroke, some inflammatory cytokines (e.g. IL-1β, TNFα) induce Ccz1, a GEF that activates Rab7a. The hyperactivated Rab7a

546 increases the rate of degradation of junctional proteins leading to increased paracellular BBB

547 permeability. Please see also **Figures S4** and **S5**.

REFERENCES

1. Liebner, S. *et al.* Functional morphology of the blood-brain barrier in health and disease. *Acta Neuropathol* **135**, 311-336 (2018).

2. Bamford, J., Sandercock, P., Dennis, M., Burn, J. & Warlow, C. A prospective study of acute cerebrovascular disease in the community: the Oxfordshire Community Stroke Project--1981-86. 2. Incidence, case fatality rates and overall outcome at one year of cerebral infarction, primary intracerebral and subarachnoid haemorrhage. *J Neurol Neurosurg Psychiatry* **53**, 16-22 (1990).

3. Bonita, R., Solomon, N. & Broad, J.B. Prevalence of stroke and stroke-related disability. Estimates from the Auckland stroke studies. *Stroke* **28**, 1898-1902 (1997).

4. Kammersgaard, L.P. *et al.* Short- and long-term prognosis for very old stroke patients. The Copenhagen Stroke Study. *Age Ageing* **33**, 149-154 (2004).

5. Rothwell, P.M. *et al.* Population-based study of event-rate, incidence, case fatality, and mortality for all acute vascular events in all arterial territories (Oxford Vascular Study). *Lancet* **366**, 1773-1783 (2005).

6. Kaur, C. & Ling, E.A. Blood brain barrier in hypoxic-ischemic conditions. *Current neurovascular research* **5**, 71-81 (2008).

7. Sandoval, K.E. & Witt, K.A. Blood-brain barrier tight junction permeability and ischemic stroke. *Neurobiol Dis* **32**, 200-219 (2008).

8. Arai, K., Jin, G., Navaratna, D. & Lo, E.H. Brain angiogenesis in developmental and pathological processes: neurovascular injury and angiogenic recovery after stroke. *The FEBS journal* **276**, 4644-4652 (2009).

9.   Hawkins, B.T. & Davis, T.P. The blood-brain barrier/neurovascular unit in health and
disease. *Pharmacological reviews* **57**, 173-185 (2005).

10.   Knowland, D. *et al.* Stepwise recruitment of transcellular and paracellular pathways
underlies blood-brain barrier breakdown in stroke. *Neuron* **82**, 603-617 (2014).

11.   Lutz, S.E. *et al.* Caveolin1 Is Required for Th1 Cell Infiltration, but Not Tight Junction
Remodeling, at the Blood-Brain Barrier in Autoimmune Neuroinflammation. *Cell Rep* **21**,
2104-2117 (2017).

12.   Doll, D.N., Barr, T.L. & Simpkins, J.W. Cytokines: their role in stroke and potential use
as biomarkers and therapeutic targets. *Aging Dis* **5**, 294-306 (2014).

13.   Rayasam, A. *et al.* Immune responses in stroke: how the immune system contributes to
damage and healing after stroke and how this knowledge could be translated to better
cures? *Immunology* **154**, 363-376 (2018).

14.   Iadecola, C., Buckwalter, M.S. & Anrather, J. Immune responses to stroke: mechanisms,
modulation, and therapeutic potential. *J Clin Invest* **130**, 2777-2788 (2020).

15.   Doyle, K.P. & Buckwalter, M.S. Immunological mechanisms in poststroke dementia. *Curr*
Opin Neurol **33**, 30-36 (2020).

16.   Endres, M. *et al.* Immune Pathways in Etiology, Acute Phase, and Chronic Sequelae of
Ischemic Stroke. *Circ Res* **130**, 1167-1186 (2022).

17.   Stamatovic, S.M., Keep, R.F., Wang, M.M., Jankovic, I. & Andjelkovic, A.V. Caveolae-
mediated internalization of occludin and claudin-5 during CCL2-induced tight junction
remodeling in brain endothelial cells. *J Biol Chem* **284**, 19053-19066 (2009).

18.   Marchiando, A.M. *et al.* Caveolin-1-dependent occludin endocytosis is required for TNF-
induced tight junction regulation in vivo. *J Cell Biol* **189**, 111-126 (2010).

19.   Drab, M. *et al.* Loss of caveolae, vascular dysfunction, and pulmonary defects in caveolin-

1 gene-disrupted mice. *Science* **293**, 2449-2452 (2001).

20.   Razani, B. *et al.* Caveolin-1 null mice are viable but show evidence of hyperproliferative

and vascular abnormalities. *J Biol Chem* **276**, 38121-38138 (2001).

21.   Jasmin, J.F. *et al.* Caveolin-1 deficiency increases cerebral ischemic injury. *Circ Res* **100**,

721-729 (2007).

22.   Li, X., Garrity, A.G. & Xu, H. Regulation of membrane trafficking by signalling on

endosomal and lysosomal membranes. *J Physiol* **591**, 4389-4401 (2013).

23.   Zerial, M. & McBride, H. Rab proteins as membrane organizers. *Nat Rev Mol Cell Biol* **2**,

107-117 (2001).

24.   Zhang, M., Chen, L., Wang, S. & Wang, T. Rab7: roles in membrane trafficking and

disease. *Biosci Rep* **29**, 193-209 (2009).

25.   Gutierrez, M.G., Munafo, D.B., Beron, W. & Colombo, M.I. Rab7 is required for the

normal progression of the autophagic pathway in mammalian cells. *J Cell Sci* **117**, 2687-

2697 (2004).

26.   Edinger, A.L., Cinalli, R.M. & Thompson, C.B. Rab7 prevents growth factor-independent

survival by inhibiting cell-autonomous nutrient transporter expression. *Developmental cell*

5, 571-582 (2003).

27.   Romero Rosales, K., Peralta, E.R., Guenther, G.G., Wong, S.Y. & Edinger, A.L. Rab7

activation by growth factor withdrawal contributes to the induction of apoptosis. *Mol Biol*

Cell **20**, 2831-2840 (2009).

28.   Ceresa, B.P. & Bahr, S.J. rab7 activity affects epidermal growth factor:epidermal growth

616   factor receptor degradation by regulating endocytic trafficking from the late endosome. *J*

617   *Biol Chem* **281**, 1099-1106 (2006).

618   29.   Peralta, E.R., Martin, B.C. & Edinger, A.L. Differential effects of TBC1D15 and

619   mammalian Vps39 on Rab7 activation state, lysosomal morphology, and growth factor

620   dependence. *J Biol Chem* **285**, 16814-16821 (2010).

621   30.   Roy, S.G., Stevens, M.W., So, L. & Edinger, A.L. Reciprocal effects of rab7 deletion in

622   activated and neglected T cells. *Autophagy* **9**, 1009-1023 (2013).

623   31.   Nordmann, M. *et al.* The Mon1-Ccz1 Complex Is the GEF of the Late Endosomal Rab7

624   Homolog Ypt7. *Current Biology* **20**, 1654-1659 (2010).

625   32.   Kinchen, J.M. & Ravichandran, K.S. Identification of two evolutionarily conserved genes

626   regulating processing of engulfed apoptotic cells. *Nature* **464**, 778-782 (2010).

627   33.   Poteryaev, D., Datta, S., Ackema, K., Zerial, M. & Spang, A. Identification of the Switch

628   in Early-to-Late Endosome Transition. *Cell* **141**, 497-508 (2010).

629   34.   Langemeyer, L. *et al.* A conserved and regulated mechanism drives endosomal Rab

630   transition. *Elife* **9** (2020).

631   35.   Kiontke, S. *et al.* Architecture and mechanism of the late endosomal Rab7-like Ypt7

632   guanine nucleotide exchange factor complex Mon1–Ccz1. *Nature Communications* **8**,

633   14034 (2017).

634   36.   Jopling, H.M. *et al.* Rab GTPase regulation of VEGFR2 trafficking and signaling in

635   endothelial cells. *Arterioscler Thromb Vasc Biol* **29**, 1119-1124 (2009).

37. Kofler, N. *et al.* The Rab-effector protein RABEP2 regulates endosomal trafficking to
mediate vascular endothelial growth factor receptor-2 (VEGFR2)-dependent signaling. *J*
Biol Chem **293**, 4805-4817 (2018).

38. Okon, I.S. *et al.* Protein kinase LKB1 promotes RAB7-mediated neuropilin-1 degradation
to inhibit angiogenesis. *The Journal of clinical investigation* **124**, 4590-4602 (2014).

39. Chichger, H., Braza, J., Duong, H., Boni, G. & Harrington, E.O. Select Rab GTPases
Regulate the Pulmonary Endothelium via Endosomal Trafficking of Vascular Endothelial-
Cadherin. *Am J Respir Cell Mol Biol* **54**, 769-781 (2016).

40. Nitta, T. *et al.* Size-selective loosening of the blood-brain barrier in claudin-5-deficient
mice. *J Cell Biol* **161**, 653-660 (2003).

41. Jiang, S.X. *et al.* Chlortetracycline and demeclocycline inhibit calpains and protect mouse
neurons against glutamate toxicity and cerebral ischemia. *J Biol Chem* **280**, 33811-33818
(2005).

42. Cantalupo, G., Alifano, P., Roberti, V., Bruni, C.B. & Bucci, C. Rab-interacting lysosomal
protein (RILP): the Rab7 effector required for transport to lysosomes. *EMBO J* **20**, 683-
693 (2001).

43. Clarkson, B.D. *et al.* T cell-derived interleukin (IL)-21 promotes brain injury following
stroke in mice. *J Exp Med* **211**, 595-604 (2014).

44. Li, L., Lou, W., Li, H., Zhu, Y. & Huang, X. Upregulated C-C Motif Chemokine Ligand 2
Promotes Ischemic Stroke via Chemokine Signaling Pathway. *Ann Vasc Surg* **68**, 476-486
(2020).

45.   Lin, Y. *et al.* Critical role of astrocytic interleukin-17 A in post-stroke survival and
658   neuronal differentiation of neural precursor cells in adult mice. *Cell Death & Disease* **7**,
659   e2273-e2273 (2016).

660   46.   Kuchitsu, Y. & Fukuda, M. Revisiting Rab7 Functions in Mammalian Autophagy: Rab7
661   Knockout Studies. *Cells* **7** (2018).

662   47.   Umeda, K. *et al.* ZO-1 and ZO-2 independently determine where claudins are polymerized
663   in tight-junction strand formation. *Cell* **126**, 741-754 (2006).

664   48.   Liu, J., Jin, X., Liu, K.J. & Liu, W. Matrix metalloproteinase-2-mediated occludin
665   degradation and caveolin-1-mediated claudin-5 redistribution contribute to blood-brain
666   barrier damage in early ischemic stroke stage. *J Neurosci* **32**, 3044-3057 (2012).

667   49.   Underly, R.G. *et al.* Pericytes as Inducers of Rapid, Matrix Metalloproteinase-9-Dependent
668   Capillary Damage during Ischemia. *J Neurosci* **37**, 129-140 (2017).

669   50.   Abdullah, Z., Rakkar, K., Bath, P.M. & Bayraktutan, U. Inhibition of TNF-alpha protects
670   in vitro brain barrier from ischaemic damage. *Mol Cell Neurosci* **69**, 65-79 (2015).

671   51.   Shi, Y. *et al.* Endothelium-targeted overexpression of heat shock protein 27 ameliorates
672   blood-brain barrier disruption after ischemic brain injury. *Proc Natl Acad Sci U S A* **114**,
673   E1243-E1252 (2017).

674   52.   Agola, J.O. *et al.* A competitive nucleotide binding inhibitor: in vitro characterization of
675   Rab7 GTPase inhibition. *ACS Chem Biol* **7**, 1095-1108 (2012).

676   53.   Wang, Y. *et al.* Ephrin-B2 controls VEGF-induced angiogenesis and lymphangiogenesis.
677   *Nature* **465**, 483-486 (2010).

678   54.   Arac, A. *et al.* Systemic augmentation of alphaB-crystallin provides therapeutic benefit

679   twelve hours post-stroke onset via immune modulation. *Proc Natl Acad Sci U S A* **108**,

680   13287-13292 (2011).

681   55.   Mazzoni, J. *et al.* The Wnt Inhibitor Apcdd1 Coordinates Vascular Remodeling and Barrier

682   Maturation of Retinal Blood Vessels. *Neuron* **96**, 1055-1069 e1056 (2017).

STAR ★ METHODS

Detailed methods are provided in the online version of this paper and include the following:

KEY RESOURCES TABLE

RESOURCE AVAILABILITY

- o Lead contact

- o Materials availability

- o Data and code availability

EXPERIMENTAL MODEL AND SUBJECT DETAILS

- o Mice

METHOD DETAILS

- o Mouse ischemic stroke model

- o Biocytin-TMR and serum IgG permeability analysis

- o Immunofluorescence

- o Western blotting

- o mBEC culture and Rab7a silencing *in vitro*

- o Rab7a-GTP pull down assay

- o Trans-endothelial electrical resistance

- o *In vitro* assays with primary mouse BECs

QUANTIFICATION AND STATISTICAL ANALYSIS

- o Image acquisition and quantification

- o Statistical analysis

708 **STAR ★ METHODS**

709 **KEY RESOURCES TABLE**

Reagent or resource	Source	Identifier
Primary antibodies		
Rabbit anti-Rab7a	Abcam	Cat #ab137029 RRID: AB_2629474
Mouse anti-ZO-1	Thermo Fisher Scientific	Cat #33-9100, RRID: AB_2533147
Rabbit anti-ZO-1	Thermo Fisher Scientific	Cat# 61-7300, RRID: AB_2533938
Rabbit anti-GLUT1	Millipore	Cat# 07-1401, RRID: AB_1587074
Rabbit anti-Claudin-5	Thermo Fisher Scientific	Cat #34-1600, RRID: AB_2533157
Rabbit anti-VE-cadherin	Abcam	Cat# ab33168, RRID: AB_870662
Rat anti-CD68	Abcam	Cat# ab53444, RRID: AB_869007
Rabbit anti-Iba1	FUJIFILM Wako Shibayagi	Cat# 019-19741, RRID: AB_839504
Mouse anti-NeuN	Sigma-Aldrich	Cat #MAB377, RRID: AB_2298772
Mouse anti-Beta-actin	Novus	Cat# NB600-501,

		RRID: AB_10077656
Rabbit anti-Caveolin 1	Abcam	Cat# ab18199, RRID: AB_444314
Mouse anti-GST	Santa Cruz Biotechnology	Cat# sc-57590, RRID: AB_783590
Rabbit anti-Occludin	Thermo Fisher Scientific	Cat# 71-1500, RRID: AB_2533977
Rabbit anti-Ccz1	Abcam	Cat# ab131311, RRID: AB_11158512
Rabbit anti-EEA1	Abcam	Cat# ab2900, RRID: AB_2262056
Rabbit anti-LAMP1	Abcam	Cat# ab24170, RRID: AB_775978
Rabbit anti-ATG5	Cell Signaling Technology	Cat# 12994, RRID: AB_2630393
Rabbit anti-p62	Abcam	Cat# ab109012, RRID: AB_2810880
Chemicals, peptides, and recombinant proteins		
Biocytin-TMR (5-(and-6)-tetramethylrhodamine biocytin)	ThermoFisher Scientific	Cat #T12921
Albumin from Bovine Serum (BSA), AlexaFluor™ 594 conjugate	ThermoFisher Scientific	Cat #A13101

Griffonia (Bandeiraea) Simplicifolia Lectin I (GSL I, BSL I), Rhodamine	Vector Labs	Cat #RL-1102-2
Lipofectamine™ 2000 Transfection Reagent	Invitrogen	Cat #11668019
Recombinant Mouse TNFα	R&D systems	Cat # 410-MT
Recombinant Mouse IL-1β	R&D systems	Cat #401-ML
LysoTracker™ Red	Thermo Fisher Scientific	Cat #L7528
Experimental Models		
C57BL/6 mouse primary brain microvascular endothelial cells	Cell Biologics	Cat #C57-6023
Rab7a$^{flox/flox}$	Dr. Aimee Edinger	RRID: IMSR_JAX:021589
Cdh5(PAC)-Cre^{ERT2} (VEC-PAC)	Gift from Dr. Jan Kitajewski	RRID: IMSR_GPT:T052686
Tg(eGFP-Claudin5)		Published in [10]
Plasmids		
pGEX-4T-3-mR7BD	Addgene	Cat #79149, RRID: Addgene_79149
Oligonucleotides		
Rab7a Stealth siRNA MSS276704, MSS276705, MSS276706	Invitrogen	Cat #1320001

Stealth RNAi™ siRNA Negative Control Med GC Duplex #3	Invitrogen	Cat #12935113
Rab7a flox forward: 5'-CTCACTCACTCCTAAATGG-3'	IDT	N/A
Rab7a flox reverse: 5'-TTAGGCTGTATGTATGTGC-3'	IDT	N/A
Rab7a null forward: 5'-GGGCTGCAGGAATTCGGATAAC-3'	IDT	N/A
Rab7a null reverse: 5'-CATGGTAACAAGTCTGTCGTCC-3'	IDT	N/A
Cre forward: 5'-GCTAAGTGCCTTCTCTACACCTGC-3'	IDT	N/A
Cre reverse: 5'- GGAAAATGCTTCTGTCCGTTTG -3'	IDT	N/A
GFP forward: 5'-CCCTGAAGTTCATCTGCACCAC-3'	IDT	N/A
GFP reverse: 5'-TTCTCGTTGGGGTCTTTGCTC-3'	IDT	N/A
Software and algorithms		
Zen microscopy software	Zeiss	RRID:SCR_013672
Prism 9	GraphPad	RRID: SCR_002798

Empiria Studio	LI-COR	RRID: SCR_022512
ImageJ/Fiji	NIH	RRID: SCR_002285

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to the lead contact,

Dritan Agalliu (da191@cumc.columbia.edu).

Materials availability

This study did not generate new unique reagents.

Data and code availability

All data are available in the manuscript and the supplementary information. The codes used in this

study have been previously published and are freely available (see Key Resources Table). Any

additional information required to reanalyze the data reported in this study will be available upon

request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mice

All experimental procedures were approved by the IACUC committees at the University of

California, Irvine and Columbia University Irving Medical Center. The following mouse strains

were used: *Tg(eGFP-Claudin5)* [10]; *Rab7a^{flox/flox}* [30]; *Cdh5(PAC)-Cre^{ERT2} (VEC-PAC)* [53], kindly

provided by Dr. Jan Kitajewski (Columbia University Irving Medical Center, New York, US). To

induce Cre^{ERT2}-mediated recombination, 4-OH-tamoxifen (Sigma-Aldrich) was dissolved in 10%

ethanol/corn oil mixture to a final concentration of 2 mg/ml, and administered by intraperitoneal

injections at a dose of 35 μg/g body weight for 5 consecutive days starting at postnatal day P1 and

5 as well as 5 consecutive days in the adult ending one week before the t-MCAO procedure.

METHOD DETAILS

Mouse ischemic stroke model

Ischemic stroke was induced in 10-14 weeks old male mice by t-MCAO for 45 minutes as

described [54]. Briefly, mice were anesthetized with 1.5-2% isoflurane, left external and common

carotid artery were ligated and a 7-0 silicon rubber-coated monofilament (Doccol) was inserted

through the common carotid artery into the internal carotid artery. Reperfusion was obtained by

removing the monofilament 45 minutes after its insertion. The body temperature of the mice was

monitored and maintained at 37°C throughout the procedure. The neurological deficits in mice

after t-MCAO were evaluated and scored as described in Jiang S.X. et al., 2005 [41].

Biocytin-TMR and serum IgG permeability analysis

Mice received a tail vein injection of 100 μl of Biocytin-TMR (1% in PBS; ThermoFisher

Scientific). The dye was allowed to circulate for 30 minutes. Animals were anesthetized with

isoflurane and perfused first with PBS, then with 4% PFA in PBS. AlexaFluor 488-conjugated

goat anti-mouse IgG (Invitrogen, 1:500) was used to visualize serum IgG leakage in brain sections.

Immunofluorescence

After dissection, PFA-perfused brains and livers were fixed in 4% PFA at 4°C for 6 hours, then

cryoprotected in 30% sucrose and embedded in Tissue-Tek OCT. For immunofluorescence

staining for Rab7a, Claudin-5 and ZO-1, brains were dissected after perfusion with PBS, fresh-

frozen in Tissue-Tek and sections were fixed with cold 95% ethanol for 30 minutes and acetone

for 1 minute followed by three 5-minute washes with PBS. Brains were sectioned in 12 μm-thick

coronal slices spanning all bregma regions of interest using a Leica cryostat. For primary mBECs

cultures, cells were fixed with cold 95% ethanol for 30 minutes and acetone for 1 minute followed

by three 5-minute washes with PBS. The following primary antibodies were used: rabbit anti-

Rab7a (1:100, Abcam), rabbit anti-ZO-1 (1:500, Invitrogen), rabbit anti-GLUT1 (1:200, Thermo

Scientific), rabbit anti-Claudin5 (1:500, Invitrogen), rabbit anti-VE-cadherin (1:250, Abcam), rat

anti-CD68 (1:500, Abcam), rabbit anti-Iba1 (1:1000, Wako), mouse anti-NeuN (1:500, Millipore).

BSL-rhodamine (1:250, Vector Laboratories) was used to label the vasculature.

Western blotting

Protein levels for endothelial markers were assessed using the Odyssey Sa infrared imaging system

(LI-COR). β-actin was used as housekeeping gene. The following primary antibodies were used:

mouse anti-Claudin5 (1:500, Invitrogen), rabbit anti-Caveolin-1 (1:2000, Abcam), rabbit anti-

Occludin (1:2000, Invitrogen), rabbit anti-VE-cadherin (1:2000, Abcam), mouse anti-ZO-1

(1:2000, Thermo Scientific), rabbit anti-Rab7a (1:1000, Abcam), mouse anti-GST (1:1000, Santa

Cruz), rabbit anti-Ccz1 (1:1000, Abcam), rabbit anti-EEA1 (1:2000, Abcam), rabbit anti-LAMP1

(1:1000, Abcam), mouse anti-β-actin (1:5000, Novus Biologicals), p62 (1:10000, Abcam) and

Atg5 (1:500, Cell Signaling). IR-Dyes 680 and 800 (1:10000, LI-COR) were used as secondary

antibodies. Protein expression levels were quantified using the Empiria Studio software (LI-COR),

normalized on their respective β-actin expression values and presented as a percentage of protein

levels in control conditions.

mBEC culture and Rab7a silencing *in vitro*

Primary mBECs were purchased from Cell Biologics and grown in endothelial cell medium

supplemented with growth factors and 5% FBS (Cell Biologics). Stealth RNAi duplexes targeting

Rab7a were purchased from Invitrogen. Stealth RNAi negative control Med CG Duplex #3

(Invitrogen) was used as negative control in all experiments. 12.5 nM of siRNA was transfected

using Lipofectamine 2000 (Invitrogen), following the manufacturer's instructions.

Rab7a-GTP pull-down assay

The plasmid pGEX-4T-3-mR7BD, which expresses a recombinant protein consisting of the Rab7a

binding domain of the murine RILP protein fused to the C terminus of GST (Addgene, cat#79149),

was transformed into Escherichia coli strain BL21. The transformed bacteria were induced to

express the GST-RILP protein, which was then purified from the bacterial cell lysates using

glutathione-Sepharose 4B beads (GE Healthcare) as described [27]. Control and cytokine/OGD

treated mBECs were lysed in pull-down buffer (20 mM HEPES, 100 mM NaCl, 5 mM $MgCl_2$, 1%

TX-100, and protease inhibitors) and sonicated. Protein levels in the mBEC lysates were quantified

via the BCA assay. Each pull-down was performed by incubating 30 μl of the GST-RILP bound

bead slurry pre-equilibrated in pull-down buffer with 300 μg of cell lysate, and rocking the bead-

lysate mixture in a nutator overnight at 4°C. Afterwards, the beads were washed thrice with cold

pull-down buffer, and bound proteins were eluted by adding sample loading buffer with SDS and

incubating at 95°C for 10 min.

Trans-endothelial electrical resistance

mBECs transfected with either siCTRL or siRab7a were plated on poly-D-lysine and collagen IV-coated gold-electrode array 96-well plates (Applied Biophysics) and grown to confluency. They were then switched to low serum medium (1% FBS) for 24 hours and treated with TNFα and IL1β as described above. The electrical resistance of the cultures was recorded every 20 minutes for the duration of the experiment using an ECIS Z- θ system (Applied Biophysics). GraphPad Prism was used to generate resistance curves and to calculate the area under the curve for each experimental condition, as previously described [55].

In vitro **assays with primary mouse BECs**

For both albumin uptake and Lysotracker assays, mBECs were grown to confluency on poly-D-lysin- and collagen IV-coated glass-bottom 24-well plates (Greiner Bio-One), switched to low serum (1% FBS) medium and treated with TNFα and IL1β as described above. Cells were incubated with either AlexaFluor 594-albumin (100 μg/ml; Thermo Fisher Scientific) or Lysotracker (50 nM; Thermo Fisher Scientific) for 1 hour at 37º C, washed with PBS and fixed in 4% PFA.

QUANTIFICATION AND STATISTICAL ANALYSIS

Image acquisition and quantification

Images of the whole brain sections were acquired with a Zeiss Axioimager fluorescence microscope, higher resolution images were acquired using a Zeiss LSM700 confocal microscope. For comparison purposes, all images of the same staining were acquired under the same settings.

All images were processes using Fiji, and all quantifications were carried out blinded. The areas

of Biocytin-TMR or IgG leakage were quantified by uniformly thresholding the brain sections and

measuring the area exceeding the threshold in each section. The volume of the leakage was

calculated as the product of cross-sectional areas and distance between sections. Intensity of the

leakage was determined by sampling identical regions in contralateral and ipsilateral cortex in each

animal, normalizing their fluorescence intensity on the average fluorescence intensity of the liver

from the same animal and calculating the ratio between the ipsilateral and the contralateral cortex.

To quantify junctional abnormalities, junctional strands, gaps and protrusions were identified as

described before [10] and the percentage of junctional gaps or protrusion over the total number of

junctional strands was calculated. To quantify neuronal viability, we calculated the ratio of NeuN

positive cells over the total of cells for each field of view. Albumin uptake and Lysotracker *in vitro*

were quantified by measuring the area covered by albumin or lysotracker respectively, and

dividing it by the number of cells in the field, as previously described [55].

Statistical analyses

All statistical analyses were performed using GraphPad Prism version 8.1 or higher. Unless

differently specified in the figure legend, data are represented as mean ± s.e.m.. The Shapiro-Wilk

test was used to verify the normality of datasets. For dataset showing normal distribution, pairwise

comparisons were performed using a two-tailed Student's t-test and multiple comparisons were

performed using one-way ANOVA with post-hoc Tukey's correction. P values lower than 0.05

were considered statistically significant (***: $p<0.001$; **: $p<0.01$; *: $p<0.05$).