Modulation of Allostery with Multiple Mechanisms by Hotspot Mutations in TetR

Jiahua Deng[a,1]**, Yuchen Yuan**[a,1]**, and Qiang Cui**[a,b,c,2]

[a]Department of Chemistry, Boston University, Boston, MA 02215; [b]Department of Biomedical Engineering, Boston University, Boston, MA 02215; [c]Department of Physics, Boston University, Boston, MA 02215

This manuscript was compiled on August 29, 2023

For many biomedical and biotechnological applications, modulating allosteric coupling offers unique opportunities. Such efforts can largely benefit from efficient prediction and evaluation of allostery hotspot residues that dictate the degree of co-operativity between distant sites. In several hotspot mutants revealed by recent deep mutational scanning (DMS) experiments of a bacterial transcription factor, tetracycline repressor (TetR), we demonstrate that effects of allostery hotspot mutations can be evaluated by judiciously combining extensive unbiased and enhanced sampling molecular dynamics simulations. The results recapitulate the qualitative effects of these mutations on abolishing the induction function of TetR and provide a semi-quantitative rationale for the different degrees of rescuability to restore allosteric coupling of the hotspot mutations observed in the DMS analysis. Free energy landscapes and variations of structural and energetic properties of different functional states of the mutants show that the hotspot mutations perturb inter-domain coupling and/or intra-domain conformational properties relative to the wild type. Several mutations clearly perturb multiple properties, also highlighting the complexity of realistic situations. Thus, the results support our mechanistic model that hotspot residues contribute to allostery through distinct molecular mechanisms, a feature that we propose to be broadly applicable to many allosteric systems. Overall, our study provides general computational strategies for the identification and evaluation of hotspot mutations modulating allostery in proteins.

protein allostery | molecular dynamics | deep mutational scanning | free energy landscape| enhanced sampling

Allostery couples distant sites by the binding of allosteric effector at the allosteric site and propagation of structural or dynamical perturbations to the functional site to modulate its activity. It is essential to the function of proteins and nucleic acids in many biological processes(1). Controlling the allosteric coupling and communications is of broad therapeutic value and offers unique opportunities(2) for biomedical and biotechnological applications. Designing molecules that target an allosteric site rather than the active site can potentially lead to higher levels of selectivity and specificity, especially for drug targets that feature highly conserved binding pockets such as kinases(3, 4). Numerous serious human diseases such as various cancers are caused by mutations that either disrupt or activate allosteric communications(5), and resistance to drugs also often arises due to mutations that perturb allostery in the target protein(6). The understanding of allosteric communications also makes it feasible to design and engineer target proteins with desired purposes by mutations, such as controlling enzyme activities for catalysis, constructing regulatory systems for gene expression, and developing biosensors and biological logic gates(7–13).

To effectively design allosteric therapeutics and engineer/re-engineer allosteric proteins, it is useful to first identify and target allostery hotspots. Hotspots are the set of key residues that dictate the co-operativity between the effector site and the active site. In addition to their locations, the specific sequence features of allostery hotspot residues may also be important to their functional significances, since their contributions are likely sensitive to the local steric or polar interactions they are engaged in. Allostery hotspots can be explored by various computational techniques, such as statistical analysis of residual co-evolution(14–16), motional network and community analysis(17–20), mapping propagation of conformational distortions(21), and perturbation analysis of collective normal modes(22–24). Experimentally, site-directed mutagenesis(25), solution NMR chemical shift perturbations and relaxation measurements(26, 27), surface plasmon resonance(28), and *in vitro* and *in vivo* biochemical essays/kinetics(29) are commonly used approaches to probe allostery hotspots. Since each methodology has its own limitation(s) in either cost or accuracy, it is still challenging to identify a complete set of allosteric hotspots, which prevents the evaluation of the individual role of each hotspot and a clear understanding of their mechanisms. A high-throughput methodology which could exhaustively and systematically analyze the sequence-function relationship is of great promise. Deep mutational scanning (DMS) analysis(30–32) has emerged as a powerful technique for functional screening of allostery hotspots in proteins in

---

Significance Statement

To understand the mutation effects on allosteric coupling, we demonstrate that the same non-inducible phenotype could be the result of perturbations in distinct structural and energetic properties in a bacterial transcription repressor. Reduced anti-cooperativity between the distal binding sites is found in the mutants to reflect the loss of function. Multiple mutations also alter the conformational ensemble of the ligand binding site for unfavorable binding, suggesting that hotspot mutations can be used to fine-tune allostery properties including effector specificity. For effectively evaluating and ranking hotspot mutations despite the prevalence of compensatory interactions in allostery systems, our work underscored the value of explicitly computing the functional free energy landscapes to provide quantitative guidance to allostery modulation for therapeutic and engineering applications.

J.D., Y. Y. and Q.C. designed research; J.D. and Y. Y. performed research; J.D., Y. Y. and Q.C. analyzed data; and J.D., Y. Y. and Q.C. wrote the paper.

The authors have no competing interest.

[1]J.D. and Y. Y. contributed equally to this work.

[2]To whom correspondence should be addressed. E-mail: qiangcui@bu.edu

a comprehensive and unbiased fashion. As demonstrated by recent applications(33–38), DMS analysis readily reveals both the location and sequence variability of allostery hotspots, leading to new insights into the unique characteristics of allostery network in various systems. However, DMS studies are only able to identify phenotypes through the binary functional readout due to its high-throughput nature. Establishing the effect of a given mutation on protein function and other quantitative properties requires additional molecular details.

Another feature that is commonly observed in allosteric systems is the network of compensatory interactions(39), which leads to near-degenerate allosteric communication pathways(40). This feature makes allostery highly adaptable during evolution and ensures robustness of functionally essential co-operativities(34); it can also facilitate re-wiring of allostery communication pathways in engineering applications(12, 41). Nevertheless, these compensatory effects have major implications to the design of allosteric inhibitors, as the effect of design might be easily abolished by additional mutations(4, 6). Therefore, robust designs should target allostery communications that, once disrupted, are difficult to restore. Such properties are costly to evaluate using experiments due to the need of sampling a large number of sequences. Providing insights into how easily perturbations due to allostery hotspot mutations are reversed using computational approaches(42) is of tremendous value but has not been widely attempted.

To comparatively evaluate the effect of hotspot mutations, it is important to establish the mechanism through which mutations of hotspot residues perturb allostery. This informs the relevant quantitative computations that can be used to compare the impacts of different mutations. Moreover, a mechanistic understanding will greatly facilitate the development of specific strategies for either enhancing or reversing the effect of mutation by either introducing additional mutations or interactions with other molecules(43). For example, allosteric communication can be tuned by modulating the stability of altered loop conformation(44, 45), targeting new surface binding pockets via small molecules (16, 43, 46, 47), or mutations of key residues along allostery pathways(12, 48).

A fitting example in this context is a bacterial transcription repressor, tetracycline repressor (TetR). TetR's normal function relies on the allosteric coupling between the ligand binding domains (LBDs) and the DNA binding domains (DBDs), which are separated by more than 20 Å (Fig. 1a). Binding of the inducer (ligand-Mg^{2+} complex) leads to the dissociation of the repressor from the promoter DNA sequence, allowing the access of RNA polymerase and therefore transcription of the repressed gene. Firmly establishing the contributions of hotspot residues has fundamental value to many aspects of prokaryotic physiology, including antibiotic resistance, and provides guidance to rational engineering of the TetR family for gene regulation(49). Therefore, allostery in TetR has been extensively studied with structural(50–53), mutagenesis(54–56) and more recently, DMS studies(34, 35). Based on propagation of structural distortions, a set of allostery hotspots abolishing the induction function in TetR were proposed(57) and verified(58, 59) to localize at the domain interface of the two binding sites and relay the effect of inducer binding to the DBDs, whose reorientation impacts the DNA binding affinity. By sharp contrast, DMS analysis observed that allostery

hotspots in the TetR family members are broadly distributed over a major portion of the protein structures(34, 35), underscoring that a mechanical view of allostery(21, 47, 60, 61), especially in TetR is incomplete. Moreover, the DMS experiments found that the loss of induction in hotspot mutants could be restored by additional mutations(34). These observations again highlighted that the allostery network has a considerable degree of degeneracy and redundancy, and not all allostery hotspots are of equal weights during evolution.

These considerations motivated our recent atomistic molecular dynamics (MD) simulations of the wild type (WT) TetR(62). Multiple local (e.g., backbone dihedral and residual contacts) and global (e.g., motional coupling and free energy landscapes) analyses established the molecular and energetic basis of the strong anti-cooperativity between the ligand and DNA binding sites through a conformational selection mechanism that underlies the allostery function of TetR. The multi-faceted computational framework we established have captured a major fraction of the hotspot residues identified in the DMS study and demonstrated that these residues contribute to allostery in distinct ways. Nevertheless, our previous simulations focused entirely on the WT protein, thus the roles and features of hotspot residues had to be inferred indirectly. Moreover, the general strategy for hotspot identification also generated a considerable number of "false positives" and missed hotspot residues in two regions, the middle of the $\alpha 8$ helix and the C-terminus of the protein(62). Evidently, the molecular mechanism for the disruption of allostery in the hotspot mutants, the non-inducible phenotype, remains unclear.

In the current study, we conduct extensive unbiased (6-60 μs) and enhanced sampling molecular dynamics (MD) simulations for several representative hotspot mutants to better define the mechanism of allostery modulation by the corresponding mutations. We selected mutants that involve hotspot residues in distinct regions of the protein (see Fig. 1b): domain interface (G102D, R49A), near the ligand binding site (L127D, G138D/H and L145A), and the C-terminal helix (I193A and G196D). Several residues (L145A, I193A and G196D) are also chosen in the regions missed entirely in our previous computational analyses in the WT to explicitly probe their mutation effects.

Results

Doubly-Bound States Are Better Stabilized in Hotspot Mutants.
In the previous structural studies of TetR(50, 52), the center-of-mass distance separation between the DBDs (DBD distance) was considered to be a key indicator for the DNA binding affinity. In the WT, ligand binding was proposed to rigidify the protein structure such that the DBD distance was too long to allow favorable binding to the DNA according to the induction function of TetR. Thus, we reason that the loss of the induction function in some, if not all, hotspot mutants is due to adopting favorable conformations in the DBDs for DNA binding even in the ligand-bound state; i.e., these mutants are able to bind tightly to both the ligand and DNA in such conformations, referred to as the doubly-bound state. To test this hypothesis, we conduct metadynamics simulations to explicitly probe how hotspot mutations modulate the free energy profile associated with the DBD distance in the ligand-bound state. The use of the DBD distance as the reaction coordi-

(a)　　　　　　　　(b)

Fig. 1. Overview of the TetR structure and experimentally-identified allostery hotspot distributions. (a) DMS studies(34) revealed a broad distribution of allostery hotspots (in red) throughout the majority of the TetR structure. In the ligand-bound crystal structure (PDB code: 4ac0), the ligand, minocycline, is bound to a Mg^{2+} ion. The ligand is converted to anhydrotetracycline in all simulations. (b) The locations of hotspot mutations studied in this work are grouped into three regions: the domain interface, near the ligand binding site, and the C-terminal helix α10.

nate is supported by previous MD simulations on the order of
50-100 ns(63), which observed that the DBD distance can be
used to distinguish induced (ligand-bound) and non-induced
(DNA-bound) conformations. We also note that the DBDs are
perturbed by the hotspot mutations in both the DNA-bound
(Fig. S1) and apo (Fig. S2) states in unbiased simulations. As
shown in Fig. S2, the DNA-bound state of the mutants tends
to span broader DBD distances compared to the WT, leading
to Kullback-Leibler (KL) divergence values in the range of
0.8-1.8 for the corresponding DBD distance distributions (Ta-
ble S2). These values reflect modest differences in the DBD
conformations between the mutants and WT. Regarding the
DBDs in the apo state, their conformations in the hotspot
mutants are overall similar to the WT, except that the DBD
separations are slightly shifted towards longer distances (Fig.
S2), most notably in G138D and G138H.

In the WT, the computed free energy profile (Fig. 2)
supports the hypothesis from the structural studies(50, 52,
55). There is a single free energy basin at $\sim$ 46 Å along
the DBD distance coordinates. The basin (ligand-only basin)
corresponds to the crystal structure of the ligand-bound TetR,
in which the DBD distance is $\sim$ 42 Å(50). As the DBD distance
decreases, the free energy increases monotonically. For the
DBD distance reaching $\sim$38 Å, the corresponding conformation
is expected for favorable DNA binding(52), which has a free
energy cost about 10 k_BT. These observations are consistent
with the normal function of TetR. The low DNA binding
affinity upon ligand binding suggests that compressing the
DBDs in the ligand-bound state is energetically unfavorable.
In all the hotspot mutants studied, by remarkable contrast,
the computed free energy profiles are double-well in nature
(Fig. 2a-c and Fig. S3a). The ligand-only basin is consistently
observed at the large DBD distance of $\sim$46 Å, similar to the
WT. In addition, a second free energy basin is observed at
a shorter DBD distance expected for favorable DNA binding
in the ligand-bound state. The location of this inner basin
is largely consistent ($\sim$38 Å) among all the various mutants
studied as well.

To further verify the nature of the second basin in the
PMFs along the DBD distance in the mutants, we measure
the KL divergence of the DBD distance distributions between

Table 1. Overlap of DBD distance distributions between the ligand-bound and DNA-bound states characterized by the Kullback-Leibler (KL) divergence [a]

	WT	G102D	R49A	G138H
unbiased	5.2	2.9	2.1	0.7
reweighted	6.8	1.2	3.0	1.7
doubly-bound basin[b]	6.0	0.4	0.5	0.1
ligand-only basin[b]	54.4	56.7	57.3	56.7
	L127D	G138D	I193A	G196D
unbiased	3.8	2.9	3.1	3.0
reweighted	2.8	3.8	2.0	3.3
doubly-bound basin[b]	0.6	0.4	0.8	1.0
ligand-only basin[b]	57.5	56.7	56.1	57.2

a. In the KL divergence calculations, the reference is taken to be the distribution from the ligand-bound simulation. DNA-bound state simulations are all unbiased simulations (see Table S1). For the ligand-bound states, results from both unbiased simulations (see Table S1) and reweighted distributions from metadynamics trajectories are used. The distributions are shown in Fig. S4-S5.

b. Reweighted distributions from metadynamics simulations are split into "doubly-bound basin" and "ligand-only basin" components with the DBD distance of 42.5 Å as the boundary.

the ligand-bound and DNA-bound state simulations (see Figs. 211
S4-S5). As shown in Table 1, the DBD distance distributions 212
of the inner basin in the ligand-bound state of the hotspot 213
mutants overlap well with the corresponding DNA-bound state 214
with KL divergence no larger than 1.0. In comparison, the WT 215
simulations lead to much larger KL divergence values of the 216
DBD distance distributions between the ligand-bound state 217
and DNA-bound-state. A smaller KL divergence value indi- 218
cates a larger similarity between the two distributions under 219
comparison, suggesting that the DBDs in the ligand-bound 220
states of the mutants are more favorable for DNA binding 221
than the DBDs in the ligand-bound WT. Another structural 222
variable thought to be important to DNA binding (affinity) 223
is the relative angle of the α4 helices at the domain interface 224
connecting the DBDs and LBDs(50, 52) (see definitions of key 225
DBD order parameters in SI). KL divergence of the α4 angle 226
distributions in the two bound states is also substantially lower 227
in the hotspot mutants than in the WT (Table S3 and Fig. S6- 228
S7), also showing that the ligand-bound states of the mutants 229
have higher populations in the DNA-binding-competent con- 230
formation than the ligand-bound state of the WT. Collectively, 231
these results quantitatively demonstrate that the free energy 232
basin at the short DBD-distance sampled in the metadynamics 233
is indeed favorable for DNA binding in the ligand-bound state. 234
Therefore, although only the ligand is explicitly included in 235
the metadynamics simulations, we refer to this basin as the 236
"doubly-bound basin" since the corresponding conformational 237
ensemble is expected to bind tightly to both the ligand and 238
DNA. 239

Among the hotspot mutants studied, the doubly-bound 240
basin is higher in free energy than the ligand-only basin at 241
the longer DBD distance by 2-7 k_BT. The only exception 242
is G102D, in which the two basins are similar in stability 243
(Fig. 2a), suggesting that G102D is highly poised to bind 244
both the ligand and the DNA. While the R49A mutation 245
is also close to the domain interface and involves a similar 246

Fig. 2. Potentials of mean force (PMF) for the DBD distance separation in the ligand-bound states from metadynamics simulations. The reaction coordinate is the center of mass separation between the two DBDs in each monomer. See the precise definition in Materials and Methods. All PMFs show a basin at ∼ 46 Å, referred to as the "ligand-only basin". All mutant PMFs also show another basin at ∼ 38 Å, referred to as the "doubly-bound basin". Subfigures are grouped by the regions shown in Fig. 1b. The solid lines are the average values and the shaded areas show the statistical error estimated from block analysis. The statistical errors are all on the order of 0.1 k_BT. For additional analysis of statistical error and convergence, see Fig. S3b in the SI.

magnitude change in net charge, the mutation effect on the DBD energetics is notably smaller. Interestingly, G138D and G138H exhibit rather different stabilities of the doubly-bound basins (Fig. 2b) despite implicating mutation of the same residue. These variations underscore the importance of the unique local interactions associated with specific mutations. For example, in G102D, the introduced aspartate D102 is engaged in new interactions (compared to the WT) with several cationic residues nearby, such as K155' and R158' in α8', and R49 in α4 (Fig. S8). By contrast, in R49A, the interaction between R49 and E157' in α8' in the WT(62) is lost. In G138D, the introduced aspartate D138 is engaged in hydrogen-bonding interactions with the hydroxyl group in the ligand and therefore has rather static orientations. While in G138H, the lack of such hydrogen-bonding interactions leaves the introduced histidine H138 flexible, with occasional flip of the sidechain to push on α5 (Fig. S8). Finally, it is worth noting that while previous WT simulations did not identify any distinct properties of the hotspot residues in the middle of α8 and C-terminal region (α10)(62), the current metadynamics simulations of the α8 mutant (V145A) and C-terminal mutants (I193A and G196D) have clearly captured the effect of the mutations on the DBDs (Fig. S3a and Fig. 2c). In these three mutants, the doubly-bound basin is distinctly visible and is only higher in free energy by ∼2-4 k_BT than the ligand-only basin.

Inter-Domain Coupling Is Reduced in Hotspot Mutants. The normal induction function of the WT TetR relies on the strong anti-cooperativity between the LBDs and DBDs. The double-well feature of the computed DBD free energy profiles of the hotspot mutants in Fig. 2 suggests that the magnitude of inter-domain coupling is substantially reduced compared to the WT. As a result, these mutants are more inclined to bind tightly to DNA even in the ligand-bound state. Importantly, the reduction of inter-domain coupling is also evident from several additional lines of analyses, further supporting the robustness of the conclusion.

First, we note that, in the ligand-bound state simulations, the ligand binding pocket in the hotspot mutants exhibits notable changes depending on the DBD distance. At a longer DBD distance in the ligand-only basin, the l6-α8 distance (the Cα distance between R104 and L134) that characterizes the compactness of the binding pocket exhibits overall similar

distributions as the WT (Fig. 3a-c), with merely shifted peak positions or modified distribution widths. In the doubly-bound basin, which features a compressed DBD distance, much larger changes in the ligand binding pocket distributions relative to the WT are observed for several mutants (Fig. 3d-f). For example, both G102D and G138H exhibit much broader distributions. The peak position is also notably shifted in L127D, the two G138 mutants G138H/D, as well as G196D (also see Fig. S9 and Table S4). These observations suggest that the coupling between the binding pocket and the DBDs is not altered much in the ligand-only basin in the hotspot mutants, but the coupling is more significantly modified in the doubly-bound basin of the mutants. In particular, the substantially expanded pair-wise distance distributions in G102D and G138H in the doubly-bound basin suggest that the coupling between LBDs and DBDs is much reduced relative to the WT. The reduced inter-domain couplings at compressed DBD distance are consistent with the higher stability of the doubly-bound basin in these hotspot mutants in comparison to the WT (Fig. 2).

The variation of inter-domain coupling among the hotspot mutants relative to the WT is also directly visualized in the free energy landscapes of the ligand-bound states projected onto the two-dimensional space spanned by the DBD distance and the l6-α8 distance (Fig. 4). In the WT, free energy landscapes of the two bound states are distinct and exhibit limited overlap along either collective variable, reflecting the strong anti-cooperativity between the LBDs and DBDs. In the free energy landscapes of the mutants, especially in the ligand-bound simulations, the DBD distance distributions are much broader than the WT. The doubly-bound basin in the two-dimensional mutant free energy landscapes is well distinguished from the ligand-only basin along the DBD distance coordinate but significantly overlaps with the ligand-only basin along the l6-α8 distance coordinate. The above observations reflect weaker inter-domain anti-cooperativity. In particular, for G102D and G138H, the binding pocket distributions are much broader with compressed DBD distance, suggesting even weaker inter-domain coupling in the doubly-bound basin in these mutants.

Finally, we examine the free energy landscapes of the apo states projected onto the two-dimensional space spanned by

Fig. 3. Binding pocket distance distributions in different conformational states of the ligand-bound states from metadynamics simulations. Subfigures are grouped by the regions shown in Fig. 1b. The reweighted distributions of the Cα distance between R104 and L134 (l6-α8 distance. l6 stands for the linker region of α6-α7.) (a-c) in the "ligand-only basin" (DBD distance range is between 46.5 Å and 47 Å), and (d-f) in the "doubly-bound basin" (DBD distance range is between 37.5 Å and 38 Å) from metadynamics simulations.

Fig. 4. Two-dimensional free energy landscapes of the DNA-bound and ligand-bound states: (a)WT, (b) G102D, (c) R49A, (d) G138H, (e) L127D, (f) G138D, (g) I193A, (h) G196D. The horizontal axis is the center-of-mass distance between the DBDs, and the vertical axis is the Cα distance between R104 and L134 (l6-α8 distance), which characterizes the compactness of the ligand binding pocket. For the DNA-bound states, the results are based on unbiased MD simulations (see Table S1 for simulation length summary). For the ligand-bound states, results based on unbiased MD simulations and reweighted distribution from metadynamics simulations are shown for comparison. While the unbiased and metadynamics simulations sample mostly similar conformational ensembles, due to the free energy barrier along the DBD distance (Fig. 2), the metadynamics simulations better capture the underlying free energy landscape and structural properties, especially for the "doubly-bound basin", which is hardly sampled in the unbiased simulations at the microsecond time scale. Color scheme: rainbow - ligand-bound state metadynamics simulations; gray - unbiased ligand-bound state simulations; copper - DNA-bound simulations. Color bars are in the unit of $k_B T$.

(a)   (b)   (c)

(d)   (e)   (f)

Fig. 5. Free energy landscapes and ligand binding pocket distance distributions of the apo states. Two-dimensional free energy landscapes of the (a) WT, (b) G102D and (c) R49A variants. Results of the remaining mutants are qualitatively similar to the WT and shown in Fig. S10. The color bar is in the unit of $k_B T$. The horizontal axis is the first principal component (PC1), and the vertical axis is the Cα distance between Q109 and E147' (α7-α8' distance). PC1 describes the complex motions of the DBDs, while the inter-helical distance characterizes the compactness of the ligand binding pocket. The WT and mutants generally feature diagonal shape landscapes, while an extra basin is also observed in the mutants (G102D and R49A). Two-dimensional free energy landscapes projected onto different order parameters are shown in Fig. S11, which also exhibit similar changes for the two domain-interface mutants (G102D and R49A). (d-f)Distributions of the α7-α8' distance in different mutants. The results are based on extensive unbiased samplings (30-60 μs) for each variant. Subfigures are grouped by the regions shown in Fig.1b.

the collective DBD motion (represented by the first principle component PC1 used in our previous work(62)), and $\alpha 7 - \alpha 8'$ (the Cα distance between Q109 and E147') distance of the ligand binding pocket (Fig. 5a-c). Projection onto other co-ordinates leads to generally similar observations (Fig. S11). Analysis of the apo landscape is complementary to the bound-state simulations discussed so far, as the shapes of the apo landscapes reflect the intrinsic features of inter-domain cou-pling in the absence of any local stabilization by the ligand or the DNA. The two-dimensional free energy landscape of the WT shown in Fig. 5a is much broader than those computed for the bound states(62), pointing to a conformational selection mechanism for TetR function. Moreover, the diagonal shape of the landscape is consistent with the strong anti-cooperativity expected for the LBDs and DBDs. For the studied hotspot mutants, the apo free energy landscapes are similarly broad, indicating that the conformational selection mechanism is still applicable. The diagonal shapes of the landscapes remain qualitatively similar to the WT for most mutants (Fig. S10), suggesting that the anti-cooperativity between the LBDs and DBDs is not significantly perturbed in the apo state of these mutants. Exceptions are the two mutations at the domain interface, G102D and R49A, of which the free energy land-scapes deviate significantly from the diagonal shape with an additional basin that features a positive PC1 value and short $\alpha 7 - \alpha 8'$ distance (compare Figs. 5b,c and 5a). As explained in the SI using a revised thermodynamic model (Fig. S12) for al-lostery in TetR(62), such changes in the free energy landscape are strong signatures for the decrease of inter-domain cou-pling. Moreover, the most evident changes in the free energy landscapes associated with decreased inter-domain coupling in G102D and R49A are also observed when the free energy

landscapes are projected onto other two-dimensional spaces spanned by the DBD distance and the $\alpha 7 - \alpha 8'$ distance (Fig. S11).

Binding Pocket Is Perturbed to Decrease Ligand Binding in Hotspot Mutants. The free energy results (Figs. 2) and analysis of inter-domain coupling suggest that the studied hotspot mutants have much stronger propensities to bind both the ligand and DNA, which is consistent with their non-inducible phenotype observed in the DMS experiment(34). However, a higher population of the doubly-bound state is not the only potential cause for the loss of induction function. Reducing TetR's apparent ligand binding affinity to be lower than the binding affinity for the DNA promoter sequence, or enhancing the DNA binding affinity would also lead to the non-inducible phenotype (see Fig. S12). In general, the apparent ligand or DNA binding affinity can be modified by hotspot mutations through perturbations in either the ligand/DNA-bound state or the apo state, or both. Therefore, we evaluate the impact of the hotspot mutations on the properties of both bound states and the apo state.

For the DNA-bound states, the ligand binding pocket re-mains loosely packed in both WT and the hotspot mutants (Fig. S13-15). For the ligand-bound states, at long DBD distances (in the ligand-only basin), the ligand binding pocket remains tightly packed and also experiences limited pertur-bation by mutations (Fig. 3a-c). With compressed DBD distances (in the doubly-bound basin), however, considerably broader binding pocket distributions are observed in several mutants (Fig. 3d-f), especially in G102D and G138H. There-fore, upon DNA binding, the binding affinity to the ligand is expected to be further weakened in these mutants as compared

to the WT, which contributes to the diminished induction activities.

In the apo states, many hotspot mutations impact the conformational ensembles of the ligand binding pocket. As shown in Fig. 5d, the WT ligand binding pocket favors the binding-competent conformation (long $\alpha7$-$\alpha8'$ distances) in the apo state. In the two mutants at the domain interface (G102D and R49A), by contrast, the ligand binding pocket is dominated by the binding-incompetent conformation (short $\alpha7$-$\alpha8'$ distances) that is favored in the DNA-bound state (see Fig. S13). Therefore, a lower apparent ligand binding affinity is expected for these mutants relative to the WT. In the other mutants, the magnitudes of the perturbation are smaller. In some cases, such as L127D (Fig. 5e), the $\alpha7$-$\alpha8'$ distance distribution becomes unimodal and the peak position shifts towards a short distance that is favored in the DNA-bound state (also see Fig. S13), suggesting considerable conformational rearrangements in the binding pocket and reduction in the ligand binding affinity. In the case of the C-terminal mutants (I193A and G196D), the general shape of the distribution remains similar to the WT, although the peak positions of the distributions are shifted (Fig. 5f), also reflecting notable perturbation of the ligand binding pocket.

Discussion

The Mechanism of Allostery Modulation by Hotspot Mutations in TetR. In our recent study(62), motivated by the distinct contributions from allostery hotspots, we emphasized the importance of analyzing multiple properties at both the local and global scales, which was demonstrated to be essential to capturing multiple hotspots. However, since only the WT protein was investigated and contributions of residues to allostery can only be inferred indirectly, those analyses generated considerably numbers of "false positives", while missing residues in certain regions, such as the C-terminal helix and the middle of $\alpha8$. By conducting extensive unbiased and enhanced simulations of multiple functional states of the hotspot mutants and characterizing their structural and energetic properties, we have gained new insights into the nature of allostery in TetR and the mechanism by which the induction function is modulated relative to the WT protein. Encouragingly, all of the mutant simulations have clearly captured the effects of the hotspot mutations, including those not identified in previous WT simulations(62). The PMF results for the DBD separation in the ligand-bound states (Fig. 2) clearly demonstrate that hotspot mutation effects can propagate from many regions in the protein to the DBDs, including the C-terminus, which is at the opposite end of the protein structure to the DBDs. The successful recapitulation of the hotspot mutation effects in this work underscores the value of conducting explicit mutant simulations for the evaluation of hotspot candidates, since mutant simulations are able to capture the cascade of changes in structure and energetics not evident in the WT. Additionally, compensation effects of nearby residues are included, and therefore "false positives" can be minimized.

This observation highlights the global nature of energetic coupling in proteins(64), and is consistent with the distributed nature of allosteric network as observed in the DMS study(34) and our previous MD analysis(62). Perturbations by these hotspot mutations are eventually manifested by the functional change of the DNA-bound state population, as the consequence of certain structural changes in the LBDs and/or DBDs. Our results explicitly illustrate that the loss of induction function in the hotspot mutants is due to either reduced inter-domain coupling, which leads to stabilization of the doubly-bound state, or perturbation of the ligand binding pocket conformation, which reduces apparent ligand binding affinity. Therefore, the current observations generally support the thermodynamic model proposed in our previous work(62). It classifies allostery hotspots into two distinct types that dictate inter-domain coupling and intra-domain energetics, respectively. The current analysis of the mutants at the atomic level highlights that a single mutation may lead to notable perturbation in both types of properties; for example, G102D clearly impacts both inter-domain coupling and conformational distributions of the ligand-binding pocket. Therefore, the type 1 & 2 hotspots discussed in the context of the thermodynamic model(62) are *idealized* classifications and the realistic situations tend to mix both effects. Such general features are broadly applicable to many allosteric systems, especially those sharing the two-domain structural topology.

The induction function of TetR relies critically on the anti-cooperativity between the ligand and DNA binding domains. In the hotspot mutants, we observe various signatures based on energetic and structural properties that consistently point to the reduction of inter-domain coupling in comparison to the WT protein. Our analysis of multiple functional states or free energy basins further reveals that the inter-domain coupling is not simply a constant value but dependent on the particular conformational and binding state of the protein. For example, in G102D and G138H, the coupling is observed to further decrease upon compression of the DBDs in the ligand-bound state. Moreover, significant decrease of inter-domain coupling in the apo state is most evident only in the two domain-interface mutants (G102D and R49A), while all studied mutants exhibit reduced anti-cooperativity in the ligand-bound states as reflected by the stabilization of the doubly-bound basin.

The observation that the ligand binding pocket can be perturbed by hotspot mutations suggests that ligand binding properties such as affinity and specificity can be tuned by residues beyond the ligand-contacting residues. In other words, mutation of hotspot residues can perturb beyond the coupling between distant sites, a notion that has implications in applications. For example, this realization offers insights into how distal mutations might affect inhibitor potency in the context of drug resistance development(6). Moreover, it also highlights additional flexibility in the strategies that can be used to fine-tune the various properties relevant to the allostery function, including effector specificity(41).

The essence of our dissection of allostery modulation in TetR concerns the analysis of inter-domain coupling and intra-domain properties that affect the apparent binding affinity of effector (inducer) on an equal footing. Since the proper function of most allostery systems relies on co-operativity between distinct domains and tight binding of the effector, the strategies and insights we have established through the analysis of TetR have major implications to the general prediction and evaluation of allostery hotspots in proteins.

Rescuability of Allostery Hotspots. In the DMS study of TetR(34), another remarkable observation was that the loss of induction function in hotspot mutants could be rescued by

additional mutation(s), also in a distributed and degenerate fashion. At the qualitative level, since the induction function can be disrupted in multiple ways by perturbing either inter-domain coupling or intra-domain energetics, we expect that restoration of the normal function can also be accomplished in degenerate fashions through reversing changes in these properties. At the quantitative level, the degree of rescuability was defined as the ability of restoring the TetR function on non-inducible (dead) mutants. The DMS experiments found that among the five hotspots analyzed experimentally (R49A, D53V, G102D, N129D and G196D), the degree of rescuability follows the order of N129D>G196D>D53V>R49A>G102D(34). As discussed in our earlier work(62), successful design of allostery inhibition(2, 4) needs to target allostery hotspots that are particularly difficult to rescue, since otherwise allostery can be readily restored with additional mutations, leading to abolishment of inhibition. Therefore, being able to evaluate the rescuability of allostery hotspots using computation is of great value.

In this context, a closely related quantity is the magnitude of perturbation on the free energy landscape. Specifically, it is interesting to compare the impact of hotspot mutations on the DBD distance free energy profile in the ligand-bound state with their degree of rescuability. As shown by the PMF calculations (Fig. 2), the impact of mutation on DBD energetics is the largest for the G102D mutant, in which the doubly-bound basin is similar in stability to the ligand-only basin; G102D was also observed to be most difficult to rescue in the DMS analysis. Although R49A is also a mutation at the domain interface and features an identical change in the net charge, the magnitude of the effect on DBD energetics is notably smaller, which is consistent with the higher degree of rescuability of R49A observed in the DMS experiment(34). Moreover, the relative stability of the doubly-bound basin and the ligand-only basin is comparable in G196D and R49A. This is qualitatively consistent with the similar degrees of rescuability of these two mutants observed in the DMS experiment. Therefore, these observations support that mutations that lead to a larger change in the DBD free energy landscapes are more difficult to rescue.

The current work demonstrates that functionally relevant PMF simulations are more effective than unbiased simulations in terms of capturing and qualitatively ranking different hotspot residues and related mutation effects. This is due in part to the fact that conformational sampling remains a general challenge, especially for the bound states, which tend to be structurally more rigid and therefore feature higher free energy barriers for conformational rearrangements. Thus, free energy simulations along the relevant coordinate(s) play critical roles for the evaluation of allostery modulation. This further highlights the importance of mechanistic analysis of allostery, which is required for the identification of proper coordinate(s) for functional free energy landscape computations. To aide the latter task, it is essential to continue the development of creative computational methodologies, such as those based on machine learning analysis of conformational ensembles associated with distinct functional states(65–68).

The general consistency between computational and experimental results regarding mutant rescuability suggests that it is feasible and meaningful to use functional free energy simulations to rank hotspot mutations for evaluating rescuability,

which is a property of interest in the context of realistic applications, such as the design of allosteric inhibitors(2, 4). Indeed, the considerable level of redundancy for the allostery network identified in DMS studies raised the questions of whether engineering allosteric modulators can be a successful path. The current work helps compare the rescuability of hotspot mutations and supports that engineering designs to modulate allostery could be tested through MD simulations. Our computational strategies take advantage of multiple, modest-length enhanced sampling simulations and therefore are scalable to other allostery systems.

Supporting Information Appendix (SI). Additional simulation details, results and schematics for a revised thermodynamic model are included in the SI.

Materials and Methods

Unbiased MD Simulations of Different Functional States. Extensive unbiased MD simulations (6-60 μs) are conducted to compare hotspot mutants and WT TetR in terms of key structural properties and construct free energy landscapes in multiple functional (apo, ligand-bound and DNA-bound) states on local NVIDIA P100 GPUs and the Anton 2 machine. WT trajectories were generated in the previous study(62). The mutants studied in this work include G102D, R49A, L127D, G138D, G138H, V145A, I193A, and G196D. All mutations are generated *in silico* based on the corresponding coordinates of the WT. Unbiased MD setup for the mutants, the definitions of various order parameters and structural properties are the same as our previous work on the WT (62). Simulation details and data analysis for the unbiased trajectories are presented in the SI.

For the ligand-bound state, we start with the mutant ligand-bound structures based on the crystal structure of the WT (PDB code: 4AC0). For the DNA-bound state, the initial structure is modeled based on the crystal structure of the TetR(D)-DNA complex (PDB code: 1QPI), in which the protein has a high level of sequence homology to TetR(B) (66%). 1 μs and 5 μs trajectories are generated for the bound states on local GPUs and Anton 2, respectively.

For the apo state, three independent models are used as the initial structures to reduce bias and facilitate the exploration of the conformational space. The first structure is the same as the ligand-bound crystal structure but without the ligand-Mg^{2+} complex; the second structure is the same as the DNA-bound structure but without the DNA; the third structure is modeled using the TetR(D) apo crystal structure (PDB code: 1BJZ) as the template. Independent simulations of apo states are carried out both on local GPUs and Anton 2 starting from different initial structures for each system. For apo state simulations on local GPUs, 3 independent trajectories of 1 μs or 3 μs are carried out starting from the aforementioned three initial structures, respectively. For the apo state simulations on Anton 2, 3 independent trajectories of 15 μs, 15 μs, and 20 μs are carried out for the WT starting from the three initial structures, respectively; 2 independent trajectories of 15 μs each are carried out for G102D starting from the first two initial structures of the apo state, respectively; 3 independent trajectories of 10 μs each are carried out for the rest mutants starting from the three initial structures, respectively.

Metadynamics Simulations of Ligand-bound States. Multiple-walker well-tempered metadynamics simulations(69, 70) are carried out using the PLUMED(71, 72)-OpenMM interface on NVIDIA A100 GPUs. The collective variable (CV), DNA-binding domain (DBD) distance, is the center-of-mass (COM) separation between the Cα of residues 37-44 in each monomer. The bias factor is set to be 10. The simulation temperature is 303.15 K. A new Gaussian biasing potential is added every 1 ps with an initial height of 1.2 kJ/mol and a width of 0.2 nm. Ten walkers with different initial velocities are used per system in parallel while sharing hill history with other walkers every 1 ps. A harmonic restraint is applied to the CV with

a force constant of 150 kJ/mol when the value of the CV is larger than 5 nm (upper wall) or smaller than 3 nm (lower wall). Other basic MD parameters are the same as the unbiased simulations. Convergence is monitored by the block-average analysis of which the details are presented in the next subsection. The simulation is stopped when adequate convergence has reached, which typically takes 110 - 140 ns per walker in each independent metadynamics run.

Data Analysis for Metadynamics Simulations. Metadynamics trajectories are analyzed using the metadynamics reweighting algorithm(73) implemented in the PLUMED package (version 2.8)(72) to obtain the unbiased distribution of the DBD distance, which is then converted into the 1D PMF. Block-average analysis is applied to assess convergence of metadynamics simulations and the associated statistical errors for each system. The average errors of the block-averaged PMF as a function of the block size are plotted as the convergence check. The final 1D PMF of each system is the block average of every 20 ns trajectories in each walker, and the shaded area in the PMF plot represents standard error of the mean (SEM) of the block-averaged PMF. The reweighted distributions of other order parameters, structural properties, as well as the 2D free energy landscapes are obtained using the same reweighting algorithm.

ACKNOWLEDGMENTS. This work is supported by NIH Grant R35-GM141930. Computational resources from ACCESS (project BIO230101) are greatly appreciated. Part of the computational work was performed on the Shared Computing Cluster which is administered by Boston University's Research Computing Services (URL: www.bu.edu/tech/support/research/). Anton 2 computer time (award MCB200062P) was provided by the Pittsburgh Supercomputing Center (PSC) through NIH Grant R01GM116961. The Anton 2 machine at PSC was generously made available by D.E. Shaw Research.

1. S. J. Wodak et al., Allostery in its many disguises: From theory to applications. *Structure* **27**, 566–578 (2019).
2. A Chatzigoulas, Z Cournia, Rational design of allosteric modulators: Challenges and successes. *WIREs Comput. Mol. Sci.* **11**, e1529 (2021).
3. R Nussinov, CJ Tsai, Allostery in disease and in drug discovery. *Cell* **153**, 293–305 (2013).
4. D Ni, et al., Along the allostery stream: Recent advances in computational methods for allosteric drug discovery. *WIREs Comput. Mol. Sci.* **11**, e1585 (2021).
5. R Nussinov, et al., Allostery: Allosteric cancer drivers and innovative allosteric drugs. *J. Mol. Biol.* **434**, 167569 (2022).
6. S Lu, et al., Emergence of allosteric drug-resistance mutations: new challenges for allosteric drug discovery. *Drug Disc. Today* **25**, 177–184 (2020).
7. S Osuna, The challenge of predicting distal active site mutations in computational enzyme design. *Wiley Interdiscip. Rev.: Comput. Mol. Sci.* **11**, e1502 (2021).
8. ND Taylor, et al., Engineering an allosteric transcription factor to respond to new ligands. *Nat. Methods* **13**, 177–183 (2016).
9. DC Sloas, JC Tran, AM Marzilli, JT Ngo, Tension-tuned receptors for synthetic mechanotransduction and intercellular force detection. *Nat. Biotech.* pp. DOI10.1038/s41587–022–01638–y (2023).
10. O Dagliyan, NV Dokholyan, KM Hahn, Engineering proteins for allosteric control by light or ligands. *Nat. Protoc.* **14**, 1863–1883 (2019).
11. M Shaaya, et al., Light-regulated allosteric switch enables temporal and subcellular control of enzyme activity. *eLife* **9**, e60647 (2020).
12. J Chen, YL Vishweshwaraiah, NV Dokholyan, Design and engineering of allosteric communications in proteins. *Curr. Opin. Struct. Biol.* **73**, 102334 (2022).
13. YL Vishweshwaraiah, J Chen, VR Chirasani, ED Tabdanov, NV Dokholyan, Two-input protein logic gate for computation in living cells. *Nat. Commun.* **12**, 6615 (2021).
14. GM Suel, SW Lockless, MA Wall, R Ranganathan, Evolutionarily conserved networks of residues mediate allosteric communication in proteins. *Nat. Struct. Biol.* **10**, 59–69 (2003).
15. KF O'Rourke, SD Gorman, DD Boehr, Biophysical and computational methods to analyze amino acid interaction networks in proteins. *Comput. Struct. Biotech. J.* **14**, 245–251 (2016).
16. KA Reynolds, RN McLaughlin, R Ranganathan, Hot spots for allosteric regulation on protein surfaces. *Cell* **147**, 1564–1575 (2011).
17. A VanWart, J Eargle, Z Luthey-Schulten, RE Amaro, Exploring residue component contributions to dynamical network models of allostery. *J. Chem. Theory Comput.* **8**, 2949–2961 (2012).
18. VA Feher, JD Durrant, A. T. Van Wart, RE Amaro, Computational approaches to mapping allosteric pathways. *Curr. Opin. Struct. Biol.* **25**, 98–103 (2014).
19. I Rivalta, V Batista, *Allostery: Methods and Protocols*, Methods in Molecular Biology, eds. L DiPaola, A Giuliani. (Springer) Vol. 2253, pp. 137–151 (2021).
20. U Doshi, MJ Holliday, EZ Eisenmesser, D Hamelberg, Dynamical network of residue-residue contacts reveals coupled allosteric effects in recognition, catalysis, and mutation. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 4735–4740 (2016).
21. J Wang, et al., Mapping allosteric communications within individual proteins. *Nat. Commun.* **11**, 3862 (2020).
22. WJ Zheng, BR Brooks, D Thirumalai, Low-frequency normal modes that describe allosteric transitions in biological nanomachines are robust to sequence variations. *Proc. Acad. Natl. Sci. U.S.A.* **103**, 7664–7669 (2006).
23. E Guarnera, IN Berezovsky, Toward comprehensive allosteric control over protein activity. *Struct.* **27**, 866–878 (2019).
24. PT Lake, RB Davidson, H Klem, GM Hocky, M. McCullagh, Residue-level allostery propagates through the effective coarse-grained hessian. *J. Chem. Theory Comput.* **16**, 3385–3395 (2020).
25. KCH Ngan, et al., Activity-based crispr scanning uncovers allostery in dna methylation maintenance machinery. *eLife* **12**, e80640 (2023).
26. GP Lisi, JP Loria, Solution nmr spectroscopy for the study of enzyme allostery. *Chem. Rev.* **116**, 6323–6369 (2016).
27. MJ Holliday, C Camilloni, GS Armstrong, M Vendruscolo, EZ Eisenmesser, Networks of dynamic allostery regulate enzyme function. *Struct.* **25**, 276–286 (2017).
28. C Köhler, et al., Dynamic allosteric communication pathway directing differential activation of the glucocorticoid receptor. *Sci. Adv.* **6**, eabb5277 (2020).
29. A Zhuravleva, EM Clerico, LM Gierasch, An interdomain energetic tug-of-war creates the allosterically active state in hsp70 molecular chaperones. *Cell* **151**, 1296–1307 (2012).
30. DM Fowler, S Fields, Deep mutational scanning: a new style of protein science. *Nat. Methods* **11**, 801–807 (2014).
31. S Raman, Systems approaches to understanding and designing allosteric proteins. *Biochemistry* **57**, 376–382 (2018).
32. CL Araya, DM Fowler, Deep mutational scanning: assessing protein function on a massive scale. *Trends Biotechnol.* **29**, 435–442 (2011).
33. AJ Faure, et al., Mapping the energetic and allosteric landscapes of protein binding domains. *Nature* **604**, 175 (2022).
34. M Leander, Y Yuan, A Meger, Q Cui, S Raman, Functional plasticity and evolutionary adaptation of allosteric regulation. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 25445–25454 (2020).
35. M Leander, Z Liu, Q Cui, S Raman, Deep mutational scanning and machine learning reveal structural and molecular rules governing allosteric hotspots in homologous proteins. *eLife* **11**, e79932 (2022).
36. G Meier, et al., Deep mutational scan of a drug efflux pump reveals its structure-function landscape. *Nat. Chem. Biol.* **19**, 440 (2022).
37. AS Garruss, KM Collins, GM Church, Deep representation learning improves prediction of LacI-mediated transcriptional repression. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2022838118 (2021).
38. KR Hanning, M Minot, AK Warrender, W Kelton, ST Reddy, Deep mutational scanning for therapeutic antibody engineering. *Trends Pharmacol. Sci.* **43**, 123–135 (2022).
39. N Azbukina, A Zharikova, V Ramensky, Intragenic compensation through the lens of deep mutational scanning. *Biophys. Rev.* **14**, 1161–1182 (2022).
40. TM Groseclose, et al., Biomolecular systems engineering: Unlocking the potential of engineered allostery via the lactose repressor topology. *Annu. Rev. Biophys.* **50**, 303–321 (2021).
41. AN Hersey, VE Kay, S Lee, MJ Realff, CJ Wilson, Engineering allosteric transcription factors guided by the laci topology. *Cell Sys.* **14**, 645–655 (2023).
42. NV Dokholyan, Controlling allosteric networks in proteins. *Chem. Rev.* **116**, 6463–6487 (2016).
43. MA Cruz, et al., A cryptic pocket in ebola vp35 allosterically controls rna binding. *Nat. Commun.* **13**, 2269 (2022).
44. E Papaleo, et al., The role of protein loops and linkers in conformational dynamics and allostery. *Chem. Rev.* **116**, 6391–6423 (2016).
45. I Galdadas, et al., Allosteric communication in class a β-lactamases occurs via cooperative coupling of loop dynamics. *eLife* **10**, e66567 (2021).
46. JG Greener, MJE Sternberg, Structure-based prediction of protein allostery. *Curr. Opin. Struct. Biol.* **50**, 1–8 (2018).
47. J Lee, et al., Surface sites for engineering allosteric control in proteins. *Science* **322**, 438–442 (2008).
48. R Kityk, M Vogel, R Schlecht, B Bukau, MP Mayer, Pathways of allosteric regulation in hsp70 chaperones. *Nat. Commun.* **6**, 8308 (2015).
49. BC Stanton, et al., Systematic transfer of prokaryotic sensors and circuits to mammalian cells. *ACS Synth. Biol.* **3**, 880–891 (2014).
50. W Hinrichs, et al., Structure of the Tet repressor-tetracycline complex and regulation of antibiotic resistance. *Science* **264**, 418–420 (1994).
51. C Kisker, W Hinrichs, K Tovar, W Hillen, W Saenger, The complex formed between tet repressor and tetracycline-mg^{2+} reveals mechanism of antibiotic resistance. *J. Mol. Biol.* **247**, 260–280 (1995).
52. P Orth, D Schnappinger, W Hillen, W Saenger, W Hinrichs, Structural basis of gene regulation by the tetracycline inducible tet repressor–operator system. *Nat. Struct. Biol.* **7**, 215–219 (2000).
53. Z Yu, SE Reichheld, A Savchenko, J Parkinson, AR Davidson, A comprehensive analysis of structural and sequence conservation in the TetR family transcriptional regulators. *J. Mol. Biol.* **400**, 847–864 (2010).
54. B Hecht, G Müller, W Hillen, Noninducible tet repressor mutations map from the operator binding motif to the c terminus. *J. Bacteriol.* **175**, 1206–1210 (1993).
55. G Müller, et al., Characterization of non-inducible Tet repressor mutants suggests conformational changes necessary for induction. *Nat. Struct. Biol.* **2**, 693–703 (1995).
56. JE Seedorff, ME Rodgers, R Schleif, Opposite allosteric mechanisms in TetR and CAP. *Protein Sci.* **18**, 775–781 (2009).
57. W Saenger, P Orth, C Kisker, W Hillen, W Hinrichs, The tetracycline repressor – a paradigm for a biological switch. *Angew. Chem. Int. Ed. Engl.* **39**, 2042–2052 (2000).
58. M Sevvana, et al., An exclusive α/β code directs allostery in TetR–peptide complexes. *J. Mol. Biol.* **416**, 46–56 (2012).
59. S Werten, J Schneider, GJ Palm, W Hinrichs, Modular organisation of inducer recognition and

allostery in the tetracycline repressor. *FEBS J.* **283**, 2102–2114 (2016).

60. MD Daily, JJ Gray, Allosteric communication occurs via networks of tertiary and quaternary
motions in proteins. *PLoS Comput. Biol.* **5**, e1000293 (2009).

61. AS Walker, WP Russ, R Ranganathan, A Schepartz, Rna sectors and allosteric function within
the ribosome. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 19879–19887 (2020).

62. Y Yuan, J Deng, Q Cui, Molecular dynamics simulations establish the molecular basis for the
broad allostery hotspots distribution in the tetracycline repressor. *J. Am. Chem. Soc.* **144**,
10870–10887 (2022).

63. F Haberl, H Lanig, T Clark, Induction of the tetracycline repressor: Characterization by
molecular-dynamics simulations. *Proteins: Struct. Funct. & Bioinf.* **77**, 857–866 (2009).

64. VJ Hilser, JO Wrabl, HN Motlagh, Structural and energetic basis of allostery. *Annu. Rev.*
Biophys. **41**, 585–609 (2012).

65. L Bonati, E Trizio, A Rizzi, M Parrinello, A unified framework for machine learning collective
variables for enhanced sampling simulations: mlcolvar. *J. Chem. Phys.* **159**, 014801 (2023).

66. S Mehdi, Z Smith, L Herron, Z Zou, P Tiwary, Enhanced sampling with machine learning: A
review, arxiv (2023), https://arxiv.org/abs/2306.09111. (accessed 27 Aug 2023).

67. MD Ward, et al., Deep learning the structural determinants of protein biochemical properties
by comparing structural ensembles with diffnets. *Nat. Commun.* **12**, 3023 (2021).

68. D Mendels, F Bylehn, T Sirk, JJ de Pablo, Systematic modification of functionality in disordered
elastic networks through free energy surface tailoring. *Sci. Adv.* **9**, eadf7541 (2023).

69. O Valsson, P Tiwary, M Parrinello, Enhancing important fluctuations: Rare events and metady-
namics from a conceptual viewpoint. *Annu. Rev. Phys. Chem.* **67**, 159–184 (2016).

70. A Barducci, G Bussi, M Parrinello, Well-tempered metadynamics: A smoothly converging and
tunable free-energy method. *Phys. Rev. Lett.* **100**, 020603 (2008).

71. M Bonomi, et al., Plumed: A portable plugin for free-energy calculations with molecular
dynamics. *Comp. Phys. Commun.* **180**, 1961–1972 (2009).

72. The PLUMED consortium, Promoting transparency and reproducibility in enhanced molecular
simulations. *Nat. Methods* **16**, 670–673 (2019).

73. P Tiwary, M Parrinello, A time-independent free energy estimator for metadynamics. *J. Phys.*
Chem. B **119**, 736–742 (2015).