Deep multiple-instance learning accurately predicts gene haploinsufficiency and deletion pathogenicity

Zhihan Liu[1,2,3*], Yi-Fei Huang[1,3*]

[1]Department of Biology, Pennsylvania State University,
University Park, PA 16802, USA

[2]Molecular, Cellular, and Integrative Biosciences Program, Pennsylvania State University,
University Park, PA 16802, USA

[3]Huck Institutes of the Life Sciences, Pennsylvania State University,
University Park, PA 16802, USA

*Correspondence: zul327@psu.edu, yuh371@psu.edu

<u>Abstract</u>

Copy number losses (deletions) are a major contributor to the etiology of severe genetic disorders.

Although haploinsufficient genes play a critical role in deletion pathogenicity, current methods for

deletion pathogenicity prediction fail to integrate multiple lines of evidence for haploinsufficiency at the

gene level, limiting their power to pinpoint deleterious deletions associated with genetic disorders. Here

we introduce DosaCNV, a deep multiple-instance learning framework that, for the first time, models

deletion pathogenicity jointly with gene haploinsufficiency. By integrating over 30 gene-level features

potentially predictive of haploinsufficiency, DosaCNV shows unmatched performance in prioritizing

pathogenic deletions associated with a broad spectrum of genetic disorders. Furthermore, DosaCNV

outperforms existing methods in predicting gene haploinsufficiency even though it is not trained on

known haploinsufficient genes. Finally, DosaCNV leverages a state-of-the-art technique to quantify the

contributions of individual gene-level features to haploinsufficiency, allowing for human-understandable

explanations of model predictions. Altogether, DosaCNV is a powerful computational tool for both

fundamental and translational research.

Background

Copy number variations (CNVs) are a major contributor to human genetic diversity[1,2]. These structural variations, which involve deletions or duplications of genomic regions ranging from 50 base pairs to several megabase pairs, result from diverse mutational mechanisms during DNA replication, recombination, and repair[3,4]. CNVs may provide adaptive advantages under certain circumstances[5,6]; however, they can also lead to diseases when critical genomic regions are affected[7]. Both common and rare CNVs (population frequency < 1%) contribute to disease development, with rare CNVs particularly implicated in rare genetic disorders[2,8–13]. In the context of disease-associated CNVs, pathogenicity often arises from alterations affecting dosage-sensitive (DS) driver genes[14,15]. This concept of dosage sensitivity refers to the emergence of deleterious phenotypes when gene copy number changes. Specifically, haploinsufficient (HI) genes are intolerant to copy number loss, whereas triplosensitive (TS) genes are intolerant to copy number gain[16,17]. Thus, identifying DS genes is essential for prioritizing clinically relevant CNVs and understanding the genetic mechanisms of associated diseases.

Efforts to discover DS genes primarily involve three frameworks: machine learning-based methods, case-control data-based statistical methods, and selection pressure-based approaches. Machine learning-based methods utilize gene-level features and various algorithms to differentiate manually curated DS genes from dosage-insensitive genes[18–23]. While these models identify predictive attributes of DS genes, the limited availability of high confidence training data restricts their ability to capture the full spectrum of DS genes. Alternatively, statistical methods can be utilized to analyze case-control CNV data to identify DS genes, assuming that these genes have a higher prevalence among affected individuals[15]. For instance, a recent approach employs a modified GWAS fine-mapping framework to estimate gene-level dosage sensitivity[24]. Although this framework can predict DS genes without labeled training data, it is challenging to pinpoint causal genes because CNVs often span multiple genes. Finally, selection pressure-based methods utilize natural selection against loss-of-function (LOF) mutations or CNVs as indicators of dosage

sensitivity[25–30]. Although dosage sensitivity is strongly correlated with selection pressure at the gene level,

exceptions do exist. For instance, alterations in pleiotropic genes critical for fitness may not lead to

significant phenotypic changes[31,32], and DS genes associated with late-onset disorders may not be subject

to strong selection pressure[33–36]. Additionally, these selection pressure-based methods may be

underpowered for detecting short DS genes due to sparse polymorphisms in the human genome[27,28].

Previous research prioritizes pathogenic CNVs using two frameworks: rule-based methods and supervised

machine learning methods. Rule-based approaches automate pathogenicity scoring systems based on

clinical CNV interpretation guidelines[37,38], such as the one proposed by the American College of Medical

Genetics and Genomics (ACMG) and ClinGen[39]. However, these methods face limitations due to rigid score

assignments and difficulties in identifying pathogenicity originating from uncatalogued genomic regions.

On the other hand, supervised machine learning methods, often trained on clinically curated CNVs, utilize

simple summary statistics aggregated across affected genomic elements, such as the mean score of

dosage sensitivity across genes, for pathogenicity prediction. However, the optimal summary statistics for

deletion pathogenicity prediction are often unknown, limiting the accuracy of supervised machine

learning methods. For instance, the mean score of dosage sensitivity across genes might not adequately

convey the significance of a pathogenic CNV harboring a single highly penetrant DS driver gene amidst

multiple dosage-insensitive genes.

Furthermore, while current methods offer valuable insights into gene dosage sensitivity and CNV

pathogenicity, they struggle to connect the two. DS gene identification methods predict gene dosage

sensitivity but lack a natural way of aggregating these predictions to infer CNV pathogenicity. Conversely,

pathogenic CNV classifiers predict CNV pathogenicity without identifying DS driver genes and evaluating

their contributions. To bridge this gap and enhance predictions at both the gene and CNV levels, we

propose that multiple-instance learning (MIL) presents a promising, unified solution for the joint

predictions of gene dosage sensitivity and CNV pathogenicity[40]. MIL is suited for problems where each

sample, or bag, in the dataset contains a varying number of observations, or instances, and is classified as

positive if at least one instance is positive[40]. The objective is to predict labels at both the bag and instance

levels, assuming that the bag-level label results from the combined effect of individual instances, even

when only bag-level labels are available during training. This approach is well-aligned with our problem,

as we can view each CNV as a bag and the affected genes within it as instances, with pathogenicity labels

given solely at the bag (CNV) level.

In this study, we introduce DosaCNV, a deep MIL framework modeling deletion pathogenicity through the

joint effect of haploinsufficiency from affected genes. We focus exclusively on deletions that affect

protein-coding genes, which have more direct functional consequences compared to deletions that only

affect non-coding regions and duplications. DosaCNV leverages over 30 gene-level features from diverse

biological processes and is trained on clinically curated deletions. It outperforms existing methods in

predicting pathogenic deletions and HI genes across various sources. DosaCNV demonstrates the potential

to leverage large-scale deletion data for enhancing the predictions of HI genes and associating these

predictions with deletion pathogenicity, establishing it as an accurate and valuable computational tool for

both fundamental and translational research.

Results

Overview of DosaCNV. DosaCNV is a supervised deep MIL model designed to simultaneously infer the

pathogenicity of coding deletions and the haploinsufficiency of genes, based on the assumption that the

joint effect of gene haploinsufficiency determines deletion pathogenicity. Utilizing a fully connected

neural network, chosen for its strong ability to automatically extract useful features, DosaCNV analyzes

coding deletions by first computing the haploinsufficiency of affected genes (Fig. 1). This fully connected

neural network processes an input matrix for each deletion, with rows representing affected genes and

columns representing gene-level features. Then, it generates a vector of predicted HI probabilities (P_{HI})

for each affected gene within the deletion (Fig. 1).

Subsequently, DosaCNV utilizes the noisy-or pooling function[41], a probabilistic variant of the logical-or

operator commonly used in Bayesian networks, to combine the predicted HI probabilities at the gene level

into a scalar output ranging from 0 to 1, representing the joint effect of gene-level haploinsufficiency (Fig.

1). This output is also the predicted probability that the given deletion overlaps with at least one HI gene

by the formalization of the noisy-or function[41], and serves as a proxy for deletion pathogenicity (Fig. 1).

Notably, due to the noisy-or pooling approach, DosaCNV assumes independent effects between affected

genes and does not account for gene-gene interactions.

DosaCNV's design addresses the challenge of limited access to curated HI genes faced by previous

supervised gene classifiers. By training the model's fully connected architecture for HI prediction on all

affected genes within the deletion data, DosaCNV potentially accesses a larger and more diverse

agglomeration of HI genes compared to currently available sets. This approach may enable DosaCNV to

capture distinct HI gene characteristics more effectively, enhancing HI gene prediction relative to other

DS gene classifiers. Combining this improvement with the probabilistic integration of HI predictions for all

affected genes within a deletion of interest, DosaCNV could potentially yield more accurate deletion-level

pathogenicity predictions.

Training DosaCNV on clinically curated deletions. We retrieved 6,180 pathogenic and 5,822 benign

deletions overlapping with gene coding sequences (CDSs) from ClinVar[42], a pan-disorder repository of

clinically curated genetic variants. Also, we collected 33 gene-level features potentially predictive of

haploinsufficiency. These gene-level features can be grouped into six categories: genomics, epigenomics,

evolutionary constraints, functional genomics, deleteriousness upon perturbation, and biological

networks (Supplementary Data 1). We used the features of affected protein-coding genes within a

deletion as the input features of DosaCNV, and used the clinically curated pathogenicity of the deletion in

ClinVar as the label. Then, we utilized a "leave-one-chromosome-out" approach to partition the data into

a training set, a validation set, and a held-out test set (Methods). We trained DosaCNV on the training set

and optimized the hyperparameters of DosaCNV on the validation set. To minimize potential bias arising

from relying on deletion length for predictions[43,44], we matched pathogenic and benign deletions based

on the number of affected protein-coding genes within the training and validation sets. The test set

remained unmatched for subsequent analysis. As a result, the final training, validation, and test sets

contained 2,308, 1,060, and 2,832 coding deletions, respectively.

Evaluating DosaCNV and alternative methods on ClinVar pathogenic deletions. We assessed DosaCNV's

performance in predicting pathogenic deletions on the held-out test set. To mitigate potential biases in

performance assessment due to class imbalance and length differences between pathogenic and benign

deletions, we benchmarked DosaCNV with five alternative methods (TADA[44], StrVCTVRE[45], HIS-LOD[18], X-

CNV[46], and CADD-SV[47]) on the unmatched held-out test set and three subsets matched for the number of

affected protein-coding genes, deletion length, and both criteria, respectively.

Within the unmatched test set, 1,211 pathogenic deletions and 1,280 benign deletions had predictions

from all five alternative methods (n = 2,491). DosaCNV achieved the highest area under the receiver-

operating-characteristic curve (AUC) of 0.968 (Fig. 2A; Supplementary Table 1). To evaluate the

performance of various methods for deletions with different lengths, we categorized the unmatched test

set into four length groups: small (<50 kb), medium (50-100 kb), large (100-1000 kb), and extra-large

(>1000 kb). DosaCNV consistently delivered superior or comparable predictions across all length groups

(Fig. 2B; Supplementary Table 2). Although other methods also performed reasonably well, their

performance might be attributed to differences in length distribution between pathogenic and benign

deletions (Supplementary Fig. 1A & B). Indeed, using either the number of affected genes or deletion

length alone as a predictor for deletion pathogenicity yielded AUC values as high as 0.886 and 0.912,

respectively (Fig. 2A).

To account for this potential confounder, we examined the influence of two forms of length information

on performance: the number of affected genes and deletion length. We generated two matched test sets

with a 1:1 ratio of pathogenic to benign deletions based on the number of affected genes (n=702) and

deletion length (n=704), respectively. DosaCNV maintained the best performance on these datasets,

achieving AUC values of 0.895 and 0.891, respectively (Supplementary Fig. 2A – 2D; Supplementary Table

1 & 3). Nevertheless, a substantial decrease in performance for all methods was observed when compared

to the unmatched test set, suggesting that differences in length distribution contributed to the overly

optimistic evaluations in unmatched data.

Furthermore, we noted that controlling for one form of length information may not necessarily account

for the other (Supplementary Fig. 1A & B), as pathogenic and benign deletions with similar nucleotide

spans might not overlap with a comparable number of genes, and vice versa. To further minimize the

impact of length information, we generated a test set matched for both deletion length and the number

of affected genes (n=618). This strategy resulted in AUC values of approximately 0.5 for both the length

features (deletion length and gene number) in the matched test set and across different length groups

(Fig. 2C & D; Supplementary Table 1 & 2), indicating that these two length features could no longer be

used for prediction. Even though the overall performance of almost all methods was further reduced,

DosaCNV remained the top-performing method (Fig. 2C). DosaCNV outperformed other methods in all

length groups, except for the small group (Fig. 2D), where XCNV's performance was marginally better but

not statistically significant (DeLong test p-value = 0.582). By comparing the AUC values of this matched

test set to the unmatched test set, we estimated each method's degree of dependency on length

information. CADD-SV demonstrated the greatest dependency with a 0.227 decrease in AUC (unmatched:

0.888, matched: 0.661), while TADA exhibited the smallest dependency with a 0.080 decrease in AUC

(unmatched: 0.923, matched: 0.843). DosaCNV had the second smallest dependency, with a 0.091

decrease in AUC (unmatched: 0.968, matched: 0.877).

DosaCNV enhanced prioritization of putative pathogenic deletions in neurodevelopmental disorder

case-control studies. To further assess DosaCNV's capacity to discriminate between pathogenic and

benign deletions, we compared its performance to alternative methods in prioritizing potential

pathogenic deletions from case-control studies of neurodevelopmental disorders (NDDs). While ground

truth labels for deletions in case-control datasets are unknown, prior research indicates that CNVs may

contribute to 15-25% of NDD cases[48,49]. As a result, an effective pathogenic CNV classifier should prioritize

these pathogenic CNVs, placing them within the top predictions among all CNVs from cases and controls,

with the top-ranked CNVs predominantly depleted in controls. Thus, we assessed the performance of each

method based on the enrichment of case-associated deletions within their top predictions, using this as a

measure of their ability to prioritize putative pathogenic deletions. Specifically, for the analysis of each

dataset, we established the number of top predictions for evaluation as 5% of the total case count

(referred to as the top 5% predictions). Based on the conservative assumption that coding CNVs contribute

to 10% of cases, with half of these CNVs being deletions.

We examined 16,048 coding deletions from Coe et al., 2014 (dbVar accession number: nstd100)[50],

comprising 29,085 intellectual disability, developmental delay, and autism spectrum disorder (ID/DD/ASD)

cases, alongside 19,584 controls. Subsequently, we evaluated 4,230 coding deletions from Zarrei et al.,

2019 (dbVar accession number: nstd173)[51], which included 2,691 NDD cases and 1,769 controls. Within

the nstd100 dataset, DosaCNV had the highest enrichment of case-associated deletions among its top 5%

predictions (odds ratio = 102.3), which significantly outperforms the second-best method HIS-LOD (odds

ratio = 31.1; z-test for log odds ratios p-value = 1.026×10^{-10}) (Fig. 3A; Supplementary Table 4). Specifically,

DosaCNV prioritized 84 more case-associated deletions among top predictions compared to HIS-LOD,

suggesting a potential to detect 5.8% (84/1,454) more putative pathogenic deletions. For the nstd173

dataset, DosaCNV again had the highest enrichment of case-associated deletions among its top 5%

predictions (odds ratio = 2.12) (Fig. 3B; Supplementary Table 4). However, this did not significantly differ

from the next best method TADA (odds ratio = 1.79; z-test for log odds ratios p-value = 0.562), likely

attributable to the smaller sample size.

Additionally, we investigated the overlap between the two case-control datasets and the ClinVar test set

to provide further evidence for DosaCNV's capacity to prioritize pathogenic deletions. We posited that

deletions with over 95% overlap should exhibit similar functional consequences and, therefore,

comparable pathogenicity. We identified deletions from the case-control datasets with over 95% overlap

with true pathogenic deletions from the ClinVar test set. Then, we assessed whether they were primarily

found in cases and prioritized as top predictions by DosaCNV. In the nstd100 dataset, we uncovered 164

overlapping deletions, with 154 observed in cases, substantiating the depletion of most likely pathogenic

deletions within controls. Furthermore, 148 of these 164 deletions were included in DosaCNV's nstd100

top predictions, demonstrating its robust prioritization capability. For the nstd173 dataset, we observed

13 deletions overlapping with true pathogenic deletions, with 11 of them present in cases. Only 4 of the

13 overlapping deletions were among the top 5% predictions of DosaCNV, possibly due to the smaller

sample size of the overlapping deletions. Notably, DosaCNV did not prioritize any deletions that

overlapped with benign ClinVar deletions as top predictions in either dataset.

Finally, we assessed DosaCNV's capacity to distinguish pathogenic deletions from benign deletions in

patients, a prevalent and challenging clinical scenario when identifying causal variants. We obtained 740

clinically curated coding deletions within 140 DD cases from Foong et al., 2015[52], which included 33

pathogenic deletions. DosaCNV effectively prioritized pathogenic deletions among benign deletions (Fig.

3C), attaining an AUC of 0.997, which was the highest among all methods (Supplementary Fig. 3). However,

the difference in AUC was not statistically significant compared to the second-best method StrVCTVRE

(AUC = 0.981; DeLong test p-value = 0.115). With a DosaCNV score threshold of 0.8, the true positive rate

reached 0.970, and the false positive rate was 0.028, indicating the accurate classification of almost all

pathogenic deletions with only a small fraction of benign deletions misclassified. The precision of DosaCNV

was also high (0.604) at this threshold. These results suggest that DosaCNV can accurately distinguish

pathogenic deletions from benign deletions in patients, and 0.8 is a reasonable threshold in the practice

of clinical variant interpretation.

DosaCNV-HI outperforms alternative methods in predicting HI genes from multiple sources. To

investigate whether the superior performance of DosaCNV at the deletion level is due to more accurate

gene haploinsufficiency estimation, we evaluated our model's ability to distinguish HI and likely HI genes

from haplosufficient (HS) genes. First, we assembled six positive sets of HI and likely HI genes, including:

1) 340 ClinGen "gold standard" HI genes[53], 2) 397 human orthologs of mouse heterozygous knockout

lethal (MHKO) genes[54], 3) 214 high-confidence SFARI ASD genes (SFARI gene score 1)[55], 4) 432 DDG2P

genes sensitive to heterozygous LOF mutations[56], 5) 72 eyeG2P genes sensitive to heterozygous LOF

mutations[56], and 6) 82 skinG2P genes sensitive to heterozygous LOF mutations[56]. We defined all other

genes as the HS gene set. To balance the number of genes and mitigate the impact of CDS length on model

performance, we used MatchIt[57] to match HI and HS genes by CDS length, retaining only matched genes

for downstream performance analysis.

To circumvent circularity, we retrained DosaCNV by excluding DDG2P and EDS scores from gene-level

input features, as a subset of DDG2P genes was used for gene-level prediction assessment, and the EDS

score was trained on a portion of our assembled HI genes. After training, the DosaCNV gene-level model,

DosaCNV-HI (Methods), generated HI probability predictions for 20,268 protein-coding genes.

We assessed DosaCNV-HI's performance against 12 alternative methods across six datasets,

encompassing two metrics for LOF mutation intolerance (DeepLOF[29], and LOEUF[28]), one for missense

mutation intolerance (Mis. OEUF[28]), two for both LOF and missense mutation intolerance (RVIS[26],

VIRLoF[58]), and seven for haploinsufficiency (HIS[18], IS[23], GHIS[19], HIPred[20], Episcore[21], EDS[22], pHaplo[24]). The

comparison included genes with predictions available from all methods. To evaluate performance robustly

within imbalanced sets, we computed the AUC for each method by sampling an equal number of HS genes

of matching CDS length as the HI or likely HI gene sets.

DosaCNV-HI significantly surpassed alternative methods in predicting ClinGen HI genes, achieving an AUC

of 0.901, which is substantially higher than the second-best method, Episcore, with an AUC of 0.854

(Delong test p-value = 0.019) (Fig. 4A; Supplementary Table 5). DosaCNV-HI consistently achieved high

rankings in the remaining five sets of likely HI genes. Notably, even in cases where DosaCNV-HI was not

the top performer (ranking second or third), the differences in the AUCs were not statistically significant

compared to those of the best methods (Fig. 4B-F; Supplementary Table 5). It is worth noting that the

performance of DosaCNV-HI could still be overly optimistic when analyzing all available genes without

considering overlap between HI/HS genes and deletions in the ClinVar training set. To address this, we

conducted a separate analysis excluding genes overlapped by deletions in the ClinVar training set, and

DosaCNV-HI maintained its high-ranking performance across all six gene sets (Supplementary Fig. 4;

Supplementary Table 6).

Evaluating the robustness of DosaCNV-HI and alternative methods in identifying challenging HI and HS

genes. Metrics based on evolutionary constraints, such as LOEUF and DeepLOF, demonstrated reasonable

performance, indicating that many HI genes are indeed subject to strong purifying selection in human

populations. However, these metrics, along with supervised HI gene scores dependent on them, may face

challenges in identifying HI genes in three particular cases: 1) short HI genes with limited statistical power

for constraint inference[27,28], 2) HI genes under low evolutionary constraint, including those involved in

late-onset disorders (e.g., *BRCA2*, *ATM*, and *ELN*)[33–36], and 3) falsely predicted HS genes under high

evolutionary constraint, such as pleiotropic genes with minor effects on single phenotypes but significant

cumulative fitness impacts[31,32]. To evaluate the robustness of our method and alternative approaches in

these scenarios, we generated three challenging gene sets.

First, we combined all six HI and likely HI gene sets to form a unified set of 988 genes, aiming to enhance

statistical power. From this combined set, we focused on the 758 genes that have predictions across all

methods for further analysis. Next, we filtered this set for genes with fewer than 10 expected LOF

mutations in human populations[29], creating a "short, likely HI" gene set (n=73). Second, using an

established LOEUF cutoff (<0.35 for constrained genes) and a similar cutoff for DeepLOF (>0.835 for

constrained genes), we generated a "low constraint, likely HI" gene set (n=278). Finally, we applied the

same cutoffs to identify a set of highly evolutionarily constrained HS genes not implicated in diseases

across G2P, Human Phenotype Ontology[59], ClinVar[42], and OMIM databases[60], yielding a "high constraint,

unlikely HI" gene set (n=209).

We identified the 2,402 top-ranked genes for each method as predicted HI genes, comparable to the

LOEUF cutoff of 0.35, and evaluated performance based on the percentage of each challenging gene set

included within the top-ranked genes. An ideal HI gene classifier should maximize the inclusion of genes

from the two likely HI gene sets while minimizing the inclusion of genes from the unlikely HI gene set.

Among all evaluated methods for the "short, likely HI" gene set, HIS exhibited the highest inclusion

percentage (0.466, 34/73), followed by Episcore (0.411, 30/73), and then our method, DosaCNV-HI (0.397,

29/73) (Fig. 4G; Supplementary Table 7). Notably, almost all methods showed a higher inclusion

percentage compared to LOEUF. This suggests that these methods, potentially due to the integration of

diverse complementary gene-level features, are less affected by the limited statistical power in detecting

short, constrained genes. For the "low constraint, likely HI" gene set, DosaCNV-HI and HIS achieved the

highest (0.439, 122/278) and second highest (0.414, 115/278) inclusion percentages, respectively (Fig. 4H;

Supplementary Table 7). This outcome may be attributed to the fact that these two methods directly

model haploinsufficiency instead of evolutionary constraints.

Remarkably, using gene CDS length alone as a predictor resulted in the lowest inclusion percentage (0.072,

15/209) for the "high constraint, unlikely HI" gene set (Fig. 4I; Supplementary Table 7). In contrast,

DosaCNV-HI exhibited the second lowest inclusion percentage (0.148, 31/209), with CDS length excluding

7.6% more unlikely HI genes (Fig. 4I). This observation emphasizes a positive correlation between gene

haploinsufficiency and CDS length for evolutionarily constrained genes, aligning with prior research that

suggests longer genes tend to be more crucial during early developmental stages[61], with

haploinsufficiency potentially serving as an inherent property of these genes[62]. In summary, while

DosaCNV-HI may not always be the top-performing method, it consistently ranks highly for predicting

challenging HI genes and difficult HS genes.

Elucidating key features and their contributions to gene haploinsufficiency prediction. We applied Deep

SHAP[63] (SHapley Additive exPlanations) alongside DosaCNV-HI to identify key features determining a

gene's probability of being HI. Based on coalition game theory, Deep SHAP fairly and uniquely allocates

an input's prediction to its features (SHAP values) by computing a weighted average of marginal

contributions across all feature combinations. We obtained SHAP values for each feature across 988 HI

and likely HI genes using Deep SHAP and sampled 1,000 HS genes as a background set. We then visualized

these values using a default SHAP summary plot[63], offering a global explanation for the entire HI and likely

HI gene set (Fig. 5A). The global feature importance is derived from the mean absolute SHAP values over

all HI and likely HI genes. Pathogenic SNV presence emerged as the most important feature, as genes

sensitive to disrupted SNVs may also be sensitive to dosage changes. Features linked to evolutionary

constraints and conservation, such as PLI, LOEUF, and PhastCons, also ranked highly, emphasizing the

importance of evolutionary constraints and conservation as common HI gene characteristics (Fig. 5A).

Notably, we observed a positive correlation between the second most important feature, the number of

transcript isoforms, and the DosaCNV-HI prediction (Fig. 5A). This finding is in line with a previous study

showing that genes with a greater number of isoforms are more likely to be essential genes[64]. The sixth

most important feature, tissue specificity (tau), also exhibited a positive correlation with the DosaCNV-HI

prediction (Fig. 5A), which implies that HI genes may have greater tissue specificity in expression than HS

genes[18]. Epigenetically, higher DosaCNV-HI predictions were associated with stronger signals for

repressive histone modifications (H3K27me3) and weaker signals for active modifications (H3K4me3 and

H3K9ac) (Fig. 5A). Considering that the majority of epigenetic data were obtained from adult

tissues/organs and most HI genes are developmental genes[15], this result suggests that many HI genes may

be epigenetically silenced outside early developmental stages. Intriguingly, we observed a positive

correlation between the H2A.Z signal and DosaCNV-HI predictions, although this histone variant is known

to facilitate both gene activation and repression[65]. Additionally, several features, such as a greater number

of enhancers, increased promoter CpG density, more extensive protein-protein interactions, elevated

mean expression levels, and fewer paralogs, contributed to higher DosaCNV-HI predictions (Fig. 5A),

aligning with previous studies[18,21,66].

Deep SHAP also offers local explanations by attributing the HI prediction to individual feature

contributions for a specific gene. To demonstrate this, we retrieved coordinates of five recurrent

pathogenic microdeletions from ClinGen[67], and obtained DosaCNV-HI predictions and corresponding

SHAP values for 30 protein-coding genes overlapping with these microdeletions (Fig. 5B). DosaCNV-HI

identified candidate disease genes with plausible haploinsufficiency mechanisms, consistent with previous

findings, such as *CHD1L* and *GJA8* for the 1q21.1 deletion[68,69], *CHRNA7* for the 15q13.3 deletion[70,71],

PMP22 for the 17p12 deletion[72], *CRHR1* and *KANSL1* for the 17q21.31 deletion[73,74], and *SHANK3* for the

22q13 deletion[75] (Fig. 5B). Comparing the SHAP values of all features for these candidate genes allows us

to understand why they are predicted to have a higher chance of being HI. For example, in the well-

documented HI gene *PMP22*, encoding the essential peripheral myelin protein 22 for myelin formation

and stability[72], key contributors to its HI prediction revealed by SHAP values include pathogenic SNVs

presence, numerous transcript isoforms, evolutionary constraints, enhancer abundance, and extensive

protein interactions (Fig. 5B). These features highlight *PMP22*'s regulatory complexity within the myelin

gene network, which implicates the necessity of precise dosage control for its role in proper neural

development and function[72]. Similarly, *KANSL1*, another established HI gene involved in chromatin

modification and gene regulation[74], demonstrated a comparable pattern of feature importance, though

adapted to its unique biological functions (Fig. 5B). These insights elucidate the rationale behind

DosaCNV's predictions for each gene and deletion, ultimately improving our capacity to explore the

biological mechanisms underpinning gene haploinsufficiency and deletion pathogenicity.

Discussion

In this study, we present DosaCNV, a novel deep MIL model that jointly predicts coding deletion

pathogenicity and gene haploinsufficiency. Prior research has examined these predictions independently,

without effectively connecting the two. Our approach, for the first time, bridges this gap in a biologically

coherent manner, which significantly enhances predictions at both the gene and deletion levels. Moreover,

this method enables the evaluation of HI genes' contribution to deletion pathogenicity and expands the

spectrum of learned HI gene features. Additionally, we employ rigorous training and data preprocessing

strategies, such as size matching and leave-one-chromosome-out, to strengthen model generalizability

and mitigate biases arising from information leakage[76].

While DosaCNV demonstrated superior performance to other prediction tools in ClinVar data, the results

also revealed a notable challenge shared by all methods, including DosaCNV: the dependence of

predictions on deletion length. While a strong positive association exists between deletion length and

pathogenicity, it is not the primary determinant of pathogenicity. Overreliance on this non-causal

association may lead to biased predictions for long benign deletions and short pathogenic deletions[43].

Furthermore, as machine-learning models can easily capture this strong correlation and maintain high

accuracy within unmatched datasets, the impact of other features directly involved in the underlying

mechanisms of pathogenic deletions might be overshadowed. This emphasizes the importance of taking

the length information into account during both model development and performance evaluation.

A robust pathogenic CNV classifier must be generalizable to effectively prioritize putative pathogenic

deletions across various disease contexts, addressing the growing accumulation of unlabeled CNV data.

DosaCNV, trained on a limited set of pan-disorder deletions, exhibited superior performance in prioritizing

putative pathogenic deletions in ID/DD/ASD cases, as evidenced by the highest enrichment of case-

associated deletions in predicted pathogenic deletions and its favorable sensitivity in prioritizing clinically

curated pathogenic deletions in a real clinical scenario. This generalizability supports DosaCNV's

underlying hypothesis that the pathogenicity of coding deletions predominantly results from affected HI

genes, which may share cross-disease commonalities captured by the gene-level features used in the

model. However, the performance may also be influenced by the enrichment of ID/DD/ASD patients in

our pan-disorder training data. While HI genes often share certain characteristics, their effects may also

differ in a disease-dependent manner based on their involvement in distinct biological processes. To

account for the variation of gene effects across diseases, DosaCNV is designed for adaptability, enabling

easy training with diverse datasets and gene-level features tailored to the specific disease context under

investigation.

Data scarcity in predicting gene haploinsufficiency poses a significant challenge for supervised machine-

learning methods, with fewer than 1,000 high-confidence HI and likely HI genes available out of an

estimated ~2,900[27,28]. DosaCNV adopts an innovative model design, learning predictive features for

haploinsufficiency from affected genes within curated deletions, without requiring gene-level labels. This

approach enables a more comprehensive capture of the HI gene landscape, allowing DosaCNV-HI to

outperform alternative methods in six HI and likely HI gene sets. However, our model remains dependent

on a limited number of clinically curated deletions, which may be subject to ascertainment bias, such as

the enrichment of certain types of genes and deletions[45]. Consequently, developing larger and less biased

datasets of curated CNVs is crucial for improving novel HI gene discovery and pathogenic deletion

prediction.

Evolutionary constraint metrics, such as pLI and LOEUF, have effectively prioritized disease-associated

genes based on the idea that deleterious mutations in disease genes are often depleted through purifying

selection. While these methods primarily focus on fitness effects, they can also suggest haploinsufficiency,

a phenotypic effect, especially for HI genes involved in early developmental processes where carrying a

null allele can result in reduced fitness. However, exceptions exist where reduced fitness does not align

with haploinsufficiency. For example, mutations in late-onset HI genes may not significantly reduce fitness,

and mutations in pleiotropic genes may lead to reduced fitness without causing discernible phenotypic

changes[32]. Moreover, evolutionary constraint metrics based on population genetics data are known to be

underpowered for short genes with sparse polymorphisms[27–29]. To address these challenges,

incorporating gene features orthogonal to evolutionary constraint metrics is crucial. By integrating over

30 gene-level features from diverse biological processes, DosaCNV-HI has demonstrated promising

performance in scenarios that pose challenges for traditional evolutionary constraint metrics (Fig. 4G, H,

I), underscoring the importance of considering a broader array of gene features to enhance the accuracy

and reliability of haploinsufficiency predictions.

To address the limited interpretability of deep learning models, we integrated DosaCNV-HI with a model-

specific explanation tool, Deep SHAP[63], to elucidate key gene features that highlighted the biological

characteristics of HI genes (Fig. 5A). Several metrics for evolutionary constraints, such as pLI and LOEUF,

emerged as top features, reinforcing the idea that HI gene disruption frequently leads to developmental

disorders and is thus subject to strong purifying selection[15,62]. Epigenetically, higher DosaCNV-HI

predictions correlated with weaker signals for active histone modification marks (H3K4me3, H3K9ac) and

stronger signals for repressive marks (H3K27me3). This supports the hypothesis that a substantial number

of HI genes function as developmental transcription factors, which are critical during specific

developmental stages and become epigenetically silenced outside these periods[77,78]. Additionally, our

results revealed a positive correlation between DosaCNV-HI predictions and H2A.Z signal, which can be

attributed to H2A.Z's prevalent occurrence in bivalent domains within promoters of developmental

genes[65], where HI genes are notably enriched. Regarding regulatory elements, higher enhancer numbers

and promoter CpG densities were associated with increased DosaCNV-HI predictions. These

characteristics have been associated with essential developmental genes[21,66,79,80], enabling intricate

spatiotemporal expression patterns associated with hypothesized mechanisms of haploinsufficiency[81].

Certain limitations in our study highlight the need for further refinement to improve the accuracy and

comprehensiveness of our method. First, we focused on coding regions in the current study, but the

potential contributions of non-coding regions to CNV pathogenicity should not be overlooked[82,83]. Ignoring

non-coding regions constrains the interpretation of non-coding CNVs and may impair coding CNV

predictions by erroneously assigning pathogenicity to seemingly innocuous coding elements. Second,

DosaCNV's assumption that independent effects of HI genes are the sole contributors to deletion

pathogenicity oversimplifies the complex interplay between genomic elements. This simplification fails to

account for the epistasis between affected genomic elements and variable penetrance/expressivity

arising from other genetic variants, such as SNVs[84–86]. Lastly, we assumed heterozygosity for all deletions

in the training and testing sets due to the absence of zygosity information in ClinVar. This assumption is

partially supported by the overrepresentation of dominant disease genes and limited overlap with

gnomAD benign SVs in ClinVar pathogenic deletions[45]. However, we observed a few pathogenic deletions

affecting only single recessive disease genes, suggesting the potential existence of homozygous deletions.

Consequently, the accuracy of our HI gene and pathogenic deletion predictions could be impacted, given

that DosaCNV was specifically designed to predict haploinsufficiency which is a dominant effect. The

future availability of fully phased CNV data will be instrumental in facilitating a more comprehensive

understanding of the relationships between the mode of inheritance of disease genes and the

pathogenicity of CNVs.

Despite the exclusive focus on deletions in this study, our proposed framework can be suitably adapted

to predict the pathogenicity of duplications. This adaptation essentially involves shifting the focus of gene-

level prediction from haploinsufficiency to triplosensitivity, achievable by simply utilizing curated

duplications as training data. Despite maintaining the same framework structure, a reassessment of

predictive features is needed, given the potential discrepancies between HI and TS genes. The integration

of appropriate input features will further elucidate the unique characteristics of TS genes.

Conclusions

In conclusion, DosaCNV is an accurate and robust deep multiple-instance learning framework that

simultaneously predicts deletion pathogenicity and gene haploinsufficiency. Traditional methods have

often treated these domains in isolation. DosaCNV, however, offers a unified approach to bridge this gap,

enhancing predictions at both the deletion and gene levels evidenced in multiple datasets. Furthermore,

by integrating the Deep SHAP technique to quantify gene-level feature contributions, DosaCNV not only

provides transparent and human-understandable explanations for its predictions but also holds promise

in further elucidating the intricate landscape of gene haploinsufficiency. However, challenges persist,

notably potential biases in clinically curated CNV datasets, with the ClinVar dataset being an example.

Despite our rigorous efforts during model development and evaluation, these biases inevitably limit our

model's precision and overall applicability. As our understanding of previously underexplored genomic

regions, such as non-coding regions, expands in tandem with rapid advancements in sequencing

technology, we anticipate the emergence of more comprehensive and unbiased CNV catalogs and

supporting resources. Such developments will be pivotal in improving the accuracy and reliability of tools

like DosaCNV, while also expanding their range of applications.

Methods

Details of DosaCNV. We denote $\mathbf{X}_i$ as the input for the i-th deletion, which is a matrix of dimensions M x

L, where M represents the number of genes and L corresponds to the number of gene-level features.

Given that the number of genes varies for each deletion, each $\mathbf{X}_i$ is automatically padded to accommodate

100 genes if M < 100 by adding dummy genes with all feature values set to zero. Consequently, $\mathbf{X}_i$ assumes

a shape of 100 x L. In the fully connected-layer, we aim to model HI probabilities for each of the M genes

within the i-th deletion, utilizing their respective gene-level feature vectors (rows of $\mathbf{X}_i$). To achieve this,

we employ two hidden layers for feature representation learning from $\mathbf{X}_i$. The calculations for these layers

can be represented by the following equations:

$$\mathbf{Z}_i^{(1)} = \mathrm{ReLU}(\mathbf{X}_i\mathbf{W}^{(1)} + \mathbf{b}^{(1)}),$$

$$\mathbf{Z}_i^{(2)} = \mathrm{Dropout}(\mathrm{ReLU}(\mathbf{Z}_i^{(1)}\mathbf{W}^{(2)} + \mathbf{b}^{(2)})),$$

where superscripts (1) and (2) denote the 1st and 2nd hidden layers, respectively. ReLU and Dropout refer

to the rectified linear layer[87] and dropout layer[88], while $\mathbf{W}$ and $\mathbf{b}$ represent the weight matrix and bias

vector, respectively. $\mathbf{Z}_i$ serves as the matrix of hidden units for the i-th deletion, with each row

constituting the hidden unit vector for a gene. Subsequently, we incorporate an additional layer to convert

$\mathbf{Z}^{(2)}$ into HI probabilities ($\mathbf{P}_{\mathrm{HI}}$) for the genes:

$$\mathbf{P}_{\mathrm{HI},i} = \sigma(\mathbf{Z}_i^{(2)}\mathbf{W}^{(3)} + \mathrm{b}^{(3)}),$$

where σ denotes the logistic function, which ensures that the HI probabilities remain within the range of

0 and 1. $\mathbf{W}^{(3)}$ and $\mathrm{b}^{(3)}$ represent the weight vector and bias term, respectively. $\mathbf{P}_{\mathrm{HI},i}$ is a vector of length

100, wherein each element serves as the gene-level prediction of haploinsufficiency (DosaCNV-HI score)

for a gene or dummy gene within the i-th deletion. To naturally aggregate probabilities within $\mathbf{P}_{\mathrm{HI},i}$ and

infer deletion pathogenicity, we introduce a pooling layer that employs the noisy-or function:

$$\hat{y}_i = 1 - \prod_{j=1}^{100} (1 - P_{HI,i,j} \times \text{Mask}_{i,j}),$$

where $\hat{y}_i$ is the predicted probability of the i-th deletion harboring at least one HI gene and serves as a proxy for deletion pathogenicity. $P_{HI,i,j}$ denotes the P_{HI} for the j-th gene or dummy gene within the i-th deletion. $\text{Mask}_{i,j}$ is a binary indicator, specifying whether the j-th gene is an actual gene ($\text{Mask}_{i,j} = 1$) or a dummy gene ($\text{Mask}_{i,j} = 0$). This mechanism prevents the propagation of information from dummy genes by setting $P_{HI,i,j}$ to zero.

We use the mean binary cross entropy loss to measure the discrepancy between the labels of deletion pathogenicity and predicted pathogenicity probabilities:

$$\text{loss}(\boldsymbol{y}, \hat{\boldsymbol{y}}) = -\frac{1}{N}\sum_{i=1}^{N}[y_i \log(\hat{y}_i) + (1 - y_i)\log(1 - \hat{y}_i)],$$

where $\boldsymbol{y}$ and $\hat{\boldsymbol{y}}$ represent the vectors of the observed pathogenicity labels and predicted pathogenicity probabilities in a mini-batch of data of size N, respectively.

For model training, we initialize all parameters using He Normal initialization[89] and employ the Adam algorithm[90] for mini-batch gradient descent to update model parameters. To mitigate overfitting, we implement early stopping and incorporate an L2 regularization term.

Following the training process, the full model is utilized for variant-level prediction of deletion pathogenicity. Concurrently, all layers of the full model are extracted, excluding the noisy-or pooling layer, to establish a model for gene-level prediction of haploinsufficiency, referred to as DosaCNV-HI.

ClinVar training, validation, and test set. We obtained the clinically curated rare SVs dataset (GRCh37; dbVar accession number: nstd102) from ClinVar[42] on May 25, 2021. We filtered for copy number losses/deletions meeting specific criteria: (1) >49 bp in length, (2) not somatic origin, (3) at least 1 bp overlap with a CDS, (4) <101 affected protein-coding genes, and (5) clinical interpretations as pathogenic, likely pathogenic, pathogenic/likely pathogenic, benign, likely benign, or benign/likely benign. Inner start

and end coordinates were also required for inclusion. We retrieved coordinates for 20,268 protein-coding

genes and their corresponding CDSs from the GENCODE[91] v38 human basic gene annotation file in GRCh37

and used BEDTools[92] to determine overlaps with deletions.

To minimize overfitting due to potential ascertainment bias, we identified redundant deletion groups with

95% reciprocal overlap using BEDTools and retained only the shortest deletion within each group. This

process yielded 12,002 deletions, which were labeled as pathogenic (1) if they had clinical interpretations

as pathogenic, likely pathogenic, or pathogenic/likely pathogenic and benign (0) if they had clinical

interpretations as benign, likely benign, or benign/likely benign (6,180 pathogenic and 5,822 benign).

To further minimize overfitting and avoid overly optimistic evaluations due to partial overlaps between

deletions in the same chromosome, we adopted a "leave-one-chromosome-out" strategy to partition the

deletions into training, validation, and testing sets. The test set included deletions from chromosomes 1,

3, 5, and 7, while the validation set comprised deletions from chromosomes 2, 4, 6, and 8, and the training

set contained deletions from the remaining chromosomes. This approach produced 6,678, 2,492, and

2,832 deletions for the training, validation, and testing sets.

We used R package MatchIt[57] to match pathogenic and benign deletions by the number of affected

protein-coding genes in the training and validation sets; only matched pairs were kept. We left the test

set unmatched for further analysis of the length impact on performance evaluation. This approach

resulted in 2,308 and 1,060 deletions in the final training and validation sets, respectively.

DosaCNV gene-level features. DosaCNV leverages 33 gene-level features of 20,268 protein-coding genes

to predict gene haploinsufficiency and subsequently infer deletion pathogenicity. The gene-level features

can be grouped into six main categories: genomics, epigenomics, evolutionary constraints, functional

genomics, deleteriousness upon perturbation, and biological networks (Supplementary Data 1). (1)

Genomics: We used the Ensembl biomaRt R package[93] to extract the number of coding exons and

transcript isoforms for each gene. We sourced promoter CpG density from a previous study[66]. We

retrieved a list of 3,543 human duplicated genes, organized into 945 groups, from the Duplicated Gene

Database (DGD)[94]. We determined the number of paralogs for each gene based on the group size of the

corresponding gene group and assigned a paralog count of zero for genes absent in the list. (2)

Epigenomics: We obtained ChIP-seq broad peaks for H2A.Z, H3K4me3, H3K9ac, and H3K27me3 in

promoter regions across cell types, as well as promoter-enhancer interactions predicted by EpiTensor[95]

from a previous study[21]. We calculated the average length of the corresponding ChIP-seq peak in a gene's

promoter region across all cell types to define input features for these epigenomic marks. Additionally,

we determined the number of enhancers associated with a gene by averaging the number of predicted

promoter-enhancer interactions across cell types. (3) Evolutionary constraint: We extracted lof_z[25],

mis_z[25], pLI[27], LOEUF[28], and Mis. OEUF[28] from the gnomAD database[28] v2.1.1. We retrieved SNP-level

PhastCons100way vertebrate and PhastCons30way mammalian scores[96] from the dbNSFP database[97] v4,

and averaged these two metrics across the entire span of each gene to obtain gene-level scores. We

sourced promoter and exonic PhastCons100way vertebrate scores from a previous study[66] and obtained

sHet values from the original publication[98]. (4) Functional genomics: We sourced GTEx mean expression[99],

GO terms for embryo development[100], and tissue specificity (tau)[101] values from a recent study[29]. We

retrieved median gene-level transcripts per million (TPMs) across 54 tissues from the v7 release of the

GTEx portal[99]. We counted the number of non-zero expression tissues to determine the number of tissues

expressed for each gene. (5) Deleteriousness upon perturbation: We retrieved SNP-level MetaLR[102], CADD

raw[103], and DANN[104] scores from the dbNSFP database[97], and calculated the average scores across all

coding exons of each gene. We downloaded a list of 2,545 DDG2P genes from the G2P database[56] and

generated a binary feature to indicate whether a gene is included in this list. We sourced EDS values from

the original publication[22], obtained CRISPR HAP1 cell knockout q-values from a previous study[105], and

downloaded SNV data from the ClinVar[42]. We created a binary feature indicating the presence or absence

of pathogenic SNVs in a gene, based on ClinVar SNV data. Additionally, for each gene, we calculated the

percentage of pathogenic SNVs relative to the total number of submitted SNVs in ClinVar. (6) Biological

networks: We downloaded a dataset of 64,006 binary protein-protein interactions involving 9,094

proteins from the Human Reference Protein Interactome Mapping Project[106]. We counted the number of

interaction partners for each gene included in the dataset and assigned a count of zero for genes absent

in the dataset. We obtained PPI degree values from a recent study[29].

We performed mean imputation for missing values and z-standardized all features, excluding binary ones,

based on the means and standard deviations derived from genes not overlapping deletions in the ClinVar

held-out test set.

Model training. We mapped 33 gene-level features to each affected gene within deletions, and padded

deletions with fewer than 100 genes using dummy genes with all zero feature values within the ClinVar

training, validation, and test sets. This process resulted in an input matrix for each deletion with a shape

of 100 x 33. We then trained DosaCNV using the ClinVar training set and tuned hyperparameters via grid

search based on the ClinVar validation set. The search space included: the number of nodes in each hidden

ReLU layer (32, 64, 128, 256); lambda for the L2 penalty (0.006, 0.02, 0.06, 0.2); batch size (16, 32, 64);

and Adam learning rate (3×10^{-5}, 10^{-4}, 3×10^{-4}, 10^{-3}). The optimal hyperparameters based on the lowest

validation loss were 128 and 64 nodes for the first and second hidden ReLU layers, 0.02 for the L2 penalty,

a batch size of 32, and a learning rate of 3×10^{-4}. Additionally, we set the dropout rate to 0.5[88].

For gene-level prediction using DosaCNV-HI, we excluded EDS and DDG2P from the gene-level features to

avoid potential circularity. DosaCNV was trained on the same ClinVar training set using the same set of

optimal hyperparameters from the DosaCNV model for deletion-level prediction.

Deletion-level evaluation. We obtained deletion-level predictions for comparison from five alternative

methods: (1) TADA v1.0.2, available at (https://github.com/jakob-he/TADA), using default settings on the

GRCh37 assembly; (2) StrVCTVRE v1.7, accessed from (https://github.com/andrewSharo/StrVCTVRE),

employing default settings with all analyzed deletions lifted to GRCh38 using the UCSC liftOver tool; (3)

HIS-LOD, in which we retrieved gene-level predictions (HIS) for all genes from the original publication and

aggregated them to deletions-level predictions based on the method described therein[18]; (4) X-CNV,

downloaded from (https://github.com/kbvstmd/XCNV), with default settings on the GRCh37 assembly;

and (5) CADD-SV v1.1 webtool (https://cadd-sv.bihealth.org/score), using default settings on the GRCh37

assembly.

To evaluate performance on the ClinVar held-out test set (n=2,832) and assess the impact of length

distribution differences, we compiled a joint set of 2,491 predictions from all five alternative methods,

denoting it as the unmatched test set. Using MatchIt, we matched pathogenic and benign deletions in the

unmatched test set based on the number of affected protein-coding genes (n=702), deletion length in

base pairs (n=704), and both criteria combined (n=618), retaining only matched pairs. To examine

performance on deletions of varying lengths, we further divided each of the four sets into four length

groups: small (<50 kb), medium (50-100 kb), large (100-1000 kb), and extra-large (>1000 kb). We obtained

AUC values and corresponding standard errors for each method on each set using the R package pROC[107].

We visualized the AUC values for all methods using R package ggplot2[108] and tested the statistical

significance for the differences between AUC values using the DeLong test, via the roc.test function in

pROC. To estimate each method's dependency on length information, we subtracted the AUC value of the

matched test set from that of the unmatched test set, considering both deletion length and the number

of affected genes.

We evaluated the prioritization ability for putative pathogenic deletions by obtaining rare SVs from case-

control studies by Coe et al., 2014[50] (nstd100) with 29,085 ID/DD/ASD cases and 19,584 controls, and

Zarrei et al., 2019[51] (nstd173) with 2,691 NDD cases and 1,769 family member controls. We applied the

same filtering conditions with the ClinVar data, excluding the condition for clinical interpretation. The

resulting nstd100 and nstd173 sets contained 16,528 and 4,500 coding deletions, respectively, with

16,048 and 4,230 deletions having predictions from all methods. Since these datasets lack pathogenicity

labels, we calculated the enrichment of case-associated deletions in each method's top predictions using

odds ratio as the performance measure. For each dataset, we defined the number of top predictions for

evaluation as 5% of the total case count (referred to as the top 5% predictions), assuming that coding

CNVs contribute to 10% of cases, half of which are deletions. This approach resulted in 1,454 and 134 top

predictions for nstd100 and nstd173, respectively. We visualized the odds ratios by ggplot2 and tested

the statistical significance for the differences between odds ratios by transforming them to log odds ratios

and then using the z-test.

To further assess our method, we determined the inclusion percentage of the nstd100 and nstd173

deletions that had over 95% reciprocal overlap with curated pathogenic deletions from ClinVar, within our

method's top predictions. Specifically, we found 164 nstd100 deletions and 13 nstd173 deletions had over

95% reciprocal overlap with pathogenic deletions in the ClinVar held-out test set.

To evaluate performance in a clinical scenario, we acquired clinically curated rare SVs from 140

developmental delay cases reported by Foong et al., 2015[52]. After filtering, 740 coding deletions, including

33 pathogenic ones, remained. We generated predictions for these deletions and visualized them in a

scatter plot using ggplot2. Additionally, 663 of the 740 deletions had predictions from all methods. Based on

the joint set of 663 deletions, we determined the AUC values for our method and the alternative methods

using pROC.

Gene-level evaluation. To evaluate the performance of DosaCNV-HI and 12 alternative methods—

including DeepLOF[29], LOEUF[28], Mis.OEUF[28], RVIS[26], VIRLoF[58], HIS[18], IS[23], GHIS[19], HIPred[20], Episcore[21], EDS[22],

and pHaplo[24]—in predicting HI and likely HI genes, we obtained LOEUF and Mis.OEUF from the gnomAD

database v2.1.1 and gene scores for all other methods from their respective original publications.

We retrieved six HI and likely HI gene sets for evaluation: (1) 340 "gold standard" HI genes with sufficient

evidence for haploinsufficiency (ClinGen HI score 3) from the ClinGen gene dosage sensitivity map[53]; (2)

397 human orthologs of mouse heterozygous knockout lethal genes from the GitHub repository for

gnomAD (https://github.com/macarthur-lab/gnomAD_lof/); (3) 214 high-confidence ASD genes (SFARI

gene score 1) from the SFARI GENE database[55]; (4) 432 DDG2P genes with the monoallelic autosomal

allelic requirement and absent gene product mutation consequence from the G2P database[56]; (5) 72

eyeG2P genes with the same requirements as DDG2P genes from the G2P database; and (6) 82 skinG2P

genes with the same requirements as DDG2P genes from the G2P database. We conducted analyses using

genes with predictions from all methods, resulting in 210, 315, 176, 360, 59, and 76 HI/likely HI genes

remaining within the six gene lists, respectively. All other protein-coding genes were considered as the HS

gene set.

To mitigate the imbalance between the six HI or likely HI gene sets and the HS gene set, and to control

the influence of gene length on performance evaluation, we adopted a matching approach. For each HI

or likely HI gene set, we used MatchIt to pair each HI gene with an HS gene of a similar CDS length to

obtain a matched gene set. Subsequently, we utilized pROC to compute the AUC values and corresponding

standard errors for each method within the matched gene sets. We visualized the AUC values for all

methods using ggplot2 and tested the statistical significance for the differences between AUC values using

the DeLong test.

For the evaluation of challenging HI and HS genes, we aggregated the six gene sets into a single HI and

likely HI gene list (n=988). From genes with predictions across all methods (n=758), we generated a "short,

likely HI" gene set (n=73) by filtering genes with fewer than 10 expected LOF mutations from the gnomAD

database. Using established LOEUF (<0.35 for constrained genes)[28] and DeepLOF (>0.835 for constrained

genes)[29] cutoffs for constrained genes, we created a "low constraint, likely HI" gene set (n=278). Applying

the same cutoffs, we generated a "high constraint, unlikely HI" gene set (n=209) for constrained HS genes

not implicated in diseases across the G2P, Human Phenotype Ontology[59], ClinVar, and OMIM databases[60].

Subsequently, we defined the top 2,402 genes predicted by each method as the top predictions for each

method (based on cutoff: LOEUF <0.35), representing the predicted HI genes. We calculated the inclusion

percentages with corresponding standard errors of the three challenging gene sets in each method's top

predictions as the performance measure and visualized these inclusion percentages using ggplot2.

Feature importance. To elucidate the key features driving DosaCNV-HI gene-level predictions, we

employed the DeepExplainer function from the SHAP v0.41.0 Python library[63]. We generated a

background set, consisting of 31 gene-level features for 1,000 randomly selected HS genes, resulting in a

1,000 x 31 matrix where each row represents a gene-level feature vector for a specific gene. The explainer

was initialized with the DosaCNV-HI model and the defined background set, allowing the calculation of

conditional expectations for each gene-level feature and the corresponding conditional expectation of HI

predictions for the 1,000 randomly sampled HS genes. Subsequently, we generated an input set

containing 31 gene-level features for 988 HI and likely HI genes, which we fed into the explainer to obtain

the SHAP values. The output was a 988 x 31 SHAP value matrix, where each row represents the SHAP

values for corresponding gene-level feature values of a particular gene. We used the default bee swarm

summary plot from the package to visualize the global importance and impact distributions of gene-level

features. The features on the y-axis were ranked in descending order of global importance, which the

package calculated as the mean absolute value for the corresponding column in the output SHAP value

matrix.

We employed the same approach to generate SHAP values for 30 protein-coding genes overlapping

pathogenic microdeletions at 1q21.1 (distal, BP3-BP4), 15q13.3 (BP4-BP5), 17p12 (HNPP), 17q21.31, and

22q13, with coordinates retrieved from the ClinGen database in the GRCh37 assembly. We visualized the

SHAP values and percentiles of DosaCNV-HI predictions for these affected genes as a heatmap using

ggplot2.

Availability of data and materials

The DosaCNV program and the precomputed DosaCNV-HI gene scores can be accessed at

https://github.com/Zhihan-Leo-Liu/DosaCNV. Datasets used in this study, along with the source code for

analysis can be found at https://github.com/Zhihan-Leo-Liu/DosaCNV/analysis/. Additionally, the sources

and download links for all gene-level features are provided in Supplementary Data 1.

Acknowledgments

The authors thank Mingfu Shao and Xinru Zhang for helpful discussions.

Funding

Research reported in this publication was supported by the National Institute of General Medical Sciences

of the National Institutes of Health under Award Number R35GM142560 and by startup funds from

Pennsylvania State University. The content is solely the responsibility of the author and does not

necessarily represent the official views of the National Institutes of Health.

Author information

Authors and Affiliations

Department of Biology, Pennsylvania State University, University Park, PA 16802, USA

Zhihan Liu & Yi-Fei Huang

Huck Institutes of the Life Sciences, Pennsylvania State University, University Park, PA 16802, USA

Zhihan Liu & Yi-Fei Huang

Molecular, Cellular, and Integrative Biosciences Program, Pennsylvania State University, University Park,

PA 16802, USA

Zhihan Liu

Contributions

ZL implemented the DosaCNV program and conducted all analyses. YH conceived and supervised the

project. ZL and YH prepared and reviewed the manuscript.

Corresponding authors

Correspondence to Zhihan Liu or Yi-Fei Huang.

Ethics declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Competing interests

The authors declare no competing interests.

Additional materials

Additional file 1

Additional file 1. docx

Supplementary Figures 1-4 and Supplementary Tables 1-7

Additional file 2

Additional file 2.xls

The sources and download links for all gene-level features used in this study. Referred to as

Supplementary Data 1 in the main text.

References

1. Zhang, F., Gu, W., Hurles, M. E. & Lupski, J. R. Copy Number Variation in Human Health, Disease, and

Evolution. *Annu. Rev. Genomics Hum. Genet.* **10**, 451–481 (2009).

2. Conrad, D. F. *et al.* Origins and functional impact of copy number variation in the human genome.

Nature **464**, 704–712 (2010).

3. Hastings, P. J., Lupski, J. R., Rosenberg, S. M. & Ira, G. Mechanisms of change in gene copy number.

Nat. Rev. Genet. **10**, 551–564 (2009).

4. Liu, P., Carvalho, C. M. B., Hastings, P. J. & Lupski, J. R. Mechanisms for recurrent and complex

human genomic rearrangements. *Curr. Opin. Genet. Dev.* **22**, 211–220 (2012).

5. Perry, G. H. *et al.* Diet and the evolution of human amylase gene copy number variation. *Nat. Genet.*

39, 1256–1260 (2007).

6. Leffler, E. M. *et al.* Resistance to malaria through structural variation of red blood cell invasion

receptors. *Science* **356**, eaam6393 (2017).

7. Malhotra, D. & Sebat, J. CNVs: Harbingers of a Rare Variant Revolution in Psychiatric Genetics. *Cell*

148, 1223–1241 (2012).

8. McCarroll, S. A. *et al.* Deletion polymorphism upstream of IRGM associated with altered IRGM

expression and Crohn's disease. *Nat. Genet.* **40**, 1107–1112 (2008).

9. McCarroll, S. A. *et al.* Donor-recipient mismatch for common gene deletion polymorphisms in graft-

versus-host disease. *Nat. Genet.* **41**, 1341–1344 (2009).

10. Cooper, G. M. *et al.* A copy number variation morbidity map of developmental delay. *Nat. Genet.*

43, 838–846 (2011).

11. Pinto, D. *et al.* Functional impact of global rare copy number variation in autism spectrum disorders.

Nature **466**, 368–372 (2010).

12. McCarthy, S. E. *et al.* Microduplications of 16p11.2 are associated with schizophrenia. *Nat. Genet.*

41, 1223–1227 (2009).

13. Shao, X. *et al.* Copy number variation is highly correlated with differential gene expression: a pan-

cancer study. *BMC Med. Genet.* **20**, 175 (2019).

14. Weischenfeldt, J., Symmons, O., Spitz, F. & Korbel, J. O. Phenotypic impact of genomic structural

variation: insights from and for human disease. *Nat. Rev. Genet.* **14**, 125–138 (2013).

15. Rice, A. M. & McLysaght, A. Dosage sensitivity is a major determinant of human copy number

variant pathogenicity. *Nat. Commun.* **8**, 14366 (2017).

16. Seidman, J. G. & Seidman, C. Transcription factor haploinsufficiency: when half a loaf is not enough.

J. Clin. Invest. **109**, 451–455 (2002).

17. Veitia, R. A. & Potier, M. C. Gene dosage imbalances: action, reaction, and models. *Trends Biochem.*

Sci. **40**, 309–317 (2015).

18. Huang, N., Lee, I., Marcotte, E. M. & Hurles, M. E. Characterising and Predicting Haploinsufficiency in

the Human Genome. *PLOS Genet.* **6**, e1001154 (2010).

19. Steinberg, J., Honti, F., Meader, S. & Webber, C. Haploinsufficiency predictions without study bias.

Nucleic Acids Res. **43**, e101 (2015).

20. Shihab, H. A., Rogers, M. F., Campbell, C. & Gaunt, T. R. HIPred: an integrative approach to

predicting haploinsufficient genes. *Bioinformatics* **33**, 1751–1757 (2017).

21. Han, X. *et al.* Distinct epigenomic patterns are associated with haploinsufficiency and predict risk

genes of developmental disorders. *Nat. Commun.* **9**, 2138 (2018).

22. Wang, X. & Goldstein, D. B. Enhancer Domains Predict Gene Pathogenicity and Inform Gene

Discovery in Complex Disease. *Am. J. Hum. Genet.* **106**, 215–233 (2020).

23. Khurana, E., Fu, Y., Chen, J. & Gerstein, M. Interpretation of Genomic Variants Using a Unified

Biological Network Approach. *PLOS Comput. Biol.* **9**, e1002886 (2013).

24. Collins, R. L. *et al.* A cross-disorder dosage sensitivity map of the human genome. *Cell* **185**, 3041-

3055.e25 (2022).

25. Samocha, K. E. *et al.* A framework for the interpretation of de novo mutation in human disease. *Nat.*

Genet. **46**, 944–950 (2014).

26. Petrovski, S., Wang, Q., Heinzen, E. L., Allen, A. S. & Goldstein, D. B. Genic Intolerance to Functional

Variation and the Interpretation of Personal Genomes. *PLoS Genet.* **9**, e1003709 (2013).

27. Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**, 285–291

(2016).

28. Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in 141,456

humans. *Nature* **581**, 434–443 (2020).

29. LaPolice, T. M. & Huang, Y.-F. *A deep learning framework for predicting human essential genes from*

population and functional genomic data. http://biorxiv.org/lookup/doi/10.1101/2021.12.21.473690

(2021) doi:10.1101/2021.12.21.473690.

30. Ruderfer, D. M. *et al.* Patterns of genic intolerance of rare copy number variation in 59,898 human

exomes. *Nat. Genet.* **48**, 1107–1111 (2016).

31. Simons, Y. B., Bullaughey, K., Hudson, R. R. & Sella, G. A population genetic interpretation of GWAS

findings for human quantitative traits. *PLOS Biol.* **16**, e2002985 (2018).

32. Fuller, Z. L., Berg, J. J., Mostafavi, H., Sella, G. & Przeworski, M. Measuring intolerance to mutation in

human genetics. *Nat. Genet.* **51**, 772–776 (2019).

33. Wooster, R. *et al.* Identification of the breast cancer susceptibility gene BRCA2. *Nature* **378**, 789–792

(1995).

34. Roy, R., Chun, J. & Powell, S. N. BRCA1 and BRCA2: different roles in a common pathway of genome

protection. *Nat. Rev. Cancer* **12**, 68–78 (2012).

35. Raybould, M. C., Birley, A. J. & Hultén, M. Molecular variation of the human elastin (ELN) gene in a

normal human population. *Ann. Hum. Genet.* **59**, 149–161 (1995).

36. Lu, S. *et al.* Atm-haploinsufficiency enhances susceptibility to carcinogen-induced mammary tumors.

Carcinogenesis **27**, 848–855 (2006).

37. Ganel, L., Abel, H. J., FinMetSeq Consortium & Hall, I. M. SVScore: an impact prediction tool for

structural variation. *Bioinformatics* **33**, 1083–1085 (2017).

38. Geoffroy, V. *et al.* AnnotSV: an integrated tool for structural variations annotation. *Bioinformatics*

34, 3572–3574 (2018).

39. Riggs, E. R. *et al.* Technical standards for the interpretation and reporting of constitutional copy

number variants: a joint consensus recommendation of the American College of Medical Genetics

and Genomics (ACMG) and the Clinical Genome Resource (ClinGen). *Genet. Med. Off. J. Am. Coll.*

Med. Genet. **22**, 245–257 (2020).

40. Amores, J. Multiple instance classification: Review, taxonomy and comparative study. *Artif. Intell.*

201, 81–105 (2013).

41. Pearl, J. *Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference*. (Morgan

Kaufmann, 1988).

42. Landrum, M. J. *et al.* ClinVar: improving access to variant interpretations and supporting evidence.

Nucleic Acids Res. **46**, D1062–D1067 (2018).

43. Skelly, A. C., Dettori, J. R. & Brodt, E. D. Assessing bias: the importance of considering confounding.

Evid.-Based Spine-Care J. **3**, 9–12 (2012).

44. Hertzberg, J., Mundlos, S., Vingron, M. & Gallone, G. TADA—a machine learning tool for functional

annotation-based prioritisation of pathogenic CNVs. *Genome Biol.* **23**, 67 (2022).

45. Sharo, A. G., Hu, Z., Sunyaev, S. R. & Brenner, S. E. StrVCTVRE: A supervised learning method to

predict the pathogenicity of human genome structural variants. *Am. J. Hum. Genet.* **109**, 195–209

(2022).

46. Zhang, L. *et al.* X-CNV: genome-wide prediction of the pathogenicity of copy number variations.

Genome Med. **13**, 132 (2021).

47. Kleinert, P. & Kircher, M. A framework to score the effects of structural variants in health and

disease. *Genome Res.* gr.275995.121 (2022) doi:10.1101/gr.275995.121.

48. Girirajan, S. & Eichler, E. E. Phenotypic variability and genetic susceptibility to genomic disorders.

Hum. Mol. Genet. **19**, R176–R187 (2010).

49. Miller, D. T. *et al.* Consensus Statement: Chromosomal Microarray Is a First-Tier Clinical Diagnostic

Test for Individuals with Developmental Disabilities or Congenital Anomalies. *Am. J. Hum. Genet.* **86**,

749–764 (2010).

50. Coe, B. P. *et al.* Refining analyses of copy number variation identifies specific genes associated with

developmental delay. *Nat. Genet.* **46**, 1063–1071 (2014).

51. Zarrei, M. *et al.* A large data resource of genomic copy number variation across

neurodevelopmental disorders. *NPJ Genomic Med.* **4**, 26 (2019).

52. Foong, J., Girdea, M., Stavropoulos, J. & Brudno, M. Prioritizing Clinically Relevant Copy Number

Variation from Genetic Interactions and Gene Function Data. *PLOS ONE* **10**, e0139656 (2015).

53. Riggs, E. R. *et al.* Copy number variant discrepancy resolution using the ClinGen dosage sensitivity

map results in updated clinical interpretations in ClinVar. *Hum. Mutat.* **39**, 1650–1659 (2018).

54. Bult, C. J., Blake, J. A., Smith, C. L., Kadin, J. A. & Richardson, J. E. Mouse Genome Database (MGD)

2019. *Nucleic Acids Res.* **47**, D801–D806 (2019).

55. Abrahams, B. S. *et al.* SFARI Gene 2.0: a community-driven knowledgebase for the autism spectrum

disorders (ASDs). *Mol. Autism* **4**, 36 (2013).

56. Thormann, A. *et al.* Flexible and scalable diagnostic filtering of genomic variants using G2P with

Ensembl VEP. *Nat. Commun.* **10**, 2373 (2019).

57. Ho, D., Imai, K., King, G. & Stuart, E. A. MatchIt: Nonparametric Preprocessing for Parametric Causal

Inference. *J. Stat. Softw.* **42**, 1–28 (2011).

58. Abramovs, N., Brass, A. & Tassabehji, M. GeVIR is a continuous gene-level metric that uses variant

distribution patterns to prioritize disease candidate genes. *Nat. Genet.* **52**, 35–39 (2020).

59. Köhler, S. *et al.* The Human Phenotype Ontology in 2021. *Nucleic Acids Res.* **49**, D1207–D1217

(2020).

60. Amberger, J., Bocchini, C. A., Scott, A. F. & Hamosh, A. McKusick's Online Mendelian Inheritance in

Man (OMIM®). *Nucleic Acids Res.* **37**, D793–D796 (2009).

61. Lopes, I., Altab, G., Raina, P. & de Magalhães, J. P. Gene Size Matters: An Analysis of Gene Length in

the Human Genome. *Front. Genet.* **12**, (2021).

62. Zug, R. Developmental disorders caused by haploinsufficiency of transcriptional regulators: a

perspective based on cell fate determination. *Biol. Open* **11**, bio058896 (2022).

63. Lundberg, S. & Lee, S.-I. A Unified Approach to Interpreting Model Predictions. Preprint at

https://doi.org/10.48550/arXiv.1705.07874 (2017).

64. Yong Ryu, J., Uk Kim, H. & Yup Lee, S. Human genes with a greater number of transcript variants

tend to show biological features of housekeeping and essential genes. *Mol. Biosyst.* **11**, 2798–2807

(2015).

65. Ku, M. *et al.* H2A.Z landscapes and dual modifications in pluripotent and multipotent stem cells

underlie complex genome regulatory functions. *Genome Biol.* **13**, R85 (2012).

66. Boukas, L., Bjornsson, H. T. & Hansen, K. D. Promoter CpG Density Predicts Downstream Gene Loss-

of-Function Intolerance. *Am. J. Hum. Genet.* **107**, 487–498 (2020).

67. Rehm, H. L. *et al.* ClinGen — The Clinical Genome Resource. *N. Engl. J. Med.* **372**, 2235–2242 (2015).

68. Harvard, C. *et al.* Understanding the impact of 1q21.1 copy number variant. *Orphanet J. Rare Dis.* **6**,

54 (2011).

69. Prokudin, I. *et al.* Exome sequencing in developmental eye disease leads to identification of causal

variants in GJA8, CRYGC, PAX6 and CYP1B1. *Eur. J. Hum. Genet.* **22**, 907–915 (2014).

70. Hoppman-Chaney, N., Wain, K., Seger, P., Superneau, D. & Hodge, J. Identification of single gene

deletions at 15q13.3: further evidence that CHRNA7 causes the 15q13.3 microdeletion syndrome

phenotype. *Clin. Genet.* **83**, 345–351 (2013).

71. Gillentine, M. & Schaaf, C. P. The Human Clinical Phenotypes of Altered CHRNA7 Copy Number.

Biochem. Pharmacol. **97**, 352–362 (2015).

72. Li, J., Parker, B., Martyn, C., Natarajan, C. & Guo, J. The PMP22 Gene and Its Related Diseases. *Mol.*

Neurobiol. **47**, 673–698 (2013).

73. Wray, C. D. 17q21.31 Microdeletion associated with infantile spasms. *Eur. J. Med. Genet.* **56**, 59–61

(2013).

74. Koolen, D. A. *et al.* Mutations in the chromatin modifier gene KANSL1 cause the 17q21.31

microdeletion syndrome. *Nat. Genet.* **44**, 639–641 (2012).

75. Bozdagi, O. *et al.* Haploinsufficiency of the autism-associated Shank3 gene leads to deficits in

synaptic function, social interaction, and social communication. *Mol. Autism* **1**, 15 (2010).

76. Whalen, S., Schreiber, J., Noble, W. S. & Pollard, K. S. Navigating the pitfalls of applying machine

learning in genomics. *Nat. Rev. Genet.* **23**, 169–181 (2022).

77. Creyghton, M. P. *et al.* H2AZ Is Enriched at Polycomb Complex Target Genes in ES Cells and Is

Necessary for Lineage Commitment. *Cell* **135**, 649–661 (2008).

78. Yang, X. *et al.* Silencing of developmental genes by H3K27me3 and DNA methylation reflects the

discrepant plasticity of embryonic and extraembryonic lineages. *Cell Res.* **28**, 593–596 (2018).

79. Frankel, N. *et al.* Phenotypic robustness conferred by apparently redundant transcriptional

enhancers. *Nature* **466**, 490–493 (2010).

80. Norris, M., Lovell, S. & Delneri, D. Characterization and Prediction of Haploinsufficiency Using

Systems-Level Gene Properties in Yeast. *G3 GenesGenomesGenetics* **3**, 1965–1977 (2013).

81. Johnson, A. F., Nguyen, H. T. & Veitia, R. A. Causes and effects of haploinsufficiency. *Biol. Rev.* **94**,

1774–1785 (2019).

82. Spielmann, M., Lupiáñez, D. G. & Mundlos, S. Structural variation in the 3D genome. *Nat. Rev.*

Genet. **19**, 453–467 (2018).

83. Bhartiya, D. & Scaria, V. Genomic variations in non-coding RNAs: Structure, function and regulation.

Genomics **107**, 59–68 (2016).

84. Carvalho, C. M. B. *et al.* Dosage Changes of a Segment at 17p13.1 Lead to Intellectual Disability and

Microcephaly as a Result of Complex Genetic Interaction of Multiple Genes. *Am. J. Hum. Genet.* **95**,

565–578 (2014).

85. Girirajan, S. *et al.* Phenotypic Heterogeneity of Genomic Disorders and Rare Copy-Number Variants.

N. Engl. J. Med. **367**, 1321–1331 (2012).

86. Davies, R. W. *et al.* Using common genetic variation to examine phenotypic expression and risk

prediction in 22q11.2 deletion syndrome. *Nat. Med.* **26**, 1912–1918 (2020).

87. Nair, V. & Hinton, G. E. Rectified Linear Units Improve Restricted Boltzmann Machines.

88. Srivastava, N., Hinton, G., Krizhevsky, A., Sutskever, I. & Salakhutdinov, R. Dropout: A Simple Way to

Prevent Neural Networks from Overfitting.

89. He, K., Zhang, X., Ren, S. & Sun, J. Delving Deep into Rectifiers: Surpassing Human-Level

Performance on ImageNet Classification. Preprint at http://arxiv.org/abs/1502.01852 (2015).

90. Kingma, D. P. & Ba, J. Adam: A Method for Stochastic Optimization. Preprint at

http://arxiv.org/abs/1412.6980 (2017).

91.   Frankish, A. *et al.* GENCODE reference annotation for the human and mouse genomes. *Nucleic Acids*

Res. **47**, D766–D773 (2019).

92.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.

Bioinformatics **26**, 841–842 (2010).

93.   Durinck, S., Spellman, P. T., Birney, E. & Huber, W. Mapping identifiers for the integration of

genomic datasets with the R/Bioconductor package biomaRt. *Nat. Protoc.* **4**, 1184–1191 (2009).

94.   Ouedraogo, M. *et al.* The Duplicated Genes Database: Identification and Functional Annotation of

Co-Localised Duplicated Genes across Genomes. *PLoS ONE* **7**, e50653 (2012).

95.   Zhu, Y. *et al.* Constructing 3D interaction maps from 1D epigenomes. *Nat. Commun.* **7**, 10812 (2016).

96.   Siepel, A. *et al.* Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes.

Genome Res. **15**, 1034–1050 (2005).

97.   Liu, X., Li, C., Mou, C., Dong, Y. & Tu, Y. dbNSFP v4: a comprehensive database of transcript-specific

functional predictions and annotations for human nonsynonymous and splice-site SNVs. *Genome*

Med. **12**, 103 (2020).

98.   Cassa, C. A. *et al.* Estimating the Selective Effects of Heterozygous Protein Truncating Variants from

Human Exome Data. *Nat. Genet.* **49**, 806–810 (2017).

99.   The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science* **369**, 1318–

1330 (2020).

100.   The Gene Ontology Resource: 20 years and still GOing strong. *Nucleic Acids Res.* **47**, D330–D338

(2019).

101.   Yanai, I. *et al.* Genome-wide midrange transcription profiles reveal expression level relationships

in human tissue specification. *Bioinformatics* **21**, 650–659 (2005).

102.   Dong, C. *et al.* Comparison and integration of deleteriousness prediction methods for

nonsynonymous SNVs in whole exome sequencing studies. *Hum. Mol. Genet.* **24**, 2125–2137 (2015).

103.   Kircher, M. *et al.* A general framework for estimating the relative pathogenicity of human

genetic variants. *Nat. Genet.* **46**, 310–315 (2014).

104.   Quang, D., Chen, Y. & Xie, X. DANN: a deep learning approach for annotating the pathogenicity

of genetic variants. *Bioinformatics* **31**, 761–763 (2015).

105.   Blomen, V. A. *et al.* Gene essentiality and synthetic lethality in haploid human cells. *Science* **350**,

1092–1096 (2015).

106.   Luck, K. *et al.* A reference map of the human binary protein interactome. *Nature* **580**, 402–408

(2020).

107.   Robin, X. *et al.* pROC: an open-source package for R and S+ to analyze and compare ROC curves.

BMC Bioinformatics **12**, 77 (2011).

108.   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer, 2009). doi:10.1007/978-0-

387-98141-3.

Figures

Figure 1. Schematic representation of the DosaCNV model. DosaCNV utilizes a fully connected architecture in conjunction with a noisy-or pooling layer to jointly infer gene haploinsufficiency and coding deletion pathogenicity. Deletions are first annotated with gene-level features of affected genes as input. For a given deletion, the fully connected architecture processes its gene-level input features and predicts the HI probability (P_{HI}) for each affected gene. Subsequently, the noisy-or pooling layer integrates these gene-level predictions, assuming independent gene effects, to compute the probability of the deletion harboring at least one HI gene, which serves as a proxy for deletion-level pathogenicity. The right side of the panel illustrates an example deletion input, along with its intermediate gene-level predictions and the noisy-or pooling calculation.

Figure 2. Performance comparison of DosaCNV and alternative methods in predicting pathogenicity of deletions from the unmatched and matched ClinVar held-out test sets. For each set, comparisons are presented for the entire set and four deletion subgroups based on length: small (<50 kb), medium (50-100 kb), large (100-1000 kb), and extra-large (>1000 kb). (A)(B) Unmatched test set (n=2,491). (C)(D) Test set matched by both the number of affected genes and deletion length (n=618). Vertical lines denote ± one standard error. Horizontal dashed lines indicate an AUC of 0.5.

Figure 3. Comparative assessment of DosaCNV and alternative methods for prioritizing potential pathogenic deletions in case-control and curated case-only datasets. (A)(B) Evaluation of prioritization performance in case-control nstd100 (16,048 deletions; 29,085 ID/DD/ASD cases and 19,584 population controls) and nstd173 (4,230 deletions; 2,691 NDD cases and 1,769 family member controls) datasets. The enrichment of case-associated deletions within the top 5% predictions (1,454 for nstd100 and 134 for nstd173) for each method was used as the performance metric. Error bars represent one standard error. (C) DosaCNV scores for 740 clinically curated deletions in a case-only dataset from Foong et al., 2015. A DosaCNV score threshold of 0.8 (indicated by the horizontal dashed line) yields a true positive rate of 0.970 and a false positive rate of 0.028.

Figure 4. Predictive power of DosaCNV-HI and 12 alternative methods in discriminating HI and likely HI genes from HS genes. AUC values for DosaCNV-HI and alternatives are shown in predicting (A) ClinGen HI genes (n=210/340; 210 out of 340 genes had predictions from all methods), (B) human orthologs of mouse heterozygous knock-out lethal genes (n=315/397), (C) high confidence SFARI ASD genes (n=176/214), (D) DDG2P genes sensitive to heterozygous LOF mutations (n=366/432), (E) eyeG2P genes sensitive to heterozygous LOF mutations (n=59/72), (F) skinG2P genes sensitive to heterozygous LOF mutations (n=76/82). Horizontal black lines denote ± one standard error. Inclusion percentages for (G) "short, likely HI", (H) "low constraint, likely HI", and (I) "high constraint, unlikely HI" gene sets within the top 2,402 genes ranked by each method; a higher percentage for "likely HI" sets or a lower percentage for "unlikely HI" sets suggests superior performance. Error bars depict one standard error.

Figure 5. Elucidating gene-level feature contributions to haploinsufficiency predictions via SHAP values. (A) The SHAP summary plot is presented with gene-level features on the y-axis, arranged by global importance (mean absolute SHAP values) in descending order. Each dot represents a specific feature value for a gene within the HI or likely HI gene set (n=988), and the color denotes the actual feature value (red: high; blue: low). The x-axis displays SHAP values, which indicate the impact of each feature on the deviation between a gene's DosaCNV-HI prediction and the average DosaCNV-HI prediction for background HS genes (n=1000). Positive SHAP values enhance the DosaCNV-HI prediction (genes more likely to be HI), whereas negative values diminish it (genes less likely to be HI). The sum of SHAP values across all features for each gene equals the difference between its predicted HI probability and the expected HI probability of a designated background gene set (HS genes in our case). (B) DosaCNV-HI percentiles are shown for genes overlapping five pathogenic microdeletions, along with their corresponding SHAP values. A gene with a lower DosaCNV-HI percentile is more likely to be HI.