An amphipathic helix facilitates direct membrane binding of Mycoplasma FtsZ

Soumyajit Dutta[1], Sakshi Poddar[2,3], Joyeeta Chakraborty[1], Ramanujam Srinivasan[2,3],

Pananghat Gayathri[1#]

[1] Biology Division, Indian Institute of Science Education and Research, Pune – 411008, India

[2] School of Biological Sciences, National Institute of Science Education and Research,

Bhubaneswar – 752050, India

[3] Homi Bhabha National Institutes (HBNI), Training School Complex, Anushakti Nagar,

Mumbai,

India 400094

Running title: Membrane Binding Amphipathic helix in Mycoplasma FtsZs

Keywords: Bacteria, cell division, mycoplasma, FtsZ, amphipathic helix

#Address correspondence to Pananghat Gayathri, gayathri@iiserpune.ac.in

Abstract

Cell division in bacteria is initiated by constriction of the Z-ring comprising two essential proteins FtsZ and FtsA. Despite our knowledge about the crucial function of the Z-ring in bacterial division, the precise roles and mechanism of how FtsZ and FtsA drive cell constriction remain elusive. FtsZ/FtsA in wall-less bacteria like mycoplasmas is an ideal model system for obtaining mechanistic insights into Z-ring constriction in the absence of cell wall machinery. In this study, we have analyzed FtsZ and FtsA sequences of 113 mycoplasma species and compared with the corresponding protein sequences in cell-walled bacteria. We report a phylogenetically distinct group of 12 species that possess FtsZs without the canonical FtsA interacting conserved C-terminal peptide (CCTP) motif. Interestingly, these FtsZs contain a putative membrane-binding amphipathic helix as an N-terminal or C-terminal extension to the globular FtsZ domain. As a proof-of-concept, we experimentally show that the proposed C-terminal amphipathic helix in *M. genitalium* FtsZ binds liposomes *in vitro* as well as localizes to *E. coli* membrane *in vivo*. Additionally, we identify a putative cholesterol recognition motif within the C-terminal amphipathic helix region of *M. genitalium* FtsZ. Our study catalogues the functional variations of membrane attachment by the FtsZ and FtsA system in cell wall-less mycoplasmas and provides a new perspective to study novel functions of FtsZ/A system in cell division.

Importance

Z-ring and peptidoglycan synthesis machinery both play crucial roles in bacterial cell division. Currently, our knowledge about how FtsZ and FtsA, the two primary components of the Z-ring, function, is limited to cell-walled bacteria where ring constriction is coupled to peptidoglycan synthesis. Cell wall-less bacterial FtsZ/A system is an excellent model to study

the mechanism of Z-ring constriction in the absence of cell wall synthesis machinery. Here,

we analysed FtsZ protein sequences across mycoplasma species and identified their

characteristic sequence features. Our study reveals a novel group of FtsZs from mycoplasma

with an inherent membrane binding and probable cholesterol sensing amphipathic motif,

which serves as a new paradigm to explore fundamental roles of FtsZ and FtsA in Z-ring

constriction during bacterial division.

Introduction

FtsZ and FtsA are the two major proteins involved in bacterial cell division (1, 2). The tubulin-like protein FtsZ forms short patches of overlapping filaments called the Z-ring which treadmills and gradually leads to constriction of the cell (3, 4). The presence of FtsZ is essential to initiate constriction but seems to be dispensable at the later stages of bacterial cell division (5, 6). Treadmilling FtsZ filaments act as a guide for incorporating peptidoglycan strands in the newly forming septum during constriction (3, 4, 7, 8). In most bacteria, the actin-homolog FtsA, with the help of its C-terminal amphipathic helix, functions as the primary membrane anchor for FtsZ (9, 10). FtsZ and FtsA interact with other cell division-related proteins to further regulate the downstream events in the constriction process (11–15). Other than FtsA, there are proteins such as ZipA (in Gram-negative gamma proteobacteria) and SepF (in Gram-positive bacteria and cyanobacteria) that act as membrane anchors for FtsZ (16, 17).

The domain organization of FtsZ, a tubulin-like protein, contains a GTP binding globular core, an N-terminal extension, a C-terminal disordered linker, and a conserved C-terminal peptide (CCTP) (Fig.1b) (18). The globular core domain contains two highly conserved tubulin signature motifs, (1) GTP binding motif- [S/A/G]–G–G–T–G–[S/A/T]–G, and (2) GTP hydrolysis motif- N–X–D–X–X–D, (where X can be any residue) (19, 20). The CCTP motif also shows a high degree of conservation across different bacterial species with a consensus sequence of D/E–I/V–P–X–F/Y–L–R/K, (where X can be any residue) (21, 22). The CCTP is required for interaction between FtsZ and many of its interacting membrane tethering proteins like FtsA, and the less conserved ZipA or SepF (23–25).

FtsA acts as the primary membrane anchor for FtsZ and it is widely conserved in most bacterial species. FtsA is an actin family protein containing a globular core domain followed

by a disordered linker and C-terminal amphipathic helix (CTAH) (Fig. 1b). Membrane

binding ability of the CTAH of FtsA helps FtsZ filaments assemble at the inner cytoplasmic

membrane and initiate divisome formation (9, 26). However, the precise role of this essential

protein in the divisome assembly is not clear yet. Although protofilament architecture of FtsA

has been captured *in vitro* for *Thermotoga maritima* (27), *Streptococcus pneumoniae* (28,

29), *Escherichia coli* (30, 31), and *Vibrio cholerae* (32), the ATPase activity of *T. maritima*

or *S. pneumoniae* has not been observed despite the presence of ATPase domain. Rapid ATP

hydrolysis for *E. coli* FtsA has been observed in presence of phospholipids (33) while *in vitro*

Z-ring dynamics does not require ATPase activity of FtsA (34). Thus, while a passive

membrane anchoring role is important, it is not yet clear if FtsA plays additional active role

through ATP hydrolysis.

Cell division is a complex process that relies on coordinated and collective action of a

number of proteins involved in Z-ring assembly, cell wall synthesis and their regulations

(35). Overall, in bacteria, there are at least twenty different proteins that play an important

role in the formation of divisome, a multiprotein complex that coordinates different steps of

the division in a highly regulated and organized manner (36). Notably, very often in bacteria,

many of the division-related proteins are transcribed from a single operon called division and

cell wall (*dcw*) cluster (37). The *dcw* operon in cell walled model organisms *E. coli* and

Bacillus subtilis comprises sixteen and seventeen genes, respectively, which include FtsZ,

FtsA and their downstream interactor proteins involved in cell wall synthesis (Fig. 1a). While

much of the information regarding the function of cell division proteins has been deciphered

based on studies carried out in walled bacterial species, our understanding of cell wall-less

systems is limited. In cell wall-less bacteria, genes related to peptidoglycan synthesis are lost

and a concomitant shorter *dcw* operon is the result of an extensive genome reduction (38). In

mycoplasmas, for example, only five genes namely *mraZ, mraW (rsmH), ftsA, ftsZ* and *sepF*

can be found in the *dcw* operon (39, 40). The highly conserved nature of these genes across the bacterial kingdom implies their important function in cell division in a cell wall less milieu. In cell-walled bacteria, cell wall synthesis by the peptidoglycan synthesis machinery is supposedly the major player for constriction force generation as the inward force generated by the Z-ring is not sufficient to overcome the turgor pressure (7). However, how constriction occurs in cell wall-less bacteria like mycoplasmas, where FtsZ and FtsA are possibly the only two proteins to constitute the Z-ring in the absence of the cell wall synthesis machinery, is an enigma.

Earlier studies on *Mycoplasma genitalium* suggest that the cell is capable of division in the absence of FtsZ without affecting its doubling rate (41). In *M. pneumoniae*, a close relative of *M. genitalium*, motility-driven mechanical force appears to facilitate constriction, a mechanism alternatively termed as cytofission (42). Other cell wall-less strains of bacteria including the L-form have been found to utilize processes such as membrane blebbing, budding, or extrusion-resolution for multiplying (43, 44). Thus, if FtsZ is not essential, why it remains conserved in genome of cell wall-less bacteria, remains unexplained. However, in *M. genitalium,* FtsZ has been found to localize at the mid cell at the onset of constriction indicating its role in cell division (45). Interestingly, a recent discovery on mycoplasma JCVI-syn3a, the minimal cell with synthetic genome, shows the requirement *ftsZ* and *sepF* for normal cell morphology and division (40). This essentially makes all five *dcw* cluster genes namely *mraZ*, *mraW*, *ftsA*, *ftsZ*, and *sepF* an integral part of the minimal gene set in JCVI-syn3a. These findings prompted us to look into characteristic sequence features of these cell division proteins across mycoplasma species to understand their role in cell wall-less bacteria.

Here we report the sequence features of the primary contractile ring forming protein FtsZ across 113 mycoplasma species. We identified a phylogenetically distinct FtsZ cluster

comprising 12 mycoplasmas having an amphipathic helix with potential membrane binding

characteristics. *M. genitalium*, the bacterium with a minimal genome, is one of the member

species of this clade, which interestingly, possesses a C-terminal amphipathic helix (CTAH).

CTAH will possibly allow a direct membrane attachment for FtsZ bypassing the requirement

of FtsA as membrane anchor. Furthermore, we have experimentally shown that the *M.*

genitalium CTAH motif binds to liposomes *in vitro* as well as localizes to *E. coli* membrane

in vivo. We also identified a cholesterol recognition amino acid consensus (CRAC) sequence

in the C-terminal of *M. genitalium* and *M. pneumoniae* FtsZ. Overall, our identification of

unique membrane binding and cholesterol interacting properties of bacterial FtsZ provide us

with examples of naturally existing minimal divisome components, a powerful handle to

obtain new insights to the fundamental mechanism of cell division.

Results

All mycoplasma species do not possess *ftsZ* and *ftsA* genes

We analysed 113 mycoplasma genomes in NCBI database for division and cell wall (*dcw*)

operon to check the presence of division related genes (Supplementary data sheet). While

searching the NCBI genome database, we found *ftsA* as an unannotated gene present in

between *mraW* and *ftsZ* in most mycoplasmas. After checking the domain organization of its

protein coding sequence, we confirmed this as FtsA (Fig. S1). Importantly, in mycoplasma,

the four division related genes namely *mraZ*, *mraW*, *ftsA*, and *ftsZ* were present in the same

sequential manner as found in cell walled bacteria. As expected, all additional genes related

to the cell wall synthesis present in the *dcw* operon in cell-walled bacteria were absent. We

observed that even FtsZ and FtsA were not conserved across all mycoplasma species.

Accordingly, based on the presence of FtsA and FtsZ, we divided them into four groups (Fig.

1c, d). Based on the comparison with the 16S rRNA based phylogenetic tree, we observed

that the presence of FtsZ/FtsA was correlated to the species organization, except for a few

species that were outliers (highlighted by '*'; Fig. 1c, d).

Majority of them i.e., 82 out of 113 species, were classified into group 1 where both FtsZ and

FtsA are present (Fig. 1d). Group 2, having only FtsZ but no FtsA, includes 10 species while

group 3 has 7 species where FtsA is present but FtsZ is absent. Finally, the 14 species of

group 4 lack both FtsZ and FtsA. Further, we analysed the sequences systematically to find

out the characteristic features of these two proteins in the various groups and to further

correlate with their role in cell division.

FtsZ sequence analysis reveals variable C-terminal sequences among mycoplasma

species

To compare the FtsZ proteins among group 1 mycoplasma species, we checked for the

conservation of three characteristic signature motifs of FtsZ: the GTP binding motif, GTPase

motif and CCTP (18, 27). We found the GTP binding and hydrolysis motifs were well

conserved across species whereas the CCTP was variable (Fig. 2a). We divided the sequences

into five subgroups based on the signature amino acid sequences of FtsZ CCTP (Fig 2a, S2).

The cladogram constructed from the FtsZ sequences interestingly matched with the CCTP

based classification (Fig. 2b). The CCTP based groups of FtsZ also forms clusters in the 16S

rRNA based cladogram (Fig. S3. 14 organisms out of 82 possessed a canonical FtsZ CCTP

and were classified into group 1A. Sequences that showed conservation of charges similar to

canonical CCTP residues, but missed the conserved proline, were classified into group 1B.

Group 1C sequences were more divergent where only proline and phenylalanine were

conserved as found in cell walled bacterial CCTPs while the negatively charged stretch of

residues were not conserved. The rest of the species did not show any conservation in their

CCTP and while comparing with the cladogram we identified two such clusters (called 1D

and 1E) separated by group 1C. Multiple sequence alignment of the C-terminal FtsZs of 1D

and 1E groups showed conservation of negative charge and some hydrophobic residues (Fig.

S2), though not belonging to the canonical motif signature. Importantly, the invariant proline

of CCTP is missing in groups 1B, 1D and 1E. We also report an extended N-terminal region

(i.e. more than 15 amino residues compared to the core globular domain of cell walled

bacteria FtsZs) in groups D and E (except *M. genitalium* and *M. pneumoniae*) FtsZ sequences

(Fig. S4, Table S1).

Mycoplasma FtsZs of Group 1E possess an amphipathic helix

One of the interesting findings of our analysis is the presence of a potential amphipathic helix

(AH) in each of the twelve FtsZ sequences of group 1E, where CCTP motifs for FtsA

interaction are probably absent. Ten FtsZ sequences were identified to possess AH at the N-

terminal and the other two (*M. genitalium* and *M. pneumoniae*) at the C terminal (Fig. S5).

The helical wheel diagram highlights the amphipathic nature of the proposed N-terminal and

C-terminal helical regions (Fig. 3a, Table S2). We also calculated two parameters related to

amphipathic helix, namely, hydrophobicity and hydrophobic moments for all the proposed

AHs and compared with known membrane binding AHs from FtsA, SepF and MinD from

cell walled bacteria. Importantly, the amphipathic parameters for all the twelve proposed

mycoplasma AHs fall in the same region indicating their probable amphipathic membrane

binding nature (Fig. 3b).

Membrane binding features of Mycoplasma FtsAs and SepFs show varied combinations of membrane attachment

The lack of canonical CCTP motif and the presence of an inherent amphipathic helix in FtsZs prompted us to check whether an amphipathic helix is present in the corresponding group 1E FtsAs. The amphipathic nature of the FtsA C-terminal helix was checked by plotting hydrophobic moment and hydrophobicity and compared with known membrane binding amphipathic helix. We found that all FtsA sequences from 1E group (except *M. iowae*) possess the canonical C-terminal amphipathic helix (Fig. S6a, b). Moreover, when we checked FtsA sequences from other groups, we found organisms from group 1A and group 1B without the C-terminal amphipathic helix which interestingly form separate clusters in the phylogenetic tree (Fig. S6a). We next checked for the presence of the alternative membrane attachment adaptor SepF and found that it is present in the genome of only six mycoplasma species of group 1A and its locus is next to the *ftsZ* gene of the *dcw* cluster (Fig. S6c). Interestingly, five out of six SepF containing species lack the C-terminal amphipathic helix in the FtsA (Fig. S6d). We also checked for potential membrane binding site at the N-terminal end of the six mycoplasma SepF sequences by helical wheel diagram. Though, no specific hydrophobic face or amphipathic helix was identified similar to that of *B. subtilis* SepF (Fig. S7 a, b), there are conserved multiple hydrophobic amino acids (e.g. phenylalanine, tryptophan etc.) which might be responsible here for hydrophobic loop mediated membrane binding (Fig. S7b). Thus, FtsZs in mycoplasma groups can anchor to the membrane in a variety of modes starting with its inherent amphipathic helix, or through FtsA amphipathic helix/hydrophobic loop as well as through SepF where FtsA AH is absent.

M. genitalium **FtsZ CTAH binds liposomes and localises to membranes in** *E. coli*

To experimentally verify the membrane binding ability of the proposed AHs, we chose *M.*

genitalium sequence as a representative FtsZ with amphipathic helix at the C-terminal end.

The proposed amphipathic helix from *M. genitalium* FtsZ i.e. the C-terminal 23 amino acid

stretch (347-366) was inserted at the C-terminal end of *Spiroplasma melliferum* (Sm) FtsZ

protein to obtain a chimeric construct (named as SmFtsZ^{MgCTAH}). We used liposomes made

with polar lipids (in molar ratio of 25% DOPG, 20% DOPC, 30% cardiolipin, and 25%

spingomyelin) according to their relative abundance in mycoplasma membrane (46).

Liposome sedimentation assay confirmed that SmFtsZ^{MgCTAH} binds liposomes as we could

see the presence of protein in the pellet fraction which increases with liposome concentration

(Fig. 3c, d). In contrast, the untagged SmFtsZ protein (where instead of *M. genitalium* FtsZ

CTAH motif, *S. melliferum* FtsZ CCTP motif is present) did not bind to liposomes as no

detectable protein band was observed in the pellet of liposomes (Fig. 3c).

Further, we tested if the predicted C-terminal amphipathic helix could localise the core region

(amino acids 1 – 366 lacking the CCTP) of EcFtsZ to membranes in *E. coli*. Earlier studies

have shown that the core EcFtsZ$_{1-366}$-mVenus fusions localise to the membranes and organise

into spiral polymers or ring-like structures only in the presence of a membrane anchor such as

the amphipathic helix (residues 254 – 270) from EcMinD. We thus created a translation

fusion of the core region of EcFtsZ$_{1-366}$-mNeonGreen (EcFtsZ$_{1-366}$-mNG) to the MinD

amphipathic helix (EcFtsZ$_{1-366}$-mNG-MinD^{AH}) or the CTAH from *M. genitalium* FtsZ

(EcFtsZ$_{1-366}$-mNG-MgFtsZ^{CTAH}) and tested the localisation in an *E. coli* strain (JKD7-1)

depleted of the native endogenous FtsZ. At 30 ºC (in the presence of the native full-length

endogenous FtsZ), cells showed normal morphology and both EcFtsZ$_{1-366}$-mNG-MinD^{AH} and

EcFtsZ$_{1-366}$-mNG-MgFtsZ^{CTAH} localised to Z-rings, possibly due to co-polymerization with

the endogenous full-length FtsZ (Fig. 4a, b). However, at 42ºC, when the cells appear

filamentous (due to depletion of endogenous FtsZ), EcFtsZ$_{1-366}$-mNG-MgFtsZ^{CTAH} was seen

localised to the membranes (Fig. 4d) and was similar to those formed by EcFtsZ$_{1-366}$-mNG-

MinD^{AH} (Fig. 4c). We thus conclude that the predicted 23 amino acid residues at the C-

terminus of MgFtsZ can function as an amphipathic helix and target localization to liposomes

and membranes.

Sequence analysis identifies a cholesterol recognition motif in *M. genitalium* and *M.*

***pneumoniae* FtsZ**

A careful analysis of the *M. genitalium* and *M. pneumoniae* FtsZ sequences identified a

possible Cholesterol Recognition Amino acid Consensus (CRAC) motif located within the C-

terminal helix (Fig. 5a). The CRAC motif is defined as (**L/V**)-X$_{1-5}$-**Y**-X$_{1-5}$-(**R/K**) where "X"

represents any residue and central tyrosine has been shown to be essential for cholesterol

binding (Jamin *et al.*, 2005). We found that *M. genitalium* contains CRAC sequence as **L^{357}**-

X-X-X-**Y^{361}**-X-**K^{363}** whereas *M. pneumoniae* has **L^{367}**-X-X-X-**Y^{371}**-X-**K^{373}** in their respective

C-terminal ends of FtsZs which may give specificity for binding to cholesterol and thereby to

the membrane where 35-50% of total membrane lipid contents is cholesterol (47). The

presence of CRAC motif was only found in *M. genitalium* and *M. pneumoniae* species with

C-terminal amphipathic helix of the FtsZ sequences.

To indeed confirm that the proposed CRAC motif confers a preference for cholesterol, we

performed pelleting assay of SmFtsZ^{MgCTAH} construct with liposomes containing cholesterol

(in molar ratio of 30% DOPG, 30% DOPC, 40% Cholesterol). We observed an increased

membrane binding in cholesterol containing liposomes compared to liposomes made with

30% DOPG and 70% DOPC (Fig. 5c, 5d) which might indicate its preference for cholesterol.

We then disrupted the proposed CRAC motif (MCRAC) by mutating three residues E356K,

Y361V and K363E to see if it abolishes cholesterol specific binding (Fig. 5b). However,

interestingly, SmFtsZ$^{\text{MgCTAH(M-CRAC)}}$ too showed increased binding similar to SmFtsZ$^{\text{MgCTAH}}$

in the liposome pelleting experiments (Fig. 5c, 5d). Based on the local secondary structure

prediction of the C-terminal 23 amino acid peptide stretch, we hypothesise two helical

regions to be present separated by a short break in the helix with residues having low helix

forming probability (Fig. 5e) while the CRAC motif containing helix binds to cholesterol by

getting inserted inside membrane. The mutant construct SmFtsZ$^{\text{MgCTAH(M-CRAC)}}$ might result

in increased binding because of the variation in membrane binding properties and it was

challenging to delineate the distinction between the two functions (cholesterol recognition

and membrane binding).

Discussion

Here, we provide evidence that a phylogenetically separate FtsZ clade among

mycoplasmas possesses an inherent amphipathic helix that might facilitate FtsA-independent

membrane attachment. We experimentally show the putative C-terminal amphipathic helix of

M. genitalium FtsZ binds liposomes. Also, when we expressed the chimeric protein in *E. coli*

cell in a FtsZ depleted condition, it showed membrane localization which further confirms

membrane binding ability of MgFtsZ$^{\text{CTAH}}$ motif. In minimal genome bacterial species like

mycoplasma, the possibility of direct membrane binding in cell division protein FtsZ,

bypassing the requirement of additional adaptor proteins, is of relevance. FtsZ with a

membrane binding helix provides an ideal minimal system for mechanistic insights into the

process of cell constriction and division. Additionally, the presence of a CRAC motif makes

this core division protein an interesting model for studying division in mycoplasmas.

In previous studies, based on protein sequence analysis, the presence of amphipathic

helix was proposed in the N-terminal tail region of archaeal FtsZ (48). Another report on a

novel species of *Atribacteria* predicted a similar amphipathic helical region at the N-terminal

extension of FtsZ, which might bind to membrane (49). However, the membrane tethering

role of the proposed amphipathic helices from FtsZ has not been experimentally

demonstrated till date. Recently eukaryotic chloroplast FtsZ1 from *Arabidopsis thaliana* has

been shown to interact with membranes via its C-terminal amphipathic β-strand motif (50). It

is interesting that eukaryotic FtsZs have also evolved a mechanism of an inherent

amphipathic helix directly binding to the membrane, making the anchoring role of FtsA

redundant.

In the case of mycoplasma FtsZs, we found both N- and C-terminal putative

membrane attachment sites. The functional significance of such oppositely oriented

membrane binding sites is currently not clear from the perspective of FtsZ filament structure

and mycoplasma cell division. Earlier, a chimeric construct of FtsZ fused to a membrane

tethering sequence (from MinD protein) was shown to constrict tubular liposomes (51). It has

been also shown that changing the position of membrane targeting sequence in FtsZ from the

conventional C-terminal end to N-terminal end results in convex bulge rather than concave

depression (52). Interestingly in both the cases bending force could be generated which lead

to the tubulation of the liposomes *in vitro*. However, experiments with membrane targeting

sequence attached with FtsZ protein did not show full constriction. Septation was only

sometimes observed upon addition of FtsA indicating the requirement of FtsA and its role

beyond Z-ring anchoring (53). Loose and Mitchison's *in vitro* experiments with *E. coli* FtsZ

and FtsA on supported lipid bilayer also suggested a dual antagonistic role of FtsA in

anchoring and assembling FtsZ filaments as well as its rapid disassembly from the membrane

(34). Similarly, variations in the stoichiometry of FtsA and FtsZ revealed that excess FtsA

can effectively destabilize FtsZ polymers *in-vitro* (33). FtsA has also been suggested to be

involved in the septation process and membrane remodelling by introducing local curvature

in the inner membrane (29). Thus, FtsA could play multiple roles as membrane anchor as

well as in remodelling and regulating FtsZ filament dynamics. In contrast to the above

experiments where FtsA was used to anchor FtsZ to the membrane, the *M. genitalium* FtsZA

proteins, as revealed from our experiment and sequence analysis, are both potentially capable

of membrane binding and possibly work in a novel way to drive cell division. We believe *M.*

genitalium and *M. pneumoniae* FtsA could potentially serve as a model to study its

fundamental role beyond just FtsZ tethering.

M. genitalium and *M. pneumoniae* among mycoplasmas were additionally found to

contain a CRAC motif within the C-terminal amphipathic helix region. Our in vitro results

also showed increased membrane binding to cholesterol containing liposomes indicating the

possibility of cholesterol recognition through CRAC. CRAC motif has been known to exist in

numerous proteins spanning across eukaryotic and prokaryotic organisms (54–56). In

bacteria, CRAC motif has been so far identified solely in pathogenic bacterial toxins that

specifically bind to the cholesterol present in the membrane of eukaryotic host cells (57, 58).

CRAC sequence containing peptides within bacterial leukotoxin has been shown previously

to have higher affinity towards cholesterol and mutation in the CRAC motif abrogates

binding (59). However, our results of $SmFtsZ^{MgCTAH(M-CRAC)}$, the mutated construct for

CRAC motif, showed unaffected membrane binding property similar to $SmFtsZ^{MgCTAH}$ in the

presence of cholesterol. $SmFtsZ^{MgCTAH}$ bound to liposomes composed of 30% DOPG and

70% DOPC, although a marked increase in binding was observed in liposomes with 30%

DOPG, 30% DOPC, and 40% cholesterol or mycoplasma polar lipid composition which

consists of 25% DOPG, 20% DOPC, 30% cardiolipin, and 25% sphingomyelin. Moreover,

our *in vivo* experiments show that $SmFtsZ^{MgCTAH}$ bind *E. coli* membrane as well although

mycoplasma and *E. coli* have very different membrane composition. We think it is possible

that the CRAC mutated $SmFtsZ^{MgCTAH(M-CRAC)}$ construct still shows increased membrane

affinity in presence of cholesterol because membrane binding can be determined by multiple other factors such as overall lipid composition, hydrophobicity of AH, charge pattern, and lipid-amino acid interactions. For example, earlier studies on a eukaryotic sterol synthesis enzyme squalene monooxygenase showed similar results where cholesterol induced regulation remined unaffected even after disrupting the central critical tyrosine and phenylalanine residue of the CRAC and CARC motif respectively present within the membrane binding loop (60, 61). Thus, from our preliminary data we believe the unique CRAC motif present in *M. genitalium* FtsZ protein is functional and can interact with cholesterol. Additional data in future will shed more light into its cholesterol binding properties. Notably, the presence of a CRAC motif within FtsZ might be of significance as this could imply its membrane localization and subsequent role in cell division can be cholesterol dependent especially when 35-50% total mycoplasma membrane lipids is composed of cholesterol. We also checked if similar cholesterol recognition sequences are present in the other Mycoplasma FtsZs (containing N-terminal AH) and FtsAs (containing C-terminal AH) and did not find any. Thus, the membrane attachment of other AH containing FtsZ and FtsAs appear to follow separate mechanisms. Future *in vivo* studies will possibly give us more information about how membrane cholesterol composition affects cellular functions of mycoplasma divisome machinery.

Overall, our analysis highlights different possible modes of FtsZ-anchoring to the mycoplasma membrane, which are shown schematically in Fig. 6a-e. Group 1A FtsZ binds to FtsA in a similar mechanism as found in cell walled bacteria with its canonical CCTP (Fig. 6a). In some species where FtsA lacks the amphipathic helix, FtsZ ring probably needs SepF for membrane anchoring. One similar example can be found in mycobacteria where SepF acts as the membrane anchor for FtsZ and become an essential protein in this organism which lacks homologs of membrane tethers like FtsA or ZipA (62). Importantly, SepF mediated

membrane binding possibly occurs through hydrophobic loops rather than amphipathic helix.

Next, group 1B and 1C have deviant type of CCTPs for binding FtsA. However, because

group 1B FtsAs do not posses a canonical amphipathic helix, its binding towards membrane

is unclear (Fig. 6b, c). It is possible that smaller motifs of one or two amino acids can

contribute to membrane binding, as has been observed for DivIVA (63) and bactofilins (64).

Finally, group 1D and 1E do not have a consensus CCTP and experimental proof of whether

the non-canonical CCTP binds with FtsA is not available (Fig 6d, e). Group 1E FtsZs with N-

or C terminal amphipathic helix raises the possibility of direct membrane attachment making

FtsA's membrane tethering function dispensable. Based on our results here, we hypothesise

that the C-terminal amphipathic helix in *M. genitalium* and *M. pneumoniae* FtsZ will help it

to directly attach to the membrane. Once binding happens, the CRAC motif gets inserted into

the lipid bilayer and binds to interact with cholesterol (Fig. 6e right). However, additional

studies are required to test and verify this hypothesis. We believe our findings in

mycoplasmas to be an interesting observation to set the stage for designing new set of

experiments and looking into a novel membrane tethering role of FtsZ as well as to study

individual and cooperative roles of FtsZ and FtsA in ring constriction in the absence of cell

wall synthesis machinery.

Materials and methods

Sequence analysis of FtsZ and FtsA

We analysed a total 113 Mycoplasma species with complete sequence information of the

FtsZ and FtsA from the NCBI genome database. The division and cell wall (*dcw*) cluster

locus was first identified by manually checking genomes of every representative strain

(detailed information of the strains is listed in supplementary data sheet) through a keyword-

based search with 'FtsZ', 'FtsA', 'rsmH' (gene annotation for MraW protein) or 'MraZ'. We

had prior information about these four mycoplasma genes present in the dcw operon from the

existing literature. We found that the upstream coding sequence to the *ftsZ* gene was

annotated as a hypothetical protein-coding gene except for *Mycoplasma mycoides,* where it is

annotated as *ftsA*. Indeed, in some earlier published literature, the hypothetical gene has been

reported as '*ftsA*' (65). Hence, we downloaded the protein sequences of the *ftsZ* gene along

with the hypothetical protein-coding gene upstream of *ftsZ* for our analysis. We verified all

the FtsA sequences by multiple sequence alignment with known and already characterized

FtsA sequences from cell-walled model bacteria, followed by manual assignment of

secondary structures present in FtsA. Multiple sequence alignment of FtsZ and FtsA

sequences were performed in Jalview (https://www.jalview.org/) using Muscle (Edgar, 2004)

with the default parameter. FtsA domain assignment was done in PROMALS3D

(http://prodata.swmed.edu/promals3d/promals3d.php). The sequence logo was created using

WebLogo (https://weblogo.berkeley.edu/logo.cgi). For secondary structure prediction of

protein sequences, we used the PSIPRED server (http://bioinf.cs.ucl.ac.uk/psipred/).

HeliQuest (https://heliquest.ipmc.cnrs.fr/cgi-bin/ComputParams.py) was used for helical

wheel representation and calculation of hydrophobic moment and hydrophobicity parameters.

Phylogenetic trees were constructed using maximum likelihood method in MEGA 11 (66)

software with default parameters and visualized in iTOL (https://itol.embl.de/) (67).

Cloning and protein purification

SmftsZ gene was PCR amplified from the chromosomal DNA of *Spiroplasma melliferum*

KC3 followed by restriction digestion and ligation at the Nde1 and BamH1 sites into a $(His)_6$

tag containing pHis17 vector as reported earlier (68). Restriction free cloning (69) was used

for cloning of untagged SmFtsZ construct using specific primers. A detailed list of plasmid

constructs and oligonucleotide primers used in our study is given in Table S3. SmFtsZ^{MgCTAH}

was created by replacing 45 base pairs from the 3' end of the untagged SmFtsZ sequence

with respective 69 base pairs of *M. genitalium ftsZ* using PCR. The amplified product was

then inserted into (His)$_6$-SUMO-pET28a vector at the Nde1 and BamH1 sites by restriction

digestion and ligation. The construction of pJSB2-EcFtsZ$_{1-366}$-mNG-MinD^{AH} (pCCD907)

will be described in detail elsewhere, but is identical to the EcFtsZ$_{1-366}$-mVenus-MinD^{AH}

described by Erickson and colleagues (52), except that mNeonGreen replaces mVenus. The

EcMinD residues corresponding to amino acids 254 – 270 in pJSB2-EcFtsZ$_{1-366}$-mNG-

MinD^{AH} were replaced with the predicted C-terminal amphipathic helix (23 amino acids

corresponding to 347 – 366) from MgFtsZ by restriction free cloning to give pJSB2-EcFtsZ$_{1-}$

$_{366}$-mNG-MgFtsZ^{CTAH} (pCCD1071). pJSB2 is pBAD derivative carrying the tightly regulated

arabinose inducible promoter (70).

Untagged SmFtsZ was expressed in *E. coli* BL21-AI cells by 0.2% arabinose induction at

OD600= 0.6-0.8 and was grown at 30°C for 5 hours. Cells were harvested by centrifugation

at 4,000xg at 4°C for 20 mins and cell pellet was flash-frozen in liquid nitrogen and stored

at -80°C until processed. On the day of purification, cell pellet was taken out from -80°C and

thawed in ice for 10 mins and then resuspended in lysis buffer (50 mM Tris-HCl pH 8.0,

50 mM NaCl, 10% glycerol) followed by sonication to lyse the cells. The total lysate was

spun at 30,000xg for 1 hour to remove the cell debris and the clear supernatant was loaded

into HiTrap Q HP 5 ml column (GE) equilibrated in Buffer A50 (50 mM Tris-HCl pH 8.0,

50 mM NaCl). Protein was eluted in 5 mL fractions using a linear KCl gradient from 50

mM to 1000 mM over 20 column volumes. Peak fractions containing proteins were

confirmed from SDS PAGE, collected and pooled together for dialysis against A50 buffer

and loaded into MonoQ column to increase the purity level. Finally, the pure fractions were

confirmed from SDS PAGE gel and pooed for dialysis in 50 mM HEPES KOH storage buffer

(pH 7.4, 50 mM KCl). After dialyzing for 2 hours, the protein was concentrated using

centricon (with 10 kDa membrane cut off), flash-frozen in aliquots and stored at -80°C.

SmFtsZ^{MgCTAH} (in pET28a SUMO vector) was expressed in *E. coli* Rosetta (DE3) pLysS at

OD600= 0.5-0.6. by inducing with 0.6 mM IPTG and grown at 18°C for 16 hours. Cells

were harvested by centrifugation at 4,000xg for 20 mins at 4°C and cell pellet was flash-

frozen in liquid nitrogen and stored at -80°C until purification. Cells were resuspended in

lysis buffer (50 mM Tris-HCl pH 8.0, 200 mM KCl, 10% glycerol) and sonicated to lyse

the cells. The lysate was centrifuged at 30,000×g at 4°C for 1 hour and the supernatant was

loaded to a 5 mL His-Trap column (GE), equilibrated with Buffer A (50 mM Tris-HCl pH

8.0, 200 mM KCl). Protein was eluted using a step gradient of 2%, 5%, 10%, 20%, 50% and

100% Buffer B (50 mM Tris-HCl pH 8.0, 200 mM KCl, 500 mM Imidazole). To cleave

the N-terminal (His)$_6$-SUMO tag, peak protein fractions were pooled and Ulp1 protease was

added in a 1:100 (protease:protein) molar ratio and dialysis was set up against the cleavage

buffer (50 mM Tris-HCl pH 8.0, 200 mM KCl, 0.5 mM DTT) for 10 hours at 4°C. (His)$_6$-

SUMO tag and uncleaved protein was removed by passing the dialysed sample mixture

through a His-Trap column and flowthrough containing the SmFtsZ^{MgCTAH} protein was

collected. Finally, the protein was dialysed in 50 mM HEPES KOH storage buffer (pH 7.4,

50 mM KCl) for 2 hours, concentrated using centricon (with 10 kDa membrane cut off),

flash-frozen in aliquots and stored at -80°C. SmFtsZ$^{MgCTAH(M-CRAC)}$ protein purification was

done using the same protocol as SmFtsZ^{MgCTAH}. All three purified proteins were run on a

12% SDS PAGE for checking the purity level and quality (Fig. S8).

Growth conditions and microscopy

E. coli strains were grown in LB medium (Lennox or Miller) at 30 ℃ containing antibiotics

as required. Carbenicillin, kanamycin and chloramphenicol were used at 100 µg/ml, 25 µg/ml

and 34 µg/ml, respectively. Ampicillin was used at 200 µg/ml. *E. coli* strain JKD7_1 (W3110

recA56 ftsZ::kan) carrying pKD3 (Amp^R, *repA^{ts}*) (CCD56; (70) and the plasmid pJSB2

(Cm^R) (pCCD434; Stricker and Erickson, 2003), pJSB2-EcFtsZ$_{1-366}$-mNG-MinD^{AH}

(pCCD907) or pJSB2-EcFtsZ$_{1-366}$-mNG-MgFtsZ^{CTAH} (pCCD1071) was grown in LB Lennox

at 30 ℃ (permissive temperatures) or at 42 ℃ (non-permissive temperature) to allow for the

loss of pKD3 plasmid. Expression of EcFtsZ$_{1-366}$-mNG-MinD^{AH} and EcFtsZ$_{1-366}$-mNG-

MgFtsZ^{CTAH} from pJSB2 derived plasmids was achieved by the addition of 0.2 % L-

arabinose. Repression was achieved by addition of 2% glucose.

For microscopy, JKD7-1/pKD3 strain carrying the various pJSB2 derived plasmids were

grown in repression media (2 % glucose) at 30 ℃ and diluted 1:100 into fresh LB Lennox

medium containing 0.2 % L-arabinose and grown further at 42 °C for 2.5 hours. The cultures

were grown to an OD$_{600}$ between 0.4 - 0.6 and 1.5 mL culture was centrifuged at 1400 x *g* for

3 minutes. The cells pellets were resuspended in 100 – 150 µL of fresh LB medium, mounted

on LB-agarose pads (1.6 %) and imaged using an epifluorescence microscope (DeltaVision^{TM}

Elite) with a 100X oil immersion phase objective (PLN100XOPH) of NA 1.25. and equipped

with an in-line CCD camera (CoolSNAP^{TM} HQ2). Solid-state illumination (Spectra 7,

Lumencor Light Engine$^©$) was used as a light source with a 1 mm diameter liquid light guide.

Excitation filters and emission filters of 488/28 nm and 525/48 nm respectively were used for

imaging mNeonGreen tagged strains. Images were processed using Fiji v2.9.0/ 1.53t (71).

Liposome preparation and sedimentation assay

Liposomes were prepared by using synthetic lipids DOPC (1,2-dioleoyl-sn-glycero-3-phosphocholine), DOPG (1,2-dioleoyl-snglycero-3-phospho-rac-1-glycerol), and DPG/cardiolipin (1,3-bis[1,2-dioleoyl-sn-glycero-3-phospho]- glycerol) and SM (sphingomyelin), all purchased as chloroform solution from Avanti Polar Lipids, Inc. Cholesterol powder was purchased from Sigma. To prepare 2 mM stock liposomes, the required volume of individual components (as per molar ratio) was first taken in a glass tube. The chloroform was evaporated under a stream of nitrogen and further kept for drying at room temperature for 15 mins. A thin layer of lipids was visible at the bottom wall of the tube once the chloroform was dried. The dried lipid mixture was hydrated with the required volume of reaction buffer containing 50 mM HEPES KOH, 50 mM KCl (pH 7.4). After vortexing well, the mixture became turbid and it was extruded 15-20 times through a 100 nm pore-sized polycarbonate membrane until the solution became translucent.

For liposome sedimentation assay, purified flash-frozen protein was taken out from -80°C, thawed in ice for 10 mins and spun at 21,000xg for 1 hour at 4°C to remove any precipitation and supernatant was used in the assay. We set up a 100 µL reaction with 2 µM of protein and a varying concentration of liposome from 0 µM (i.e., no liposome control) to 1000 µM. The reaction mixture was incubated at 30°C for 20 mins and spun at 21,000xg for 40 mins to pellet down the liposomes (for pelleting cholesterol containing liposomes, ultracentrifugation at 100,000xg was used for 30 mins). Immediately after the spin, we took out the supernatant, washed the pellet with 100 µL of buffer and resuspended it in the same volume. Next, the supernatant and the pellet samples were mixed with 2X Laemmli buffer and equal amounts of both were loaded and run into a 12% SDS-PAGE gel followed by staining with Coomassie Brilliant Blue to check the presence of protein on the gel. Experiments were performed in triplicates and representative gel images are shown. The percentage intensities of protein bands in pellet and supernatant fractions were calculated as intensity of pellet (or supernatant)

528 divided by sum of the two intensities. The intensity of the bands were quantified using the

529 ImageJ software (Rueden *et al.*, 2017) by measuring the area of the respective bands and

530 percentage of protein in pellet fractions were calculated by dividing the pellet band intensity

531 by the sum of the pellet and supernatant band intensity. All the plots were generated in

532 GraphPad Prism software.

Author Contributions

SD and PG conceived the project and designed the experiments. SD conducted all

experiments and analysed the data except mentioned otherwise. JC contributed to cloning

and protein purification of untagged SmFtsZ. SMP carried out and analysed the *in vivo*

experiments related to imaging of JKD7-1 strains, while RS supervised, analysed the work

and edited the manuscript. SD and PG wrote the manuscript. PG supervised the project and

acquired funding. All authors read and approved the final submitted version of the

manuscript.

Acknowledgments

All authors sincerely thank IISER Pune for all the facilities to carry out this research. We

thank Ajay Kumar Sharma for the construction of pJSB2-EcFtsZ1-366-mNG-MinD[AH]. We

thank Harold Erickson (Duke University, USA) for strains and plasmids. We acknowledge

fellowships from CSIR to SD, IISER Pune to JC and NISER, DAE to SMP. Research work in

the lab of PG was supported by Department of Biotechnology (DBT) Membrane Structural

Biology Program grant (BT/PR28833/BRB/10/1705/2018), SERB CRG (CRG/2018/003795)

and IISER Pune. Work in RS lab was supported by SERB CRG (CRG/2021/000337) and

intra-mural funding from Department of Atomic Energy (DAE).

References

554   1.   Margolin W. 2005. FtsZ and the division of prokaryotic cells and organelles. 11. Nature Reviews
555   Molecular Cell Biology 6:862–871.

556   2.   Adams DW, Errington J. 2009. Bacterial cell division: assembly, maintenance and disassembly
557   of the Z ring. 9. Nature Reviews Microbiology 7:642–653.

558   3.   Bisson-Filho AW, Hsu Y-P, Squyres GR, Kuru E, Wu F, Jukes C, Sun Y, Dekker C, Holden S,
559   VanNieuwenhze MS, Brun YV, Garner EC. 2017. Treadmilling by FtsZ filaments drives
560   peptidoglycan synthesis and bacterial cell division 6.

561   4.   Yang X, Lyu Z, Miguel A, McQuillen R, Huang KC, Xiao J. 2017. GTPase activity–coupled
562   treadmilling of the bacterial tubulin FtsZ organizes septal cell wall synthesis. 6326. Science
563   355:744–747.

564   5.   Whitley KD, Jukes C, Tregidgo N, Karinou E, Almada P, Cesbron Y, Henriques R, Dekker C,
565   Holden S. 2021. FtsZ treadmilling is essential for Z-ring condensation and septal constriction
566   initiation in Bacillus subtilis cell division. 1. Nat Commun 12:2448.

567   6.   Squyres GR, Holmes MJ, Barger SR, Pennycook BR, Ryan J, Yan VT, Garner EC. 2021. Single-
568   molecule imaging reveals that Z-ring condensation is essential for cell division in Bacillus
569   subtilis. Nat Microbiol 6:553–562.

570   7.   Xiao J, Goley ED. 2016. Redefining the roles of the FtsZ-ring in bacterial cytokinesis. Current
571   Opinion in Microbiology 34:90–96.

572   8.   Coltharp C, Xiao J. 2017. Beyond force generation: Why is a dynamic ring of FtsZ polymers
573   essential for bacterial cytokinesis? 1. BioEssays 39:e201600179.

9.   Pichoff S, Lutkenhaus J. 2005. Tethering the Z ring to the membrane through a conserved

membrane targeting sequence in FtsA. 6. Molecular Microbiology 55:1722–1734.

10.   Juarez JR, Margolin W. 2012. A bacterial actin unites to divide bacterial cells: A bacterial actin

unites to divide bacterial cells. 10. The EMBO Journal 31:2235–2236.

11.   Corbin BD, Geissler B, Sadasivam M, Margolin W. 2004. Z-ring-independent interaction

between a subdomain of FtsA and late septation proteins as revealed by a polar recruitment

assay. J Bacteriol 186:7736–7744.

12.   Busiek KK, Eraso JM, Wang Y, Margolin W. 2012. The Early Divisome Protein FtsA Interacts

Directly through Its 1c Subdomain with the Cytoplasmic Domain of the Late Divisome Protein

FtsN. J Bacteriol 194:1989–2000.

13.   Du S, Pichoff S, Lutkenhaus J. 2016. FtsEX acts on FtsA to regulate divisome assembly and

activity. Proceedings of the National Academy of Sciences 113:E5052–E5061.

14.   Vega DE, Margolin W. 2019. Direct Interaction between the Two Z Ring Membrane Anchors

FtsA and ZipA. J Bacteriol 201:e00579-18.

15.   Park K-T, Du S, Lutkenhaus J. 2020. Essential Role for FtsL in Activation of Septal Peptidoglycan

Synthesis. mBio 11:e03012-20.

16.   Hale CA, de Boer PAJ. 1997. Direct Binding of FtsZ to ZipA, an Essential Component of the

Septal Ring Structure That Mediates Cell Division in E. coli. Cell 88:175–185.

17.   Hamoen LW, Meile J-C, Jong WD, Noirot P, Errington J. 2006. SepF, a novel FtsZ-interacting

protein required for a late step in cell division. 3. Molecular Microbiology 59:989–999.

18.   Löwe J, Amos LA. 1998. Crystal structure of the bacterial cell-division protein FtsZ. 6663.

Nature 391:203–206.

19.   Erickson HP. 1995. FtsZ, a prokaryotic homolog of tubulin? Cell 80:367–370.

20.   Nogales E, Downing KH, Amos LA, Löwe J. 1998. Tubulin and FtsZ form a distinct family of
598   GTPases. 6. Nat Struct Mol Biol 5:451–458.

599   21.   Ma X, Margolin W. 1999. Genetic and Functional Analyses of the Conserved C-Terminal Core
600   Domain of Escherichia coli FtsZ. J Bacteriol 181:7531–7544.

601   22.   Yan K, Pearce KH, Payne DJ. 2000. A Conserved Residue at the Extreme C-Terminus of FtsZ Is
602   Critical for the FtsA-FtsZ Interaction in Staphylococcus aureus. 2. Biochemical and Biophysical
603   Research Communications 270:387–392.

604   23.   Ma X, Sun Q, Wang R, Singh G, Jonietz EL, Margolin W. 1997. Interactions between
605   heterologous FtsA and FtsZ proteins at the FtsZ ring. 21. Journal of Bacteriology 179:6788–
606   6797.

607   24.   Mosyak L, Zhang Y, Glasfeld E, Haney S, Stahl M, Seehra J, Somers WS. 2000. The bacterial cell-
608   division protein ZipA and its interaction with an FtsZ fragment revealed by X-ray
609   crystallography. EMBO J 19:3179–3191.

610   25.   Duman R, Ishikawa S, Celik I, Strahl H, Ogasawara N, Troc P, Lowe J, Hamoen LW. 2013.
611   Structural and genetic analyses reveal the protein SepF as a new membrane anchor for the Z
612   ring. 48. Proceedings of the National Academy of Sciences 110:E4601–E4610.

613   26.   Busiek KK, Margolin W. 2015. Bacterial Actin and Tubulin Homologs in Cell Growth and
614   Division. 6. Current Biology 25:R243–R254.

615   27.   Szwedziak P, Wang Q, Freund SM, Löwe J. 2012. FtsA forms actin-like protofilaments: FtsA
616   forms actin-like protofilaments. 10. The EMBO Journal 31:2249–2260.

617   28.   Lara B, Rico AI, Petruzzelli S, Santona A, Dumas J, Biton J, Vicente M, Mingorance J, Massidda

618   O. 2005. Cell division in cocci: localization and properties of the Streptococcus pneumoniae

619   FtsA protein. Mol Microbiol 55:699–711.

620   29.   Krupka M, Cabré EJ, Jiménez M, Rivas G, Rico AI, Vicente M. 2014. Role of the FtsA C Terminus

621   as a Switch for Polymerization and Membrane Association. 6. mBio 5:e02221-14.

622   30.   Krupka M, Rowlett VW, Morado D, Vitrac H, Schoenemann K, Liu J, Margolin W. 2017.

623   Escherichia coli FtsA forms lipid-bound minirings that antagonize lateral interactions between

624   FtsZ protofilaments. Nature Communications 8:15957.

625   31.   Morrison JJ, Conti J, Camberg JL. 2022. Assembly and architecture of Escherichia coli divisome

626   proteins FtsA and FtsZ. Journal of Biological Chemistry 298:101663.

627   32.   Nag D, Chatterjee A, Chakrabarti G. 2020. FtsA-FtsZ interaction in Vibrio cholerae causes

628   conformational change of FtsA resulting in inhibition of ATP hydrolysis and polymerization.

629   International Journal of Biological Macromolecules 142:18–32.

630   33.   Conti J, Viola MG, Camberg JL. 2018. FtsA reshapes membrane architecture and remodels the

631   Z-ring in *Escherichia coli*: FtsA reshapes membrane architecture. 4. Molecular Microbiology

632   107:558–576.

633   34.   Loose M, Mitchison TJ. 2014. The bacterial cell division proteins FtsA and FtsZ self-organize into

634   dynamic cytoskeletal patterns. Nat Cell Biol 16:38–46.

635   35.   McQuillen R, Xiao J. 2020. Insights into the Structure, Function, and Dynamics of the Bacterial

636   Cytokinetic FtsZ-Ring. 1. Annual Review of Biophysics 49:309–341.

637   36.   den Blaauwen T, Hamoen LW, Levin PA. 2017. The divisome at 25: the road ahead. Current

638   Opinion in Microbiology 36:85–94.

639   37.   Margolin W. 2000. Themes and variations in prokaryotic cell division. 4. FEMS Microbiol Rev

640   24:531–548.

641   38.   Bové JM. 1993. Molecular Features of Mollicutes. Clinical Infectious Diseases 17:S10–S31.

642   39.   Alarcón F, Vasconcelos ATR de, Yim L, Zaha A. 2007. Genes involved in cell division in

643   mycoplasmas. 1 suppl. Genetics and Molecular Biology 30:174–181.

644   40.   Pelletier JF, Sun L, Wise KS, Assad-Garcia N, Karas BJ, Deerinck TJ, Ellisman MH, Mershin A,

645   Gershenfeld N, Chuang R-Y, Glass JI, Strychalski EA. 2021. Genetic requirements for cell division

646   in a genomically minimal cell. Cell 184:2430-2440.e16.

647   41.   Lluch-Senar M, Querol E, Piñol J. 2010. Cell division in a minimal bacterium in the absence of

648   ftsZ. 2. Molecular Microbiology 78:278–289.

649   42.   Hasselbring BM, Jordan JL, Krause RW, Krause DC. 2006. Terminal organelle development in

650   the cell wall-less bacterium Mycoplasma pneumoniae. 44. PNAS 103:16478–16483.

651   43.   Briers Y, Walde P, Schuppler M, Loessner MJ. 2012. How did bacterial ancestors reproduce?

652   Lessons from L-form cells and giant lipid vesicles. BioEssays 34:1078–1084.

653   44.   Mercier R, Kawai Y, Errington J. 2014. General principles for the formation and proliferation of

654   a wall-free (L-form) state in bacteria. eLife 3:e04629.

655   45.   Martínez-Torró C, Torres-Puig S, Marcos-Silva M, Huguet-Ramón M, Muñoz-Navarro C, Lluch-

656   Senar M, Serrano L, Querol E, Piñol J, Pich OQ. 2021. Functional Characterization of the Cell

657   Division Gene Cluster of the Wall-less Bacterium Mycoplasma genitalium. Frontiers in

658   Microbiology 12:2521.

46.   Gaspari E, Malachowski A, Garcia-Morales L, Burgos R, Serrano L, Martins dos Santos VAP,

Suarez-Diez M. 2020. Model-driven design allows growth of Mycoplasma pneumoniae on

serum-free media. npj Syst Biol Appl 6:1–11.

47.   Dahl J. 1993. The Role of Cholesterol in Mycoplasma Membranes, p. 167–188. *In* Rottem, S,

Kahane, I (eds.), Mycoplasma Cell Membranes. Springer US, Boston, MA.

48.   Liao Y, Ithurbide S, Evenhuis C, Löwe J, Duggin IG. 2021. Cell division in the archaeon Haloferax

volcanii relies on two FtsZ proteins with distinct functions in division ring assembly and

constriction. 5. Nat Microbiol 6:594–605.

49.   Katayama T, Nobu MK, Kusada H, Meng X-Y, Hosogi N, Uematsu K, Yoshioka H, Kamagata Y,

Tamaki H. 2020. Isolation of a member of the candidate phylum 'Atribacteria' reveals a unique

cell membrane structure. 1. Nature Communications 11:6381.

50.   Liu X, An J, Wang L, Sun Q, An C, Wu B, Hong C, Wang X, Dong S, Guo J, Feng Y, Gao H. 2022. A

novel amphiphilic motif at the C-terminus of FtsZ1 facilitates chloroplast division. The Plant Cell

34:419–432.

51.   Osawa M, Anderson DE, Erickson HP. 2008. Reconstitution of Contractile FtsZ Rings in

Liposomes. Science 320:792–794.

52.   Osawa M, Anderson DE, Erickson HP. 2009. Curved FtsZ protofilaments generate bending

forces on liposome membranes. 22. The EMBO Journal 28:3476–3484.

53.   Osawa M, Erickson HP. 2013. Liposome division by a simple bacterial division machinery. 27.

PNAS 110:11000–11004.

54.   Wood WG, Schroeder F, Igbavboa U, Avdulov NA, Chochina SV. 2002. Brain membrane

cholesterol domains, aging and amyloid beta-peptides. Neurobiology of Aging 23:685–694.

55.   Jafurulla Md, Tiwari S, Chattopadhyay A. 2011. Identification of cholesterol recognition amino

acid consensus (CRAC) motif in G-protein coupled receptors. Biochemical and Biophysical

Research Communications 404:569–573.

56.   Fantini J, Carlus D, Yahi N. 2011. The fusogenic tilted peptide (67–78) of α-synuclein is a

cholesterol binding domain. Biochimica et Biophysica Acta (BBA) - Biomembranes 1808:2343–

2351.

57.   Boesze-Battaglia K, Brown A, Walker L, Besack D, Zekavat A, Wrenn S, Krummenacher C,

Shenker BJ. 2009. Cytolethal Distending Toxin-induced Cell Cycle Arrest of Lymphocytes Is

Dependent upon Recognition and Binding to Cholesterol*. Journal of Biological Chemistry

284:10650–10658.

58.   Brown AC, Balashova NV, Epand RM, Epand RF, Bragin A, Kachlany SC, Walters MJ, Du Y,

Boesze-Battaglia K, Lally ET. 2013. Aggregatibacter actinomycetemcomitans leukotoxin utilizes

a cholesterol recognition/amino acid consensus site for membrane association. J Biol Chem

288:23607–23621.

59.   Koufos E, Chang EH, Rasti ES, Krueger E, Brown AC. 2016. Use of a Cholesterol Recognition

Amino Acid Consensus Peptide To Inhibit Binding of a Bacterial Toxin to Cholesterol.

Biochemistry 55:4787–4797.

60.   Chua NK, Howe V, Jatana N, Thukral L, Brown AJ. 2017. A conserved degron containing an

amphipathic helix regulates the cholesterol-mediated turnover of human squalene

monooxygenase, a rate-limiting enzyme in cholesterol synthesis. Journal of Biological

Chemistry 292:19959–19973.

61.   Chua NK, Coates HW, Brown AJ. 2020. Squalene monooxygenase: a journey to the heart of

cholesterol synthesis. Progress in Lipid Research 79:101033.

62. Gola S, Munder T, Casonato S, Manganelli R, Vicente M. 2015. The essential role of SepF in
mycobacterial division. 3. Mol Microbiol 97:560–576.

63. Oliva MA, Halbedel S, Freund SM, Dutow P, Leonard TA, Veprintsev DB, Hamoen LW, Löwe J.
2010. Features critical for membrane binding revealed by DivIVA crystal structure. The EMBO
Journal 29:1988–2001.

64. Deng X, Gonzalez Llamazares A, Wagstaff JM, Hale VL, Cannone G, McLaughlin SH, Kureisaite-
Ciziene D, Löwe J. 2019. The structure of bactofilin filaments reveals their mode of membrane
binding and lack of polarity. 12. Nat Microbiol 4:2357–2368.

65. Fisunov GY, Evsyutina DV, Semashko TA, Arzamasov AA, Manuvera VA, Letarov AV, Govorun
VM. 2016. Binding site of MraZ transcription factor in Mollicutes. Biochimie 125:59–65.

66. Tamura K, Stecher G, Kumar S. 2021. MEGA11: Molecular Evolutionary Genetics Analysis
Version 11. Molecular Biology and Evolution 38:3022–3027.

67. Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. Nucleic Acids Research 49:W293–W296.

68. Chakraborty J, Poddar SM, Dutta S, Bahulekar V, Harne S, Srinivasan R, Gayathri P. 2022.
Structures of FtsZ from a cell wall-less bacterium Spiroplasma provide a mechanism for kinetic
polarity. bioRxiv https://doi.org/10.1101/2022.10.13.512043.

69. van den Ent F, Löwe J. 2006. RF cloning: A restriction-free method for inserting target genes
into plasmids. 1. Journal of Biochemical and Biophysical Methods 67:67–74.

70. Stricker J, Erickson HP. 2003. In Vivo Characterization of Escherichia coli ftsZ Mutants: Effects
on Z-Ring Structure and Function. Journal of Bacteriology 185:4796–4805.

71.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,

Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A.

2012. Fiji: an open-source platform for biological-image analysis. Nat Methods 9:676–682.

Figure legends

Fig. 1 A scheme for classification of mycoplasma species based on the presence of *ftsA*

and *ftsZ* in their genomes

(a) Genomic locus of division and cell wall cluster (*dcw*) operon genes of *Escherichia coli*,

Bacillus subtilis, and cell wall-less bacteria. The arrowhead indicates gene orientation 5′ →

3′.

(b) Domain architecture of FtsA, FtsZ.

(c) 16S rRNA-based tree of 113 mycoplasma species is constructed using maximum

likelihood analysis and the distribution of FtsZ and FtsA proteins are highlighted. Solid red

(for FtsZ) and solid blue (for FtsA) circles represent the presence of the proteins and

transparent colored circles indicate their absence. Outlier species which are away from

clusters of the respective groups are marked as yellow colored star (*).

(d) Grouping of mycoplasmas based on the presence of FtsZ and FtsA. The number of

organisms consisting each group is mentioned within brackets.

Fig. 2 Classification based on conserved C-terminal peptide (CCTP) sequences matches

with the phylogenetic relationship of mycoplasma FtsZs

(a) Group 1 mycoplasmas having FtsZ and FtsA are grouped into five categories based on the

various types of CCTP present. Groupwise WebLogos are shown for three FtsZ motifs: GTP

binding, hydrolysis, and CCTP which are found to be conserved in cell-walled bacteria. The

number within the bracket for a group represents number of sequences used for generating

the Weblogo.

(b) Cladogram of FtsZ from group 1A mycoplasma species is constructed using the

maximum likelihood method. Different clusters are highlighted and labeled as 1A – E as per

our annotated groups based on CCTP motifs presents at the C-terminal end of FtsZ.

Fig. 3 Mycoplasma FtsZ representative of Group 1E binds to membrane directly

through an inherent amphipathic helix

(a) Helical wheel projections show the amphipathic nature of the helix-forming amino acids

at the FtsZ terminals. Hydrophobic residues are depicted in yellow showing the amphipathic

face. Arrows indicate the hydrophobic moments of the respective helices.

(b) Hydrophobic moment plot of the putative N-terminal and C-terminal α-helices of group

1E FtsZs. The hydrophobic moment is plotted against hydrophobicity to predict its potential

to bind to the membrane. Known amphipathic helices from FtsA, MinD, and SepF also

plotted simultaneously for comparison.

(c) A representative 12% SDS-PAGE gel showing the presence of SmFtsZ^{MgCTAH} and

Untagged SmFtsZ protein in the supernatant (S) and pellet fraction (P) with increasing

concentrations of liposomes from 0 μM (i.e. no liposome control) to 1000 μM.

(d) Quantification plot for SmFtsZ^{MgCTAH} protein in the pellet fraction. The fraction of protein

was calculated from the SDS PAGE gel band intensity in the pellet fraction divided by the

sum of band intensities in pellet and supernatant. Error bar denotes standard deviation

calculated from three independent experiments. ND: Not detectable.

Fig. 4 C-terminal amphipathic helix of MgFtsZ can bind to membrane *in vivo* in *E. coli*

cells

The predicted C-terminal amphipathic helix of MgFtsZ can localise EcFtsZ$_{1-366}$ (lacking

CCTP) to membranes in *E. coli*. Localization of (i) EcFtsZ$_{1-366}$-mNG-MinD^{AH} and (ii)

EcFtsZ$_{1-366}$-MgFtsZ^{CTAH} to Z-rings formed by endogenous native full-length FtsZ at 30 ℃.

Localization of (iii) EcFtsZ$_{1-366}$-mNG-MinD^{AH} and (iv) EcFtsZ$_{1-366}$-MgFtsZ^{CTAH} to inner

membrane of filamentous cells due to depletion of native full-length FtsZ at 42 ℃.

Arrowheads mark the regions of membrane localisation. Scale bars represent 10 µm.

Fig. 5 Cholesterol Recognition Amino acid Consensus (CRAC) motif in *M. genitalium*

FtsZ provides it a preference for cholesterol rich liposome binding

(a) Sequence alignment of C-terminal end region of *M. genitalium* and *M. pneumoniae* FtsZ

shows the presence of CRAC motif specific conserved residues (marked with '*'). The

consensus motif is mentioned below the alignment.

(b) Sequence of SmFtsZ^{MgCTAH} (denoted as CRAC) and SmFtsZ$^{MgCTAH(M-CRAC)}$ (denoted as

MCRAC) C-terminal motifs. The consensus amino acid sequence specific to CRAC motif are

highlighted in bold. The mutated residues in MCRAC are underlined.

(c) Representative 12% SDS-PAGE gels showing cholesterol mediated induction of

membrane binding in CRAC (i.e SmFtsZ^{MgCTAH}) and MCRAC (i.e. SmFtsZ$^{MgCTAH(M-CRAC)}$) in

liposome pelleting assay. Liposome containing cholesterol (+Chol) has higher amount of

protein in pellet fraction compared to the liposome without cholesterol (-Chol).

(d) Quantification of protein amount in pellet fraction in liposome pelleting assay. CTAH and

MCRAC both show more binding with liposomes containing cholesterol with a significant

difference in 1000 µM liposome conditions. Error bars indicate standard deviation from three

independent experiments. p values are obtained from an unpaired two tailed t-test using the

function 'T.TEST' in Microsoft Excel. Significance level is denoted by *, $p < 0.05$; **, $p <$

0.01; ***, $p < 0.005$); ND, not detected.

(e) Local secondary structure of the C-terminal end region of *M. genitalium* FtsZ was

predicted using PEP-FOLD server (https://bioserv.rpbs.univ-paris-diderot.fr/services/PEP-

FOLD/). Two helical regions have been identified of which the Helix 2 contains the

hypothetical CRAC motif sequence that could get inserted inside membrane to interact with

cholesterol. The schematic shown below the plot represents the same.

Fig. 6 Modes of FtsZA interactions across different mycoplasma groups. The possible

modes of FtsZ-FtsA interaction and their membrane-binding potentials are shown

schematically for five mycoplasma groups.

Fig. 1

Tree scale: 1

Fig. 2

a

Conserved FtsZ motifs			Group
GTP binding [S/A/G]–G–G–T–G–[S/A/T]–G	**GTP hydrolysis** N–X–D–X–X–D	**CCTP** D/E–I/V–P–X–F/Y–L–R/K	

 GGGTGTG | NVDFAD | D.LD.P.F.R | Walled (6) |
GGGTGTG	NLDFAD	SP..PP.FP.FLK	1A (14)
GGGTGTG	NLDFAD	DDLA.D.FK	1B (20)
GGGTGTG	NLDFND	ENP..P.FF	1C (21)
GGGTGTG	N|DYAD	TE.NSYE	1D (15)
GKGTGTG	NLDEAD	SK.F.D	1E (12)

b

Fig. 3

a

b

Hydrophobic moment vs **Hydrophobicity**

Legend:
- ■ Mycoplasma FtsZ
- ● FtsA
- ▲ MinD
- ▼ SepF

c

Liposome (µM): 0, 50, 200, 600, 1000 (S / P each)

SmFtsZ^MgCTAH ◄ 50 kDa

Untagged SmFtsZ ◄ 50 kDa

d

% Protein vs Liposome conc. (µM)
- □ Supernatant
- ■ Pellet
- ND at 0

Fig. 4

a   30°C (+EcFtsZ_FL)

b

c   42°C (EcFtsZ_FL Depleted)

d

$\mathbf{EcFtsZ_{1\text{-}366}\text{-}MinD^{AH}}$ $\mathbf{EcFtsZ_{1\text{-}366}\text{-}MgFtsZ^{CTAH}}$

Fig. 5

a

										10									20			

M.genitalium/347-369 K L K L L D E L K E L G M K Y V K H Q N Q I - Y

M.pneumoniae/357-380 P F Q V L N D L K E L G L K Y V K Q Q T G F N Y

-**L/V**–(X)$_{1-5}$–**Y**–(X)$_{1-5}$–**R/K**–

L-X-X-X-**Y**-X-**K**

b

	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369
CRAC	K	L	K	L	L	D	E	L	K	E	L	G	M	K	Y	V	K	H	Q	N	Q	I	Y
MCRAC	K	L	K	L	L	D	E	L	K	K	L	G	M	K	V	V	E	H	Q	N	Q	I	Y

c

e

helical (red), coil (blue), extended (green)

d

Fig. 6

Group 1A Group 1B Group 1C

a b c

Group 1D Group 1E

d e

Amphipathic helix

Hydrophobic loop

Different classes CCTPs

CRAC motif

Cholesterol

Lipids

? Unknown interaction