Title:

Netrin signaling mediates survival of dormant epithelial ovarian cancer cells

Authors:

Pirunthan Perampalam[1,2], James I. MacDonald[1,3], Komila Zakirova[1,3], Daniel T. Passos[1,3], Sumaiyah Wasif[1,3], Yudith Ramos-Valdes[1,4], Maëva Hervieu[5], Patrick Mehlen[5,6], Rob Rottapel[7,8], Benjamin Gibert[5], Rohann Correa[1,9], Trevor G. Shepherd[1,4,9,10,11], Frederick A. Dick[1,3,9,12,13]

Affiliations:

1. London Regional Cancer Program, London Health Sciences Centre Research Institute, London, Ontario, Canada, N6A 4L6.
2. Department of Biochemistry, University of Western Ontario, London, ON, Canada, N6A 5C1.
3. Department of Pathology and Laboratory Medicine, University of Western Ontario, London, ON, Canada, N6A 5C1.
4. The Mary and John Knight Translational Ovarian Cancer Research Unit, London Regional Cancer Program, London, Ontario, Canada, N6A 4L6.
5. Apoptosis, Cancer and Development Laboratory - Equipe labellisée 'La Ligue', LabEx DEVweCAN, Institut Convergence PLAsCAN, Centre de Recherche en Cancérologie de Lyon (CRCL), INSERM U1052-CNRS UMR5286, Université de Lyon, Université Claude Bernard Lyon1, Centre Léon Bérard, 69008 Lyon, France.
6. Netris Pharma, 69008 Lyon, France
7. Princess Margaret Cancer Centre, University Health Network, 101 College Street, Toronto, Ontario, Canada, M5G 1L7
8. Department of Medical Biophysics, University of Toronto, 1 King's College Circle, Toronto, Ontario, Canada, M5S 1A8
9. Department of Oncology, Western University, London, Ontario, Canada, N6A 4L6.
10. Department of Obstetrics and Gynecology, Western University, London, Ontario, Canada, N6A 5W9.
11. Department of Anatomy and Cell Biology, Western University, London, Ontario, Canada, N6A 5C1.
12. Children's Health Research Institute, London, Ontario, Canada, N6A 4V2.
13. Correspondence: fdick@uwo.ca

Keywords:
Netrin, high grade serous ovarian cancer, DYRK1A, dormancy, quiescence, spheroids

Conflict of interest statement:
P. Mehlen is the founder and CEO of Netris Pharma and has an ownership interest including stocks and patents. The other authors have declared that no conflict of interest exists.

Abstract

Dormancy in cancer is a clinical state in which residual disease remains undetectable for a prolonged duration. At a cellular level, rare cancer cells cease proliferation and survive chemotherapy and disseminate disease. We created a suspension culture model of high grade serous ovarian cancer (HGSOC) dormancy and devised a novel CRISPR screening approach to identify survival genes in this context. In combination with RNA-seq, we discovered the Netrin signaling pathway as critical to dormant HGSOC cell survival. We demonstrate that Netrin-1, -3, and its receptors are essential for low level ERK activation to promote survival, and that Netrin activation of ERK is unable to induce proliferation. Deletion of all UNC5 family receptors blocks Netrin signaling in HGSOC cells and compromises viability during the dormancy step of dissemination in xenograft assays. Furthermore, we demonstrate that Netrin-1 and -3 overexpression in HGSOC correlates with poor outcome. Specifically, our experiments reveal that Netrin overexpression elevates cell survival in dormant culture conditions and contributes to greater spread of disease in a xenograft model of abdominal dissemination. This study highlights Netrin signaling as a key mediator HGSOC cancer cell dormancy and metastasis.

Introduction

Relapse from initially effective cancer treatment represents one of the greatest barriers to improving outcomes (1). Dormant or minimal residual disease that is not clinically detectable is a common source of relapse, particularly in a metastatic setting (2). This has motivated the search for, and study of, disseminated cancer cells. Cellular dormancy has been characterized in cancer cells derived from a variety of primary disease sites including breast, prostate, melanoma, and others (3, 4). Disseminated cells often occupy specific cellular niches in perivascular space or in bone marrow (5). They adopt catabolic metabolism (2), and withdraw from the cell cycle (4). Stem like phenotypes and epithelial to mesenchymal transition also characterize the biology of these cells as they survive dissemination and lay quiescent in distant tissues (1, 3, 5). Across a variety of disease sites this dormant state is characterized by diminished Ras-MAPK signaling and elevated p38 activity (3, 6). To develop treatments specifically tailored for dormant cells, a systematic search for vulnerabilities unique to dormancy is needed. To this end, high grade serous ovarian cancer (HGSOC) disseminates in ascites as growth arrested cellular aggregates (7). This has been modeled in culture (8), and studies confirm active induction of cell cycle arrest mechanisms (9, 10), induction of autophagy (11, 12), and other known characteristics of dormancy (13). This suggests that HGSOC spheroids are ideal to search for survival dependencies specific to dormant cells.

High grade serous ovarian cancer (HGSOC) is the deadliest gynecologic malignancy, accounting for more than 70% of cases (14). Its poor prognosis is largely attributed to late-stage diagnosis with metastases already present. Metastatic spread is facilitated by multicellular clusters of tumor cells, called spheroids, that are shed into the peritoneal cavity and disseminate to nearby organs (15). The non-proliferative state of spheroid cells renders them chemoresistant

and contributes to disease recurrence, emphasizing the need to understand the biology of these

cells (16). Dormant HGSOC cells are characterized by similar metabolic changes, EMT, and

cytokine signaling as dormant cancer cells from other disease sites (3, 13). Mechanisms that

mediate their dormant properties include AMPK-LKB1 signaling that induces autophagy (17,

18), STAT3 and FAK for survival and chemoresistance (19, 20), and DYRK1A signaling to

arrest proliferation (10). Unfortunately, therapeutic approaches to target these pathways in

ovarian cancer have yet to emerge, indicating that more candidates are needed.

Netrins are a family of secreted factors that reside in the extracellular matrix and were

first identified in axon guidance in the developing nervous system (21, 22). Together with their

receptors, Netrins have since been implicated in cancer progression (23). Netrin-1 is the most

studied and its misregulation in cancer leads to pro-survival signals (24). Netrin interactions

with DCC and UNC5 homologs (UNC5H, UNC5A-D) convert pro-death signals by these

dependence receptors to survival signals (25). Netrin signals through PI3K-AKT and ERK-

MAPK are known to participate in orientation decisions in development by providing guidance

cues (26-28), while loss of ligand binding leads DCC or UNC5H to stimulate dephosphorylation

of DAPK1 that sends pro-death signals (29, 30). The gene encoding DCC is frequently deleted

in cancer (24), and Netrin-1 blocking antibodies are in Phase II clinical trials for endometrial and

cervical carcinomas to inhibit metastatic growth (25). Alternative receptors for Netrins such as

NEO, DSCAM, and Integrins also mediate their signals (23), creating a myriad of possible

pathways for Netrins to regulate cancer cells. Lastly, functional roles for Netrin-3, -4, and -5 are

still relatively unexplored compared to Netrin-1 (31), although their expression patterns suggest

roles in advanced forms of some cancers (32). A functional role for Netrins and their receptors

in dormancy, or other aspects of HGSOC spheroid biology have yet to be reported.

In this study we utilized a suspension culture system to interrogate survival mechanisms in dormancy using HGSOC spheroids. To maximize discovery of gene loss events that compromise viability in dormancy, we devised a novel genome-wide CRISPR screen approach that we term 'GO-CRISPR'. In addition to the standard CRISPR workflow, GO-CRISPR utilizes a parallel screen in which 'guide-only' cells lacking Cas9 are used to control for stochastic changes in gRNA abundance in non-proliferative cells. This screen identified multiple Netrin ligands, DCC, UNC5Hs, and downstream MAPK signaling components as supporting survival of HGSOC cells in dormant culture conditions. A transcriptomic analysis of HGSOC spheroids also revealed Netrin signaling components were enriched in HGSOC spheroids compared to adherent cells, and that in the absence of the DYRK1A survival kinase Netrin transcriptional increases were lost. We show that multiple Netrins, UNC5 receptors, and the downstream MEK-ERK axis is crucial for survival signaling in dormant cell culture. Importantly, Netrin activation of ERK is restricted to spheroid cultures and is unable to stimulate proliferation, consistent with its role in dormant survival. We blocked Netrin signaling by deleting UNC5 receptors and using xenograft approaches demonstrated that loss of Netrin signaling compromises survival of HGSOC cells, specifically in the dormant stage of dissemination. Furthermore, we show that overexpression of Netrin-1 and -3 are associated with poor prognosis in HGSOC and their overexpression in culture increases survival in suspension culture. Netrin overexpression in xenografts contributes to increased disease dissemination in a model of metastasis. Our study highlights Netrin-ERK signaling as a requirement for spheroid survival and suggests it may be a potential therapeutic target for eradicating dormant HGSOC cells.

Results

Axon guidance genes are essential factors for HGSOC spheroid viability

To dissect genetic dependencies of dormant ovarian cancer cells, we utilized a suspension culture system that encourages cellular aggregation, leading to non-proliferative spheroids that resemble dormant aggregates found in ovarian cancer patient ascites (7, 10, 13, 33). **Fig. 1A** illustrates withdrawal from cellular proliferation by HGSOC cells in suspension as evidenced by reduced expression of Ki67 and increased expression of the quiescence marker p130 (34). Furthermore, phospho-western blots for ERK demonstrate reduced but not absent phosphorylation, and phospho-p38 either remains consistent or is increased. Reduced phospho-ERK and elevated phospho-p38 are indicative of the characteristic signaling switch reported to underlie most paradigms of cancer cell dormancy (5). These findings are relatively consistent among the three HGSOC cell lines; iOvCa147, OVCAR8, TOV1946 selected for use in our screen. To discover genes required for survival in spheroids, we utilized a new approach called GO-CRISPR (Guide Only CRISPR) that we developed specifically for survival gene discovery in three-dimensional spheroid cell culture conditions (**Fig. 1B**). GO-CRISPR incorporates sgRNA abundances from non-Cas9-expressing cells to control for stochastic (non-Cas9 dependent) effects that are a challenge in three-dimensional culture conditions with slow or arrested proliferation (35). We used sequencing to determine sgRNA identity and abundance in the library, initially infected cells, and in cells following 48 hours of suspension culture (**Fig. 1B**). Read count data was analyzed using a software package that we developed to analyze GO-CRISPR data (summarized in **Fig. 1 -Fig. Sup. 1A&B**). We selected genes that had a Spheroid/Adherent Enrichment Ratio (ER) less than 1 ($P<0.05$) because some genes become depleted in adherent growth before being transferred to suspension culture conditions and this

ratio selects for genes with a stronger effect in suspension when cells are not proliferating (**Fig.**

1B). To assess reliability of screen data, we investigated the classification of non-targeting

control sgRNAs. We computed a receiver operator characteristic curve (ROC) and determined

the area under the curve for each cell line (**Fig. 1 -Fig. Sup. 1C**). Greater than 95% of sgRNAs

in iOVCA147 and OVCAR8 screens were classified as non-essential, while 83% of sgRNAs in

TOV1946 cells were non-essential. In addition, we compared GO-CRISPR screen ER values

with viability data from siRNA knock down for 35 individual genes we have previously studied

in spheroid survival assays in OVCAR8 and iOVCA147 cells (**Fig. 1 -Fig. Sup. 1D**)(10). This

revealed a statistically significant correlation between the two sets of data, suggesting that genes

we previously knew supported spheroid viability (or were dispensable) were classified similarly

by our screen. These controls suggested our screen data was reliable for identifying new survival

dependencies.

We found 6,717 genes with an ER below 1 in iOvCa147 cells; and 7,637 genes in

TOV1946; and 7,640 genes in OVCAR8 cells. Among these genes; 1,382 were common to all

three cell lines (**Fig. 1D**). We performed pathway enrichment analysis on these common genes

and identified several significantly enriched pathways. These include primary and secondary

Reactome gene sets such as signal transduction, metabolism, and signaling by GPCR (**Fig. 1E**).

In addition, we identified a tertiary category called 'axon guidance,' that includes numerous

components of the Netrin signaling pathway that scored highly.

Expression of Netrin ligands and receptors are enriched in HGSOC spheroids

DYRK1A was previously discovered to support survival of ovarian cancer cells in

suspension culture (10). Since DYRK1A phosphorylates RNA polymerase II in transcription

initiation (36), we sought to determine if it regulates gene expression in spheroid survival. We

created $DYRK1A^{-/-}$ iOvCa147 cells and confirmed their deficiency for survival in suspension

(**Fig. 2 -Fig. Sup. 1A-E**). Using iOvCa147 and $DYRK1A^{-/-}$ derivatives, we performed RNA-seq

on adherent and spheroid cells. We collected spheroids following a 6-hour incubation in

suspension because overexpression of known cell cycle transcriptional targets due to DYRK1A

loss were first evident at this time point, but loss of viability due to DYRK1A deficiency had not

yet occurred (10). We compared iOvCa147 adherent cells to 6-hour suspension culture to

identify transcriptional changes that occur as these cells transitioned from adherent conditions to

suspension conditions (**Fig. 2A**). We identified 1,834 genes that were upregulated, and pathway

enrichment analysis identified many of the same categories as our CRISPR screen including

axon guidance (**Fig. 2B**). We then compared iOvCa147 spheroid cells to $DYRK1A^{-/-}$ spheroid

cells to identify mRNA expression changes that were lost through DYRK1A deficiency (**Fig.**

2C). We identified 744 genes that were downregulated and these also belonged to many of the

same pathways, including axon guidance (**Fig. 2D**).

We compared enriched pathways from these two RNA-seq analyses with those that were

identified in our GO-CRISPR screens to identify gene expression programs in spheroids whose

products are essential for cell survival. 78 Reactome pathways were commonly enriched across

the GO-CRISPR screens and transcriptomic analyses, including 78 of the 83 pathways identified

in RNA-seq from $DYRK1A^{-/-}$ spheroid cells (**Fig. 2E**). The axon guidance pathway was one of

the most enriched among these common pathways (**Fig. 2F**). These data reveal that our GO-

CRISPR screen and transcriptomic analyses converge on this highly specific tertiary Reactome

category of axon guidance alongside much broader primary or secondary gene sets. This

category includes Netrin ligands, receptors, and downstream signaling components, suggesting a

previously unappreciated role in HGSOC spheroid viability.

We separately sought to confirm elevated expression of Netrins and some of their

signaling components in HGSOC spheroid cell culture. We examined expression levels of

ligands and receptors in the three cell lines used in our GO-CRISPR screen using RT- qPCR

(**Fig. 3A-C**). The precise identities of upregulated genes varies between cell lines, however each

cell line increases expression of some Netrins, at least one UNC5, and at least one of DCC,

DSCAM, and Neogenin. This suggests that the Netrin pathway components are expressed in

HGSOC cells may play a previously unappreciated role in dormancy and disease progression.

Since Netrin signaling has not been previously associated with dormancy we examined

expression of Netrin-1 and -3 ligands in spheroids isolated directly from patient ascites by

immunohistochemistry. First, we assessed proliferative markers in HGSOC tumor tissue and

spheroids by staining for Ki67 and p130. As expected, solid HGSOC tumors display uniform

Ki67 positivity and relative absence of p130 (**Fig. 4A-B**). Conversely, serial sections of

spheroids demonstrated rare Ki67 nuclear staining and abundant and uniform p130 staining.

This indicates that cells in these patient derived spheroids are quiescent (**Fig. 4C-E**). To

investigate the presence of Netrins, we initially tested Netrin-1 and -3 antibody specificity using

spheroids generated from OVCAR8 cells bearing knock out or over expression of Netrins (**Fig. 4**

-Fig. Sup. 1A-F). Dormant patient derived spheroids were stained for Netrins and epithelial

markers Cytokeratin and EpCAM. Since HGSOC is characterized by point mutations in p53 and

its accumulation in the nucleus, sections were also stained for p53 to confirm cancer cell identity

within these spheroids by high level nuclear staining (**Fig. 4F-H**). Overall, this confirms Netrin

expression occurs in dormant HGSOC patient spheroids, precisely the context our screen

indicates that they are critical for cell survival.

Netrin signaling stimulates ERK to support HGSOC spheroid survival.

Netrin ligands and their dependence receptors are best known for their roles in axon

guidance (37), but are appreciated to regulate apoptosis in cancer (23). **Fig. 5A** illustrates

components of Netrin signaling and the frequency with which they were discovered to have an

ER of <1 in our screens. In addition, intracellular signaling molecules known to be downstream

of Netrin receptors that are shared with other enriched pathways identified in our screens, are

also illustrated. To confirm screen findings, we independently knocked out *NTN1*, *NTN3*, *NTN4*,

NTN5, the UNC5H receptor homologs (*UNC5A*, *UNC5B*, *UNC5C*, and *UNC5D*), as well as

DCC, *DSCAM*, and *NEO1* alone or in multigene combinations using lentiviral delivery of

sgRNAs and Cas9 (**Fig. 5 -Fig. Sup. 1A-C**). We assessed the effect of their loss on spheroid

survival in six HGSOC cell lines (iOvCa147, TOV1946, OVCAR3, OVCAR4, OVCAR8, and

COV318). Loss of at least one Netrin ligand resulted in reduced spheroid cell survival in five of

six cell lines (**Fig. 5B**). Notably, Netrin-3 loss showed the most pronounced effect on survival.

Cell lines such as OVCAR8 and COV318 were relatively resistant to the deletion of Netrin

pathway components as individual UNC5, *DCC*, and *DSCAM* genes rarely affected viability

(**Fig. 5B**). In these instances, combined deletions of UNC5 family members, or

DCC;DSCAM;NEO1 (DDN for short) were either inviable or highly sensitive to suspension

culture induced cell death, suggesting redundancy in this pathway in some cell lines. Other cell

lines, such as OVCAR3, were highly sensitive to the loss of multiple individual ligands and any

individual UNC5H receptor was highly sensitizing to death induction in suspension (**Fig. 5B**).

Overall, these illustrate a robust role for Netrins in survival in this dormant cell culture assay.

Importantly, loss of receptor expression does not elevate viability as expected for a dependence

receptors, nor does it induce DAPK1 dephosphorylation (**Fig. 5 -Fig. Sup. 1D**), suggesting that

Netrin receptors in HGSOC spheroids transmit positive survival signals.

To determine if Netrin dependent signaling is active in HGSOC spheroids, we stimulated

cells in suspension with recombinant Netrin-1 (**Fig. 5C**). Western blotting for phospho-ERK, as

a measure of activation, revealed stimulation of MEK-ERK signaling in each of the cell lines

tested. Because MAPK signaling is best known for its role in proliferation, we also tested Netrin-

1 stimulation of quiescent, adherent cells and compared this with stimulation in spheroid culture.

Both conditions display similar, low levels of phospho-ERK, but Netrin-1 was only able to

stimulate ERK phosphorylation in suspension (**Fig. 5D**). Importantly, stimulation of adherent

cells with an equivalent concentration of EGF induced pERK levels commonly seen in

proliferating cells. Collectively, the context and magnitude of pERK stimulation by Netrin-1 is

inconsistent with proliferative signaling, suggesting it may contribute to survival. This suggested

that Netrin is a candidate pathway to provide low level, but essential survival signals through the

MAPK pathway in dormant spheroids. We investigated the dependence of ERK phosphorylation

on UNC5H and DDN receptors and discovered that it is lower when receptors are deleted (**Fig.**

5E). Furthermore, Netrin-1 stimulation of suspension cultures of OVCAR8 cells, or derivatives

deleted for UNC5 family members or DDN receptors, indicated that Netrin receptors are

essential to activate ERK in suspension culture (**Fig. 5F**). To confirm that Netrin-MEK-ERK

signaling contributes to survival in suspension culture, we used two different MEK inhibitors,

PD184352 and Trametinib, to treat OVCAR8 suspension cultures and observed that it causes

loss of cell viability in suspension culture (**Fig. 5G**). Since we routinely use reattachment of

spheroids to adherent plastic as a test of viability, also tested viability using trypan blue dye

exclusion to confirm that Trametinib is truly killing cells in suspension and it revealed the same

conclusion (**Fig. 5H**). As expected, treatment of cells in suspension with Trametinib lead to

lower phospho-ERK levels (**Fig. 5I**). The simplest explanation of this data is that Netrins signal

through a heterodimer or multi-receptor complex containing both an UNC5 and DDN component

that provides low level stimulation for MEK and ERK specifically in dormant culture conditions.

Netrin signaling is essential for dormant spheroid survival during metastatic dissemination

To investigate the role of Netrin signaling in the context of residual disease and relapse, we

xenografted mice with control OVCAR8L cells, or UNC5 4KO cells, by intraperitoneal injection

(**Fig. 6A**). Two weeks following injection, mice were euthanized, and spheroids were obtained

by washing the abdominal cavity with PBS. Importantly, at this early stage of engraftment, solid

tumor lesions were not yet observable, indicating that we are modeling minimal residual disease

where dormancy is most relevant. Extraction of RNA from control OVCAR8 cell spheroids

recovered at this time point indicates a reduction in expression of pro-proliferative genes

compared to the same cells prior to engraftment (**Fig. 6B**). Furthermore, comparison with RT-

qPCR measurements of these same markers in suspension culture induced OVCAR8 spheroids

indicates that spheroids from abdominal washes are equivalently withdrawn from the cell cycle

(**Fig. 6, -Fig. Sup. 1**). This suggests spheroids at this stage are dormant. Furthermore, recovery

of spheroids by replating on adherent plastic indicates that deletion of UNC5 receptors to block

Netrin signaling extensively reduces survival under these circumstances (**Fig. 6C**). We also

quantitated cancer cells in these abdominal washes by qPCR using human Alu repeat detection to

count cells and this also demonstrates that cancer cell abundance is reduced in this dormant state

by loss of UNC5 receptors. Further xenografts were performed to investigate the long term

consequences of reduced survival of UNC5 4KO cells in mice. This demonstrated that mice

engrafted with UNC5 4KO cells had a considerably longer survival than control OVCAR8

recipient mice that received the same number of cells (**Fig. 6E,F**). Collectively, this

demonstrates that inhibition of Netrin signaling compromises viability and eventual disease

progression in a model of dormancy and dissemination.

Netrin overexpression is correlated with poor outcomes in HGSOC.

Genomic studies indicate that HGSOC is one of the most aneuploid cancers (38). In

addition to the frequency of copy number variants per tumor, the locations of gains and losses

are dissimilar between cases, making this a highly heterogenous disease. Evaluation of TCGA

genomic data for HGSOC indicates rare amplifications or deletions of Netrin or Netrin-receptor

genes (**Fig. 7 -Fig. Sup. 1A**). Because of redundancy of Netrins and their receptor families this

data suggests HGSOC cases with deficiency for Netrin signaling are rare. This is consistent with

our data that all six cell lines tested reveal evidence of Netrin component dependence for

survival (**Fig. 5B**). Some TCGA HGSOC cases display elevated gene expression for Netrins or

their receptors, but only rare cases display low expression (**Fig. 7 -Fig. Sup. 1B**). This suggests

that Netrin signaling in HGSOC is retained despite high copy number variation in this cancer

type and there is occasional overexpression.

In agreement with our experimental data that multiple Netrin ligands can contribute to

spheroid survival in HGSOC (**Fig. 5A&B**), sorting cases by high level expression of Netrin-1

and -3 ligands reveals a significantly shorter overall survival compared to cases with low level

expression and this trend is also evident in the cases that only overexpress Netrin-3 (**Fig. 7A**).

This genomic data supports the interpretation that Netrin signaling is retained in HGSOC and its

elevation is deleterious to patient outcome.

We investigated overexpression of Netrin-1 and -3 in spheroid survival by generating

OVCAR8 derivatives that stably overexpress these proteins (**Fig. 7B**). We then utilized these

cells in suspension culture assays for spheroid survival as before (**Fig. 7C**). This data indicates

that overexpression of either Netrin-1 or -3 increases OVCAR8 cell survival in dormant spheroid

culture conditions. Examination of fixed spheroids isolated from this experiment further

indicates that elevated survival in this experiment is explained by an increase in abundance of

individual spheroids that overexpress either Netrin-1 or -3 (**Fig. 7D**). These experiments reveal

the significance of a previously undiscovered role for Netrins in HGSOC. High level expression

correlates with poor outcome and in suspension culture, overexpression confers oncogenic

properties on HGSOC cells.

Netrin overexpression induces abdominal spread of HGSOC.

Our findings from CRISPR screens and cell culture experiments indicate that Netrin

signaling supports dormant HGSOC cell survival. Our analysis of TCGA is also predictive of a

worse outcome for patients. We sought to solidify these findings by searching for how Netrin

signaling affects HGSOC disease pathogenesis.

We utilized a xenograft assay in which OVCAR8 cells overexpressing Netrin-1 or -3

were injected into the peritoneal cavity and compared with OVCAR8 cells bearing an empty

vector as control. Mice were harvested following 35 days to analyze disease dissemination and

characterize the effect of Netrins in this model of HGSOC metastatic spread (**Fig. 8A**). Necropsy

of all animals at endpoint allowed for the identification and quantitation of tumor nodules within

the abdomen (**Fig. 8B**). Petal plots illustrate the frequency of specific organ and tissue sites

displaying tumor nodules (**Fig. 8B**). This revealed that spread to the omentum was similar across

all genotypes of xenografted cells and it occurred in all mice (**Fig. 8B**). In contrast, tumor

nodules were detected with increased abundance on the diaphragm, the liver, and mesenteries

that support the uterus (mesometrium) (**Fig. 8B**). In addition to more frequent disease spread to

these locations caused by Netrin overexpressing cells, we note that the quantity of nodules was

also increase in comparison with OVCAR8 control cells (**Fig. 8C-F**). We confirmed the source

of these lesions to be the xenografted cells using immunohistochemical staining for human

cytokeratin (**Fig. 8G,H**). Lastly, Netrin-1 and -3 histochemical staining confirms its presence in

the local tumor microenvironment (**Fig. 8G,H**). The similarity of increased tumor nodules seen

in xenografts and elevated spheroid formation and viability seen in culture suggests that Netrin

signaling improves cell viability in spheroid aggregates in vivo leading to increased disease

spread.

Discussion

In this report we investigated survival mechanisms in a model of HGSOC dormancy. We

utilized a modified CRISPR screening approach that was adapted to discover gene loss events in

largely non-proliferative conditions. We compared essential genes discovered in the screen with

gene expression increases identified by RNA-seq in dormant HGSOC cultures. This implicated

the Netrin signaling pathway in HGSOC spheroid survival. We independently validated the role

of Netrins, their receptors, and downstream intracellular targets MEK and ERK in the survival

biology of HGSOC cells in dormant culture and in vivo in xenografts. Furthermore, we reveal

that overexpression of Netrin-1 and -3 are correlated with poor clinical outcome in HGSOC and

over expression of these ligands induces elevated survival in dormant HGSOC culture conditions

and increased disease spread in a xenograft model of dissemination. Our work implicates

Netrins, their receptors, and MEK-ERK as a previously unappreciated, but critical signaling

network in HGSOC survival.

HGSOC spheroid cell biology has presented a unique challenge to cancer chemotherapy.

Dormant spheroids are most abundant in patients with late-stage disease and they contribute to

therapeutic resistance and seed metastases (39-41). Hence, there is a critical unmet need to

understand HGSOC spheroid dependencies and target them therapeutically. Our data suggests

that Netrin signaling is a universally active mechanism in spheroid cell survival. HGSOC is

characterized by extensive genomic rearrangements and copy number variants (38), and we

speculate that Netrin signaling is retained because of the extensive redundancy of components of

this pathway. Netrin-1 and -3 can both contribute to survival signaling and knock out of three or

four different Netrin receptors is necessary to compromise Netrin signaling in survival. A key

goal for the future is to identify Netrin-1 or -3 overexpression patients and compare their specific

disease progression patterns with our xenograft model. This will be most informative of the

effects of Netrin ligand overexpression in scenarios of dormancy, metastasis, and chemotherapy

resistance.

This study demonstrates that Netrins function to stimulate UNC5H and fibronectin repeat

containing receptors such as DCC to activate survival signals in HGSOC spheroids. This is

distinct from the known role of UNC5H and DCC as dependence receptors where Netrin inhibits

their pro-apoptotic signaling (25). Our data does not show elevated survival of spheroid cells in

response to deletions of receptors, even in instances where families of Netrin receptors are co-

deleted. Consistent with this observation, over expression of Netrin-1 or -3 in OVCAR8 cells

stimulates higher level survival in suspension. In colon cancer, deletion of DCC confers a

survival advantage and underscores the dependence receptor concept (42, 43). In HGSOC DCC

or UNC5H deletions are relatively rare (38). This suggests that ovarian cancer cells utilize Netrin

signals for positive survival signaling instead of to neutralize inherent death signals from

dependence receptors. Our data suggests that Netrin activates MEK and ERK to support cell

survival indicating that targeting ligands, receptors or downstream at MEK are all possible

approaches to inhibit this survival pathway. Netrin-1 inhibition in solid tumor xenograft models

of melanoma have been shown to be effective in combination with chemotherapy (44), but its

effects on a dormant scenario are unknown. Identifying agents that cooperate with standard

HGSOC chemotherapeutic agents such as carboplatin would offer an attractive strategy to

eradicate dormant cells while combating progressive disease. Alternatively, a novel approach to

targeting dormancy discovered in this study is, ironically, MEK inhibition. While MAPK

signaling in cancer is universally known for its role in proliferation (45), and MEK inhibitors are

used to block proliferation (46, 47), it is noteworthy that Netrin stimulation of MAPK signaling

is reported to provide survival signals and to guide axon outgrowth in a non-proliferative context

(26, 48). Dormancy is characterized by low phospho-ERK levels (3, 6), indicating low MEK

activity, however low levels of RAS-MAPK signaling are known to provide survival signals

(45). A challenge with deploying MEK inhibitors as an anti-dormancy setting in HGSOC therapy

is its incompatibility with standard chemotherapies that rely on cell proliferation. Overall, our

study aimed to identify potential new therapeutic targets in dormancy and Netrins and MEK are

potential new candidates for dormant HGSOC disease.

Current literature offers little to connect Netrins with HGSOC beyond the observation

that *NTN1* expression is increased in tumors compared to benign lesions (49). Similarly,

dormancy literature does not include a role for Netrins (3-6, 50). However, we expect that Netrin

signaling may play an unappreciated role in dormancy in other cancer disease site paradigms. As

stated in the introduction, dormancy is characterized by acquisition of stem like properties, EMT,

and in most paradigms it involves rare cells establishing themselves in perivascular or bone

marrow niches and Netrins have also been implicated in these processes (51-55). Inhibition of

Netrin-1 with NP137 and treatment with decitabine has been shown to reduce lung seeding

suggesting that Netrins are relevant in a metastatic scenario that may include dormant

intermediate steps in dissemination (56). Furthermore, Netrin signaling through UNC5B is

known to direct vascular branching (57). This suggests that Netrins are available in the

perivascular niche where dormant cells often reside. Lastly, Netrin-1 has been shown to be a

component of the bone marrow niche that suppresses hematopoietic stem cell proliferation (51).

While not a cancer dormancy scenario, it indicates that Netrins support some solitary dormant

cell properties and are expressed by resident cells in the appropriate dormant cell niches beyond

just HGSOC. For these reasons we expect Netrins will have a significant role in tumor dormancy

beyond ovarian cancer.

Methods

Cell Lines and Engineering Cas9+ cells: High-grade serous ovarian cancer (HGSOC) cell lines OVCAR3, OVCAR4, OVCAR8, COV318, TOV1946 and iOvCa147 were used in this study. The iOvCa147 cells have previously been characterized and reported (58). All cell lines were verified by STR analysis. iOvCa147, OVCAR8, and TOV1946 cells were engineered to express Cas9 using pLentiCas9-Blast and clones were isolated. High Cas9 editing efficiency was determined by viability studies using sgRNAs targeting selected fitness genes (PSMD1, PSMD2, EIF3D) and a non-targeting control (LacZ) as previously described (59).

Lentiviral sgRNA library preparation: HEK293T cells were transfected with the combined A and B components of the GeCKO v2 whole genome library (123,411 sgRNAs in total) along with plasmids encoding lentiviral packaging proteins (60). Media was collected 2-3 days later, cells and debris were pelleted by centrifugation, and supernatant containing viral particles was filtered through a 0.45 µM filter and stored at −80°C with 1.1 g/100 mL BSA.

GO-CRISPR screens in iOvCa147, OVCAR8, and TOV1946 cells: Cas9-positive and Cas9-negative cells were separately transduced with lentiviruses at a multiplicity of infection of 0.3 and with a predicted library coverage of >1000-fold. Cells were selected in puromycin and maintained in complete media (DMEM/F12 with 10% FBS, 1% pen-strep-glutamine), containing puromycin in all following steps. More than 10^9 cells were collected and split into three groups consisting of approximately 3.0×10^8 cells. Triplicate samples of 6.2×10^7 cells were saved for sgRNA sequence quantitation (T_0). The remaining cells were seeded at 2.0×10^6 cells/mL in 20 x 10 cm ULA plates. Following 2 days of culture, media containing spheroids was transferred to 10 x 15 cm adherent tissue culture plates. The next day unattached spheroid cells were collected and re-plated onto additional 15 cm plates. This process was repeated for a total of 5 days to

maximally capture viable cells. Attached cells were collected and pooled for DNA extraction

(T_f). Genomic DNA was extracted using the QIAmp DNA Blood Maxi Kit (Qiagen, #51194).

PCR amplification and barcoding performed as described (Hart paper, Cell, 2015) using

NEBNext Q5 Ultra mastermix (NEB, # MO544). PCR products were extracted from agarose

using a Monarch DNA Gel Extraction Kit (NEB, # T1020), quantitated and next generation

sequencing was performed using an Illumina NextSeq 75 cycle High Output kit on an Illumina

NextSeq platform.

TRACS analysis: The library reference file containing a list of all sgRNAs and their sequences

(CSV file), raw reads for the pooled sgRNA library (FASTQ files (L0) and raw reads (FASTQ

files) for all T_0 and T_f timepoints and replicates for Cas9-positive and Cas9-negative cells were

loaded into TRACS (https://github.com/developerpiru/TRACS)(61). TRACS then automatically

trimmed the reads using Cutadapt (v1.15), built a Bowtie2 (v2.3.4.1) index and aligned the reads

using Samtools (v1.7). TRACS used the MAGeCK read count function (v0.5.6) to generate read

counts and incremented all reads by 1 to prevent zero counts and division-by-zero errors. The

TRACS algorithm was then run using this read count file to determine the Library Enrichment

Score (ES), Initial ES, Final ES and the Enrichment Ratio (ER) for each gene. VisualizeTRACS

was used to generate graphs of Initial and Final ratios.

Pathway enrichment analyses: For GO-CRISPR screens and multi-omics comparisons, we used

the filtered list of genes obtained using TRACS and performed gene ontology and pathway

enrichment analysis using the ConsensusPathDB enrichment analysis test (Release 34

(15.01.2019), http://cpdb.molgen.mpg.de/) for top-ranked genes of interest. p_{adj} values and ER

values for each gene were used as inputs. The minimum required genes for enrichment was set to

45 and the FDR-corrected p_{adj} value cutoff was set to < 0.01. The Reactome pathway dataset was

used as the reference. For each identified pathway, ConsensusPathDB provides the number of

enriched genes and a q value (p_{adj}) for the enrichment. Bar plots were generated in R 3.6.2 using

these values to depict the significant pathways identified.

Generating DYRK1A knockout cells: A double cutting CRISPR/Cas9 approach with a pair of

sgRNAs (sgRNA A and B) was used to completely excise exon 2 (322 bp region) of *DYRK1A*

using a px458 vector (Addgene #48138) that was modified to express the full CMV promoter.

PCR primers that flank the targeted region of *DYRK1A* were used to verify deletion. Single-cell

clones were generated by liming dilutions and evaluated for DYRK1A status by PCR, western

blot, and sequencing. See Supplemental Table 1 for sgRNA and PCR primer sequences.

DYRK1A IP kinase assay: Whole cell lysates from adherent parental iOvCa147 cells and

DYRK1A$^{-/-}$ cells were extracted using complete RIPA buffer and incubated overnight with

DYRK1A antibody (Cell Signaling Technology anti-DYRK1A rabbit antibody #8765). Samples

were then washed with buffer and Dynabeads (Thermo Fisher Scientific Dynabeads Protein G

#10003D) were added for 2 hours. Samples were then washed with buffer and recombinant Tau

protein (Sigma recombinant Tau protein #T0576) and ATP (Sigma #A1852) were added and

samples were incubated for 30 minutes at 37°C. 5x SDS was then added and samples were

resolved by SDS-PAGE. Antibodies used for western blotting were phosphospecific Tau

antibody (Cell Signaling Technology phospho-Tau Ser-404 rabbit antibody #20194) and Tau

protein (Cell Signaling Technology anti-Tau rabbit antibody #46687).

RNA preparation and RNA-sequencing: Total RNA was collected from parental iOvCa147 cells

and *DYRK1A$^{-/-}$* cells from 24-hour adherent or 6-hour spheroid conditions using the Monarch

Total RNA Miniprep Kit (NEB #T2010S) as described above. RNA was collected in three

replicates for each condition (adherent and spheroids). Samples were quantitated using Qubit 2.0

Fluorometer (Thermo Fisher Scientific) and quality control analysis using Agilent 2100

Bioanalyzer (Agilent Technologies #G2939BA) and RNA 6000 Nano Kit (Agilent Technologies

#5067-1511). Ribosomal RNA removal and library preparation was performed using ScriptSeq

Complete Gold Kit (Illumina #BEP1206). High-throughput sequencing was performed on an

Illumina NextSeq 500 platform (mid-output, 150-cycle kit).

RNA-seq analyses: Raw FASTQ data was downloaded from Illumina BaseSpace. Reads were

aligned using STAR 2.6.1a(62) to the human genome

(Homo_sapiens.GRCh38.dna.primary_assembly.fa; sjdbGTFfile:

Homo_sapiens.GRCh38.92.gtf) to generate read counts. DESeq2 was run with BEAVR(63) and

used to analyze read counts (settings: False discovery rate (FDR): 10%; drop genes with less

than 1 reads) to identify differentially expressed genes ($\log_2$ fold change > 1 for upregulated

genes; or $\log_2$ fold change < -1 for downregulated genes; p_{adj} < 0.05) for the following

comparisons: parental iOvCa147 adherent cells vs. parental iOvCa147 spheroid cells; or parental

iOvCa147 spheroid cells vs. *DYRK1A*$^{-/-}$ spheroid cells. For RNA-seq, we formed pathway

analyses in BEAVR and downloaded the pathway enrichment table to construct bar plots and

determine overlapping pathways using R 3.6.2.

Western blots: Adherent cells were washed in 1x PBS and lysed on the plate by the addition of

complete RIPA buffer with protease (Cell Signaling Technology #5871S) and phosphatase (Cell

Signaling Technologies #5870S) inhibitors and incubated for 5-10 minutes on ice. Spheres were

collected by gentle centrifugation at 300 rpm, washed gently in ice cold PBS, centrifuged again

and then resuspended in RIPA buffer as above. Samples were then centrifuged at 14,000 RPM at

4°C for 10 minutes. The supernatant liquid for each sample was collected and the pellets re-

extracted with RIPA containing the above inhibitor cocktails. Following centrifugation, the

supernatants for each treatment were collected and pooled. Protein was determined by Bradford

assay. To assess the effect of Netrin 1 on downstream signaling, spheres incubated under non

adherent conditions for 5 days were challenged with the addition of 500 ng/ml human

recombinant Netrin-1 (R&D Systems #6419-N1-025) for a period of 20 minutes before collection.

Lysates were mixed with 5x SDS loading dye buffer and resolved using standard SDS-PAGE

protocols. Antibodies used for blotting were p42/44 MAPK (Erk1/2) (Cell Signaling Technology

rabbit antibody #4695), Phospho-p44/42 MAPK(Erk1/2) T202/Y204 (Cell Signaling

Technology; rabbit antibody #4370), p38 MAPK (Cell Signaling Technology; rabbit antibody

#9212), p38 MAPK T180/Y182 (Cell Signaling Technology; rabbit antibody #9211),

p130(RBL2, Santa Cruz Biotechnology, rabbit antibody sc-317 discontinued), Ki67 (Abcam

rabbit antibody, # ab16667), DYRK1A (Cell Signaling Technology anti-DYRK1A rabbit

antibody #8765), Netrin-1 (Abcam rabbit antibody #ab126729), Netrin-3 (Abcam rabbit antibody

#ab185200), α-tubulin (Cell Signaling Technology anti-αTubulin rabbit antibody #2125) and β-

actin (Millipore Sigma rabbit antibody #A2066).

RT-qPCR analysis of gene expression: Cells from a minimum of three biological replicates were

triturated off the plate using Versene (0.5mM EDTA in PBS), collected, counted and 10^6 cells

plated onto Ultra-Low Attachment cell culture dishes (Corning) to induce sphere formation. The

spheres were collected after 24 h and RNA extracted using the Monarch Total RNA Miniprep

Kit (New England Biolabs #T2010S). RNA was extracted from adherent cells directly from the

plate. Extracted RNA was quantitated using a NanoDrop and stored at -80º until used. First

strand cDNA was generated from isolated RNA using the iScript cDNA Synthesis Kit (Bio-Rad,

#1708890) and RT-qPCR performed on the resulting cDNA (diluted 5-fold) using the iQ SYBR

Green Supermix (Bio-Rad, #1708882) and following the manufactures instructions. Reactions

were performed in 0.2ml non-skirted low profile 96 well PCR plates (Thermo Scientific, #AD-

0700) using a CFX Connect Real Time System thermo-cycler (Bio-Rad).

TCGA patient data analyses: We used the cBioPortal for Cancer Genomics

(https://www.cbioportal.org)(64) to analyze Netrin pathway alterations in Ovarian Serous

Cystadenocarcinoma patients (TCGA, PanCancer Atlas) for genomic alterations and mRNA

expression. OncoPrints were generated for mRNA expression (mean z-score threshold ± 1.5),

and mutations and putative copy number alterations for the following genes: *NTN1, NTN3,*

NTN4, NTN5, UNC5A, UNC5B, UNC5C, UNC5D, DCC, DSCAM. Survival analysis of *NTN1,*

and *-3* utilized patients from TCGA, PanCancer Atlas dataset and RNA-seq expression (high

expression z-score threshold > 1.2 and low is below 1.2).

Generation of gene knockout cell lines: Gibson Assembly (NEB #E2611) was used to clone

sgRNAs sequences derived from the GeCKO v2 library per gene of interest (*NTN1, NTN3,*

NTN4, NTN5, DSCAM, DCC, NEO1, UNC5A, UNC5B, UNC5C, UNC5D, and *EGFP* control

(See Supplemental Table 1 for sgRNA sequences for each)) into lentiCRISPR v2 (Addgene

#52961). For individual gene knockouts pooled libraries of up to four sgRNAs were generated.

For multiple gene knockouts pooled libraries of two sgRNAs/gene were constructed. To each 20-

nucleotide guide (Supplemental table I) was appended the 5' sequence, 5'-

TATCTTGTGGAAAGGACGAAACACC-3' and the 3' sequence, 5'-

GTTTTAGAGCTAGAAATAGCAAGTTAAAAT-3' (65). A 100 bp fragment was then

generated using the forward primer, 5'-

GGCTTTATATATCTTGTGGAAAGGACGAAACACCG-3', and reverse primer, 5'-

CTAGCCTTATTTTAACTTGCTATTTCTAGCTCTAAAAC-3'. The resulting gel purified

fragment was quantitated by Qubit and cloned into the lentiviral vector LentiCRISPR-V2

previously digested with BsmBI (New England Biolabs, # R0739) as described (65). To generate

virus particles, HEK293T cells were transfected with the assembled plasmid along with plasmids

encoding lentiviral packaging proteins. Media was collected after 2-3 days and any cells or

debris were pelleted by centrifugation at 500 x g. Supernatant containing viral particles was

filtered through a low protein binding 0.45 µM filter. For each knockout, it was important to

ensure that each target cell was transduced with a full complement of guides. To ensure this, the

cells were transduced at relatively high MOI. Briefly, 25,000 cells were plated and transduced

with fresh HEK293T conditioned media containing virus. While we did not titer the viral

preparations for these experiments, previous experience indicated that we could generate

sufficiently high virus titers using our protocol. Under these conditions, we found a lack of

puromycin sensitive cells following 2-3 days in media containing 2-4 µg/ml puromycin

indicating that every cell received at least one virus particle. Multi-gene knock outs were created

through multiple rounds of infection with the relevant culture supernatants followed by one

round of puromycin selection.

Generation of gene overexpression cell lines: Plasmids encoding Netrin1 and Netrin3 were

obtained from Addgene (Ntn1-Fc-His, #72104 and Ntn3-Fc-His, #72105). Netrin1 was PCR

amplified using the forward primer 5'-

GCTATCGATATCCCAAACGCCACCATGATGCGCGCTGTGTGGGAGGCGCTG-3' and

reverse 5'-GCTATCTCTAGAGGCCTTCTTGCACTTGCCCTTCTTCTCCCG-3' and cloned

into the EcoRV/XbaI sites of pcDNA3.1/myc-HisA. For lentiviral expression, pcDNA NTN1

was amplified using the primers, forward 5'-

CACTGTAGATCTCCAAACGCCACCATGATGCGCGCTGTGTGGGAGGCGCTG-3' and,

reverse 5'-ACGCGTGAATTCTTATCAACCGGTATGCATATTCAGATCCTCTTCTGAGAT-

3' while NTN3 was amplified from pcDNA NTN3 using, forward 5'-

CTCGAGAGATCTGCGGCCGCCACCATGCCTGGCTGGCCCTGG-3' and the same reverse

primer as for NTN1. In this way both constructs carry a C-terminal Myc tag. Both cDNA's were

inserted into the BamHI/EcoRI sites of the lentiviral expression vector FUtdTW (Addgene

#22478). Construction of lentiviral particle and transduction into the target cells was as described

above.

Spheroid formation and reattachment viability assays: For each targeted gene or cDNA

expressed in iOvCa147, OVCAR3, OVCAR4, OVCAR8, COV318, or TOV1946 cells, spheroid

viability was assayed as follows: Three to ten biological replicate cultures of knockout cells were

cultured for 72 hours in suspension conditions using ULA plasticware (2×10^6 cells per well) to

allow spheroid formation. Addition of therapeutics (Tramatinib, Selleckchem, #S4484) was

made at the time of plating. Spheroids were collected and transferred directly to standard

plasticware to facilitate reattachment for 24 hours, or dissociated and mixed with trypan blue.

Reattached cells were fixed in 25% methanol in PBS for 3 minutes. Fixed cells were incubated

for 30 minutes with shaking in 0.5% crystal violet, 25% methanol in PBS. Plates were carefully

immersed in water to remove residual crystal violet and destained in 10% acetic acid in 1x PBS

for 1 hour with shaking to extract crystal violet from cells. Absorbance of crystal violet was

measured at 590 nm using a microplate reader (Perkin Elmer Wallac 1420).

Xenografts: All animal experiments were approved by the Animal Care Committee at Western

University (protocol #2020-039). ARRIVE principles were followed in carrying out xenograft

experiments. Female NOD-SCID mice were purchased from JAX (#001303). They were

randomly selected prior to being engrafted between six and eight weeks of age with 4 x10^6

OVCAR8 control or modified derivatives. Five to ten mice were housed for 14 days, 35 days, or

until humane endpoint for each OVCAR8 or modified derivative. Following euthanasia all mice

were subjected to a necropsy to analyze disease spread. Tumor lesions were photographed and

quantitated by individuals unaware of the genotypes. Abdominal organs and associated tumor

lesions were fixed in 10% formalin overnight and transferred to 70% ethanol and later

embedded, sectioned and stained as described below. In cases where abdominal washes were

performed, an incision was made in the peritoneal membrane and PBS was injected, the

abdominal region was agitated and PBS was withdrawn using a P1000 pipet tip and aliquoted for

downstream analysis in cell culture and qPCR.

Patient derived spheroids: Patient derived spheroids were isolated from ascites fluid obtained

from debulking surgical procedures. These HGSOC samples were obtained with permission and

institutional research ethics oversight (Project ID #115904). Spheroids were prepared by filtering

the ascites fluid through a 100μm mesh filter and washing the captured spheroids from the filter

with PBS. The spheres were collected by gravity on ice in PBS, washed twice with ice cold PBS

and allowed to settle on ice. Spheres were then fixed for 20 minutes in 10% formalin, washed,

and resuspended in PBS before being embedded in paraffin.

Histology and immunohistochemistry: Fixed organs from xenografts or spheroids were

embedded in paraffin, sectioned, and stained with H&E using standard methods. For

immunohistochemistry, sections were deparaffinized by standard procedures and antigen

retrieval was performed in 10mM citrate, pH6.0. The sections were blocked with 10% goat

serum/3% BSA/0.3% Triton X-100 and stained with antibodies to Netrin-1 (Abcam #ab126729,

1:200), Cytokeratin (human specific cytokeratin 7 and 8, ZETA Corporation #Z2018ML, 1:200),

EpCAM (Cell Signaling Technology, #14452, 1:250), p53 (Cell Signaling Technologies, #2527,

1:125), Netrin-3 (1:250), Ki67 (Cell Signaling Technology, #12202, 1:400) and p130 (RBL2,

Santa Cruz Biotechnology, SC-317, 1:150). Netrin-3 antibodies were generated using immunized

rabbits and their generation and purification will be published elsewhere. Primary antibodies

were followed with a biotinylated anti-rabbit (Vector Labs, # BA-1000) or anti-mouse (Jackson

ImmunoResearch, #115-067-003) IgG as a secondary antibody (1:500). Proteins were visualized

using DAB chemistry (ImmPACT DAB, Vector Labs, #SK-4105).

Statistical analyses: Specific statistical tests used are indicated in the figure legends for each

experiment. Analysis was done using GraphPad Prism 6.

Materials availability

All previously unpublished reagents created in this study are available on request from the corresponding author.

Data availability

All sequencing data have been deposited in GEO under accession codes GSE190294 and GSE150246. Source data for graphs and original blots are available in source data files, or from public databases as described in the methods section.

References

1. Giancotti FG. Mechanisms governing metastatic dormancy and reactivation. Cell. 2013;155(4):750-64.

2. Massague J, Ganesh K. Metastasis-Initiating Cells and Ecosystems. Cancer Discov. 2021;11(4):971-94.

3. Sosa MS, Bragado P, Aguirre-Ghiso JA. Mechanisms of disseminated cancer cell dormancy: an awakening field. Nat Rev Cancer. 2014;14(9):611-22.

4. Phan TG, Croucher PI. The dormant cancer cell life cycle. Nat Rev Cancer. 2020;20(7):398-411.

5. Goddard ET, Bozic I, Riddell SR, Ghajar CM. Dormant tumour cells, their niches and the influence of immunity. Nat Cell Biol. 2018;20(11):1240-9.

6. Yeh AC, Ramaswamy S. Mechanisms of Cancer Cell Dormancy--Another Hallmark of Cancer? Cancer Res. 2015;75(23):5014-22.

7. Correa RJ, Peart T, Valdes YR, DiMattia GE, Shepherd TG. Modulation of AKT activity is associated with reversible dormancy in ascites-derived epithelial ovarian cancer spheroids. Carcinogenesis. 2012;33(1):49-58.

8. Lengyel E, Burdette JE, Kenny HA, Matei D, Pilrose J, Haluska P, et al. Epithelial ovarian cancer experimental models. Oncogene. 2014;33(28):3619-33.

9. Hu J, Nakhla H, Friedman E. Transient arrest in a quiescent state allows ovarian cancer cells to survive suboptimal growth conditions and is mediated by both Mirk/dyrk1b and p130/RB2. Int J Cancer. 2011;129(2):307-18.

10. MacDonald J, Ramos-Valdes Y, Perampalam P, Litovchick L, DiMattia GE, Dick FA. A Systematic Analysis of Negative Growth Control Implicates the DREAM Complex in Cancer Cell Dormancy. Mol Cancer Res. 2017;15(4):371-81.

11. Correa RJ, Valdes YR, Peart TM, Fazio EN, Bertrand M, McGee J, et al. Combination of AKT inhibition with autophagy blockade effectively reduces ascites-derived ovarian cancer cell viability. Carcinogenesis. 2014;35(9):1951-61.

12. Correa RJ, Valdes YR, Shepherd TG, DiMattia GE. Beclin-1 expression is retained in high-grade serous ovarian cancer yet is not essential for autophagy induction in vitro. J Ovarian Res. 2015;8:52.

13. Shepherd TG, Dick FA. Principles of dormancy evident in high-grade serous ovarian cancer. Cell Div. 2022;17(1):2.

14. Lheureux S, Gourley C, Vergote I, Oza AM. Epithelial ovarian cancer. Lancet. 2019;393(10177):1240-53.

15. Lengyel E. Ovarian cancer development and metastasis. Am J Pathol. 2010;177(3):1053-64.

16. Bowtell DD, Bohm S, Ahmed AA, Aspuria PJ, Bast RC, Jr., Beral V, et al. Rethinking ovarian cancer II: reducing mortality from high-grade serous ovarian cancer. Nat Rev Cancer. 2015;15(11):668-79.

17. Peart T, Ramos Valdes Y, Correa RJ, Fazio E, Bertrand M, McGee J, et al. Intact LKB1 activity is required for survival of dormant ovarian cancer spheroids. Oncotarget. 2015;6(26):22424-38.

18. Buensuceso A, Ramos-Valdes Y, DiMattia GE, Shepherd TG. AMPK-Independent LKB1 Activity Is Required for Efficient Epithelial Ovarian Cancer Metastasis. Mol Cancer Res. 2019.

19.   Chen MW, Yang ST, Chien MH, Hua KT, Wu CJ, Hsiao SM, et al. The STAT3-miRNA-
92-Wnt Signaling Pathway Regulates Spheroid Formation and Malignant Progression in Ovarian
Cancer. Cancer Res. 2017;77(8):1955-67.
20.   Diaz Osterman CJ, Ozmadenci D, Kleinschmidt EG, Taylor KN, Barrie AM, Jiang S, et
al. FAK activity sustains intrinsic and acquired ovarian cancer resistance to platinum
chemotherapy. Elife. 2019;8.
21.   Rajasekharan S, Kennedy TE. The netrin protein family. Genome Biol. 2009;10(9):239.
22.   Kryza D, Wischhusen J, Richaud M, Hervieu M, Sidi Boumedine J, Delcros JG, et al.
From netrin-1-targeted SPECT/CT to internal radiotherapy for management of advanced solid
tumors. EMBO Mol Med. 2023;15(4):e16732.
23.   Mehlen P, Delloye-Bourgeois C, Chedotal A. Novel roles for Slits and netrins: axon
guidance cues as anticancer targets? Nat Rev Cancer. 2011;11(3):188-97.
24.   Arakawa H. Netrin-1 and its receptors in tumorigenesis. Nat Rev Cancer.
2004;4(12):978-87.
25.   Brisset M, Grandin M, Bernet A, Mehlen P, Hollande F. Dependence receptors: new
targets for cancer therapy. EMBO Mol Med. 2021;13(11):e14495.
26.   Forcet C, Stein E, Pays L, Corset V, Llambi F, Tessier-Lavigne M, et al. Netrin-1-
mediated axon outgrowth requires deleted in colorectal cancer-dependent MAPK activation.
Nature. 2002;417(6887):443-7.
27.   Larrieu-Lahargue F, Welm AL, Thomas KR, Li DY. Netrin-4 induces lymphangiogenesis
in vivo. Blood. 2010;115(26):5418-26.
28.   Yin K, Wang L, Zhang X, He Z, Xia Y, Xu J, et al. Netrin-1 promotes gastric cancer cell
proliferation and invasion via the receptor neogenin through PI3K/AKT signaling pathway.
Oncotarget. 2017;8(31):51177-89.
29.   Llambi F, Lourenco FC, Gozuacik D, Guix C, Pays L, Del Rio G, et al. The dependence
receptor UNC5H2 mediates apoptosis through DAP-kinase. EMBO J. 2005;24(6):1192-201.
30.   Castets M, Coissieux MM, Delloye-Bourgeois C, Bernard L, Delcros JG, Bernet A, et al.
Inhibition of endothelial cell apoptosis by netrin-1 during angiogenesis. Dev Cell.
2009;16(4):614-20.
31.   Bruikman CS, Zhang H, Kemper AM, van Gils JM. Netrin Family: Role for Protein
Isoforms in Cancer. J Nucleic Acids. 2019;2019:3947123.
32.   Jiang S, Richaud M, Vieugue P, Rama N, Delcros JG, Siouda M, et al. Targeting netrin-3
in small cell lung cancer and neuroblastoma. EMBO Mol Med. 2021;13(4):e12878.
33.   Shepherd TG, Theriault BL, Campbell EJ, Nachtigal MW. Primary culture of ovarian
surface epithelial cells and ascites-derived ovarian cancer cells from patients. Nat Protoc.
2006;1(6):2643-9.
34.   Litovchick L, Chestukhin A, DeCaprio JA. Glycogen synthase kinase 3 phosphorylates
RBL2/p130 during quiescence. Molecular and cellular biology. 2004;24(20):8970-80.
35.   Zanoni M, Piccinini F, Arienti C, Zamagni A, Santi S, Polico R, et al. 3D tumor spheroid
models for in vitro therapeutic screening: a systematic approach to enhance the biological
relevance of data obtained. Sci Rep. 2016;6:19103.
36.   Di Vona C, Bezdan D, Islam AB, Salichs E, Lopez-Bigas N, Ossowski S, et al.
Chromatin-wide profiling of DYRK1A reveals a role as a gene-specific RNA polymerase II
CTD kinase. Mol Cell. 2015;57(3):506-20.
37.   Meijers R, Smock RG, Zhang Y, Wang JH. Netrin Synergizes Signaling and Adhesion
through DCC. Trends Biochem Sci. 2020;45(1):6-12.

38.   Network CGA. Integrated genomic analyses of ovarian carcinoma. Nature.
2011;474(7353):609-15.
39.   Ahmed N, Stenvers KL. Getting to know ovarian cancer ascites: opportunities for
targeted therapy-based translational research. Front Oncol. 2013;3:256.
40.   Narod S. Can advanced-stage ovarian cancer be cured? Nat Rev Clin Oncol.
2016;13(4):255-61.
41.   Keyvani V, Farshchian M, Esmaeili SA, Yari H, Moghbeli M, Nezhad SK, et al. Ovarian
cancer stem cells and targeted therapy. J Ovarian Res. 2019;12(1):120.
42.   Fearon ER, Cho KR, Nigro JM, Kern SE, Simons JW, Ruppert JM, et al. Identification of
a chromosome 18q gene that is altered in colorectal cancers. Science. 1990;247(4938):49-56.
43.   Mehlen P, Rabizadeh S, Snipas SJ, Assa-Munt N, Salvesen GS, Bredesen DE. The DCC
gene product induces apoptosis by a mechanism requiring receptor proteolysis. Nature.
1998;395(6704):801-4.
44.   Boussouar A, Tortereau A, Manceau A, Paradisi A, Gadot N, Vial J, et al. Netrin-1 and
Its Receptor DCC Are Causally Implicated in Melanoma Progression. Cancer Res.
2020;80(4):747-56.
45.   Downward J. Targeting RAS signalling pathways in cancer therapy. Nat Rev Cancer.
2003;3(1):11-22.
46.   Barbosa R, Acevedo LA, Marmorstein R. The MEK/ERK Network as a Therapeutic
Target in Human Cancer. Mol Cancer Res. 2021;19(3):361-74.
47.   Flaherty KT, Infante JR, Daud A, Gonzalez R, Kefford RF, Sosman J, et al. Combined
BRAF and MEK inhibition in melanoma with BRAF V600 mutations. N Engl J Med.
2012;367(18):1694-703.
48.   Chen J, Du H, Zhang Y, Chen H, Zheng M, Lin P, et al. Netrin-1 Prevents Rat Primary
Cortical Neurons from Apoptosis via the DCC/ERK Pathway. Front Cell Neurosci. 2017;11:387.
49.   Papanastasiou AD, Pampalakis G, Katsaros D, Sotiropoulou G. Netrin-1 overexpression
is predictive of ovarian malignancies. Oncotarget. 2011;2(5):363-7.
50.   Recasens A, Munoz L. Targeting Cancer Cell Dormancy. Trends Pharmacol Sci.
2019;40(2):128-41.
51.   Renders S, Svendsen AF, Panten J, Rama N, Maryanovich M, Sommerkamp P, et al.
Niche derived netrin-1 regulates hematopoietic stem cell dormancy via its receptor neogenin-1.
Nat Commun. 2021;12(1):608.
52.   Ozmadenci D, Feraud O, Markossian S, Kress E, Ducarouge B, Gibert B, et al. Netrin-1
regulates somatic cell reprogramming and pluripotency maintenance. Nat Commun.
2015;6:7398.
53.   Lengrand J, Pastushenko I, Vanuytven S, Song Y, Venet D, Sarate RM, et al.
Pharmacological targeting of netrin-1 inhibits EMT in cancer. Nature. 2023;620(7973):402-8.
54.   Ducarouge B, Redavid AR, Victoor C, Chira R, Fonseca A, Hervieu M, et al. Netrin-1
blockade inhibits tumor associated Myeloid-derived suppressor cells, cancer stemness and
alleviates resistance to chemotherapy and immune checkpoint inhibitor. Cell Death Differ. 2023.
55.   Cassier PA, Navaridas R, Bellina M, Rama N, Ducarouge B, Hernandez-Vargas H, et al.
Netrin-1 blockade inhibits tumour growth and EMT features in endometrial cancer. Nature.
2023;620(7973):409-16.
56.   Grandin M, Meier M, Delcros JG, Nikodemus D, Reuten R, Patel TR, et al. Structural
Decoding of the Netrin-1/UNC5 Interaction and its Therapeutical Implications in Cancers.
Cancer Cell. 2016;29(2):173-85.

57.   Lu X, Le Noble F, Yuan L, Jiang Q, De Lafarge B, Sugiyama D, et al. The netrin
receptor UNC5B mediates guidance events controlling morphogenesis of the vascular system.
Nature. 2004;432(7014):179-86.
58.   Tong JG, Valdes YR, Sivapragasam M, Barrett JW, Bell JC, Stojdl D, et al. Spatial and
temporal epithelial ovarian cancer cell heterogeneity impacts Maraba virus oncolytic potential.
BMC Cancer. 2017;17(1):594.
59.   Hart T, Chandrashekhar M, Aregger M, Steinhart Z, Brown KR, MacLeod G, et al. High-
Resolution CRISPR Screens Reveal Fitness Genes and Genotype-Specific Cancer Liabilities.
Cell. 2015;163(6):1515-26.
60.   Sanjana NE, Shalem O, Zhang F. Improved vectors and genome-wide libraries for
CRISPR screening. Nat Methods. 2014;11(8):783-4.
61.   Perampalam P, MacDonald JI, Dick FA. GO-CRISPR; a highly controlled workflow to
discover gene essentiality. bioRxiv. 2020;doi: https://doi.org/10.1101/2020.06.04.134841.
62.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.
63.   Perampalam P, Dick FA. BEAVR: a browser-based tool for the exploration and
visualization of RNA-seq data. BMC Bioinformatics. 2020;21(1):221.
64.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer
genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer
Discov. 2012;2(5):401-4.
65.   Wang T, Lander ES, Sabatini DM. Large-Scale Single Guide RNA Library Construction
and Use for CRISPR-Cas9-Based Genetic Screens. Cold Spring Harb Protoc. 2016;2016(3):pdb
top086892.

 Figure Legends

 Figure 1. GO-CRISPR screens implicate axon guidance pathways as supporting HGSOC
spheroid cell viability. A) iOvCa147, TOV1946, or OVCAR8 cells were cultured under
adherent conditions (Adh) or in suspension to induce spheroid formation (Sph). Lysates were
prepared and analyzed by western blotting for the proliferation marker Ki67 and the quiescence
marker p130, and phosphorylated and total levels of ERK and p38. Tubulin was blotted as a
loading control. **B)** Flow chart of GO-CRISPR screening used for each of iOvCa147, TOV1946,
or OVCAR8 control cells and derivatives of each that express Cas9. Cas9-positive cells (top
row) and Cas9-negative cells (bottom row) were transduced with the GeCKO v2 pooled sgRNA
library. After antibiotic selection, cells were expanded under adherent culture conditions (0
hours) before being transferred to suspension culture conditions to induce spheroid formation
and select for cell survival. After 48 hours spheroids were transferred to standard plasticware to
isolate viable cells. Red arrows indicate the relevant comparisons of sgRNA sequence abundance
that were made to analyze screen outcomes. Genes with relatively greater effect on viability in
suspension were selected by comparing their scores between adherent and suspension conditions
and considering genes with an enrichment ratio (ER) of <1. **C)** Scatter plots representing the
spheroid score on y-axis and adherent score on the x-axis calculated by TRACS for each gene in
each cell line (iOvCa147, TOV1946, OVCAR8). Colored data points represent genes with ER <
1 and $p_{adj} < 0.05$ (paired t-test). **D)** Venn diagram illustrating overlap of genes identified as
supporting cell viability in suspension culture from iOvCa147, TOV1946, and OVCAR8 cells.
E) Graph depicting enriched pathways from ConsensusPathDB using the 1382 commonly
identified genes from D. Categories are ranked by q-value. Tiers of Reactome categories are
indicated by shading.

Figure 1 - Figure Supplement 1. CRISPR screen analysis details and internal controls. A)
Flowchart of data analysis used to identify genes that are most relevant to ovarian cancer cell
survival in suspension. **B)** Mathematical formulas to define gene scores, enrichment scores
(n=guides per gene), and enrichment ratios used by TRACS. **C)** Reader-operator curves were
generated to assess TRACS categorization of 1000 non-targeting control sgRNAs in the
GeCKOv2 library for each of the three cell lines screened. **D)** Spheroid cell viability levels from
low throughput siRNA gene knock downs for 35 genes in iOvCa147 and OVCAR8 cells was
plotted against the enrichment ratio for the same genes in the same two cell lines from this
screen. These independent approaches to measure viability were compared using linear
regression.

Figure 2. Axon guidance pathway components are upregulated in iOvCa147 spheroid cells
in a DYRK1A dependent manner. A) RNA was isolated from iOvCa147 cells following
culture under adherent conditions or in suspension conditions to induce spheroid formation for 6
hours. Triplicate independent cultures were processed for RNA-seq. A volcano plot shows
differentially expressed genes in spheroid cells compared to adherent. 1,937 genes were found to
be downregulated in iOvCa147 spheroid cells ($\log_2$ fold change < 1, $p_{adj} < 0.05$ (Wald test), FDR
10%, highlighted in grey) and 1,834 genes were upregulated ($\log_2$ fold change > 1, $p_{adj} < 0.05$,
FDR 10%, highlighted in blue). **B)** Top 15 most significantly enriched pathways ($p_{adj} < 0.05$)
whose genes were upregulated in suspension culture compared to adherent in RNA-seq analysis.
C) A volcano plot showing differentially expressed genes in $DYRK1A^{-/-}$ spheroid cells compared

to iOvCa147 spheroid cells. 744 genes were found to be downregulated in *DYRK1A$^{-/-}$* spheroid
cells (log$_2$ fold change < 1, p$_{adj}$ < 0.05 (Wald test), FDR 10%, highlighted in red) and 96 genes
were upregulated (log$_2$ fold change > 1, p$_{adj}$ < 0.05 (Wald test), FDR 10%, highlighted in grey).
D) Top 15 most significantly enriched pathways that were represented by downregulated genes
in *DYRK1A$^{-/-}$* suspension culture compared to control cells in suspension. **E)** Venn diagram
depicting overlapping enriched pathways identified in GO-CRISPR screens in green; enriched
pathways identified in upregulated genes in parental iOvCa147 spheroid cells in blue; and
enriched pathways identified in downregulated genes in *DYRK1A$^{-/-}$* spheroid cells in red. 78
pathways were commonly enriched in all three datasets (shown in yellow). **F)** Top 10 most
significantly enriched pathways among the 78 identified in C.

Figure 2 - Figure Supplement 1. Generation of iOvCa147 cells deficient for DYRK1A and
loss of viability in suspension. A) Strategy to generate *DYRK1A$^{-/-}$* cells using a pair of sgRNAs
that flank exon 2. PCR For and PCR Rev primers flank exon 2 and are used to detect deletion
events. **B)** Agarose gel showing PCR products for iOVCA147 cells and putative *DYRK1A$^{-/-}$*
cells. The full length amplicon containing exon 2 was detected in parental iOvCa147 cells (1,348
bp). A smaller amplicon (1,026 bp) was detected in *DYRK1A$^{-/-}$* cells, indicting successful
excision of the 322 bp region encompassing exon 2. **C)** Western blot comparing DYRK1A
expression in iOVCA147 cells and *DYRK1A$^{-/-}$* cells. **D)** IP kinase assay to evaluate DYRK1A
activity in *DYRK1A$^{-/-}$* cells. Anti-DYRK1A antibodies or IgG was used to immunoprecipitate
from iOVCA147 and *DYRK1A$^{-/-}$* cells. Precipitates were incubated with ATP and Tau protein.
Samples were resolved by SDS-PAGE and probed with pSer404-Tau antibody. **E)** Parental
iOvCa147 or *DYRK1A$^{-/-}$* cells were incubated in suspension conditions for 24 hours, 72 hours, or
4 days to induce spheroid formation, and then re-plated in adherent conditions for 24 hours to
allow reattachment. Reattached spheroid cells were stained with crystal violet and absorbance
was quantified. Alternatively, cells in suspension were isolated and utilized for Cell TitreGLO
cell viability measurements. *DYRK1A$^{-/-}$* spheroid cells had impaired survival compared to
parental iOvCa147 spheroid cells at each time point and reattached and crystal violet stained cell
viability matched Cell TitreGLO measurements of viability (one way anova, ****P<0.0001).

Figure 3. Expression of Netrin ligands and their dependence receptors is increased in
suspension culture. A-C) RT-qPCR was performed to quantitate mRNA expression levels of
Netrin ligands and receptors in three different HGSOC cell lines. Relative expression of the
indicated transcripts is shown for suspension culture conditions compared with adherent. All
experiments were performed in at least triplicate biological replicates. Means were compared
with the same gene in adherent culture using a one way anova (*P<0.05, ****P<0.0001).

Figure 4. Netrins are expressed in dormant patient derived spheroids. Spheroids were
isolated from ascites of HGSOC patients and processed for immunohistochemical staining. **A-B)**
Serial sections from solid HGSOC tumors were stained with antibodies to p130, Ki67, or a no
primary antibody control (NP). Scale bar = 100μm for upper panels and 2mm for lower panels.
C-E) Serial spheroid sections from the indicated patients were stained with Ki67 and p130.

Sections with dense positive nuclear staining are indicated with white arrows. Scale bar =
100μm. **F-H)**. Immunohistochemical staining was performed for the indicated proteins on serial
sections of ascites derived spheroids. Omission of primary antibody was used a control for
background staining for each patient sample (NP). Scale bar = 100μm

Figure 4 - Figure Supplement 1. Specificity of Netrin-1 and Netrin-3 IHC staining of
OVCAR8 spheroids. A) OVCAR8 cells were allowed to form spheroids for six hours before
fixation in formalin and embedding in paraffin for sectioning. Serial sections were stained with
Netrin-1 Abs or a no primary antibody control. **B)** OVCAR8 cells overexpressing Netrin-1 were
used to prepare spheroids as in A and stained in parallel to samples in A. **C)** OVCAR8 deleted
for Netrin-1 expression were used to prepare spheroids as in A and serial sections were stained
for Netrin-1 or with a no primary antibody control and counter stained. **D)** OVCAR8 cells were
allowed to form spheroids for six hours before fixation in formalin and embedding in paraffin for
sectioning. Serial sections were stained with Netrin-3 Abs or a no primary antibody control. **E)**
OVCAR8 cells overexpressing Netrin-3 were used to prepare spheroids as in D and stained in
parallel to samples in D. **F)** OVCAR8 deleted for Netrin-3 expression were used to prepare
spheroids as in D and serial sections were stained for Netrin-3 or with a no primary antibody
control. All scale bars = 100μm

Figure 5. Netrin ligands and their receptors are required for spheroid cell survival. A)
Illustration of netrin ligands, receptors, and other intracellular signaling molecules that are
included in the Axon Guidance pathway category. The frequency of their identification in
CRISPR screens is illustrated by shading and indicates how many cell lines had an enrichment
ratio <1 for a given component. **B)** The indicated ovarian cancer cell lines were infected with
lentiviruses expressing sgRNAs directed against the indicated Netrin signaling genes. Cells were
transferred to suspension culture conditions to induce spheroid formation for 72 hours and then
returned to adherent conditions for 24 hours to facilitate reattachment. Re-attached cells were
stained with Crystal Violet and retained dye was extracted and quantitated to measure relative
survival. Each cell-gene combination was assayed in at least three biological replicates,
averaged, and viability is displayed as a bubble plot. Mean survival for a given cell-gene
combination was compared with GFP control gRNA transduced cells using one way anova and
significance levels are illustrated by bubble size. Inviable cell-gene combinations are depicted as
empty spaces. **C)** Cultures of the indicated cell lines were cultured in suspension for five days
and stimulated with 0.5 μg/mL Netrin-1. Netrin-1 signaling was analyzed by SDS-PAGE and
western blotting for phospho-ERK, ERK, and tubulin. **D)** Quiescent adherent OVCAR8 cells
were stimulated with 0.5 μg/mL Netrin-1 or 0.5 μg/mL EGF, and compared with OVCAR8 cells
in suspension stimulated as in C. Extracts were prepared and blotted for phospho-ERK, ERK,
and tubulin. **E)** Suspension cultures of OVCAR8, or knock out derivatives, were harvested and
analyzed for relative phosphorylation levels of ERK by western blotting. Total ERK and tubulin
blotting serve as expression and loading controls. **F)** Netrin-1 signaling in OVCAR8, UNC5
4KO and DDN 3KO derivatives was tested by transferring cells to suspension and stimulating
with Netrin-1 as before. Western blotting for phospho-ERK, ERK were also as before. **G)**

OVCAR8 cells were seeded in suspension culture and treated with the MEK inhibitors PD184352, Trametinib or DMSO vehicle for 72 hours. Mean viability was determined by re-attachment and compared by one way anova (****$P<0.0001$). **H)** OVCAR8 cells were cultured in suspension and treated with Trametinib or DMSO vehicle as in G. Viability was determined by trypan blue dye exclusion and compared by one way anova (**$P<0.01$). **I)** Extracts were prepared from Trametinib and control treated spheroid cells and blotted for phospho-ERK, ERK, and tubulin. **J)** Model summarizing the roles of Netrin ligands, receptors, and downstream targets MEK and ERK in dormant survival signaling.

Figure 5 - Figure Supplement 1. Evaluation of target gene transcript levels in multigene knock out cells. A) RT-qPCR was performed on RNA isolated from control sgRNA expressing OVCAR8 cells and UNC5 4KO to assess expression of UNC5 family members. Graphs depict relative expression levels for each indicted gene in control and knock out cell lines. Mean expression levels of targeted cells were compared to background measurements using one way anova (****$P<0.0001$). **B)** RT-qPCR was used to measure mRNA levels of NTN1 and NTN3 transcripts in their respective CRISPR targeted, and OVCAR8 controls. Mean values were compared by one way anova (****$P<0.0001$). **C)** Levels of mRNA corresponding to DCC, DSCAM, and NEO1 were determined by RT-qPCR in DDN 3KO cells and control OVCAR8. Mean values were compared by one way anova (****$P<0.0001$). **D)** The indicated genotypes of cells were subjected to suspension culture and protein extracts were prepared at the indicated times. Proteins were resolved by SDS-PAGE, blotted, and probed with the indicated antibodies.

Figure 6. Loss of Netrin signaling reduces spheroids and prolongs survival. A) Control OVCAR8L (negative control CRISPR targeted cells) and UNC5 4KO cells were injected into the peritoneal space of female NOD/SCID mice. After two weeks, animals were euthanized and engrafted cells were collected with abdominal washes. **B)** RT-qPCR was used to compare gene expression of the indicated cell cycle markers in RNA extracted from proliferating cells in culture before engraftment and compared with RNA obtained from spheroids in xenografted mice. Technical replicates of RNA derived from three different mice is shown. **C)** Spheroids obtained from abdominal washes from each mouse were plated in adherent conditions to compare their abundance between control and UNC5 4KO genotypes. One example of each genotype of cell is shown. Crystal violet staining and dye extraction were used to quantitate biomass and averages were compared by one way anova (n=6, *$P<0.05$). **D)** DNA was extracted from cells collected in abdominal washes and human Alu repeats were detected by qPCR to quantitate and compare human cancer cells. Means were compared by one way anova (n=6, **$P<0.01$). **E)** Control OVCAR8 and UNC5 4KO cells were xenografted as above and mice were monitored until humane endpoint. **F)** Kaplan-Meier analysis of survival for mice engrafted with the indicated genotypes of cells. Survival was compared by logrank test.

Figure 6 - Figure Supplement 1. Evaluation of dormancy arrest in cell culture by RT-qPCR. RNA was extracted from asynchronously proliferating cells and suspension cultures 72 hours post transfer from adherent. RT-qPCR was used to compare gene expression of the indicated cell cycle markers in iOvCa147, OVCAR8, TOV1946.

Figure 7. Netrin ligand overexpression is associated with poor clinical outcome in HGSOC.
A) TCGA RNA-seq data for HGSOC patients (TCGA PanCancer Atlas study) was used to
identify high Netrin-1 or -3 and low Netrin-1 or -3 expressing patients (high expressing are
above z-score 1.2). Overall survival was used to construct Kaplan-Meier plots and survival was
compared using a logrank test. **B)** OVCAR8 cells were stably transduced with lentiviral
constructs to overexpress epitope tagged Netrin-1 or -3. Western blotting for Netrins, Myc-tags,
and Actin were used to determine relative expression levels of both Netrins in these cell
populations and vector controls. **C)** Control and Netrin overexpressing cells were transferred to
suspension culture conditions to form spheroids, and replated to assay for viability. Mean
viability and standard deviation is shown for each. One way anova was used to compare survival
(* $P<0.05$, *** $P<0.001$). **D)** Reattached spheroids were fixed and stained with Crystal Violet to
examine size and abundance in control and Netrin-1 or -3 overexpression.

Figure 7 - Figure Supplement 1. Frequency of Netrin ligand and receptor deletions,
mutations or expression changes in high grade serous ovarian cancer. A) Oncoprint of
genetic alterations in 10 netrin signaling related genes in 398 patients. Alteration types are
defined on the right. Only the first 154 patients are shown. **B)** Oncoprint illustrating gene
expression outliers among 10 netrin signaling related genes in 201 patients (first 122 patients
shown). Genomic data used is from Ovarian Serous Cystadenocarcinoma (TCGA, PanCancer
Atlas) (201 samples with RNA seqV2, z-score cut off of 1.5). **C)** TCGA RNA-seq data for
HGSOC patients (TCGA PanCancer Atlas study) was used to identify high Netrin-1 or -3 and
low Netrin-1 or -3 expressing patients (high expressing are above z-score 1.2). Progression free
survival was used to construct Kaplan-Meier plots and survival was compared using a logrank
test.

Figure 8. Netrin overexpression causes increased dissemination of tumor nodules. A)
Netrin overexpressing and control OVCAR8 cells were injected into the intraperitoneal space of
female NOD/SCID mice. Mice were euthanized following 35 days and analyzed for disease
burden by necropsy and histopathology. **B)** The spread of cancer to the diaphragm, liver,
omentum, and mesometrium was determined from necropsies and colors used in the anatomical
schematic correspond with petal plots for each genotype of cells. Petal plot radius illustrates
frequency of mice bearing disease spread to a particular location. The radius of color fill is
proportional to the total number of animals with tumor nodules found in that location. **C)**
Photographs of necropsy findings in the mesometrium. Locations of ovaries (*), oviduct (yellow
arrow), and uterine horn (green arrow) are indicated in each case. Tumor nodules are indicated
by white arrows. **D)** The number of mesometrium associated tumor nodules was determined for
each mouse. Mean values are indicated and differences between genotype were determined by
one way anova (* $P<0.05$, ** $P<0.01$). **E)** Photographs of necropsy findings in the liver.
Location of sternum is indicated (#) in each image. Tumor nodules are indicated by white
arrows. **F)** The number of liver associated tumor nodules was determined for each mouse. Mean
values are indicated and differences between genotype were determined by one way anova (**
$P<0.01$, *** $P<0.001$). **G** and **H)** Histology of mesometrial tumor nodules are shown. Serial

1004 sections were stained with H&E, or with the indicated antibodies for immunohistochemistry.
1005 Ovaries are indicated (*), as are the oviduct (yellow arrow), the uterus (green arrow), and the
1006 tumor nodule (white arrow). Scale bar = 2mm

Figure 1

A

TRACS (Toolset for the Ranked Analysis of GO-CRISPR Screens) algorithm:

All Enrichment Scores (ES) for all genes → Discard genes with Library ES in bottom quartile → Compare Spheroid ES and Adherent ES → Select genes with negative log2 Spheroid/Adherent ratio, *P<0.05*

B

Gene scoring formula

Gene score:
$$GS_j = \sum_{i=1}^{s} \left(\log_2 \frac{[normalized\ sgRNA\ abundance]_{Cas9+}}{[avg\ normalized\ sgRNA\ abundance]_{Cas9-}} \right)$$

Enrichment score:
$$ES_j = \frac{\frac{1}{n}\sum_{i=1}^{n} GS\ rank}{s}$$

Enrichment ratio:
$$ER_j = \log_2 \frac{ES_{j_{final}}}{ES_{j_{initial}}}$$

C   **D**

Figure 1 – Figure Supplement 1

Figure 2

Figure 2 – Figure Supplement 1

A

Figure 3

Figure 4

Figure 4 – Figure Supplement 1

Figure 5

Figure 5 – Figure Supplement 1

Figure 6

A iOvCa147

B OVCAR8

C TOV1946

Figure 6 – Figure Supplement 1

Figure 7

Figure 7 – Figure Supplement 1

Figure 8