Title:

Netrin signaling mediates survival of dormant epithelial ovarian cancer cells

Authors:

Pirunthan Perampalam[1,2], James I. MacDonald[1,3], Komila Zakirova[1,3], Daniel T. Passos[1,3], Yudith
Ramos-Valdes[1,4], Maëva Hervieu[5], Patrick Mehlen[5,6], Rob Rottapel[7,8], Benjamin Gibert[5], Rohann
Correa[1,9], Trevor G. Shepherd[1,4,9,10,11], Frederick A. Dick[1,3,9,12,13]

Affiliations:

1. London Regional Cancer Program, London Health Sciences Centre Research Institute,
London, Ontario, Canada, N6A 4L6.
2. Department of Biochemistry, University of Western Ontario, London, ON, Canada, N6A
5C1.
3. Department of Pathology and Laboratory Medicine, University of Western Ontario,
London, ON, Canada, N6A 5C1.
4. The Mary and John Knight Translational Ovarian Cancer Research Unit, London
Regional Cancer Program, London, Ontario, Canada, N6A 4L6.
5. Apoptosis, Cancer and Development Laboratory - Equipe labellisée 'La Ligue', LabEx
DEVweCAN, Institut Convergence PLAsCAN, Centre de Recherche en Cancérologie de
Lyon (CRCL), INSERM U1052-CNRS UMR5286, Université de Lyon, Université
Claude Bernard Lyon1, Centre Léon Bérard, 69008 Lyon, France.
6. Netris Pharma, 69008 Lyon, France
7. Princess Margaret Cancer Centre, University Health Network, 101 College Street,
Toronto, Ontario, Canada, M5G 1L7
8. Department of Medical Biophysics, University of Toronto, 1 King's College Circle,
Toronto, Ontario, Canada, M5S 1A8
9. Department of Oncology, Western University, London, Ontario, Canada, N6A 4L6.
10. Department of Obstetrics and Gynecology, Western University, London, Ontario,
Canada, N6A 5W9.
11. Department of Anatomy and Cell Biology, Western University, London, Ontario,
Canada, N6A 5C1.
12. Children's Health Research Institute, London, Ontario, Canada, N6A 4V2.
13. Correspondence: fdick@uwo.ca

Keywords:
Netrin, high grade serous ovarian cancer, DYRK1A, dormancy, quiescence, spheroids
37

38 **Conflict of interest statement:**
39 P. Mehlen is the founder and CEO of Netris Pharma and has an ownership interest including stocks
40 and patents. The other authors have declared that no conflict of interest exists.

Abstract

Dormancy in cancer is a clinical state in which residual disease remains undetectable for a prolonged duration. At a cellular level, rare cancer cells cease proliferation and survive chemotherapy and disseminate disease. We utilized a suspension culture model of high grade serous ovarian cancer (HGSOC) cell dormancy and devised a novel CRISPR screening approach to identify genetic requirements for cell survival under growth arrested and spheroid culture conditions. In addition, multiple RNA-seq comparisons were used to identify genes whose expression correlates with survival in dormancy. Combined, these approaches discover the Netrin signaling pathway as critical to dormant HGSOC cell survival. We demonstrate that Netrin-1 and -3, UNC5H receptors, DCC and other fibronectin receptors induce low level ERK activation to promote survival in dormant conditions. Furthermore, we determine that Netrin-1 and -3 overexpression is associated with poor prognosis in HGSOC and demonstrate their overexpression elevates cell survival in dormant conditions. Lastly, Netrin-1 or -3 overexpression contributes to greater spread of disease in a xenograft model of abdominal dissemination. This study highlights Netrin signaling as a key mediator HGSOC cancer cell dormancy and metastasis.

Introduction

Relapse from initially effective cancer treatment represents one of the greatest barriers to improving outcomes (1). Dormant or minimal residual disease that is not clinically detectable is a common source of relapse, particularly in a metastatic setting (2). This has motivated the search for, and study of, disseminated cancer cells. Cellular dormancy has been characterized in cancer cells derived from a variety of primary disease sites including breast, prostate, melanoma, and others (3, 4). Disseminated cells often occupy specific cellular niches in perivascular space or in bone marrow (5), they adopt catabolic metabolism (2), and withdraw from the cell cycle (4). Stem like phenotypes and epithelial to mesenchymal transition also characterize the biology of these cells as they survive dissemination and lay quiescent in distant tissues (1, 3, 5). Across a variety of disease sites this dormant state is characterized by diminished Ras-MAPK signaling and elevated p38 activity (3, 6). To develop treatments specifically tailored for dormant cells, a systematic search for vulnerabilities unique to dormancy is needed. To this end, high grade serous ovarian cancer disseminates as growth arrested cellular aggregates that are amenable to modeling in culture (7), suggesting they are ideal to search for survival dependencies of dormant cells.

High grade serous ovarian cancer (HGSOC) is the deadliest gynecologic malignancy, accounting for more than 70% of cases (8). Its poor prognosis is largely attributed to late-stage diagnosis with metastases already present. Metastatic spread is facilitated by multicellular clusters of tumor cells, called spheroids, that are shed into the peritoneal cavity and disseminate to nearby organs (9). The non-proliferative state of spheroid cells renders them chemoresistant and contributes to disease recurrence, emphasizing the need to understand the biology of these cells (10). Dormant HGSOC cells are characterized by similar metabolic changes, EMT, and

cytokine signaling as dormant cancer cells from other disease sites (3, 11). Mechanisms that

mediate their dormant properties include AMPK-LKB1 signaling that induces autophagy (12,

13), STAT3 and FAK for survival and chemoresistance (14, 15), and DYRK1A signaling to

arrest proliferation (16). Unfortunately, therapeutic approaches to target these pathways in

ovarian cancer have yet to emerge, indicating that more candidates are needed.

Netrins are a family of secreted factors that reside in the extracellular matrix around cells

and were first identified in axon guidance in the developing nervous system (17, 18). Together

with their receptors, Netrins have since been implicated in cancer progression (19). Netrin-1 is

the most studied and its misregulation in cancer leads to pro-survival signals (20). Netrin

interactions with DCC and UNC5 homologs (UNC5H, UNC5A-D) convert pro-death signals by

these dependence receptors to survival signals (21). Netrin signals through PI3K-AKT and

ERK-MAPK are known to participate in growth and orientation by providing guidance cues (22-

24), while loss of ligand binding leads DCC or UNC5H to stimulate dephosphorylation of

DAPK1 that sends pro-death signals (25, 26). The gene encoding DCC is frequently deleted in

cancer (20), and Netrin-1 blocking antibodies are in Phase II clinical trials for endometrial and

cervical carcinomas to inhibit metastatic growth (21). Alternative receptors for Netrins such as

NEO, DSCAM, and Integrins also mediate their signals (19), creating a myriad of possible

signaling pathways for Netrins to regulate cancer cells. Lastly, functional roles for Netrin-3, -4,

and -5 are still relatively unexplored compared to Netrin-1 (27), although their expression

patterns suggest roles in advanced forms of some cancers (28). A functional role for Netrins and

their receptors in dormancy, or other aspects of HGSOC spheroid biology have yet to be

reported.

In this study we utilized a suspension culture system to interrogate survival mechanisms

in dormancy using HGSOC spheroids. To maximize discovery of gene loss events that

compromise viability in dormancy, we devised a novel genome-wide CRISPR screen approach

that we term 'GO-CRISPR'. In addition to the standard CRISPR workflow, GO-CRISPR

utilizes a parallel screen in which 'guide-only' cells lacking Cas9 are used to control for

stochastic changes in gRNA abundance in non-proliferative cells. This screen identified multiple

Netrin ligands, DCC, UNC5Hs, and downstream MAPK signaling components as supporting

survival of HGSOC cells in dormant culture conditions. A transcriptomic analysis of HGSOC

spheroids also revealed Netrin signaling components were enriched in HGSOC spheroids

compared to adherent cells, and that in the absence of the DYRK1A survival kinase Netrin

transcriptional increases were lost. We show that multiple Netrins, UNC5 receptors, and the

downstream MEK-ERK axis is crucial for survival signaling in dormancy. Overexpression of

Netrin-1 and -3 are associated with poor prognosis in HGSOC and their overexpression in

culture increases survival in dormant suspension culture. Lastly, we demonstrate that Netrin

signaling contributes to disease dissemination in a xenograft model of metastasis. Our study

highlights Netrin-ERK signaling as a requirement for spheroid survival and suggests it may be a

potential therapeutic target for eradicating dormant HGSOC cells.

Results

Axon guidance genes are essential factors for EOC spheroid viability

To dissect genetic dependencies of dormant ovarian cancer cells, we utilized a suspension

culture system that encourages cellular aggregation, leading to non-proliferative spheroids that

resemble dormant aggregates found in ovarian cancer patient ascites. **Fig. 1A** shows phospho-

western blots for ERK and p38 from three high grade serous EOC cell lines; iOvCa147,

OVCAR8, TOV1946. Reduced phospho-ERK and elevated phospho-p38 are indicative of the

characteristic signaling switch reported to underlie most paradigms of cancer cell dormancy (5).

To discover genes required for survival in spheroids, we utilized a new approach called GO-

CRISPR (Guide Only CRISPR) that we developed specifically for survival gene discovery in

three-dimensional organoid/spheroid cell culture conditions (**Fig. 1B**). GO-CRISPR incorporates

sgRNA abundances from non-Cas9-expressing cells to control for stochastic (non-Cas9

dependent) effects that are a challenge in three-dimensional culture conditions with slow or

arrested proliferation. We used sequencing to determine sgRNA identity and abundance in the

library, initially infected cells, and in cells following 48 hours of suspension culture (**Fig. 1B**).

Read count data was analyzed using a software package that we developed to analyze GO-

CRISPR data (summarized in **Sup. Fig. 1A&B**). We filtered the results for genes that had a

Spheroid/Adherent Enrichment Ratio (ER) less than 1, with a $P<0.05$, which is indicative of

genes that were most depleted in dormant suspension conditions (**Fig. 1B**). To assess reliability

of screen data, we investigated the classification of non-targeting control sgRNAs. We computed

a receiver operator characteristic curve (ROC) and determined the area under the curve for each

cell line (**Sup. Fig. 1C**). Greater than 95% of sgRNAs in iOVCA147 and OVCAR8 screens were

classified as non-essential, while 83% of sgRNAs in TOV1946 cells were non-essential. In

addition, we compared GO-CRISPR screen ER values with viability data from siRNA knock

down for 35 individual genes we have previously studied in spheroid survival assays (**Sup. Fig.**

1D)(16). This revealed a statistically significant correlation between the two sets of data,

suggesting that genes we previously knew supported spheroid viability (or were dispensable)

were classified similarly by our screen. These controls suggested our screen data was reliable for

identifying new survival dependencies.

We found 6,717 genes with an ER below 1 in iOvCa147 cells; and 7,637 genes in

TOV1946; and 7,640 genes in OVCAR8 cells. Among these genes; 1,382 were commonly

depleted in suspension culture in all three cell lines (**Fig. 1D**). We performed pathway

enrichment analysis on these common genes and identified several significantly enriched

pathways. These include primary Reactome gene sets such as signal transduction, metabolism,

and signaling by GPCR (**Fig. 1E**). In addition, we identified a tertiary category called 'axon

guidance,' that includes numerous components of the Netrin signaling pathway.

Expression of Netrin ligands and receptors are enriched in EOC spheroids

DYRK1A was previously discovered to support survival of ovarian cancer cells in

suspension culture (16). Since DYRK1A phosphorylates RNA polymerase II in transcription

initiation (29), we sought to determine if it regulates gene expression in spheroid survival. We

created *DYRK1A$^{-/-}$* iOvCa147 cells and confirmed their deficiency for survival in suspension

(**Sup. Fig. 2A-E**). Using iOvCa147 and *DYRK1A$^{-/-}$* derivatives, we performed RNA-seq on

adherent and spheroid cells. We collected spheroids following a 6-hour incubation in suspension

because overexpression of known cell cycle transcriptional targets due to DYRK1A loss were

first evident at this time point, but loss of viability due to DYRK1A deficiency had not yet

occurred (16). We compared iOvCa147 adherent cells to 6-hour suspension culture to identify

transcriptional changes that occur as these cells transitioned from adherent conditions to

suspension conditions (**Fig. 2A**). We identified 1,834 genes that were upregulated, and pathway

enrichment analysis identified many of the same categories as our CRISPR screen including

axon guidance (**Fig. 2B**). We then compared iOvCa147 spheroid cells to *DYRK1A*$^{-/-}$ spheroid

cells to identify mRNA expression changes caused by DYRK1A deficiency (**Fig. 2C**). We

identified 744 genes that were downregulated and these also belonged to many of the same

pathways, including axon guidance (**Fig. 2D**).

We compared enriched pathways from these two RNA-seq analyses with those that were

identified in our GO-CRISPR screens to identify gene expression programs in spheroids whose

products are essential for cell survival. 78 Reactome pathways were commonly enriched across

the GO-CRISPR screens and transcriptomic analyses, including 78 of the 83 pathways identified

in RNA-seq from *DYRK1A*$^{-/-}$ spheroid cells (**Fig. 2E**). The axon guidance pathway was one of

the most enriched among these common pathways (**Fig. 2F**). These data reveal that our GO-

CRISPR screen and transcriptomic analyses converge on this highly specific tertiary Reactome

category of axon guidance alongside much broader primary or secondary gene sets. This

category includes Netrin ligands, receptors, and downstream signaling components, suggesting a

previously unappreciated role in HGSOC spheroid viability.

We separately sought to confirm expression of Netrins and their signaling components in

HGSOC spheroids. We examined expression levels of ligands and receptors in the three cell

lines used in our GO-CRISPR screen using RT- qPCR (**Fig. 3A-C**). The precise identities of

upregulated genes varies between cell lines, however each cell line increases expression of some

Netrins, at least one UNC5, and at least one of DCC, DSCAM, and Neogenin. This suggests that

the Netrin pathway may play a previously unappreciated role in HGSOC dormancy and disease

progression. Since Netrin signaling has not been previously reported in HGSOC dormancy we

examined expression of Netrin-1 and -3 ligands in spheroids isolated directly from patient ascites

by immunohistochemistry (**Fig. 3D-F**). We confirmed Netrin-1 and -3 antibody specificity using

spheroids generated from OVCAR8 cells bearing knock out or over expression of these Netrins

(**Sup. Fig. 3**). Spheroids were paraffin embedded and serial sections were also stained with

epithelial markers Cytokeratin and EpCAM, as well as p53 to confirm cancer cell identity within

spheroids that co-stain for Netrins (**Fig. 3D-F**). Overall, this confirms Netrin expression occurs

in HGSOC patient spheroids in precisely the context our screen indicates that it is critical for cell

survival.

Netrin signaling stimulates ERK to support HGSOC spheroid survival.

Netrin ligands and their dependence receptors are best known for their roles in axon guidance

(30), but are appreciated to regulate apoptosis in cancer (19). **Fig. 4A** illustrates components of

Netrin signaling and the frequency with which they were discovered to have an ER of <1 in our

screens. In addition, intracellular signaling molecules known to be downstream of Netrin

receptors, and that are shared with other enriched pathways identified in our screens are also

illustrated. To confirm screen findings, we independently knocked out *NTN1*, *NTN3*, *NTN4*,

NTN5, the UNC5H receptor homologs (*UNC5A*, *UNC5B*, *UNC5C*, and *UNC5D*), as well as

DCC, *DSCAM*, and *NEO1* alone or in multigene combinations using lentiviral delivery of

sgRNAs and Cas9 (**Fig. 4B and Sup. Fig. 4**). We assessed their loss on spheroid survival in six

HGSOC cell lines (iOvCa147, TOV1946, OVCAR3, OVCAR4, OVCAR8, and COV318). Loss

of at least one netrin ligand resulted in reduced spheroid cell survival in five of six cell lines

(**Fig. 4B**). Notably, Netrin-3 loss showed the most pronounced effect on survival. Cell lines such

as OVCAR8 and COV318 were relatively resistant to the deletion of netrin pathway components

as individual UNC5H, *DCC*, and *DSCAM* genes rarely affected viability (**Fig. 4B**). In these

instances, combined deletions of UNC5H family members, or *DCC;DSCAM;NEO1* (DDN for

short) were either inviable or highly sensitive to suspension culture induced cell death suggesting

redundancy in this pathway in some cell lines. Other cell lines, such as OVCAR3, were highly

sensitive to the loss of multiple individual ligands and any individual UNC5H receptor was

highly sensitizing to death induction in suspension culture (**Fig. 4B**). Overall, these illustrate a

robust and redundant role for Netrins in survival in this dormant cell culture assay. Importantly,

loss of receptor expression does not elevate viability as expected for a dependence receptor,

suggesting that Netrin receptors in HGSOC simply transmit positive survival signals.

To determine if Netrin dependent signaling is active in EOC spheroids, we transferred

serum deprived adherent cells to suspension and stimulated with recombinant Netrin-1 (**Fig. 4C**).

Western blotting for phospho-ERK, as a measure of activation, reveals stimulation of MEK-ERK

signaling in each of the cell lines tested under spheroid culture conditions. Because MEK and

ERK family members were identified in our screens (**Fig. 4A**), Netrin is a candidate pathway to

provide low level, but essential survival signals through the MAPK pathway. We investigated

the dependence of ERK phosphorylation on UNC5H and DDN receptors and demonstrate it is

lower when receptors are deleted (**Fig. 4D**). Activation using Netrin-1 stimulation of suspension

cultures of OVCAR8 cells, or derivatives deleted for UNC5 family members or DDN receptors,

indicates that Netrin receptors are essential to activate ERK in dormant suspension culture (**Fig.**

4E). Inhibition of MEK using the MEK1/2 inhibitor PD184352 causes loss of cell viability in

suspension culture (**Fig. 4F**), emphasizing that MEK-ERK signaling is key to cell survival. The

simplest explanation of this data is that Netrins signal through a heterodimer or multi-receptor

complex containing both an UNC5 and DDN component that provides stimulation for MEK and

ERK in the absence of mitogenic signals in dormant culture conditions.

Netrin overexpression is correlated with poor outcome in HGSOC.

Genomic studies indicate that HGSOC is one of the most aneuploid cancers (31). In

addition to the frequency of copy number variants per tumor, the locations of gains and losses

are dissimilar between cases, making this a highly heterogenous disease. Evaluation of TCGA

genomic data for HGSOC indicates rare amplifications or deletions of Netrin or Netrin-receptor

genes (**Sup. Fig. 5A**). Because of redundancy of Netrins and their receptor families this data

suggests HGSOC cases with deficiency for Netrin signaling are rare. This is consistent with our

data that all six cell lines tested reveal evidence of Netrin component dependence for survival

(**Fig. 4B**). Some TCGA HGSOC cases display elevated gene expression for Netrins or their

receptors, but only rare cases display low expression (**Sup. Fig. 5B**). Collectively, this suggests

that Netrin signaling in HGSOC is retained despite high copy number variation in this cancer

type and there is occasional overexpression.

In agreement with our experimental data that multiple Netrin ligands can contribute to

spheroid survival in HGSOC (**Fig. 4A&B**), sorting cases by high level expression of Netrin-1

and -3 ligands reveals a significantly shorter survival compared to cases with low level

expression and this trend is evident in the cases that only overexpress Netrin-3 (**Fig. 5A**). This

genomic data supports the interpretation that Netrin signaling is retained in HGSOC and its

elevation is deleterious to patient outcome.

We investigated overexpression of Netrin-1 and -3 in spheroid survival by generating

OVCAR8 derivatives that stably overexpress these proteins (**Fig. 5B**). We then utilized these

cells in suspension culture assays for spheroid survival as before (**Fig. 5C**). This data indicates

that overexpression of either Netrin-1 or -3 increases OVCAR8 cell survival in dormant spheroid

culture conditions. Examination of fixed spheroids isolated from this experiment further

indicates that elevated survival in this experiment is explained by an increase in abundance of

individual spheroids that overexpress either Netrin-1 or -3 (**Fig. 5D**). These experiments reveal

the significance of a previously undiscovered role for Netrins in HGSOC. High level expression

correlates with poor outcome and in suspension culture, overexpression confers oncogenic

properties on HGSOC cells.

Netrin overexpression induces abdominal spread of HGSOC.

Our findings from CRISPR screens and cell culture experiments indicate that Netrin

signaling supports dormant HGSOC cell survival. Our analysis of TCGA is also predictive of a

worse outcome for patients. We sought to solidify these findings by searching for how Netrin

signaling affects HGSOC disease pathogenesis.

We utilized a xenograft assay in which OVCAR8 cells overexpressing Netrin-1 or -3

were injected into the peritoneal cavity and compared with OVCAR8 cells bearing an empty

vector as control. Mice were harvest following 35 days to analyze disease dissemination and

characterize the effect of Netrins in this model of HGSOC metastatic spread (**Fig. 6A**). Necropsy

of all animals at endpoint allowed for the identification and quantitation of tumor nodules within

the abdomen (**Fig. 6B**). Petal plots illustrate the frequency of specific organ and tissue sites

displaying tumor nodules (**Fig. 6B**). This revealed spread to the omentum was similar across all

genotypes of xenografted cells and occurred in all mice (**Fig. 6B**). In contrast, tumor nodules

were detected with increased abundance on the diaphragm, the liver, and mesenteries that

support the uterus (mesometrium) (**Fig. 6B**). In addition to more frequent disease spread to these

locations caused by Netrin overexpressing cells, we note that the quantity of nodules was also

increase in comparison with OVCAR8 control cells (**Fig. 6C-F**). We confirmed the source of

these lesions to be the xenografted cells using immunohistochemical staining for human

cytokeratin (**Fig. 6G,H**). Lastly, Netrin-1 and -3 histochemical staining confirms its presence in

the local tumor microenvironment (**Fig. 6G,H**). The similarity of increased tumor nodules seen

in xenografts and elevated spheroid formation and viability seen in culture suggests that Netrin

improve cell viability in spheroid aggregates in vivo leading to increased disease spread.

Discussion

In this report we investigated survival mechanisms in a model of HGSOC dormancy. We

utilized a modified CRISPR screening approach that was adapted to discover gene loss events in

largely non-proliferative conditions. We compared essential genes discovered in the screen with

gene expression increases identified by RNA-seq in dormant HGSOC cultures. This implicated

the Netrin signaling pathway in HGSOC spheroid survival. We independently validated the role

of Netrins, their receptors, and downstream intracellular targets MEK and ERK in the survival

biology of HGSOC cells in dormant culture. Furthermore, we reveal that overexpression of

Netrin-1 and -3 are correlated with poor clinical outcome in HGSOC and over expression of

these ligands induces elevated survival in dormant HGSOC culture conditions and disease spread

in a xenograft model of dissemination. Our work implicates Netrins, their receptors, and MEK-

ERK as a previously unappreciated, but critical signaling network in HGSOC survival.

HGSOC spheroid cell biology has presented a unique challenge to cancer chemotherapy.

Dormant spheroids are most abundant in patients with late-stage disease and they contribute to

therapeutic resistance and seed metastases (32-34). Hence, there is a critical unmet need to

understand HGSOC spheroid dependencies and target them therapeutically. Our data suggests

that Netrin signaling is a universally active mechanism in spheroid cell survival. HGSOC is

characterized by extensive genomic rearrangements and copy number variants (31), and we

speculate that Netrin signaling is retained because of the extensive redundancy of components of

this pathway. Netrin-1 and -3 can both contribute to survival signaling and knock out of three or

four different Netrin receptors is necessary to compromise Netrin signaling in survival. A key

goal for the future is to identify Netrin-1 or -3 overexpression patients and compare their specific

disease progression patterns with our xenograft model. This will be most informative of the

effects of Netrin ligand overexpression in scenarios of dormancy, metastasis, and chemotherapy

resistance.

This study demonstrates that Netrins function to stimulate UNC5H and fibronectin repeat

containing receptors such as DCC to activate survival signals in HGSOC spheroids. This is

distinct from the known role of UNC5H and DCC as dependence receptors where Netrin inhibits

their pro-apoptotic signaling (21). Our data does not show elevated survival of spheroid cells in

response to deletions of receptors, even in instances where families of Netrin receptors are co-

deleted. Consistent with this observation, over expression of Netrin-1 or -3 in OVCAR8 cells

stimulates higher level survival in suspension. In colon cancer, frequent deletions of DCC confer

a survival advantage and underscore the dependence receptor concept (35, 36). In HGSOC DCC

or UNC5H deletions are relatively rare (31). This suggests that ovarian cancer cells utilize Netrin

signals for positive survival signaling instead of to neutralize inherent death signals from

dependence receptors. Our data suggests that Netrin activates MEK and ERK to support cell

survival indicating that targeting ligands, receptors or downstream at MEK are all possible

approaches to inhibit this survival pathway. Netrin-1 inhibition in solid tumor xenograft models

of melanoma have been shown to be effective in combination with chemotherapy (37), but its

effects on a dormant scenario are unknown. Identifying agents that cooperate with standard

HGSOC chemotherapeutic agents such as carboplatin would offer an attractive strategy to

eradicate dormant cells while combating progressive disease. Alternatively, a novel approach to

targeting dormancy discovered in this study is, ironically, MEK inhibition. Dormancy is

characterized by low phospho-ERK levels (3, 6), indicating low MEK activity, yet low levels of

RAS-MAPK signaling are known to provide survival signals (38). A challenge with deploying

MEK inhibitors as an anti-dormancy setting in HGSOC therapy is its incompatibility with

standard chemotherapies that rely on cell proliferation. Overall, our study aimed to identify

potential new therapeutic targets in dormancy and Netrins and MEK are feasible new candidates

for dormant HGSOC disease.

Current literature offers little to connect Netrins with HGSOC beyond the observation

that *NTN1* expression is increased in tumors compared to benign lesions (39). Similarly, recent

dormancy literature beyond HGSOC does not include a role for Netrins (3-6, 40). We expect that

Netrin signaling may play an unappreciated role in dormancy in other cancer disease site

paradigms. As stated in the introduction, dormancy is characterized by acquisition of stem like

properties, EMT, and in most paradigms it involves rare cells establishing themselves in

perivascular or bone marrow niches. A chick chorioallantoic membrane assay of metastasis in

which MDA-MB-231 cells were seeded was monitored for lung dissemination. Inhibition of

Netrin-1 with NP137 and treatment with decitabine has been shown to reduce lung seeding

suggesting that Netrins are relevant in a metastatic scenario that may include dormant

intermediate steps in dissemination (41). Furthermore, Netrin signaling through UNC5B is

known to direct vascular branching (42). This suggests that Netrins are available in the

perivascular niche where dormant cells often reside. Lastly, Netrin-1 has been shown to be a

component of the bone marrow niche that suppresses hematopoietic stem cell proliferation (43).

While not a cancer dormancy scenario, it indicates that Netrins support some solitary dormant

cell properties and are expressed by resident cells in the appropriate dormant cell niches beyond

just HGSOC. For these reasons we expect Netrins will have a significant role in tumor dormancy

beyond ovarian cancer.

Methods

Cell Lines and Engineering Cas9+ cells: High-grade serous ovarian cancer (HGSOC) cell lines
OVCAR3, OVCAR4, OVCAR8, COV318, TOV1946 and iOvCa147 were used in this study.
The iOvCa147 cells have previously been characterized and reported (44). All cell lines were
verified by STR analysis. iOvCa147, OVCAR8, and TOV1946 cells were engineered to express
Cas9 using pLentiCas9-Blast and clones were isolated. High Cas9 editing efficiency was
determined by viability studies using sgRNAs targeting selected fitness genes (PSMD1, PSMD2,
EIF3D) and a non-targeting control (LacZ) as previously described (45).

Lentiviral sgRNA library preparation: HEK293T cells were transfected with the combined A
and B components of the GeCKO v2 whole genome library (123,411 sgRNAs in total) along
with plasmids encoding lentiviral packaging proteins (46). Media was collected 2-3 days later,
cells and debris were pelleted by centrifugation, and supernatant containing viral particles was
filtered through a 0.45 µM filter and stored at −80°C with 1.1 g/100 mL BSA.

GO-CRISPR screens in iOvCa147, OVCAR8, and TOV1946 cells: Cas9-positive and Cas9-
negative cells were separately transduced with lentiviruses at a multiplicity of infection of 0.3
and with a predicted library coverage of >1000-fold. Cells were selected in puromycin and
maintained in complete media (DMEM/F12 with 10% FBS, 1% pen-strep-glutamine), containing
puromycin in all following steps. More than 10^9 cells were collected and split into three groups
consisting of approximately 3.0×10^8 cells. Triplicate samples of 6.2×10^7 cells were saved for
sgRNA sequence quantitation (T_0). The remaining cells were seeded at 2.0×10^6 cells/mL in 20
x 10 cm ULA plates. Following 2 days of culture, media containing spheroids was transferred to
10 x 15 cm adherent tissue culture plates. The next day unattached spheroid cells were collected
and re-plated onto additional 15 cm plates. This process was repeated for a total of 5 days to

maximally capture viable cells. Attached cells were collected and pooled for DNA extraction

(T_f). Genomic DNA was extracted using the QIAmp DNA Blood Maxi Kit (Qiagen, #51194).

PCR amplification and barcoding was performed as described (Hart paper, Cell, 2015). Next

generation sequencing was performed using an Illumina MiSeq 50 cycle kit on an Illumina

MiSeq platform.

TRACS analysis: The library reference file containing a list of all sgRNAs and their sequences

(CSV file), raw reads for the pooled sgRNA library (FASTQ files (L0) and raw reads (FASTQ

files) for all T_0 and T_f timepoints and replicates for Cas9-positive and Cas9-negative cells were

loaded into TRACS (https://github.com/developerpiru/TRACS). TRACS then automatically

trimmed the reads using Cutadapt (v1.15), built a Bowtie2 (v2.3.4.1) index and aligned the reads

using Samtools (v1.7). TRACS used the MAGeCK read count function (v0.5.6) to generate read

counts and incremented all reads by 1 to prevent zero counts and division-by-zero errors. The

TRACS algorithm was then run using this read count file to determine the Library Enrichment

Score (ES), Initial ES, Final ES and the Enrichment Ratio (ER) for each gene. VisualizeTRACS

was used to generate graphs of Initial and Final ratios.

Pathway enrichment analyses: For GO-CRISPR screens and multi-omics comparisons, we used

the filtered list of genes obtained using TRACS and performed gene ontology and pathway

enrichment analysis using the ConsensusPathDB enrichment analysis test (Release 34

(15.01.2019), http://cpdb.molgen.mpg.de/) for top-ranked genes of interest. p_{adj} values and ER

values for each gene were used as inputs. The minimum required genes for enrichment was set to

45 and the FDR-corrected p_{adj} value cutoff was set to < 0.01. The Reactome pathway dataset was

used as the reference. For each identified pathway, ConsensusPathDB provides the number of

enriched genes and a q value (p_{adj}) for the enrichment. Bar plots were generated in R 3.6.2 using

these values to depict the significant pathways identified.

Generating DYRK1A knockout cells: A double cutting CRISPR/Cas9 approach with a pair of

sgRNAs (sgRNA A and B) was used to completely excise exon 2 (322 bp region) of *DYRK1A*

using a px458 vector (Addgene #48138) that was modified to express the full CMV promoter.

PCR primers that flank the targeted region of *DYRK1A* were used to verify deletion. Single-cell

clones were generated by liming dilutions and evaluated for DYRK1A status by PCR, western

blot, and sequencing. See Supplemental Table 1 for sgRNA and PCR primer sequences.

DYRK1A IP kinase assay: Whole cell lysates from adherent parental iOvCa147 cells and

DYRK1A$^{-/-}$ cells were extracted using complete RIPA buffer and incubated overnight with

DYRK1A antibody (Cell Signaling Technology anti-DYRK1A rabbit antibody #8765). Samples

were then washed with buffer and Dynabeads (Thermo Fisher Scientific Dynabeads Protein G

#10003D) were added for 2 hours. Samples were then washed with buffer and recombinant Tau

protein (Sigma recombinant Tau protein #T0576) and ATP (Sigma #A1852) were added and

samples were incubated for 30 minutes at 37°C. 5x SDS was then added and samples were

resolved by SDS-PAGE. Antibodies used for western blotting were phosphospecific Tau

antibody (Cell Signaling Technology phospho-Tau Ser-404 rabbit antibody #20194) and Tau

protein (Cell Signaling Technology anti-Tau rabbit antibody #46687).

RNA preparation and RNA-sequencing: Total RNA was collected from parental iOvCa147 cells

and *DYRK1A$^{-/-}$* cells from 24-hour adherent or 6-hour spheroid conditions using the Monarch

Total RNA Miniprep Kit (NEB #T2010S) as described above. RNA was collected in three

replicates for each condition (adherent and spheroids). Samples were quantitated using Qubit 2.0

Fluorometer (Thermo Fisher Scientific) and quality control analysis using Agilent 2100

Bioanalyzer (Agilent Technologies #G2939BA) and RNA 6000 Nano Kit (Agilent Technologies

#5067-1511). Ribosomal RNA removal and library preparation was performed using ScriptSeq

Complete Gold Kit (Illumina #BEP1206). High-throughput sequencing was performed on an

Illumina NextSeq 500 platform (mid-output, 150-cyclekit).

RNA-seq analyses: Raw FASTQ data was downloaded from Illumina BaseSpace. Reads were

aligned using STAR 2.6.1a(47) to the human genome

(Homo_sapiens.GRCh38.dna.primary_assembly.fa; sjdbGTFfile:

Homo_sapiens.GRCh38.92.gtf) to generate read counts. DESeq2 was run with BEAVR(48) and

used to analyze read counts (settings: False discovery rate (FDR): 10%; drop genes with less

than 1 reads) to identify differentially expressed genes ($\log_2$ fold change > 1 for upregulated

genes; or $\log_2$ fold change < -1 for downregulated genes; $p_{adj} < 0.05$) for the following

comparisons: parental iOvCa147 adherent cells vs. parental iOvCa147 spheroid cells; or parental

iOvCa147 spheroid cells vs. *DYRK1A*$^{-/-}$ spheroid cells. For RNA-seq, we formed pathway

analyses in BEAVR and downloaded the pathway enrichment table to construct bar plots and

determine overlapping pathways using R 3.6.2.

Western blots: Adherent cells were washed in 1x PBS and lysed on the plate by the addition of

complete RIPA buffer with protease (Cell Signaling Technology #5871S) and phosphatase (Cell

Signaling Technologies #5870S) inhibitors and incubated for 5-10 minutes on ice. Spheres were

collected by gentle centrifugation at 300 rpm, washed gently in ice cold PBS, centrifuged again

and then resuspended in RIPA buffer as above. Samples were then centrifuged at 14,000 RPM at

4°C for 10 minutes. The supernatant liquid for each sample was collected and the pellets re-

extracted with RIPA containing the above inhibitor cocktails. Following centrifugation, the

supernates for each treatment were collected and pooled. Protein was determined by Bradford

assay. To assess the effect of Netrin 1 on downstream signaling, spheres incubated under non

adherent conditions for 5 days were challenged with the addition of 500 ng/ml human

recombinant Netrin-1 (R&D Systems #6419-N1-025) for a period of 20 minutes before collection.

Lysates were mixed with 6x SDS loading dye buffer and resolved using standard SDS-PAGE

protocols. Antibodies used for blotting were p42/44 MAPK (Erk1/2) (Cell Signaling Technology

rabbit antibody #4695), Phospho-p44/42 MAPK(Erk1/2) T202/Y204 (Cell Signaling

Technology; rabbit antibody #4370), p38 MAPK (Cell Signaling Technology; rabbit antibody

#9212), p38 MAPK T180/Y182 (Cell Signaling Technology; rabbit antibody #9211), DYRK1A

(Cell Signaling Technology anti-DYRK1A rabbit antibody #8765), Netrin-1 (Abcam rabbit

antibody #ab126729), Netrin-3 (Abcam rabbit antibody #ab185200), α-tubulin (Cell Signaling

Technology anti-αTubulin rabbit antibody #2125) and β-actin (Millipore Sigma rabbit antibody

#A2066).

RT-qPCR analysis of gene expression: Cells were triturated off the plate using Versene (0.5mM

EDTA in PBS), collected, counted and 10^6 cells plated onto Ultra-Low Attachment cell culture

dishes (Corning) to induce sphere formation. The spheres were collected after 24 h and RNA

extracted using the Monarch Total RNA Miniprep Kit (New England Biolabs #T2010S). RNA

was extracted from adherent cells directly from the plate. Extracted RNA was quantitated using a

NanoDrop and stored at -80º until used. First strand cDNA was generated from isolated RNA

using the iScript cDNA Synthesis Kit (Bio-Rad, #1708890) and RT-qPCR performed on the

resulting cDNA (diluted 5-fold) using the iQ SYBR Green Supermix (Bio-Rad, #1708882) and

following the manufactures instructions. Reactions were performed in 0.2ml non-skirted low

profile 96 well PCR plates (Thermo Scientifi, #AD-0700) using a CFX Connect Real Time

System thermo-cycler (Bio-Rad).

TCGA patient data analyses: We used the cBioPortal for Cancer Genomics

(https://www.cbioportal.org)(49) to analyze Netrin pathway alterations in Ovarian Serous

Cystadenocarcinoma patients (TCGA, PanCancer Atlas) for genomic alterations and mRNA

expression. OncoPrints were generated for mRNA expression (mean z-score threshold ± 1.5),

and mutations and putative copy number alterations for the following genes: *NTN1, NTN3,*

NTN4, NTN5, UNC5A, UNC5B, UNC5C, UNC5D, DCC, DSCAM. Survival analysis of *NTN1,*

and -*3* utilized patients from TCGA, PanCancer Atlas dataset and RNA-seq expression (high

expression z-score threshold > 1.2 and low is below 1.2).

Generation of gene knockout cell lines: Gibson Assembly (NEB #E2611) was used to clone

sgRNAs sequences derived from the GeCKO v2 library per gene of interest (*NTN1, NTN3,*

NTN4, NTN5, DSCAM, DCC, NEO1, UNC5A, UNC5B, UNC5C, UNC5D, and *EGFP* control

(See Supplemental Table 1 for sgRNA sequences for each)) into lentiCRISPR v2 (Addgene

#52961). For individual gene knockouts pooled libraries of up to four sgRNAs were generated.

For multiple gene knockouts pooled libraries of two sgRNAs/gene were constructed. To each 20-

nucleotide guide (Supplemental table I) was appended the 5' sequence, 5'-

TATCTTGTGGAAAGGACGAAACACC-3' and the 3' sequence, 5'-

GTTTTAGAGCTAGAAATAGCAAGTTAAAAT-3' (50). A 100 bp fragment was then

generated using the forward primer, 5'-

GGCTTTATATATCTTGTGGAAAGGACGAAACACCG-3', and reverse primer, 5'-

CTAGCCTTATTTTAACTTGCTATTTCTAGCTCTAAAAC-3′. The resulting fragment was

gel purified, quantitated by Qubit and cloned into the lentiviral vector as described (50). To

generate virus particles, HEK293T cells were transfected with the assembled plasmid along with

plasmids encoding lentiviral packaging proteins. Media was collected after 2-3 days and any

cells or debris were pelleted by centrifugation at 500 x g. Supernatant containing viral particles

was filtered through a low protein binding 0.45 µM filter. For each knockout, it was important to

ensure that each target cell was transduced with a full complement of guides. To ensure this, the

cells were transduced at relatively high MOI. Briefly, 25,000 cells were plated and transduced

with fresh HEK293T conditioned media containing virus. While we did not titer the viral

preparations for these experiments, previous experience indicated that we could generate

sufficiently high virus titers using our protocol. Under these conditions, we found a lack of

puromycin sensitive cells following 2-3 days in media containing 2-4 µg/ml puromycin

indicating that every cell received at least one virus particle. Multi-gene knock outs were created

through multiple rounds of infection with the relevant culture supernatants followed by one

round of puromycin selection.

Generation of gene overexpression cell lines: Plasmids encoding Netrin1 and Netrin3 were

obtained from Addgene (Ntn1-Fc-His, #72104 and Ntn3-Fc-His, #72105). Netrin1 was PCR

amplified using the forward primer 5'-

GCTATCGATATCCCAAACGCCACCATGATGCGCGCTGTGTGGGAGGCGCTG-3' and

reverse 5'-GCTATCTCTAGAGGCCTTCTTGCACTTGCCCTTCTTCTCCCG-3' and cloned

into the EcoRV/XbaI sites of pcDNA3.1/myc-HisA. For lentiviral expression, pcDNA NTN1

was amplified using the primers, forward 5'-

CACTGTAGATCTCCAAACGCCACCATGATGCGCGCTGTGTGGGAGGCGCTG-3' and,

reverse 5'-ACGCGTGAATTCTTATCAACCGGTATGCATATTCAGATCCTCTTCTGAGAT-

3' while NTN3 was amplified from pcDNA NTN3 using, forward 5'-

CTCGAGAGATCTGCGGCCGCCACCATGCCTGGCTGGCCCTGG-3' and the same reverse

primer as for NTN1. In this way both constructs carry a C-terminal Myc tag. Both cDNA's were

inserted into the BamHI/EcoRI sites of the lentiviral expression vector FUtdTW (Addgene

#22478). Construction of lentiviral particle and transduction into the target cells was as described

above. Selection was with Zeocin (ThermoFisher #R25001).

Spheroid formation and reattachment assays: For each targeted gene in iOvCa147, OVCAR3,

OVCAR4, OVCAR8, COV318, or TOV1946 cells, spheroid viability was assayed as follows:

Knockout cells were cultured for 72 hours in suspension conditions using ULA plasticware (2 ×

10^6 cells per well) to allow spheroid formation. Addition of therapeutics (PD184352) was made

at the time of plating. Spheroids were collected and transferred directly to standard plasticware to

facilitate reattachment for 24 hours. Reattached cells were fixed in 25% methanol in PBS for 3

minutes. Fixed cells were incubated for 30 minutes with shaking in 0.5% crystal violet, 25%

methanol in PBS. Plates were carefully immersed in water to remove residual crystal violet and

destained in 10% acetic acid in 1x PBS for 1 hour with shaking to extract crystal violet from

cells. Absorbance of crystal violet was measured at 590 nm using a microplate reader (Perkin

Elmer Wallac 1420). Each knockout was normalized to the EGFP control. Because crystal violet

binds protein, a linear relationship between the absorbance and the number of cells present is

obtained and may be represented as the percent of spheroid cells surviving suspension.

Xenografts: Female NOD-SCID mice were purchased from JAX (#001303). They were

engrafted between six and eight weeks of age with 4 x10^6 OVCAR8 control or modified

derivatives. Mice were housed for 35 days and euthanized and subjected to a necropsy to analyze

disease spread. Tumor lesions were photographed and quantitated. Abdominal organs and

associated tumor lesions were fixed in 10% formalin overnight and transferred to 70% ethanol

until embedded.

Patient derived spheroids: Patient derived spheroids were isolated from ascites fluid obtained

from debulking surgical procedures. These HGSOC samples were obtained with permission and

institutional REB oversight. Spheroids were prepared by filtering the ascites fluid through a

100μm mesh filter and washing the captured spheroids from the filter with PBS. The spheres

were collected by gravity on ice in PBS, washed twice with ice cold PBS and allowed to settle on

ice. Spheres were then fixed for 20 minutes in 10% formalin, washed, and resuspended in PBS

before being embedded in paraffin.

Histology and immunohistochemistry: Fixed organs from xenografts or spheroids were

embedded in paraffin, sectioned, and stained with H&E. Sections were deparaffinized by

standard procedures and antigen retrieval was performed in 10mM citrate, pH6.0. The sections

were blocked with 10% goat serum/3% BSA/0.3% Triton X-100 and stained with antibodies to

Netrin-1 (Abcam #ab126729, 1:200), Cytokeratin (human specific cytokeratin 7 and 8, ZETA

Corporation #Z2018ML, 1:200), EpCAM (Cell Signaling Technologies, #14452), p53 (Cell

Signaling Technologies, #2527) and Netrin-3 (1:250). Netrin-3 antibodies were generated using

immunized rabbits and their generation and purification will be published elsewhere. Primary

antibodies were followed with a biotinylated anti-rabbit (Vector Labs, # BA-1000) or anti-mouse

(Jackson ImmunoResearch, #115-067-003) IgG as a secondary antibody (1:500). Proteins were

visualized using DAB chemistry (ImmPACT DAB, Vector Labs, #SK-4105).

Statistical analyses:　Specific statistical tests used are indicated in the figure legends for each

experiment.　Analysis was done using GraphPad Prism 6.

References

1. Giancotti FG. Mechanisms governing metastatic dormancy and reactivation. Cell.
2013;155(4):750-64.
2. Massague J, Ganesh K. Metastasis-Initiating Cells and Ecosystems. Cancer Discov.
2021;11(4):971-94.
3. Sosa MS, Bragado P, Aguirre-Ghiso JA. Mechanisms of disseminated cancer cell
dormancy: an awakening field. Nat Rev Cancer. 2014;14(9):611-22.
4. Phan TG, Croucher PI. The dormant cancer cell life cycle. Nat Rev Cancer.
2020;20(7):398-411.
5. Goddard ET, Bozic I, Riddell SR, Ghajar CM. Dormant tumour cells, their niches and the
influence of immunity. Nat Cell Biol. 2018;20(11):1240-9.
6. Yeh AC, Ramaswamy S. Mechanisms of Cancer Cell Dormancy--Another Hallmark of
Cancer? Cancer Res. 2015;75(23):5014-22.
7. Lengyel E, Burdette JE, Kenny HA, Matei D, Pilrose J, Haluska P, et al. Epithelial
ovarian cancer experimental models. Oncogene. 2014;33(28):3619-33.
8. Lheureux S, Gourley C, Vergote I, Oza AM. Epithelial ovarian cancer. Lancet.
2019;393(10177):1240-53.
9. Lengyel E. Ovarian cancer development and metastasis. Am J Pathol. 2010;177(3):1053-
64.
10. Bowtell DD, Bohm S, Ahmed AA, Aspuria PJ, Bast RC, Jr., Beral V, et al. Rethinking
ovarian cancer II: reducing mortality from high-grade serous ovarian cancer. Nat Rev Cancer.
2015;15(11):668-79.
11. Shepherd TG, Dick FA. Principles of dormancy evident in high-grade serous ovarian
cancer. Cell Div. 2022;17(1):2.
12. Peart T, Ramos Valdes Y, Correa RJ, Fazio E, Bertrand M, McGee J, et al. Intact LKB1
activity is required for survival of dormant ovarian cancer spheroids. Oncotarget.
2015;6(26):22424-38.
13. Buensuceso A, Ramos-Valdes Y, DiMattia GE, Shepherd TG. AMPK-Independent
LKB1 Activity Is Required for Efficient Epithelial Ovarian Cancer Metastasis. Mol Cancer Res.
2019.
14. Chen MW, Yang ST, Chien MH, Hua KT, Wu CJ, Hsiao SM, et al. The STAT3-miRNA-
92-Wnt Signaling Pathway Regulates Spheroid Formation and Malignant Progression in Ovarian
Cancer. Cancer Res. 2017;77(8):1955-67.
15. Diaz Osterman CJ, Ozmadenci D, Kleinschmidt EG, Taylor KN, Barrie AM, Jiang S, et
al. FAK activity sustains intrinsic and acquired ovarian cancer resistance to platinum
chemotherapy. Elife. 2019;8.
16. MacDonald J, Ramos-Valdes Y, Perampalam P, Litovchick L, DiMattia GE, Dick FA. A
Systematic Analysis of Negative Growth Control Implicates the DREAM Complex in Cancer
Cell Dormancy. Mol Cancer Res. 2017;15(4):371-81.
17. Rajasekharan S, Kennedy TE. The netrin protein family. Genome Biol. 2009;10(9):239.
18. Kryza D, Wischhusen J, Richaud M, Hervieu M, Sidi Boumedine J, Delcros JG, et al.
From netrin-1-targeted SPECT/CT to internal radiotherapy for management of advanced solid
tumors. EMBO Mol Med. 2023;15(4):e16732.
19. Mehlen P, Delloye-Bourgeois C, Chedotal A. Novel roles for Slits and netrins: axon
guidance cues as anticancer targets? Nat Rev Cancer. 2011;11(3):188-97.

20.   Arakawa H. Netrin-1 and its receptors in tumorigenesis. Nat Rev Cancer.
2004;4(12):978-87.
21.   Brisset M, Grandin M, Bernet A, Mehlen P, Hollande F. Dependence receptors: new
targets for cancer therapy. EMBO Mol Med. 2021;13(11):e14495.
22.   Forcet C, Stein E, Pays L, Corset V, Llambi F, Tessier-Lavigne M, et al. Netrin-1-
mediated axon outgrowth requires deleted in colorectal cancer-dependent MAPK activation.
Nature. 2002;417(6887):443-7.
23.   Larrieu-Lahargue F, Welm AL, Thomas KR, Li DY. Netrin-4 induces lymphangiogenesis
in vivo. Blood. 2010;115(26):5418-26.
24.   Yin K, Wang L, Zhang X, He Z, Xia Y, Xu J, et al. Netrin-1 promotes gastric cancer cell
proliferation and invasion via the receptor neogenin through PI3K/AKT signaling pathway.
Oncotarget. 2017;8(31):51177-89.
25.   Llambi F, Lourenco FC, Gozuacik D, Guix C, Pays L, Del Rio G, et al. The dependence
receptor UNC5H2 mediates apoptosis through DAP-kinase. EMBO J. 2005;24(6):1192-201.
26.   Castets M, Coissieux MM, Delloye-Bourgeois C, Bernard L, Delcros JG, Bernet A, et al.
Inhibition of endothelial cell apoptosis by netrin-1 during angiogenesis. Dev Cell.
2009;16(4):614-20.
27.   Bruikman CS, Zhang H, Kemper AM, van Gils JM. Netrin Family: Role for Protein
Isoforms in Cancer. J Nucleic Acids. 2019;2019:3947123.
28.   Jiang S, Richaud M, Vieugue P, Rama N, Delcros JG, Siouda M, et al. Targeting netrin-3
in small cell lung cancer and neuroblastoma. EMBO Mol Med. 2021;13(4):e12878.
29.   Di Vona C, Bezdan D, Islam AB, Salichs E, Lopez-Bigas N, Ossowski S, et al.
Chromatin-wide profiling of DYRK1A reveals a role as a gene-specific RNA polymerase II
CTD kinase. Mol Cell. 2015;57(3):506-20.
30.   Meijers R, Smock RG, Zhang Y, Wang JH. Netrin Synergizes Signaling and Adhesion
through DCC. Trends Biochem Sci. 2020;45(1):6-12.
31.   Network CGA. Integrated genomic analyses of ovarian carcinoma. Nature.
2011;474(7353):609-15.
32.   Ahmed N, Stenvers KL. Getting to know ovarian cancer ascites: opportunities for
targeted therapy-based translational research. Front Oncol. 2013;3:256.
33.   Narod S. Can advanced-stage ovarian cancer be cured? Nat Rev Clin Oncol.
2016;13(4):255-61.
34.   Keyvani V, Farshchian M, Esmaeili SA, Yari H, Moghbeli M, Nezhad SK, et al. Ovarian
cancer stem cells and targeted therapy. J Ovarian Res. 2019;12(1):120.
35.   Fearon ER, Cho KR, Nigro JM, Kern SE, Simons JW, Ruppert JM, et al. Identification of
a chromosome 18q gene that is altered in colorectal cancers. Science. 1990;247(4938):49-56.
36.   Mehlen P, Rabizadeh S, Snipas SJ, Assa-Munt N, Salvesen GS, Bredesen DE. The DCC
gene product induces apoptosis by a mechanism requiring receptor proteolysis. Nature.
1998;395(6704):801-4.
37.   Boussouar A, Tortereau A, Manceau A, Paradisi A, Gadot N, Vial J, et al. Netrin-1 and
Its Receptor DCC Are Causally Implicated in Melanoma Progression. Cancer Res.
2020;80(4):747-56.
38.   Downward J. Targeting RAS signalling pathways in cancer therapy. Nat Rev Cancer.
2003;3(1):11-22.
39.   Papanastasiou AD, Pampalakis G, Katsaros D, Sotiropoulou G. Netrin-1 overexpression
is predictive of ovarian malignancies. Oncotarget. 2011;2(5):363-7.

40.   Recasens A, Munoz L. Targeting Cancer Cell Dormancy. Trends Pharmacol Sci.
2019;40(2):128-41.
41.   Grandin M, Meier M, Delcros JG, Nikodemus D, Reuten R, Patel TR, et al. Structural
Decoding of the Netrin-1/UNC5 Interaction and its Therapeutical Implications in Cancers.
Cancer Cell. 2016;29(2):173-85.
42.   Lu X, Le Noble F, Yuan L, Jiang Q, De Lafarge B, Sugiyama D, et al. The netrin
receptor UNC5B mediates guidance events controlling morphogenesis of the vascular system.
Nature. 2004;432(7014):179-86.
43.   Renders S, Svendsen AF, Panten J, Rama N, Maryanovich M, Sommerkamp P, et al.
Niche derived netrin-1 regulates hematopoietic stem cell dormancy via its receptor neogenin-1.
Nat Commun. 2021;12(1):608.
44.   Tong JG, Valdes YR, Sivapragasam M, Barrett JW, Bell JC, Stojdl D, et al. Spatial and
temporal epithelial ovarian cancer cell heterogeneity impacts Maraba virus oncolytic potential.
BMC Cancer. 2017;17(1):594.
45.   Hart T, Chandrashekhar M, Aregger M, Steinhart Z, Brown KR, MacLeod G, et al. High-
Resolution CRISPR Screens Reveal Fitness Genes and Genotype-Specific Cancer Liabilities.
Cell. 2015;163(6):1515-26.
46.   Sanjana NE, Shalem O, Zhang F. Improved vectors and genome-wide libraries for
CRISPR screening. Nat Methods. 2014;11(8):783-4.
47.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.
48.   Perampalam P, Dick FA. BEAVR: a browser-based tool for the exploration and
visualization of RNA-seq data. BMC Bioinformatics. 2020;21(1):221.
49.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer
genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer
Discov. 2012;2(5):401-4.
50.   Wang T, Lander ES, Sabatini DM. Large-Scale Single Guide RNA Library Construction
and Use for CRISPR-Cas9-Based Genetic Screens. Cold Spring Harb Protoc. 2016;2016(3):pdb
top086892.

Figure 1. GO-CRISPR screens implicate axon guidance pathways as supporting EOC spheroid cell viability.
A) iOvCa147, TOV1946, or OVCAR8 cells were cultured under adherent conditions (Adh) or in suspension to induce spheroids (Sph). Lysates were prepared and analyzed by western blotting for phosphorylated and total levels of ERK and p38. **B)** Flow chart of GO-CRISPR screening used for each of iOvCa147, TOV1946, or OVCAR8 control cells or that express Cas9. Cas9-positive cells (top row) and Cas9-negative cells (bottom row) were transduced with the GeCKO v2 pooled sgRNA library. After antibiotic selection, cells were expanded under adherent culture conditions (T_0) before being transferred to suspension culture conditions to induce spheroid formation and select for cell survival. After 48 hours spheroids were transferred to standard plasticware to isolate viable cells (T_f). Red arrows indicate the relevant comparisons of sgRNA sequence abundance that were made to analyze screen outcomes. **C)** Scatter plots representing spheroid score (T_f) on y-axis and adherent (T_0) score on x-axis calculated by TRACS for each gene in each cell line (iOvCa147, TOV1946, OVCAR8). Colored data points represent genes with ER < 1 and p_{adj} < 0.05. **D)** Venn diagram illustrating overlap of genes identified as supporting cell viability in suspension culture from iOvCa147, TOV1946, and OVCAR8 cells. **E)** Graph depicting enriched pathways from ConsensusPathDB using the 1382 commonly identified genes from D. Categories are ranked by q-value.

Figure 2. Axon guidance pathway components are upregulated in iOvCa147 spheroid cells in a DYRK1A dependent manner. A) RNA was isolated from iOvCa147 cells following culture under adherent conditions or in suspension conditions to induce spheroid formation for 6 hours. Triplicate independent cultures were processed for RNA-seq. A volcano plot shows differentially expressed genes in spheroid cells compared to adherent. 1,937 genes were found to be downregulated in iOvCa147 spheroid cells ($\log_2$ fold change < 1, p_{adj} < 0.05, FDR 10%, highlighted in grey) and 1,834 genes were upregulated ($\log_2$ fold change > 1, p_{adj} < 0.05, FDR 10%, highlighted in blue). **B)** Top 15 most significantly enriched pathways (p_{adj} < 0.05) whose genes were upregulated in suspension culture compared to adherent in RNA-seq analysis. **C)** A volcano plot showing differentially expressed genes in $DYRK1A^{-/-}$ spheroid cells compared to iOvCa147 spheroid cells. 744 genes were found to be downregulated in $DYRK1A^{-/-}$ spheroid cells ($\log_2$ fold change < 1, p_{adj} < 0.05, FDR 10%, highlighted in red) and 96 genes were upregulated ($\log_2$ fold change > 1, p_{adj} < 0.05, FDR 10%, highlighted in grey). **D)** Top 15 most significantly enriched pathways (p_{adj} < 0.05) that were represented by downregulated genes in DYRK1A-/- suspension culture compared to control cells in suspension. **E)** Venn diagram depicting overlapping enriched pathways identified in GO-CRISPR screens in green; enriched pathways identified in upregulated genes in parental iOvCa147 spheroid cells in blue; and enriched pathways identified in downregulated genes in $DYRK1A^{-/-}$ spheroid cells in red. 78 pathways were commonly enriched in all three datasets (shown in yellow). **F)** Top 10 most significantly enriched pathways (p_{adj} < 0.05) among the 78 identified in C.

Figure 3. Expression of Netrin ligands and their dependence receptors is increased in suspension culture. A-C) RT-qPCR was performed to quantitate mRNA expression levels of Netrin ligands and receptors in three different HGSOC cell lines. Relative expression of the indicated transcripts is shown for suspension culture conditions. All experiments for performed in at least triplicate replicates. **D-F)**. Immunohistochemical staining was performed for the indicated proteins using spheroids isolated from HGSOC patient ascites that was fixed and paraffin embedded. Omission of primary antibody was used a control for background staining for each patient sample. Scale bar = 100μm

Figure 4. Netrin ligands and their receptors are required for spheroid cell survival. **A)** Illustration of netrin ligands, receptors, and other intracellular signaling molecules that are included in the Axon Guidance pathway category. The frequency of their identification in CRISPR screens is illustrated by shading and indicates how many cell lines had a negative enrichment ratio for a given component. **B)** The indicated ovarian cancer cell lines were infected with lentiviruses expressing sgRNAs directed against the indicated Netrin signaling genes. Cells were transferred to suspension culture conditions to induce spheroid formation for 72 hours and then returned to adherent conditions for 24 hours to facilitate reattachment. Re-attached cells were stained with Crystal Violet and retained dye was extracted and quantitated to measure relative survival. Each cell-gene combination was assayed in at least three biological replicates, averaged, and viability is displayed as a bubble plot. Mean survival for a given cell-gene combination was compared with GFP control gRNA transduced cells using anova and significance levels are illustrated by bubble size. Inviable cell-gene combinations are depicted as empty white spaces. **C)** Cultures of the indicated cell lines were serum starved and transferred to suspension culture conditions and stimulated. Netrin-1 signaling was analyzed by SDS-PAGE and western blotting for phospho-ERK, ERK, and tubulin. **D)** Suspension cultures of OVCAR8, or knock out derivatives, were harvested and analyzed for relative phosphorylation levels of ERK by western blotting. Total ERK and Tubulin blotting serves as expression and loading controls. **E)** Netrin-1 signaling in OVCAR8 cells deleted for all UNC5 receptors (UNC5 4KO) was tested by serum starving cells, transferring them to suspension, and stimulating with Netrin-1 as before. Western blotting for phospho-ERK, ERK, and tubulin were as before. **F)** OVCAR8 cells were seeded in suspension culture and treated with the MEK inhibitor PD184352 (PD) or DMSO vehicle for up to 72 hours. Mean viability was compared by anova (***$P<0.005$). **G)** Model summarizing the roles of Netrin ligands, receptors, and downstream targets MEK and ERK in dormant survival signaling.

Figure 5. Netrin ligand overexpression is associated with poor clinical outcome in HGSOC. A) TCGA RNA-seq data for HGSOC patients (TCGA PanCancer Atlas study) was used to identify high Netrin-1 or -3 and low Netrin-1 or -3 expressing patients (high expressing are above z-score 1.2 and low are below 1.2). Overall survival was used to construct Kaplan-Meier plots and survival was compared using a logrank test. **B)** OVCAR8 cells were stably transduced with lentiviral constructs to overexpress epitope tagged Netrin-1 or -3. Western blotting for Netrins, Myc-tags, and tubulin were used to determine relative expression levels of both Netrins in these cell populations and vector controls. **C)** Control and Netrin overexpressing cells were transferred to suspension culture conditions to form spheroids, and replated to assay for viability. Mean viability and standard deviation is shown for each. Anova was used to compare survival (* P<0.05, *** P<0.001). **D)** Reattached spheroids were fixed and stained with Crystal Violet to examine size and abundance in control and Netrin-1 or -3 overexpression.

Figure 6. Netrin overexpression causes increased dissemination of tumor nodels. A) Netrin overexpressing and control OVCAR8 cells were injected into the intraperitoneal space of female NOD/SCID mice. Mice were euthanized following 35 days and analyzed for disease burden by necropsy and histopathology. **B)** The spread of cancer to the diaphragm, liver, omentum, and mesometrium was determined from necropsies and colors used in the anatomical schematic correspond with petal plots for each genotype of cells. Petal plots illustrate frequency of mice bearing disease spread to a particular location. The radius of color fill is proportional to the total number of animals with tumor nodules found in that location. **C)** Photographs of necropsy findings in the mesometrium. Locations of ovaries (*), oviduct (yellow arrow), and uterine horn (green arrow) are indicated in each case. Tumor nodules are indicated by white arrows. **D)** The number of mesometrium associated tumor nodules was determined for each mouse. Mean values are indicated and differences between genotype were determined by anova (* P<0.05, ** P<0.01). **E)** Photographs of necropsy findings in the liver. Location of sternum is indicated (#) in each image. Tumor nodules are indicated by white arrows. **F)** The number of liver associated tumor nodules was determined for each mouse. Mean values are indicated and differences between genotype were determined by anova (* P<0.05, ** P<0.01). **G** and **H)** Histology of mesometrial tumor nodules are shown. Serial sections were stained with H&E, or with the indicated antibodies for immunohistochemistry. Ovaries are indicated (*), as are the oviduct (yellow arrow), the uterus (green arrow), and the tumor nodule (white arrow). Scale bar = 2mm