Elimination of subtelomeric repeat sequences exerts little

effect on telomere essential functions in *Saccharomyces*

cerevisiae

Can Hu[1], Xue-Ting Zhu[1], Ming-Hong He[1], Yangyang Shao[2], Zhongjun Qin[2], Zhi-Jing Wu[1,3*] and

Jin-Qiu Zhou[1,3*]

[1] The State Key Laboratory of Molecular Biology, CAS Center for Excellence in Molecular Cell

Science, Shanghai Institute of Biochemistry and Cell Biology, Chinese Academy of Sciences;

University of Chinese Academy of Sciences, Shanghai, 200031, China;

[2] Key Laboratory of Synthetic Biology, CAS Center for Excellence in Molecular Plant Sciences,

Shanghai Institute of Plant Physiology and Ecology, Chinese Academy of Sciences; University

of Chinese Academy of Sciences, Shanghai, 200032, China;

[3] Key Laboratory of Systems Health Science of Zhejiang Province, Hangzhou Institute for

Advanced Study, University of Chinese Academy of Sciences, Hangzhou, 310024, China.

*To whom correspondence should be addressed. Tel: +86-21-54921076; Fax: +86-21-

54921075; E-mail: jqzhou@sibcb.ac.cn or wuzhijing6@sibcb.ac.cn

ABSTRACT

Telomeres, which are chromosomal end structures, play a crucial role in maintaining genome stability and integrity in eukaryotes. In the baker's yeast *Saccharomyces cerevisiae*, the X- and Y'-elements are subtelomeric repetitive sequences found in all thirty-two and seventeen telomeres, respectively. While the Y'-elements serve as a backup for telomere functions in cells lacking telomerase, the function of the X-elements remains unclear. This study utilized the *S. cerevisiae* strain SY12, which has three chromosomes and six telomeres, to investigate the role of X-elements (as well as Y'-elements) in telomere maintenance. Deletion of Y'-elements (SY12$^{Y\Delta}$), X-elements (SY12$^{XY\Delta+Y}$), or both X- and Y'-elements (SY12$^{XY\Delta}$) did not impact the length of the terminal TG$_{1-3}$ tracks or telomere silencing. However, inactivation of telomerase in SY12$^{Y\Delta}$, SY12$^{XY\Delta+Y}$, and SY12$^{XY\Delta}$ cells resulted in cellular senescence and the generation of survivors. These survivors either maintained their telomeres through homologous recombination-dependent TG$_{1-3}$ track elongation or underwent microhomology-mediated intra-chromosomal end-to-end joining. Our findings indicate the non-essential role of subtelomeric X- and Y'-elements in telomere regulation in both telomerase-proficient and telomerase-null cells and suggest that these elements may represent remnants of *S. cerevisiae* genome evolution. Furthermore, strains with fewer or no subtelomeric elements exhibit more concise telomere structures and offer potential models for future studies in telomere biology.

INTRODUCTION

Telomeres, specialized nucleoprotein structures located at the end of linear chromosomes in eukaryotic cells, are crucial for maintaining genomic stability and protecting chromosomal ends from being perceived as DNA breaks (Wellinger and Zakian, 2012). In the budding yeast *Saccharomyces cerevisiae*, telomeric DNA consists of approximately ~300±75 base pairs of $C_{1-3}A/TG_{1-3}$ repeats with a 3' G-rich single-stranded overhang (Wellinger and Zakian, 2012). Adjacent to the telomeric TG_{1-3} repeats, there are subtelomeric repeat elements known as X- and Y'-elements, which vary between telomeres, as well as strains (Chan and Tye, 1983a, b; Louis, 1995). The Y'-elements, immediately internal to the telomeric repeats, are present as a tandem array of zero to four copies, they fall into two major size classes, 6.7 kb Y'-long (Y'-L) and 5.2 kb Y'-short (Y'-S) (Chan and Tye, 1983a, b). Y'-elements are highly conserved with only ~2% divergence between strains (Louis and Haber, 1992). One entire Y'-element contains two large open reading frames (ORFs), an ARS consensus sequence (ACS) and a STAR element (Subtelomeric anti-silencing regions) consisting of binding sites for Tbf1 and Reb1 (Chan and Tye, 1983a, b; Fourel et al., 1999; Louis and Haber, 1992). The X-element, a much more heterogeneous sequence abutting Y'-elements or telomeric repeats, contains the 473 bp "core X" sequence and the subtelomeric repeats (STRs) A, B, C and D (Louis and Haber, 1991; Louis et al., 1994). The STRs are found in some chromosome ends, while the "core X" sequence is shared by all chromosomes. Recent long-read sequencing shows that subtelomeric regions display high evolutionary plasticity and are rich in a various of structure

variants such as reciprocal translocations, transpositions, novel insertions, deletions and

duplications (O'Donnell et al., 2023).

Telomeric DNA elongation primarily relies on telomerase, an enzyme comprising a reverse

transcriptase, an RNA component, and accessory factors (Palm and de Lange, 2008; Wellinger

and Zakian, 2012). In *S. cerevisiae*, the telomerase holoenzyme consists of the reverse

transcriptase Est2, the RNA template TLC1, and accessory factors Est1, Est3,

Pop1/Pop6/Pop7 proteins (Lemieux et al., 2016; Lendvay et al., 1996; Lundblad and Szostak,

1989; Singer and Gottschling, 1994). In the absence of telomerase, homologous recombination

can take place to replicate telomeres, resulting in telomerase-deficient "survivors" (Lundblad

and Blackburn, 1993; Teng and Zakian, 1999). These survivors are broadly categorized into

Type I and Type II based on distinct telomere structures (Lundblad and Blackburn, 1993; Teng

and Zakian, 1999). Type I survivors possess tandem amplified Y'-elements (both Y'-L and Y'-

S) and very short TG$_{1-3}$ tracts, indicating that Y'-elements serve as substrates for homologous

recombination. Type II survivors display long heterogeneous TG$_{1-3}$ tracts. On solid medium,

approximately 90% of the survivors are Type I, while 10% are Type II (Teng et al., 2000).

However, in liquid culture, Type II survivors grow faster and eventually dominate the population

(Teng and Zakian, 1999). The proteins required for Type I and Type II survivor formation appear

to be different. Type I survivors depend on Rad51, Rad54, Rad55, Rad57, and Pif1 (Chen et

al., 2001; Hu et al., 2013; Le et al., 1999). while the formation of Type II survivors requires the

Mre11/Rad50/Xrs2 (MRX) complex, KEOPS complex, Rad59, Sgs1, and Rad6, most of which

are critical for DNA resection (Chen et al., 2001; He et al., 2019; Hu et al., 2013; Johnson et al.,

2001; Le et al., 1999; Louis, 2001; Nicolette et al., 2010; Teng et al., 2000; Wellinger and Zakian,

2012; Wu et al., 2017). Although Type I and Type II pathway are working independently, Kockler

et al. found that the proteins involved in each pathway can work together via two sequential

steps and contribute to a unified ALT (Alternative lengthening of telomeres) process (Kockler

et al., 2021).

The amplification of Y'-elements represents a significant feature of telomere recombination

in telomerase-null Type I survivors (Lundblad and Blackburn, 1993; Teng and Zakian, 1999),

and as a result, extrachromosomal Y' circular DNAs have been observed in Type I survivors

(Larrivee and Wellinger, 2006). Additionally, Y'-element acquisition has been observed in the

initiation step of pre-senescence, suggesting a potential role for Y'-elements in Type II survivor

formation (Churikov et al., 2014). Furthermore, Y'-elements are mobilized through a

transposition-like RNA-mediated process involving Ty1 activity in telomerase-negative

survivors (Maxwell et al., 2004). Y'-elements also express potential DNA helicases, Y'-Help, in

telomerase-null survivors (Yamada et al., 1998). Thus, Y'-elements play a significant role as

donors in homologous recombination-mediated telomere maintenance. The functions of X-

elements, on the other hand, are less clear. The "core X" sequence consists of an ACS element

and, in most cases, an Abf1 binding site (Louis, 1995), and acts as a protosilencer (Lebrun et

al., 2001). In contrast, STRs and Y'-STAR possess anti-silencing properties that limit the

spreading of heterochromatin (Fourel et al., 1999). Interestingly, a previous study demonstrated

that telomeres with X-only ends (containing only X-elements) were more efficiently elongated

compared to those with X-Y' ends (containing both X- and Y'-elements) in *tel1Δ rif1Δ* strains

(Craven and Petes, 1999). Moreover, subtelomeric elements (including X-elements) and

associated factors like Reb1 and Tbf1 antagonize telomere anchoring at the nuclear envelope

(Hediger et al., 2006). However, considering that X-elements are present in all telomeres while

Y'-elements are not, the specific functions of X- and Y'-elements in genome integrity after the

evolution of telomerase have long been a subject of questioning (Jager and Philippsen, 1989;

Zakian and Blanton, 1988).

In wild-type yeast strain BY4742, there are eight Y'-S and eleven Y'-L elements at the thirty-

two telomere loci. Additionally, each telomere locus contains one X-element. The genetic

deletion of all X- and Y'-elements to directly investigate the roles of X- and Y'-elements in

genome integrity is a challenging and complex task. In this study, we utilized recently reported

chromosome-fused budding yeast strains (Shao et al., 2018) to eliminate both X- and Y'-

elements completely. This approach allows us to reinvestigate the roles of X- and Y'-elements

at telomeres.

RESULTS

Telomere recombination in telomerase-null chromosome-fused yeast strains SY1 to

SY12

The functions of Y'-elements have been previously linked to telomere recombination (Churikov

et al., 2014; Larrivee and Wellinger, 2006; Lundblad and Blackburn, 1993; Teng and Zakian,

1999). To further investigate the role of Y'-elements in telomere recombination, we utilized a

series of chromosome-fused budding yeast strains derived from the wild-type BY4742 strain,

including SY1, SY3, SY5, SY7, SY8, SY9, SY10, SY11, SY12 and SY13 (also referred to as

SYn for convenience) (Figure 1A) (Shao et al., 2018). The remaining subtelomeric elements in

SY8 to SY13 strains are listed in supplementary file 2. We excluded SY14 from these

experiments since the presence of circular chromosome was prominent in SY14 *tlc1Δ* cells

(one fused chromosome) (Wu et al., 2020), We generated haploid SYn *tlc1Δ TLC1* strains by

deleting the chromosomal copy of the *TLC1* gene and introducing a plasmid-borne wild-type

TLC1 gene (pRS316-*TLC1*). Clones that lost the pRS316-*TLC1* plasmid (containing the *URA3*

marker) were identified upon counter-selection on 5'-fluoroorotic-acid (5'-FOA) plates and were

subsequently re-streaked on YPD plates for at least 9 cycles for survivor formation (referred to

as the "multiple-colony streaking assay" in Methods). The telomere patterns of the survivors

were then determined through Southern blotting assay (Figure 1B-D).

The canonical telomerase-independent survivors can be broadly categorized into two types:

Type I and Type II survivors, based on the restriction fragments generated after XhoI digestion

(Lundblad and Blackburn, 1993; Teng and Zakian, 1999). Type I survivors exhibit tandem

duplication of Y'-elements and very short TG_{1-3} tracts, while Type II survivors contain long

heterogeneous TG_{1-3} sequences. Consistent with previous reports, BY4742 *tlc1Δ* cells

generated both Type I (subtelomeric Y'-element recombination) and Type II (TG_{1-3}

recombination) survivors (Figure 1B) (Hu et al., 2013). Intriguingly, as the number of

chromosomes decreased, the frequency of Type II survivors gradually diminished, while Type

I survivors became the predominant type (Figure 1B-D). Furthermore, non-canonical survivors

with distinct patterns from Type I or Type II emerged in SY9 *tlc1Δ* (six chromosomes), SY10

tlc1Δ (five chromosomes), SY11 *tlc1Δ* (four chromosomes), SY12 *tlc1Δ* (three chromosomes)

and SY13 *tlc1Δ* (two chromosomes) (Figure 1C and 1D indicated by triangles at the bottom of

the panels). Notably, the Y'-telomere band of ~1.2 kb was not detected in two clones of SY11

tlc1Δ cells (clones 2 and 5), the majority of clones of SY12 *tlc1Δ* cells (except for clones 9, 14,

and 15), and the majority of clones of SY13 *tlc1Δ* cells (except for clones 1, 4, 8 and10) (Figure

1D). We speculate that either the Y'-elements have eroded or the chromosomal ends containing

Y'-elements have fused with other ends in these non-canonical survivors. These findings

suggest that the ratio of survivor types is influenced by the number of chromosomes.

Figure 1. Telomere structures in SYn *tlc1*Δ survivors. Telomere Southern blotting assay was performed

to examine telomere structure. The genomic DNA extracted from BY4742 (wild type) and SYn strains

(labeled on top) was digested with XhoI and subjected to Southern hybridization with a TG$_{1-3}$ probe. (**A**)

Telomerase-proficient strains (labeled on top), whose chromosome numbers are labeled at the bottom.

Two independent clones of each strain were examined. (**B-D**) SYn *tlc1*Δ survivors generated on plates.

Four (BY4742 *tlc1*Δ) and fifteen (SYn *tlc1*Δ) individual survivor clones (labeled on top of each panel) of

each strain were examined. "+" at the bottom indicates Type I survivors. "Δ" marks the survivors which

are non-canonical Type I or Type II.

Characterizing the survivor pattern in SY12

To determine the chromosomal end structures of the non-canonical survivors shown in Figure

1, we selected SY12 *tlc1*Δ survivors for further analysis. In the SY12 strain, there are six

telomeres corresponding to the native chromosomes I-L, X-R, XIII-L, XI-R, XVI-L, and XIV-R.

We employed Southern blotting after NdeI digestion to validate the telomere and subtelomere

structures (Figure 2—figure supplement 1A). The results revealed that, in the SY12 strain used

in our study, only the XVI-L telomere contained a single copy of the Y'-element, while all

telomeres harbored X-elements (Figure 2—figure supplement 1B). For simplicity, we referred

to the chromosomes containing the original I, XIII, and XVI as chromosome 1, 2, and 3,

respectively (Figure 2A, left panel).

We conducted a re-examination of telomere recombination upon telomerase inactivation in

SY12 cells. Deletion of *TLC1* in SY12 cells resulted in cell senescence, and different clones

recovered at various time points in liquid medium (Figure 2B). Telomere Southern blotting

analysis showed progressive shrinking of the telomeric XhoI fragments over time, and TG_{1-3}

recombination occurred to maintain telomeres (Figure 2C). Since the liquid culture contained a

mixture of different colonies, we employed a multiple-colony streaking assay and Southern

blotting analysis to examine the telomere patterns of fifty independent SY12 *tlc1Δ* survivors

(Figure 2D and Figure 2—figure supplement 2). Among these survivors, eight clones (labeled

in red, 16% of the survivors tested) exhibited the typical Type I telomere structure characterized

by Y'-element amplification (Figure 2D and 2E). This was confirmed by Southern blotting

analysis using a Y' probe (Figure 2—figure supplement 2). The emergence of Type I survivor

in SY12 strain which only contain one Y'-element indicates that multiple Y'-elements in tandem

are not strictly required for Type I formation. Clone 1 (labeled in orange, 2% of the survivors

tested) displayed heterogeneous telomeric TG_{1-3} tracts (Figure 2D and 2E), indicating it was a

Type II survivor. This was further confirmed by restoring the telomere length to the level

observed in SY12 cells through the re-introduction of the *TLC1* gene into one representative

clone (named SY12 *tlc1Δ*-T1) and subsequent passaging on yeast complete (YC) medium

lacking uracil (Ura-) for 20 cycles (Figure 2—figure supplement 3A).

Notably, ten of the examined clones (labeled in blue, 20% of the survivors tested) displayed

no telomere signals associated with canonical Type I or Type II survivors (Figure 2D and 2E).

Their hybridization patterns were strikingly similar to those of SY14 *tlc1Δ* survivors (Wu et al.,

2020), which survived through intra-chromosomal circularization. To investigate whether the

three chromosomes in these SY12 *tlc1*Δ survivors had undergone intra-chromosomal fusions,

we selected a clone, namely SY12 *tlc1*Δ-C1 and performed PCR-mapping assay to determine

the erosion points of each chromosome end, as previously described (Wu et al., 2020). A PCR

product of the predicted length would be obtained only if the corresponding chromosome region

was intact. The PCR-mapping assay precisely identified the borders of telomere erosion for the

three chromosomes in SY12 *tlc1*Δ-C1 cells. For chromosome 1 (Figure 2A, left panel), the

chromosome regions approximately 3.3 kb and 1.9 kb proximal to telomere I-L and X-R,

respectively, had been lost (Figure 2—figure supplement 4 and supplementary file 3).

Regarding chromosome 2, the terminal ~3.8 kb of telomere XIII-L and ~2.5 kb of telomere XI-

R remained intact (Figure 2—figure supplement 4 and supplementary file 3). For chromosome

3, the terminal ~0.1 kb of telomere XVI-L was intact, while the terminal ~3.4 kb of telomere XIV-

R was preserved (Figure 2—figure supplement 4 and supplementary file 3). To confirm the

chromosome fusion events, we performed PCR-sequencing analysis. If a given pair of primers,

oriented to different chromosome ends, produced PCR products, it indicated that the

corresponding arms had fused. The results revealed that the three chromosomes in SY12

*tlc1*Δ-C1 cells had undergone intra-chromosomal fusions through microhomology-mediated

end joining (MMEJ) (Wu et al., 2020), resulting in the formation of circular chromosomes (Figure

2F and supplementary file 3). Notably, the fusion junctions of the three chromosomes in SY12

*tlc1*Δ-C1 cells differed in nucleotide sequence and length (22 bp, 8 bp, and 5 bp in

chromosomes 1, 2, and 3, respectively). Moreover, the sequences involved in the ends-fusion

were not perfectly complementary (Figure 2F). For example, the fusion sequence of

chromosome 3 was 5 bp long and contained one mismatch. To further verify the chromosome

structure in the "circular survivors" SY12 *tlc1*Δ-C1 (Figures 2F), we performed the pulsed-field

gel electrophoresis (PFGE) analysis. Control strains included SY12 (three linear chromosomes)

and SY15 (one circular chromosome). The PFGE result confirmed that like the single circular

chromosome in SY15 cells, the circular chromosome in the SY12 *tlc1*Δ-C1 survivors couldn't

enter the gel, while the linear chromosomes in SY12 were separated into distinct bands, as

expected (Figure 2—figure supplement 5). Thus, the survivors shown in Figure 2D, which

displayed an identical hybridization pattern to the SY12 *tlc1*Δ-C1 clone, were all likely "circular

survivors". Consistently, the telomere signals detected in the SY12 strain were still not observed

in the SY12 *tlc1*Δ-C1 survivor after reintroducing a plasmid-borne wild-type *TLC1* gene (Figure

2—figure supplement 3B).

Twelve clones of SY12 *tlc1*Δ survivors (labeled in green, 24% of the survivors tested)

exhibited no Y'-telomere signals compared to SY12 cells but displayed different lengths of TG_{1-3}

tracts (Figure 2D and 2E). Due to their non-canonical telomere structures, characterized by

the absence of both Y'- amplification and superlong TG_{1-3} sequences, we designated these

SY12 *tlc1*Δ survivors (labeled in green, Figure 2D) as Type X. In Type X survivors, the DNA

bands with sizes of approximately 2.3 kb, 5.1 kb, 15.3 kb, 18.5 kb, and 21.9 kb were roughly

comparable to the telomeres of XI-R, X-R, I-L, XIII-L, and XIV-R in SY12 cells (indicated on the

left in the panel). The newly emerged band at approximately 4.3 kb likely originated from the

XVI-L telomere (indicated by the red arrow on the right in the panel) (Figure 2D), where the Y'-

elements had been eroded, leaving only the TG_{1-3} tracts at the very ends (Figure 2A, right

panel). It remains unclear whether Y'-element erosion is common in telomerase-null BY4742

type II survivors. However, in SY12 *tlc1Δ* cells, the remaining single copy of the Y'-element

couldn't find homology sequences to repair telomeres, whereas the multicopy X-element could

easily find homology sequences to repair telomeres and form the type X survivors. To verify

this notion, we reintroduced the *TLC1* gene into one representative clone (named SY12 *tlc1Δ*-

X1) and examined the telomere length. As expected, the telomeres of X-R and XI-R were

restored to the lengths observed in wild-type SY12 cells, and accordingly, the newly emerged

4.3 kb band was also elongated (Figure 2G). Given that the restriction fragments of telomeres

I-L (15.3 kb), XIII-L (18.5 kb), and XIV-R (21.9 kb) were quite long, detecting minor changes in

telomere length was challenging under the assay conditions of Southern blotting. To determine

the chromosomal end structure of the Type X survivor, we randomly selected a typical

Type X survivor, and performed PCR-sequencing analysis. The results revealed the intact

chromosome ends for I-L, X-R, XIII-L, XI-R, and XIV-R, albeit with some mismatches

compared with the *S. cerevisiae* S288C genome (http://www.yeastgenome.org/), which

possibly arising from recombination events that occurred during survivor formation. Notably,

the sequence of the Y'-element in XVI-L could not be detected, while the X-element

remained intact (Figure 2—figure supplement 6). These data indicated that Type X survivors

possess linear chromosomes with telomeres terminating in TG_{1-3} repeats, while the Y'-element

has been eroded (Figure 2A, right panel). Consistently, no Y' signals were detected in these

twelve Type X survivors (labeled in green, Figure 2—figure supplement 2), suggesting that the

Y'-element has not been translocated to other telomeres and is not essential for yeast cell

viability.

In addition to the aforementioned Type I, Type II, circular, and Type X survivors, there were

some clones (labeled in black, 38% of the survivors tested) which exhibited non-uniform

telomere patterns and were not characterized (Figure 2D and Figure 2E). We speculated that

combinations of diverse mechanisms were occurring within each "uncharacterized survivor".

For instance, in the case of two survivors (clones 9 and 18, Figure 2D) in which only one

hybridization signal could be detected, pointing to the possibility that two chromosomes

underwent intra-chromosomal fusions while one retained its ends through TG_{1-3} recombination.

However, the sizes of the two telomere restriction fragments on the linear chromosome were

too close to be distinguished and separated, resulting in only one hybridization signal.

Alternatively, it is also plausible that three chromosomes experienced intra-chromosomal

fusions, with one fusion point containing TG_{1-3} repeats. For the uncharacterized clones 4, 5, 7,

15, and 43, they exhibited significant amplification of TG_{1-3} sequences, and the telomeres of

these survivors did not resolve into distinct bands (Figure 2D). We hypothesize that the

observed telomere patterns in these survivors could be attributed to extensive TG_{1-3}

recombination. However, we cannot exclude the possibility of coexisting diverse mechanisms

within a survivor, such as telomere elongation through TG_{1-3} amplification, as well as intra- and

inter-chromosomal fusions. Since we couldn't figure out the telomere structures in these

survivors, we classified them as "uncharacterized survivors".

To further determine the genetic requirements for survivors in SY12, we constructed the

SY12 *tlc1Δ rad52Δ* pRS316-*TLC1* strain. The plasmid-borne wild-type *TLC1* gene (pRS316-

TLC1) was counter-selected on 5'-FOA plates. SY12 *tlc1Δ rad52Δ* cells were measured by the

cell viability assay (see Materials and methods). The results showed double deletion of *TLC1*

and *RAD52* in SY12 strain could slightly accelerate senescence, and SY12 *tlc1Δ rad52Δ*

survivors could be generated but took much longer to recover than the SY12 *tlc1Δ* survivors

(Figure 2—figure supplement 7A), suggesting that Rad52 is not strictly required for survivor

generation in the SY12 strain in liquid. We also passaged SY12 *tlc1Δ rad52Δ* cells on solid

medium until survivor emerged. Southern blotting of twenty-five clones revealed that neither

Type I nor Type II survivors were found, and instead circular survivors except clone 20 were

obtained (labeled in blue, Figure 2—figure supplement 7B). We conclude that the formation of

circular survivors in the SY12 *tlc1Δ rad52Δ* strain is mediated by MMEJ as observed in the

SY14 *tlc1Δ rad52Δ* strain (Wu et al., 2020), but not *RAD52* mediate pathways. Since no Type

X survivor was detected in SY12 *tlc1Δ rad52Δ* strain, we constructed the SY12 *tlc1Δ rad51Δ*

pRS316-*TLC1* and SY12 *tlc1Δ rad50Δ* pRS316-*TLC1* strain to investigate on which pathway

the formation of the Type X survivor relied. After being counter-selected on 5'-FOA plates, cells

were passaged on solid medium until survivor arose. Southern blotting assay indicated the

emergence of Type X survivors even in the absence of Rad51 (labeled in green, clone 2, 5,

11 and 18, Figure 2—figure supplement 8A). In contrast, no Type X survivor was detected

in the SY12 *tlc1Δ rad50Δ* strain (Figure 2—figure supplement 8B). These data suggest that

the formation of the Type X survivor depends on Rad50-mediated Type II pathway.

Taken together, our results indicate that telomerase inactivation in SY12 cells leads to cell

senescence and the emergence of survivors with diverse telomere patterns, including Y'-

amplification (Type I), elongated TG$_{1-3}$ tracts (Type II), intra-chromosomal end-to-end joining

(circular), Y'- loss (Type X), and uncharacterized.

Figure 2

Figure 2. Survivor formation in SY12 *tlc1*Δ strain. (**A**) Schematic representation of chromosome (and

telomere) structures (not drawn to scale) in the SY12 strain (left panel) and the Type X survivor (right

panel). The Roman numerals, native chromosomes; the Arabic numerals on the left, chromosome

numbers of SY12; yellow box, X-element; green box, Y'-element; tandem grey triangles, telomeres;

black circles, centromere; vertical arrows and numbers, positions and lengths of the terminal XhoI

digestion fragments detected by the telomeric TG_{1-3} probe. Chromosome numbers are omitted in the

Type X survivor (right panel). (**B**) Cell viability assay in liquid medium. The growth of SY12 (labeled in

black) and SY12 *tlc1*Δ (three clones labeled in blue, purple and green respectively) strains were

monitored every 24 hr for 12 days. (**C**) Telomeric Southern blotting assay of SY12 *tlc1*Δ survivors.

Genomic DNAs prepared from SY12 *tlc1*Δ survivors assayed in (B) were digested with XhoI and

subjected to Southern blotting with a TG_{1-3} probe. (**D**) Telomere Southern blotting assay of SY12 *tlc1*Δ

survivors obtained on solid medium. Genomic DNA from fifty independent SY12 *tlc1*Δ clones (labeled on

top) was digested with XhoI and hybridized to a telomere-specific TG_{1-3} probe. Type II survivors: in

orange; Type I survivors: in red; circular survivors: in blue; Type X survivors: in green; uncharacterized

survivors: in black. Theoretical telomere restriction fragments of the SY12 strain are indicated on the left.

The red arrows indicate the new band of about 4.3 kb emerged in Type X survivors. The asterisks

indicate the non-specific bands. Genomic DNA stained with Gelred was used as a relative loading

control (LC). (**E**) The ratio of survivor types in SY12 *tlc1*Δ strain. n=50; Type I, in red; Type II, in orange;

Type X, in green; uncharacterized survivor, in grey; circular survivor, in blue. (**F**) Schematic of three

circular chromosomes and fusion sequences in the SY12 *tlc1*Δ-C1 survivor. The sequence in blue

indicates the sequences of X-R, XI-R or XIV-R, the sequence in red indicates the sequences of I-L, XIII-

L or XVI-L. Bases in green are mis-paired. The numbers above or below the schematic line

(chromosome) indicate the distance to the corresponding telomeres. (**G**) Telomere Southern blotting

analysis of an SY12 *tlc1*Δ Type X survivor at the 20[th] re-streak after *TLC1* reintroduction. The red arrows

indicate the new band of about 4.3 kb emerged in Type X survivors. LC: loading control.

Figure 2—figure supplement 1. Characterization of SY12 strain. (**A**) Schematic representation of

chromosome (and telomere) structures (not drawn to scale) in the SY12 strain. The Roman numerals,

native chromosomes; the Arabic numerals on the left, chromosome numbers of SY12; yellow box, X-

element; green box, Y'-element; tandem grey triangles, telomeres; black circles, centromere; vertical

arrows and numbers, positions and lengths of the terminal NdeI digestion fragments detected by the

telomeric TG$_{1-3}$ probe. (**B**) Southern blotting analysis of telomere length in BY4742 and SY12 (labeled

on top) cells. Genomic DNA prepared from two independent clones of BY4742 and SY12 strains was

digested with NdeI, and then subjected to Southern blotting with a TG$_{1-3}$ probe (left panel). The numbers

in brackets indicate the telomere length of the corresponding chromosomes. The blot was then stripped

and reprobed with a Y' probe (right panel). The asterisk indicates the non-specific band.

Figure 2—figure supplement 2. Telomere Southern blot with a Y'-element probe examining SY12

*tlc1*Δ survivors. The blot in Figure 2D was then stripped and reprobed with a Y' probe.

Figure 2—figure supplement 3. Southern blotting results of reintroduce *TLC1* into SY12 *tlc1*Δ

survivors. (**A**) Southern blotting result of SY12 *tlc1*Δ Type II survivor at the 20[th] re-streaks after *TLC1*

reintroduction. LC: loading control. (**B**) Southern blotting result of SY12 *tlc1*Δ circular survivor at the 20th

re-streaks after *TLC1* reintroduction. LC: loading control.

Figure 2—figure supplement 4. Borders of erosion of the SY12 *tlc1*Δ-C1 survivor are defined by PCR

mapping. (**A**) Upper panel, a schematic diagram of the subtelomeric region of 0-60 kb proximal to I-L

telomere is shown. Primer pairs (No. 1^L to 17^L) are aligned and indicated at their corresponding

subtelomeric loci. Lower panel, the genotype is listed on the left, and primer pairs are listed on top; solid

circles mean positive PCR products and open circles mean no PCR products with corresponding primer

pairs. (**B**) Upper panel, a schematic diagram of the subtelomeric region of 0-50 kb proximal to X-R

telomeric TG_{1-3} sequence is shown. Primer pairs (No. 1^R to 42^R) are aligned and indicated at their

corresponding subtelomeric loci. Lower panel, the genotype is listed on the left, and primer pairs are

listed on top. (**C**) Upper panel, a schematic diagram of the subtelomeric region of 0-10 kb proximal to

XIII-L telomeric TG_{1-3} sequence is shown. Primer pairs (No. 1^L to 6^L) are aligned and indicated at their

corresponding subtelomeric loci. Lower panel, the genotype is listed on the left, and primer pairs are

listed on top. (**D**) Upper panel, a schematic diagram of the subtelomeric region of 0-65 kb proximal to XI-

R telomeric TG_{1-3} sequence is shown. Primer pairs (No. 1^R to 25^R) are aligned and indicated at their

corresponding subtelomeric loci. Lower panel, the genotype is listed on the left, and primer pairs are

listed on top. (**E**) Upper panel, a schematic diagram of the subtelomeric region of 0-35 kb proximal to

XVI-L telomeric TG_{1-3} sequence is shown. Primer pairs (No. 1^L to 29^L) are aligned and indicated at their

corresponding subtelomeric loci. Lower panel, the genotype is listed on the left, and primer pairs are

listed on top. (**F**) Upper panel, a schematic diagram of the subtelomeric region of 0-60 kb proximal to

XIV-R telomeric TG_{1-3} sequence is shown. Primer pairs (No. 1^R to 21^R) are aligned and indicated at their

corresponding subtelomeric loci. Lower panel, the genotype is listed on the left, and primer pairs are

listed on top.

Figure 2—figure supplement 5. PFGE result of circular survivors. Chromosomal DNA analysis of

"circular survivors" SY12 *tlc1*Δ-C1, SY12^{YΔ} *tlc1*Δ-C1, SY12^{XYΔ} *tlc1*Δ-C1, SY12$^{XYΔ+Y}$ *tlc1*Δ-C1 by PFGE.

The *S. cerevisiae* strain SY15 (with a single circular chromosome) was used as control. As marker, the

size of three chromosomes in wild type *S. pombe* strain is indicated on right.

Figure 2—figure supplement 6. Telomere structure determination of Type X survivor. Schematic

representation of chromosome (and telomere) structures (not drawn to scale) in the Type X survivor.

The Roman numerals, native chromosomes; the Arabic numerals on the left, chromosome numbers of

SY12; yellow box, X-element; green box, Y'-element; tandem grey triangles, telomeres; black circles,

centromere. The sequence in yellow box belongs to X-element. Bases in rad are mis-paired. The dotted

line in green represents the sequence of Y'-element is lost.

Figure 2—figure supplement 7. Survivor formation in SY12 *tlc1Δ rad52Δ* strain. (**A**) Cell viability assay

in liquid culture. The growth of SY12 (labeled in black), SY12 *tlc1Δ* (two clones labeled in purple and

blue respectively) and SY12 *tlc1Δ rad52Δ* (three clones labeled in violet, pink and carmine respectively)

were monitored every 24 hr for 12 days. (**B**) Telomere Southern blotting assay of SY12 *tlc1Δ rad52Δ*

survivors obtained on solid medium. Genomic DNA from twenty-five independent clones (labeled on top)

was digested with XhoI and hybridized to a telomere-specific $TG_{1–3}$ probe. Circular survivors: in blue;

uncharacterized survivors: in black. Theoretical telomere restriction fragments of the SY12 strain are

indicated on the left. The asterisks indicate the non-specific bands. Genomic DNA stained with Gelred

was used as a relative loading control (LC).

Figure 2—figure supplement 8. Southern blotting result of SY12 *tlc1Δ rad51Δ* and SY12 *tlc1Δ rad50Δ*

survivors. (**A**) Genomic DNAs of twenty-five independent SY12 *tlc1Δ rad51Δ* clones were digested with

XhoI and subjected to Southern blotting with a TG_{1-3} probe. Circular survivors: in blue; Type X survivors:

in green; uncharacterized survivors: in black. Theoretical telomere restriction fragments of the SY12

strain are indicated on the left. The red arrows indicate the new band of about 4.3 kb emerged in Type X

survivors. The asterisks indicate the non-specific bands. Genomic DNA stained with Gelred was used as

a relative loading control (LC). (**B**) Genomic DNAs of twenty-five independent SY12 *tlc1Δ rad50Δ* clones

were digested with XhoI and subjected to Southern blotting with a TG_{1-3} probe. Type I survivors: in rad;

Circular survivors: in blue; uncharacterized survivors: in black. Theoretical telomere restriction fragments

of the SY12 strain are indicated on the left.

Deletion of all of the X- and Y'-elements in the SY12 strain

We aimed to determine whether the subtelomeric X-elements are dispensable or not. In the

SY12 strain, there are six X-elements distributed among six telomeres (Figure 2A, left panel).

To precisely delete all X- and Y'-elements in SY12 strains, we employed a method that

combines the efficient CRISPR-Cas9 cleavage system with the robust homologous

recombination activity of yeast, as previously described (Shao et al., 2018; Shao et al., 2019).

Briefly, the Cas9 nuclease cleaved the unique DNA sequences adjacent to the subtelomeric

region (site S1) with the guidance of gRNA1. The resulting chromosome break was repaired

through homologous recombination (HR) using the provided chromosome ends which

excluding the X- and Y'-elements. Subsequently, the *URA3* marker and the guide RNA

expression plasmid (pgRNA) were eliminated by inducing gRNA2 expression on pCas9 using

galactose (Figure 3—figure supplement 1). This approach allowed us to initially delete the Y'-

element and X-element in XVI-L, generating the SY12$^{Y\Delta}$ strain (Figure 3A and supplementary

file 4). Subsequently, through five successive rounds of deletions, we removed all remaining X-

elements, resulting in the SY12$^{XY\Delta}$ strain (Figure 3A and supplementary file 4). To confirm the

series of deletions, we performed PCR analysis using a primer located within the deletion region

and another primer annealing upstream of the region (indicated by purple arrows in Figure 3—

figure supplement 1). This analysis verified the complete deletion of the subtelomeric X- and

Y'-elements (Figure 3B, rows 3 to 7). Additionally, we conducted a separate PCR analysis using

primers specific to either X- or Y'-elements, which confirmed the absence of both X- and Y'-

elements in the SY12$^{XY\Delta}$ strain (Figure 3B, rows 8). Subsequently, we inserted a Y'-long

element (cloned from the native XVI-L sequence, which does not contain the centromere-

proximal short telomere sequence) into the left arm of chromosome 3 in the SY12$^{XY\Delta}$ strain,

resulting in the SY12$^{XY\Delta+Y}$ strain containing a single Y'-element but no X-element (Figure 3A

and supplementary file 4). The successful insertion was confirmed by PCR analysis (Figure 3B,

lane 9).

Figure 3. Characterization of SY12$^{Y\Delta}$, SY12$^{XY\Delta}$ and SY12$^{XY\Delta+Y}$ strains. (**A**) Schematic of chromosome

structures in the SY12, SY12$^{Y\Delta}$, SY12$^{XY\Delta}$ and SY12$^{XY\Delta+Y}$ strains. Yellow box, X-element; green box, Y'-

element; tandem grey triangles, telomeres. Vertical arrows and numbers indicate the positions and sizes

of the sites and length of XhoI and PaeI-digested terminal fragments. (**B**) PCR analyses of the

engineered sites of the individual telomeres (labeled on left) in BY4742, SY12, SY12$^{Y\Delta}$, SY12$^{Y\Delta+1X\Delta}$,

SY12$^{Y\Delta+2X\Delta}$, SY12$^{Y\Delta+3X\Delta}$, SY12$^{Y\Delta+4X\Delta}$, SY12$^{XY\Delta}$ and SY12$^{XY\Delta+Y}$ strains (labeled on top). Primer

sequences for the PCR analyses are listed in supplementary file 1. *RAP1* was an internal control. (**C**)

Morphology of BY4742, SY12, SY12$^{Y\Delta}$, SY12$^{XY\Delta}$ and SY12$^{XY\Delta+Y}$ cells in the exponential growth phase

(30°C in YPD). Shown are DIC images. Scale bar, 2 μm. (**D**) Growth analysis of the SY12, SY12$^{Y\Delta}$,

SY12$^{XY\Delta}$ and SY12$^{XY\Delta+Y}$ strains. Several clones of the SY12, SY12$^{Y\Delta}$, SY12$^{XY\Delta}$ and SY12$^{XY\Delta+Y}$ strains

were re-streaked on YPD plates 61 times at intervals of two days. Shown were the 3rd, 31st and 61st re-

streaks. (**E**) Growth analysis of BY4742, SY12, SY12$^{Y\Delta}$, SY12$^{XY\Delta}$ and SY12$^{XY\Delta+Y}$ cells in liquid culture.

Error bars represent standard deviation (s.d.), n = 3. (**F**) FACS analysis of DNA content of BY4742,

SY12, SY12$^{Y\Delta}$, SY12$^{XY\Delta}$ and SY12$^{XY\Delta+Y}$ cells. (**G**) Dotting assays on YPD plates at low (24°C) and high

(37°C) temperatures, or on YPD plates containing methyl methane sulfonate (MMS), camptothecin

(CPT) or hydroxyurea (HU) at the indicated concentrations. The BY4742 *mre11Δ* haploid strain serves

as a negative control because Mre11 is involved in the repair of double-stranded breaks (Lewis et al.,

2004).

Figure 3—figure supplement 1. Schematics of CRISPR–Cas9-mediated deletion of X- and Y'-elements

on individual chromosomes in SY12 cells. The specific DNA sequences centromere-proximal to the

subtelomeric region (site S1) were cleaved by the Cas9 nuclease with the guidance of gRNA1.

Homologous recombination between the broken ends and the provided donors led to the deletion of X-

and Y'-elements. Galactose induction of gRNA2 on pCas9 caused the cleavage at site S2 and the *URA3*

marker was counter-selected on 5'-FOA plates. Deletion of X- and Y'-elements were determined by PCR

analysis with a primer located within the deletion region and another upstream of the region (indicated by

purple arrows).

───

Subtelomeric X- and Y'-elements are dispensable for cell proliferation, various stress

responses, telomere length control and telomere silencing

The SY12$^{Y\Delta}$, SY12$^{XY\Delta}$, and SY12$^{XY\Delta+Y}$ cells, cultured in YPD medium at 30°C, exhibited the

same cell morphology as the parental strains SY12 and BY4742 (Figure 3C). To assess the

stability of their genomes, we restreaked several clones of SY12$^{Y\Delta}$, SY12$^{XY\Delta}$, and SY12$^{XY\Delta+Y}$

strains on YPD plates for a total of 61 times at two-day intervals (Figure 3D). Similar to the

SY12 strain, the progeny colonies of SY12$^{Y\Delta}$, SY12$^{XY\Delta}$, and SY12$^{XY\Delta+Y}$ grew robustly on solid

medium (Figure 3D). Moreover, SY12$^{Y\Delta}$, SY12$^{XY\Delta}$, and SY12$^{XY\Delta+Y}$ cells exhibited growth rates

comparable to those of SY12 and BY4742 cells in liquid medium (Figure 3E). FACS analysis

revealed that SY12$^{Y\Delta}$, SY12$^{XY\Delta}$, and SY12$^{XY\Delta+Y}$ had the same 1C and 2C DNA content as wild-

type cells (Figure 3F), indicating that the X- and Y'-elements are not necessary for cell

proliferation under normal conditions. Additionally, the growth of SY12$^{Y\Delta}$, SY12$^{XY\Delta}$, and

SY12$^{XY\Delta+Y}$ cells at different temperatures (24 and 37°C) (Figure 3G, upper panel) closely

resembled that of SY12 and BY4742 cells. Furthermore, SY12$^{Y\Delta}$, SY12$^{XY\Delta}$, SY12$^{XY\Delta+Y}$, SY12,

and BY4742 cells exhibited similar sensitivities to various genotoxic agents, including

hydroxyurea (HU), camptothecin (CPT), and methyl methanesulfonate (MMS) (Figure 3G,

lower panel). These results indicate that the X- and Y'-elements are dispensable for cellular

responses to cold or heat treatment and DNA damage challenges, consistent with a recent

study of "synthetic yeast genome project", namely Sc2.0, showing that thousands of genome-

wide edits, including the deletion of subtelomeric repetitive sequences, deletion of introns,

and relocation of tRNAs genes, yielded a strain that displays comparable growth with wild

type strain (Richardson et al., 2017; Zhao et al., 2023).

Next, we examined the effects of X- and Y'-element elimination on telomeres. Southern

blotting assay revealed that SY12$^{Y\Delta}$, SY12$^{XY\Delta}$, and SY12$^{XY\Delta+Y}$ cells maintained stable telomeres

at a length of approximately 300 bp, comparable to that in SY12 cells (Figure 4A), indicating

that the X- and Y'-elements are not required for telomere length regulation. To determine

whether the deletion of X- and Y'-elements abolishes telomere silencing, we constructed

haploid strains of SY12$^{Y\Delta}$ *sir2*Δ, SY12$^{XY\Delta}$ *sir2*Δ, SY12$^{XY\Delta+Y}$ *sir2*Δ, SY12 *sir2*Δ, and BY4742 *sir2*Δ.

We then performed real-time RT-PCR to quantify the expression of the *MPH3* and *HSP32*

genes, located near the subtelomeric region of X-R (X-only end) and XVI-L (X-Y' end),

respectively (Figure 4B), and found that the increase of the *MPH3* or *HSP32* expression upon

SIR2 deletion in SY12$^{Y\Delta}$, SY12$^{XY\Delta}$ and SY12$^{XY\Delta+Y}$ strains was more significant than that in the

BY4742 or the SY12 strain, indicating that telomere silencing remains effective in the absence

of X-and Y'-elements (Figure 4B). These findings align with previous studies showing that

telomeres without an X- or Y'-element exert a position effect on the transcription of neighboring

genes (Aparicio et al., 1991), and that X- and Y'-elements function as modulators of TPE (Fourel

et al., 1999; Lebrun et al., 2001; Ottaviani et al., 2008).

In conclusion, the SY12$^{Y\Delta}$, SY12$^{XY\Delta}$, and SY12$^{XY\Delta+Y}$ strains behave similarly to the wild-type

SY12 strain under all tested conditions (Figures 3 and 4). Their simplified telomere structure

makes them potentially useful tools for telomere studies.

Figure 4

Figure 4. Telomere length and telomere silencing analyses of SY12$^{Y\Delta}$, SY12$^{XY\Delta}$ and SY12$^{XY\Delta+Y}$ strains.

(**A**) Southern blotting analysis of telomere length in SY12, SY12$^{Y\Delta}$, SY12$^{XY\Delta}$ and SY12$^{XY\Delta+Y}$ (labeled on

top) cells. Genomic DNA prepared from three independent clones of SY12, SY12$^{Y\Delta}$, SY12$^{XY\Delta}$ and

SY12$^{XY\Delta+Y}$ strains were digested with XhoI and PaeI, and then subjected to Southern blotting with a TG$_{1-}$

$_3$ probe. The numbers in brackets indicate the telomere length of the corresponding chromosomes. (**B**)

Expressions of *MPH3* and *HSP32* in BY4742, SY12, SY12$^{Y\Delta}$, SY12$^{XY\Delta}$ and SY12$^{XY\Delta+Y}$ cells were

detected by qRT-PCR. The numbers above the schematic line (lower panels) indicate the distance to

the corresponding subtelomeric elements or telomeres. The RNA levels of *MPH3* and *HSP32* were

normalized by *ACT1*. The wild-type value is arbitrarily set to 1. Error bars represent standard deviation

(s.d.), n = 3. 'ns', p > 0.5 (Student's t-test); *, $P < 0.05$ (Student's t-test); **, $P < 0.01$ (Student's t-test).;

****, $P < 0.0001$ (Student's t-test).

Y'-elements are not strictly required for the formation of Type II survivors

The BY4742 strain harbors nineteen Y'-elements distributed among seventeen telomere loci.

Numerous studies have emphasized the significance of Y'-elements in telomere recombination.

For instance, Type I survivors exhibit significant amplification of Y'-elements (Lundblad and

Blackburn, 1993; Teng and Zakian, 1999) and survivors show a marked induction of the

potential DNA helicase Y'-Help1 encoded by Y'-elements (Yamada et al., 1998). Additionally,

the acquisition of Y'-elements by short telomeres delays the onset of senescence (Churikov et

al., 2014).

To investigate the requirement of Y'-elements in survivor formation, we deleted *TLC1* in

SY12$^{Y\Delta}$ cells and conducted a cell viability assay. The results demonstrated that three individual

colonies underwent senescence and subsequently recovered at different passages in liquid

media (Figure 5A). Further analysis through Southern blotting revealed that the telomeres of

SY12$^{Y\Delta}$ *tlc1*Δ cells underwent progressive shortening with each passage until reaching critically

short lengths. Subsequently, TG$_{1-3}$ recombination occurred, leading to abrupt telomere

elongation (Figure 5B).

Next, we examined the telomere patterns of fifty independent SY12$^{Y\Delta}$ *tlc1*Δ survivors using a

multiple-colony streaking assay and Southern blotting analysis. Out of the fifty clones analyzed,

no Type I survivors were detected due to the deletion of Y'-elements in SY12$^{Y\Delta}$ strain (Figure

5C). Two clones (labeled in orange, 4% of the survivors tested) displayed heterogeneous

telomere tracts (Figure 5C and Figure 5D). Reintroduction of *TLC1* into a representative clone

(named SY12$^{Y\Delta}$ *tlc1*Δ-T1) resulted in telomere length restoration similar to SY12$^{Y\Delta}$ cells (Figure

5—figure supplement 1A), indicating their classification as Type II survivors. Twenty-six clones

(labeled in blue, 52% of the survivors tested) exhibited patterns identical to that of the SY12

*tlc1*Δ circular survivors (Figure 5C, Figure 5D and Figure 2D). Further mapping of erosion

borders and sequencing of fusion junctions (Figure 5E, Figure 5—figure supplement 2, and

supplementary file 3) confirmed that three chromosomes from a randomly selected clone

(named SY12$^{Y\Delta}$ *tlc1*Δ-C1) underwent intra-chromosomal fusions mediated by microhomology

sequences. The erosion sites and fusion sequences differed from those observed in SY12

*tlc1*Δ-C1 cells (Figure 2F), suggesting the stochastic nature of intra-chromosome end fusion by

MMEJ. As expected, the telomere Southern blotting pattern (XhoI digestion) of the SY12$^{Y\Delta}$

*tlc1*Δ-C1 survivor remained unchanged following telomerase reintroduction (Figure 5—figure

supplement 1B). Further PFGE analysis confirmed that the chromosomes in SY12^{YΔ} *tlc1*Δ-C1

were circulated (Figure 2—figure supplement 5). Notably, a significant proportion of the

survivors displayed telomere signals that were different from those of either the Type II or

circular survivors (labeled in black, 44% of the survivors tested, Figure 5C and Figure 5D), and

they were uncharacterized survivors. Further deletion of *RAD52* in the SY12^{YΔ} *tlc1*Δ cells

affected, but did not eliminate, survivor generation (Figure 5—figure supplement 3A). Southern

blotting assay confirmed that most of the recovered clones were circular survivors, and two

were uncharacterized survivors (clone 9 and 16, labeled in black, Figure 5—figure supplement

3B) suggesting that survivor formation in SY12^{YΔ} *tlc1*Δ rad52Δ cells does not strictly rely on the

homologous recombination. Overall, these findings indicate that Y'-elements are not strictly

required for Type II survivor formation (Churikov et al., 2014).

Figure 5

Figure 5. Survivor analysis of SY12$^{Y\Delta}$ *tlc1*Δ strain. (**A**) Cell viability assay in liquid medium. The growth

of SY12$^{Y\Delta}$ (labeled in black) and SY12$^{Y\Delta}$ *tlc1*Δ (three clones labeled in blue, purple and green

respectively) strains were monitored every 24 hr for 12 days. (**B**) Telomeric Southern blotting assay of

SY12$^{Y\Delta}$ *tlc1*Δ survivors. Genomic DNAs prepared from SY12$^{Y\Delta}$ *tlc1*Δ survivors assayed in (A) were

digested with XhoI and subjected to Southern blotting with a TG_{1-3} probe. (**C**) Telomere Southern

blotting analysis of SY12^{YΔ} *tlc1*Δ survivors obtained on solid medium. Genomic DNAs of fifty

independent survivors (labeled 1 to 50 on top) were digested with XhoI and hybridized by a TG_{1-3} probe.

Type II survivors: in orange; circular survivors: in blue; uncharacterized survivors: in black. Theoretical

telomere restriction fragments of the SY12^{YΔ} strain are indicated on left. LC: loading control. (**D**) The

ratio of survivor types in SY12^{YΔ} *tlc1*Δ strain. n=50; Type II, in orange; uncharacterized survivor, in grey;

circular survivor, in blue. (**E**) Schematic of three circular chromosomes and fusion sequences in the

SY12^{YΔ} *tlc1*Δ-C1 survivor. The sequence in blue indicates the sequences of X-R, XI-R or XIV-R, the

sequence in red indicates the sequences of I-L, XIII-L or XVI-L. Bases in green are mis-paired, dashes

are deleted. The numbers above or below the schematic line (chromosome) indicate the distance to the

corresponding telomeres.

Figure 5—figure supplement 1. Southern blotting results of reintroducing *TLC1* into SY12^{YΔ} *tlc1*Δ

survivors. (**A**) Southern blotting result of SY12^{YΔ} *tlc1*Δ Type II survivor at the 20th re-streaks after *TLC1*

reintroduction. LC: loading control. (**B**) Southern blotting result of SY12^{YΔ} *tlc1*Δ circular survivor at the

20th re-streaks after *TLC1* reintroduction. LC: loading control.

Figure 5—figure supplement 2. PCR mapping of the borders of erosion in SY12^{YΔ} *tlc1*Δ-C1 cell. Red

lines indicate the regions which have been deleted in the SY12^{YΔ} strain.

Figure 5—figure supplement 3. Survivor formation in SY12^{YΔ} *tlc1*Δ *rad52*Δ strain. (**A**) Cell viability

assay in liquid culture. The growth of SY12^{YΔ} (labeled in black), SY12^{YΔ} *tlc1*Δ (two clones labeled in

purple and blue respectively) and SY12^{YΔ} *tlc1*Δ *rad52*Δ (three clones labeled in violet, pink and carmine

respectively) strains were monitored every 24 hr for 12 days. (**B**) Telomere Southern blotting assay of

SY12^{YΔ} *tlc1*Δ *rad52*Δ survivors obtained on solid medium. Genomic DNA from twenty-five independent

clones (labeled on top) was digested with XhoI and hybridized to a telomere-specific TG_{1-3} probe.

Circular survivors: in blue; uncharacterized survivors: in black. Theoretical telomere restriction fragments

of the SY12^{YΔ} strain are indicated on the left. The asterisks indicate the non-specific bands. Genomic

DNA stained with Gelred was used as a relative loading control (LC).

X-elements are not strictly necessary for survivor generation

To investigate the contribution of X-elements to telomere recombination, we employed the SY12$^{XYΔ+Y}$ strain, which contains only one Y'-element in the subtelomeric region, and the SY12^{XYΔ} *tlc1*Δ strain, which lacks both the X- and Y'-elements. Subsequently, we deleted *TLC1* in the SY12$^{XYΔ+Y}$ and SY12^{XYΔ} strains and conducted a cell viability assay. Consistently, the deletion of *TLC1* in SY12$^{XYΔ+Y}$ and SY12^{XYΔ} resulted in telomere shortening, senescence, and the formation of Type II survivors (Figure 6—figure supplement 1). Then, fifty independent clones of SY12$^{XYΔ+Y}$ *tlc1*Δ or SY12^{XYΔ} *tlc1*Δ survivors were examined using Southern blotting (Figure 6A and 6B).

Among the SY12$^{XYΔ+Y}$ survivors analyzed, twenty-two clones underwent chromosomal circularization (labeled in blue, 44% of the survivors tested, Figure 6A and Figure 6C). We randomly selected a clone named SY12$^{XYΔ+Y}$ *tlc1*Δ-C1, and the results of erosion-border mapping and fusion junction sequencing showed that it had undergone intra-chromosomal fusions mediated by microhomology sequences (Figure 6D, Figure 6—figure supplement 2, and supplementary file 3). Subsequently, Southern blotting revealed that the chromosome structure of SY12$^{XYΔ+Y}$ *tlc1*Δ-C1 remained unchanged after *TLC1* reintroduction (Figure 6— figure supplement 3), and PFGE analysis confirmed the circular chromosome structure in SY12$^{XYΔ+Y}$ *tlc1*Δ-C1 (Figure 2—figure supplement 5). Additionally, seven clones utilized the Type II recombination pathway and exhibited heterogeneous telomeric TG$_{1-3}$ tracts (labeled in orange, 14% of the survivors tested, Figure 6A and Figure 6C). Reintroduction of *TLC1* into a

representative clone (named SY12$^{XYΔ+Y}$ *tlc1*Δ-T1) restored the telomere length to normal

(Figure 6—figure supplement 3). These findings indicate that the majority of cells underwent

intra-chromosomal circularization or TG$_{1-3}$ recombination. While even though there is a Y'-

element, no Type I survivors were generated in SY12$^{XYΔ+Y}$ *tlc1*Δ survivors (Figure 6A). We

speculated that the short TG$_{1-3}$ repeats located centromere-proximal to the Y'-elements play a

crucial role in strand invasion and subsequent Y'-recombination. This speculation is consistent

with a previous report stating that Type I events are virtually absent in the yeast strain Y55,

which lacks TG$_{1-3}$ repeats centromere-proximal to the Y'-element (Louis, 2001). We also

observed some clones displayed non-canonical telomere signals like SY12 *tlc1*Δ

"uncharacterized" survivors (labeled in black, 42% of the survivors tested, Figure 6A and Figure

6C). Overall, these data suggest that X-elements are not strictly necessary for survivor

formation.

Among the SY12^{XYΔ} survivors, twenty-four displayed a "circular survivor" pattern (labeled in

blue, 48% of the survivors tested, Figure 6B and Figure 6E). Additional PCR-sequencing

assays and PFGE analysis of the SY12^{XYΔ} *tlc1*Δ-C1 cells confirmed the occurrence of intra-

chromosomal fusions mediated by microhomology sequences (Figure 6F, Figure 6—figure

supplement 4, supplementary file 3 and Figure 2—figure supplement 4). Reintroduction of *TLC1*

into a representative clone named SY12^{XYΔ} *tlc1*Δ-C1 could restore its telomere length to WT

level (Figure 6—figure supplement 5A). Four of fifty survivors harbored Type II telomere

structure (labeled in orange, 8% of the survivors tested, Figure 6B and Figure 6E).

Reintroduction of *TLC1* into a representative clone named SY12^{XYΔ} *tlc1*Δ-T1 could restore its

telomere length to WT level (Figure 6—figure supplement 5B). Some of the survivors (labeled

in black, 44% of the survivors tested, Figure 6B and Figure 6E) were not characterized. Like in

SY12 *tlc1*Δ cells, Rad52 is not strictly required for the formation of circular survivors in SY12^{XYΔ}

*tlc1*Δ *rad52*Δ and SY12$^{XYΔ+Y}$ *tlc1*Δ *rad52*Δ strains (Figure 6—figure supplement 6A and B). To

investigate whether Type I-specific mechanisms are still utilized in the survivor formation in Y'-

less strain, we deleted *RAD51* in SY12^{XYΔ} *tlc1*Δ, and found that SY12^{XYΔ} *tlc1*Δ *rad51*Δ strain

was able to generate three types of survivors, including Type II survivor, circular survivor and

uncharacterized survivor (Figure 6—figure supplement 7A), similar to the observations in

SY12^{XYΔ} *tlc1*Δ strain (Figure 6B). Notably, the proportions of circular and uncharacterized

survivors in the SY12^{XYΔ} *tlc1*Δ *rad51*Δ strain were 36% (9/25) and 32% (8/25) (Figure 6—figure

supplement 7B and supplementary file 5), respectively, lower than 48% and 44% in the SY12^{XYΔ}

*tlc1*Δ strain (Figure 6E and supplementary file 5). Accordingly, the ratio of Type II survivor in

SY12^{XYΔ} *tlc1*Δ *rad51*Δ was (32% of the survivors tested, Figure 6—figure supplement 7B and

supplementary file 5) was higher than SY12^{XYΔ} *tlc1*Δ strain (8% of the survivors tested, Figure

6E and supplementary file 5), suggesting that Type I-specific mechanisms still contribute to the

survivor formation even in the Y'-less strain SY12^{XYΔ}. Collectively, the aforementioned data

suggest that X-elements, as well as Y'-elements, are not essential for the generation of Type II

survivors.

Figure 6

Figure 6. Survivor analysis of SY12$^{XY\Delta}$ *tlc1*Δ and SY12$^{XY\Delta+Y}$ *tlc1*Δ strains. (**A**) and (**B**) Telomere

Southern blotting analysis of SY12$^{XY\Delta+Y}$ *tlc1*Δ (A) and SY12$^{XY\Delta}$ *tlc1*Δ (B) survivors obtained on solid

medium. fifty independent survivors (labeled 1 to 50 on top) were randomly picked, and their genomic

DNAs were digested with XhoI and subjected to the Southern blotting assay with a TG$_{1-3}$ probe. Type II

survivors: in orange; circular survivors: in blue; uncharacterized survivors: in black. The sizes of

individual telomere restriction fragments of the SY12$^{XY\Delta+Y}$ and SY12$^{XY\Delta}$ strain are indicated on the left.

LC: loading control. (**C**) and (**E**) The percentage of survivor types in SY12$^{XY\Delta+Y}$ *tlc1*Δ (C) and SY12$^{XY\Delta}$

*tlc1*Δ (E) strains. n=50; Type II, in orange; uncharacterized survivor, in grey; circular survivor, in blue.

(**D**) and (**F**) Schematic of three circular chromosomes and fusion sequences in the SY12$^{XY\Delta+Y}$ *tlc1*Δ-C1

(D) and SY12$^{XY\Delta}$ *tlc1*Δ-C1 (F) survivors, respectively. The sequence in blue indicates the sequences of

X-R, XI-R or XIV-R, the sequence in red indicates the sequences of I-L, XIII-L or XVI-L. Bases in green

are mis-paired, dashes are deleted. The numbers above or below the schematic line (chromosome)

indicate the distance to the corresponding telomeres.

Figure 6—figure supplement 1. SY12$^{XY\Delta+Y}$ *tlc1*Δ and SY12$^{XY\Delta}$ *tlc1*Δ strains form Type II survivors in

liquid culture. (**A**) Cell viability assay in liquid medium: The growth of SY12$^{XY\Delta+Y}$ (labeled in black) and

SY12$^{XY\Delta+Y}$ *tlc1*Δ (three clones labeled in blue, purple and green respectively) strains were monitored

every 24 hr for 12 days. (**B**) Telomeric Southern blotting assay of SY12$^{XY\Delta+Y}$ *tlc1*Δ survivors. Genomic

DNAs prepared from survivors in (A) were digested with XhoI and subjected to Southern blotting with a

TG$_{1-3}$ probe. (**C**) Cell viability assay in liquid medium: The growth of SY12$^{XY\Delta}$ (labeled in black) and

SY12$^{XY\Delta}$ *tlc1*Δ (three clones labeled in blue, purple and green respectively) strains were monitored every

24 hr for 12 days. (**D**) Telomeric Southern blotting assay of SY12$^{XY\Delta}$ *tlc1*Δ survivors. Genomic DNAs

prepared from survivors in (C) were digested with XhoI and subjected to Southern blotting with a TG$_{1-3}$

probe.

Figure 6—figure supplement 2. PCR mapping of the borders of erosion in SY12$^{XY\Delta+Y}$ *tlc1*Δ-C1 cell.

Red lines indicate the regions which are absent in the SY12$^{XY\Delta+Y}$ strain.

Figure 6—figure supplement 3. Southern blotting results of an SY12$^{XY\Delta+Y}$ *tlc1*Δ circular survivor and

an SY12$^{XY\Delta+Y}$ *tlc1*Δ Type II survivor at the 20th re-streaks after *TLC1* reintroduction. LC: loading control.

Figure 6—figure supplement 4. PCR mapping of the borders of erosion in SY12$^{XY\Delta}$ *tlc1*Δ-C1 cell. Red

lines indicate the regions which have been deleted in the SY12$^{XY\Delta}$ strain.

Figure 6—figure supplement 5. Southern blotting results of reintroducing *TLC1* into SY12$^{XY\Delta}$ *tlc1*Δ

survivors. (**A**) Southern blotting result of SY12$^{XY\Delta}$ *tlc1*Δ circular survivor at the 20th re-streaks after *TLC1*

reintroduction. LC: loading control. (**B**) Southern blotting result of SY12$^{XY\Delta}$ *tlc1*Δ Type II survivor at the

20th re-streaks after *TLC1* reintroduction. LC: loading control.

Figure 6—figure supplement 6. Survivor formation in SY12$^{XY\Delta+Y}$ *tlc1*Δ *rad52*Δ and SY12$^{XY\Delta}$ *tlc1*Δ

*rad52*Δ strains. (**A**) Cell viability assay in liquid culture. The growth of SY12$^{XY\Delta+Y}$ (labeled in black),

SY12$^{XY\Delta+Y}$ *tlc1*Δ (two clones labeled in purple and blue respectively) and SY12$^{XY\Delta+Y}$ *tlc1*Δ *rad52*Δ (three

clones labeled in violet, pink and carmine respectively) strains shown in the left panel; the survivor

formation in SY12$^{XY\Delta}$ (labeled in black), SY12$^{XY\Delta}$ *tlc1*Δ (two clones labeled in purple and blue

respectively) and SY12$^{XY\Delta}$ *tlc1*Δ *rad52*Δ (three clones labeled in violet, pink and carmine respectively)

strains shown in the right panel. They were monitored every 24 hr for 12 days. (**B**) Telomere Southern

blotting assay of SY12$^{XY\Delta+Y}$ *tlc1*Δ *rad52*Δ (upper) and SY12$^{XY\Delta}$ *tlc1*Δ *rad52*Δ (lower) survivors obtained

on solid medium. Genomic DNA from twenty-five independent clones (labeled on top) was digested with

XhoI and hybridized to a telomere-specific TG$_{1-3}$ probe. Circular survivors: in blue; uncharacterized

survivors: in black. Theoretical telomere restriction fragments of each strain are indicated on the left.

The asterisks indicate the non-specific bands. Genomic DNA stained with Gelred was used as a relative

loading control (LC).

Figure 6—figure supplement 7. Survivor formation in SY12$^{XY\Delta}$ *tlc1*Δ *rad51*Δ strain. (**A**) Southern

blotting result of SY12$^{XY\Delta}$ *tlc1*Δ *rad51*Δ strain. Twenty-five independent survivors (labeled 1 to 25 on top)

were randomly picked, and their genomic DNAs were digested with XhoI and subjected to the Southern

blotting assay with a TG$_{1-3}$ probe. Type II survivors: in orange; circular survivors: in blue;

uncharacterized survivors: in black. (**B**) The percentage of survivor types in SY12$^{XY\Delta}$ *tlc1Δ rad51Δ*

strain. n=25; Type II, in orange; uncharacterized survivor, in grey; circular survivor, in blue.

____

Discussion

The wild-type yeast strain BY4742, commonly used in laboratories, possesses nineteen Y'-

elements at seventeen telomere loci and thirty-two X-elements at thirty-two telomere loci. This

abundance of Y'-elements and X-elements poses challenges for loss-of-function studies,

highlighting the need for a strain lacking all Y'-elements and X-elements. Fortunately, we have

previously constructed the single-chromosome yeast strain SY14, which contains only one copy

of Y'-element and two copies of X-element (Shao et al., 2018), and could have been an ideal

tool. However, the telomerase-null survivors of SY14 mainly bypassed senescence through

chromosomal circularization, providing limited insights into the roles of Y'- and X-elements in

telomere maintenance (Wu et al., 2020). Therefore, in this study, we employed the SY12 strain,

which has three chromosomes, to investigate the functions of Y'- and X-elements at telomeres

(Figure 2A, left panel).

We constructed the SY12$^{Y\Delta}$, SY12$^{XY\Delta+Y}$, and SY12$^{XY\Delta}$ strains, which lack the Y'-element, X-

elements, and both X- and Y'-elements, respectively (Figure 3A). Surprisingly, the SY12$^{Y\Delta}$,

SY12$^{XY\Delta}$, and SY12$^{XY\Delta+Y}$ strains exhibited minimal defects in cell proliferation, genotoxic

sensitivity, and telomere homeostasis (Figures 3 and 4). These results demonstrate, for the first

time, that both X- and Y'-elements are dispensable for cellular functions. Thus, the SY12$^{Y\Delta}$,

SY12$^{XY\Delta}$, and SY12$^{XY\Delta+Y}$ strains established in this study, with their simplified telomere

structures, are valuable resources for telomere biology research.

Subtelomeric regions are known to be highly variable and often contain species-specific

homologous DNA sequences. In the case of fission yeast, subtelomeric regions consist of

subtelomeric homologous (SH) and telomere-distal sequences. Previous studies have shown

that subtelomeric homologous sequences in fission yeast do not significantly impact telomere

length, mitotic cell growth, or stress responses. However, they do play a role in buffering against

the spreading of silencing signals from the telomere (Tashiro et al., 2017). Though the "core X"

sequence acts as a protosilencer (Lebrun et al., 2001), the X-STRs and Y'-STAR possess anti-

silencing properties that limit the spreading of heterochromatin in budding yeast (Fourel et al.,

1999), the telomere position effect remains effective in the strains that lack both X- and Y'-

elements (Figure 4B). Given the remarkable differences in both sequence and size between

the subtelomeric regions of budding yeast and fission yeast, it is difficult to compare the extent

to which subtelomeric elements affect telomere silencing.

Amplification of Y'-element(s) is a characteristic feature of canonical Type I survivors. Type I

survivor emerged in SY12 strain indicating that multiple Y'-elements in tandem is not strictly

required for type I recombination (Figure 2D). Interestingly, the telomerase-null SY12$^{Y\Delta}$ and

SY12$^{XY\Delta}$ cells, lacking Y'-elements, failed to generate Type I survivors but could generate Type

II survivors, indicating that the acquisition of Y'-elements is not a prerequisite for Type II survivor

formation (Figure 5C and Figure 6B). These observations support the notion that Type I and

Type II survivors form independently, although both may utilize a common alternative telomere-

lengthening pathway (Kockler et al., 2021). Moreover, a subset of SY12 *tlc1*Δ, SY12^{YΔ} *tlc1*Δ,

SY12$^{XYΔ+Y}$ *tlc1*Δ, and SY12^{XYΔ} *tlc1*Δ cells could escape senescence and become survivors

through microhomology-mediated intra-chromosomal end-to-end fusion (chromosome

circularization) (Figure 2D, Figure 5C, Figure 6A and 6B, labeled in blue). Notably, the survivors

with all circular chromosomes were readily recovered from the telomerase-null SY11 to SY14,

but not SY1 to SY10 cells (Figure 1). Several reasons could account for this. First, a smaller

number of telomeres provides fewer recombination donors and acceptors, resulting in less

efficient inter-chromosomal homologous recombination (e.g., TG$_{1-3}$ tracts recombination or Y'-

element acquisition). Second, the continuously shortened telomeres of linear chromosomes

may trigger another round of senescence, while survivors with circular chromosomes do not

encounter end-replication problems and therefore exhibit greater stability. Third, the presence

of homologous sequences at both chromosome ends appears to be a minimum requirement

for microhomology-mediated intra-chromosomal end-to-end fusion. With fewer homologous

sequences, the probability of chromosome circularization decreases, and with more

chromosomes, the likelihood of circularizing each chromosome within a cell diminishes. Fourth,

in cells with fewer telomeres, intra-chromosomal telomere fusions are more likely to occur, while

lethal inter-chromosomal fusions are competed out. However, we can speculate that in

telomerase-null cells with eroded chromosome ends, stochastic repair mechanisms such as

homologous recombination, microhomology-mediated end joining, and inter- and intra-

chromosomal fusions operate simultaneously. Only those survivors that maintain a relatively

stable genome and robust growth can be experimentally recovered.

Saccharomyces cerevisiae (budding yeast) and *Schizosaccharomyces pombe* (fission yeast)

are the most commonly used laboratory systems, separated by approximately 1 Gya (billion

years ago) according to molecular-clock analyses (Hedges, 2002). Despite both species having

genomes are both over 12 megabases in length, haploid *S. cerevisiae* contains 16

chromosomes, while *S. pombe* has only 3 chromosomes (Forsburg, 2005). The telomerase-

independent mechanisms for maintaining chromosome ends differ between these two yeasts.

In budding yeast, homologous recombination is the primary mode of survival in telomerase-

deficient cells, resulting in the generation of Type I or Type II survivors (McEachern and Haber,

2006). Telomerase- and recombination-deficient cells occasionally escape senescence through

the formation of palindromes at chromosome ends in the absence of *EXO1* (Maringele and

Lydall, 2004). Fission yeast cells lacking telomerase can also maintain their chromosome

termini by recombining persistent telomere sequences, and survivors with all intra-circular

chromosomes (Nakamura et al., 1998) or intermolecular fusions (Tashiro et al., 2017; Wang

and Baumann, 2008) have been observed. In our research, some SY12 *tlc1Δ* cells, which have

three chromosomes, also bypassed senescence by circularizing their chromosomes (Figure

2D), suggesting that a lower chromosome number increases the likelihood of recovering

survivors containing circular chromosomes.

While most eukaryotes employ telomerase for telomere replication, some eukaryotes lack

telomerase and utilize recombination as an alternative means to maintain telomeres

(Biessmann and Mason, 1997). In *Drosophila*, telomeres are replicated through a

retrotransposon mechanism (Levis et al., 1993; Louis, 2002). The structure and distribution of

Y'-elements in *S. cerevisiae* suggest their origin from a mobile element (Jager and Philippsen,

1989; Louis and Haber, 1992), and Y'-elements can be mobilized through a transposition-like

RNA-mediated process (Maxwell et al., 2004). In telomerase-deficient yeast cells, homologous

recombination can acts as a backup mechanism for telomere replication (Lundblad and

Blackburn, 1993), and the reintroduction of telomerase efficiently inhibits telomere

recombination and dominates telomere replication (Chen et al., 2009; Peng et al., 2015; Teng

and Zakian, 1999), These findings suggest that subtelomeric region amplification mediated by

recombination and/or transposition may represent ancient telomere maintenance mechanisms

predating the evolution of telomerase (de Lange, 2004). Therefore, subtelomeric X- and Y'-

elements might be considered as evolutionary "fossils" in the *S. cerevisiae* genome, and their

elimination has little impact on telomere essential functions and genome stability.

MATERIALS AND METHODS

Yeast strains and plasmids

Yeast strains used in this study are listed in supplementary file 6. The plasmids for gene deletion

and endogenous expression of *TLC1* were constructed based on the pRS series as described

previously (Sikorski and Hieter, 1989). We use PCR to amplify the upstream and downstream

sequence adjacent to the target gene, and then the PCR fragments were digested with different

restriction enzymes and inserted into pRS plasmids. Plasmids were introduced into budding

yeast by standard procedures, and transformants were selected on auxotrophic medium (Orr-

Weaver et al., 1981).

Multiple-colony streaking assay

Single clones of indicated yeast strains were randomly picked and streaked on extract-peptone-

dextrose (YPD) plates. Thereafter, several clones of their descendants were passaged by

successive re-streaks at 30°C. This procedure was repeated dozens of times every two days.

Telomere Southern blotting

Southern blotting was performed as previously described (Hu et al., 2013). Yeast genomic DNA

was extracted by a phenol chloroform method. Restriction fragments were separated by

electrophoresis in 1% agarose gel, transferred to Amersham Hybond-N$^+$ membrane (GE

Healthcare) and hybridized with α-^{32}P dCTP labeled probe.

Cell viability assay

Cell viability assay was performed as previously described with few modifications (Le et al.,

1999). Three independent single colonies of indicated strains were grown to saturation at 30°C.

Then the cell density was measured every 24 hours by spectrometry (OD_{600}), and the cultures

were diluted to the density at $OD_{600} = 0.01$. This procedure was repeated several times to allow

the appearance of survivors. The genomic DNA samples at indicated time points were

harvested for telomere length analysis.

Molecular analysis of circular chromosomes

Fusion events were determined by PCR amplification and DNA sequencing. Genomic DNA was

extracted by phenol chloroform. Frist, we use primers pairs located at different sites of each

chromosome arm at an interval of 1 kb (listed in Supplementary file 1) to determine the erosion

site of each chromosome; PCR was performed as standard procedures in 10 µl reactions by

TaKaRa Ex Taq. To amplify the sequence of fusion junction we use pairs of primers oriented

to different arm of each chromosome; PCR was performed as standard procedures in 50 µl

reactions by TaKaRa LA Taq. The fragments were purified by kit (QINGEN), then they were

sequenced directly or cloned into the pMD18-T Vector (TaKaRa) for sequencing.

CRISPR-Cas9 mediated X- and Y'-elements deletion

X- and Y'-elements were deleted as described (Shao et al., 2018; Shao et al., 2019). Briefly,

pgRNA and a DNA targeting cassette, containing a selection marker, a homology arm (DR1),

a direct repeat (DR2), and telomeric repeats, were co-introduced into indicated cells harboring

pCas9. pCas9 nuclease was directed to a specific DNA sequence centromere-proximal to the

subtelomeric region with the guidance of gRNA1, where it induces a double-stranded break.

Homologous recombination between the broken chromosome and the provided DNA targeting

cassette caused the deletion of X- and Y'-elements. The positive transformants identified by

PCR were transferred into the galactose-containing liquid medium, which induces the

expression of the gRNA2 on pCas9 to cut at the target site near the *URA3* gene and on the

backbone of pgRNA. Then the culture was plated on the medium containing 5'-FOA to select

for eviction of the *URA3* marker.

Cell growth assay

Three individual colonies of the indicated strains were inoculated into 5 ml liquid medium and

incubated at 30°C. The cell cultures were then diluted in 30 ml of fresh YPD medium to the

density at OD_{600} = 0.1. Then the density of cells was measured by spectrometry (OD_{600}) hourly.

Fluorescence-activated cell sorting (FACS) assay

The FACS analysis was performed as previously described (He et al., 2019). Yeast cells were

cultured at 30°C until the log phase, and then 1 ml of the cells was harvested. The cells were

washed with cold sterile ddH_2O and fixed with 70% ethanol overnight at 4°C. The following day,

the cells were washed with 50 mM sodium citrate buffer (pH 7.2) and then digested with 0.25

mg/ml RNase A at 37°C for 2-3 hours, followed by 0.2 mg/ml Protease K at 50°C for 1 hour.

Both RNase A and Protease K were diluted in sodium citrate buffer. The cells were

resuspended in 500 µl sodium citrate buffer and then sonicated for 45 seconds at 100% power.

The DNA of the cells was stained with 20 µg/ml propidium iodide (PI) at 4°C overnight or at

room temperature for 1 hour. FACS analysis was performed on a BD LSRII instrument.

Serial dilution assay

A single colony per strain was inoculated into 3 ml liquid medium and incubated at 30°C. The

cell cultures were then adjusted to a concentration of OD_{600} ~0.5. Five-fold serially diluted cells

were spotted on the indicated plates. The plates were incubated at 30°C for the appropriate

time prior to photography.

RNA extraction and RT-qPCR

Three independent single colonies of indicated strains were grown to log phase at 30°C. Yeast

pellets from a 1 ml cell culture were digested with Zymolyase 20 T (MP Biomedicals, LLC) to

obtain spheroplasts. RNA was extracted with RNeasy mini kit (Qiagen) followed by reverse

transcription using the Fastquant RT kit (Tiangen). Real-time PCR was carried out using

SYBR Premix Ex Taq™ II (Takara) on the Applied Biosystems StepOne Real-Time PCR

System. Primer pairs used in RT-qPCR were listed in supplementary file 1. The gene

expression levels were normalized to that of *ACT1* and the wild-type value is arbitrarily set to

1.

Pulsed-field gel electrophoresis (PFGE) analysis

DNA plugs for pulse-field gel electrophoresis (PFGE) were prepared according to the

manufacturer's instructions (Bio-Rad) and Ishii et al (Ishii et al., 2008). Fresh yeast cells were

inoculated in 50 ml YPD and incubated at 30°C until the OD600 reached approximately 1.0.

The cells were subsequently harvested, washed twice with cold EDTA buffer (50 mM, pH 8.0),

and resuspended in 300 µl of CSB buffer (10 mM pH 7.2 Tris-Cl, 20 mM NaCl, 100 mM pH 8.0

EDTA, 4 mg/ml lyticase) and blended with 300 µl of 2% low-melt agarose (Bio-Rad). Then, 100

µl of resuspended cells were added to each plug and incubated at 4°C for 30 min until the

agarose plugs were solidified. The solidified agarose plugs were incubated in lyticase buffer

(10 mM pH 7.2 Tris-Cl, 100 mM pH 8.0 EDTA, 1 mg/ml lyticase) at 37°C for 3 hours, followed

by incubation in Proteinase K Reaction Buffer (100 mM pH 8.0 EDTA, 0.2% sodium

deoxycholate, 1% sodium lauryl sarcosine) containing 1 mg/ml Proteinase K at 50°C for 12

hours. The plugs were washed four times in 25 ml of wash buffer (20 mM Tris, pH 8.0, 50 mM

EDTA) for 1 hour each time at room temperature with gentle agitation. The plugs were then

fixed into a pulsed field agarose gel (Bio-Rad), and the CHEF-DR II Pulsed Field

Electrophoresis System (Bio-Rad) was used for gel electrophoresis. The electrophoresis

conditions for separation were as follows: 0.8% agarose gel, 1×TBE buffer, 14°C temperature,

first run: initial switch time 1200 s; final switch time 1200 s; run time 24 hours; voltage gradient

2 V/cm; angle 96°; second run: initial switch time 1500 s; final switch time 1500 s; run time 24

hours; voltage gradient 2 V/cm; angle 100°; third run: initial switch time 1800 s; final switch time

1800 s; run time 24 hours; voltage gradient 2 V/cm; angle 106°. The gel was stained with

GelstainRedTM nucleic acid dye (US EVERBRIGHT), and PFGE Gels were imaged by Tanon

2500.

ACKNOWLEDGMENTS

We thank members of Zhou lab for discussions and suggestions for this project.

AUTHOR CONTRIBUTIONS

Conceptualization: Zhi-Jing Wu, Jin-Qiu Zhou.

Funding acquisition: Jin-Qiu Zhou.

Investigation: Can Hu, Zhi-Jing Wu, Xue-Ting Zhu.

Methodology: Zhi-Jing Wu, Yangyang Shao, Zhongjun Qin.

Supervision: Jin-Qiu Zhou.

Writing-original draft: Zhi-Jing Wu.

Writing-review & editing: Zhi-Jing Wu, Can Hu, Xue-Ting Zhu, Ming-Hong He, Jin-Qiu Zhou.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

Aparicio, O.M., Billington, B.L., and Gottschling, D.E. (1991). Modifiers of position effect are shared
between telomeric and silent mating-type loci in S. cerevisiae. Cell *66*, 1279-1287.

Biessmann, H., and Mason, J.M. (1997). Telomere maintenance without telomerase. Chromosoma *106*,
63-69.

Chan, C.S.M., and Tye, B.K. (1983a). A Family of Saccharomyces-Cerevisiae Repetitive Autonomously
Replicating Sequences That Have Very Similar Genomic Environments. Journal of Molecular Biology
168, 505-523.

Chan, C.S.M., and Tye, B.K. (1983b). Organization of DNA-Sequences and Replication Origins at Yeast
Telomeres. Cell *33*, 563-573.

Chen, Q., Ijpma, A., and Greider, C.W. (2001). Two survivor pathways that allow growth in the absence
of telomerase are generated by distinct telomere recombination events. Mol Cell Biol *21*, 1819-1827.

Chen, X.F., Meng, F.L., and Zhou, J.Q. (2009). Telomere recombination accelerates cellular aging in
Saccharomyces cerevisiae. PLoS Genet *5*, e1000535.

Churikov, D., Charifi, F., Simon, M.N., and Geli, V. (2014). Rad59-facilitated acquisition of Y' elements
by short telomeres delays the onset of senescence. PLoS Genet *10*, e1004736.

Craven, R.J., and Petes, T.D. (1999). Dependence of the regulation of telomere length on the type of
subtelomeric repeat in the yeast Saccharomyces cerevisiae. Genetics *152*, 1531-1541.

de Lange, T. (2004). T-loops and the origin of telomeres. Nat Rev Mol Cell Biol *5*, 323-329.

Forsburg, S.L. (2005). The yeasts Saccharomyces cerevisiae and Schizosaccharomyces pombe:
models for cell biology research. Gravit Space Biol Bull *18*, 3-9.

Fourel, G., Revardel, E., Koering, C.E., and Gilson, E. (1999). Cohabitation of insulators and silencing
elements in yeast subtelomeric regions. Embo Journal *18*, 2522-2537.

He, M.H., Liu, J.C., Lu, Y.S., Wu, Z.J., Liu, Y.Y., Wu, Z., Peng, J., and Zhou, J.Q. (2019). KEOPS
complex promotes homologous recombination via DNA resection. Nucleic Acids Res *47*, 5684-5697.

Hedges, S.B. (2002). The origin and evolution of model organisms. Nat Rev Genet *3*, 838-849.

Hediger, F., Berthiau, A.S., van Houwe, G., Gilson, E., and Gasser, S.M. (2006). Subtelomeric factors
antagonize telomere anchoring and Tel1-independent telomere length regulation. EMBO J *25*, 857-867.

Hu, Y., Tang, H.-B., Liu, N.-N., Tong, X.-J., Dang, W., Duan, Y.-M., Fu, X.-H., Zhang, Y., Peng, J.,
Meng, F.-L.*, et al.* (2013). Telomerase-null survivor screening identifies novel telomere recombination
regulators. PLoS genetics *9*, e1003208.

Ishii, K., Ogiyama, Y., Chikashige, Y., Soejima, S., Masuda, F., Kakuma, T., Hiraoka, Y., and Takahashi,
K. (2008). Heterochromatin integrity affects chromosome reorganization after centromere dysfunction.
Science *321*, 1088-1091.

Jager, D., and Philippsen, P. (1989). Many Yeast Chromosomes Lack the Telomere-Specific Y'
Sequence. Molecular and Cellular Biology *9*, 5754-5757.

Johnson, F.B., Marciniak, R.A., McVey, M., Stewart, S.A., Hahn, W.C., and Guarente, L. (2001). The
Saccharomyces cerevisiae WRN homolog Sgs1p participates in telomere maintenance in cells lacking
telomerase. EMBO J *20*, 905-913.

Kockler, Z.W., Comeron, J.M., and Malkova, A. (2021). A unified alternative telomere-lengthening
pathway in yeast survivor cells. Molecular cell *81*, 1816-1829 e1815.

Larrivee, M., and Wellinger, R.J. (2006). Telomerase- and capping-independent yeast survivors with

alternate telomere states. Nat Cell Biol *8*, 741-747.

Le, S., Moore, J.K., Haber, J.E., and Greider, C.W. (1999). RAD50 and RAD51 define two pathways that

collaborate to maintain telomeres in the absence of telomerase. Genetics *152*, 143-152.

Lebrun, E., Revardel, E., Boscheron, C., Li, R., Gilson, E., and Fourel, G. (2001). Protosilencers in

Saccharomyces cerevisiae subtelomeric regions. Genetics *158*, 167-176.

Lemieux, B., Laterreur, N., Perederina, A., Noel, J.F., Dubois, M.L., Krasilnikov, A.S., and Wellinger,

R.J. (2016). Active Yeast Telomerase Shares Subunits with Ribonucleoproteins RNase P and RNase

MRP. Cell *165*, 1171-1181.

Lendvay, T.S., Morris, D.K., Sah, J., Balasubramanian, B., and Lundblad, V. (1996). Senescence

mutants of Saccharomyces cerevisiae with a defect in telomere replication identify three additional EST

genes. Genetics *144*, 1399-1412.

Levis, R.W., Ganesan, R., Houtchens, K., Tolar, L.A., and Sheen, F.M. (1993). Transposons in place of

telomeric repeats at a Drosophila telomere. Cell *75*, 1083-1093.

Lewis, L.K., Storici, F., Van Komen, S., Calero, S., Sung, P., and Resnick, M.A. (2004). Role of the

nuclease activity of Saccharomyces cerevisiae Mre11 in repair of DNA double-strand breaks in mitotic

cells. Genetics *166*, 1701-1713.

Louis (2001). SGS1 is required for telomere elongation in the absence of telomerase.

Louis, E.J. (1995). The chromosome ends of Saccharomyces cerevisiae. Yeast *11*, 1553-1573.

Louis, E.J. (2002). Are Drosophila telomeres an exception or the rule? Genome biology *3*,

REVIEWS0007.

Louis, E.J., and Haber, J.E. (1991). Evolutionarily recent transfer of a group I mitochondrial intron to

telomere regions in Saccharomyces cerevisiae. Curr Genet *20*, 411-415.

Louis, E.J., and Haber, J.E. (1992). The structure and evolution of subtelomeric Y' repeats in

Saccharomyces cerevisiae. Genetics *131*, 559-574.

Louis, E.J., Naumova, E.S., Lee, A., Naumov, G., and Haber, J.E. (1994). The chromosome end in

yeast: its mosaic nature and influence on recombinational dynamics. Genetics *136*, 789-802.

Lundblad, V., and Blackburn, E.H. (1993). An alternative pathway for yeast telomere maintenance

rescues est1- senescence. Cell *73*, 347-360.

Lundblad, V., and Szostak, J.W. (1989). A mutant with a defect in telomere elongation leads to

senescence in yeast. Cell *57*, 633-643.

Maringele, L., and Lydall, D. (2004). Telomerase- and recombination-independent immortalization of

budding yeast. Genes Dev *18*, 2663-2675.

Maxwell, P.H., Coombes, C., Kenny, A.E., Lawler, J.F., Boeke, J.D., and Curcio, M.J. (2004). Ty1

mobilizes subtelomeric Y' elements in telomerase-negative Saccharomyces cerevisiae survivors. Mol

Cell Biol *24*, 9887-9898.

McEachern, M.J., and Haber, J.E. (2006). Break-induced replication and recombinational telomere

elongation in yeast. Annu Rev Biochem *75*, 111-135.

Nakamura, T.M., Cooper, J.P., and Cech, T.R. (1998). Two modes of survival of fission yeast without

telomerase. Science *282*, 493-496.

Nicolette, M.L., Lee, K., Guo, Z., Rani, M., Chow, J.M., Lee, S.E., and Paull, T.T. (2010). Mre11-Rad50-
Xrs2 and Sae2 promote 5' strand resection of DNA double-strand breaks. Nat Struct Mol Biol *17*, 1478-
1485.

O'Donnell, S., Yue, J.-X., Saada, O.A., Agier, N., Caradec, C., Cokelaer, T., De Chiara, M., Delmas, S.,
Dutreux, F., Fournier, T.*, et al.* (2023). Telomere-to-telomere assemblies of 142 strains characterize the
genome structural landscape in Saccharomyces cerevisiae. Nat Genet *55*, 1390-1399.

Orr-Weaver, T.L., Szostak, J.W., and Rothstein, R.J. (1981). Yeast transformation: a model system for
the study of recombination. Proc Natl Acad Sci U S A *78*, 6354-6358.

Ottaviani, A., Gilson, E., and Magdinier, F. (2008). Telomeric position effect: from the yeast paradigm to
human pathologies? Biochimie *90*, 93-107.

Palm, W., and de Lange, T. (2008). How shelterin protects mammalian telomeres. Annu Rev Genet *42*,
301-334.

Peng, J., He, M.H., Duan, Y.M., Liu, Y.T., and Zhou, J.Q. (2015). Inhibition of telomere recombination by
inactivation of KEOPS subunit Cgi121 promotes cell longevity. PLoS Genet *11*, e1005071.

Richardson, S.M., Mitchell, L.A., Stracquadanio, G., Yang, K., Dymond, J.S., DiCarlo, J.E., Lee, D.,
Huang, C.L.V., Chandrasegaran, S., Cai, Y.*, et al.* (2017). Design of a synthetic yeast genome. Science
(New York, NY) *355*, 1040-1044.

Shao, Y., Lu, N., Wu, Z., Cai, C., Wang, S., Zhang, L.L., Zhou, F., Xiao, S., Liu, L., Zeng, X.*, et al.*
(2018). Creating a functional single-chromosome yeast. Nature *560*, 331-335.

Shao, Y.Y., Lu, N., Xue, X.L., and Qin, Z.J. (2019). Creating functional chromosome fusions in yeast
with CRISPR-Cas9. Nat Protoc *14*, 2521-2545.

Sikorski, R.S., and Hieter, P. (1989). A system of shuttle vectors and yeast host strains designed for
efficient manipulation of DNA in Saccharomyces cerevisiae. Genetics *122*, 19-27.

Singer, M.S., and Gottschling, D.E. (1994). TLC1: template RNA component of Saccharomyces
cerevisiae telomerase. Science *266*, 404-409.

Tashiro, S., Nishihara, Y., Kugou, K., Ohta, K., and Kanoh, J. (2017). Subtelomeres constitute a
safeguard for gene expression and chromosome homeostasis. Nucleic Acids Res *45*, 10333-10349.

Teng, S.C., Chang, J., McCowan, B., and Zakian, V.A. (2000). Telomerase-independent lengthening of
yeast telomeres occurs by an abrupt Rad50p-dependent, Rif-inhibited recombinational process.
Molecular cell *6*, 947-952.

Teng, S.C., and Zakian, V.A. (1999). Telomere-telomere recombination is an efficient bypass pathway
for telomere maintenance in Saccharomyces cerevisiae. Molecular and Cellular Biology *19*, 8083-8093.

Wang, X., and Baumann, P. (2008). Chromosome fusions following telomere loss are mediated by
single-strand annealing. Molecular cell *31*, 463-473.

Wellinger, R.J., and Zakian, V.A. (2012). Everything you ever wanted to know about Saccharomyces
cerevisiae telomeres: beginning to end. Genetics *191*, 1073-1105.

Wu, Z., Liu, J., Zhang, Q.D., Lv, D.K., Wu, N.F., and Zhou, J.Q. (2017). Rad6-Bre1-mediated H2B
ubiquitination regulates telomere replication by promoting telomere-end resection. Nucleic Acids Res *45*,
3308-3322.

Wu, Z.J., Liu, J.C., Man, X., Gu, X., Li, T.Y., Cai, C., He, M.H., Shao, Y., Lu, N., Xue, X.*, et al.* (2020).
Cdc13 is predominant over Stn1 and Ten1 in preventing chromosome end fusions. Elife *9*.

Yamada, M., Hayatsu, N., Matsuura, A., and Ishikawa, F. (1998). Y'-Help1, a DNA helicase encoded by

the yeast subtelomeric Y' element, is induced in survivors defective for telomerase. J Biol Chem *273*,

33360-33366.

Zakian, V.A., and Blanton, H.M. (1988). Distribution of telomere-associated sequences on natural

chromosomes in Saccharomyces cerevisiae. Mol Cell Biol *8*, 2257-2260.

Zhao, Y., Coelho, C., Hughes, A.L., Lazar-Stefanita, L., Yang, S., Brooks, A.N., Walker, R.S.K., Zhang,

W., Lauer, S., Hernandez, C*., et al.* (2023). Debugging and consolidating multiple synthetic

chromosomes reveals combinatorial genetic interactions. Cell *186*, 5220-5236 e5216.

1020

1021

1022   **Figure supplements**

1023

1024

1025   **Figure 2-figure supplement 1.**

1026

1027

1028 **Figure 2-figure supplement 2**

1029

1030 **Figure 2-figure supplement 3**

1031

Figure 2-figure supplement 4

1033

Figure 2-figure supplement 5

1035

1036 **Figure 2-figure supplement 6**

1037

1038   **Figure 2-figure supplement 7**

A

B

1039

1040 **Figure 2-figure supplement 8**

1041

Figure 3-figure supplement 1

1043

Figure 5-figure supplement 1

1045

1046 **Figure 5-figure supplement 2**

1047

1048 **Figure 5-figure supplement 3**

1049

1050 **Figure 6-figure supplement 1**

1051

Figure 6-figure supplement 2

1053

Figure 6-figure supplement 3

1055

Figure 6-figure supplement 4

1057

Figure 6-figure supplement 5

A

B

1059

1060 **Figure 6-figure supplement 6**

A

B

1061

Figure 6-figure supplement 7

1062

1063