Title: Lineage maintenance due to assortative mating but not ecological divergence in a

sympatric sibling species pair

Mitchell Irvine[1]*, Zachary Stewart[1], Nagalingam Kumaran[2], Chapa G. Manawaduge[1],

Solomon Balagawi[3], Brendan Missenden[4], Melissa Starkie[4], Anthony R. Clarke[1], David

Hurwood[1], and Peter Prentis[1]

* Correspondence: mitchell.irvine@hdr.qut.edu.au

Mitchell Irvine (ORCID ID – 0009-0004-2338-8463)

[1]School of Biological and Environmental Sciences, Queensland University of Technology,

GPO Box 2434, Brisbane, Qld, Australia

Abstract

Background

Lineage theory suggests that the development of traits associated with reproductive isolation coupled with traits associated with ecological differentiation are essential for the maintenance of sympatric lineages. However, the relative importance of these factors have rarely been tested simultaneously. *Bactrocera tryoni* and *B. neohumeralis* are a tephritid fruit fly species pair that have significant overlap in geographic range and host use, with time of male mating the only known difference in their mating systems. Using this system, we tested the relative importance of ecological differentiation versus assortative mating in sympatric lineage maintenance.

Results

Genome-wide SNP analyses found strong genetic differentiation between the species with no evidence for hybridization in the field. Most outlier SNPs were restricted to narrow regions towards the centromeres and telomeres of chromosomes. Enrichment of annotation terms indicated an overabundance of genes with the 'abnormal neuroanatomy' term. Terms of interest associated with sleep and circadian rhythm, potentially important to the allochronic reproductive barrier, were non-enriched. Ecological data found no evidence for ecological divergence or competitive displacement between the two species based on significant positive correlations between species numbers trapped at different times of the year, trapped in different habitats within a region, or when reared from fruit.

Conclusions

Our study highlights the significance of assortative mating over ecological differentiation for sympatric lineage maintenance of the *B. tryoni/B. neohumeralis* sibling pair. The paper

represents one of the most well-characterized examples of the importance of genomic

divergence in the coexistence of two closely related species.

Keywords 3-10 words

Bactrocera tryoni, *Bactrocera neohumeralis*, lineage maintenance, coexistence, Tephritidae,

competitive displacement, recombination

Background

Species are increasingly defined as separately evolving lineages [1-4]. If lineages remain

separate, then attributes used to define a species, such as unique mate recognition, distinct

phenotypes and adaptation to specific habitats, may (or may not) emerge through selection,

genetic drift, or pleiotropic change [5, 6]. However, while lineage theory may be coming

widely accepted, the questions faced by evolutionary biologists of what drives lineage

divergence and then keeps lineages separate in sympatry, remain [7, 8].

Central to our understanding of lineage maintenance, particularly between sympatric or

parapatric lineages, are the opposing roles of natural selection and recombination on the

formation or dissolution of lineage boundaries. Natural selection promotes the development

of lineage boundaries through the formation of locally adapted gene complexes; while free

recombination acts to dissolve them through breaking down the associations of such

complexes [9-11]. A reduction in recombination rates is considered necessary for lineage

boundaries to persist and strengthen over time [12]. Traits associated with reproductive

isolation, such as those involved in assortative mating, are considered effective mechanisms

for recombination reduction and essential for lineage divergence in the presence of gene flow

when coupled with the development of traits associated with local adaptation [10, 13].

On an ecological scale, for two lineages to persist in sympatry is considered to be reliant on

two criteria; their ability to overcome the effects of hybridization through the development of

reproductive isolation, and the development of niche partitioning to avoid inter-lineage

competition asymmetries[11, 14, 15]. Without meeting these criteria, theory suggests that

lineages will likely dissolve through hybridization or cease to exist in sympatry through

competitive displacement [14, 16]. Few empirical studies have examined evidence for

assortative mating and competitive displacement simultaneously in the same species pair.

The Australian frugivorous tephritids (Diptera: Tephritidae), *Bactrocera tryoni* (Froggatt) and

Bactrocera neohumeralis (Hardy), are a fruit fly sibling pair which have been studied in

terms of their speciation histories and genetic stability for over 60 years [17-26]. Genetically,

B. tryoni and *B. neohumeralis* are very similar [23, 27, 28], with the sharing of

polymorphisms in coding and non-coding regions indicating continued genetic exchange or

recent separation [22, 24, 27]. The species show only minor variation in their morphology;

the humeral calli is bright yellow in *B. tryoni* and brown in *B. neohumeralis* [29]. While

intermediate colour variants are well documented, genetic analysis using microsatellites

suggest that intermediates represent phenotypic variation within either species rather than

hybrid individuals [21, 24, 30, 31]. While *B. tryoni* is typically more abundant than *B.*

neohumeralis, the geographic ranges of the two species extensively overlap, with the endemic

range of *B. neohumeralis* being entirely encompassed by the range of *B. tryoni* except for a

few islands in the Torres Strait north of Queensland (Fig. 1a,b) [32]. Both *B. tryoni* and *B.*

neohumeralis are highly polyphagous, using fruit from multiple plant families for breeding,

with host utilization data [presented as 'simple' lists of host fruit that either species have been

reared from at least once with little or no further information] showing significant overlap

between the pair (Fig. 1c) [33-35].

Figure 1. Range and host overlap between *Bactrocera tryoni* and *B. neohumeralis*. A)

Distribution of both species in Australia. B) Distribution of both species in the Cape York

Peninsula and Torres Strait Islands. C) Host usage overlap between species. Numbers in

circles represent the known number of host species utilized. Range and host data gathered

from Hancock, E.L. [33], Leblanc, Vueti [34], QDPC [35]. Note: Recent expansion of *B.*

tryoni into central and western Victoria is not shown because it is still under regulatory

control in many places and is not associated with the species' evolutionary history.

So closely related are *B. tryoni* and *B. neohumeralis* that only one known feature

unambiguously defines them as separate biological species: their time of mating [19, 36, 37].

Bactrocera neohumeralis mates during the middle of the day at high light intensity over a

mating window of three to seven hours, while *B. tryoni* initiates mating during dusk at low

light intensity in a mating window of approximately one hour [38]. This difference in mating

time is the only known mechanism maintaining reproductive isolation between the species

[18-20, 26, 37], with other components of their mating system not known to be different [38-

40], including an absence of post-zygotic barriers [41]. Recent work shows that female *B.*

neohumeralis will, like *B. tryoni*, also mate at dusk, at which time *B. tryoni* males do not

discriminate between females of either species [41]. This caused Yeap et al [41]. to conclude

"*the mating time difference may be a weaker reproductive isolating barrier than once*

assumed".

If mating time is a weak reproductive barrier between the two species, then other factors must

be playing a role in maintaining separate lineages; for example, genomic incompatibility [42,

43]. Additionally, for herbivorous insects such as fruit flies, lineage divergence is largely

considered to be driven by host switching and host specialization that reduces the effects of

competition and natural enemies [44-48] and so ecological factors also need to be considered

when investigating lineage maintenance. Conversely the lineages may not be evolutionarily

stable, in which case evidence for genetic recombination should be expected from sympatric

field collections of the two taxa.

In this study we conducted a comprehensive investigation of how the *B. tryoni* and *B.*

neohumeralis lineages have been maintained in sympatry through dense geographic sampling

of nuclear genomes and rigorously collected ecological data. Furthermore, we determined

whether specific genomic regions are associated with reproductive isolation between the two

species. Despite utilizing the same resources in their environment, we found no evidence of

competitive displacement between the two species. Notably, our findings demonstrated that

B. tryoni and *B. neohumeralis* exhibit strong genomic differentiation that can be attributed to

specific genomic regions that likely contribute to maintaining the two as distinct lineages in

sympatry.

Results

Molecular Data

Read processing, alignment and SNP calling

After demultiplexing samples, we obtained between 2,596,430 and 5,228,647 high-quality

sequence reads for each sample. Alignment to the *B. tryoni* genome assembly using BWA-

MEM indicated that most samples had high mapping rates (median alignment percentage =

74.98%), with a small number of samples having low sequence alignment (minimum

alignment percentage = 0.38%); see Table S2 and S3 for full details of read counts and

alignment percentages. 28,334 single nucleotide polymorphisms (SNPs) were identified using

Freebayes from the two species that met our threshold for inclusion.

Populations cluster by species and not geography

We found that populations clustered based on species and not geography in all analyses.

Principal Component Analysis (PCA) (Fig. 2) and fastSTRUCTURE (Fig. 3) both resolved

species-specific groups, demonstrating clear genetic delineation of the two species and an

absence of intermediate genotypes between the species. Several *B. neohumeralis* samples

(genotype codes 526, 528, 529, 530, and 531 from the NS location) appear to be outliers

based on PCA (See Figure S1), but one sample from the same location (527) grouped with

the majority of *B. neohumeralis* samples in ordination space. Additionally, the chooseK.py

function of fastSTRUCTURE predicted three population clusters, with the outlier samples

from the PCA representing the third cluster (See Fig. S2). These outlier samples had poor

genotype calling rates (min = 27.9%, max = 32.3%) compared to sample 527 (91.3%) and

most other samples (median = 89.1%). As such, we believe these to be outliers to result from

sequencing issues and were excluded from further analysis.

Patterns of differentiation among populations (F_{ST}) were also lower between populations

within a species (*B. neohumeralis* = 0.0025796 when ignoring outlier samples, *B. tryoni* =

0.0036269) and two orders of magnitude higher between species (0.10843). Identity-By-State

(IBS) analysis also provided insight into the genetic homogeneity of the two species, with *B.*

tryoni having a mean value of 0.797 and *B. neohumeralis* having a mean value of 0.790

(species with a value closer to 1 are more identical) suggesting that both species are

approximately equally homogenous.

Figure 2. Principal component analysis (PCA) of single nucleotide polymorphisms

(SNPs) for *B. neohumeralis* and *B. tryoni*. Values from the first two eigenvectors plotted for

each sample. Eigenvectors one (horizontal) and two (vertical) explain 10.204% and 1.68% of

the variance in the data, respectively.

Figure 3. fastSTRUCTURE analysis for *B. neohumeralis* and *B. tryoni*. Results at cluster

size (K) equals two for single nucleotide polymorphisms (SNPs) for both species grouped by

collection site. Red and blue colouration of the bars indicates the inferred population

admixture proportions as a percentage (numbers not shown). Three-digit numbers along the

bottoms of each row of bars are genotype ID codes. The intermediate grouping includes

samples with an intermediate phenotype. All three intermediate samples (genotypes 803, 805

and 807) presented as *B. neohumeralis* genotypes.

Outlier loci are clustered in specific chromosome segments between the species

Using outlier detection in BayeScan, we found 357 of the 28,334 SNPs (1.26%) displayed a

significantly elevated level of differentiation between the species. The distribution of outlier

SNPs along autosomal chromosomes (Fig. 4, NC_ prefix) was largely restricted to ~ 16

narrow genomic regions. These narrow genomic regions with outlier loci were not uniformly

distributed across the five autosomal chromosomes with 10 occurring on chromosome ends

and two near the centre of chromosomes. A relatively small number of outlier SNPs were

mapped to 'unplaced contigs' (See Fig. S3).

Figure 4. Outlier plot representation of SNPs for *B. neohumeralis* and *B. tryoni*

identified on the five autosomal chromosomes. F_{ST} values for each SNP were computed

and are plotted as the Y-axis value (left) for each chromosome, with the X-axis representing

the position in base pairs along each chromosome (scaled proportionately); these

chromosomes and contigs are shown as separate rows in the figure, with their NCBI sequence

identifier shown to the right. Outlier SNPs are highlighted in blue.

We investigated genes that contained outlier SNPs or were in close proximity to outlier

SNPs, resulting in the identification of 266 genes. Enrichment of annotation terms indicates

an overabundance of genes with the 'abnormal neuroanatomy' term (n=11 genes, p= 0.0164)

within these candidate genes. Several neurology-related terms which were not enriched but

still of interest included 'nervous system development', 'central nervous system

development', 'sensory system', and 'sensory organ development'. Additional non-enriched

terms of interest included sleep and circadian rhythm-related terms, including: 'abnormal

sleep', 'circadian rhythm', 'abnormal circadian rhythm', and 'abnormal circadian behaviour'.

Inositol-related terms were statistically enriched, which included 'inositol biosynthetic

process' (n=1 gene, p= 0.0115) and 'inositol-1,4,5-trisphosphate 3-kinase activity' (n=1 gene,

p= 0.0467). The 19 genes annotated with these terms are shown in Table S4.

Ecological Data

No differences in seasonal activity between the species

Trap data collected over several years showed no difference in seasonal activity between *B.*

neohumeralis and *B. tryoni* (Fig. 5 & Fig. S4). For any given sampling event there were

generally significantly more *B. tryoni* than *B. neohumeralis* trapped (See Table S5), but the

abundance of both species over time were highly linearly correlated across nine sites with

correlation values ranging from: $r = 0.776$ (Ayr, $p < 0.001$) to $r = 0.993$ (Maryborough, $p <$

0.001) (See Table S6). South Johnstone was the only site with a non-significant result ($r =$

0.926; $p = 0.074$) however, limited collections were recorded at this site ($n = 4$).

Figure 5. Correlation of seasonal abundance of *B. tryoni* and *B. neohumeralis*. Samples

were collected at multiple locations throughout Queensland, Australia. Scatterplots were

developed using monthly mean of trap catches over 2 - 7 years. Data from May [49].

No differences in habitat use between the species

There was a significant correlation between *B. tryoni* and *B. neohumeralis* abundances in

their habitat use (Fig. 6). Within a habitat type, there was a significant correlation in the two

species monthly abundances for all habitat types, with the correlation ranging from r = 0.54

(dry forest, p <0.001) to r = 0.90 (mixed farming, p < 0.001) (See Table S8).

Figure 6. Correlation in habitat use of *Bactrocera tryoni* and *B. neohumeralis*. Samples

were collected from different human-defined habitat types within the Bundaberg district of

Queensland. Each data point is the adult trap catch of both species for a trap run for three

consecutive days. For each habitat type, nine traps were run per month for seven months.

Limited differences in host use between species

One-thousand, six hundred and seven fruits were sampled from the field and the emergence

of *B. tryoni* and *B. neohumeralis* adults counted. For white sapote, mulberry, peach, plum,

carambola and nectarine, sufficient samples were collected to carry out formal analysis.

Highly significant correlations between the number of *B. tryoni* and *B. neohumeralis* adults

reared from each of these fruit types was found (correlation ranging from r = 0.904, p <0.001

for white sapote to r = 0.952, p < 0.001 for peach) (Fig. 7 and Table S9). For guava, two

collections occurred: a single *B. tryoni* and no *B. neohumeralis* emerged from the September

2013 collection while 1609 *B. tryoni* and 1353 *B. neohumeralis* emerged from the October

2013 collection; for feijoa a single collection occurred with equal numbers [71 flies each] of

B. tryoni and *B. neohumeralis* emerging. For sapodilla, *Syzigium* spp, grumichama, hog plum,

jabotica, white mulberry, avocado, black sapote, longan, mango and cashew, *B. tryoni* but no

B. neohumeralis emerged.

Figure 7. Correlation in host use of *Bactrocera tryoni* and *B. neohumeralis*. Each data

point is the number of adult flies reared from an individual collection of multiple fruit pieces

from a research orchard at Nambour, Queensland.

Discussion

Assortative mating and reduced recombination in the absence of local niche adaptation

Genetic comparisons of *B. tryoni* and *B. neohumeralis* individuals collected throughout the

overlapping range showed no evidence of hybridization between the species. This was

despite samples of each species being collected from the same traps, i.e., in immediate

sympatry. Several intermediate phenotypes were collected, however, these samples presented

strongly as *B. neohumeralis* genotypes with no indication of hybridization. This is consistent

with the findings of prior studies of the species pair and supports the theory that strong

reproductive barriers exist in non-laboratory conditions [21, 24, 41]. While Yeap et al.[41]

suggest the allochronic reproductive barrier between these taxa may not be as strong as

initially proposed due to their ability to produce viable hybrid offspring under laboratory

conditions, the lack of hybrid individuals identified from natural populations in this study

suggest that assortative mating is effective in maintaining species boundaries.

The linkage of genes involved in assortative mating with those associated with habitat

preference and performance is considered necessary for sympatric speciation [50, 51] and has

been revealed to have involvement in host switching and specialization of phytophagous

insects such as apple maggot flies and pea aphids [52, 53]. In these studies, the genes

involved appear to map tightly together along narrow regions of the chromosome that is

consistent with the patterns observed in our data. These tightly mapped genes have been

theorized as a mechanism for speciation in the presence of geneflow and are generally

considered to occur through chromosomal rearrangements, such as inversions, where the

genes are brought into physical linkage [50, 51]. A small region of modifier alleles can also

act to reduce recombination between populations in the absence of chromosomal

arrangements [12, 13]. Regardless of the mechanism causing genetic variation, the regions of

differentiation are expected to involve locally adapted alleles or gene complexes and result in

reduced recombination between populations [11, 13]. In our data however, there is an

apparent lack of genes relating to local adaptation in the narrow regions containing outlier

SNPs. Instead, genes related to mating behaviour and circadian rhythm have been identified

that we propose are responsible for separating the two species and forming the time-of-day

premating reproductive barrier through assortative mating. While ecological divergence is

generally required for sympatric speciation, sexual selection alone may be sufficient in

forming species boundaries [54, 55].

In the absence of locally adapted genes/gene complexes under selection pressure, it is likely

that the positioning of these narrow regions of genetic variation along the chromosome had

the greatest influence on reducing recombination. Most outlier SNPs observed in our data

were largely restricted to narrow regions located towards the telomeres or centromeres of

chromosomes (Fig. 4). Reduced recombination has been observed in these regions in several

taxa, including *Drosophila*, mice, rabbits, and humans [50, 56-60]. The combination of genes

associated with assortative mating, and their chromosomal positioning may have sufficiently

suppressed recombination, preventing gene flow between species without the need for

selection pressure on locally adapted genes.

A lack of competitive displacement still allows co-existence

Competitive displacement has long been suggested as an important mechanism facilitating

divergence and speciation [61-63]. The ability of two nascent lineages to coexist by

minimizing competition, or by utilizing non-competitive ecological space, should be

considered central to early lineage divergence and persistence [16]. Where two herbivore

populations share resources, then displacement can occur in time (i.e. seasonal activity),

location or host usage [64, 65]. However, our data shows no evidence for this in the *B.*

tryoni/*B. neohumeralis* sibling pair. Both species were active at the same time of the year and

in the same habitats within a landscape. *Bactrocera tryoni* was recovered from a greater

range of host fruits than *B. neohumeralis* and that may be seen as evidence for competitive

displacement (i.e. competitive advantage by *B. neohumeralis* causing *B. tryoni* to use novel

hosts), but we consider it more likely that this pattern simply reflects *B. tryoni*'s greater

abundance in the environment and so greater likelihood of being sampled, rather than a

competition avoidance mechanism. This is because when both species were reared from a

fruit, their numbers were closely correlated. In cases where competition between fruit flies

has been known to cause changes in field host use patterns, one species will dominate in a

given host, pushing a second species to host fruits that the first only irregularly uses [66, 67]:

we did not observe this pattern. The positive correlation in the number of emerging adults of

both species from fruit also offers field data supporting the work of Kay [68], who in

laboratory studies found no interspecific larval competition between the species. Our data

suggests that, contrary to classical niche theory, ecology has little influence on speciation and

lineage maintenance in our sibling pair and that lineage maintenance is largely driven by

genetic factors and assortative mating.

Our study provides a comprehensive comparison of both molecular and ecological data,

between two sympatric sister taxa. High levels of niche conservatism and a lack of

competitive displacement suggests that the formation and persistence of these taxa occurred

in the absence of ecological selection pressures. Further, the development of assortative

mating and the reduction of recombination of tightly linked sites within chromosomal

telomeres and centromeres appears to have sufficiently reduced geneflow between the species

to maintain the lineage boundaries in sympatry, suggesting that chromosomal architecture

and reproductive isolation in the form of assortative mating, had a greater influence on

lineage divergence than ecological factors.

Methods

Molecular data sample collection

Individuals of *B. tryoni* and *B. neohumeralis* were collected in December 2020 and December

2021, using cue lure baited Steiner [69], Lynfield [70] or Paton traps [71]. These traps were

deployed by Queensland Department of Agriculture and Fisheries (QDAF) as part of ongoing

Biosecurity Queensland (BQ) surveillance. A subsample of the trapping sites were selected

from the existing network to represent the geographic range of both species (Fig. 1). Up to 10

individuals of each species were collected per site with a total of 81 *B. tryoni* and 105 *B.*

neohumeralis individuals collected from 19 sites. Detailed trapping records are included in

Table S1. Identity of the samples were confirmed following the descriptions in fruit fly

identification handbooks [29, 72], and the samples were stored at -20°C in 100% ethanol.

Extraction and sequencing

DNA was extracted using the QIAGEN DNeasy® Blood & Tissue Kit following the

manufacturer's protocol. Samples were screened for quality and quantity, using both gel

electrophoresis and Qubit assay, and then sent to Diversity Arrays Technology Pty Ltd,

Canberra (DArT P/L), for DArTseq high-density genotyping. A PstI/SphI, restriction enzyme

combination was used by DArT P/L, and the fragments (up to 90bp) were sequenced on an

Illumina Hiseq2500 as single end reads.

Data sourcing

All relevant analyses made use of the *B. tryoni* genome assembly (Genbank accession =

GCF_016617805.1).

Read alignment and SNP calling

Reads were demultiplexed using process_radtags as part of Stacks v2.60 [73]. Read mapping

took place using BWA-MEM [74] to produce a SAM alignment file with appropriate read

group specification for each sample. Subsequently, samtools [75] produced sorted and

indexed BAM alignment files. Freebayes v1.3.6 [76] was used in an iterative process to

predict SNPs in all samples. Firstly, each sample had SNP calling performed individually

using freebayes; output Variant Call Format (VCF) files were left-aligned using BCFtools

[77] and block substitutions were decomposed using vt [78]. Resulting VCF files were

merged using BCFtools, and this VCF file was used as input for a second round of SNP

calling using freebayes (-@ *file.vcf --only-use-input-alleles*) to ensure all samples were

genotyped at the same locations. The variant calls were filtered using a method based upon

James, Arenas-Castro [79] wherein SNPs were removed if they had missing data in > 50% of

the population, had a quality score < 30, a depth < 3 read alignments, and a minor allele count

< 1; we also filtered SNPs with a minor allele frequency < 5%.

Population structure

To investigate whether admixture was occurring between the two species, we used

fastSTRUCTURE v1.0 [80]. A simple prior of K=1-10 was used, after which the chooseK.py

utility program provided with fastSTRUCTURE was used to assess the optimal number of

genetic clusters to explain the structure in the species populations. Plots were generated using

pophelper v2.3.1 [81] in R.

Additionally, we ran PCA to provide further visualisation of the variations within and

between the two species, using the SNPRelate v1.24.0 [82] package in R. The linkage

disequilibrium-based SNP pruning function provided with this software was run with default

parameters to mitigate any bias associated with linked SNP clusters upon the downstream

PCA. We similarly performed this SNP pruning prior to calculation of IBS proportions for

samples using SNPRelate.

Mean F_{ST} values [83] were calculated both within and between the two species using

VCFtools [84]. Within species comparisons were conducted for each species separately and

samples from each collection site were considered as separate populations; for between

species comparison, the samples were grouped by species and entered as two separate

populations.

Outlier SNPs and gene candidates

Variant predictions were converted from VCF to GESTE format using PGDSpider v2.1.1.5

[85], then outlier SNPs were predicted using Bayescan v2.1 [86] with default parameters.

Genes containing outlier SNPs within their gene model (exons and/or introns), as well as

genes who were the nearest to an intergenic outlier SNP were identified using a custom

script.

Candidate genes were queried against the *Drosophila melanogaster* gene models

[87](FlyBase release FB2023_01) using MMseqs2, and the FlyBase automated gene

summaries of the best hit were parsed for functional and phenotypic annotations which we

attributed to our sequences if a minimum E-value of $1e^{-10}$ was achieved.

Seasonal abundance

A historical dataset [49] was used for the comparison of seasonal abundance of *B.*

neohumeralis with *B. tryoni*. This dataset has been previously used to report *B. tryoni* [but not

B. neohumeralis] seasonal phenology [88], and so the phenology patterns *per se,* are not

explored here. Rather, we focus on evidence for temporal niche segregation in *B.*

neohumeralis and *B. tryoni* through paired t-tests and Pearson correlation analysis using the R

package ggpubr [89]. We predict that if there is temporal niche displacement between the two

species, then there should be no, or even negative correlation between their seasonal

abundances. We interpret significant positive correlation between their populations as a lack

of evidence of temporal niche displacement.

Detailed methodology on the fly trapping dataset is described elsewhere [49, 88]. In

summary, 15 trapping sites across Queensland, Australia, were used, ranging from temperate

Stanthorpe in the south to tropical Cairns, ~1500km to the north (See Table S5). Fly trapping

was carried out for up to six years per location using a fruit-based liquid lure trap [90]. At

each location ten traps were maintained, with traps cleared weekly. The data in May [49]

varies between sites in how it is presented, and modifications were needed in order to be able

to directly compare *B. tryoni* and *B. neohumeralis* populations and six sites were excluded as

seasonal data were not available (see Table S5 for details). The t-tests and correlation

analysis between *B. tryoni* and *B. neohumeralis* populations was conducted on a site-by-site

basis (i.e., data were never collated across sites), so the different manipulations performed on

the site-specific data sets did not affect results.

Habitat use

This study compared abundance of *B. tryoni* and *B. neohumeralis* across different human-

defined habitat types to determine habitat use patterns. Six habitat types, grassland, suburbia,

horticultural farming, mixed sugarcane and other crops farming (mixed farming hereafter),

dry forest, and wet forest were chosen for the study in the Bundaberg region of Queensland

(~350 km north of Brisbane). Specific descriptions of each habitat type are outlined in Table

S7. For each habitat type there were nine replicate fruit fly traps. Trapping commenced in

September 2010 (spring) and was completed in March 2011 using cuelure baited modified

Steiner traps [69]. The trapping period covers the period of peak yearly activity for *B. tryoni*

[91]. The trapping program involved setting up the traps on a monthly basis at each trap site,

removing the traps at each site three days after setting up and replacing the traps with fresh

wicks and attractants in the following month. Flies collected over the three days in each

month were counted and identified to species level using a fruit fly identification guide [29].

Paired t-tests and Pearson correlation analysis on the total trapping dataset for a habitat type

was performed to compare the relationship in abundance of the two different species at that

habitat type.

Host fruit use

Fruits were collected from the QDAF Maroochy Research Facility, Nambour, Queensland

(~100 km north of Brisbane) from separate orchard blocks that consisted of either mixed

tropical fruit varieties, stone fruit or mangoes, as well as some isolated trees. Small numbers

of ripe or mature green fruit were sampled on a weekly or fortnightly basis between

December 2012 and October 2015. Fruit were selected from a number of trees within the

orchard block and picked from various heights and aspects to obtain a random sample. Fruits

collected were: white sapote (*Casimiroa edulis*), mulberry (*Morus nigra*), peach (*Prunus*

persica), plum (*Prunus domestica*), carambola (*Averrhoa carambola*), nectarine (*Prunus*

persica), guava (*Psidium guajava*), sapodilla (*Manilkara zapota*), *Syzigium* spp, grumichama

(*Eugenia brasiliensis*), hog plum (*Spondias mombin*), jabotica (*Plinia cauliflora*), white

mulberry (*Morus alba*), avocado (*Persea americana*), black sapote (*Diospyros nigra*), longan

(*Dimocarpus longan*), mango (*Mangifera indica*) and cashew (*Anacardium occidentale*).

Fruit were placed in paper bags and transported to the QDAF Brisbane laboratories. Fruit

were counted and weighed and then placed on gauzed plastic containers over vermiculite, in

plastic boxes with gauzed lids to allow ventilation. Boxes were held in a Controlled

Environment Room (26°C and 70%RH) to allow insects to develop through to the pupal

stage. The vermiculite was sieved weekly until all insects had exited fruit and pupated, and

then fruit was inspected before being discarded. Fruit fly pupae that were reared out of fruit

samples were placed into small plastic boxes with gauzed lids containing vermiculite. Once

adult fruit flies had emerged from pupation, they were identified to species level using

morphological taxonomic characters. The abundance of the two fly species from a host was

compared using a paired t-test and Pearson correlation analysis.

List of abbreviations

BQ – Biosecurity Queensland

IBS – Identity by State

PCA – Principal Component Analysis

QDAF – Queensland Department of Agriculture and Fisheries

SNP – Single Nucleotide Polymorphism

VCF – Variant Call Format

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All data generated or analysed during this study are included in this published article [and its

supplementary information files].

Competing interests

The authors declare that they have no competing interests.

Funding

This research was partially funded by the ARC ITTC project IC150100026, Centre for Fruit

Fly Biosecurity Innovation awarded to A.R.C. and P.P.

Authors' contributions

A.R.C. and P.P. conceived the study. N.K., C.G.M., S.B., B.M. and, L.S. conducted field

work and collected samples for molecular and ecological analysis. M.I., N.K., A.R.C. and,

P.P. designed, collated, and analyzed ecological data. Z.S. and P.P. designed and performed

bioinformatics and SNP analysis. M.I., Z.S., P.P. collated analyzed molecular data. M.I., Z.S.,

N.K., C.G.M., M.S., A.R.C., D.H. and P.P. contributed to the writing of the manuscript. All

authors have read and approved the final version.

Acknowledgements

We thank J. Ryan for assisting with generating the distribution map, E. Vueti who carried out

much of the Bundaberg sampling as part of an uncompleted postgraduate study, and L. Senior

(QDAF) for her assistance with rearing flies from fruit that provided data for the host

utilization study. We also thank The Southern Downs Regional Council, The Granite Belt

Growers Association, B. Doohan, The Greenhoose, Lockhart River, D. Masters, J. Kerr and J.

Whistler for their assistance in collecting samples for the molecular analysis.

Author details

[1]School of Biology and Environmental Science, Queensland University of Technology, GPO

Box 2434, Brisbane, Queensland, Australia. [2]Commonwealth Scientific and Industrial

Research Organization (CSIRO), Ecosciences Precinct, Dutton Park, Queensland, Australia.

[3]Horticulture and Forestry Science, Department of Agriculture and Fisheries, P.O. Box 652,

Cairns, Queensland, Australia. [4]Department of Agriculture and Fisheries, Ecosciences

Precinct, Dutton Park, Queensland, Australia

References

1.   de Queiroz K. The general lineage concept of species and the defining properties of

the species category. In: Wilson RA, editor. Species: New interdisciplinary essays.

Cambridge, Massachusetts: MIT Press; 1999. p. 48-89.

2.   Fišer C, Robinson CT, Malard F. Cryptic species as a window into the paradigm shift

of the species concept. Molecular Ecology. 2018;27:613-35.

3.   Shirley MH, Vliet KA, Carr AN, Austin JD. Rigorous approaches to species

delimitation have significant implications for African crocodilian systematics and

conservation. Proceedings of the Royal Society B. 2014;281.

4.   Mayden RL. A hierarchy of species concepts: the denouement in the saga of the

species problem. In: Claridge MF, Dawah HA, Wilson MR, editors. Species: the Units of

Biodiversity. London: Chapman & Hall Ltd.; 1997. p. 381-424.

5.   de Queiroz K. Different species problems and their resolutions. BioEssays.

2005;27(1263-1269).

6.   de Queiroz K. Species concepts and species delimitation. Systematic Biology.

2007;56:879-86.

7.   Saxenhofer M, Schmidt S, Ulrich RG, Heckel G. Secondary contact between diverged

host lineages entails ecological speciation in a European hantavirus. PLoS Biology. 2019;17.

8.   Martin CH, Richards EJ. The paradox behind the pattern of rapid adaptive radiation:

How can the speciation process sustain itself through an early burst? Annual Review of

Ecology, Evolution, and Systematics. 2019;50:569-93.

9.   Hey J. Genes, Categories, and Species : The Evolutionary and Cognitive Cause of the

Species Problem. Oxford, UNITED STATES: Oxford University Press, Incorporated; 2001.

10.   Schluter D, Rieseberg LH. Three problems in the genetics of speciation by selection.

Proceedings of the National Academy of Sciences - PNAS. 2022;119(30):1-e2122153119.

11.   Felsenstein J. Skepticism towards Santa Rosalia, or why are there so few kinds of

animals? Evolution. 1981;35(1):124-38.

12.   Butlin RK. Recombination and speciation. Molecular Ecology. 2005;14(9):2621-35.

13.   Ortiz-Barrientos D, Engelstädter J, Rieseberg LH. Recombination rate evolution and

the origin of species. Trends in Ecology & Evolution. 2016;31(3):226-36.

14.   Weber MG, Strauss SY. Coexistence in Close Relatives: Beyond Competition and

Reproductive Isolation in Sister Taxa. Annual Review of Ecology, Evolution, and

Systematics. 2016;47(1):359-81.

15.   Coyne JA, Orr HA. Speciation: Sinauer Associates Sunderland, MA; 2004.

16.   Germain RM, Hart SP, Turcotte MM, Otto SP, Sakarchi J, Rolland J, et al. On the

origin of coexisting species. Trends in Ecology & Evolution. 2021;36:284-93.

17.   Birch LC. Natural selection between two species of tephritid fruit flies of the genus

Dacus. Evolution. 1961;15:360-74.

18.   Smith PH. Genetic manipulation of the circadian clock's timing of sexual behaviour in

the Queensland fruit flies, *Dacus tryoni* and *Dacus neohumeralis*. Physiological Entomology.

1979;4:71-8.

19.   Lewontin RC, Birch LC. Hybridization as a source of variation for adaptation to new

environments. Evolution. 1966;20(3):315-36.

20.   Meats A, Pike N, An X, Raphael K, Wang WYS. The effects of selection for early

(day) and late (dusk) mating lines of hybrids of *Bactrocera tryoni* and *Bactrocera*

neohumeralis. Genetica. 2003;119(3):283-93.

21.   Pike N. Natural incidence of fruit flies with character states intermediate to those of

the sibling species *Bactrocera tryoni* (Froggatt) and *B. neohumeralis* (Hardy) (Diptera:

Tephritidae). Australian journal of entomology. 2004;43(1):23-7.

22.   Gilchrist AS, Cameron EC, An X, Meats A, Frommer M, Raphael KA. The tryoni

complex of tephritid flies in Australia. Proceedings of 2nd FAO/IAEA International

Conference on Area-Wide Control of Insect Pests: Integrating the Sterile Insect and Related

Nuclear and Other Techniques,. Vienna Austria: IAEA; 2005.

23.   Gilchrist AS, Shearman DCA, Frommer M, Raphael KA, Deshpande NP, Wilkins

MR, et al. The draft genome of the pest tephritid fruit fly Bactrocera tryoni: resources for the

genomic analysis of hybridising species. Bmc Genomics. 2014;15.

24.   Gilchrist AS, Ling AE. DNA microsatellite analysis of naturally occurring colour

intermediates between *Bactrocera tryoni* (Froggatt) and *Bactrocera neohumeralis* (Hardy)

(Diptera: Tephritidae). Australian journal of entomology. 2006;45(2):157-62.

25.   Yeap HL, Lee SF, Robinson F, Mourant RG, Sved JA, Frommer M, et al. Separating

two tightly linked species-defining phenotypes in Bactrocera with hybrid recombinant

analysis. BMC Genetics. 2020;21:132

26.   Raphael KA, Sved JA, Pearce S, Oakeshott JG, Gilchrist AS, Sherwin WB, et al.

Differences in gene regulation in a tephritid model of prezygotic reproductive isolation.

Insect Molecular Biology. 2019;28:689-702.

27.   Wang Y, Yu H, Raphael K, Gilchrist AS. Genetic delineation of sibling species of the

pest fruit fly *Bactocera* (Diptera : Tephritidae) using microsatellites. Bulletin of

Entomological Research. 2003;93(4):351-60.

28.   Morrow JL, Scott L, Congdon B, Yeates D, Frommer M, Sved J. Close genetic

similarity between two sympatric species of tephritid fruit fly reproductively isolated by

mating time. Evolution. 2000;54(3):899-910.

29.   Drew RAI. The tropical fruit flies (Diptera: Tephritidae: Dacinae) of the Australasian

and Oceanian regions. Memoirs of the Queensland Museum. 261989. p. 1-521.

30.   Wolda H. Reproductive isolation between two closely related species of the

Queensland Fruit Fly, *Dacus tryoni* (Frogg) and *D. neohumeralis* Hardy (Diptera :

Tephritidae) : II Genetic variation in humeral callus pattern in each species as compared with

laboratory-bred hybrids. Australian Journal of Zoology. 1967;15:515-39.

31.   Wolda H. Reproductive isolation between two closely related species of the

Queensland Fruit Fly, *Dacus tryoni* (Frogg) and *D. neohumeralis* Hardy (Diptera :

Tephritidae) : I Variation in humeral callus pattern and the occurence of intermediate colour

forms in the wild. Australian Journal of Zoology. 1967;15:501-13.

32.   Osborne R, Meats A, Frommer M, Sved JA, Drew RAI, Robson MK. Australian

distribution of 17 species of fruit flies (Diptera: Tephritidae) caught in cue lure traps in

February 1994. Australian Journal of Entomology. 1997;36(1):45-50.

33.   Hancock DL, E.L. H, Lloyd AC, Elson-Harris MM. The distribution and host plants

of fruit flies (Diptera: Tephritidae) in Australia. Brisbane: Queensland Department of Primary

Industries; 2000.

34.   Leblanc L, Vueti ET, Drew RA, Allwood AJ. Host plant records for fruit flies

(Diptera: Tephritidae: Dacini) in the Pacific Islands. Proceedings of the Hawaiian

Entomological Society. 2012;44:11-53.

35.   QDPC. Queensland Primary Industries Insect Collection Database. In: Fisheries

QDoAa, editor. Queensland Primary Industries Insect Collection Database2021.

36.   Vogt W. A Re-Evaluation of Introgression Between *Dacus Tryoni* and *Dacus*

Neohumeralis (Diptera: Tephritidae). Australian Journal of Zoology. 1977;25(1):59-69.

37.   Birch LC, Vogt WG. Plasticity of taxonomic characters of the Queensland fruit flies

Dacus tryoni and *Dacus neohumeralis* (Tephritidae). Evolution. 1970;24:320-43.

38.   Ekanayake WMTD, Jayasundara MSH, Peek T, Clarke AR, Schutze MK. The mating

system of the true fruit fly *Bactrocera tryoni* and its sister species, *Bactrocera neohumeralis*.

Insect Science. 2017;24(3):478-90.

39.   Bellas T, Fletcher BS. Identification of the major components in the secretion from

the rectal pheromone glands of the Queensland fruit flies, *Dacus tryoni* and *Dacus*

neohumeralis (Diptera: Tephritidae). Journal of Chemical Ecology. 1979;5(5):795-803.

40.   Mankin RW, Lemon M, Harmer AMT, Evans CS, Taylor PW. Time-pattern and

frequency analyses of sounds produced by irradiated and untreated male *Bactrocera tryoni*

(Diptera : Tephritidae) during mating behavior. Annals of the Entomological Society of

America. 2008;101(3):664-74.

41.   Yeap HL, Lee SF, Robinson F, Mourant RG, Sved JA, Frommer M, et al. Separating

two tightly linked species-defining phenotypes in *Bactrocera* with hybrid recombinant

analysis. BMC Genetics. 2020;21(2):132.

42.   Dobzhansky T. Studies on hybrid sterility. II. Localization of sterility factors in

Drosophila pseudoobscura hybrids. Genetics. 1936;21 113-35.

43.   Turelli M, Barton NH, Coyne JA. Theory and speciation. Trends in Ecology &

Evolution. 2001;16:330-43.

44.   Gavrilets S, Losos JB. Adaptive radiation: contrasting theory with data. Science.

2009;323(5915):732-7.

45.   Jeffries MJ, Lawton JH. Enemy free space and the structure of ecological

communities. Biological Journal of the Linnean Society. 1984;23(4):269-86.

46.   Janz N, Nylin S. The oscillation hypothesis of host-plant range and speciation. In:

Tilmon K, editor. Specialization, speciation, and radiation: The evolutionary biology of

herbivorous insects: University of California Press Berkeley; 2008.

47.   Schluter D. The ecology of adaptive radiation: OUP Oxford; 2000.

48.   Yoder JB, Clancey E, Des Roches S, Eastman JM, Gentry L, Godsoe W, et al.

Ecological opportunity and the origin of adaptive radiations. Journal of Evolutionary

Biology. 2010;23(8):1581-96.

49.   May AWS. A taxonomic and ecological study of the Dacinae (fam. Trypetidae) in

Queensland. [Thesis (Ph.D.)]: University of Queensland; 1961.

50.   Ortíz-barrientos D, Reiland J, Hey J, Noor MA. Recombination and the divergence of

hybridizing species. Genetica. 2002;116(2-3):167-78.

51.   Yeaman S. Genomic rearrangements and the evolution of clusters of locally adaptive

loci. Proceedings of the National Academy of Sciences. 2013;110(19):E1743-E51.

52.   Feder JL, Berlocher SH, Roethele JB, Dambroski H, Smith JJ, Perry WL, et al.

Allopatric genetic origins for sympatric host-plant shifts and race formation in *Rhagoletis*.

Proceedings of the National Academy of Sciences. 2003;100(18):10314-9.

53.   Hawthorne DJ, Via S. Genetic linkage of ecological specialization and reproductive

isolation in pea aphids. Nature. 2001;412(6850):904-7.

54.   Servedio MR, Boughman JW. The role of sexual selection in local adaptation and

speciation. Annual Review of Ecology, Evolution, and Systematics. 2017;48:85-109.

55.   Hebets EA, Bern M, McGinley RH, Roberts A, Kershenbaum A, Starrett J, et al.

Sister species diverge in modality-specific courtship signal form and function. Ecology and

Evolution. 2021;11(2):852-71.

56.   Carneiro M, Ferrand N, Nachman MW. Recombination and speciation: loci near

centromeres are more differentiated than loci near telomeres between subspecies of the

European rabbit (*Oryctolagus cuniculus*). Genetics. 2009;181(2):593-606.

57.   Carpenter AT. Synaptonemal complex and recombination nodules in wild-type

Drosophila melanogaster females. Genetics. 1979;92(2):511-41.

58.   Payseur BA, Nachman MW. Microsatellite variation and recombination rate in the

human genome. Genetics. 2000;156(3):1285-98.

59.   Nachman MW, Churchill GA. Heterogeneity in rates of recombination across the

mouse genome. Genetics. 1996;142(2):537-48.

60. Lindsley DL, Sandler L, Counce SJ, Chandley C, Lewis KR, Riley R, et al. The

genetic analysis of meiosis in female *Drosophila melanogaster*. Philosophical Transactions

of the Royal Society of London B, Biological Sciences. 1977;277(955):295-312.

61. Rice WR, Salt GW. Speciation via disruptive selection on habitat preference:

experimental evidence. The American Naturalist. 1988;131:911-7.

62. Thoday JM, Gibson JB. Isolation by disruptive selection. Nature. 1962;193:1164.

63. Via S. Sympatric speciation in animals: the ugly duckling grows up. Trends in

Ecology & Evolution. 2001;16:381-90.

64. Facon B, Hafsi A, Charlery de la Masselière M, Robin S, Massol F, Dubart M, et al.

Joint species distributions reveal the combined effects of host plants, abiotic factors and

species competition as drivers of species abundances in fruit flies. Ecology Letters.

2021;24(9):1905-16.

65. Clarke AR, Measham PF. Competition: A Missing Component of Fruit Fly (Diptera:

Tephritidae) Risk Assessment and Planning. Insects. 2022;13.

66. Charlery de la Masselière M, Ravigné V, Facon B, Lefeuvre P, Massol F, Quilici S, et

al. Changes in phytophagous insect host ranges following the invasion of their community:

Long-term data for fruit flies. Ecology and Evolution. 2017;7(14):5181-90.

67. Hassani IM, Delatte H, Ravaomanarivo LHR, Nouhou S, Duyck P-F. Niche

partitioning via host plants and altitude among fruit flies following the invasion of Bactrocera

dorsalis. Agricultural and Forest Entomology. 2022;24(4):575-85.

68. Kay BJ. The complexity of competition between three native Australian fruit fly

species, Bactrocera tryoni, B. neohumeralis and B. jarvisi in a changing environment [Thesis

(PhD)]: Queensland University of Technology; 2023.

69. Steiner LF. Low-cost plastic fruit fly trap. Journal of Economic Entomology.

1957;50(4):508-9.

70.   Cowley JM, Page FD, Nimmo PR, Cowley DR. Comparison of the effectiveness of

two traps for *Bactrocera tryoni* (Froggatt)(Diptera:Tephritidae) and implications for

quarantine surveillance systems. Australian Journal of Entomology. 1990;29(3):171-6.

71.   Plant Health Australia. Fruit Fly ID Australia: Methods for detection 2023 [Available

from: https://www.fruitflyidentification.org.au/surveillance-and-detection/methods-for-

detection/.

72.   Plant Health Australia. The Australian handbook for the identification of Fruit Flies.

3.0 ed. Canberra, ACT: Plant Health Australia; 2018.

73.   Rochette NC, Rivera Colón AG, Catchen JM. Stacks 2: Analytical methods for

paired end sequencing improve RADseq based population genomics. Molecular ecology.

2019;28(21):4737-54.

74.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. arXiv preprint arXiv:13033997. 2013.

75.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.

76.   Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing.

2012.

77.   Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve

years of SAMtools and BCFtools. Gigascience. 2021;10(2).

78.   Tan A, Abecasis GR, Kang HM. Unified representation of genetic variants.

BIOINFORMATICS. 2015;31(13):2202-4.

79.   James ME, Arenas-Castro H, Groh JS, Allen SL, Engelstädter J, Ortiz-Barrientos D.

Highly Replicated Evolution of Parapatric Ecotypes. Molecular biology and evolution.

2021;38(11):4805-21.

80.   Raj A, Stephens M, Pritchard JK. FastSTRUCTURE: Variational inference of

population structure in large SNP data sets. Genetics (Austin). 2014;197(2):573-89.

81.   Francis RM. pophelper: an R package and web app to analyse and

visualize population structure. Molecular ecology resources. 2017;17(1):27-32.

82.   Zheng X, Levine D, Shen J, Gogarten SM, Laurie C, Weir BS. A high-performance

computing toolset for relatedness and principal component analysis of SNP data.

BIOINFORMATICS. 2012;28(24):3326-8.

83.   Weir BS, Cockerham CC. Estimating F-Statistics for the Analysis of Population

Structure. Evolution. 1984;38(6):1358-70.

84.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The

variant call format and VCFtools. Bioinformatics. 2011;27(15):2156-8.

85.   Lischer HEL, Excoffier L. PGDSpider: an automated data conversion tool for

connecting population genetics and genomics programs. BIOINFORMATICS.

2012;28(2):298-9.

86.   Foll M, Gaggiotti O. A genome-scan method to identify selected loci appropriate for

both bominant and codominant markers: a bayesian perspective. Genetics (Austin).

2008;180(2):977-93.

87.   Larkin A, Marygold SJ, Antonazzo G, Attrill H, dos Santos G, Garapati PV, et al.

FlyBase: updates to the *Drosophila melanogaster* knowledge base. Nucleic acids research.

2021;49(D1):D899-D907.

88.   Muthuthantri S, Maelzer D, Zalucki MP, Clarke AR. Seasonal phenology of

Bactrocera tryoni (Froggatt) (Diptera: Tephritidae) in Queensland. Australian journal of

entomology. 2010;49(3):221-33.

89.   Kassambara A. ggpubr: 'ggplot2' based publication ready plots 2023 [Available from:

https://rpkgs.datanovia.com/ggpubr/.

90.   May AWS, Caldwell NEH. Fruit fly control. Queensland Agricultural Journal.

1944;58:224-9.

91.   Clarke AR, Leach P, Measham PF. The Fallacy of Year-Round Breeding in

Polyphagous Tropical Fruit Flies (Diptera: Tephritidae): Evidence for a Seasonal

Reproductive Arrestment in *Bactrocera* Species. Insects. 2022;13:NA.