Stumpy forms are the predominant transmissible forms of *Trypanosoma brucei*.

Jean Marc Tsagmo Ngoune[1], Parul Sharma[1,2], Aline Crouzols[1] and Brice Rotureau[1,3*]

[1] Trypanosome Transmission Group, Trypanosome Cell Biology Unit, Institut Pasteur,

Université Paris Cité, INSERM U1201, Paris, France.

[2] Sorbonne Université, ED515 Complexité du Vivant, Paris, France

[3] Parasitology Unit, Institut Pasteur of Guinea, Conakry, Guinea.

*Corresponding authors: brice.rotureau@pasteur.fr

Keywords: *Trypanosoma brucei*, stumpy, slender, tsetse, transmission

Abstract

Recently, Schuster *et al.* demonstrated that bloodstream slender forms of African

trypanosomes are readily transmissible to young tsetse flies where they can complete

their complex life cycle (1). In their experimental conditions, a single slender parasite

was sufficient for productive infection. Here, we compared the infectivity of slender and

stumpy bloodstream forms in adult flies with a mature immune system, and without

using any chemical compounds that would alter the insect immune response and/or

promote the infection. After ingestion of slender forms, infected flies were observed

only in one out of 24 batches of non-immunocompetent teneral flies and with a high

number of parasites. In contrast, infected flies were detected in 75% (18/24) of the

batches infected with stumpy parasites, and as few as 10 stumpy parasites produced

mature infections in immune adult flies. We discuss that, although Schuster *et al.* have

unequivocally demonstrated the intrinsic capacity of slender form trypanosomes to

infect young and naive tsetse flies, highlighting the remarkable plasticity and

adaptability of these protists, this phenomenon is unlikely to significantly contribute to

the epidemiology of African trypanosomiases. According to both experimental and field

observations, stumpy forms appear to be the most adapted forms for African

trypanosome transmission from the mammalian host to the tsetse fly vector in natural

conditions.

Introduction

Protist parasites of the *Trypanosoma brucei* group cause Human African Trypanosomiasis (HAT), or sleeping sickness in humans, and nagana in cattle (2). They are transmitted by the blood feeding tsetse fly following a long (at least 3 weeks) and complex (at least 9 distinct stages) cyclical development (review in (3)). In the mammalian host's blood circulation, proliferating slender trypanosomes differentiate into cell cycle-arrested stumpy cells upon quorum sensing when they reach high parasite densities (4-7). This differentiation is thought not only to regulate the parasite load in the reservoir host (8), but also to provide transmissible parasites adapted to pursue the life cycle in the vector host (9). Indeed, stumpy forms express several transcripts and proteins necessary to the next developmental stage in the insect, the procyclic form, including the Protein Associated to Differentiation 1 or PAD1 (5). For decades, arrest of the cell cycle and differentiation to the stumpy stage were presumed essential for the developmental progression of bloodstream trypanosomes to the insect stages.

Recently, Schuster *et al.* demonstrated that slender trypanosomes can also present some intrinsic characteristic of transmissible forms (PAD1 mRNAs and proteins) and are readily transmissible to both young male and female tsetse flies, where they can complete their complex life cycle (1), yet with a lower efficiency than stumpy forms (10). In their experimental conditions, a single slender parasite was sufficient for productive infection. However, these laboratory conditions are significantly different from what is encountered in the field. First, only young teneral flies (1-3 days post-eclosion) with an immature immune system were used. Second, in some experiments, chemical compounds altering the insect immune response (glutathione) and/or promoting the infection (N-acetyl-glucosamine) were added to the infective meal. To assess the importance of these parameters, we challenged the infectivity of slender bloodstream forms in adult tsetse flies, i.e. in conditions closer to the natural situation.

Material and methods

Strains, culture and *in vitro* differentiation

The AnTat 1.1E Paris pleomorphic strain of *Trypanosoma brucei brucei* was derived from a strain originally isolated from a bushbuck in Uganda in 1966 (11). Bloodstream form trypanosomes were cultivated in HMI-9 medium supplemented with 10% (v/v) FBS (12) at 37°C in 5% CO_2. Proliferative slender cells were maintained at densities

lower than 5.10^5 parasites/ml to prevent their natural quorum-sensing-dependent

differentiation into stumpy forms. For *in vitro* slender to stumpy BSF differentiation, we

used 8-pCPT-2′-O-Me-5′-AMP, a nucleotide analogue of 5′-AMP (BIOLOG Life

Science Institute, Germany). Briefly, $2x10^6$ pleomorphic AnTat 1.1E slender forms

were incubated with 8-pCPT-2′-O-Me-5′-AMP (5 µM) for 48 h (13). Freshly

differentiated stumpy forms and slender cells were then centrifuged at 1,400 × g for 10

minutes and resuspended at the appropriate densities in SDM-79 medium

supplemented with 10% FBS. Cells were resuspended at either 10^3, 10^4 or 10^5

parasites / ml. Assuming individual bloodmeal volumes ranging between 10-100 µl,

this would correspond to ingestions of 10-100, 100-1,000 or 1,000-10,000 parasites

per condition.

Tsetse fly maintenance, infection and dissection

Glossina morsitans morsitans tsetse flies were maintained in Roubaud cages at 27°C

and 70% hygrometry and fed through a silicone membrane with fresh mechanically

defibrinated sheep blood (BCL, France). Adult (between 2 and 3 weeks after

emergence) or teneral males (between 24h and 72h post-emergence) were allowed to

ingest parasites through a silicone membrane and without any chemical supplement.

A total of 3 to 5 independent biological replicates per condition were performed with

batches of 50 flies per condition.

Flies were starved for at least 24 hours before being dissected blindly 28 to 31 days

post-ingestion for isolation of all stages from the midgut and salivary glands. For

recovery of all tsetse organs, after rapid isolation of the salivary glands in a first drop

of phosphate buffer saline (PBS), whole tsetse alimentary tracts, from the distal part of

the foregut to the Malpighian tubules, were dissected and arranged lengthways in

another drop of PBS as previously described (14, 15). Isolated organs were then

scrutinized under a microscope at 40x magnification by two independent readers and

infection rates per organ were scored (16).

Immuno-fluorescence analysis (IFA)

Cultured parasites were washed in TDB and spread onto poly-L-lysine coated slides.

For flash methanol fixation, slides were air-dried for 10 min, fixed in methanol at -20°C

for 5 seconds and rehydrated for 20 min in PBS. For immunodetection of stumpy forms,

slides were incubated for 1 h at 37°C with a rabbit polyclonal anti-PAD1 antibody

(kindly provided by Keith Matthews, University of Edinburgh) (5) diluted at 1:300 in

PBS containing 0.1% Bovine Serum Albumin (BSA). After 3 consecutive 5 min washes in PBS, a species and subclass-specific secondary antibody coupled to the Alexa 488 fluorochrome (Jackson ImmunoResearch) diluted at 1:1000 in PBS containing 0.1% BSA was applied for 1 h at 37°C. After washing in PBS, slides were finally stained with 4',6-diamidino-2-phenylindole (DAPI, 1 µg/ml) for visualization of kinetoplast and nuclear DNA content and mounted under coverslips with ProLong antifade reagent (Invitrogen), as previously described (14). Slides were observed under an epifluorescence DMI4000 microscope (Leica) with a 100x oil objective (NA 1.4) to assess the proportion of PAD1-positive cells in the infective meals (n > 100 cells / condition).

Results

Pleomorphic *T. b. brucei* bloodstream forms were either maintained in culture at a density lower than 5.10^5 parasites / ml to prevent quorum-sensing-induced differentiation and obtain only slender forms or induced for differentiation with a 5'-AMP nucleotide analogue to obtain mostly stumpy forms. The expression of PAD1 at the cell surface was assessed by immunofluorescence analysis prior to each experimental infection: no PAD1 expression was detected in the slender group, whereas an average of 63% (52% to 71%, n = 12 replicates) of the induced cells were expressing PAD1. Batches of 50 teneral (< 72h) or adult (2-3 weeks) male tsetse flies were fed in parallel with either slender or stumpy forms at densities corresponding to individual ingestion of about 10, 100 or 1,000 parasites per bloodmeal. In total, 1,384 flies from 12 distinct experimental infections were dissected about 4 weeks (28 to 31 days) after parasite ingestion. Infection rates in midguts and salivary glands were quantified and plotted for each condition (Figure 1 and Table S1).

After ingestion of slender forms, infected flies were observed in only 1 batch out of 24. This occurred in not yet fully immunocompetent teneral flies and with the highest number of ingested parasites (1,000 to 10,000 parasites). In contrast, midgut and salivary glands infected flies were observed in 75% (18/24) and 62.5% (15/24) of the batches infected with stumpy parasites, respectively. As few as 10 stumpy parasites produced mature infections in immunocompetent adult flies and the infection rates were similar whatever the amounts of stumpy forms ingested. However, in more susceptible non-immune teneral flies, the infection rates were increasing with the number of stumpy forms ingested.

Figure 1. Stumpy forms are the predominant transmissible forms. Comparison of
(**A**) midgut and (**B**) salivary gland infection rates in teneral (T) Vs. adult (A) tsetse flies
(batches of 50 flies) infected with 10 to 100 (circles, 3 independent experiments), 100
to 1,000 (squares, 4 independent experiments) or 1,000 to 10,000 (triangles, 5
independent experiments) parasites in the slender (SL in blue) or stumpy (ST in red)
forms.

Discussion

The recent findings of Schuster *et al.* (1) have opened a debate on the traditional view
of the trypanosome life cycle where slender trypanosomes are considered as non-
competent for cyclical development in the insect vector (17). The authors proposed
that their observations could provide a solution to a long-lasting paradox, namely the
successful transmission of parasites in chronic infections, despite low parasitemia.

However, Schuster *et al.* performed all their experimental infections in laboratory
conditions that have been optimized for maximum transmission efficiency, with the use
of teneral flies, and for some experiments, the addition of *N*-acetylglucosamine and
glutathione. Teneral flies remain unfed up to 3-5 days after emergence from their
puparium (1) and are known to be significantly more susceptible to trypanosome
infection as they are immunologically immature (weak immune system and, leaky
peritrophic matrix) (18-21). However, in the wild, tsetse flies remain teneral for 1 to 4
days maximum (before they take their first blood meal), although they subsequently
live as adult flies for up to 9 months (22), which attenuates the possible epidemiological

impact of teneral flies on trypanosome transmission. In addition, all infective meals

were supplemented with 60 mM *N*-acetylglucosamine, an inhibitor of tsetse midgut

lectins enhancing trypanosome infection (23). For infections with monomorphic

parasites, the addition of 12.5 mM glutathione, an antioxidant that reduces the midgut

environment, protects trypanosomes from cell death induced by reactive oxygen

species (24). These chemical compounds inhibit the (already weak) immune response

in teneral flies, and substantially enhance the chances for trypanosomes to develop in

the insect vector.

Conditions are far less favourable in the field; hence we investigated and compared

the infection potential of slender and stumpy forms in adult and teneral flies without the

addition of chemicals. We observed that slender forms were not infective to adult tsetse

flies and only at densities higher than 10^5 parasites / ml. However, in endemic areas,

especially in Western Africa, parasitemia in confirmed cases are usually very low (<

10^4 parasites / ml in Guinea for instance) and it is necessary to concentrate parasites

in blood prior to microscopic examination to increase sensitivity of parasitological

diagnosis (2). Hence, the possible transmission of a few slender trypanosomes from

the blood of individuals with a chronic infection is unlikely to explain the maintenance

of the parasite circulation in tsetse populations.

By contrast, we observed that as few as 10 stumpy parasites are enough to produce

mature infections in both teneral and adult flies, already with a significant efficiency. In

patients with low parasitemia, the quorum-sensing-triggered differentiation of slender

to stumpy forms could be compatible with, or even more adapted to, extravascular

forms present in some tissues and organs, with a limited dilution of the parasites and

parasite factors remaining concentrated locally. Indeed, extravascular PAD1-positive

trypanosomes were detected in high numbers at least in adipose tissues (25) and in

the dermis (26) of experimentally infected mice. The presence of PAD1-positive

extravascular trypanosomes was also assessed in the skin of confirmed gambiense

HAT cases and unconfirmed seropositive individuals in endemic areas (27) (and

unpublished data). This suggests that stumpy trypanosomes accessible to tsetse flies

are likely more abundant than previously estimated in individuals with low parasitemia.

Schuster *et al.* have unequivocally demonstrated the intrinsic capacity of slender form

trypanosomes to infect young and naive tsetse flies, highlighting the remarkable

plasticity and adaptability of these protists. The fine understanding of the underlying

cellular mechanisms and / or transient adaptations involved in this process remains an

exciting challenge. However, this event is unlikely to contribute to the epidemiology of African trypanosomiases in natural settings. According to both experimental and field observations, stumpy forms appear to be the most adapted forms for African trypanosome transmission from the mammalian host to the tsetse fly vector in natural conditions.

Acknowledgements

We thank K. Matthews for providing the anti-PAD1 antibody and P. Bastin for his critical reading of the manuscript.

Funding

This work was supported by the Institut Pasteur, the Programme Investissement d'Avenir of the French Government, Laboratoire d'Excellence, ANR-10-LABX-62-IBEID and ANR-11-LABX-0024-PARAFRAP. This work was supported the French National Agency for Scientific Research projects ANR-18-CE15-0012 TrypaDerm and ANR-19-CE15-0004-02 AdipoTryp. None of these funding sources has a direct scientific or editorial role in the present study.

Author contributions

AC, JMTN and PS performed the experiments. BR designed the study, analysed the data, and wrote the manuscript. JMTN, PS and BR discussed the manuscript.

Competing interest

All authors declare no financial relationships with any organizations that might have an interest in the submitted work in the previous three years, no other relationships or activities that could appear to have influenced the submitted work, and no other relationships or activities that could appear to have influenced the submitted work.

References

1.   Schuster S, Lisack J, Subota I, Zimmermann H, Reuter C, Mueller T, et al. Unexpected plasticity in the life cycle of *Trypanosoma brucei*. eLife. 2021;10.

2.   Buscher P, Cecchi G, Jamonneau V, Priotto G. Human African trypanosomiasis. Lancet. 2017.

3.   Rotureau B, Van Den Abbeele J. Through the dark continent: African
trypanosome development in the tsetse fly. Frontiers in cellular and infection
microbiology. 2013;3:53.

4.   Vassella E, Reuner B, Yutzy B, Boshart M. Differentiation of African
trypanosomes is controlled by a density sensing mechanism which signals cell cycle
arrest via the cAMP pathway. J Cell Sci. 1997;110(21):2661-71.

5.   Dean S, Marchetti R, Kirk K, Matthews KR. A surface transporter family conveys
the trypanosome differentiation signal. Nature. 2009;459(7244):213-7.

6.   Mony BM, MacGregor P, Ivens A, Rojas F, Cowton A, Young J, et al. Genome-
wide dissection of the quorum sensing signalling pathway in *Trypanosoma brucei*.
Nature. 2014;505(7485):681-5.

7.   Rojas F, Silvester E, Young J, Milne R, Tettey M, Houston DR, et al.
Oligopeptide Signaling through TbGPR89 Drives Trypanosome Quorum Sensing. Cell.
2019;176(1-2):306-17 e16.

8.   Turner CM, Aslam N, Dye C. Replication, differentiation, growth and the
virulence of *Trypanosoma brucei* infections. Parasitology. 1995;111 (Pt 3):289-300.

9.   Rico E, Rojas F, Mony BM, Szoor B, Macgregor P, Matthews KR. Bloodstream
form pre-adaptation to the tsetse fly in *Trypanosoma brucei*. Frontiers in cellular and
infection microbiology. 2013;3:78.

10.   Matthews KR, Larcombe S. Comment on 'Unexpected plasticity in the life cycle
of *Trypanosoma brucei*'. eLife. 2022;11.

11.   Le Ray D, Barry JD, Easton C, Vickerman K. First tsetse fly transmission of the
"AnTat" serodeme of *Trypanosoma brucei*. Ann Soc Belg Med Trop. 1977;57(4-5):369-
81.

12.   Hirumi H, Hirumi K. Continuous cultivation of *Trypanosoma brucei* blood stream
forms in a medium containing a low concentration of serum protein without feeder cell
layers. J Parasitol. 1989;75(6):985-9.

13.   Barquilla A, Saldivia M, Diaz R, Bart JM, Vidal I, Calvo E, et al. Third target of
rapamycin complex negatively regulates development of quiescence in *Trypanosoma*
brucei. Proc Natl Acad Sci U S A. 2012;109(36):14399-404.

14.   Rotureau B, Subota I, Bastin P. Molecular bases of cytoskeleton plasticity during
the Trypanosoma brucei parasite cycle. Cell Microbiol. 2011;13(5):705-16.

15. Rotureau B, Subota I, Buisson J, Bastin P. A new asymmetric division
contributes to the continuous production of infective trypanosomes in the tsetse fly.
Development. 2012;139(10):1842-50.

16. Rotureau B, Blisnick T, Subota I, Julkowska D, Cayet N, Perrot S, et al. Flagellar
adhesion in *Trypanosoma brucei* relies on interactions between different skeletal
structures in the flagellum and cell body. J Cell Sci. 2014;127(Pt 1):204-15.

17. Guegan F, Figueiredo L. A two-stage solution. eLife. 2021;10.

18. Wijers DJ. Factors that may influence the infection rate of *Glossina palpalis* with
Trypanosoma gambiense. I. The age of the fly at the time of the infected feed. Ann
Trop Med Parasitol. 1958;52(4):385-90.

19. Walshe DP, Lehane MJ, Haines LR. Post eclosion age predicts the prevalence
of midgut trypanosome infections in Glossina. PLoS ONE. 2011;6(11):e26984.

20. Aksoy S, Gibson WC, Lehane MJ. Interactions between tsetse and
trypanosomes with implications for the control of trypanosomiasis. Adv Parasitol.
2003;53:1-83.

21. Weiss BL, Wang J, Maltz MA, Wu Y, Aksoy S. Trypanosome infection
establishment in the tsetse fly gut is influenced by microbiome-regulated host immune
barriers. PLoS Pathog. 2013;9(4):e1003318.

22. Challier A. The Ecology of Tsetse (Glossina Spp) (Diptera, Glossinidae) - a
Review (1970-1981). Insect Sci Appl. 1982;3(2-3):97-143.

23. Peacock L, Ferris V, Bailey M, Gibson W. Multiple effects of the lectin-inhibitory
sugars D-glucosamine and N-acetyl-glucosamine on tsetse-trypanosome interactions.
Parasitology. 2006;132(Pt 5):651-8.

24. MacLeod ET, Maudlin I, Darby AC, Welburn SC. Antioxidants promote
establishment of trypanosome infections in tsetse. Parasitology. 2007;134(Pt 6):827-
31.

25. Trindade S, Rijo-Ferreira F, Carvalho T, Pinto-Neves D, Guegan F, Aresta-
Branco F, et al. *Trypanosoma brucei* Parasites Occupy and Functionally Adapt to the
Adipose Tissue in Mice. Cell Host Microbe. 2016.

26. Capewell P, Cren-Travaille C, Marchesi F, Johnston P, Clucas C, Benson RA,
et al. The skin is a significant but overlooked anatomical reservoir for vector-borne
African trypanosomes. eLife. 2016;5.

27.   Camara M, Soumah AM, Ilboudo H, Travaille C, Clucas C, Cooper A, et al.
Extravascular Dermal Trypanosomes in Suspected and Confirmed Cases of
gambiense Human African Trypanosomiasis. Clin Infect Dis. 2021;73(1):12-20.
285