1

Sexual antagonism and sex determination in three syngnathid species alongside a male pregnancy gradient

Arseny Dubin[1], Jamie Parker[1], Astrid Boehne[2], Olivia Roth[1]

[1]Marine Evolutionary Biology, Zoological Institute, Kiel University, Am Botanischen Garten 1-9,
24118 Kiel, Germany;

[2]Center for Molecular Biodiversity Research, Zoological Research Museum Alexander Koenig,
Bonn, Germany

Abstract:

The allocation of energy towards gamete production, parental care, mate choice, sex roles, and
sexual dimorphism generates divergence in selection between sexes, leading to sexual
antagonism (SA). Due to the shared genetic makeup a single genomic locus can contain a gene
or allele with differing fitness impacts on each sex. This intralocus sexual conflict is resolved
through intersex bias in gene expression and the formation of sex-linked genomic regions
evolving towards sex determination (SD) systems. SD encompasses environmental SD (ESD),
monogenic SD, and polygenic SD. The latter is believed to represent a transitional state that
occurs during the shift from one SD locus to another. While the precise mechanisms driving
these shifts remain unclear, SA is expected to be a major contributor. To investigate the link
between SA and SD, we selected three species of the syngnathids representing pipefishes and
seahorses along the gradient of their unique male pregnancy evolution, displaying varying sex
roles and sexual dimorphism. By identifying regions of genetic divergence (Fst) and sex-biased
expression, we uncovered that sex role and mate competition, rather than male pregnancy,
primarily drive SA. Processes related to non-coding RNAs and biased allele expression were
detected as crucial mediators of SA. The causative relationship between SD and SA varies
depending on sex role and mate competition dynamics. Most notably, we discovered
intraspecies sex chromosome polymorphism in *Hippocampus erectus*. Overall, our findings
shed light on the interplay between SA and SD, suggesting that understanding SA and its
resolving mechanism is crucial for unraveling the evolution of SD in diverse species.

1. Introduction

A phenotype that directly or indirectly enhances an individual´s reproductive success (i.e., fitness) is expected to increase in frequency within a population through natural selection (Gregory 2009). The selection acting on a phenotype arises from a dynamic interplay between an organism and its environment. The environment influences an organism´s gene expression and embryonic development, thereby shaping its development (Hamdoun and Epel 2007; Bosch et al. 2014; Moczek 2015). Subsequently, this affects its reproduction and survival. In turn, organisms actively interact with their surroundings, selecting and modifying their environment to adjust their probability of survival and reproduction (Laland et al. 1999; Trappes et al. 2022). This renders an organism both subject and object of its own evolution (Levins and Lewontin 1985; Svensson 2018; Erik I Svensson 2023). Such bi-directional cause-and-effect dynamics extend beyond the organism and its abiotic environment to encompass interactions also between sexes of the same species — where one sex selects for certain phenotypes of the opposite sex and vice versa (Svensson 2018; Svensson 2019; Erik I Svensson 2023; Erik I. Svensson 2023).

The emergence of sexual reproduction fostered the formation of male and female gametes in most metazoans, resulting in a division of sexes (Bachtrog et al. 2014; Furman et al. 2020). The allocation of energy into gamete production, parental care, mate choice, sex roles and sexual dimorphism all contribute to differences in natural and sexual selection between the sexes (Parsch and Ellegren 2013; Bachtrog et al. 2014; Mank 2017; Rowe et al. 2018; Svensson 2019; Furman et al. 2020; Erik I. Svensson 2023). This variation in selection can lead to opposing male and female fitness strategies, where the optimal strategy for one sex negatively impacts the other, resulting in sexual antagonism (SA) (Rice and Holland 1997; Lucotte et al. 2016; Svensson 2018; Iversen et al. 2019; Svensson 2019; Tosto et al. 2023). This conflict transforms the relationship between the sexes into a perpetual "tug of war," a reciprocal cause-and-effect dynamic fueled by SA. Due to the shared genome sequence between males and females, a single genomic locus can harbor a gene or allele with divergent fitness effects on the sexes. This divergence in fitness effects of a gene/allele is termed intralocus sexual conflict, and loci that harbour these genes/alleles are defined as sexually antagonistic loci (Bachtrog et al. 2014; Mank 2017; Smith et al. 2023).

Intralocus sexual conflict can be resolved through two primary mechanisms: intersex bias in

gene expression or the confinement of conflicting alleles to the benefiting sex (Bachtrog et al.

2014; Cheng and Kirkpatrick 2016; Wright et al. 2018; Furman et al. 2020; Lichilín et al. 2021;

Smith et al. 2023; Tosto et al. 2023). This confinement can be achieved through reduced

recombination rates and ultimately the formation of sex-linked genomic regions (Bachtrog et al.

2014; Furman et al. 2020; Smith et al. 2023). While sex-biased gene expression can alleviate

intralocus sexual conflict, sex-biased gene expression has been deemed insufficient for its

complete resolution. Genes exhibiting sex-biased expression frequently display genetic

differentiation between the sexes, as evidenced by elevated intersex Fst values (Cheng and

Kirkpatrick 2016; Wright et al. 2018). Consequently, the resolution of intralocus sexual conflict

appears to depend on a balance of differential gene expression and allele fixation (Bachtrog et

al. 2014; Cheng and Kirkpatrick 2016; Wright et al. 2018; Furman et al. 2020; Smith et al. 2023;

Tosto et al. 2023). This association underscores the link between intralocus sexual conflict

resolution and the evolution of sex determination (SD) systems, which often involve formation of

sex-linked genomic regions with suppressed recombination.

The mechanisms underlying SD exhibit remarkable diversity across clades and even closely

related species (Bachtrog et al. 2014; Furman et al. 2020). Mammals and birds rely on generally

well-conserved genetic SD system involving XY and ZW sex chromosomes, respectively

(Bachtrog et al. 2014; Furman et al. 2020). In contrast, reptiles and teleost fish have evolved a

broader range of SD strategies, ranging from environmental SD (ESD), to monogenic SD with a

single master sex determination gene, and polygenic SD (Mank et al. 2006; Mank et al. 2006;

Bachtrog et al. 2014; Furman et al. 2020). The latter is believed to represent a transitional state

that occurs during the shift from one SD locus to another, often referred to as a "turnover" event

(Schartl et al. 2023). The observed variation in SD loci among closely related species indicates

a turnover event in their evolutionary history. The frequency of turnover events varies across

taxa, with some, like the teleost clade *Cichlidae*, experiencing them more frequently than others

(Lichilín et al. 2021). Interestingly, novel SD loci, despite evolving independently, often are co-

opted from a limited subset of genes involved in the classic sex differentiation network. An

exception to this pattern is the de novo evolved salmonid master SD gene (sdY), which arose

from a duplication of IFN regulatory factor 9 (irf9) (Bertho et al. 2018). The sdY gene product,

however, interacts with the female-determining transcription factor foxl2, which belongs to the

classic sex differentiation network. This interaction stops the female differentiation pathway,

facilitating testis development. This confirms the notion that selection for novel SD loci in a

turnover event is restricted to genes within the classic sex differentiation cascade or genes that

interact with it (Sander Van Doorn 2014; Bertho et al. 2018).

While the precise mechanisms driving turnover events remain unclear, SA selection is thought

to be a major contributor. Currently, one of the main research gaps revolves around the

question of precedence: what initiates selection, a sex determination (SD) gene or a sexual

antagonistic conflict (SA)? It is unknown whether an SD gene emerges first, leading to a region

of recombination suppression and subsequent accumulation of SA loci within the region, or

whether SA loci promote the emergence of an SD locus by selecting for a region of

recombination suppression (Sander Van Doorn 2014; Schartl et al. 2023; Smith et al. 2023). SD

research has proposed several models describing the relationships between SD and SA loci

and the roles these relationships play in turnover events, each with varying degrees of support

(Schartl et al. 2023; Smith et al. 2023).

The teleost family *Syngnathidae* is famous for their unconventional morphology and unique form

of parental care – male pregnancy. The degree of parental investment evolved on a gradient

within the family, ranging from simple egg attachment to sealed brood pouch with a placenta-

like structure to supply oxygen, and potentially nutrients, to the developing embryos (Wilson

2001; Skalkos et al. 2020). In addition to differences in parental care among syngnathid males,

there is also variation in sex roles across the species. Some species, e.g., most seahorses,

adhere to conventional sex roles in which the female choose their mate. In other species, e.g.,

most pipefishes, females compete for access to mates and carry costly secondary sexual

signals, while the male chooses the mate and cares for the offspring (Wilson et al. 2003a). Both

varying sex roles with sexual dimorphism as well as differences in parental investment are

expected to contribute to strong divergence in selection between the sexes, which in turn should

be reflected in the underlying genomic and transcriptomic profile.

To investigate the link between sexual antagonistic conflict and sex determination, we selected

three syngnathid species along the gradient of male pregnancy that evolved with different sex

roles and intensities of sexual dimorphism. The species with the most advanced male

pregnancy *Hippocampus erectus* (sealed brood pouch and a placenta-like structure to nurture

the eggs) has evolved with traditional sex role, monogamy, and larger males (Foster and

Vincent 2004). *Syngnathus typhle* and *Nerophis ophidion* represent species with reversed sex

roles and polygamy. *N. ophidion* has a strong sexual dimorphism with large decorated females

(Wilson et al. 2003a). In *S. typhle*, in contrast, males and females are difficult to differentiate
when out of their mating season. *S. typhle* has a more advanced form of male pregnancy with
eggs protected by an inverted brood pouch, whereas *N. ophidion* males have external egg
attachment on the ventral side (Wilson et al. 2003b). Given the profound influence of male
pregnancy on morphology, physiology, and life history, we hypothesized that male pregnancy is
the primary driver of SA relationships in these species. We anticipated either a strong genetic
sex-linkage or a pronounced divergence in expression patterns of genes associated with male
pregnancy adaptation, e.g., those involved in osmotic regulation, oxygen transport, and immune
function. We expected the extent of genetic and expression divergence to correlate with the
gradient of male pregnancy complexity. Alternatively, a different pattern would suggest that
other factors, such as dimorphism and sex roles, also play significant roles. To look for
signatures of SA, we used RNA-seq data obtained from head kidney and gill of males and
females of each species. In addition, we investigated the relationship between SA loci and an
SD system by searching for regions of elevated genetic divergence between the sexes (Fst) and
the accumulation of genes/alleles showing bias in expression in these regions.
144
This study sheds light on sex-linked genomic regions in three syngnathid species, the
underlying processes involved in SA and the relationships between the loci under SA and sex-
linked regions.

2. Methods

2.1 Sample collection, nucleic acid isolation and sequencing

An adult *Syngnathus typhle* male and an adult *Nerophis ophidion* female, were taken from our
animal facility at GEOMAR, Germany. Large animals were chosen to achieve sufficient amounts
of high molecular weight (HMW) DNA for Pacific Biosciences library preparation. *N. ophidion*
displays a strong sexual size dimorphism with large females and small males. We thus sampled
a female *N. ophidion* for genome sequencing.
DNA extraction, library preparation, and PacBIO sequencing was performed by the Norwegian
Sequencing Centre, Oslo, Norway (sequencing.uio.no).

The DNA samples were processed according to the Pacific Biosciences Express library
preparation protocol without prior fragmentation. The final libraries were size-selected with a
BluePippin with 11 kb cut-off and sequenced on a Sequel II instrument using a Sequel II Binding
kit v2.0 with Sequencing chemistry v2.0. Circular Consensus Sequences (CCS) were generated
using the CCS pipeline (SMRT Link v10.1.0.119588) with default settings.

Hippocampus erectus individuals were bred at the GEOMAR aquaria facilities, Kiel, for several
generations prior to the experiment. *Syngnathus typhle* and *Nerophis ophidion* were wild-caught
in the Baltic Sea (54°23′N; 10°11′E, Germany) and subsequently kept in aquaria facilities at
GEOMAR Helmholtz Centre for Ocean Research, Kiel, Germany. Ten males of each species
were used. Five females were used in the case of *N. ophidion* and *S. typhle*, and seven in the
case of *H. erectus*. All fishes were euthanized with an overdose of MS-222 (Tricaine, 500 mg
L−1; Sigma-Aldrich). After dissection, head kidney and gill samples were collected from each
individual and placed in an RNAlater solution (Invitrogen) and incubated at 4°C for one week.
Samples were then stored at −20°C until extraction. Total RNA was extracted using a RNeasy
Mini Kit (Qiagen) according to the manufacturer's protocol. Libraries were prepared using
Illumina TruSeq stranded mRNA kit (2×150bp) and sequenced on an Illumina NovaSeq 6000
instrument at the Competence Centre for Genomic Analysis (CCGA), Kiel.

2.2 Genome and transcriptome assembly and annotation

2.2.1 Genome assembly

The *S. typhle* and *N. ophidion* HiFi reads (CCS reads with >99% accuracy) were assembled
using the Genome Assembly pipeline (SMRT Link 10.1.0.119588) with default settings. Gene
space completeness of the assemblies was assessed with BUSCO (v5.2.2) (Manni et al. 2021)
using the Actinopterygii reference protein set. Continuity of the assemblies was assessed with
Quast (v5.0.2) (Mikheenko et al. 2018). The chromosome-level assembly of *H. erectus* and
supporting annotation used in this study was previously published by Li et.al (PRJNA613176) (Li
et al. 2021).

2.2.2 Transcriptome assembly and gene prediction

The RNA reads generated in this study were quality trimmed with Fastp (v0.20.1) (Chen et al.
2018) with default settings. Reads from both organs were pooled and assembled with Trinity
(v2.8.1) (Grabherr et al. 2011), to generate a single transcriptome assembly per species. In
order to reduce redundancy of transcripts per gene, the resulting assemblies were subjected to
several filtering steps. First, assemblies were filtered with Transrate (v1.0.3) (Smith-Unna et al.
2016) using raw reads as evidence. The software assigns a quality score for the entire
assembly and per contig on the basis of various metrics, like the base quality scores, number of
reads mapped per contig etc., determines an optimal threshold and filters the assembly.
Transrate-filtered assemblies were subjected to CD-Hit (v4.8.1) (Li and Godzik 2006) clustering
transcripts with 98% similarity threshold into one. These contigs were then used to identify
putative protein sequences encoded within the assemblies with TransDecoder (v5.5.0).
Obtained protein sequences were clustered with CD-Hit (v4.8.1) (Li and Godzik 2006) with 95%
similarity threshold. Sequence identifiers of these filtered proteins were used to filter the
Transdecoder CDS fasta file to reduce redundancy by only selecting the CDS sequences whose
protein product passed a similarity filter. We used these filtered cDNA sequences as our
transcriptome assemblies in all subsequent analyses.

2.2.3 Repeat masking and genome annotation

Prior to genome annotation, repetitive elements were identified and masked. EDTA (v2.0) (Xu
and Wang 2007; Gremme et al. 2013; Xiong et al. 2014; Ou and Jiang 2018; Ou et al. 2019; Shi
and Liang 2019; Su et al. 2019; Zhang et al. 2022) was used to produce a *de novo* repeat
library for *S. typhle* and *N. ophidion* assemblies. The resulting repeat libraries were used to
mask the corresponding assemblies with RepeatMasker (v4.1.2-p1) using the settings "-a -gff -
pa 32 -u -xsmall -nolow".
In order to use RNA-sequencing data as evidence for gene prediction, we had to align them to
the genome. For both *S. typhle* and *N. ophidion,* RNA reads were thus aligned to their
respective species' genome assemblies with STAR (v2.7.9a) (Dobin et al. 2013) in two-pass
mode. Only uniquely mapped reads were considered. Filtered transcriptome assemblies were

aligned to the genomes with GMAP (version 2017-11-15) (Wu and Watanabe 2005) with the
following options "-B 5 -t 32 --input-buffer-size=1000000 --output-buffer-size=1000000 -f
samse".

The *S. typhle* and *N. ophidion* genome assemblies were annotated using BRAKER (v2.1.6)
(Lomsadze 2005; Stanke et al. 2006; Gotoh 2008; Stanke et al. 2008; Li et al. 2009; Barnett et
al. 2011; Iwata and Gotoh 2012; Lomsadze et al. 2014; Buchfink et al. 2015; Hoff et al. 2016;
Hoff et al. 2019; Brůna et al. 2020; Brůna et al. 2021). We ran BRAKER twice. First with RNA,
both the unassembled reads and the transcriptome assemblies, then with OrthoDB proteins
(odb10_vertebrata_fasta) (Kriventseva et al. 2019) as evidence. Afterwards, we combined the
annotations with TSEBRA (v1.0.3) (Gabriel et al. 2021), giving the RNA evidence more weight
(pref_braker1.cfg). The cDNA and amino acid sequences were obtained from the final BRAKER
GTF with AGAT (v0.8.0) (Dainat et al. 2023) toolkit using agat_sp_extract_sequences.pl script
with "--aa --clean_final_stop –clean_internal_stop" options to get amino acid sequences, and "-t
exon –merge" to get cDNA. The putative function of predicted protein sequences was
determined with Orthofinder (v2.4.0) (Emms and Kelly 2019) using sequences from the
following fish species:
Danio rerio (GRCz11), Gasterosteus aculeatus (BROADS1), Hippocampus comes
(H_comes_QL1_v1, GCA_001891065.1), Hippocampus erectus (PRJNA613176), Lepisosteus
oculatus (LepOcu1), Oryzias latipes (ASM223467v1), Syngnathus acus
(GCF_901709675.1_fSynAcu1.2), Takifugu rubripes (rubripes.fTakRub1.2), Syngnathus
rostellatus (GCF_901709675.1).

2.2.4 Transcriptome functional annotation

The transcriptomes were functionally annotated in a similar manner to the genomes. We used
predicted proteins from *H. erectus*, *S. typhle* and *N. ophidion* transcriptome assemblies together
with the previously mentioned eight fish species to run orthology analysis with Orthofinder
(v2.4.0) (Emms and Kelly 2019).

250 2.3 Variant calling

251 2.3.1 Genome-independent approach

Following a similar approach as described in El Taher et al. (El Taher et al. 2021), KisSplice
(v2.4.0) (Sacomoto et al. 2012; Lopez-Maestre et al. 2016; Benoit-Pilven et al. 2018) was used
to call SNPs directly from raw RNA reads without prior genome mapping. The SNP data was
then processed with the kissDE R package (v1.14.0) (Sacomoto et al. 2012; Lopez-Maestre et
al. 2016; Benoit-Pilven et al. 2018) to identify SNPs that showed differential expression between
sexes. To predict their functional impact and identify the genes that exhibited differential allele
expression, the SNPs were placed on the corresponding transcriptome assemblies with BLAT
(v36) (Kent 2002) using "–minIdentity=80" option and KisSplice2refTranscriptome (v1.3.3)
(Sacomoto et al. 2012; Lopez-Maestre et al. 2016; Benoit-Pilven et al. 2018). At this step, we
utilized KisSplice2refTranscriptome flags to filter the data, which excludes SNPs that occurred in
multiple assembled genes (SNP_in_mutliple_assembled_genes flag) and could simply
represent a sequencing error (Possible_sequencing_error flag). Then, we filtered the dataset to
include only bi-allelic SNPs.
As we wanted to investigate both, sex-biased allele expression and identify sex-linked regions
within the genome, the aforementioned SNPs were divided into two categories: i) differentially-
expressed alleles (DEA) – alleles that show a bias in expression, i.e., differences in read counts
for pairs of variants, between the sexes. ii) Sex-specific alleles (SSA) (male or female) – were
defined as SNPs where all individuals of one sex are homozygous at this locus, whereas at
least 75% of individuals of the opposite sex are heterozygous at this locus. We considered
these as fixed differences but point out that due to the usage of RNA sequencing data those
could also represent extremes of differential allele expression. However, the KisSplice
developers previously showed that expressed allele frequencies very closely reflect true allele
frequency differences as would be derived from DNA sequencing (Sacomoto et al. 2012; Lopez-
Maestre et al. 2016). In the following, with SNPs of category ii) we considered two potential sex
determination systems – an XY system with a male-specific Y and ZW system with a female-
specific W chromosome.

2.3.2 Genome-based approach

The RNA reads were aligned to the annotated genomes with STAR (v2.7.9a) (Dobin et al. 2013)
in two-pass mode with the options "--outFilterIntronMotifs RemoveNoncanonical --
outSAMunmapped None --outFilterMultimapNmax 1". The alignment data was then processed
with the Genome Analysis Toolkit (GATK) (v4.2.2.0-17) (Poplin et al. 2017) and Picard (v2.25.4)
(Broad Institute 2019). First, duplicate reads were marked (MarkDuplicates) and reads with Ns
in the cigar strings were split into multiple alignments (SplitNCigarReads). Then, potential
variants were called on each sample in gVCF mode (–ERC GVCF option). gVCF files for each
sample were combined into a single gVCF per species and the variants were called with
GATK's joint genotyping method.
As GATK guidelines do not recommend to use standard hard filtering criteria when working with
SNPs called from RNA-seq data, the final VCFs were filtered with "QD < 2.0" and "FS > 30.0"
thresholds applied, and using "--select-type-to-include SNP --restrict-alleles-to BIALLELIC"
options to keep only high quality bi-allelic SNPs.

2.3.3 Fst calculation

Using SNPs resulting from the genome-based approach, we calculated intersex Weir and
Cockerham Fst values for each of the genomes in windows of three sizes – 1Kb, 10Kb, 100Kb
with VCFtools (v0.1.14) (Danecek et al. 2011). To minimize bias introduced by uneven numbers
of males and females, we used data from only 5 individuals and 2 organs of each sex and set a
missing data threshold to 20 percent.

2.3.4 Overlap between approaches

In order to minimize software-related bias, we investigated the overlap between SNPs called by
KisSplice and GATK. To place KisSplice-called variants on a genome, we aligned KisSplice
SNPs to the genome using STAR (v2.7.9a) (Dobin et al. 2013) and processed these alignments
with KisSplice2refgenome (v2.0.5) (Sacomoto et al. 2012; Lopez-Maestre et al. 2016; Benoit-
Pilven et al. 2018). Alleles and their locations were compared using a custom python script.

2.3.5 Differential gene expression analysis

Transcript abundance was calculated with Kallisto (v0.46.1) (Bray et al. 2016) and imported into
DESeq2 (v1.22.2) (Love et al. 2014) with tximport (v1.22.0) (Soneson et al. 2015). Genes with
total counts less than 10 across all samples were removed from the dataset. Expression
between sexes was analyzed using DESeq2 per species and organ with sex representing the
model contrast. The difference in expression was considered significant if an adjusted p-value
was lower than 0.01. The resulting gene lists were used in conjunction with Fst scans to identify
genes that are differentially expressed and located within a region of elevated Fst.

2.3.6 Intersex SNP density analysis – kmerGWAS analysis

For both SNP density calculations and kmerGWAS analysis, we followed the SexFindR
workflow suggested by Grayson and colleagues (Grayson et al. 2022).
The SNP density per sample was calculated with vcftools in 10kb windows on bi-allelic SNPs
called by GATK from previous steps. The results were analyzed using
SNPdensity_permutations_fugu.R script, provided within the SexFindR workflow.
The script calculates the mean number of SNPs in males and females per window and then
calculates the difference between the male and female means.
The KmerGWAS analysis was performed according to the SexFindR recommendations with no
modifications, using kmersGWAS library (Voichek and Weigel 2020), PLINK (v1.07) (Purcell et
al. 2007), ABySS (v2.0.2.) (Jackman et al. 2017), and BLAST (v2.5.0) (Altschul et al. 1990;
Camacho et al. 2009; Boratyn et al. 2012). Results were visualized with R.

3. Results

3.1 Genome assemblies and annotation

To identify sex-linked genomic regions and provide additional support for genome-independent
SNP calling, we generated draft genome assemblies and their annotations for two of the studied
species. While the here used *H. erectus* dataset was already published (Li et al. 2021), the *N.*
ophidion and *S. typhle* genomes were newly generated for this study. From here, the species

will be referred to as HE (*Hippocampus erectus*), NO (*Nerophis ophidion*), and ST (*Syngnathus*
typhle).

The NO genome was sequenced to ~22x coverage with PacBio HiFi reads of 14.9 Kb mean
read length. The resulting assembly length was 1.7 Gb with an N50 of 1.2 Mb. For the ST
genome, the coverage with PacBio HiFi was ~40x with a mean read length of 14.7 Kb. The total
length of the assembly was 405 Mb with an N50 of 2.3 Mb. While we did not reach
chromosomal resolution with these assemblies, the N50 of 1.2 and 2.3 Mb allowed us to detect
large sex-linked regions, blocks of genes with sex-biased allele expression patterns and their
synteny (i.e., corresponding linked locations).

The BUSCO gene space completeness scores revealed high levels of completeness with the
Actinopterygii dataset for HE, NO, and ST assemblies were:

HE: Complete:96.6%/Missing:2.3%,   NO: Complete:93.9%/Missing:4.0%,   and   ST:
Complete:94.4%/Missing:4.1% respectively. The BRAKER2 workflow resulted in 27'499 genes
annotated in ST (BUSCO C:93.7%/M:4.8%), and 38'598 genes in NO (BUSCO
C:93.2%/M:4.3%). For comparison, the current HE annotation contains 20'137 genes (BUSCO
C:97.4%/M:1.7%). The inflated gene number in NO could be attributed to the extremely high
repeat content of the genome which makes gene prediction difficult even when the genome is
masked (Roth et al. 2020), resulting in erroneous predictions. This inflated gene number bias
was to some extent corrected by the orthology assignment. 19'762 genes (98.1%) of HE were
attached to orthogroups, 29'875 (77.44%) of NO, and 23'907 (86.9%) of ST. To facilitate
comparisons across species and their genomes, in the subsequent analyses we only used
genes that had orthologs to the zebrafish *Danio rerio* – 21'685 in HE, 21'801 in NO, and 20'362
in ST. This reduced the bias in gene numbers across our species set.

3.2 Alignment of variant calling approaches

We used SNP data to assess allele frequency differences between the sexes and identify
putative sex-linked regions. For each part, however, a different variant calling approach was
applied. We wanted to avoid that assembly and annotation quality influence differential allele
expression results and thus opted for a genome-independent variant calling approach in this
part of the study. The KisSplice software allows to identify SNPs and alternative splicing events
from raw RNA-seq reads without prior genome information (Lopez-Maestre et al. 2016). The
SNPs were then placed on transcripts assembled from the same RNA-seq datasets. KisSplice

was previously demonstrated to derive true allele frequency differences from expression

differences (Sacomoto et al. 2012; Lopez-Maestre et al. 2016).

The GATK RNA variant call workflow is a well-established, rather conservative protocol that

relies on genome information. GATK-derived reference genome-based data was used to search

for sex-linked regions within the genome and provided additional support for SNPs called with

the genome-independent approach. To compare the two approaches, we placed pre-filtered bi-

allelic SNPs called with KisSplice on the genome. With a custom python script, we compared

both, location of the SNP and the called allele. We found that around 79 to 83 % of KisSplice

SNPs were confirmed by the GATK bi-allelic SNPs suggesting high congruence.

380 3.3 Differential allele expression and sex-linked SNPs

381 3.3.1 Differential allele expression between the sexes

We here considered a scenario where an intralocus sexual conflict would cause a difference in

allele expression. Sex differences in allelic expression point towards an emerging or ongoing

sexual conflict. The accumulation of sex-specific alleles (SSAs) at certain locations in the

genome supports actual sex-linkage of the underlying SNPs/alleles linked to the evolution of a

sex chromosome. Sex chromosomes can ultimately resolve sexual conflicts by confining SA

genes/alleles to the sex they benefit. Species with XY sex determination systems are expected

to have considerably more or only male-specific SNPs due to their male Y chromosome, while

ZW species are expected to have more or only female-specific SNPs.

Here, allele level differential expression between the sexes was looked at specifically. We

categorized bi-allelic SNPs into SSAs male or female, and differentially expressed alleles

(DEAs).

All three examined syngnathid species showed signs of sex-specific allele expression. The

number of DEAs was considerably higher in NO than in the two other species assessed (Ratio:

3:1). Further, we looked at species-specific distributions of male and female SSAs. While male

and female SSAs were identified in NO and ST, HE only harbored male SSAs (Figure 1. Panels

A in each species). Overall, in our analysis ST exhibits the weakest divergence between the

sexes in both DEA and SSA numbers. On the other side of a spectrum, HE had ~100 times

more male SSAs than ST and ~17 times more than NO. To examine the genomic locations of

these male SSAs identified using transcriptome assemblies, genome and transcriptome

annotations were matched. The HE *chr14* appeared to harbor ~2.4 times more genes with

male-specific alleles than the next best hit (124 vs. 52). With the exception of *chr14*, other

notable hits came from unplaced scaffolds. Upon further examination, it appeared that both the

chr14 and the aforementioned scaffolds share all their 1–1 orthologs with *chr1* of *Hippocampus*

abdominalis (HA) suggesting that they all belong to the same chromosome. With that in mind,

the expanded *chr14* would be the size of ~34.3Mb (vs 13.1Mb original length), which is similar

to ~36 Mb of its supposed homologue – the *chr1* of HA.

The expanded *chr14* would contain 327 genes that bear male-specific alleles (~9.5 per Mb)

compared to the now next best 15 hits from *chr4* (~0.7 per Mb). This pattern marks HE as a

candidate XY species.

To assess biological processes and pathways that might be affected by differential allele

expression, we applied an over-representation analysis (ORA) to gene lists that contained DEAs

(Figure 1. Panels B, for each species). The analysis suggested "Cell Cycle", "ncRNA metabolic

process", and "Signaling by Rho GTPases" to be overrepresented in gene lists with DEA in all

three species. The broad KEGG term "Cell Cycle" includes 469 genes involved in cell replication

and nuclear division. The pathways involved in RNA processing and metabolism also appeared

overrepresented in all species. The ncRNA metabolic pathway is affected by differential allele

expression in all studied species, and ncRNA processing only in NO and ST. Some more broad

GO-terms "RNA degradation", "Metabolism of RNA", "tRNA metabolic process", however,

appeared only in some species. The third overrepresented pathway – "Signaling by Rho

GTPases" is involved in many cellular processes and helps cells to respond to external and

internal stimuli via Rho GTPases – a family of molecular-switch proteins (G proteins) (Bar-Sagi

and Hall 2000; Jaffe and Hall 2005; Hodge and Ridley 2016; Perry and Maddox 2019; Boueid et

al. 2020). Several members of the G proteins family were found to be crucial for

spermatogenesis (as examined in mammals) (Lui et al. 2003; Zhang et al. 2010; Zhu et al.

2023).

Processes related to noncoding RNAs (ncRNA metabolic process and ncRNA processing) were

found to be influenced by DEA between sexes in all species. Noncoding RNAs have an

important role in pregnancy progression. miRNAs regulate the physiological processes in a

sexually dimorphic manner, ensuring normal fetal development, successful pregnancy, and

susceptibility to disease (Varì et al. 2021). ncRNAs are implicated in promoting sex-differences,

e.g., X inactivation-specific transcript (XIST) is a long ncRNA that mediates X chromosome

inactivation (Brown et al. 1991). Epigenetic mechanisms are also responsible for governing

gene-hormonal-environmental interactions. There is a considerable amount of data from teleost

fish showing that miRNAs and lncRNAs are involved in sex determination processes and

gonadal differentiation (Bizuayehu et al. 2012; Juanchich et al. 2013; Jing et al. 2014; Lau et al.

2014; Xiao et al. 2014; Juanchich et al. 2016; Tao et al. 2016; Wang et al. 2017; Pinhal et al.

2018; Qiu et al. 2018; Yan et al. 2021). Studies have also brought up potential roles of ncRNAs

in regulation of immune responses in teleost fish (Wang et al. 2018).

HE and NO share a higher number of pathways affected by sex-biased allele expression –

mainly cell maintenance, signal transduction, intracellular transport, and various metabolic

processes (Figure 1. Panels B, for mentioned species). While ST had some overrepresented

pathways involved in cell cycle and metabolism, the main affected pathways appeared to be

related to embryonic morphogenesis and embryo development. Such a reproduction-related

pathway overrepresentation among DEAs might be particularly relevant to understanding sexual

conflict in ST, considering that ST exhibits the weakest differences between sexes in both DEA

and SSA numbers. It is difficult to judge, however, what roles these pathways play in the

sampled organs.

The NOD-like receptor signaling pathway is enriched in genes with DEAs in NO and ST. Among

them were NLR family member X1 (*nlrx1*), nucleotide-binding oligomerization domain-containing

protein 2 (*nod2*) and TNF receptor associated factor 2b (*traf2b*) in NO, TNF receptor-associated

factor 5 (*traf5*) and interleukin-1 beta (*il1b*) in ST. The nucleotide-binding oligomerization

domain-like (NOD-like) receptors are a type of pattern recognition receptors that play a key role

in innate immune system responses (Jones et al. 2016; Chou et al. 2023).

3.3.2 Sex-Specific Alleles (SSAs)

The only species that had enough genes with SSAs to conduct a meaningful enrichment

analysis was HE. From a list of 353 genes with male SSAs, 5 pathways appeared to be

overrepresented; from GO:BP – glycerol-3-phosphate metabolic process, from KEGG – VEGF

signaling pathway, C-type lectin receptor signaling pathway, Herpes simplex virus 1 infection,

and from Reactome – Adaptive Immune System.

Genes from "glycerol-3-phosphate metabolic process" were two glycerol-3-phosphate

dehydrogenases and glycerol kinase 3. Most of the genes in overrepresented adaptive and

innate immune pathways were related to signal transduction, metabolism and cell cycle, and

only indirectly to the immune system. However, some genes with SSAs could support the

suggested sexual immune dimorphism (Roth et al. 2011; Pappert et al. 2023), namely: integrin

alpha 5 (*itga5*), caspase-9 (*casp9*), mechanistic target of rapamycin kinase (*mtor*), nuclear

factor of activated T cells 2a (*nfatc2a*), and cluster of differentiation 40 (*cd40*).

Both ST and NO had genes with male and female SSAs. In NO – 24M/24F genes, and in ST

6F/4M. Since a meaningful enrichment analysis is impossible with the low number of genes

involved, we scanned the lists manually and looked for genes that might be involved in SA.

Considering sampled tissues functions, we expected to see signatures of SA in genes related to

the innate and adaptive immune systems in both tissues, and cellular transport in gill, since

fathers might need more oxygen when they are pregnant (Goncalves et al. 2015). Notable hits

included: huntingtin (*htt*, NO-F), cryptochrome circadian regulator 2 (*cry2*, NO-F), macroH2A.1

histone (*macroh2a1*, NO-F), T cell immune regulator 1, ATPase H+ transporting V0 subunit a3b

(*tcirg1b*, NO-M), transporter associated with antigen processing 1 (*tap1*, NO-M), cathepsin K

(*ctsk*, NO-M), ATP binding cassette subfamily B member 9 (*abcb9*, NO-M), and integrin beta 4

(*itgb4*, ST-F).

3.4 The search for sex-linked genome regions

We next integrated results from our sex-biased SNPs analysis above with whole-genome

intersex Fst scans. Genes were sorted according to their Fst and SNP density across the

genome. We further compared our results to the study by Long et al. that focused on sex

chromosomes in two seahorse species, *Hippocampus erectus* and *Hippocampus abdominalis*,

and the pipefish species *Syngnathus scovelli* (Long et al. 2023). This required additional

kmerGWAS analysis.

3.4.1 Genome-wide picture

We calculated intersex Fst values for all three genomes in windows of 1kb, 10kb and 100kb.

The 10kb and 100kb scans provided an overall look at the genome and facilitated the

identification of regions or chromosomes with elevated Fst values that might be indicative of

sex-linkage. If a sex chromosome or sex-linked region within a genome has accumulated

sufficient fixed differences between the sexes, we would expect to see a distinct Fst peak at that

location. For Weir Fst estimation we expected 0.5 Fst values for a fixed difference between the

homozygous and the heterozygous sex (Gammerdinger et al. 2020). For 10kb and 100kb

windows we used the threshold of 0.4. We considered a lower 0.4 threshold since values were

calculated over large genomic regions that might include sites with low Fst, hence lowering the

overall Fst in a given window. Each species exhibited a unique distribution pattern of Fst values

across the genome, as shown in Figure 2A. ST exhibited the lowest overall Fst values, with

regions of elevated Fst being sparsely distributed along the genome (33 10Kb windows with Fst

>=0.4 or ~0.08 per Mb). This finding aligns with our observations on sex-biased allele

expression and complements the recently published data by Long et al., which focused on a

different species within the same genus, *Syngnathus scovelli*. Both NO and HE had similar

overall Fst levels, with NO having 1222 10Kb >=0.4 windows (~0.7 per Mb) and HE 721 10Kb

>=0.4 windows (~1.7 per Mb).

However, in NO the elevated regions were scattered throughout the genome, while HE had

three chromosome-size scaffolds differentiating from the rest of the genome with *chr14* having

the highest number of windows with elevated Fst (183 windows or ~13.9 per Mb) and the next

two *chr1* and *chr18* with fewer hits (31 and 21 windows or ~1.2 and ~2.1 per Mb respectively).

The fragmented nature of the genome assembly and the genome size could be responsible for

sparse distribution of elevated Fst windows in NO, i.e., most of the contig hits can potentially

come from a single chromosome. But considering that NO possesses both male and female

SSAs (like ST), we think it is unlikely that there is a single sex-linked region or chromosome.

3.4.2 Elevated Fst gene scan

The results from a 1kb-window scan were used to identify individual genes that overlap with

elevated Fst windows. Using this more sensitive approach putative sex-linked genes can be

identified. We used the same approach to determine the Fst threshold as with previous scans.

For HE and NO we used 0.5 threshold for the reasons described above and for ST we

decreased it to 0.33 to see slight bias in allele frequencies. Whenever an annotation of a

particular gene overlapped with an elevated Fst window, this gene was marked and the window

was annotated as belonging to this gene.

Matching the above-described scans, there was no distinct region of elevated Fst. Both NO and

HE had genes with intersex Fst >0.9, which implies near-complete fixation of alleles in each

sex, i.e., both sexes are homozygous at this locus and possess different alleles. Specifically, in

NO, three genes demonstrated this pattern: proteolipid protein 1b (plp1b – 1.0 Fst, genomic

location – ctg.000011F), tetratricopeptide repeat protein 3 (ttc3 – 0.92 Fst, genomic location –

ctg.005052F) and transcobalamin beta b (tcnbb – 0.91 Fst, genomic location – ctg.000011F).

Similarly, in HE, near-complete allele fixation was observed for two genes: interferon regulatory

factor 2 (irf2 – 0.92Fst, genomic location - chr18), and most notably anti-Müllerian hormone

(amh – 0.91 Fst, genomic location – chr13). Neither of those genes belong to the same contigs,

or form a distinct cluster of genes with elevated Fst. In ST, the gene with the highest value was

trafficking kinesin protein 2 (*trak2* – 0.8 Fst, genomic location – *ctg.000074F*).

3.4.3 Fst-SBGs overlap

Genes that show intersex bias in expression were also found to have elevated intersex Fst. In
humans and flies the Fst scores of SBGs follow a "twin peaks" pattern, i.e., genes with an
intermediate bias in expression show the highest intersex Fst, interpreted as targets of ongoing
sex-specific selection on viability (Cheng and Kirkpatrick 2016).

Resolution of sexual conflict may be insufficient via bias in expression in some genes, causing
alleles to diverge between the sexes. SBGs with elevated intersex Fst in guppies, however,
were suggested to relate to the formation of sex-specific genetic architecture rather than
intralocus sexual conflict (Wright et al. 2018).

To identify genomic signatures of SAS, we investigated an overlap between SBGs and the
windows with elevated Fst. Further, to assess sex-linked regions, we investigated clustering of
genomic locations of SBGs with elevated intersex Fst. Because differential gene expression
analysis was done using transcriptome assemblies and Fst scans using genome annotations,
the results had to be compared using orthology assignment. The search was first executed per
organ and then for both organs combined.
The results support the previous Fst window-scan approach. As such, in HE, the same three
chromosomal scaffolds appeared to contain the most SBGs with elevated Fst: *chr14* (196 hits or
~14.9 per Mb), *chr18* (46 hits or ~4.6 per Mb), and *chr1* (28 hits or ~1.15 per Mb). Among
scaffolds with high number of SBGs and elevated Fst, we identified many unplaced scaffolds
e.g., *ctg37* (128 hits), *ctg26* (92 hits), *ctg27* (67 hits), *ctg33* (54 hits), *ctg39* (19 hits). However,
since our synteny analysis with the HA genome showed that all these scaffolds share 1-1
orthologs with *chr1* of HA, it is most likely that they belong to the same HE chromosome –
chr14. The number of overlapping genes declined from here onwards. Notably, one of the well-
known sex-determination genes, anti-Mullerian hormone (*amh*), appears among the most male-
biased genes in both gill and head kidney of HE, and is also among the genes with the highest
Fst in the genome – 0.91 (Figure 2B). The *amh* gene region, however, does not belong to either
of the two candidate sex-linked scaffolds. As such, the *amh*-containing scaffold appeared during
Fst scans of the genome, but was among candidates with low number of elevated Fst-windows
(>300 in the top scaffolds vs 5 in the *amh*-containing scaffold).
In NO the number of SBGs with elevated Fst was not as significant as in HE, and there were
more contigs that contained similar numbers of overlapping genes, which could be a

consequence of a fragmented genome. There were no known sex-determining genes in the
overlap.

ST showed the smallest number of SBGs with elevated Fst, and they could only be identified in
the differential gene expression analysis from the head kidney. As in other species, we could
not narrow it down to a single distinct region or contig enriched in overlapping genes. Most of
the overlapping genes were involved in various metabolic processes, cellular transport, and
aging. However, one of the most male-biased genes with elevated Fst (0.41) was doublesex
and mab-3 related transcription factor 1 (*dmrt1*) – another well-known sex-determination gene in
teleosts.

As we sampled blood-rich organs, the expression profile should not solely cover the specialized
processes in the organs themselves but also show some of the blood expression patterns.

We examined the lists of SBGs with elevated Fst manually for all species and organs. Their
putative functions were deduced based on the information on the Zfin database (Bradford et al.
2022) followed by a literature review. Important to note is that it is difficult to determine their role
in SA relationship due to interdependence of cellular processes and pleiotropic nature of genes
(Stearns 2010). Furthermore, the roles of SBGs and the reasons for differential expression in a
particular organ might not be directly related to the reasons for elevated Fst values at these loci.
When discussing the functions of the genes, we thus payed more attention to their potential
roles in SA relationship.

In HE, among the SBGs with the highest log fold-change and Fst in both organs was *amh* – one
of the key genes involved in vertebrate gonadal differentiation pathway (Morrish and Sinclair
2002; Yatsu et al. 2016; Nagahama et al. 2021; Wagner et al. 2023). Besides that, *amh* also
appears to have functions in the brain (Pfennig et al. 2015; Silva and Giacobini 2021).

Among the SBGS two genes are involved in spermatogenesis. The DEAD (Asp-Glu-Ala-Asp)
box polypeptide 4 (*ddx4*) appeared within SBG/Fst lists in both organs. It is an RNA helicase,
mostly known as a germline factor (Raz 2000; Hickford et al. 2011; Hansen and Pelegri 2021),
has functions outside of germ cell-related processes (Yajima and Wessel 2011; Lasko 2013;
Schwager et al. 2015; Yajima and Wessel 2015). Interestingly, *ddx4* was found to be highly
expressed in certain lung cancer cells (Noyes et al. 2023). The exosome component 10
(*exosc10*) was shown to be important for spermatogenesis. Mutant mice with inactivated
exosc10 in male germ cells had small testes, showed impaired germ cell differentiation and
were subfertile (Jamin et al. 2017). The *exosc10* is also involved in inactivation of X
chromosome in somatic cells via regulation of Xist expression (Ciaudo et al. 2006).

Several genes under SAS were involved in synaptic and neuronal processes.

The low-density lipoprotein receptor-related protein 4 (*lrp4*) is one of the regulators of the Wnt/β-

catenin signaling pathway and is important in the vertebrate neuromuscular junction (Shen et al.

2015; Mosca et al. 2017; Walker et al. 2021). The gene has other roles, namely in bone

homeostasis as shown in mice (Chang et al. 2014; Shen et al. 2015), and roles in tooth

development as shown in zebrafish (Ahn et al. 2017). The interleukin 1 receptor accessory

protein-like 1b (*il1rapl1b*) belongs to a family of IL1/Toll receptors, regulates dendrite and

synapse formation and is responsible for the X-linked intellectual disabilities in humans

(Yamagata et al. 2015; Montani et al. 2018). *Il1rapl1*-knockout mice showed mildly impaired

working memory and remote fear memory, high locomotor activity, and decreased open-space

and height anxiety (Yasumura et al. 2014). In zebrafish *il1rapl1b* is involved in synapse

formation in olfactory sensory neurons (Yoshida and Mishina 2008). Another gene in the list

involved in regulation of the Wnt/β-catenin signaling – Van Gogh-like (Vangl) planar cell polarity

protein 1 (*vangl1*) (Mentink et al. 2018). Certain variants of vangl1 are associated with neural

tube defects in zebrafish, human and mice (Torban et al. 2008; Kibar et al. 2009; Reynolds et al.

2010; Yang et al. 2017). The Triple-A syndrome (*aaas*) gene was among the genes under SAS

in HE. Mutations in *aaas* are responsible for the Allgrove syndrome in humans which is

associated with multiple neurological abnormalities (Handschug 2001; Brooks et al. 2004;

Bizzarri et al. 2013).

Two immune-related genes also appeared among genes under SAS, namely protein tyrosine

phosphatase non-receptor type 22 (*ptpn22*) – negative regulator of T cell receptor signaling

variants of which are associated with severe autoimmune diseases (Fousteri et al. 2013;

Armitage et al. 2021; Tizaoui et al. 2021), and limb development membrane protein 1-like

(*lmbr1l*) – a gene that was shown to be essential for lymphopoiesis in mice, and mutations in

which causing severe immunodeficiency (Choi et al. 2019). Interestingly, *lmbr1l* modulates

lymphopoiesis via negative regulation of the Wnt/β-catenin signaling pathway.

In NO, there were several genes potentially under SAS involved in DNA break repair, and

chromosome maintenance, inactivation, and DNA replication. The tonsoku-like (*tonsl*) gene is a

DNA repair protein, certain variants of which causing abnormalities in skeletal development

(Burrage et al. 2019). Moreover, *tonsl* was found to be regulating epithelial cell immortalisation

in breast cancer (Khatpe et al. 2023). The minichromosome maintenance complex component 4

(*mcm4*) as part of the minichromosome maintenance (MCM) complex is responsible for DNA

replication (Sheu et al. 2014). In humans, mutation in *mcm4* causes genomic instability and

natural killer cell deficiency (Gineau et al. 2012). The partner and localizer of BRCA2 (*palb2*) –

is an important DNA double-strand break gene in humans, mutations in *palb2* are associated

with breast cancers (Wu et al. 2020). In mice *palb2* male mutants showed reduced fertility

(Simhadri et al. 2014), and in zebrafish *palb2* (*fancn* in the publication) knockout caused female-

to-male sex reversal phenotype (Ramanagoudr-Bhojappa et al. 2018).

The H2A.X variant histone (*h2ax*) – a histone variant that is one of the key actors in DNA break

repair and chromatin remodeling (Pinto and Flaus 2010). The *h2ax* is involved in sex

chromosome inactivation during meiosis in male mice (Fernandez-Capetillo et al. 2003; Turner

et al. 2004).

The leucine-rich repeats and WD repeat domain containing 1 (*lrwd1*) is presumed to be the part

of origin recognition complex (ORC). ORC is responsible for initiation of DNA replication and

DNA repair (Shen et al. 2010; Vermeulen et al. 2010; Liu et al. 2023).

The final gene in this group, RAD51 paralog D (*rad51d*), is a member of the RAD51

recombinases that are highly conserved across species (Lin et al. 2006; Greenhough et al.

2023). For successful recombination RAD51 is required to interact with *palb2*, that assists

localization of it to DNA damage sites (Zhang et al. 2009; Greenhough et al. 2023). Mutations in

rad51d are associated with high risk of ovarian cancer in humans (Suszynska et al. 2020;

Greenhough et al. 2023).

Besides DNA damage repair and replication, one of the genes – ATPase

sarcoplasmic/endoplasmic reticulum Ca2+ transporting 3 (*atp2a3*) was found to be among

genes under selection related to hypoosmotic regulation in Takifugu species (Zhang et al.

2020).

3.4.4 Intersex SNP densities and kmerGWAS results in *Hippocampus erectus*

Our analyses identified HE as the most prominent candidate for containing a sex chromosome

among the three investigated species. However, we identified three potential candidates for a

sex chromosome: *chr14*, *chr18*, and *chr1*. The *chr14* has the highest numbers of SSAs and

DEAs as well as the highest number of elevated Fst windows, which makes it the most likely

sex-linked chromosome candidate. At the same time, the classical sex-determination gene –

amh, was among genes with the highest Fst values and showed highest expression bias in

males. The *amh* is located on scaffold *chr13* with an overall low Fst.

In order to give additional support to *chr14* as HE sex chromosome, we compared male/female

SNP densities across the genome and ran a kmerGWAS analysis to exclude that scans on

unphased genomes might have driven the outcome of this study. We run kmerGWAS, since the

latter does not rely on genome information. To do so, we followed the approach suggested by

the SexFindR workflow. Both analyses provided similar results, highlighting *chr14* as a definite

sex-chromosome candidate. The results from male/female SNP density scans were much

noisier. The KmerGWAS analysis resulted in a clear peak for *chr14* (Figure 3B).

3.4.5 Long et al. data comparison

Long and colleagues had recently published a study investigating sex chromosomes in three

syngnathids, including HE. In that study, the HE genome was reconstructed using the genome

of *Hippocampus abdominalis* (HA), a species phylogenetically more derived compared to HE

(Betancur-R et al. 2017) with a larger genome that underwent an additional chromosome fusion

(He et al. 2022; Long et al. 2023).

To compare our results to Long et al. 2022, we performed a synteny analysis between the HA

genome used by Long and the HE genome at our disposal (PRJNA613176) (Li et al. 2021). For

this purpose, we used annotation data from these genomes, identified 1–1 orthologs, and used

locations of these orthologs in each genome to identify homologous chromosomes (Figure 3C).

The sex chromosome that we identified (*chr14*) and *H. abdominalis chr4* identified by Long et al.

were not homologous. For additional tests, the data published by Long et al. 2022 were used.

We chose kmerGWAS because we obtained the cleanest signal using this method, and

because identification of sex-linked kmers is not affected by assembly quality. The resulting sex-

associated kmers were aligned to HE genome. The results of kmerGWAS confirmed Long's

finding, although the signal was not as pronounced as in HE using our data (Figure 3A). This

suggests that in HE, just like in some other teleost species, sex chromosomes are non-stable

and could be population/strain dependent.

4. Discussion

To explore the connection between sexual antagonism (SA) and the formation of sex

determination (SD) systems, we analyzed both the signatures of SA and the mechanisms of SD

across three syngnathid species, which exhibit varying degrees of male pregnancy and sex-role

reversal. Our objective was to determine if the pronounced SA pressure, as exemplified by male

pregnancy and shifts in sex roles, directly influences the emergence of distinct sex-linked

genomic regions, thereby shaping SD systems. Alternatively, we sought to understand whether

SA and SD are related phenomena that evolve in tandem without a direct causal relationship

between the two. Here we provide a comprehensive examination of the dynamics within SA

relationships and their potential impact on the evolution of SD systems in these species.

4.1 Sexual antagonism (SA) in three syngnathid species along the male pregnancy gradient.

In our investigation of SA within the Syngnathidae family, we hypothesized based on prior SA research in other species that there would be a subset of genes experiencing unresolved intralocus sexual conflict, meaning that the divergent fitness effects these genes have on males and females could not be reconciled solely through one of the mechanisms such as differential gene expression or genetic fixation. Such an unresolved conflict at specific loci suggests an involvement in ongoing SA relationships. We focused on two main indicators of ongoing SA : I) Intersex expression bias – differences in gene expression between males and females, highlighting a intralocus conflict that is resolved via sex-specific optimal expression levels. II) Intersex genetic divergence – quantified by the fixation index (Fst), where a high Fst value signifies genetic divergence between sexes as a way to resolve intralocus conflict. Identifying genes that exhibit both indicators is key to understanding SA's role in genetic divergence between sexes and the underlying biological processes that drive the SA relationships. We propose that this SA-induced genetic divergence may influence the evolution of sex determination (SD) systems in syngnathids.

Our analysis also included differentially expressed alleles (DEAs) and sex-specific allele (SSA) expression, offering a more comprehensive view of how genetic and expression differences contribute to SA resolution. These mechanisms often remain unnoticed in SA and SD studies, which primarily concentrate on biased gene expression and genetic divergence signatures alone. Instead of restricting conflicting alleles to one sex through recombination suppression, each sex could express only the beneficial allele, eliminating the need for suppression.

Male pregnancy, alongside with sex-role reversal, likely heightens the divergence in fitness strategies between sexes, thus amplifying SA pressures. By examining SA markers alongside male pregnancy gradient and varying sex roles, we aimed to discern whether SA significantly influences the development of SD systems. The unique syngnathid physiology offers a novel framework to investigate the interplay between SA and SD.

4.1.1 Genes under ongoing SA. SBG-Fst overlap.

In our study, we investigated three syngnathid species alongside the male pregnancy gradient:
Nerophis ophidion (NO) with basic external egg attachment, *Syngnathus typhle* (ST) with an
inverted brood pouch and a placenta-like structure, and *Hippocampus erectus* (HE) with a
sealed brood pouch and a placenta-like structure. We hypothesized the degree of male
pregnancy to be the key driver of SA relationship and consequently affect the formation and
evolution of SD systems within these species. We expected NO to show the least active SA
indicators, SBGs with elevated Fst values or fewer DEAs. In contrast, HE, with the most
advanced pregnancy form, was anticipated to show the strongest SA signals, including a large
sex-linked genomic region rich in pregnancy-related genes. ST was considered intermediate,
with a moderate number of active SA markers and possibly a small sex-linked region or loci.

As an alternative hypothesis, we also considered the potential impact of varying sex roles on SA
and SD systems. In species with traditional sex roles, like HE, we anticipated that males would
be under stronger selection, which might result in a higher count of active SA markers with male
bias and signs of an XY SD system. Conversely, for species where sex roles are reversed (ST
and NO), females could exhibit more SA markers with male bias, potentially resulting in ZW SD
system. This consideration becomes more complex when factoring in mating strategies
(monogamous or polygamous) and other unknown factors, suggesting there might be many
alternative patterns.

Furthermore, whether SA and SD formation processes are linked also remains to be
determined. It is logical to assume that if SA pressure is strong enough and could not be
resolved by bias in expression, this would lead to formation of a sex-linked genomic region.
However, if such mechanisms exist, then SA and SD system formation processes would
probably be unlinked.

The data, however, suggest a more nuanced reality than our initial two hypotheses proposed.

The least divergence between the sexes was observed in ST, as indicated by both the overall
intersex Fst levels across the genome and the numbers of SBGs with elevated intersex Fst
values. This underscores that the resolution of sexual antagonistic conflict primarily occurs
through sex-specific gene and/ or allele expression, or through other unknown mechanisms. An
elevated Fst of 0.4 for *dmrt1*, a known SD gene (Kobayashi et al. 2013), suggests ongoing
sexual conflict at the locus of this SBG. Despite other genes exhibiting higher Fst values in
different genomic locations and the nearly equal numbers of SSAs between the sexes, *dmrt1*
remained a robust SD gene candidate for ST. This is not only because SD predominantly

involves co-option of the same set of genes, but also as its Fst value is close to the expected

0.5 for a fixed difference between sexes (Gammerdinger et al. 2020).

In HE, SBGs with elevated Fst clustered on a single chromosome-length scaffold, suggesting a

putative sex chromosome. Notably, several genes associated with synaptic and neuronal

processes were found among these SBGs. This reminds of a recent study on signature of SAS

on sex chromosome of *Gasterosteus nipponicus* (Japan Sea stickleback) that identified genes

associated to mental disorders and neurological human development near regions of high

intersex Fst on the sex chromosome (Dagilis et al. 2022). The role of these genes in SA,

however, remains as yet undiscovered.

In NO, the SBGs with elevated Fst were scattered throughout the genome and did not cluster

within a particular genomic region. The genome itself showed overall elevated intersex Fst

levels suggesting reduced recombination landscape in this species. The notable group of genes

implicated in SAS, primarily identified through the overlap of Fst and DEG, included genes

associated with DNA break repair, chromosome maintenance, inactivation, and DNA replication.

While it is tempting to speculate that their involvement in SA could be linked to

spermatogenesis, given the importance of these genes in many cellular processes, it remains

difficult to ultimately define their roles in SA relationships in NO.

Our analysis of SBGs with elevated intersex Fst unveils that species with traditional sex roles

and the most advanced form of male pregnancy (HE) showed the strongest SA signature and

male bias which is consistent with both our hypotheses. In contrast, ST with its intermediate

male pregnancy and sex-role reversal, shows the weakest SA signal, while NO, featuring the

basic form of male pregnancy and extreme sex-role reversal, presents a moderate SA signal

alongside notable genetic divergence between sexes. While mate choice is somewhat biased

towards the male sex in both ST and NO (e.g., (Berglund and Rosenqvist 1993), females in ST

have a more pronounced role in mate choice compared to NO females. This proposes sexual

selection on females in ST to be less skewed than in NO, possibly allowing to deal with SA loci

via expression bias alone. Differences in mate choice between NO and ST may be attributed to

their distinct egg production regimes (Sogabe and Ahnesjö 2011), highlighting the impact of

reproductive traits on mating strategies and preferences. In the case of NO, the mating system

may prioritize traits that signal high immediate fecundity, a consequence of the cyclical

readiness of eggs for fertilization. On the other hand, in ST with its steady egg production, a

different set of traits can be under selection, reflecting the ongoing opportunity for spawning.

Our findings suggest that neither male pregnancy as the sole driver of SA selection across

these species nor sex roles. Rather, the dynamics of sex roles a mate choice are pivotal, with

the sex subjected to higher competition being under stronger selection and showing higher

signatures of active SA.

Furthermore, elevated genetic divergence observed in NO suggests alternative mechanisms

may mediate SA selection, underscoring the complexity of these evolutionary dynamics.

4.1.2 Non-coding RNAs as mediators of sexual antagonism in syngnathids. Evidence

from DEAs.

In all three syngnathid species, non-coding RNA processing and metabolism were among the

pathways linked to differential allele expression between the sexes. Enrichment of the GO term

"ncRNA metabolic process" as identified in all examined species, suggests a sex-specific allele

expression of genes associated to chemical reactions and pathways related to non-coding RNA

transcripts (Ashburner et al. 2000; The Gene Ontology Consortium et al. 2023). We observed a

similar enrichment pattern in ncRNA-related pathways when assessing intersex differences in

gene expression in old and young ST individuals (Pappert et al. 2023). This hints towards an

active role of ncRNAs in SA processes in syngnathids. The roles of ncRNAs in SA processes

could be especially important in species with low genetic divergence between the sexes like in

ST.

Non-coding RNAs do not code for a protein product and include ribosomal RNAs (rRNAs) and

transfer RNAs (tRNAs). The latter are involved in a translation process, while regulatory RNAs

are associated to various cellular processes: long-noncoding RNAs (lncRNAs), microRNAs

(miRNAs), piwi-interacting RNAs (piRNAs), small nuclear RNAs (snRNAs) and small nucleolar

RNAs (snoRNAs) (Frías-Lasserre and Villagra 2017; Bartel 2018; Kopp and Mendell 2018;

Anon 2022; Mattick et al. 2023). A plethora of reports discuss effects of regulatory ncRNA

expression between the sexes on development, physiology, immunity and disease (Zhou et al.

2014; Gershoni and Pietrokovski 2017; Zhang et al. 2017; Gal-Oz et al. 2019; Rosspopoff et al.

2023a; Wang et al. 2023). A known role of ncRNA roles in sexual development is the

inactivation of the with XIST in humans and other placental mammals (Brown et al. 1991; Lee et

al. 1999; Dupont and Gribnau 2013; Posynick and Brown 2019; Rosspopoff et al. 2023b).

However, much of the literature on sexually antagonistic co-evolution and intralocus sexual

conflict in non-model organisms is focused on either fixed genetic differences or differential

gene expression as ways to resolve intralocus sexual conflict, while the roles of ncRNAs in SA

relationships remain as yet understudied.

Besides sex-specific effects via regulatory ncRNA, the potential roles of ncRNAs in mediating

SA could be related to expression and processing of rRNA and tRNA. The 5S and 18S rRNA

abundance, and 5S–18S rRNA ratio have been used as sex markers in fish (Shen et al. 2017).

Though Shen et al discuss development of ovaries and testes, variation in rRNA transcript

abundance can have differential importance for sexes in other tissues. Differential rRNA allele

expression could be another mechanism mediating SA relationships in syngnathids. Tissue-

specific rRNA allele expression has been recently described in humans and mice (Parks et al.

2018). Not only differential transcript abundance and allele expression, but also RNA's ability to

form specific structures is hypothesized to have an impact on epigenetic differences between

the sexes as exemplified by a study on variation in R-loop formation in *Drosophila melanogaster*

(Stanek et al. 2022).

4.1.3 Highly divergent alleles may reflect allele-specific expression, not fixation.

Our Weir Fst analysis, based on 10kb windows, is expected to show Fst values around 0.5 for

completely fixed differences between homozygous and heterozygous sexes (Gammerdinger et

al. 2020). While we expected slightly lower mean Fst due to windowing, surprisingly, genes like

irf2 and amh (HE), plp1b, ttc3, tcnbb in (NO), and trak2 (ST) displayed Fst values exceeding

0.8, suggesting near-complete fixation of opposite alleles in each sex. However, the absence of

correlation between read counts and Fst indicates that these values might not reflect true allele

frequency differences. Instead, highly biased or allele-specific expression (ASE) is a more likely

explanation. ASE can arise from chromosome inactivation, regulatory elements, nonsense-

mediated decay, or imprinting (Castel et al. 2015; Fan et al. 2020). While extensively studied in

human pathology (Langmyhr et al. 2021; Rao et al. 2021; Van Beek et al. 2023), sex-specific

ASE in the context of sexually antagonistic selection (SA) remains underexplored.

Understanding the role of these highly divergent but potentially non-fixed alleles in syngnathid

SA relationships presents a significant challenge.

4.2 Unraveling sex determination systems in three syngnathids.

This study aimed to identify sex-determination (SD) systems in three syngnathid fish species
(NO, ST, and HE) and investigate their relationship to loci under ongoing SA. In HE, we
observed a striking over-accumulation of SBGs with and without elevated Fst, DEAs, and SSAs
on a single chromosome scaffold, suggesting a putative sex chromosome. However, in the
other two species (NO and ST), our investigation of loci under SA did not show any similar
accumulation within specific genomic regions.

4.2.1 Case for ESD in pipefishes *Nerophis ophidion* and *Syngnathus typhle*.

Genetic sex determination (SD) systems typically involve a chromosomal region or entire
chromosome linked to one sex, often containing master SD genes. Exceptions exist, like the
single-allele system in *Takifugu rubripes* (Kamiya et al. 2012). Regardless, identifying these
systems usually involves scans for intersex Fst, SBG overaccumulation, or a combination.

Despite relatively high genome-wide intersex Fst in NO, distinct elevated Fst regions and
regions with accumulation of SBGs were absent. Notably, several genes showed extreme Fst
values, but they were scattered throughout the genome lacking known SD genes, suggesting
allele-specific expression. NO also possessed both male and female SSAs, which points
towards an ESD system in NO.
ST showed the lowest genetic differentiation between sexes with no distinct regions of elevated
divergence, lowest number of SBGs, and the lowest number of DEAs among studied species.
These patterns hint towards an Environmental Sex Determination (ESD) system, as reinforced
by similar patterns found in *Syngnathus scovelli*. However, the presence of dmrt1, a well-known
teleost SD gene, with an Fst of 0.4 complicates the picture. This Fst suggests that one sex is
homozygous while the other is heterozygous, potentially pointing towards an XY sex
determination system with dmrt1 as the master gene. Determining ST's SD system requires
further investigation.

Adding another layer to the complexity of SD mechanisms, recent research on *Ictalurus*

punctatus (channel catfish) (Bao et al. 2019) revealed a novel mechanism where sex is

determined by the expression of a male-specific isoform of the breast cancer anti-resistance 1

(BCAR1) gene during early developmental stages. This discovery opens up the possibility that

molecular mechanisms like DEA and ASE, that traditionally not considered in SD studies, might

be at play in species with putative ESD systems, such as NO. Notably, while the overall genetic

divergence between sexes is high in NO, relatively few SBGs show elevated Fst. This

discrepancy suggests that sex-biased allele expression might be the primary mechanism for

resolving ongoing SA in this species.

4.2.2 Intraspecies sex chromosome polymorphism in HE and multi-chromosome SA

supergene

Our study explored signatures of ongoing sexually antagonistic (SA) selection in three

syngnathid fish species. In HE, we discovered a link between SA and sex determination (SD).

Analysis revealed a single chromosome-level scaffold (*chr14*) accumulating the majority of

SBGs with elevated Fst, DEAs, and SSAs. Fst values near 0.5 or higher suggest an XY system.

This pattern underscores that in HE, ongoing SA relationships might be the driving force behind

the increased genetic divergence between sexes, ultimately leading to the formation of a

homomorphic sex chromosome.

Comparing the other two species (ST and NO) to HE, we observed no similar accumulation of

SA signatures on specific scaffolds or regions. This suggests that in ST and NO, SA

relationships and SD system formation are likely uncoupled.

Furthermore, as discussed in the end of the section 3.1.1, our findings suggest that sex roles

and mate competition, not male pregnancy, are the main drivers of SA in all three species. In

HE, this competition might further influence the SD system by promoting sex chromosome

formation.

A sex-linked chromosome was recently already described in HE (Long et al. 2022). The sex-

linked chromosome that we identified, however, was not homologous to the homomorphic sex-

linked chromosomes described in *Hippocampus erectus* (chr11 or homologue of *Hippocampus*

abdominalis chr4) and *Hippocampus abdominalis* (chr6) (HA) with low sex-specific sequence

divergence suggesting a recent evolutionary trajectory (Long et al. 2022). The identified

discrepancies in HE sex chromosomes may have either appeared due to different

methodological approaches taken in the two studies, or alternatively indicate real biological

differences of the two HE breeds.

Methodological differences in the two studies become evident in several aspects i) the type of

sequence data utilized – this study employed RNA data for identifying the sex-linked

chromosome, whereas Long et al. used DNA; ii) the genome used for mapping the generated

sequencing data – this study used the chromosome-level HE genome, whereas Long et al. used

the genome of the more distant HA; and iii) the tools and metrics utilized for identifying sex-

linkage – we focused on overaccumulation of SA signature on particular genomic region, while

the other study relied exclusively on Fst and GWAS analyses. The bioinformatic tools that were

used to conduct the analysis were different, which could contribute to differing results.

To enhance the comparability of the two datasets and potentially mitigate the impact of

methodological differences as explanations for the disparities in HE sex chromosome

identification, we re-analyzed the data from HE males and females as presented in Long et al.,

employing the chromosome-level assembly of HE from (Li et al. 2021) with kmerGWAS. Using

this approach, we have discovered compelling evidence supporting the convergent evolution of

two distinct sex-linked chromosome systems in the two breeds of HE, facilitating the resolution

of SA. Previously, differences in sex-linked regions between captive breeds and wild-types have

been observed in various species, such as zebrafish, Siamese fighting fish and cichlids (Wilson

et al. 2014; Panthum et al. 2022; Wang et al. 2022; Schartl et al. 2023; Lichilín et al. 2023). For

instance, the Z chromosome of the zebrafish wildtype ZW sex determination system was

hypothesized to have been lost during domestication (Wilson et al. 2014). The zebrafish WW

individuals develop either as females or neomales, however, their SD mechanism remains as

yet unknown. Additionally, in Siamese fighting fish, some breeds possess a monogenic SD

system with dmrt1 as a master SD gene, while others exhibit polygenic SD mechanisms

(Panthum et al. 2022; Wang et al. 2022).

The convergent evolution of sex chromosome in two distinct lineages of aquaculture-origin HEs

suggests an even younger evolutionary age of the sex-linked chromosomes, as previously

proposed by Long et al. (Long et al. 2022). The HEs used in Long et al. originated from a

cultured population in Fujian (China), with a seed population of 2,000 – 3,000 wild-caught

individuals. In contrast, HEs used in this study were obtained from a German aquarium breeder,

with unknown but likely smaller population size. The emergence of the two sex chromosomes

may have occurred in a similar scenario as reported for captive zebrafish, where domestication

selected for two sex-linked chromosomes in two separate captive populations. However, the

incomplete linkage in both sex chromosomes, as indicated by the Fst in our case (less than 0.4)

and by the GWAS in Long et al. (Long et al. 2022) (male homozygotes were found within a sex-

linked region), raises questions about the existence of a master SD locus, while the

recombination landscape could predate a turnover event as proposed by (Long et al. 2022).

Genes with sex-biased expression patterns often exhibit elevated intersex Fst values (Cheng

and Kirkpatrick 2016; Lichilín et al. 2021; Wright et al. 2018). In humans and flies, this

relationship follows a "Twin Peaks" pattern. Genes with intermediate bias in expression

demonstrate the highest Fst values, indicating ongoing sexual selection, whereas significant

bias in expression aids in resolving sexual conflict (Cheng and Kirkpatrick 2016). However, in

cichlids and guppies, the intersex Fst increases with bias in expression, thus, the most extreme

sex-biased genes display the highest intersex Fst values (Lichilín et al. 2021; Wright et al.

2018). These data indicate that a bias in expression alone may not be sufficient to fully resolve

sexual conflict, independent of the underlying mechanism.

Despite its young age, we observed an enrichment of SBGs on the HE sex-linked chromosome

compared to the rest of the genome. This, combined with an elevated intersex Fst pattern, as

revealed by kmerGWAS or GWAS as in Long's study, suggests the presence of a true sex

chromosome. The elevated intersex Fst values likely result from ancestral low recombination

rates, and Long et al. proposed several promising SD candidate genes in this context (Long et

al. 2022).

In this study we were unable to identify SD candidates related to the canonical vertebrate

gonadal differentiation pathway on our sex-linked chromosome. Instead, we observed extreme

intersex Fst (0.9) and extreme bias in expression for amh, a well-known sex determination

gene. This observation suggests a transient state during a turnover event, as recently

suggested for multi-locus SD systems (Schartl et al. 2023).

In a plausible evolutionary scenario, the chromosome containing the amh gene as an SD locus

was originally the only sex-linked chromosome in HE. Amh falls within the canonical vertebrate

gonadal differentiation pathway and genes related to amh, like anti-Müllerian hormone type II

receptor (amhr2), that was identified as SD loci in other syngnathid species (Qu et al. 2021).

Domestication in closed aquarium systems, devoid of natural predators and subjected to

selection distinct from those in the natural population, likely amplified the effects of SAS. This

led to an enrichment of the now identified sex-linked chromosome for SBGs with elevated Fst

values. Although unusual for such a young sex chromosome, selection for elevated Fst values

was likely facilitated by ongoing intralocus sexual conflict that caused bias in expression

promoting genetic linkage. Fixation must have been accelerated by captive breeding with a

limited number of mates resulting in an overall reduced recombination landscape within the

genome that facilitated linkage of chr14 (or chr11 in the other HE breed). While a new

chromosome became sex-linked, the master SD locus (amh) remained the same, creating a

multi-locus SD system. This could be similar to recently described multi-chromosome

supergene in ants that contains socially antagonistic alleles (Scarparo et al. 2023), however, in

case of HE it rather contains sexually antagonistic alleles.

Several questions remain unanswered, including the speed of such transformation, the exact

mechanisms for recombination slowdown, the relationships between bias in expression and

elevated Fst, and the role of SAS in this process. However, if confirmed, seahorse populations

of the species *Hippocampus erectus* offer an intriguing system to study the effects of SAS on

genome evolution.

Conclusion

In the complex interactions between sexual antagonism (SA) and sex determination (SD)

systems across three syngnathid species, predominantly sex roles and mate competition, rather

than male pregnancy, drive SAS. In species with the most derived form of male pregnancy,

such as *Hippocampus erectus*, this trait may amplify SAS, potentially influencing the formation

of sex chromosomes. However, the observation that the sex-linked chromosome in these

species is not specifically enriched in pouch-related genes suggests that, while male pregnancy

might contribute to the amplification of SA, it is not the sole driver of the evolution of sex

chromosomes. Intriguingly, non-coding RNAs and allele-specific expression emerge as crucial

mediators of SA, highlighting these processes as vital for future SA research. Our data suggest

a mix of environmental and genetic sex determination across the species studied, with evidence

of co-evolution between SA pressures and SD mechanisms in *Hippocampus erectus*. Notably,

in HE there appears to be convergent evolution of sex chromosomes within different

populations, indicating the dynamic nature of SD system evolution under SAS. This study

significantly advances our understanding of how reproductive strategies and selection influence

genetic divergence and the evolution of SD systems in syngnathids and beyond.

1013

1014

Ethics statement

Work was carried out in accordance with German animal welfare law and with the ethical approval given by the Ministerium für Energiewende, Landwirtschaft, Umwelt, Natur und Ditgitalisierung (MELUND) Schleswig-Holstein (permit no. V242-57983/2018). No wild endangered species were used in this investigation.

Authors' contributions

A.D., O.R. and A.B. planned the project. J.P. collected and processed samples. A.D. analyzed the data and interpreted the results with input from O.R. and A.B.. A.D. & O.R. wrote the manuscript with input from all coauthors.

Funding

This project was supported by funding from the German Research Foundation (RO-4628/4-2) and from the European Research Council (ERC) under the European Union´s Horizon research and innovation program (MALEPREG: eu-repo/grantAgreement/EC/H2020/755659) to OR.

Data Access

All sequencing data generated in this project can be found under BioProject ID PRJNA947442.

References

Ahn Y, Sims C, Murray MJ, Kuhlmann PK, Fuentes-Antrás J, Weatherbee SD, Krumlauf R. 2017. Multiple modes of Lrp4 function in modulation of Wnt/β-catenin signaling during tooth development. *Development* [Internet]:dev.150680. Available from: https://journals.biologists.com/dev/article/doi/10.1242/dev.150680/264498/Multiple-modes-of-Lrp4-function-in-modulation-of

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J. Mol. Biol.* [Internet] 215:403–410. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0022283605803602

Armitage LH, Wallet MA, Mathews CE. 2021. Influence of PTPN22 Allotypes on Innate and Adaptive Immune Function in Health and Disease. *Front. Immunol.* [Internet] 12:636618. Available from: https://www.frontiersin.org/articles/10.3389/fimmu.2021.636618/full

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. 2000. Gene Ontology: tool for the unification of biology. *Nat. Genet.* [Internet] 25:25–29. Available from: https://www.nature.com/articles/ng0500_25

Bachtrog D, Mank JE, Peichel CL, Kirkpatrick M, Otto SP, Ashman T-L, Hahn MW, Kitano J, Mayrose I, Ming R, et al. 2014. Sex Determination: Why So Many Ways of Doing It? *PLoS Biol.* [Internet] 12:e1001899. Available from: https://dx.plos.org/10.1371/journal.pbio.1001899

Barnett DW, Garrison EK, Quinlan AR, Strömberg MP, Marth GT. 2011. BamTools: a C++ API and toolkit for analyzing and managing BAM files. *Bioinformatics* [Internet] 27:1691–1692. Available from: https://academic.oup.com/bioinformatics/article/27/12/1691/255399

Bar-Sagi D, Hall A. 2000. Ras and Rho GTPases. *Cell* [Internet] 103:227–238. Available from: https://linkinghub.elsevier.com/retrieve/pii/S009286740000115X

Benoit-Pilven C, Marchet C, Chautard E, Lima L, Lambert M-P, Sacomoto G, Rey A, Cologne A, Terrone S, Dulaurier L, et al. 2018. Complementarity of assembly-first and mapping-first approaches for alternative splicing annotation and differential analysis from RNAseq data. *Sci. Rep.* [Internet] 8:4307. Available from: https://www.nature.com/articles/s41598-018-21770-7

Berglund A, Rosenqvist G. 1993. Selective males and ardent females in pipefishes. *Behav. Ecol. Sociobiol.* [Internet] 32. Available from: http://link.springer.com/10.1007/BF00183788

Bertho S, Herpin A, Branthonne A, Jouanno E, Yano A, Nicol B, Muller T, Pannetier M, Pailhoux E, Miwa M, et al. 2018. The unusual rainbow trout sex determination gene hijacked the canonical vertebrate gonadal differentiation pathway. *Proc. Natl. Acad. Sci.* [Internet] 115:12781–12786. Available from: https://pnas.org/doi/full/10.1073/pnas.1803826115

Betancur-R R, Wiley EO, Arratia G, Acero A, Bailly N, Miya M, Lecointre G, Ortí G. 2017. Phylogenetic classification of bony fishes. *BMC Evol. Biol.* [Internet] 17:162. Available from: https://bmcevolbiol.biomedcentral.com/articles/10.1186/s12862-017-0958-3

Bizuayehu TT, Lanes CF, Furmanek T, Karlsen BO, Fernandes JM, Johansen SD, Babiak I. 2012. Differential expression patterns of conserved miRNAs and isomiRs during Atlantic halibut development. *BMC Genomics* [Internet] 13:11. Available from: https://bmcgenomics.biomedcentral.com/articles/10.1186/1471-2164-13-11

Bizzarri C, Benevento D, Terzi C, Huebner A, Cappa M. 2013. Triple A (Allgrove) syndrome: an unusual association with syringomyelia. *Ital. J. Pediatr.* [Internet] 39:39. Available from: http://ijponline.biomedcentral.com/articles/10.1186/1824-7288-39-39

Boratyn GM, Schäffer AA, Agarwala R, Altschul SF, Lipman DJ, Madden TL. 2012. Domain enhanced lookup time accelerated BLAST. *Biol. Direct* [Internet] 7:12. Available from: http://biologydirect.biomedcentral.com/articles/10.1186/1745-6150-7-12

Bosch TCG, Adamska M, Augustin R, Domazet☐Loso T, Foret S, Fraune S, Funayama N, Grasis J, Hamada M, Hatta M, et al. 2014. How do environmental factors influence life cycles and development? An experimental framework for early☐diverging metazoans. *BioEssays* [Internet] 36:1185–1194. Available from: https://onlinelibrary.wiley.com/doi/10.1002/bies.201400065

Boueid M-J, Mikdache A, Lesport E, Degerny C, Tawk M. 2020. Rho GTPases Signaling in Zebrafish Development and Disease. *Cells* [Internet] 9:2634. Available from: https://www.mdpi.com/2073-4409/9/12/2634

Bradford YM, Van Slyke CE, Ruzicka L, Singer A, Eagle A, Fashena D, Howe DG, Frazer K, Martin R, Paddock H, et al. 2022. Zebrafish information network, the knowledgebase for *Danio rerio* research.Wood V, editor. *Genetics* [Internet] 220:iyac016. Available from: https://academic.oup.com/genetics/article/doi/10.1093/genetics/iyac016/6528852

Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq quantification. *Nat. Biotechnol.* [Internet] 34:525–527. Available from: https://www.nature.com/articles/nbt.3519

Broad Institute. 2019. Picard toolkit. *Broad Inst. GitHub Repos.* [Internet]. Available from: https://broadinstitute.github.io/picard/

Brooks BP, Kleta R, Caruso RC, Stuart C, Ludlow J, Stratakis CA. 2004. Triple-A syndrome with prominent ophthalmic features and a novel mutation in the AAASgene: a case report. *BMC Ophthalmol.* [Internet] 4:7. Available from: http://bmcophthalmol.biomedcentral.com/articles/10.1186/1471-2415-4-7

Brown CJ, Ballabio A, Rupert JL, Lafreniere RG, Grompe M, Tonlorenzi R, Willard HF. 1991. A gene from the region of the human X inactivation centre is expressed exclusively from the inactive X chromosome. *Nature* [Internet] 349:38–44. Available from: https://www.nature.com/articles/349038a0

Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M. 2021. BRAKER2: automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein

database. *NAR Genomics Bioinforma.* [Internet] 3:lqaa108. Available from: https://academic.oup.com/nargab/article/doi/10.1093/nargab/lqaa108/6066535

Brůna T, Lomsadze A, Borodovsky M. 2020. GeneMark-EP+: eukaryotic gene prediction with self-training in the space of genes and proteins. *NAR Genomics Bioinforma.* [Internet] 2:lqaa026. Available from: https://academic.oup.com/nargab/article/doi/10.1093/nargab/lqaa026/5836691

Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using DIAMOND. *Nat. Methods* [Internet] 12:59–60. Available from: https://www.nature.com/articles/nmeth.3176

Burrage LC, Reynolds JJ, Baratang NV, Phillips JB, Wegner J, McFarquhar A, Higgs MR, Christiansen AE, Lanza DG, Seavitt JR, et al. 2019. Bi-allelic Variants in TONSL Cause SPONASTRIME Dysplasia and a Spectrum of Skeletal Dysplasia Phenotypes. *Am. J. Hum. Genet.* [Internet] 104:422–438. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0002929719300072

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* [Internet] 10:421. Available from: https://bmcbioinformatics.biomedcentral.com/articles/10.1186/1471-2105-10-421

Chang M-K, Kramer I, Huber T, Kinzel B, Guth-Gundel S, Leupin O, Kneissel M. 2014. Disruption of Lrp4 function by genetic deletion or pharmacological blockade increases bone mass and serum sclerostin levels. *Proc. Natl. Acad. Sci.* [Internet] 111. Available from: https://pnas.org/doi/full/10.1073/pnas.1413828111

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* [Internet] 34:i884–i890. Available from: https://academic.oup.com/bioinformatics/article/34/17/i884/5093234

Cheng C, Kirkpatrick M. 2016. Sex-Specific Selection and Sex-Biased Gene Expression in Humans and Flies.Nachman MW, editor. *PLOS Genet.* [Internet] 12:e1006170. Available from: https://dx.plos.org/10.1371/journal.pgen.1006170

Choi JH, Zhong X, McAlpine W, Liao T-C, Zhang D, Fang B, Russell J, Ludwig S, Nair-Gill E, Zhang Z, et al. 2019. LMBR1L regulates lymphopoiesis through Wnt/β-catenin signaling. *Science* [Internet] 364:eaau0812. Available from: https://www.science.org/doi/10.1126/science.aau0812

Chou W-C, Jha S, Linhoff MW, Ting JP-Y. 2023. The NLR gene family: from discovery to present day. *Nat. Rev. Immunol.* [Internet]. Available from: https://www.nature.com/articles/s41577-023-00849-x

Ciaudo C, Bourdet A, Cohen-Tannoudji M, Dietz HC, Rougeulle C, Avner P. 2006. Nuclear mRNA Degradation Pathway(s) Are Implicated in Xist Regulation and X Chromosome Inactivation.Reik W, editor. *PLoS Genet.* [Internet] 2:e94. Available from: https://dx.plos.org/10.1371/journal.pgen.0020094

Dagilis AJ, Sardell JM, Josephson MP, Su Y, Kirkpatrick M, Peichel CL. 2022. Searching for signatures of sexually antagonistic selection on stickleback sex chromosomes. *Philos. Trans. R. Soc. B Biol. Sci.* [Internet] 377:20210205. Available from: https://royalsocietypublishing.org/doi/10.1098/rstb.2021.0205

Dainat J, Hereñú D, Dr. K. D. Murray, Davis E, Crouch K, LucileSol, Agostinho N, Pascal-Git, Zollman Z, Tayyrov. 2023. NBISweden/AGAT: AGAT-v1.2.0. Available from: https://zenodo.org/record/3552717

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST, et al. 2011. The variant call format and VCFtools. *Bioinformatics* [Internet] 27:2156–2158. Available from: https://academic.oup.com/bioinformatics/article/27/15/2156/402296

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* [Internet] 29:15–21. Available from: https://academic.oup.com/bioinformatics/article/29/1/15/272537

El Taher A, Ronco F, Matschiner M, Salzburger W, Böhne A. 2021. Dynamics of sex chromosome evolution in a rapid radiation of cichlid fishes. *Sci. Adv.* [Internet] 7:eabe8215. Available from: https://www.science.org/doi/10.1126/sciadv.abe8215

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol.* [Internet] 20:238. Available from: https://genomebiology.biomedcentral.com/articles/10.1186/s13059-019-1832-y

Fernandez-Capetillo O, Mahadevaiah SK, Celeste A, Romanienko PJ, Camerini-Otero RD, Bonner WM, Manova K, Burgoyne P, Nussenzweig A. 2003. H2AX Is Required for Chromatin Remodeling and Inactivation of Sex Chromosomes in Male Mouse Meiosis. *Dev. Cell* [Internet] 4:497–508. Available from: https://linkinghub.elsevier.com/retrieve/pii/S1534580703000935

Foster SJ, Vincent ACJ. 2004. Life history and ecology of seahorses: implications for conservation and management. *J. Fish Biol.* [Internet] 65:1–61. Available from: https://onlinelibrary.wiley.com/doi/10.1111/j.0022-1112.2004.00429.x

Fousteri G, Liossis S-NC, Battaglia M. 2013. Roles of the protein tyrosine phosphatase PTPN22 in immunity and autoimmunity. *Clin. Immunol.* [Internet] 149:556–565. Available from: https://linkinghub.elsevier.com/retrieve/pii/S1521661613002647

Furman BLS, Metzger DCH, Darolti I, Wright AE, Sandkam BA, Almeida P, Shu JJ, Mank JE. 2020. Sex Chromosome Evolution: So Many Exceptions to the Rules.Fraser B, editor. *Genome Biol. Evol.* [Internet] 12:750–763. Available from: https://academic.oup.com/gbe/article/12/6/750/5823304

Gabriel L, Hoff KJ, Brůna T, Borodovsky M, Stanke M. 2021. TSEBRA: transcript selector for BRAKER. *BMC Bioinformatics* [Internet] 22:566. Available from: https://bmcbioinformatics.biomedcentral.com/articles/10.1186/s12859-021-04482-0

Gammerdinger WJ, Toups MA, Vicoso B. 2020. Disagreement in F_{ST} estimators: A case study from sex chromosomes. *Mol. Ecol. Resour.* [Internet] 20:1517–1525. Available from: https://onlinelibrary.wiley.com/doi/10.1111/1755-0998.13210

Gineau L, Cognet C, Kara N, Lach FP, Dunne J, Veturi U, Picard C, Trouillet C, Eidenschenk C, Aoufouchi S, et al. 2012. Partial MCM4 deficiency in patients with growth retardation, adrenal insufficiency, and natural killer cell deficiency. *J. Clin. Invest.* [Internet] 122:821–832. Available from: http://www.jci.org/articles/view/61014

Goncalves IB, Ahnesjö I, Kvarnemo C. 2015. Embryo oxygenation in pipefish brood pouches: novel insights. *J. Exp. Biol.* [Internet] 218:1639–1646. Available from: https://journals.biologists.com/jeb/article/218/11/1639/13760/Embryo-oxygenation-in-pipefish-brood-pouches-novel

Gotoh O. 2008. A space-efficient and accurate method for mapping and aligning cDNA sequences onto genomic sequence. *Nucleic Acids Res.* [Internet] 36:2630–2638. Available from: https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gkn105

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q, et al. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat. Biotechnol.* [Internet] 29:644–652. Available from: https://www.nature.com/articles/nbt.1883

Grayson P, Wright A, Garroway CJ, Docker MF. 2022. SexFindR: A computational workflow to identify young and old sex chromosomes. Evolutionary Biology Available from: http://biorxiv.org/lookup/doi/10.1101/2022.02.21.481346

Greenhough LA, Liang C-C, Belan O, Kunzelmann S, Maslen S, Rodrigo-Brenni MC, Anand R, Skehel M, Boulton SJ, West SC. 2023. Structure and function of the RAD51B–RAD51C–RAD51D–XRCC2 tumour suppressor. *Nature* [Internet] 619:650–657. Available from: https://www.nature.com/articles/s41586-023-06179-1

Gregory TR. 2009. Understanding Natural Selection: Essential Concepts and Common Misconceptions. *Evol. Educ. Outreach* [Internet] 2:156–175. Available from: https://evolution-outreach.biomedcentral.com/articles/10.1007/s12052-009-0128-1

Gremme G, Steinbiss S, Kurtz S. 2013. GenomeTools: A Comprehensive Software Library for Efficient Processing of Structured Genome Annotations. *IEEE/ACM Trans. Comput. Biol. Bioinform.* [Internet] 10:645–656. Available from: http://ieeexplore.ieee.org/document/6529082/

Hamdoun A, Epel D. 2007. Embryo stability and vulnerability in an always changing world. *Proc. Natl. Acad. Sci.* [Internet] 104:1745–1750. Available from: https://pnas.org/doi/full/10.1073/pnas.0610108104

Handschug K. 2001. Triple A syndrome is caused by mutations in AAAS, a new WD-repeat protein gene. *Hum. Mol. Genet.* [Internet] 10:283–290. Available from: https://academic.oup.com/hmg/article-lookup/doi/10.1093/hmg/10.3.283

Hansen CL, Pelegri F. 2021. Primordial Germ Cell Specification in Vertebrate Embryos: Phylogenetic Distribution and Conserved Molecular Features of Preformation and Induction. *Front. Cell Dev. Biol.* [Internet] 9:730332. Available from: https://www.frontiersin.org/articles/10.3389/fcell.2021.730332/full

He L, Long X, Qi J, Wang Z, Huang Z, Wu S, Zhang X, Luo H, Chen X, Lin J, et al. 2022. Genome and gene evolution of seahorse species revealed by the chromosome□level genome of *Hippocampus abdominalis*. *Mol. Ecol. Resour.* [Internet] 22:1465–1477. Available from: https://onlinelibrary.wiley.com/doi/10.1111/1755-0998.13541

Hickford DE, Frankenberg S, Pask AJ, Shaw G, Renfree MB. 2011. DDX4 (VASA) Is Conserved in Germ Cell Development in Marsupials and Monotremes1. *Biol. Reprod.* [Internet] 85:733–743. Available from: https://academic.oup.com/biolreprod/article-lookup/doi/10.1095/biolreprod.111.091629

Hodge RG, Ridley AJ. 2016. Regulating Rho GTPases and their regulators. *Nat. Rev. Mol. Cell Biol.* [Internet] 17:496–510. Available from: https://www.nature.com/articles/nrm.2016.67

Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. 2016. BRAKER1: Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS. *Bioinformatics* [Internet] 32:767–769. Available from: https://academic.oup.com/bioinformatics/article/32/5/767/1744611

Hoff KJ, Lomsadze A, Borodovsky M, Stanke M. 2019. Whole-Genome Annotation with BRAKER. In: Kollmar M, editor. Gene Prediction. Vol. 1962. Methods in Molecular Biology. New York, NY: Springer New York. p. 65–95. Available from: http://link.springer.com/10.1007/978-1-4939-9173-0_5

Iversen LL, Svensson EI, Christensen ST, Bergsten J, Sand-Jensen K. 2019. Sexual conflict and intrasexual polymorphism promote assortative mating and halt population differentiation. *Proc. R. Soc. B Biol. Sci.* [Internet] 286:20190251. Available from: https://royalsocietypublishing.org/doi/10.1098/rspb.2019.0251

Iwata H, Gotoh O. 2012. Benchmarking spliced alignment programs including Spaln2, an extended version of Spaln that incorporates additional species-specific features. *Nucleic Acids Res.* [Internet] 40:e161–e161. Available from: https://academic.oup.com/nar/article/40/20/e161/2414522

Jackman SD, Vandervalk BP, Mohamadi H, Chu J, Yeo S, Hammond SA, Jahesh G, Khan H, Coombe L, Warren RL, et al. 2017. ABySS 2.0: resource-efficient assembly of large genomes using a Bloom filter. *Genome Res.* [Internet] 27:768–777. Available from: http://genome.cshlp.org/lookup/doi/10.1101/gr.214346.116

Jaffe AB, Hall A. 2005. RHO GTPASES: Biochemistry and Biology. *Annu. Rev. Cell Dev. Biol.* [Internet] 21:247–269. Available from: https://www.annualreviews.org/doi/10.1146/annurev.cellbio.21.020604.150721

Jamin SP, Petit FG, Kervarrec C, Smagulova F, Illner D, Scherthan H, Primig M. 2017. EXOSC10/Rrp6 is post-translationally regulated in male germ cells and controls the onset of spermatogenesis. *Sci. Rep.* [Internet] 7:15065. Available from: https://www.nature.com/articles/s41598-017-14643-y

Jing J, Wu J, Liu W, Xiong S, Ma W, Zhang J, Wang W, Gui J-F, Mei J. 2014. Sex-Biased miRNAs in Gonad and Their Potential Roles for Testis Development in Yellow Catfish.Qu L-H, editor. *PLoS ONE* [Internet] 9:e107946. Available from: https://dx.plos.org/10.1371/journal.pone.0107946

Jones JDG, Vance RE, Dangl JL. 2016. Intracellular innate immune surveillance devices in plants and animals. *Science* [Internet] 354:aaf6395. Available from: https://www.science.org/doi/10.1126/science.aaf6395

Juanchich A, Bardou P, Rué O, Gabillard J-C, Gaspin C, Bobe J, Guiguen Y. 2016. Characterization of an extensive rainbow trout miRNA transcriptome by next generation sequencing. *BMC Genomics* [Internet] 17:164. Available from: http://www.biomedcentral.com/1471-2164/17/164

Juanchich A, Le Cam A, Montfort J, Guiguen Y, Bobe J. 2013. Identification of Differentially Expressed miRNAs and Their Potential Targets During Fish Ovarian Development1. *Biol. Reprod.* [Internet] 88. Available from: https://academic.oup.com/biolreprod/article-lookup/doi/10.1095/biolreprod.112.105361

Kamiya T, Kai W, Tasumi S, Oka A, Matsunaga T, Mizuno N, Fujita M, Suetake H, Suzuki S, Hosoya S, et al. 2012. A Trans-Species Missense SNP in Amhr2 Is Associated with Sex Determination in the Tiger Pufferfish, Takifugu rubripes (Fugu).Peichel CL, editor. *PLoS Genet.* [Internet] 8:e1002798. Available from: https://dx.plos.org/10.1371/journal.pgen.1002798

Kent WJ. 2002. BLAT —The BLAST -Like Alignment Tool. *Genome Res.* [Internet] 12:656–664. Available from: http://genome.cshlp.org/lookup/doi/10.1101/gr.229202

Khatpe AS, Dirks R, Bhat-Nakshatri P, Mang H, Batic K, Swiezy S, Olson J, Rao X, Wang Y, Tanaka H, et al. 2023. TONSL Is an Immortalizing Oncogene and a Therapeutic Target in Breast Cancer. *Cancer Res.* [Internet] 83:1345–1360. Available from: https://aacrjournals.org/cancerres/article/83/8/1345/725108/TONSL-Is-an-Immortalizing-Oncogene-and-a

Kibar Z, Bosoi CM, Kooistra M, Salem S, Finnell RH, De Marco P, Merello E, Bassuk AG, Capra V, Gros P. 2009. Novel mutations in *VANGL1* in neural tube defects. *Hum. Mutat.* [Internet] 30:E706–E715. Available from: https://onlinelibrary.wiley.com/doi/10.1002/humu.21026

Kobayashi Y, Nagahama Y, Nakamura M. 2013. Diversity and Plasticity of Sex Determination and Differentiation in Fishes. *Sex. Dev.* [Internet] 7:115–125. Available from: https://www.karger.com/Article/FullText/342009

Kriventseva EV, Kuznetsov D, Tegenfeldt F, Manni M, Dias R, Simão FA, Zdobnov EM. 2019. OrthoDB v10: sampling the diversity of animal, plant, fungal, protist, bacterial and viral genomes for evolutionary and functional annotations of orthologs. *Nucleic Acids Res.* [Internet] 47:D807–D811. Available from: https://academic.oup.com/nar/article/47/D1/D807/5160989

Laland KN, Odling-Smee FJ, Feldman MW. 1999. Evolutionary consequences of niche construction and their implications for ecology. *Proc. Natl. Acad. Sci.* [Internet] 96:10242–10247. Available from: https://pnas.org/doi/full/10.1073/pnas.96.18.10242

Lasko P. 2013. The DEAD-box helicase Vasa: Evidence for a multiplicity of functions in RNA processes and developmental biology. *Biochim. Biophys. Acta BBA - Gene Regul. Mech.* [Internet] 1829:810–816. Available from: https://linkinghub.elsevier.com/retrieve/pii/S1874939913000680

Lau K, Lai KP, Bao JYJ, Zhang N, Tse A, Tong A, Li JW, Lok S, Kong RYC, Lui WY, et al. 2014. Identification and Expression Profiling of MicroRNAs in the Brain, Liver and Gonads of Marine Medaka (Oryzias melastigma) and in Response to Hypoxia.Antoniewski C, editor. *PLoS ONE* [Internet] 9:e110698. Available from: https://dx.plos.org/10.1371/journal.pone.0110698

Levins R, Lewontin RC. 1985. The dialectical biologist. Cambridge (Mass.) London: Harvard university press

Li C, Olave M, Hou Y, Qin G, Schneider RF, Gao Z, Tu X, Wang X, Qi F, Nater A, et al. 2021. Genome sequences reveal global dispersal routes and suggest convergent genetic adaptations in seahorse evolution. *Nat. Commun.* [Internet] 12:1094. Available from: https://www.nature.com/articles/s41467-021-21379-x

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* [Internet] 25:2078–2079. Available from: https://academic.oup.com/bioinformatics/article/25/16/2078/204688

Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* [Internet] 22:1658–1659. Available from: https://academic.oup.com/bioinformatics/article/22/13/1658/194225

Lichilín N, El Taher A, Böhne A. 2021. Sex-biased gene expression and recent sex chromosome turnover. *Philos. Trans. R. Soc. B Biol. Sci.* [Internet] 376:20200107. Available from: https://royalsocietypublishing.org/doi/10.1098/rstb.2020.0107

Lin Z, Kong H, Nei M, Ma H. 2006. Origins and evolution of the *recA / RAD51* gene family: Evidence for ancient gene duplication and endosymbiotic gene transfer. *Proc. Natl. Acad. Sci.* [Internet] 103:10328–10333. Available from: https://pnas.org/doi/full/10.1073/pnas.0604232103

Liu D, Sonalkar J, Prasanth SG. 2023. ORChestra coordinates the replication and repair music. *BioEssays* [Internet] 45:2200229. Available from: https://onlinelibrary.wiley.com/doi/10.1002/bies.202200229

Lomsadze A. 2005. Gene identification in novel eukaryotic genomes by self-training algorithm. *Nucleic Acids Res.* [Internet] 33:6494–6506. Available from: https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gki937

Lomsadze A, Burns PD, Borodovsky M. 2014. Integration of mapped RNA-Seq reads into automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Res.* [Internet] 42:e119–e119. Available from: https://academic.oup.com/nar/article/42/15/e119/2434516

Long X, Charlesworth D, Qi J, Wu R, Chen M, Wang Z, Xu L, Fu H, Zhang X, Chen X, et al. 2023. Independent Evolution of Sex Chromosomes and Male Pregnancy–Related Genes in Two Seahorse Species.Wilson M, editor. *Mol. Biol. Evol.* [Internet] 40:msac279. Available from: https://academic.oup.com/mbe/article/doi/10.1093/molbev/msac279/6964685

Lopez-Maestre H, Brinza L, Marchet C, Kielbassa J, Bastien S, Boutigny M, Monnin D, Filali AE, Carareto CM, Vieira C, et al. 2016. SNP calling from RNA-seq data without a reference genome: identification, quantification, differential analysis and impact on the protein sequence.

Nucleic Acids Res. [Internet]:gkw655. Available from: https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gkw655

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* [Internet] 15:550. Available from: http://genomebiology.biomedcentral.com/articles/10.1186/s13059-014-0550-8

Lucotte EA, Laurent R, Heyer E, Ségurel L, Toupance B. 2016. Detection of Allelic Frequency Differences between the Sexes in Humans: A Signature of Sexually Antagonistic Selection. *Genome Biol. Evol.* [Internet] 8:1489–1500. Available from: https://academic.oup.com/gbe/article-lookup/doi/10.1093/gbe/evw090

Lui W-Y, Lee WM, Cheng CY. 2003. Rho GTPases and spermatogenesis. *Biochim. Biophys. Acta BBA - Mol. Cell Res.* [Internet] 1593:121–129. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0167488902003488

Mank JE. 2017. Population genetics of sexual conflict in the genomic era. *Nat. Rev. Genet.* [Internet] 18:721–730. Available from: https://www.nature.com/articles/nrg.2017.83

Mank JE, Hall DW, Kirkpatrick M, Avise JC. 2006. Sex chromosomes and male ornaments: a comparative evaluation in ray-finned fishes. *Proc. R. Soc. B Biol. Sci.* [Internet] 273:233–236. Available from: https://royalsocietypublishing.org/doi/10.1098/rspb.2005.3334

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes.Kelley J, editor. *Mol. Biol. Evol.* [Internet] 38:4647–4654. Available from: https://academic.oup.com/mbe/article/38/10/4647/6329644

Mentink RA, Rella L, Radaszkiewicz TW, Gybel T, Betist MC, Bryja V, Korswagen HC. 2018. The planar cell polarity protein VANG-1/Vangl negatively regulates Wnt/β-catenin signaling through a Dvl dependent mechanism.Chisholm AD, editor. *PLOS Genet.* [Internet] 14:e1007840. Available from: https://dx.plos.org/10.1371/journal.pgen.1007840

Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A. 2018. Versatile genome assembly evaluation with QUAST-LG. *Bioinformatics* [Internet] 34:i142–i150. Available from: https://academic.oup.com/bioinformatics/article/34/13/i142/5045727

Moczek AP. 2015. Re-evaluating the environment in developmental evolution. *Front. Ecol. Evol.* [Internet] 3. Available from: http://journal.frontiersin.org/Article/10.3389/fevo.2015.00007/abstract

Montani C, Gritti L, Beretta S, Verpelli C, Sala C. 2018. The synaptic and neuronal functions of the X□linked intellectual disability protein Interleukin□1 receptor accessory protein like 1 (IL 1

RAPL 1). *Dev. Neurobiol.* [Internet]:dneu.22657. Available from:
https://onlinelibrary.wiley.com/doi/10.1002/dneu.22657

Morrish B, Sinclair A. 2002. Vertebrate sex determination: many means to an end. *Reproduction* [Internet]:447–457. Available from: https://rep.bioscientifica.com/view/journals/rep/124/4/447.xml

Mosca TJ, Luginbuhl DJ, Wang IE, Luo L. 2017. Presynaptic LRP4 promotes synapse number and function of excitatory CNS neurons. *eLife* [Internet] 6:e27347. Available from:
https://elifesciences.org/articles/27347

Nagahama Y, Chakraborty T, Paul-Prasanth B, Ohta K, Nakamura M. 2021. Sex determination, gonadal sex differentiation, and plasticity in vertebrate species. *Physiol. Rev.* [Internet] 101:1237–1308. Available from: https://journals.physiology.org/doi/10.1152/physrev.00044.2019

Noyes C, Kitajima S, Li F, Suita Y, Miriyala S, Isaac S, Ahsan N, Knelson E, Vajdi A, Tani T, et al. 2023. The germline factor DDX4 contributes to the chemoresistance of small cell lung cancer cells. *Commun. Biol.* [Internet] 6:65. Available from: https://www.nature.com/articles/s42003-023-04444-7

Ou S, Jiang N. 2018. LTR_retriever: A Highly Accurate and Sensitive Program for Identification of Long Terminal Repeat Retrotransposons. *Plant Physiol.* [Internet] 176:1410–1422. Available from: https://academic.oup.com/plphys/article/176/2/1410-1422/6117145

Ou S, Su W, Liao Y, Chougule K, Agda JRA, Hellinga AJ, Lugo CSB, Elliott TA, Ware D, Peterson T, et al. 2019. Benchmarking transposable element annotation methods for creation of a streamlined, comprehensive pipeline. *Genome Biol.* [Internet] 20:275. Available from:
https://genomebiology.biomedcentral.com/articles/10.1186/s13059-019-1905-y

Pappert FA, Dubin A, Torres G, Roth O. 2023. Sex or sex role – unravelling the evolutionary route of sex-specific senescence. Evolutionary Biology Available from:
http://biorxiv.org/lookup/doi/10.1101/2023.05.02.539036

Parsch J, Ellegren H. 2013. The evolutionary causes and consequences of sex-biased gene expression. *Nat. Rev. Genet.* [Internet] 14:83–87. Available from:
https://www.nature.com/articles/nrg3376

Perry JA, Maddox AS. 2019. Uncovering the secret life of Rho GTPases. *eLife* [Internet] 8:e53276. Available from: https://elifesciences.org/articles/53276

Pfennig F, Standke A, Gutzeit HO. 2015. The role of Amh signaling in teleost fish – Multiple functions not restricted to the gonads. *Gen. Comp. Endocrinol.* [Internet] 223:87–107. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0016648015002695

Pinhal D, Bovolenta LA, Moxon S, Oliveira AC, Nachtigall PG, Acencio ML, Patton JG, Hilsdorf AWS, Lemke N, Martins C. 2018. Genome-wide microRNA screening in Nile tilapia reveals

pervasive isomiRs' transcription, sex-biased arm switching and increasing complexity of expression throughout development. *Sci. Rep.* [Internet] 8:8248. Available from: https://www.nature.com/articles/s41598-018-26607-x

Pinto DMS, Flaus A. 2010. Structure and Function of Histone H2AX. In: Nasheuer H-P, editor. Genome Stability and Human Diseases. Vol. 50. Subcellular Biochemistry. Dordrecht: Springer Netherlands. p. 55–78. Available from: http://link.springer.com/10.1007/978-90-481-3471-7_4

Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Van Der Auwera GA, Kling DE, Gauthier LD, Levy-Moonshine A, Roazen D, et al. 2017. Scaling accurate genetic variant discovery to tens of thousands of samples. Genomics Available from: http://biorxiv.org/lookup/doi/10.1101/201178

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, De Bakker PIW, Daly MJ, et al. 2007. PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *Am. J. Hum. Genet.* [Internet] 81:559–575. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0002929707613524

Qiu W, Zhu Y, Wu Y, Yuan C, Chen K, Li M. 2018. Identification and expression analysis of microRNAs in medaka gonads. *Gene* [Internet] 646:210–216. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0378111917311319

Ramanagoudr-Bhojappa R, Carrington B, Ramaswami M, Bishop K, Robbins GM, Jones M, Harper U, Frederickson SC, Kimble DC, Sood R, et al. 2018. Multiplexed CRISPR/Cas9-mediated knockout of 19 Fanconi anemia pathway genes in zebrafish revealed their roles in growth, sexual development and fertility.Barsh GS, editor. *PLOS Genet.* [Internet] 14:e1007821. Available from: https://dx.plos.org/10.1371/journal.pgen.1007821

Raz E. 2000. The function and regulation of vasa-like genes in germ-cell development. *Genome Biol.* [Internet] 1:reviews1017.1. Available from: http://genomebiology.biomedcentral.com/articles/10.1186/gb-2000-1-3-reviews1017

Reynolds A, McDearmid JR, Lachance S, Marco PD, Merello E, Capra V, Gros P, Drapeau P, Kibar Z. 2010. VANGL1 rare variants associated with neural tube defects affect convergent extension in zebrafish. *Mech. Dev.* [Internet] 127:385–392. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0925477309015007

Rice WR, Holland B. 1997. The enemies within: intergenomic conflict, interlocus contest evolution (ICE), and the intraspecific Red Queen. *Behav. Ecol. Sociobiol.* [Internet] 41:1–10. Available from: http://link.springer.com/10.1007/s002650050357

Roth O, Scharsack JP, Keller I, Reusch TBH. 2011. Bateman's principle and immunity in a sex-role reversed pipefish: Sexual immune dimorphism in pipefish. *J. Evol. Biol.* [Internet] 24:1410–1420. Available from: https://onlinelibrary.wiley.com/doi/10.1111/j.1420-9101.2011.02273.x

Roth O, Solbakken MH, Tørresen OK, Bayer T, Matschiner M, Baalsrud HT, Hoff SNK, Brieuc MSO, Haase D, Hanel R, et al. 2020. Evolution of male pregnancy associated with remodeling of canonical vertebrate immunity in seahorses and pipefishes. *Proc. Natl. Acad. Sci.* [Internet] 117:9431–9439. Available from: https://pnas.org/doi/full/10.1073/pnas.1916251117

Rowe L, Chenoweth SF, Agrawal AF. 2018. The Genomics of Sexual Conflict. *Am. Nat.* [Internet] 192:274–286. Available from: https://www.journals.uchicago.edu/doi/10.1086/698198

Sacomoto GA, Kielbassa J, Chikhi R, Uricaru R, Antoniou P, Sagot M-F, Peterlongo P, Lacroix V. 2012. KIS SPLICE: de-novo calling alternative splicing events from RNA-seq data. *BMC Bioinformatics* [Internet] 13:S5. Available from: https://bmcbioinformatics.biomedcentral.com/articles/10.1186/1471-2105-13-S6-S5

Sander Van Doorn G. 2014. Patterns and Mechanisms of Evolutionary Transitions between Genetic Sex-Determining Systems. *Cold Spring Harb. Perspect. Biol.* [Internet] 6:a017681–a017681. Available from: http://cshperspectives.cshlp.org/lookup/doi/10.1101/cshperspect.a017681

Scarparo G, Palanchon M, Brelsford A, Purcell J. 2023. Social antagonism facilitates supergene expansion in ants. *Curr. Biol.* [Internet] 33:5085-5095.e4. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0960982223014501

Schartl M, Georges A, Marshall Graves JA. 2023. Polygenic sex determination in vertebrates – is there any such thing? *Trends Genet.* [Internet] 39:242–250. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0168952522003110

Schwager EE, Meng Y, Extavour CG. 2015. vasa and piwi are required for mitotic integrity in early embryogenesis in the spider Parasteatoda tepidariorum. *Dev. Biol.* [Internet] 402:276–290. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0012160614004382

Shen C, Xiong W-C, Mei L. 2015. LRP4 in neuromuscular junction and bone development and diseases. *Bone* [Internet] 80:101–108. Available from: https://linkinghub.elsevier.com/retrieve/pii/S8756328215001830

Shen Z, Sathyan KM, Geng Y, Zheng R, Chakraborty A, Freeman B, Wang F, Prasanth KV, Prasanth SG. 2010. A WD-Repeat Protein Stabilizes ORC Binding to Chromatin. *Mol. Cell* [Internet] 40:99–111. Available from: https://linkinghub.elsevier.com/retrieve/pii/S1097276510007495

Sheu Y-J, Kinney JB, Lengronne A, Pasero P, Stillman B. 2014. Domain within the helicase subunit Mcm4 integrates multiple kinase signals to control DNA replication initiation and fork progression. *Proc. Natl. Acad. Sci.* [Internet] 111. Available from: https://pnas.org/doi/full/10.1073/pnas.1404063111

Shi J, Liang C. 2019. Generic Repeat Finder: A High-Sensitivity Tool for Genome-Wide De Novo Repeat Detection. *Plant Physiol.* [Internet] 180:1803–1815. Available from: https://academic.oup.com/plphys/article/180/4/1803-1815/6117697

Silva MSB, Giacobini P. 2021. New insights into anti-Müllerian hormone role in the hypothalamic–pituitary–gonadal axis and neuroendocrine development. *Cell. Mol. Life Sci.* [Internet] 78:1–16. Available from: https://link.springer.com/10.1007/s00018-020-03576-x

Simhadri S, Peterson S, Patel DS, Huo Y, Cai H, Bowman-Colin C, Miller S, Ludwig T, Ganesan S, Bhaumik M, et al. 2014. Male Fertility Defect Associated with Disrupted BRCA1-PALB2 Interaction in Mice. *J. Biol. Chem.* [Internet] 289:24617–24629. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0021925820319931

Skalkos ZMG, Van Dyke JU, Whittington CM. 2020. Paternal nutrient provisioning during male pregnancy in the seahorse Hippocampus abdominalis. *J. Comp. Physiol. B* [Internet] 190:547–556. Available from: https://link.springer.com/10.1007/s00360-020-01289-y

Smith SH, Hsiung K, Böhne A. 2023. Evaluating the role of sexual antagonism in the evolution of sex chromosomes: new data from fish. *Curr. Opin. Genet. Dev.* [Internet] 81:102078. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0959437X23000588

Smith-Unna R, Boursnell C, Patro R, Hibberd JM, Kelly S. 2016. TransRate: reference-free quality assessment of de novo transcriptome assemblies. *Genome Res.* [Internet] 26:1134–1144. Available from: http://genome.cshlp.org/lookup/doi/10.1101/gr.196469.115

Sogabe A, Ahnesjö I. 2011. The ovarian structure and mode of egg production in two polygamous pipefishes: a link to mating pattern. *J. Fish Biol.* [Internet] 78:1833–1846. Available from: https://onlinelibrary.wiley.com/doi/10.1111/j.1095-8649.2011.02973.x

Soneson C, Love MI, Robinson MD. 2015. Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. *F1000Research* [Internet] 4:1521. Available from: https://f1000research.com/articles/4-1521/v1

Stanke M, Diekhans M, Baertsch R, Haussler D. 2008. Using native and syntenically mapped cDNA alignments to improve *de novo* gene finding. *Bioinformatics* [Internet] 24:637–644. Available from: https://academic.oup.com/bioinformatics/article/24/5/637/202844

Stanke M, Schöffmann O, Morgenstern B, Waack S. 2006. Gene prediction in eukaryotes with a generalized hidden Markov model that uses hints from external sources. *BMC Bioinformatics*

[Internet] 7:62. Available from:
https://bmcbioinformatics.biomedcentral.com/articles/10.1186/1471-2105-7-62

Stearns FW. 2010. One Hundred Years of Pleiotropy: A Retrospective. *Genetics* [Internet]
186:767–773. Available from: https://academic.oup.com/genetics/article/186/3/767/6063631

Su W, Gu X, Peterson T. 2019. TIR-Learner, a New Ensemble Method for TIR Transposable
Element Annotation, Provides Evidence for Abundant New Transposable Elements in the Maize
Genome. *Mol. Plant* [Internet] 12:447–460. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S1674205219300620

Suszynska M, Ratajska M, Kozlowski P. 2020. BRIP1, RAD51C, and RAD51D mutations are
associated with high susceptibility to ovarian cancer: mutation prevalence and precise risk
estimates based on a pooled analysis of ~30,000 cases. *J. Ovarian Res.* [Internet] 13:50.
Available from: https://ovarianresearch.biomedcentral.com/articles/10.1186/s13048-020-00654-
3

Svensson EI. 2018. On Reciprocal Causation in the Evolutionary Process. *Evol. Biol.* [Internet]
45:1–14. Available from: http://link.springer.com/10.1007/s11692-017-9431-x

Svensson EI. 2019. Eco□evolutionary dynamics of sexual selection and sexual conflict.Deere J,
editor. *Funct. Ecol.* [Internet] 33:60–72. Available from:
https://onlinelibrary.wiley.com/doi/10.1111/1365-2435.13245

Svensson Erik I. 2023. Phenotypic selection in natural populations: what have we learned in 40
years? *Evolution* [Internet] 77:1493–1504. Available from:
https://academic.oup.com/evolut/article/77/7/1493/7146186

Svensson Erik I. 2023. Why high-condition males might be bad news for females and
populations. *Proc. Natl. Acad. Sci.* [Internet] 120:e2301989120. Available from:
https://pnas.org/doi/10.1073/pnas.2301989120

Tao W, Sun L, Shi H, Cheng Y, Jiang D, Fu B, Conte MA, Gammerdinger WJ, Kocher TD,
Wang D. 2016. Integrated analysis of miRNA and mRNA expression profiles in tilapia gonads at
an early stage of sex differentiation. *BMC Genomics* [Internet] 17:328. Available from:
http://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-016-2636-z

The Gene Ontology Consortium, Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ,
Ebert D, Feuermann M, Gaudet P, Harris NL, et al. 2023. The Gene Ontology knowledgebase in
2023. *Genetics* [Internet] 224:iyad031. Available from: https://doi.org/10.1093/genetics/iyad031

Tizaoui K, Terrazzino S, Cargnin S, Lee KH, Gauckler P, Li H, Shin JI, Kronbichler A. 2021. The
role of PTPN22 in the pathogenesis of autoimmune diseases: A comprehensive review. *Semin.*

Arthritis Rheum. [Internet] 51:513–522. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0049017221000342

Torban E, Patenaude A-M, Leclerc S, Rakowiecki S, Gauthier S, Andelfinger G, Epstein DJ, Gros P. 2008. Genetic interaction between members of the Vangl family causes neural tube defects in mice. *Proc. Natl. Acad. Sci.* [Internet] 105:3449–3454. Available from: https://pnas.org/doi/full/10.1073/pnas.0712126105

Tosto NM, Beasley ER, Wong BBM, Mank JE, Flanagan SP. 2023. The roles of sexual selection and sexual conflict in shaping patterns of genome and transcriptome variation. *Nat. Ecol. Evol.* [Internet] 7:981–993. Available from: https://www.nature.com/articles/s41559-023-02019-7

Trappes R, Nematipour B, Kaiser MI, Krohs U, Van Benthem KJ, Ernst UR, Gadau J, Korsten P, Kurtz J, Schielzeth H, et al. 2022. How Individualized Niches Arise: Defining Mechanisms of Niche Construction, Niche Choice, and Niche Conformance. *BioScience* [Internet] 72:538–548. Available from: https://academic.oup.com/bioscience/article/72/6/538/6581356

Turner JMA, Aprelikova O, Xu X, Wang R, Kim S, Chandramouli GVR, Barrett JC, Burgoyne PS, Deng C-X. 2004. BRCA1, Histone H2AX Phosphorylation, and Male Meiotic Sex Chromosome Inactivation. *Curr. Biol.* [Internet] 14:2135–2142. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0960982204009194

Varì R, Scazzocchio B, Filardi T, Citarella A, Bellenghi M, Masella R, Santangelo C. 2021. Significance of Sex Differences in ncRNAs Expression and Function in Pregnancy and Related Complications. *Biomedicines* [Internet] 9:1509. Available from: https://www.mdpi.com/2227-9059/9/11/1509

Vermeulen M, Eberl HC, Matarese F, Marks H, Denissov S, Butter F, Lee KK, Olsen JV, Hyman AA, Stunnenberg HG, et al. 2010. Quantitative Interaction Proteomics and Genome-wide Profiling of Epigenetic Histone Marks and Their Readers. *Cell* [Internet] 142:967–980. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0092867410009517

Voichek Y, Weigel D. 2020. Identifying genetic variants underlying phenotypic variation in plants without complete genomes. *Nat. Genet.* [Internet] 52:534–540. Available from: https://www.nature.com/articles/s41588-020-0612-7

Wagner S, Whiteley SL, Castelli M, Patel HR, Deveson IW, Blackburn J, Holleley CE, Marshall Graves JA, Georges A. 2023. Gene expression of male pathway genes sox9 and amh during early sex differentiation in a reptile departs from the classical amniote model. *BMC Genomics* [Internet] 24:243. Available from: https://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-023-09334-0

Walker LJ, Roque RA, Navarro MF, Granato M. 2021. Agrin/Lrp4 signal constrains MuSK-dependent neuromuscular synapse development in appendicular muscle. *Development* [Internet] 148:dev199790. Available from: https://journals.biologists.com/dev/article/148/21/dev199790/272655/Agrin-Lrp4-signal-constrains-MuSK-dependent

Wang F, Jia Y, Wang P, Yang Q, Du Q, Chang Z. 2017. Identification and profiling of Cyprinus carpio microRNAs during ovary differentiation by deep sequencing. *BMC Genomics* [Internet] 18:333. Available from: http://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-017-3701-y

Wang M, Jiang S, Wu W, Yu F, Chang W, Li P, Wang K. 2018. Non-coding RNAs Function as Immune Regulators in Teleost Fish. *Front. Immunol.* [Internet] 9:2801. Available from: https://www.frontiersin.org/article/10.3389/fimmu.2018.02801/full

Wilson AB. 2001. Male Pregnancy in Seahorses and Pipefishes (Family Syngnathidae): Rapid Diversification of Paternal Brood Pouch Morphology Inferred From a Molecular Phylogeny. *J. Hered.* [Internet] 92:159–166. Available from: https://academic.oup.com/jhered/article-lookup/doi/10.1093/jhered/92.2.159

Wilson AB, Ahnesjo I, Vincent ACJ, Meyer A. 2003a. THE DYNAMICS OF MALE BROODING, MATING PATTERNS, AND SEX ROLES IN PIPEFISHES AND SEAHORSES (FAMILY SYNGNATHIDAE). *Evolution* [Internet] 57:1374–1386. Available from: https://academic.oup.com/evolut/article/57/6/1374/6756757

Wilson AB, Ahnesjo I, Vincent ACJ, Meyer A. 2003b. THE DYNAMICS OF MALE BROODING, MATING PATTERNS, AND SEX ROLES IN PIPEFISHES AND SEAHORSES (FAMILY SYNGNATHIDAE). *Evolution* [Internet] 57:1374–1386. Available from: https://academic.oup.com/evolut/article/57/6/1374/6756757

Wright AE, Fumagalli M, Cooney CR, Bloch NI, Vieira FG, Buechel SD, Kolm N, Mank JE. 2018. Male-biased gene expression resolves sexual conflict through the evolution of sex-specific genetic architecture. *Evol. Lett.* [Internet] 2:52–61. Available from: https://academic.oup.com/evlett/article/2/2/52/6697426

Wu S, Zhou J, Zhang K, Chen H, Luo M, Lu Y, Sun Y, Chen Y. 2020. Molecular Mechanisms of PALB2 Function and Its Role in Breast Cancer Management. *Front. Oncol.* [Internet] 10:301. Available from: https://www.frontiersin.org/article/10.3389/fonc.2020.00301/full

Wu TD, Watanabe CK. 2005. GMAP: a genomic mapping and alignment program for mRNA and EST sequences. *Bioinformatics* [Internet] 21:1859–1875. Available from: https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/bti310

Xiao J, Zhong H, Zhou Y, Yu F, Gao Y, Luo Y, Tang Z, Guo Z, Guo E, Gan X, et al. 2014. Identification and Characterization of MicroRNAs in Ovary and Testis of Nile Tilapia (Oreochromis niloticus) by Using Solexa Sequencing Technology.Guan X-Y, editor. *PLoS ONE* [Internet] 9:e86821. Available from: https://dx.plos.org/10.1371/journal.pone.0086821

Xiong W, He L, Lai J, Dooner HK, Du C. 2014. HelitronScanner uncovers a large overlooked cache of *Helitron* transposons in many plant genomes. *Proc. Natl. Acad. Sci.* [Internet] 111:10263–10268. Available from: https://pnas.org/doi/full/10.1073/pnas.1410068111

Xu Z, Wang H. 2007. LTR_FINDER: an efficient tool for the prediction of full-length LTR retrotransposons. *Nucleic Acids Res.* [Internet] 35:W265–W268. Available from: https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gkm286

Yajima M, Wessel GM. 2011. The DEAD-box RNA helicase Vasa functions in embryonic mitotic progression in the sea urchin. *Development* [Internet] 138:2217–2222. Available from: https://journals.biologists.com/dev/article/138/11/2217/44377/The-DEAD-box-RNA-helicase-Vasa-functions-in

Yajima M, Wessel GM. 2015. Essential elements for translation: the germline factor Vasa functions broadly in somatic cells. *Development* [Internet] 142:1960–1970. Available from: https://journals.biologists.com/dev/article/142/11/1960/46767/Essential-elements-for-translation-the-germline

Yamagata A, Yoshida T, Sato Y, Goto-Ito S, Uemura T, Maeda A, Shiroshima T, Iwasawa-Okamoto S, Mori H, Mishina M, et al. 2015. Mechanisms of splicing-dependent trans-synaptic adhesion by PTPδ–IL1RAPL1/IL-1RAcP for synaptic differentiation. *Nat. Commun.* [Internet] 6:6926. Available from: https://www.nature.com/articles/ncomms7926

Yan H, Liu Q, Jiang J, Shen X, Zhang L, Yuan Z, Wu Y, Liu Y. 2021. Identification of sex differentiation-related microRNA and long non-coding RNA in Takifugu rubripes gonads. *Sci. Rep.* [Internet] 11:7459. Available from: https://www.nature.com/articles/s41598-021-83891-w

Yang W, Garrett L, Feng D, Elliott G, Liu X, Wang N, Wong YM, Choi NT, Yang Y, Gao B. 2017. Wnt-induced Vangl2 phosphorylation is dose-dependently required for planar cell polarity in mammalian development. *Cell Res.* [Internet] 27:1466–1484. Available from: https://www.nature.com/articles/cr2017127

Yasumura M, Yoshida T, Yamazaki M, Abe M, Natsume R, Kanno K, Uemura T, Takao K, Sakimura K, Kikusui T, et al. 2014. IL1RAPL1 knockout mice show spine density decrease, learning deficiency, hyperactivity and reduced anxiety-like behaviours. *Sci. Rep.* [Internet] 4:6613. Available from: https://www.nature.com/articles/srep06613

Yatsu R, Miyagawa S, Kohno S, Parrott BB, Yamaguchi K, Ogino Y, Miyakawa H, Lowers RH, Shigenobu S, Guillette LJ, et al. 2016. RNA-seq analysis of the gonadal transcriptome during Alligator mississippiensis temperature-dependent sex determination and differentiation. *BMC Genomics* [Internet] 17:77. Available from: https://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-016-2396-9

Yoshida T, Mishina M. 2008. Zebrafish orthologue of mental retardation protein IL1RAPL1 regulates presynaptic differentiation. *Mol. Cell. Neurosci.* [Internet] 39:218–228. Available from: https://linkinghub.elsevier.com/retrieve/pii/S1044743108001656

Zhang F, Ma J, Wu J, Ye L, Cai H, Xia B, Yu X. 2009. PALB2 Links BRCA1 and BRCA2 in the DNA-Damage Response. *Curr. Biol.* [Internet] 19:524–529. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0960982209007234

Zhang H, Hou J, Liu H, Zhu H, Xu G, Xu J. 2020. Adaptive evolution of low-salinity tolerance and hypoosmotic regulation in a euryhaline teleost, Takifugu obscurus. *Mar. Biol.* [Internet] 167:90. Available from: https://link.springer.com/10.1007/s00227-020-03705-x

Zhang N, Liang J, Tian Y, Yuan L, Wu L, Miao S, Zong S, Wang L. 2010. A Novel Testis-specific GTPase Serves as a Link to Proteasome Biogenesis: Functional Characterization of RhoS/RSA-14-44 in Spermatogenesis.Kaibuchi K, editor. *Mol. Biol. Cell* [Internet] 21:4312–4324. Available from: https://www.molbiolcell.org/doi/10.1091/mbc.e10-04-0310

Zhang R-G, Li G-Y, Wang X-L, Dainat J, Wang Z-X, Ou S, Ma Y. 2022. TEsorter: An accurate and fast method to classify LTR-retrotransposons in plant genomes. *Hortic. Res.* [Internet] 9:uhac017. Available from: https://academic.oup.com/hr/article/doi/10.1093/hr/uhac017/6532211

Zhu H, Wen Z, Zhang A, Liu D, Wang H, Cheng Y, Yang X, Xiao Y, Li J, Sun D, et al. 2023. RhoGDIα regulates spermatogenesis through Rac1/cofilin/F-actin signaling. *Commun. Biol.* [Internet] 6:214. Available from: https://www.nature.com/articles/s42003-023-04579-7

Figure 1. Panels - *Hippocampus erectus*, *Nerophis ophidion*, *Syngnathus typhle*. **Subpanel A** - male vs female allele frequencies of Kissplice-called SNPs. Coloured grey are the variants that show neither sex-specificity, nor differential expression between sexes. Marked with blue are male-specific SNPs, marked with pink - female specific SNPs. **Subpanel B** - results of an enrichment analysis for 3 data sources: GO BP (red), KEGG (green) and Reactome (blue). Each point represents an enriched term, with name of the term located on the Y-axis, and padj values on the X-axis. The size of the point represents a relationship between the number of genes from the list that belong to this term vs the term size. Meaning, point size represents a proportion of the term is that is represented but the overlap genes.

Figure 2. **Panel A** - Fst values per 10 kb windows across the genome in *H.erectus*, *N.ophidion*, *S.typhle*. Y-axis - Fst value, X-axis -position within a genome. Colored points - ones that pass the 0.4 threshold. On HE plot, grey rounded rectangle marks candidate sex chromosome. **Panel B** - the overlap between genes with elevated Fst and DEG in *H.erectus*, *N.ophidion*, *S.typhle*. Blue colored circles represent gill DEGs, red colored circles represent head kidney DEGs, purple - the overlap between lists of genes.Each panel with gene lists is connected to the appropriate overlap sector with a dotted line. The lists are ordered based on *padj* value and colored based on sex (red - up-regulation in males, blue - up-regulation in females).

Figure 3. **Panels A and B** - results of kmerGWAS analysis. Y-axis - number of ABYSS contigs that aligned to the genome with BLASTn per 1Mb (i.e. adjusted for size), X-axis - the chromosome. **Panel C** - synteny analysis between *H.abdominalis* and *H.erectus* genome. Each line connects 1-1 orthologues in HA and HE. HA chromosomes were ordered based on their length, HE chromosomes were ordered by their relation to HA genome. Red marks sex-linked chr identified in our study. Blue - HE sex-linke chr identified by Long, Bold Black - *H.abdominalis* sex-linked chr.