Rapid redirection of auxin fluxes during root gravitropism by translocation of NGR

proteins driving polarization of PIN-activating kinases

Ivan Kulich*, Julia Schmid*, Anastasia Teplova*, Linlin Qi, Jiří Friml[✉]

Institute of Science and Technology Austria (ISTA), Klosterneuburg, Austria

* These authors contributed equally and are listed alphabetically.

[✉]e-mail: jiri.friml@ist.ac.at

Abstract

Root gravitropic bending represents a fundamental aspect of terrestrial plant physiology.

Gravity is perceived by sedimentation of starch-rich plastids (statoliths) to the bottom of the

central root cap cells. Following gravity perception, intercellular auxin transport is redirected

downwards leading to an asymmetric auxin accumulation at the lower root side causing

inhibition of cell expansion, ultimately resulting in downwards bending. How gravity-induced

statoliths repositioning is translated into asymmetric auxin distribution remains unclear

despite PIN auxin efflux carriers and the Negative Gravitropic Response of roots (NGR)

proteins polarize along statolith sedimentation, thus providing a plausible mechanism for

auxin flow redirection. In this study, using a functional NGR1-GFP construct, we visualized

the NGR1 localization on the statolith surface and plasma membrane (PM) domains in close

proximity to the statoliths, correlating with their movements. We determined that NGR1

binding to these PM domains is indispensable for NGR1 functionality and relies on cysteine

acylation and adjacent polybasic regions as well as on lipid and sterol PM composition.

Detailed timing of the early events following graviperception suggested that both NGR1

repolarization and initial auxin asymmetry precede the visible PIN3 polarization. This

discrepancy motivated us to unveil a rapid, NGR-dependent translocation of PIN-activating

AGCVIII kinase D6PK towards lower PMs of gravity-perceiving cells, thus providing an

attractive model for rapid redirection of auxin fluxes following gravistimulation.

Introduction

Plants are capable of sensing gravity and adjust their organs' growth direction accordingly.

This phenomenon is called gravitropism, with positive gravitropism referring to the root

growing down towards the soil and negative gravitropism describing the upward growth of

the shoot (Kawamoto and Morita, 2022; Kolesnikov et al., 2016; Takahashi et al., 2021).

The model plant *Arabidopsis thaliana* senses gravity in the shoot endodermal cells and root

cap columella cells, in special gravity sensing cells, which contain starch enriched plastids

- amyloplasts (Kiss et al., 1997, 1996, 1989; Morita, 2010). Due to their higher density,

amyloplasts thus act as statoliths and sediment within cells depending on the direction of

gravity (Sack, 1991). Following amyloplast sedimentation, the PIN auxin efflux carriers

(PIN3 and PIN7) polarize towards the bottom cell sides, thus creating a downwards

polarized auxin flow, which subsequently results in the asymmetric auxin accumulation at

the lower side of the organ mediating the shoot/root curvature (Band et al., 2012; Friml et

al., 2002; Kleine-Vehn et al., 2010; Rakusová et al., 2011). While sedimenting amyloplasts

clearly serve as a signal for asymmetric auxin distribution, the precise mechanism remains

elusive (Han et al., 2021; Kiss et al., 1989).

Recently, the LAZY protein family has been identified as a crucial player in the

process of auxin redistribution and normal tropic response for both roots and shoots

following gravistimulation (Jiao et al., 2021; Li et al., 2007; Yoshihara et al., 2013; Yoshihara

and Iino, 2007). LZY1, LZY2, and LZY3 function redundantly in shoot gravitropism; LZY2,

LZY3, and LZY4 - also referred to as NGR1, NGR3, and NGR2 (NEGATIVE GRAVITROPIC

RESPONSE OF ROOTS) - regulate the root gravitropism (Kawamoto and Morita, 2022).

ngr1,2,3 triple mutant plants display an anti-gravitropic phenotype with the root growing

slightly upwards (Ge and Chen, 2016). This phenotype is accompanied by an inverted PIN3

polarization, thus redirecting the auxin flow to the upper side of the gravistimulated root (Ge

and Chen, 2019; Taniguchi et al., 2017). Additionally, it was recently demonstrated that

following gravistimulation, NGR3 localizes asymmetrically to the plasma membrane (PM)

in columella cells of lateral roots (Furutani et al., 2020). However, the structure and

molecular function of NGR proteins, as well as specific domains that could allow for PM

association and polarization are still unknown. Polarization of NGR3 further acts as a

precursor of PIN3 relocalization (Furutani et al., 2020). Additionally, PIN transport activity

is finetuned by phosphorylation mediated by AGCVIII serine/threonine kinases like D6

PROTEIN KINASE (D6PK) (Barbosa and Schwechheimer, 2014). Whether additional

factors play a role in this signaling cascade and how the underlying molecular mechanism

connecting gravity sensing to asymmetric auxin distribution may look like, remains to be

investigated.

Here, we provide novel insights into mechanisms of gravity-induced NGR1

polarization and downstream processes leading to PIN-dependent auxin fluxes activation

and ultimately, gravitropic bending. We show the crucial function of polybasic regions and

acylation of NGR1 for its binding to the PM and amyloplasts in columella cells, which is

essential for its function in gravitropic bending. Pharmacological approaches show that the

membrane composition affects NGR1 PM binding and polarization. Finally, we show rapid

translocation of PIN activator - D6 protein kinase as an event downstream of NGR signaling.

These observations suggest a model where statolith sedimentation is accompanied by

rapid NGR relocation, which mediates similar relocation of the D6PK activator of PIN
transporters, providing means to rapidly redirect asymmetric auxin fluxes in response to
gravity.
77
78 **Results**
79
NGR1 localizes to statoliths and their sedimentation is accompanied by NGR1
polarization at the lower PM
To observe NGR localization in the columella cells, we cloned *NGR1-GFP* under its native
promoter (*NGR1p::NGR1-GFP*) and transformed it into the *ngr1/2/3* triple mutants. This
construct fully rescued the agravitropic bending phenotype of the *ngr1/2/3* triple mutant (see
further). NGR1-GFP displayed a highly specific columella expression, which was most
prominent at the PM and the statolith periphery. NGR1-GFP signal from the PM was not
evenly distributed, but rather polarized to the lower side of the columella cells in the vicinity
of the sedimented statoliths (Fig. 1A).
Constitutive overexpression of NGR1-GFP under 35S promoter (*35Sp::NGR1-GFP*)
confirmed that in multiple other cell types, NGR1-GFP localizes to the PM suggesting direct
membrane binding mechanism. Additionally, NGR1 is associated with other plastid types
such as chloroplasts in guard cells (Fig. S1A). To separate the GFP signal from the
chlorophyll, we used fluorescence lifetime imaging, which confirmed that NGR1-GFP
localizes at the chloroplast periphery (Fig. S1B). This implies a general plastid targeting
mechanism of NGR. Notably, chloroplasts in the vicinity of the PM strongly correlated with
NGR1 accumulating at the PM nearby, similar to the scenario in columella (Fig. S1A,C).
This shows that NGR1-GFP localizes to the plastids and PM in their vicinity
providing a mechanism for how gravity-induced statolith sedimentation can polarize NGR1-
GFP to the bottom PM.
100
Timing of events during the gravitropic response
Next, we tracked the timing of events during gravitropic bending. We used a vertical stage
microscope and minimized the time required for rotating the sample. 180° rotation was
performed during the imaging so that precise timing could be determined from the individual
frame timestamps. Following gravitropic stimulus, NGR1 at the PM relocalized dynamically
with the falling statoliths (Fig. 1B). First, the signal at the top membrane decreased and then
reached the bottom PM together with statoliths, around 5 minutes after the stimulation (Fig.
1C,D). If the root was rotated 180° from the vertical position, the time required for the
statolith sedimentation has increased due to the cell's elongated shape and the presence
of vacuole. In such a scenario, it took more than 15 minutes for the statoliths to fall. Again,

the increase of the NGR1 signal at the bottom side correlated with the statolith occurrence,
implying that statolith vicinity is the region where NGR1 signal gets enriched (Fig. S1D).
When monitored for an extended period of time, NGR1 signal was always decorating the
PM proximal to the statolith (Supplementary movie 1).

To put these early gravitropic events in the context of the auxin gradient formation,
we tracked PIN3 localization dynamics, intracellular calcium transients, and auxin reporter
DII-Venus (Brunoud et al., 2012). Calcium transients are one of the fastest known auxin
response events (Monshausen et al., 2011) and we monitored them using plants stably
transformed with GCaMP3 fluorescent sensor (Zariwala et al., 2012). While the signal
increase at the bottom side of the root was visible within a minute after statoliths touched
the bottom membrane, its decrease at the top PM started immediately after rotation. This
implies that the first signaling event in the root gravitropic bending is the statolith removal
from the top membrane, rather than its arrival at the bottom (Fig. 1E). While subsequent
changes in the DII-Venus degradation were consistent with the calcium-based timing, the
first visible PIN3 polarization to the bottom columella cell sides was detectable only after 15
minutes (Fig. 1F,G). Nonetheless, already the earliest auxin asymmetry is likely generated
by PIN-mediated polar auxin transport as it was completely blocked by the treatment with
N-1-naphthylphthalamic acid (NPA) (Fig. 1E), a PIN inhibitor (Abas et al., 2021).

To summarize, statolith sedimentation is accompanied by rapid NGR1 relocation to
the PM subdomain proximal to the NGR1-containing statoliths. The first event in gravitropic
response is the NGR1 leaving the top membrane accompanied by decrease in the auxin
signaling at the upper root side, followed by NGR1 arrival at the bottom membrane and
increase in auxin signaling at the bottom side. This initial auxin asymmetry is mediated by
PIN-dependent auxin transport, despite visible polarization of PIN3 can be detected only
later.

Fig. 1 | NGR1-GFP relocalization in the root columella.

A) Colocalization of NGR1-GFP (yellow) with membrane stain FM4-64 (magenta). The
statolith periphery is also decorated by the NGR1-GFP signal. NGR1-GFP is
significantly polarized to the bottom area as quantified using upper/lower signal ratio
(1=even distribution)

B) NGR1-GFP ~30 seconds and 5 minutes after gravitropic bending. Orange arrows
depict areas where the signal increased after contact with the statoliths. White stars
in the transmission channel depict statoliths.

C) Time course of individual columella cell expressing NGR1-GFP. Numbers depict
minutes from the 180° rotation.

D) Top: schematic depiction of the experiment design (valid for D-G). Bottom:
Quantification of the NGR1-GFP signal at the lower and upper membranes. Similar
setup was used for E. and F. Blue stripes in D-F represent intervals at which
statoliths touched the bottom membrane.

E) GCaMP signal after root gravitropic bending without (top) and with 20µM NPA
(bottom).

F) PIN3 enrichment at the bottom of the central columella cells.

G) DII Venus signal changes after root gravitropic bending.

Blue lines represent approximate time of statolith contact with the new bottom membrane.

Bars = 10µm

Fig. S1 | NGR1 localization in columella and guard cells.

A)   35Sp::NGR1-GFP colocalization with chloroplasts in guard cells. Contact sites with PM are depicted by white arrowheads.

B)   35Sp::NGR1-GFP colocalization with chloroplasts in guard cells performed by fluorescence lifetime imaging to separate chlorophyll autofluorescence from GFP.

C)   35Sp::NGR1-GFP in guard cells imaged using airyscan superresolution. White arrowheads point to areas of contact with the PM.

D)   NGR1-GFP localization in the columella cell rotated from vertical position upside down (180°). Numbers indicate minutes after rotation.

Bars = 5µm

NGR1 polarization does not require BFA-sensitive trafficking

To gain more insight into the dynamics of the NGR1 localization to the PM and amyloplasts, we tested different inhibitors to alter membrane composition and interfere with endocytic trafficking. Recently, it has been shown that RCC1-like domain (RLD) proteins directly interact with NGRs and play a crucial role in polar auxin transport during gravity signaling (Furutani et al., 2020). As RLD trafficking is known to be sensitive to the ARF-GEF inhibitor Brefeldin A (BFA) (Furutani et al., 2020; Wang et al., 2022), we examined if the drug also affects the localization and polarization of NGR1. 5-day-old *NGR1p::NGR1-GFP* seedlings were incubated with 50µM BFA together with the protein synthesis inhibitor, cycloheximide (CHX, 50µM) and the endocytic tracer FM4-64 (2µM) as described previously (Geldner et al., 2001). Interestingly, while FM4-64-stained membranes accumulated into so-called BFA compartments, NGR1-GFP did not and still showed association with the PM and amyloplasts (Fig. 2A-C). Furthermore, the polarization of NGR1 following gravity stimulation was not disrupted in the presence of BFA. Similar effects were observed in overexpressed

NGR1 as it also did not show co-localization with BFA aggregates (Supplementary Fig. 2A-
C).

This shows that NGR1 localization and gravity-induced polarization does not
undergo BFA-sensitive endocytic recycling mediated by the ARF-GEF GNOM as it has
been shown for other proteins involved in gravity response like RLDs and PINs (Geldner et
al., 2001; Naramoto et al., 2014; Wang et al., 2022).

NGR1 PM association requires specific membrane composition

NGR1 displays PM localization although transmembrane domains are not predicted in its
structure and it does not undergo BFA-sensitive trafficking. This implies the existence of
posttranslational modifications such as S-acylation to associate with PM. To test for the
presence of S-acylation sites, we utilized the reducing properties of Dithiothreitol (DTT),
which interferes with the formation of the thioester bond thus effectively preventing S-
acylation (Sabol et al., 2017). Following previously described conditions, NGR1-GFP
showed an increase in cytoplasmic signal and reduction of PM association (Fig. 2D). Similar
effect of DTT was observed when expressing NGR1 under the 35S promoter, suggesting
that there is indeed an acylation site (Fig. S2D-E). Interestingly, NGR1 still localizes to
amyloplasts, indicating a different way of plastid association compared to the PM.

Next, we tested whether apart from the NGR1 S-acylation, the membrane
composition itself might be essential for the NGR1 PM association. Previously it has been
shown that polybasic regions (PBRs) of proteins can target them to anionic phospholipids
like PtdIns(4)P (PI4P) (Hammond et al., 2012; McLaughlin and Murray, 2005). We tested
whether the dissociation of PI4P from the PM will affect the PM association. *NGR1p::NGR1-*
GFP was incubated for 30 minutes with 30 µM phenylarsine oxide (PAO), a PtdIns(4) kinase
inhibitor (Hammond et al., 2012; Vermeer et al., 2009). The NGR1-GFP cytosolic signal
was significantly increased, indicating that the interplay between PBRs and phospholipids
is important for NGR1 PM but not amyloplast association (Fig. 2D).

Next, we wanted to determine whether sterols might influence the PM association
of NGR1. NGR1-GFP seedlings were grown on fenpropimorph (FEN), an inhibitor of sterol
biosynthesis that decreases the number of sterols and can trigger their conversion into
cyclopropyl sterols (Frescatada-Rosa et al., 2014). NGR1-GFP showed an increase in
cytoplasmic signal and membrane localization was largely lost (Fig. 2D). Comparable
effects were also observed after treatment with another sterol synthesis inhibitor lovastatin
(LOV) (Fig. 2D; Fig. S2F). However, similarly to DTT and PAO, amyloplast localization could
not be disturbed.

These observations show that NGR1 PM but not necessarily plastid localization
depends likely on acylation, phospholipid, and sterol membrane composition.

Fig. 2 | Role of trafficking and PM composition in NGR1 localization.

    A) NGR1-GFP roots treated with 50μM CHX, 50μM BFA, and 2μM FM4-64; mock: DMSO; NGR1-GFP in yellow, FM4-64 in magenta

    B) Fluorescence intensity of FM4-64 and NGR1-GFP at the PM, amyloplasts and BFA bodies normalized to the average fluorescence intensity of FM4-64 at the PM. Statistical measurements were done with one-way ANOVA; *** equals adj. p-value of <0.0001

    C) Cytofluorogram of colocalization of FM4-64 and NGR1-GFP using Pearson's coefficient

    D) NGR1-GFP roots treated with FEN, PAO, and DTT. PM/cytoplasm ratio was calculated for mock, FEN, LOV, PAO, and DTT treated seedlings. Using one-way ANOVA, *b* is significantly different from *a* with an adj. p-value of <0.0001, *c* is significantly different from *a* with an adj. p-value of 0.006.

Bars = 10μm

Fig. S2 | Pharmacological treatments of 35Sp::NGR1-GFP seedlings.

A) BFA treatment of 35Sp::NGR1-GFP seedlings: pretreatment with 50µM CHX, followed by treatment with 50µM BFA for 90min and 2µM FM4-64 for 10min. mock: DMSO. Bars = 20µm

240
241
242

B) BFA treatment of 35Sp::NGR1-GFP seedlings: fluorescence intensity of FM4-64 and 35Sp::NGR1-GFP -GFP at PM and BFA bodies normalized to intensity of FM4-64 PM; statistical analysis with one-way ANOVA: *** equals p-value <0.0001

243
244
245

C) Cytofluorogram of JACoP colocalization analysis; intensity of pixels of FM4-64 on x-axis and 35Sp::NGR1-GFP on y-axis

246
247

D) 50mM DTT treatment of 35Sp::NGR1-GFP plants for 3h followed by 10min co-treatment with FM4-64; profile plot measured along white line. Bars = 20µm

248
249

E) PM/cytoplasm signal ratio for untreated and 50mM DTT treated 35Sp::NGR1-GFP seedlings; statistical analysis with unpaired t-test; *** equals p-value <0.0001

250
251

F) Representative image of 1µM LOV treated NGR1-GFP seedlings; NGR1p::NGR1-GFP in yellow, FM4-64 in magenta. Bars = 10µm

252
253

NGR1 PM localization is synergistically mediated by polybasic regions and a palmitoylation site

255
256

To further study NGR1 membrane localization, we examined the amino acid sequence of the NGR1 to understand which sites of the protein mediate its association with the membrane. We identified 2 polybasic regions $K^{113}K^{114}R^{115}K^{116}$ (PBR1) and $K^{186}K^{187}K^{188}R^{189}$ (PBR2) – these strands of positively charged amino acids may mediate the interaction of NGR1 with negatively charged phospholipids. Also, we noticed a conserved Cys^{206} residue

– a potential S-palmitoylation (PALM) site according to the GPS-Lipid post-translational
modification prediction tool (Xie et al., 2016).

To test the role of these sites in the NGR1 membrane localization we performed
site-directed mutagenesis based on the NGR1-GFP construct. We introduced mutations
$K^{114}E\&R^{115}E$ (-PBR1), and $K^{187}E\&K^{188}E$ (-PBR2) to the polybasic regions of the NGR1 to
neutralize their positive charge, and mutated the conserved cysteine $C^{206}A$ (-PALM) to
abolish the potential palmitoylation site (Fig. 3A). NGR1^{-PALM}-GFP, NGR1^{-PBR2}-GFP, and
NGR1$^{-PBR1\&2}$-GFP under the native NGR1 promoter were transformed into the *ngr1/2/3*
background. These mutant versions of the NGR1 restored the anti-gravitropic root
phenotype of the *ngr1/2/3* triple mutant similar to the NGR1-GFP version without mutations
(Fig. S3A). The NGR1^{-PALM} version resembled the reorientation of the NGR1^{WT}-GFP version
and NGR^{-PBR2} and NGR1$^{-PBR1\&2}$ versions demonstrated a slight delay in the main root angle
restoration after reorientation in the vertical scanner setup (Fig. S3B). Then we performed
confocal imaging of the root columella cells to check the localization of the NGR1^{-PALM}-GFP,
NGR1^{-PBR2}-GFP, and NGR1$^{-PBR1\&2}$-GFP proteins (Fig. S3C). To compare the different
mutant versions we quantified the PM/cytoplasm signal ratio. The PM/cytoplasm ratio was
slightly decreased for NGR1^{-PALM}-GFP, NGR1^{-PBR2}-GFP, and NGR1$^{-PBR1\&2}$-GFP mutants
compared to NGR1^{WT}-GFP (Fig. S3D).

As the mutations only in the palmitoylation site or only in the polybasic regions of
the NGR1 did not lead to the significant disruption of the plant gravitropic response and the
dissociation of the protein from the membrane, we combined these mutations to check their
collective effect. We generated NGR1$^{-PALM\&PBR2}$-GFP with a mutation in the PALM site and
the PBR2, and NGR1$^{-PALM\&PBR1\&2}$ with a mutation in both PBR1 and PBR2 and PALM site.
NGR1p::NGR1$^{-PALM\&PBR2}$-*GFP* and *NGR1p::NGR1*$^{-PALM\&PBR1\&2}$-*GFP* could not fully
complement *ngr1/2/3* triple mutant. The main root of the 5 d.o. seedlings demonstrated
inclination from the vertical angle (Fig. S3E). In contrast to NGR1-GFP complemented
ngr1/2/3 plants, complementation with NGR1$^{-PALM\&PBR2}$ displayed partially compromised
restoration of the main root angle upon reorientation, while the NGR1$^{-PALM\&PBR1\&2}$ root
bending was completely disrupted (Fig. 3B). With confocal microscopy we observed the
dissociation of the NGR1$^{-PALM\&PBR2}$ and NGR1$^{-PALM\&PBR1\&2}$ from the PM (Fig. 3C).
PM/cytoplasm ratio was significantly decreased for both NGR1$^{-PALM\&PBR2}$ and NGR1^{-}
$^{PALM\&PBR1\&2}$ compared to non-mutated NGR1-GFP (Fig. 3D). Taken together, the
combination of mutations in polybasic regions and palmitoylation site resulted in the
decrease of NGR1-GFP PM binding and loss of the protein functionality.

If the selected polybasic regions and palmitoylation site represent solely the mean
of the PM anchoring, then bringing the protein back to the PM by different targeting signal
should restore its functionality. To test this hypothesis, we introduced another membrane-

association site to the mutant versions of NGR1-GFP. It was shown that the N-terminus of

LZY1 mediates its membrane localization potentially via a predicted N-myristoylation site

(Yoshihara and Spalding, 2020). This site is not conserved in the NGR1. Remarkably, LZY1

does not possess the palmitoylated Cys residue conserved in NGR1 (Fig. S4A). We

changed the N-terminus of all the NGR1 mutant versions carrying the mutation $C^{206}A$ to

introduce the LZY1 N-terminal myristoylation site. The obtained $NGR1^{Nmyr\ -PALM}$-GFP,

$NGR1^{Nmyr\ -PALM\&PBR2}$-GFP, $NGR1^{Nmyr\ -PALM\&PBR1\&2}$-GFP versions under NGR1 native promoter

were transformed to the *ngr1/2/3*.

As an alternative, we used a well-characterized membrane-association sequence

from the C-terminus of ROP2 (ROP-CT) carrying a polybasic region and geranyl-

geranylation site (Fig. S4B) (Lin et al., 1996; Yalovsky et al., 2008). Similarly, as for the N-

myristoylation site, we added the ROP-CT on the C-terminus after the GFP tag to all the

variants with the mutated PALM site. The resulting $NGR1p::NGR1^{-PALM\ \ ROP-CT}$-GFP,

$NGR1p::NGR1^{-PALM\&PBR2\ \ \ \ ROP-CT}$-GFP, $NGR1p::NGR1^{-PALM\&PBR1\&2\ \ \ \ ROP-CT}$-GFP were

transformed in *ngr1/2/3* background. PM localization and gravitropic response of the

versions with another acylation site were restored gradually (Fig. S4C-H).

We examined the localization of the GFP signal in the columella of both restored

lines (NGR1$^{Nmyr\ -PALM\&PBR1\&2}$-GFP and NGR1$^{-PALM\&PBR1\&2\ ROP-CT}$-GFP). NGR1$^{Nmyr\ -PALM\&PBR1\&2}$

membrane association was partially restored, and NGR1$^{-PALM\&PBR1\&2\ ROP-CT}$ membrane

association was fully restored (Fig. 3E,F). In accordance with PM localization, the

gravitropic response of those plants was also restored (Fig. 3G,H).

Thus NGR1 PM localization is mediated synergistically by polybasic regions and

palmitoylation site, and the NGR1 PM association is crucial for NGR1 functioning in root

gravitropic response.

Fig. 3 | Polybasic regions and a palmitoylation site mediate NGR1 membrane localization.

A) Scheme of mutagenesis. K114E&R115E (-PBR1), K187E&K188E (-PBR2), and C206A (-PALM) were introduced to disrupt NGR1 polybasic regions and palmitoylation site. Different combinations of mutations were tested.

B) Main root angle restoration upon reorientation in the vertical scanner setup of the *ngr1/2/3* triple mutant complemented with NGR1p::NGR1[-PALM&PBR2]-GFP and NGR1p::NGR1[-PALM&PBR1&2]-GFP in comparison to NGR1p::NGR1[WT]-GFP, Col-0 and *ngr1/2/3* triple mutant.

C) Representative confocal microscopy images of the root columella cells.

D) PM/cytoplasm signal ratio for NGR1p::NGR1-GFP, NGR1p::NGR1[-PALM&PBR2]-GFP, and NGR1p::NGR1[-PALM&PBR1&2]-GFP variants. Mutation in the polybasic region and palmitoylation site leads to the dissociation of NGR1 protein from the PM. For

statistical analysis one-way ANOVA test was used, 'a' is significantly different from 'b' with a p-value<0.0001. The difference between the PM/Cytoplasm signal ratio of NGR1p::NGR1$^{-PALM\&PBR2}$-GFP and NGR1p::NGR1$^{-PALM\&PBR1\&2}$-GFP is not statistically significant.

E) Representative pictures of the root columella cells of NGR1$^{Nmyr\ -PALM\&PBR1\&2}$-GFP and NGR1$^{-PALM\&PBR1\&2\ ROP-CT}$-GFP. Membrane localization of NGR1$^{-PALM\&PBR1\&2}$-GFP is restored upon the addition of LZY1 N-terminal myristoylation site (left) or C-terminal region of ROP2 bearing polybasic region and geranyl-geranylation site (right).

F) PM/cytoplasm ratio quantified for the versions with another acylation site. Different letters designate the statistically significant difference with p<0.0001 in a one-way ANOVA test.

G) Rescued main root angle restoration upon reorientation in the vertical scanner setup of the *NGR1p::NGR1$^{Nmyr\ -PALM\&PBR1\&2}$-GFP* in *ngr1/2/3* line, after the addition of LZY1 N-myristoylation site to the NGR1 version with mutated polybasic regions and palmitoylation site.

H) Rescued gravitropic response of the *NGR1p::NGR1$^{-PALM\&PBR1\&2\ ROP-CT}$-GFP* in *ngr1/2/3*, with the addition of the C-terminus of the ROP2 after the GFP tag.

Fig. S3 | Mutations only in PALM site or PBRs are insufficient to disrupt NGR1 function and localization.

A) Phenotypes of the 5 d.o. seedlings of *ngr1/2/3* complemented with NGR1p::NGR1-GFP, NGR1p::NGR1^{-PALM}-GFP, NGR1p::NGR1^{-PBR2}-GFP , NGR1p::NGR1$^{-PBR1\&2}$-GFP.

B) Restoration of the main root angle upon reorientation in the vertical scanner setup.

C) Representative pictures of the root columella cells expressing different versions of NGR1p::NGR1^{mut}-GFP.

D) Quantification of the PM/cytoplasm ratio for NGR1p::NGR1^{mut}-GFP variants.

E) Phenotypes of the 5 d.o. seedlings of the *NGR1p::NGR1$^{-PALM\&PBR2}$-GFP* in *ngr1/2/3* and *NGR1p::NGR1$^{-PALM\&PBR1\&2}$-GFP* in *ngr1/2/3*.

Fig. S4 | Recovery of NGR1^{mut}-GFP upon addition of another acylation site.

A) Alignment of NGR/LZY homologs. LZY1 has the unique N-terminal myristoylation

motif. Palmitoylated Cys conserved in other NGRs is not present in LZY1.

B) The sequence of the ROP2 C-terminus with a polybasic region and geranyl-

geranylation site.

C) Phenotypes of the plants. *ngr1/2/3* complemented with different versions of

NGR1p::NGR1^{mut}-GFP with additional LZY1 N-myristoylation site (upper row) or

ROP2 C-terminus under native NGR1 promoter (lower row).

D) Representative pictures of the root columella cells expressing different versions of NGR1p::NGR1^{mut}-GFP with additional LZY1 N-myristoylation site (upper row) or ROP2 C-terminus (lower row).

E) Quantification of the PM/Cyt ratio for NGR1p::NGR1$^{Nmyr-mut}$-GFP variants.

F) Quantification of the PM/Cyt ratio for NGR1p::NGR1$^{mut\ ROP-CT}$-GFP variants.

G) Gravitropic response of the *NGR1p::NGR1$^{Nmyr\ mut}$-GFP* in *ngr1/2/3* variants with added N-myristoylation site. The restoration of the main root angle upon reorientation in the vertical scanner setup.

H) Gravitropic response in the vertical scanner setup of *ngr1/2/3* complemented with the *NGR1p::NGR1$^{mut\ ROP-CT}$-GFP* variants with added ROP2 C-terminus.

D6PK relocalizes in columella following gravistimulation in NGR1-dependent manner

NGR1 protein relocalization occurs simultaneously with statolith sedimentation and is rapidly followed by PIN-dependent asymmetric auxin accumulation at the lower root side. These events, however, precede a visible PIN3 polarization towards the bottom columella cell sides (Fig. 1D-G). Alternatively, we hypothesized that asymmetric PIN activation at this position may precede the PIN polarization there and mediate the early auxin asymmetry. To test this, we analyzed the localization of D6PK, a well-established PIN3 activating kinase (Willige et al., 2013). We monitored D6PK under its native promoter with N-terminal YFP and mCHERRY (mCH) tags. This revealed a highly dynamic polarized D6PK localization following gravistimulation. Following 180° rotation, YFP-D6PK and mCH-D6PK displayed a very clear translocation to the new bottom side of the columella cells (Fig. 4A-C). To resolve whether this polarization depends on NGR proteins, we transformed mCHERRY-D6PK (mCH-D6PK) to both *ngr1/2/3* triple mutants and Col-0 wild type plants. Notably, following gravistimulation, D6PK polarization was absent in *ngr1/2/3* triple mutants (Fig. 5C-D, Supplementary movie 2). Surprisingly, other cell types such as meristematic cells or epidermal cells still displayed undisturbed basal (WT-like) D6PK polarity in *ngr1/2/3* triple mutant suggesting a role of NGRs in D6PK targeting specifically during gravity response (Fig. 4C).

Thus, we identified D6PK, a PIN activating kinase to be translocated towards the bottom columella cell sides in an NGR-dependent manner. This provides a plausible mechanism for rapidly redirecting auxin fluxes prior to PIN relocation during root gravitropic response.

413
414

Fig.4 | NGR-Dependent D6 protein kinase transcytosis

A) Temporal color-coded image of YFP-D6PK transcytosis induced by 180°rotation.

B) Single gravistimulated (180°) columella cell undergoing YFP-D6PK transcytosis. White numbers indicate time after rotation (min).

C) mCH-D6PK 10 min after gravistimulation in WT and *ngr1/2/3* triple mutant, same temporal color code as in A. Yellow arrow depicts the new bottom membrane. Red arrowhead depicts meristematic cells, which displayed basal D6PK polarity in both genotypes.

D) Quantification of D6PK signal at the columella cell bottom PM upon 180° rotation. Blue strip depicts approximate time of statolith contact with the bottom PM.

bars = 50μm (A,C) 10μm (B).

Discussion

In this study, we provide several new crucial insights into the mechanism through which plant roots adapt their growth in response to gravity: (i) we mapped the early events after gravistimulation with unprecedented time resolution challenging the current gravitropic models; (ii) we observed real-time relocalization of the key gravitropic regulator, NGR1 on and alongside the descending statoliths and (iii) mapped the conditions for NGR1 crucial association with the PM; and, finally, (iv) we detected rapid, NGR1-dependent relocation of

434 D6PK auxin transport activator. This altogether suggests a new mechanism for rapid
435 redirection of auxin fluxes following gravistimulation.
436
Early events during root gravitropic response
The temporal correlation between NGR1 relocation and auxin asymmetry dynamics was
tested by monitoring auxin-dependent calcium transients. Interestingly, while an auxin-
induced calcium wave can be observed following statolith descent to the bottom side of the
gravity-sensing cells, preceding this, a decline in calcium transients occurs at the upper root
side. These observations suggest that the first signaling event in root gravitropic bending
may not be statolith contact with the cell bottom, but rather the departure of statoliths from
the upper PM, closely followed by NGR1 and D6PK translocation to the bottom cell sides.
Only significantly later, similar polarization of PIN auxin exporters can be confidently
observed. This may be reflected by actual onset of the root bending, as there is growing
evidence, that the actual root bending begins much sooner than previously reported, in fact,
some degree of bending was observed already within the first 5 minutes following
gravistimulation (Dubey et al., 2023; Serre et al., 2023).
450
Insights into NGR membrane association and gravity-induced polarization
NGR-GFP relocation to the new bottom membrane happens rapidly and is not affected
when ARF GEF-mediated endomembrane trafficking is inhibited. Although the precise
mechanism of NGR1 dynamic relocation following gravitropic stimulus still remains elusive,
this study brings some new insights. The NGR1 membrane anchoring mechanism is
dependent on cysteine acylation and adjacent polybasic stretches, which interact with
charged lipids at the membrane surface. Such a mechanism is typical for a plethora of
proteins (Hemsley and Grierson, 2008). Since we could restore the protein functionality by
adding two different membrane anchors on the NGR1 N and C terminus, we conclude that
the mechanism of membrane targeting is rather universal. Point mutated NGR1 proteins
which lost their membrane binding ability still retained the statolith binding, implying an
independent targeting mechanism. Therefore we can conclude that statolith-bound NGR1
is not sufficient for the NGR1 function in gravitropism.
Our pharmacological experiments indicate that both sterols and charged lipids play
a role in the NGR membrane binding. This resembles other highly polarized structures such
as root hairs, where sterols and phosphatidylinositol 4,5-bisphosphate are vital for the
polarity (Champeyroux et al., 2020; Ovecka et al., 2010). Furthermore, PIN2 polarity was
defective in sterol synthesis mutants, implying the role of sterols in planar polarity as well
(Men et al., 2008). Whether there is mobile sterol-rich domain moving alongside the statolith
or the membrane composition is uniform remains to be elucidated.

Gravity-induced D6PK kinase polarization as mechanism for rapid auxin fluxes redirection

PIN auxin transporters were shown to polarize following gravistimulation to the bottom of gravity-sensing cells (Friml et al., 2002; Furutani et al., 2020; Kleine-Vehn et al., 2010) providing a mechanism for redirection of auxin fluxes downwards. Although PIN polarization has been shown to be reliant on NGR proteins (Ge and Chen, 2019), the timing of early gravitropic events resolved here suggests that it occurs too late to account for the initial auxin asymmetry. Either, we are unable to detect early, less-pronounced PIN asymmetry or there is an additional mechanism preceding the auxin flux redirection via PIN relocation.

Our discovery of similar gravity-induced polarization of D6PK AGCVIII kinases, which is contingent on NGR activity, bridges this critical knowledge gap and suggests that, prior to relocation of PIN transporters, AGCVIII-mediated PIN activation at the lower cell sides rapidly redirects auxin fluxes following gravity perception. Notably, PIN3 phosphorylation is important for gravity response (Grones et al., 2018) and higher-order D6PK mutants display mild gravitropic bending phenotypes (Zourelidou et al., 2009), which are, however, weaker than *ngr1/2/3* mutants. This observation might be attributed to the multiplicity of the AGC kinase family and/or an additional mechanism governing the differential accumulation of PIN proteins, following D6PK polarization, thus contributing to the regulation of auxin flux.

Similarly to NGRs, D6PK membrane localization depends on polybasic regions and sterol-rich environments (Barbosa et al., 2016; Stanislas et al., 2015). However, in contrast to NGR proteins, D6PK localization is dependent on BFA-sensitive trafficking regulator GNOM (Barbosa et al., 2016, 2014) and thus is likely delivered to the PM by vesicle trafficking. RLD proteins, implicated in the regulation of the GNOM-dependent PIN trafficking pathway, are directly recruited by NGR homolog NGR3 (LZY3) (Furutani et al., 2020). Notably, RLD proteins contain the BRX domain, which is responsible for the NGR interaction (Furutani et al., 2020). In the different cellular context, BRX protein was shown to be an inhibitor of AGC kinase (PAX) induced PIN3 activation and an interactor of multiple D6PK proteins (Marhava et al., 2018). This implies that BRX containing proteins may act as negative feedback on PIN phosphorylation after gravistimulation. Indeed, our observations indicate that the D6PK signal at the new bottom membrane reaches its maximum approximately 7-8 minutes after gravistimulation and subsequently diminishes, hinting at the existence of a negative feedback mechanism. The involvement of RLDs in this process remains to be investigated.

Taken together, an updated model of early events in root gravity-sensing cells emerges: (i) Statoliths sediment and bring associated NGR proteins to the lower PM; by a

mechanism requiring electrostatic interaction-based association of NGR with the PM; (ii)

dependent on these events, the D6PK translocates almost concomitantly also to the bottom

cell sides, likely by GNOM-dependent trafficking mechanism; (iii) D6PK activates PINs at

the lower side, auxin transport get redirected downwards reflected by a decrease of auxin

response at the upper side followed by auxin response increase at the lower root side; (iv)

D6PK asymmetry gradually diminishes, possibly by action of BRX domain-containing

proteins and the whole PIN-dependent auxin transport machinery gradually relocates to the

lower cell side for more sustained redirection of auxin fluxes. There are still many

mechanistic links to be confirmed and clarified in this model but it is consistent with all

observations on the sequence of events following gravistimulation and with all additional

data from other developmental contexts.

Materials and methods

Plant cultivation

All Arabidopsis mutants and transgenic lines which were used in this project are in the

Columbia-0 (Col-0) background. NGR1p::NGR1-GFP, 35Sp::NGR1-GFP, and

D6PK::mCH-D6PK were generated as part of this research project. D6PK::YFP-D6PK has

been described previously (Zourelidou et al., 2009). Overnight seed sterilization was

executed with chlorine gas. Seeds were sown on solid agarose media – half-strength

Murashige and Skoog (0.5x MS) medium supplemented with 1% sucrose and 0.8% phyto

agar (pH=5.9) – stratified at 4°C for 2 days and subsequently grown vertically at 21°C with

16h light/8h dark cycles.

Cloning strategy

NGR1 promoter was amplified from genomic DNA using the primer pairs NGR1p-B4-FP

and NGR1p-B1r-RP, and was cloned into pDONR-P4P1r. *NGR1-GFP* fusion fragment was

obtained by overlapping PCR using the prime pairs NGR1g-B1-FP and NGR1g-L-RP to

amplify the *NGR1* protein-coding region with genomic DNA as the template and the primer

pairs eGFP-L-FP and eGFP-B2-RP to amplify the *eGFP* coding sequence, with a PKPA

protein linker inserted between NGR1 and GFP, and the resulting fragment was cloned into

pDONR221. The resulting pENTRY clones were recombined into pH7m24GW to get the

final expression construct *NGR1p::NGR1-GFP*. The pENTR clone NGR1-GFP in

pDONR221 was recombined with pH2GW7 to get the construct *35Sp::NGR1-GFP*.

To obtain desired point mutations in polybasic regions and acylation sites,

Mutagenesis of *NGR1p::NGR1-GFP* was performed using Gibson assembly (NEB HiFi

DNA assembly E2621L), overlapping compatible cohesive ends were modified. For ROP

C-terminus, an oligo-bridge Gibson assembly was performed with an oligonucleotide
containing the entire ROP2 hypervariable region. Plants were transformed with floral dip
method (Zhang et al., 2006)

NGR1 transformants were selected on plates containing 30µg/ml hygromycin.
NGR1p::NGR1-GFP plants with genotyped with primers gNGRmut-NGR1prom-fwd and
gNGRmut-T35S-rev, and the middle part of the NGR1 was sequenced to confirm the
presence of the mutations.

D6PKp::mCH-D6PK was produced using the Greengate cloning strategy.
Greengate entry blocks were generated with the use of the Gibson assembly (pGGA-D6PK
promotor) and restriction-ligation assembly (pGGC-D6PK). 2KB fragment upstream of
D6PK start codon was used and the BsaI site was removed using point mutated compatible
cohesive ends for the Gibson assembly between fragments A and B. Other building blocks
used were: pGGB-mCHERRY-linker (pGGB001), pGGD-Dummy (pGGD002), pGGE
UBQ10 terminator (pGGE009), pGGF-d-AlanineR (pGGF003), destination vector
(pGGZ001 modified to contain bacterial kanamycin resistance; Lampropoulos et al., 2013),
plasmids were obtained from Addgene or were kindly provided by Dr. Andrea Bleckmann.
pD6PK::mCH-D6PK transformants bearing DAO-selection cassette were selected on the 3
mM D-Alanine.

Microscopy
For imaging, vertically mounted Zeiss LSM800 was used with air objectives Plan
Apochromat 20x/0.8 M27 and Plan Apochromat 10x/0.45 M27 (Fig. 1G). Air objective Plan
Apochromat 20x/0.8 M27 and water objective Plan-Apochromat 40x/1.2 M27 were used in
Fig. 2 and Fig. S1, S2. Plan apochromat 60x 1.4 Oil objective was used for the airyscan
imaging (Fig. S1C). For fluorescence lifetime imaging (Fig. S1B), Leica Stellaris 5 with Plan-
Apochromat 40x/1.2 M27 was used. GCaMP signal was imaged using GFP emission (500-
540nm) using 405nm and 488nm excitation lasers. Signal obtained by 405nm excitation
was used as a ratiometric reference. For the gravitropic experiments, brightfield images
were always taken together with the fluorescent probes to determine the statolith contact
with basal PM. This was determined as a time at which statolith movement relative to the
cell surface stopped.

Plant treatments
The following inhibitors were dissolved in DMSO: Brefeldin A (Sigma); Cycloheximide
(Sigma); PAO (Sigma); Fenpropimorph (Sigma); Lovastatin (Sigma); NPA (DUCHEFA
N0926). BFA treatment was performed as described previously (Geldner et al., 2001). In
brief, seedlings were submerged in liquid MS medium and pre-treated with 50µM CHX for

30 minutes. Subsequently, they were incubated with 50μM BFA and 50μM CHX for 75
minutes. Then, 2μM FM4-64 was added and co-incubated with 50μM BFA and 50μM CHX
for 15 minutes. Treatment with PAO was executed as described previously (Barbosa et al.,
2016); seedlings were incubated in 30μM PAO-containing liquid MS medium for 20 minutes,
followed by the addition of 2μM FM4-64 and another 10-minute incubation. For
quantification purposes, 5-day-old seedlings grown on AM+ plates with 1μM LOV and
250μg/ml FEN were utilized (Stanislas et al., 2015). Representative images were captured
of 5-day-old seedlings that were initially grown on AM+ and subsequently transferred to
1μM LOV and 50μg/ml FEN for a duration of 2 hours. To treat with DTT, seedlings were
incubated in liquid MS medium supplemented with 50mM DTT (Sigma; dissolved in H2O)
for 3 hours, followed by a 10-minute incubation with 2μM FM4-64 (Sabol et al., 2017).
Unless otherwise stated, treated seedlings were mounted in liquid MS medium for live
imaging. As a control for used inhibitors, solvent control treatments were performed in
parallel.
596

Data analysis
To quantify signals in the individual cells, time series were stabilized by Correct 3D drift Fiji
plugin. All image data was obtained in 16-bit format. ROI were selected and measured
using Microsoft Excel and GraphPad Prism. Data was normalized so that the first value =
1 (Fig. 1 A,D,F, 4D) or by using reference channel (Fig 1E, 2B)
In order to analyze subcellular colocalization we utilized the tool JACoP (Just
Another Co-localization Plugin), that allows for a correlation of pixels of dual-channel
images based on Pearson's coefficient (Bolte and Cordelières, 2006).
Gravitropic bending experiments were performed in the vertically oriented scanner
setup. 4 d.o. seedlings were transferred to a new solid agarose media plate and aligned so
that the roots were straightened. Images obtained with 30 min intervals were registered with
the Fiji StackReg plugin (Schindelin et al., 2012). Roots were manually tracked and angle
increments were calculated using Microsoft Excel.
For the PM/cytoplasm ratio quantification, the background of the confocal images
was subtracted; with the multi-point selection tool 15-50 points on the PM, and 15-50 points
in the cytoplasm were selected, and the intensity of the GFP signal was measured.
The location of the PM membrane was confirmed with the FM4-64 staining. The PM region
was selected on the bottom side of the cells, as NGR1 is localized polarly. The cytoplasmic
region was selected close to the selected membrane region and did not include the region
with amyloplasts and the vacuole. For individual cells the averaged PM signal was divided
by the averaged cytoplasmic signal. The PM/cytoplasm ratio for the lines is quantified as
the average of the PM/cytoplasm ratios for individual cells of this line.

If not stated otherwise, all experiments were performed in triplicate. 3 or more independent
transgenic lines were always used with similar results.

List of primers used in the study

Primer name	Primer sequence
Transgenic line preparation	
NGR1p-B4-FP	GGGGACAACTTTGTATAGAAAAGTTGGAAGAGGAGAAGGTGGGAGAGC
NGR1p-B1r-RP	GGGGACTGCTTTTTTGTACAAACTTGTGTTTCTTTTTTTCTGACAATTGACTG
NGR1p-S-F1	ACTAAACAACCCCTTTTGAAACC
NGR1p-S-F2	CAAGATTGAAAAACTATTTGCCCT
NGR1g-B1-FP	GGGGACAAGTTTGTACAAAAAAGCAGGCTCCATGAAGTTCTTCGGGTGGATG
NGR1g-L-RP	CACCATAGCAGGCTTAGGTATCTCGAGAACTATGACTGTAATCA
eGFP-L-FP	GAGATACCTAAGCCTGCTATGGTGAGCAAGGGCGAGGAG
eGFP-B2-RP	GGGGACCACTTTGTACAAGAAAGCTGGGTGCTACTTGTACAGCTCGTCCATGCC
pGGC_D6PK_for	AGAAGTGAAGCTTGGTCTCAGGCTCCATGATGGCTTCAAAAACTCC
pGGC_D6PK_rev	AGGGCGAGAATTCGGTCTCACTGATCAGAAGAAATCAAACTCAAGATA
pGGA_D6PKpromA_for	GTGAAGCTTGGTCTCAACCTCTGTTGAACCATTTCTAAAAAAAC
pGGA_D6PKpromA_rev	GAGAGAGAGTCCCAATAAATCGTTACCTG
pGGA_D6PKpromB_for	ATTTATTGGGACTCTCTCTCTCTCTCTC
pGGA_D6PKpromB_rev	CGAGAATTCGGTCTCATGTTTAACACAGAGCAATCTTAAAC
pGGA_backbone_for	AACATGAGACCGAATTCTC
pGGA_backbone_rev	AGGTTGAGACCAAGCTTC
Genotyping	
ngr1-F199G07-GT-FP	GCGCAGACAAAAATCTTCTTG
ngr1-F199G07-GT-RP	TTGGTGGACTCGTTTGCTTAC
FLAG-LB4	CGTGTGCCAGGTGCCCACGGAATAGT
ngr2-SAIL723H11-GT-FP	TTTGGTTTTATGGACCCAACC
ngr2-SAIL723H11-GT-RP	AAGAGCTTTCTTCCTCCGATG
SAIL-LB3	TAGCATCTGAATTTCATAACCAATCTCGATACAC
ngr3-GK479C08-GT-FP	GCAAACAGATTTTCTTCACCAC
ngr3-GK479C08-GT-RP	GCACAAGTGGCTTCAAAACTC
GABI-O8409	ATATTGACCATCATACTCATTGC
gNGRmut-NGR1prom-fwd	GTAGTCAAAGTTTGGAACTTGAACACC
gNGRmut-T35S-rev	CTGGGAACTACTCACACATTATTCTGG
Mutagenesis	
mutatePBR2-fwd	TAAGAATAACAAGGAAGAAAGAGACATAAGCAAGAACTCTG

mutatePBR2-rev	TCTTTCTTCCTTGTTATTCTTACTCTCTATTGATATCTC
PBR1&2-fragment1-bb-fwd	TAAGAATAACAAGGAAGAAAGAGACATAAGCAAGAACTCTG
PBR1&2-fragment1-bb-rev	TCACATCAGACTTTTCTTCCTTAGTTCTTGACAAGAGCTTC
PBR1&2-fragment2-insert-fwd	AAGGAAGAAAAGTCTGATGTGAATAGAGAGC
PBR1&2-fragment2-insert-rev	TCTTTCTTCCTTGTTATTCTTACTCTCTATTGATATCTC
mutatePALM_C206A-fwd	CAAGAAGATTTTTGTCGCTGCAGATGGG
mutatePALM_C206A-rev	CAGCGACAAAAATCTTCTTGAAAAGATATGAGACAGAG
add-Nmyr-fwd	CCACAAGTTCAGGGGGGATCATAACAGAACAAGCACTTCC
add-Nmyr-rev	CCTGAACTTGTGGTGCATCCACCCCCAGAACTTCATGGAGCC
ROP2-CT-oligo-bridge	GCTATAAAAGTGGTGCTTCAGCCACCAAAGCAGAAGAAGAAGAAAAAGAATAAG
NGR-ROP2_CT-after-GFP-fwd	ACCGTTGCGCGTTCTTGTGACACCCAGCTTTCTTGTAC
NGR-ROP2_CT-after-GFP-rev	TGAAGCACCACTTTTATAGCCTTGTACAGCTCGTCCATG

Acknowledgements

The research leading to these results has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme grant agreement No 742985 and Austrian Science Fund (FWF): I3630-775 B25 to J.F. This research was also supported by the Lab Support Facility (LSF) and the Imaging and Optics Facility (IOF) of IST Austria, namely Tereza Bělinová for her help with the imaging. J.S. was supported by FemTECH fellowship.

References

Abas L, Kolb M, Stadlmann J, Janacek DP, Lukic K, Schwechheimer C, Sazanov LA, Mach L, Friml J, Hammes UZ. 2021. Naphthylphthalamic acid associates with and inhibits PIN auxin transporters. *Proc Natl Acad Sci U S A* **118**. doi:10.1073/pnas.2020857118

Band LR, Wells DM, Larrieu A, Sun J, Middleton AM, French AP, Brunoud G, Sato EM, Wilson MH, Péret B, Oliva M, Swarup R, Sairanen I, Parry G, Ljung K, Beeckman T, Garibaldi JM, Estelle M, Owen MR, Vissenberg K, Hodgman TC, Pridmore TP, King JR, Vernoux T, Bennett MJ. 2012. Root gravitropism is regulated by a transient lateral auxin gradient controlled by a tipping-point mechanism. *Proc Natl Acad Sci U S A* **109**:4668–4673.

Barbosa ICR, Shikata H, Zourelidou M, Heilmann M, Heilmann I, Schwechheimer C. 2016. Phospholipid composition and a polybasic motif determine D6 PROTEIN KINASE polar association with the plasma membrane and tropic responses.

Development **143**:4687–4700.

Barbosa ICR, Zourelidou M, Willige BC, Weller B, Schwechheimer C. 2014. D6 PROTEIN

KINASE activates auxin transport-dependent growth and PIN-FORMED

phosphorylation at the plasma membrane. *Dev Cell* **29**:674–685.

Barbosa IC, Schwechheimer C. 2014. Dynamic control of auxin transport-dependent

growth by AGCVIII protein kinases. *Curr Opin Plant Biol* **22**:108–115.

Bolte S, Cordelières FP. 2006. A guided tour into subcellular colocalization analysis in

light microscopy. *J Microsc* **224**:213–232.

Brunoud G, Wells DM, Oliva M, Larrieu A, Mirabet V, Burrow AH, Beeckman T, Kepinski

S, Traas J, Bennett MJ, Vernoux T. 2012. A novel sensor to map auxin response and

distribution at high spatio-temporal resolution. *Nature* **482**:103–106.

Champeyroux C, Stoof C, Rodriguez-Villalon A. 2020. Signaling phospholipids in plant

development: small couriers determining cell fate. *Curr Opin Plant Biol* **57**:61–71.

Dubey SM, Han S, Stutzman N, Prigge MJ, Medvecká E, Platre MP, Busch W, Fendrych

M, Estelle M. 2023. The AFB1 auxin receptor controls the cytoplasmic auxin

response pathway in Arabidopsis thaliana. *Mol Plant* **16**:1120–1130.

Frescatada-Rosa M, Stanislas T, Backues SK, Reichardt I, Men S, Boutté Y, Jürgens G,

Moritz T, Bednarek SY, Grebe M. 2014. High lipid order of Arabidopsis cell-plate

membranes mediated by sterol and DYNAMIN-RELATED PROTEIN1A function.

Plant J **80**:745–757.

Friml J, Wiśniewska J, Benková E, Mendgen K, Palme K. 2002. Lateral relocation of auxin

efflux regulator PIN3 mediates tropism in Arabidopsis. *Nature* **415**:806–809.

Furutani M, Hirano Y, Nishimura T, Nakamura M, Taniguchi M, Suzuki K, Oshida R,

Kondo C, Sun S, Kato K, Fukao Y, Hakoshima T, Morita MT. 2020. Polar recruitment

of RLD by LAZY1-like protein during gravity signaling in root branch angle control.

Nat Commun **11**:76.

Ge L, Chen R. 2019. Negative gravitropic response of roots directs auxin flow to control

root gravitropism. *Plant Cell Environ* **42**:2372–2383.

Ge L, Chen R. 2016. Negative gravitropism in plant roots. *Nat Plants* **2**:16155.

Geldner N, Friml J, Stierhof YD, Jürgens G, Palme K. 2001. Auxin transport inhibitors

block PIN1 cycling and vesicle trafficking. *Nature* **413**:425–428.

Grones P, Abas M, Hajný J, Jones A, Waidmann S, Kleine-Vehn J, Friml J. 2018.

PID/WAG-mediated phosphorylation of the Arabidopsis PIN3 auxin transporter

mediates polarity switches during gravitropism. *Sci Rep* **8**:10279.

Hammond GRV, Fischer MJ, Anderson KE, Holdich J, Koteci A, Balla T, Irvine RF. 2012.

PI4P and PI(4,5)P2 are essential but independent lipid determinants of membrane

identity. *Science* **337**:727–730.

Han H, Adamowski M, Qi L, Alotaibi SS, Friml J. 2021. PIN-mediated polar auxin transport
regulations in plant tropic responses. *New Phytol* **232**:510–522.

Hemsley PA, Grierson CS. 2008. Multiple roles for protein palmitoylation in plants. *Trends*
Plant Sci **13**:295–302.

Jiao Z, Du H, Chen S, Huang W, Ge L. 2021. LAZY Gene Family in Plant Gravitropism.
Front Plant Sci **11**:606241.

Kawamoto N, Morita MT. 2022. Gravity sensing and responses in the coordination of the
shoot gravitropic setpoint angle. *New Phytol* **236**:1637–1654.

Kiss JZ, Guisinger MM, Miller AJ, Stackhouse KS. 1997. Reduced gravitropism in
hypocotyls of starch-deficient mutants of Arabidopsis. *Plant Cell Physiol* **38**:518–525.

Kiss JZ, Hertel R, Sack FD. 1989. Amyloplasts are necessary for full gravitropic sensitivity
in roots of Arabidopsis thaliana. *Planta* **177**:198–206.

Kiss JZ, Wright JB, Caspar T. 1996. Gravitropism in roots of intermediate-starch mutants
of Arabidopsis. *Physiol Plant* **97**:237–244.

Kleine-Vehn J, Ding Z, Jones AR, Tasaka M, Morita MT, Friml J. 2010. Gravity-induced
PIN transcytosis for polarization of auxin fluxes in gravity-sensing root cells. *Proc Natl*
Acad Sci U S A **107**:22344–22349.

Kolesnikov YS, Kretynin SV, Volotovsky ID, Kordyum EL, Ruelland E, Kravets VS. 2016.
Molecular mechanisms of gravity perception and signal transduction in plants.
Protoplasma **253**:987–1004.

Lampropoulos A, Sutikovic Z, Wenzl C, Maegele I, Lohmann JU, Forner J. 2013.
GreenGate---a novel, versatile, and efficient cloning system for plant transgenesis.
PLoS One **8**:e83043.

Lin Y, Wang Y, Zhu JK, Yang Z. 1996. Localization of a Rho GTPase Implies a Role in Tip
Growth and Movement of the Generative Cell in Pollen Tubes. *Plant Cell* **8**:293–303.

Li P, Wang Y, Qian Q, Fu Z, Wang M, Zeng D, Li B, Wang X, Li J. 2007. LAZY1 controls
rice shoot gravitropism through regulating polar auxin transport. *Cell Res* **17**:402–
410.

Marhava P, Bassukas AEL, Zourelidou M, Kolb M, Moret B, Fastner A, Schulze WX,
Cattaneo P, Hammes UZ, Schwechheimer C, Hardtke CS. 2018. A molecular
rheostat adjusts auxin flux to promote root protophloem differentiation. *Nature*
558:297–300.

McLaughlin S, Murray D. 2005. Plasma membrane phosphoinositide organization by
protein electrostatics. *Nature* **438**:605–611.

Men S, Boutté Y, Ikeda Y, Li X, Palme K, Stierhof Y-D, Hartmann M-A, Moritz T, Grebe M.
2008. Sterol-dependent endocytosis mediates post-cytokinetic acquisition of PIN2
auxin efflux carrier polarity. *Nat Cell Biol* **10**:237–244.

Monshausen GB, Miller ND, Murphy AS, Gilroy S. 2011. Dynamics of auxin-dependent
Ca2+ and pH signaling in root growth revealed by integrating high-resolution imaging
with automated computer vision-based analysis. *Plant J* **65**:309–318.
Morita MT. 2010. Directional gravity sensing in gravitropism. *Annu Rev Plant Biol* **61**:705–
720.
Naramoto S, Otegui MS, Kutsuna N, de Rycke R, Dainobu T, Karampelias M, Fujimoto M,
Feraru E, Miki D, Fukuda H, Nakano A, Friml J. 2014. Insights into the localization
and function of the membrane trafficking regulator GNOM ARF-GEF at the Golgi
apparatus in Arabidopsis. *Plant Cell* **26**:3062–3076.
Ovecka M, Berson T, Beck M, Derksen J, Samaj J, Baluska F, Lichtscheidl IK. 2010.
Structural sterols are involved in both the initiation and tip growth of root hairs in
Arabidopsis thaliana. *Plant Cell* **22**:2999–3019.
Rakusová H, Gallego-Bartolomé J, Vanstraelen M, Robert HS, Alabadí D, Blázquez MA,
Benková E, Friml J. 2011. Polarization of PIN3-dependent auxin transport for
hypocotyl gravitropic response in Arabidopsis thaliana. *Plant J* **67**:817–826.
Sabol P, Kulich I, Žárský V. 2017. RIN4 recruits the exocyst subunit EXO70B1 to the
plasma membrane. *J Exp Bot* **68**:3253–3265.
Sack FD. 1991. Plant Gravity SensingInternational Review of Cytology, International
Review of Cytology. Elsevier. pp. 193–252.
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,
Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image
analysis. *Nat Methods* **9**:676–682.
Serre NBC, Wernerová D, Vittal P, Dubey SM, Medvecká E, Jelínková A, Petrášek J,
Grossmann G, Fendrych M. 2023. The AUX1-AFB1-CNGC14 module establishes a
longitudinal root surface pH profile. *Elife* **12**. doi:10.7554/eLife.85193
Stanislas T, Hüser A, Barbosa ICR, Kiefer CS, Brackmann K, Pietra S, Gustavsson A,
Zourelidou M, Schwechheimer C, Grebe M. 2015. Arabidopsis D6PK is a lipid
domain-dependent mediator of root epidermal planar polarity. *Nat Plants* **1**:15162.
Takahashi K, Takahashi H, Furuichi T, Toyota M, Furutani-Seiki M, Kobayashi T,
Watanabe-Takano H, Shinohara M, Numaga-Tomita T, Sakaue-Sawano A, Miyawaki
A, Naruse K. 2021. Gravity sensing in plant and animal cells. *NPJ Microgravity* **7**:2.
Taniguchi M, Furutani M, Nishimura T, Nakamura M, Fushita T, Iijima K, Baba K, Tanaka
H, Toyota M, Tasaka M, Morita MT. 2017. The Arabidopsis LAZY1 Family Plays a
Key Role in Gravity Signaling within Statocytes and in Branch Angle Control of Roots
and Shoots. *Plant Cell* **29**:1984–1999.
Vermeer JEM, Thole JM, Goedhart J, Nielsen E, Munnik T, Gadella TWJ Jr. 2009.

Imaging phosphatidylinositol 4-phosphate dynamics in living plant cells. *Plant J*
57:356–372.

Wang L, Li D, Yang K, Guo X, Bian C, Nishimura T, Le J, Morita MT, Bergmann DC, Dong
J. 2022. Connected function of PRAF/RLD and GNOM in membrane trafficking
controls intrinsic cell polarity in plants. *Nat Commun* **13**:7.

Willige BC, Ahlers S, Zourelidou M, Barbosa ICR, Demarsy E, Trevisan M, Davis PA,
Roelfsema MRG, Hangarter R, Fankhauser C, Schwechheimer C. 2013. D6PK
AGCVIII kinases are required for auxin transport and phototropic hypocotyl bending
in Arabidopsis. *Plant Cell* **25**:1674–1688.

Xie Y, Zheng Y, Li H, Luo X, He Z, Cao S, Shi Y, Zhao Q, Xue Y, Zuo Z, Ren J. 2016.
GPS-Lipid: a robust tool for the prediction of multiple lipid modification sites. *Sci Rep*
6:28249.

Yalovsky S, Bloch D, Sorek N, Kost B. 2008. Regulation of membrane trafficking,
cytoskeleton dynamics, and cell polarity by ROP/RAC GTPases. *Plant Physiol*
147:1527–1543.

Yoshihara T, Iino M. 2007. Identification of the gravitropism-related rice gene LAZY1 and
elucidation of LAZY1-dependent and -independent gravity signaling pathways. *Plant*
Cell Physiol **48**:678–688.

Yoshihara T, Spalding EP. 2020. Switching the Direction of Stem Gravitropism by Altering
Two Amino Acids in AtLAZY1. *Plant Physiol* **182**:1039–1051.

Yoshihara T, Spalding EP, Iino M. 2013. AtLAZY1 is a signaling component required for
gravitropism of the Arabidopsis thaliana inflorescence. *Plant J* **74**:267–279.

Zariwala HA, Borghuis BG, Hoogland TM, Madisen L, Tian L, De Zeeuw CI, Zeng H,
Looger LL, Svoboda K, Chen T-W. 2012. A Cre-dependent GCaMP3 reporter mouse
for neuronal imaging in vivo. *J Neurosci* **32**:3131–3141.

Zhang X, Henriques R, Lin S-S, Niu Q-W, Chua N-H. 2006. Agrobacterium-mediated
transformation of Arabidopsis thaliana using the floral dip method. *Nat Protoc* **1**:641–
646.

Zourelidou M, Müller I, Willige BC, Nill C, Jikumaru Y, Li H, Schwechheimer C. 2009. The
polarly localized D6 PROTEIN KINASE is required for efficient auxin transport in
Arabidopsis thaliana. *Development* **136**:627–636.