Gene count normalization in single-cell imaging-based spatially resolved transcriptomics

Authors:

Lyla Atta[1,2], Kalen Clifton[1,2], Manjari Anant[2,3], Jean Fan[1,2]

Affiliations:

1. Department of Biomedical Engineering, Johns Hopkins University, Baltimore, MD 21218,

USA

2. Center for Computational Biology, Whiting School of Engineering, Johns Hopkins University,

Baltimore, MD 21211, USA

3. Department of Neuroscience, Johns Hopkins University, Baltimore, MD 21218, USA

Correspondence should be addressed to:

Jean Fan (jeanfan@jhu.edu)

The authors declare that they have no competing interests.

Abstract

Recent advances in imaging-based spatially resolved transcriptomics (im-SRT) technologies now

enable high-throughput profiling of targeted genes and their locations in fixed tissues.

Normalization of gene expression data is often needed to account for technical factors that may

confound underlying biological signals. Here, we investigate the potential impact of different

gene count normalization methods with different targeted gene panels in the analysis and

interpretation of im-SRT data. Using different simulated gene panels that overrepresent genes

expressed in specific tissue anatomical regions or cell types, we find that normalization methods

that use scaling factors derived from gene counts differentially impact normalized gene

expression magnitudes in a region- or cell type-specific manner. We show that these

normalization-induced effects may reduce the reliability of downstream differential gene

expression and fold change analysis, introducing false positive and false negative results when

compared to results obtained from gene panels that are more representative of the gene

expression of the tissue's component cell types. These effects are not observed without

normalization or when scaling factors are not derived from gene counts, such as with cell volume

normalization. Overall, we caution that the choice of normalization method and gene panel may

impact the biological interpretation of the im-SRT data.

Keywords

Normalization, scaling factor, spatial transcriptomics, differential expression

Introduction

Imaging-based spatially resolved transcriptomics (im-SRT) represent a new set of technologies that enable the high-throughput detection of targeted RNA species in fixed tissues [1]–[3]. While im-SRT data is generally acquired at a single-molecule resolution, it is often aggregated into single-cell resolution by counting RNA molecules within cell regions identified by cell segmentation [4]–[7]. This achieves single-cell resolution spatially resolved gene expression profiling to enable the interrogation of cellular function and organization within tissues to enhance our understanding of cellular spatial microenvironments in biological processes such as tissue development, disease progression, and drug response [1], [2], [8].

In order to discover biological insights, normalization of gene expression data is often needed to account for technical differences that may confound underlying biological signals [4], [9], [10]. Common normalization methods for gene expression count data include scaling by a sample's total detected gene counts (also called size factor normalization, relative counts normalization, and library size normalization). This normalization approach is commonly used in the analysis of single-cell RNA-sequencing (scRNA-seq) data to account for technical differences in RNA capture rates between cells and has also been applied to im-SRT data [11]. Alternative normalization methods such as DESeq2 and TMM consider differences in gene expression composition between samples. These normalization approaches are commonly used in analysis of bulk RNA-seq data [9], [10]. Specific to im-SRT data, volume or cell area normalization has been previously applied to account for partially imaged cell volumes [12].

Given the capacities of current im-SRT technologies, a few hundred unique gene species can

typically be measured simultaneously in a given experiment. Because gene detection probes are

designed based on a target gene's sequence, these gene species must be pre-selected as part of a

gene panel in the experimental design [2], [4]. These gene panels may be designed with the

intention of spatially profiling cell-types in a tissue of interest and may thus include genes that

are consistently expressed in and/or are canonical markers of cell types of interest based on

domain knowledge or informed by scRNA-seq [12]–[15]. In recent years, commercial options

for im-SRT have become available, where off-the-shelf gene panels with probes for a pre-

selected set of genes can be used. These gene panels may focus on specific cell-types such as

immune cells or neurons, or specific pathways and processes [16]. As such, these gene panels

may be skewed to enable sensitive detection and characterization of the heterogeneity of cell

types of interest, potentially at the expense of sensitivity to other cell types present in the tissue.

Here, we investigate how different gene count normalization procedures may impact downstream

transcriptomic analyses for im-SRT data with different gene panels. Ideally, biological conclusions

drawn from im-SRT studies in the same biological system would be robust to experimental design

choices such as gene panel composition. For example, if a gene is differentially expressed in a

particular region or cell type in a given biological system probed with one gene panel, it should

also be differentially expressed given a different gene panel, if that the gene is in both panels.

However, we show using im-SRT data with simulated skewed gene panels that different

normalization methods may introduce false positive and false negative results when identifying

significantly differentially expressed genes as well as opposite fold change directions (Figure 1A).

We further demonstrate that these concerns generalize across im-SRT technologies that have

different technical features, such as gene panel size and capture efficiency. Finally, using simulated

gene panels of different sizes, we find that these concerns can be mitigated as gene panels increase

in size. Overall, we caution that the choice of normalization method and gene panel may impact

the biological interpretation of the im-SRT data.

Results

To investigate the potential impact of normalizing im-SRT data with different gene panels, we

simulated im-SRT datasets with different gene panels. We first focused on a multiplexed error-

robust fluorescent *in situ* hybridization (MERFISH) im-SRT dataset of a coronal section of the

mouse brain that originally profiled 483 genes representative of diverse cell-types (Supp Fig 1A)

[5], [17].

As a proof of principle, we created different gene panels that could allow us to better characterize

different mouse brain regions (Supp Fig 1B-E). Many brain regions are enriched for specific cell

types and in turn differentially express specific genes when compared to other brain regions. To

achieve this, we annotated mouse brain regions by aligning and transferring region annotations

from the Allen Brain Atlas using STalign [18], [19]. Then, for a brain region of interest, we

randomly sample 100 genes that are significantly overexpressed in this region, thereby simulating

a 100-gene gene panel skewed to our brain region of interest. For both the original full gene panel

and the region-skewed panel, we compared library size, DESeq2, TMM, and cell volume

normalization, in addition to no normalization.

Skewed gene panels with different normalization methods result in region-specific effects on

scaling factors and normalized gene expression magnitudes in im-SRT data

To investigate the potential impact of normalization procedures on downstream transcriptomic

analyses, we first compared the scaling factors derived from the full and skewed gene panels for

each normalization method (Figure 1B). Given that these scaling factors are applied to detected

cell gene counts to obtain the normalized gene expression magnitudes used in downstream

analysis, we expect that differences in the scaling factors between gene panels will result in

differences in the normalized gene expression magnitudes, which may in turn impact downstream

analyses. For normalization methods where scaling factors are derived based on gene count

information (library size, DESeq2, TMM), we further expect that the scaling factors will differ

depending on the gene panel used.

Under library size normalization, we found scaling factors to generally be smaller with the skewed

gene panel than with the full gene panel. This is expected given that the skewed gene panel includes

fewer genes than the full gene panel. However, we note that the shift in scaling factors is less

pronounced for some cells than others depending on brain region. Expectedly, given a gene panel

skewed towards genes enriched in the ventricle region, cells in the ventricle region have scaling

factors that are more consistent in magnitude compared to scaling factors derived from the full

gene panel than cells outside the ventricle region. We likewise find that these results are consistent

for gene panels designed for other brain regions including the habenula, fiber tracts, and dentate

gyrus (Supp Fig 2). Similar to the ventricle region, these other brain regions are enriched for

specific cell types and in turn differentially express specific genes when compared to other brain

regions. As such, differences in scaling factors for regions outside of those to which a gene panel

is skewed are driven by the underlying differences in cell type composition and resulting gene

expression differences. We also see this region-specific effect with DESeq2 and TMM

normalizations. These results indicate that in im-SRT data with region-skewed gene panels,

normalization using scaling factors derived based on gene counts may introduce region-specific

impacts on downstream analyses depending on the normalization procedure used. Alternatively,

as cell volume is independent of the number of gene counts detected in a cell, the scaling factor is

the same with the skewed and full gene panels and therefore does not result in region-specific

impacts on downstream analyses.

Next, we normalized gene counts using these various scaling factors. We evaluated how consistent

normalized gene expression magnitudes are when using the skewed gene panel compared to

normalized gene expression magnitudes with the full gene panel. To do this, we computed the

correlation in normalized gene expression magnitudes between the skewed panel and the full gene

panel (Figure 1C). Normalized gene expression magnitude correlations are the lowest for TMM

normalization, ranging from around 0.5 to 0.98 for different genes. Library size normalization

resulted in similarly low correlations between normalized gene expression magnitudes between

skewed and full gene panels. DESeq2 resulted in comparatively higher correlations, all above 0.9.

Volume normalization and no normalization resulted in perfectly correlated normalized gene

expression magnitudes, as expected given that scaling factors are unchanged between different

gene panels. We again find that these results are consistent for gene panels skewed towards other

brain regions (Supp Fig 3).

To further understand the source of differences in normalized gene expression magnitude

correlations between skewed and full gene panels with different normalization methods, we looked

at the normalized gene expression magnitudes for select highly expressed genes (Figure 1D).

Notably, given a gene panel skewed towards genes enriched in the ventricle region, under library

normalization, normalized gene expression magnitudes are similar between the full and the skewed

gene panels for cells in the ventricle region, as expected given similar scaling factors. However,

for other regions, normalized gene expression magnitudes are higher with the skewed gene panel

compared to the full gene panel. This is consistent with the smaller scaling factors for cells in these

regions with the skewed gene panel compared to the full gene panel. As with the scaling factors,

region specific differences between normalized counts in skewed and full gene panels are less

pronounced with the DESeq2 and TMM normalizations than with library normalization. These

examples show that the region-specific impacts on scaling factors when using region-skewed gene

panels with count-based normalizations can in turn differentially impact gene expression

magnitudes in a region-specific manner resulting in distortions in the relationship between

normalized gene expression magnitudes across cells.

Skewed gene panels with different normalization methods lead to region-specific biases in

differential gene expression results

Next, to evaluate the impact of these different normalization approaches with different gene panels

on downstream analysis, we applied these scaling factors and performed differential expression

analysis. To do this, we compared the p-value of a Wilcoxon rank sum test of differential

expression as well as the gene fold change between cells in each brain region and all other regions

based on normalized gene expression magnitudes achieved with the skewed gene panel versus the

full gene panel for each normalization method. When using library size normalization, ventricle-

specific genes have less significant p-values with the ventricle-skewed gene panel compared to the

full gene panel. In contrast, other gene p-values are similar when using the ventricle-skewed gene

panel and the full gene panel. This region-specific difference is less pronounced with TMM and

DESeq2 normalization. Under no normalization and volume normalization, the p-values are equal

between the skewed gene panel and the full gene panel as the normalized gene expression

magnitudes are the same (Figure 2A).

Likewise, with respect to gene expression fold changes, under library normalization, ventricle-

specific genes have lower $\log_2$ fold changes when using the skewed gene panel compared to the

full gene panel with some genes having negative log fold changes in the skewed gene panel and

positive $\log_2$ fold changes in the full gene panel. Genes not in the ventricle region specific gene

panel have similar fold changes between the two panels. These region-specific effects are also seen

with TMM normalization but are less pronounced and are not seen with DESeq2 normalization

(Figure 2B). As before, under no normalization and volume normalization, the fold changes are

equal between the skewed gene panel and the full gene panel as the normalized counts are the

same.

To understand the source of differences in $\log_2$ fold changes between skewed and full gene panels

after library size normalization, we looked at the normalized gene expression magnitudes for

individual highly expressed genes (Figure 2C). We observe that for the gene *Gpr75*, library size

normalization with the full gene panel results in normalized gene expression magnitudes that are

between cells in the ventricle region and cells in other regions. In contrast, library size

normalization with the ventricle skewed gene panel results in normalized gene expression

magnitudes in cells in the ventricle region that are lower in cells in the ventricle region compared

to cells in other regions in the data. This results in a negative $\log_2$ fold change with the ventricle

skewed gene panel that was not observed with the full gene panel. This region-specific impact is

consistent with the differences in library size scaling factors resulting from the skewed gene panel,

where cells in the ventricle region have larger scaling factors than those in other regions because

of their higher expression of genes included in the gene panel. Conversely, we observe that for the

gene *Htr2c*, library size normalization with the full gene panel results in normalized gene

expression magnitudes that are lower in cells in the habenula region compared to cells in other

regions in the data. In contrast, library size normalization with the ventricle skewed gene panel

results in normalized gene expression magnitudes in cells in the habenula region that are more

similar in magnitude to those in cells in other regions. This results in a $\log_2$ fold change that is

positive with the ventricle skewed gene panel compared to a negative $\log_2$ fold change with the

full gene panel. These examples show that the region-specific impacts on scaling factors when

using region-skewed gene panels with library size normalization can result in distortions in

normalized gene expression distributions, which in turn lead to inconsistent fold change and

differential gene expression results with different gene panels. Again, results are consistent for

gene panels designed for other brain regions (Supp Fig 4-5). These results show that the region-

specific effects of gene count-based normalization in im-SRT with skewed gene panels may result

in false negatives in differential gene expression testing, as well as fold changes with opposite

directions.

To further investigate if these observations extend to im-SRT data acquired by other technologies,

we performed the same analyses with another im-SRT dataset of a coronal section of the mouse

brain assayed by STARmap PLUS [20] using three gene panels skewed towards ventricles, fiber

tracts, and dentate gyrus respectively and observed similar results to those seen with the MERFISH

data (Supp Fig 6-8). This suggests that region-specific differences in transcriptomics analysis

results using skewed gene panels also applies to im-SRT data from other imaging-based spatial

transcriptomics technologies.

Normalization-induced biases can be mitigated with more representative or larger gene panels

Next, we investigated if impacts on the reliability of differential gene expression testing and fold

change analysis resulting from region-skewed gene panels can be mitigated with gene panels that

are more representative of the gene expression of component regions in the tissue. To do this, we

simulated a non-skewed gene panel of the same size as the region-specific gene panels used in the

previous simulations and compared scaling factors, and results of differential expression testing

and fold change analysis between the full 483-gene gene panel and non-skewed 100-gene gene

panel after normalization with the four normalization methods (Supp Fig 9). We first computed

scaling factors for each of the four normalization methods and compared to those computed with

the full gene panel. As expected, given the smaller gene panel size, library size scaling factors

were smaller with the non-skewed gene panel than with the full gene panel. In contrast to the

skewed gene panel, the decrease in library size scaling factor affected all cells similarly and did

not exhibit a region-specific pattern. Similar results were observed for DESeq2 and TMM. As

before, scaling factors for volume normalization are independent of gene panel and therefore the

same between the non-skewed and full gene panels. We then performed differential expression

testing and found that both p-values and fold changes were well correlated between the non-

skewed and the full gene panels for all genes. These results indicate that region-specific impacts

on differential gene expression testing and fold change analysis resulting from count-based

normalization can be mitigated by designing gene panels that are more representative of the gene

expression of component regions and cell types in the tissue.

We anticipate that as im-SRT technologies improve, more gene species will become

simultaneously detectable in a single experiment, reducing the need to design smaller region- or

cell type-specific gene panels. We therefore sought to investigate the extent to which cell type-

specific biases in transcriptomics analysis results will be observed with larger gene panels. To

evaluate the impact of the size of the skewed gene panel on the extent of cell type or region-specific

effects observed, we used full-transcriptome scRNA-seq data collected from sorted peripheral

blood mononuclear cells [21] to simulate skewed gene panels of different sizes. Specifically, we

simulated gene panels of 50, 100, 500, 1000, 5000 genes overexpressed in monocytes in addition

to a panel including all genes that passed quality filters. We also simulated gene panels of the same

sizes consisting of randomly selected genes. When using library size normalization, cell type-

specific differences are observed in the skewed gene panels of all sizes. However, the extent of

these differences varies depending on the size of the panel. For the monocyte specific gene panel,

larger differences are observed with the smaller gene panels compared to the larger panels (Figure

3). Again, the random gene panels that are representative of all cell-types do not show these cell

type-specific differences in results of transcriptomic analyses after normalization with either

library size, TMM, or DESeq2. Overall, these results indicate that when normalization scaling

factors are derived based on gene counts, region- or cell type-specific impacts on downstream

analysis may persist with gene panels as large as 5000 genes, though the extent of these impacts

will depend on the normalization method used as well as the degree of region- or cell type-specific

gene expression overrepresentation in the gene panel.

Discussion

Here, we investigate the impact of different normalization methods with different gene panels in

the analysis of im-SRT data. In particular, we simulated gene panels that overrepresent the gene

expression of specific component tissue anatomical regions or cell types. Given im-SRT data

with these skewed gene panels, we find that normalization methods that use count scaling factors

derived from detected gene counts such as library size, DESeq2, or TMM differentially impact

normalized gene expression magnitudes in component regions or cell types. We show that these

region or cell type-specific effects may reduce the reliability of downstream differential gene

expression results when compared to results obtained from gene panels that are more

representative of the gene expression of the tissue's component cell types.

Notably, these gene count-based normalization methods were developed for use with full-

transcriptome RNA sequencing data and rely on assumptions that may be violated in targeted

transcriptome profiling such as in im-SRT data, particularly with skewed gene panels. For

example, library size normalization assumes that there are no genes that are only highly

expressed in one subset of samples; that is, all samples have similar total gene expression [9],

[10]. This assumption is inherently violated by designing gene panels to include canonical cell

type markers for cell types of interest. Other normalization methods that attempt to account for

such compositional differences in gene expression, such as those used in the DESeq2 and edgeR

packages, rely on other assumptions. For example, DESeq2 assumes that fewer than half of

genes in the data are differentially expressed between samples. Similarly, by choosing one

sample to be a reference, TMM also assumes that most measured genes are not differentially

expression between samples [10]. This is often not the case in im-SRT gene panels as they are

often selected by identifying differentially expressed genes from prior scRNAseq data. Although

our simulated gene panels were intentionally skewed for demonstration purposes, the approach

by which these genes were selected can be used to design gene panels for im-SRT experiments,

which would result in unintentionally skewed gene panels.

Despite this, count-based normalization can still provide robust insights when gene panels are

well-designed to be representative of most of the component cell types in the tissue being

assayed. To this end, we recommend the use of non-spatially resolved single cell resolution, full

transcriptome profiling approaches such as scRNA-seq data to inform gene panel design. This

can be achieved by generating scRNA-seq and im-SRT datasets from the same tissue [12].

Furthermore, as collective atlasing efforts such as the Human BioMolecular Atlas Program

(HuBMAP) [22], the Human Cell Atlas [23], and others continue to generate tissue-specific gene

expression data, we anticipate these resources may help provide a more untargeted view of gene

expression across cell types in tissues of interest and enable researchers to design im-SRT gene

panels that are representative of most of tissues' component cell types.

In general, count normalization methods have been developed to account for variation across

cells due to technical factors. One of these technical factors that has motivated the development

of previous methods is variation in RNA capture rate in scRNA-seq. In im-SRT, we expect RNA

capture rate to be similar across cells in the same experiment, provided that the entire cell

volume is captured in the imaged region. For cells that lie at the boundaries of the imaged

regions, not all the cell volume will be captured resulting in a smaller gene count that is reflective

of technical factors rather than biological ones. If gene expression differences due to partial

volume capture compose a large proportion of total gene expression variation, count

normalization within individual im-SRT experiments may be needed to allow for reliable

downstream analysis. Volume or cell area normalization may be used to account for such partial

cell capture within imaged regions. Our results show that these normalization approaches may be

preferred when working with skewed gene panels as scaling factors are independent of gene

counts and as such gene panel differences do not impact downstream differential expression and

fold change analyses. However, the reliability of volume or cell area normalization is contingent

upon the accuracy of cell segmentation. Additional investigation may be needed to evaluate the

impact of cell segmentation accuracy in the analysis of normalized gene expression when

volume or cell area normalization is used.

Ultimately, normalization of im-SRT data should enable the removal of the effects of systematic

technical variation in detected gene counts. Accomplishing this goal without introducing

additional biases will help ensure that normalized gene expression magnitudes reflect underlying

biological heterogeneity to enable reliable and reproducible interpretation of im-SRT data in a

way that allows investigators to extract meaningful biological insights.

Methods

Datasets

Vizgen MERFISH Mouse Brain Receptor Map data was downloaded from the Vizgen and data

from slice 2 replicate 2 was used [17]. Brain anatomical region annotations for each cell were

obtained by transferring region annotations from the Allen Brain Atlas Common Coordinate

Framework (CCF) using STalign [18], [19]. STalign was applied using the 3D reconstructed Nissl

image from the Allen CCF atlas as a source and the MERFISH cell position data as a target.

Anatomical regions were combined to obtain coarse anatomical annotations as specified in

Supplementary Table 1. Cells in the ventricle, fiber tract, dentate gyrus, and habenula regions were

used for further analysis.

STARmap PLUS data downloaded from the Broad Single Cell Portal and data from well 11

(well11raw_expression_pd.csv.gz) was used [24]. Brain anatomical region annotations were

obtained from previous manually created annotations (well11_spatial.csv.gz) [25]. Anatomical

regions were combined to obtain coarse anatomical annotations. Fiber_tracts_A, Fiber_tracts_C,

and Fiber_tracts_D were combined to form the fiber tracts region; DG_A and DG_B formed the

dentate gyrus region; VS_A and VS_B formed the ventricle region. Cells in the ventricle, fiber

tract, and dentate gyrus regions were used for further analysis.

Single-cell RNA sequencing data was obtained from the 10X Genomics website [21]. Specifically

data from the following datasets was downloaded and combined: human CD14+ monocytes,

CD19+ B-cells, CD8+ cytotoxic T-cells, CD4+ helper T-cells, CD4+/CD45RO+ memory T-cells,

CD4+/CD25+ Regulatory T-cells, CD8+/CD45RA+ Naive Cytotoxic T-cells,

CD4+/CD45RA+/CD25- naive T-cells, and CD56+ natural killer cells.

Scaling factors and gene expression normalization

Library size scaling factors were computed as the sum of all gene counts in a cell. DESeq2 scaling

factors were computed using the estimateSizeFactorsForMatrix function from the DESeq2

package (version 1.38.0), with method=poscounts. TMM scaling factors were computed as the

inverse of the output of the calcNormFactors function from the edgeR package (version 3.40.0),

method=TMMwsp. For the MERFISH data, cell volume scaling factors were included in the

downloaded data. To obtain normalized gene expression, gene counts for each gene in each cell

were divided by that cell's scaling factor.

Differential gene expression testing and fold change analysis

To test for differential gene expression, for each gene in each region, a one-sided Wilcoxon rank

sum test was performed between cells in that region and all other cells in the data. Significantly

differentially expressed genes were identified using a p-value cutoff of 0.05 after a Benjamini-

Hochberg multiple hypothesis correction. For each gene in each region, $\log_2$ fold change (*LFC*)

was computed as follows:

$$LFC_{A,g} = log_2 \frac{\frac{1}{N_A} \sum_i^{N_A} x_{g,i}}{\frac{1}{N_{A'}} \sum_j^{N_{A'}} x_{g,j}}$$

where $LFC_{A,g}$ is the log_2 fold change for gene g between cells in region A and all other cells in the

data, $x_{g,i}$ is the normalized gene expression of gene g in cell i, N_A is the number of cells in region

A, $N_{A'}$ and is the number of cells in data in regions other than in region A.

Simulation of skewed gene panels in im-SRT data

Differential gene expression testing was performed on the normalized counts using the original,

full gene panel for each normalization method as well as no normalization. After differential gene

expression testing, significantly differentially expressed genes for a region were identified as those

genes with a p-value <0.05 after multiple hypothesis correction and an absolute log_2 fold change

>0.25 under any normalization procedure. The skewed gene panel was simulated by randomly

selecting 100 genes from those identified as significantly differentially expressed. The random

gene panel was simulated by randomly selecting 100 genes from all the genes in the data.

Simulation of skewed gene panels in scRNA-seq data

Differential gene expression testing was performed on the normalized counts using the original,

full gene panel for each normalization method as well as no normalization. Genes were ranked

according to their multiple hypothesis corrected p-value. Skewed gene panels were selected by

choosing the most significantly differentially expressed genes.

Acknowledgements

Research reported in this publication was supported by the National Institute of General Medical

Sciences of the National Institutes of Health under Award Number R35-GM142889. Figure 1A

was created using images from BioRender.com.

References

[1]   M. Asp, J. Bergenstråhle, and J. Lundeberg, "Spatially Resolved Transcriptomes—Next

416   Generation Tools for Tissue Exploration," *BioEssays*, vol. 42, no. 10, p. 1900221, 2020,

417   doi: 10.1002/bies.201900221.

418   [2]   M. Zhang *et al.*, "Spatial molecular profiling: platforms, applications and analysis tools,"

419   *Briefings in Bioinformatics*, vol. 22, no. 3, p. bbaa145, May 2021, doi:

420   10.1093/bib/bbaa145.

421   [3]   X. Zhuang, "Spatially resolved single-cell genomics and transcriptomics by imaging," *Nat

422   Methods*, vol. 18, no. 1, Art. no. 1, Jan. 2021, doi: 10.1038/s41592-020-01037-8.

423   [4]   L. Heumos *et al.*, "Best practices for single-cell analysis across modalities," *Nat Rev Genet*,

424   vol. 24, no. 8, Art. no. 8, Aug. 2023, doi: 10.1038/s41576-023-00586-w.

425   [5]   K. H. Chen, A. N. Boettiger, J. R. Moffitt, S. Wang, and X. Zhuang, "Spatially resolved,

426   highly multiplexed RNA profiling in single cells," *Science*, vol. 348, no. 6233, Apr. 2015,

427   doi: 10.1126/science.aaa6090.

428   [6]   E. Lubeck, A. F. Coskun, T. Zhiyentayev, M. Ahmad, and L. Cai, "Single-cell in situ RNA

429   profiling by sequential hybridization," *Nat Methods*, vol. 11, no. 4, Art. no. 4, Apr. 2014,

430   doi: 10.1038/nmeth.2892.

431   [7]   X. Wang *et al.*, "Three-dimensional intact-tissue sequencing of single-cell transcriptional

432   states," *Science*, vol. 361, no. 6400, Jul. 2018, doi: 10.1126/science.aat5691.

433   [8]   R. Dries, J. Chen, N. del Rossi, M. M. Khan, A. Sistig, and G.-C. Yuan, "Advances in

434   spatial transcriptomic data analysis," *Genome Res.*, vol. 31, no. 10, pp. 1706–1718, Oct.

435   2021, doi: 10.1101/gr.275224.121.

[9] C. Evans, J. Hardin, and D. M. Stoebel, "Selecting between-sample RNA-Seq

normalization methods from the perspective of their assumptions," *Briefings in*

Bioinformatics, vol. 19, no. 5, pp. 776–792, Sep. 2018, doi: 10.1093/bib/bbx008.

[10] M.-A. Dillies *et al.*, "A comprehensive evaluation of normalization methods for illumina

high-throughput rna sequencing data analysis," *Briefings in Bioinformatics*, vol. 14, no. 6,

pp. 671–683, Nov. 2013, doi: 10.1093/bib/bbs046.

[11] C. A. Vallejos, D. Risso, A. Scialdone, S. Dudoit, and J. C. Marioni, "Normalizing single-

cell RNA sequencing data: challenges and opportunities," *Nat Methods*, vol. 14, no. 6, Art.

no. 6, Jun. 2017, doi: 10.1038/nmeth.4292.

[12] J. R. Moffitt *et al.*, "Molecular, spatial, and functional single-cell profiling of the

hypothalamic preoptic region," *Science*, vol. 362, no. 6416, Nov. 2018, doi:

10.1126/science.aau5324.

[13] L. B. Kuemmerle *et al.*, "Probe set selection for targeted spatial transcriptomics." bioRxiv,

p. 2022.08.16.504115, Aug. 17, 2022. doi: 10.1101/2022.08.16.504115.

[14] M. Zhang *et al.*, "Spatially resolved cell atlas of the mouse primary motor cortex by

MERFISH," *Nature*, vol. 598, no. 7879, Art. no. 7879, Oct. 2021, doi: 10.1038/s41586-

021-03705-x.

[15] R. Chen *et al.*, "Decoding molecular and cellular heterogeneity of mouse nucleus

accumbens," *Nat Neurosci*, vol. 24, no. 12, Art. no. 12, Dec. 2021, doi: 10.1038/s41593-

021-00938-x.

[16] "Pre-designed Xenium Gene Expression Panels - Official 10x Genomics Support," *10x*

Genomics. https://www.10xgenomics.com/support/in-situ-gene-

expression/documentation/steps/panel-design/pre-designed-xenium-gene-expression-panels (accessed Jul. 25, 2023).

[17] "Vizgen MERFISH Mouse Receptor Map." https://info.vizgen.com/mouse-brain-map (accessed Jul. 24, 2023).

[18] K. Clifton *et al.*, "Alignment of spatial transcriptomics data using diffeomorphic metric mapping." bioRxiv, p. 2023.04.11.534630, Apr. 12, 2023. doi: 10.1101/2023.04.11.534630.

[19] Q. Wang *et al.*, "The Allen Mouse Brain Common Coordinate Framework: A 3D Reference Atlas," *Cell*, vol. 181, no. 4, pp. 936-953.e20, May 2020, doi: 10.1016/j.cell.2020.04.007.

[20] H. Zeng *et al.*, "Integrative in situ mapping of single-cell transcriptional states and tissue histopathology in a mouse model of Alzheimer's disease," *Nat Neurosci*, vol. 26, no. 3, Art. no. 3, Mar. 2023, doi: 10.1038/s41593-022-01251-x.

[21] "Datasets," *10x Genomics*. https://www.10xgenomics.com/resources/datasets (accessed Jul. 29, 2021).

[22] S. Jain *et al.*, "Advances and prospects for the Human BioMolecular Atlas Program (HuBMAP)," *Nat Cell Biol*, pp. 1–12, Jul. 2023, doi: 10.1038/s41556-023-01194-w.

[23] A. Regev *et al.*, "The Human Cell Atlas," *eLife*, vol. 6, p. e27041, Dec. 2017, doi: 10.7554/eLife.27041.

[24] "Spatial Atlas of Molecular Cell Types and AAV Accessibility across the Whole Mouse Brain - Single Cell Portal." https://singlecell.broadinstitute.org/single_cell/study/SCP1830/spatial-atlas-of-molecular-cell-types-and-aav-accessibility-across-the-whole-mouse-brain (accessed Jul. 26, 2023).

[25] H. Shi *et al.*, "Spatial Atlas of the Mouse Central Nervous System at Molecular

Resolution." bioRxiv, p. 2022.06.20.496914, Jun. 22, 2022. doi:

10.1101/2022.06.20.496914.

Main Figures

Figure 1. Impact of region-skewed gene panel on normalization scaling factors and

normalized gene expression magnitudes

A. Schematic of im-SRT normalization evaluation workflow.

(1) Im-SRT data is collected, cells are segmented to derive single cell gene counts, and region

annotations are derived via manual labeling or label transfer.

(2) This yields a cell-by-gene matrix and cell region meta data. Different gene panels can be used

to investigate tissue samples from the same biological systems. Ideally, biological conclusions

would be robust to gene panel choice.

(3) Gene count normalization approaches include ones where cell-specific gene count scaling

factors are derived based on gene counts such as in library size, DESeq2, and TMM

normalization or independently from gene counts such as in cell volume or cell area

normalization. Normalized gene expression is derived by dividing detected gene counts by the

cell-specific scaling factors. Downstream analysis includes differential gene expression testing

and fold change analysis between tissue regions to identify genes with potential biological

importance and results can be impacted by the choice of gene panel and normalization method.

B. For a coronal section of the mouse brain assayed by MERFISH, scatterplots of $\log_{10}$(scaling

factors) for each cell based on different normalization methods (library size normalization,

DESeq2 normalization, TMM normalization, and cell volume normalization) for the full gene

panel versus the ventricle-skewed gene panel. Cells within the ventricle brain region are shown

in purple. Cells in the fiber tract, habenula, and dentate gyrus brain regions are in grey. Red line

indicates x=y.

C. Boxplot of Pearson correlation coefficients (*r*) across genes for normalized gene expression

magnitudes with the full gene panel versus the ventricle region-skewed gene panel across

different normalization methods.

D. Scatterplot of normalized gene expression magnitudes when normalization was performed

with the full gene panel versus the ventricle skewed gene panel for two highly expressed genes,

Htr2c and *Gpr75*. Cells within the ventricle brain region are shown in purple. Cells in the fiber

tract, habenula, and dentate gyrus regions are in grey. Red line indicates x=y.

Figure 2. Impact of region-skewed gene panel on differential expression analysis and fold

change evaluation

A. Wilcoxon rank sum test p-values for one region-vs-all differential gene expression test with

normalized gene expression from the full gene panel versus the ventricle-skewed gene panel.

Red line indicates x=y.

B. One region-vs-all gene fold change with normalized gene expression from the full gene panel

and the ventricle gene panel. Black lines indicate $0 \log_2$ fold change and grey lines indicate 0.5

$\log_2$ fold change.

C. Library size normalized gene expression magnitudes in cells in region of interest vs cells in

other regions with the full gene panel and the ventricle skewed gene panel for two representative

highly expressed genes: *Htr2c* (left) and *Gpr75* (right).

In all panels, purple indicates ventricle region-specific genes as determined by a one-region-vs-

all differential gene expression test using the full gene panel. Grey indicates habenula, fiber tract,

or dentate gyrus region-specific genes, as determined by a one-region-vs-all differential gene

expression test using the full gene panel. Red line indicates x=y.

Figure 3. Library size normalization and downstream analysis with increasing gene panel

size

A. Scaling factors with monocyte skewed gene panels of different sizes and full gene panel.

Green indicates monocyte cells and grey indicates B cells, T cells, or NK cells.

B. Wilcoxon rank sum test p-values adjusted for multiple hypothesis testing for one region-vs-all

differential gene expression tests with monocyte skewed gene panels of different sizes and full

gene panel.

C. One region-vs-all gene $\log_2$ fold change with monocyte skewed gene panels of different sizes

and full gene panel.

In B and C, green indicates monocyte specific genes in monocyte-vs-all tests and grey indicates

B cells, T cell-, or NK cell- specific genes in B cell-, T cell-, or NK cell-vs-all tests.

Supplementary Figures

Supplementary Figure 1. Gene expression of brain anatomical regions in full and simulated

MERFISH gene panel

A. Heatmap showing scaled mean gene expression of 481 genes in full MERFISH panel in the

four analyzed brain regions.

B. Heatmap showing scaled mean gene expression of the 100 genes in 100-gene gene panel

skewed towards ventricle brain region for the four analyzed brain regions.

C. Heatmap showing scaled mean gene expression of the 100 genes in 100-gene gene panel

skewed towards habenula brain region for the four analyzed brain regions.

D. Heatmap showing scaled mean gene expression of the 100 genes in 100-gene gene panel

skewed towards fiber tract brain regions for the four analyzed brain regions.

E. Heatmap showing scaled mean gene expression of the 100 genes in 100-gene gene panel

skewed towards dentate gyrus brain regions for the four analyzed brain regions.

F. Heatmap showing scaled mean gene expression of the 100 genes in 1100-gene non-skewed

gene panel for the four analyzed brain regions.

In all panels, purple, green, blue, and red indicate the ventricles, habenula, fiber tracts, and

dentate gyrus regions, respectively.

Supplementary Figure 2: Scaling factors with MERFISH gene panels skewed towards the

habenula, fiber tract, and dentate gyrus regions

A. For a coronal section of the mouse brain assayed by MERFISH, scatterplots of $\log_{10}$(scaling

factors) for each cell based on different normalization methods (library size normalization,

DESeq2 normalization, TMM normalization, and cell volume normalization) for the full gene

panel versus the habenula region-skewed gene panel. Cells within the habenula brain region are

shown in green. Cells in the fiber tract, ventricle, and dentate gyrus brain regions are shown in

grey. Red line indicates x=y.

B. Scatterplots of $\log_{10}$(scaling factors) for each cell based on different normalization methods

(library size normalization, DESeq2 normalization, TMM normalization, and cell volume

normalization) for the full gene panel versus the fiber tract region-skewed gene panel. Cells

within the fiber tract brain regions are shown in blue. Cells in the habenula, ventricle, and dentate

gyrus brain regions are shown in grey. Red line indicates x=y.

C. Scatterplots of $\log_{10}$(scaling factors) for each cell based on different normalization methods

(library size normalization, DESeq2 normalization, TMM normalization, and cell volume

normalization) for the full gene panel versus the dentate gyrus region-skewed gene panel. Cells

within the dentate gyrus brain region are shown in red. Cells in the habenula, ventricle, and fiber

tract brain regions are shown in grey. Red line indicates x=y.

Supplementary Figure 3: Normalized gene expression correlations between full and region-

skewed MERFISH gene panels

A. Boxplot of Pearson correlation coefficients (*r*) across genes for normalized gene expression

magnitudes with the full gene panel versus the habenula region-skewed gene panel across

different normalization methods evaluated.

B. Boxplot of Pearson correlation coefficients (*r*) across genes for normalized gene expression

magnitudes with the full gene panel versus the fiber tract region-skewed gene panel across

different normalization methods evaluated.

C. Boxplot of Pearson correlation coefficients (*r*) across genes for normalized gene expression

magnitudes with the full gene panel versus the dentate gyrus region-skewed gene panel across

different normalization methods evaluated.

Supplementary Figure 4. Differential gene expression p-values with full and region-skewed

MERFISH gene panels

A. Wilcoxon rank sum test p-values adjusted for multiple hypothesis testing for one region-vs-all

differential gene expression test with normalized gene expression from the full gene panel versus

the habenula region-skewed gene panel. Green indicates habenula region-specific genes, as

determined by a one-region-vs-all differential gene expression test using the full gene panel.

Grey indicates ventricle, fiber tract, or dentate gyrus region-specific genes, as determined by a

one-region-vs-all differential gene expression test using the full gene panel. Red line indicates

x=y.

B. Wilcoxon rank sum test p-values adjusted for multiple hypothesis testing for one region-vs-all

differential gene expression test with normalized gene expression from the full gene panel versus

the fiber tract region-skewed gene panel. Blue indicates fiber tract region-specific genes, as

determined by a one-region-vs-all differential gene expression test using the full gene panel.

Grey indicates ventricle, habenula, or dentate gyrus region-specific genes, as determined by a

one-region-vs-all differential gene expression test using the full gene panel. Red line indicates

x=y.

C. Wilcoxon rank sum test p-values adjusted for multiple hypothesis testing for one region-vs-all

differential gene expression test with normalized gene expression from the full gene panel versus

the dentate gyrus region-skewed gene panel. Red indicates dentate gyrus region-specific genes,

as determined by a one-region-vs-all differential gene expression test using the full gene panel.

Grey indicates ventricle, fiber tract, or habenula region-specific genes, as determined by a one-

region-vs-all differential gene expression test using the full gene panel. Red line indicates x=y.

In all panels, green, blue, and red indicate habenula, fiber tract, and dentate gyrus-specific genes,

respectively.

Supplementary Figure 5. Gene log$_2$ fold change with full and region-skewed MERFISH

gene panels

A. One region-vs-all gene fold change with normalized gene expression from the full gene panel

versus the habenula region-skewed gene panel. Green indicates habenula region-specific genes,

as determined by a one-region-vs-all differential gene expression test using the full gene panel.

Grey indicates ventricle, fiber tract, or dentate gyrus region-specific genes, as determined by a

one-region-vs-all differential gene expression test using the full gene panel.

B. One region-vs-all gene fold change with normalized gene expression from the full gene panel

versus the fiber tract region-skewed gene panel. Blue indicates fiber tract region-specific genes,

as determined by a one-region-vs-all differential gene expression test using the full gene panel.

Grey indicates ventricle, habenula, or dentate gyrus region-specific genes, as determined by a

one-region-vs-all differential gene expression test using the full gene panel.

C. One region-vs-all gene fold change with normalized gene expression from the full gene panel

versus the dentate gyrus region-skewed gene panel. Red indicates dentate gyrus region-specific

genes, as determined by a one-region-vs-all differential gene expression test using the full gene

panel. Grey indicates ventricle, fiber tract, or habenula region-specific genes, as determined by a

one-region-vs-all differential gene expression test using the full gene panel.

In all panels, black lines indicate 0 $\log_2$ fold change, grey lines indicate 0.5 $\log_2$ fold change, red

line indicates x=y.

Supplementary Figure 6. Scaling factors with full and region-skewed STARmap gene

panels

A. For a coronal section of the mouse brain assayed by STARmap, scatterplots of $\log_{10}$(scaling

factors) for each cell based on different normalization methods (library size normalization,

DESeq2 normalization, and TMM normalization) for the full gene panel versus the ventricle

region-skewed gene panel. Cells within the ventricle brain region are shown in purple. Cells in

the fiber tract and dentate gyrus brain regions are shown in grey.

B. Scatterplots of $\log_{10}$(scaling factors) for each cell based on different normalization methods

(library size normalization, DESeq2 normalization, and TMM normalization) for the full gene

panel versus the fiber tract region-skewed gene panel. Cells within the fiber tract brain regions

are shown in blue. Cells in the ventricle and dentate gyrus brain regions are shown in grey.

C. Scatterplots of $\log_{10}$(scaling factors) for each cell based on different normalization methods

(library size normalization, DESeq2 normalization, and TMM normalization) for the full gene

panel versus the dentate gyrus region-skewed gene panel. Cells within the dentate gyrus brain

region are shown in red. Cells in the ventricle and fiber tract brain regions are shown in grey.

In all panels, red line indicates x=y.

Supplementary Figure 7. Differential gene expression p-values with full and region-skewed

STARmap gene panels

A. Wilcoxon rank sum test p-values adjusted for multiple hypothesis testing for one region-vs-all

differential gene expression test with normalized gene expression from the full gene panel versus

the ventricle region-skewed gene panel. Purple indicates ventricle region-specific genes, as

determined by a one-region-vs-all differential gene expression test using the full gene panel.

Grey indicates fiber tract or dentate gyrus region-specific genes, as determined by a one-region-

vs-all differential gene expression test using the full gene panel.

B. Wilcoxon rank sum test p-values adjusted for multiple hypothesis testing for one region-vs-all

differential gene expression test with normalized gene expression from the full gene panel versus

the fiber tract region-skewed gene panel. Blue indicates fiber tract region-specific genes, as

determined by a one-region-vs-all differential gene expression test using the full gene panel.

Grey indicates ventricle or dentate gyrus region-specific genes, as determined by a one-region-

vs-all differential gene expression test using the full gene panel.

C. Wilcoxon rank sum test p-values adjusted for multiple hypothesis testing for one region-vs-all

differential gene expression test with normalized gene expression from the full gene panel versus

the dentate gyrus region-skewed gene panel. Red indicates dentate gyrus region-specific genes,

as determined by a one-region-vs-all differential gene expression test using the full gene panel.

Grey indicates ventricle or fiber tract region-specific genes, as determined by a one-region-vs-all

differential gene expression test using the full gene panel.

In all panels, red line indicates x=y.

Supplementary Figure 8. Gene $\log_2$ fold change with full and region-skewed STARmap

gene panels

A. One region-vs-all gene $\log_2$ fold change with normalized gene expression from the full gene panel and the ventricle region-skewed gene panel. Purple indicates ventricle region-specific genes, as determined by a one-region-vs-all differential gene expression test using the full gene panel. Grey indicates fiber tract or dentate gyrus region-specific genes, as determined by a one-region-vs-all differential gene expression test using the full gene panel.

B. One region-vs-all gene $\log_2$ fold change with normalized gene expression from the full gene panel versus the habenula region-skewed gene panel. Blue indicates fiber tract region-specific genes, as determined by a one-region-vs-all differential gene expression test using the full gene panel. Grey indicates ventricle or dentate gyrus region-specific genes, as determined by a one-region-vs-all differential gene expression test using the full gene panel.

C. One region-vs-all gene $\log_2$ fold change with normalized gene expression from the full gene panel versus the dentate gyrus region-skewed gene panel. Red indicates dentate gyrus region-specific genes, as determined by a one-region-vs-all differential gene expression test using the full gene panel. Grey indicates ventricle or fiber tract region-specific genes, as determined by a one-region-vs-all differential gene expression test using the full gene panel.

In all panels, black lines indicate 0 $\log_2$ fold change, grey lines indicate 0.5 $\log_2$ fold change, red line indicates x=y.

Supplementary Figure 9. Non-skewed MERFISH gene panel

A. For a coronal section of the mouse brain assayed by MERFISH, scatterplots of $\log_{10}$(scaling factors) for each cell based on different normalization methods (library size normalization, DESeq2 normalization, TMM normalization, and cell volume normalization) for the full gene panel versus the non-skewed 100-gene gene panel. Cells within the ventricle, habenula, fiber

tract, and dentate gyrus brain regions are shown in purple, green, blue, and red, respectively. Red

line is x=y.

B. Pearson correlation coefficient (r) distribution across genes for normalized gene expression

magnitudes from the full gene panel versus the non-skewed gene panel across different

normalization methods evaluated.

C. Wilcoxon rank sum test p-values adjusted for multiple hypothesis testing for one region-vs-all

differential gene expression test with normalized gene expression from the full gene panel versus

the non-skewed 100-gene gene panel. Purple, green, blue, and red indicate ventricle, habenula,

fiber tract, and dentate gyrus region-specific genes, respectively, as determined by a one-region-

vs-all differential gene expression test using the full gene panel. Red line indicates x=y.

D. One region-vs-all gene fold change with normalized gene expression from the full gene panel

versus the non-skewed 100-gene gene panel. Purple, green, blue, and red indicate ventricle,

habenula, fiber tract, and dentate gyrus region-specific genes, respectively, as determined by a

one-region-vs-all differential gene expression test using the full gene panel. Black lines indicate

0 $\log_2$ fold change, grey lines indicate 0.5 $\log_2$ fold change, red line indicates x=y.

Figure 1

Figure 2

A   Differential expression -log$_{10}$(p-value) correspondence

Library Size   DESeq2   TMM   Cell Volume   No Normalization

B   log$_2$(fold change) correspondence

Library Size   DESeq2   TMM   Cell Volume   No Normalization

C

Htr2c

Skewed gene panel
log$_2$(fold change): 0.85
adjusted p-value: 1.72e-22

Full gene panel
log$_2$(fold change): -0.92
adjusted p-value: 2.57e-21

Gpr75

Skewed gene panel
log$_2$(fold change): -0.73
adjusted p-value: 0.35

Full gene panel
log$_2$(fold change): 0.87
adjusted p-value: 1.23e-08

Figure 3

Supplementary Figure 1

A Original
gene panel

B Ventricle region-
skewed gene panel

C Habenula region-
skewed gene panel

D Fiber tract region-
skewed gene panel

E Dentate gyrus region-
skewed gene panel

F Non-skewed
gene panel

Genes

Regions

Ventricles   Habenula   Fiber tracts   Dentate gyrus

Mean log$_{10}$(gene counts+1) Z−score

Supplementary Figure 2

Supplementary Figure 3

A   **Habenula region-skewed gene panel**

B   **Fiber tract region-skewed gene panel**

C   **Denate gyrus region-skewed gene panel**

Supplementary Figure 4

A   Habenula region-skewed gene panel

B   Fiber tract region-skewed gene panel

C   Dentate gyrus region-skewed gene panel

Supplementary Figure 5

A

 Habenula region-skewed gene panel

B Fiber tract region-skewed gene panel

C Dentate gyrus region-skewed gene panel

Supplementary Figure 6

A Ventricle region-skewed gene panel

B Fiber tract region-skewed gene panel

C Dentate gyrus region-skewed gene panel

Supplementary Figure 7

A

<div align="center">Ventricle region-skewed gene panel</div>

B

<div align="center">Fiber tract region-skewed gene panel</div>

C

<div align="center">Dentate gyrus region-skewed gene panel</div>

Supplementary Figure 8

A   Ventricle region-skewed gene panel

B   Fiber tract region-skewed gene panel

C   Dentate gyrus region-skewed gene panel

Supplementary Figure 9