White Matter Microstructural Plasticity Associated with Educational Intervention in Reading Disability

Steven L. Meisler[1,2*] (smeisler@g.harvard.edu, ORCID: 0000-0002-8888-1572)
John D. E. Gabrieli[1,2,3] (gabrieli@mit.edu, ORCID: 0000-0003-1158-5692)
Joanna A. Christodoulou[2,3,4] (jchristodoulou@mghihp.edu, ORCID: 0000-0001-8167-8021)

1: Program in Speech and Hearing Bioscience and Technology, Harvard Medical School, 260 Longwood Avenue, Boston, MA 02115, USA;
2: Department of Brain & Cognitive Sciences, Massachusetts Institute of Technology, 43 Vassar Street, Cambridge, MA 02139, USA;
3: McGovern Institute for Brain Research, 43 Vassar Street, Cambridge, MA 02139, USA;
4: Department of Communication Sciences and Disorders, MGH Institute of Health Professions, 36 First Ave, Charlestown, MA 02129, USA

*: Corresponding author (smeisler@g.harvard.edu)

Abstract

Children's reading progress typically slows during extended breaks in formal education, such as summer vacations. This stagnation can be especially concerning for children with learning disabilities, such as dyslexia, because of the potential to exacerbate the skills gap between them and their peers. Reading interventions can prevent skill loss and even lead to appreciable gains in reading ability during the summer. White matter connections enable communication among the brain regions required for proficient reading. Longitudinal neuroimaging studies relating intervention response to white matter microstructural changes are scarce but can reveal educationally relevant insights into rapid learning-driven brain plasticity. We collected reading scores and diffusion-weighted images at the beginning and end of summer for 41 children with reading disabilities who had completed either 1st or 2nd grade. Children were randomly assigned to receive an intensive reading intervention (n = 26) or a waiting control group (n = 15), enabling us to analyze how white matter properties varied across a wide spectrum of skill development and regression trajectories. On average, the intervention group had larger gains in reading compared to the non-intervention group, who declined in reading scores. Improvements on a proximal (but not distal) reading measure were associated with decreases in mean diffusivity within core reading brain circuitry (left arcuate fasciculus and left inferior longitudinal fasciculus) and increases in fractional anisotropy in the left corticospinal tract. Our findings suggest that responses to intensive reading instruction are related predominantly to white matter plasticity in tracts most associated with reading.

Keywords: DWI; White Matter; Reading; Dyslexia; Intervention; Plasticity

1. Introduction

Reading disabilities are the most common learning disability (Shaywitz, 1998), impacting as many as 20% of children (Wagner et al., 2020). Formal reading instruction begins at school entry (around 6 years old) for most children in the United States, and readers continue developing their skills in and out of school contexts. However, during extended formal education breaks such as summer vacation, typically an 8-9 week period in U.S. schools, reading progress typically slows (Cooper et al., 1996; Entwisle et al., 1997; von Hippel et al., 2018). Extended suspension of formal schooling can exacerbate achievement gaps among vulnerable readers who do versus do not participate in reading instruction (Christodoulou et al., 2017) as well as between vulnerable readers and their typically reading peers, as was observed during COVID-19 school disruptions (Kuhfeld et al., 2023). Reading interventions over the summer can halt reading skill loss and even lead to appreciable gains in reading abilities for struggling readers (Christodoulou et al., 2017; Donnelly et al., 2019).

Fluent reading is enabled by coordination of a network of regions in the brain. This system is typically left-lateralized (Murphy et al., 2019), consistent with the frequent left-lateralization of language processing (Enge et al., 2020; Fedorenko et al., 2011; Vigneau et al., 2006). A set of white matter tracts propagate information within this network (Ben-Shachar et al., 2007). Among these tracts are the inferior longitudinal fasciculus (ILF), which connects primary visual cortices to their ipsilateral anterior temporal lobes (Herbet et al., 2018), and the arcuate fasciculus (AF), which connects ipsilateral temporal, parietal, and frontal regions (Catani et al., 2005). Surgical resection and lesion mapping studies of these tracts support their necessity for fluent reading; individuals with damage to these regions show dysfluent reading (Epelbaum et al., 2008; Herbet et al., 2018; Ng et al., 2021; Zemmoura et al., 2015). The ILF is predominantly involved in early processes of word reading, such as orthographic recognition, by carrying primary visual signals to the posterior visual word form area (VWFA; see Dehaene & Cohen, 2011) in the (typically left) ventral occipitotemporal cortex (Lerma-Usabiaga et al., 2018; Yeatman et al., 2013; Yeatman & White, 2021). The ILF's anterior temporal projections also support semantic

processing (Duffau et al., 2013; Herbet et al., 2018; Saur et al., 2008; Shin et al., 2019).

The AF's connections from the anterior VWFA to temporoparietal and frontal regions

(e.g., linking Wernicke's and Broca's area) are involved in abstracting phonological and

higher-order language representations from printed text (Catani et al., 2005; Catani &

Mesulam, 2008; Weiner et al., 2017).

Learning and memory are thought to drive long-term plasticity in white matter

(Fields, 2015; Fields et al., 2014; Sampaio-Baptista & Johansen-Berg, 2017; Xin & Chan,

2020). Such changes may manifest as alterations to axonal geometry (e.g., diameter

modulations or axonal pruning/branching), myelin remodeling driven by oligodendrocyte

proliferation and differentiation, or variations to extra-axonal glial cells and vascular

systems (Sampaio-Baptista & Johansen-Berg, 2017). White matter micro- and macro-

structural properties can be inferred *in vivo* non-invasively with diffusion-weighted imaging

(DWI; Basser et al., 1994). Longitudinal DWI studies have related skill acquisition to white

matter changes in behavior-relevant bundles in animals (Blumenfeld-Katzir et al., 2011;

Sampaio-Baptista et al., 2013) and humans (Metzler-Baddeley et al., 2017; Scholz et al.,

2009), providing evidence for DWI's utility in quantifying white matter plasticity. Most DWI

studies of plasticity have used metrics from the diffusion tensor imaging (DTI) model,

including fractional anisotropy (FA) and mean diffusivity (MD). FA measures the degree

to which water molecule movement is directionally dependent (1 - water moves only along

a single axis, 0 - water moves equally as well in all directions), while MD is related to the

magnitude of water movement in all directions. Higher FA and lower MD are thought to

indicate well-myelinated white matter (however, see the *Discussion* for limitations

surrounding these metrics). Given the high test-retest reliability of DTI measures and

modest rates of developmental change (Behler et al., 2021; Wu et al., 2022; Yu et al.,

2020), observing significant and rapid changes in these metrics is encouraging in

suggesting that white matter alterations are occurring and manifest at a level resolvable

by conventional MRI.

Reading is an appropriate and educationally relevant domain for investigations of

learning-driven neural plasticity. First, reading is a skill that has been socio-culturally

introduced too recently to be a product of evolution or natural selection pressure. Second,

reading must be explicitly taught, and highly stable measures exist to gauge reading

performance (Torgesen et al., 2012; Woodcock, 2011). Longitudinal DWI studies of long-

term reading development have shown that trajectories of white matter properties and

reading skills are significantly linked, especially in the left AF, such that increases in tract

volume (Myers et al., 2014) and FA (Roy et al., 2022; Van Der Auwera et al., 2021; Wang

et al., 2017; Yeatman et al., 2012) accompany improvements in reading among children

with diverse reading abilities, although the opposite trend for FA has also been reported

for children with reading disabilities (Yeatman et al., 2012). A comparison of illiterate and

ex-illiterate adults suggests that developing literacy is associated with higher FA in the

left AF (Thiebaut de Schotten et al., 2014). Similarly, higher FA in core reading tracts have

predicted subsequently better future reading outcomes in children (Borchers et al., 2019;

Davison et al., 2022; Gullick & Booth, 2015), as well as better reading-adjacent skills,

such as phonological awareness, among pre-readers (Saygin et al., 2013; Zuk, Yu, et al.,

2021).

In comparing populations with and without reading disabilities, there has not only

been a focus on left-hemispheric core reading tracts, which have exhibited lower FA

among pre-readers with familial risk of dyslexia (Langer et al., 2017; Vandermosten et al.,

2015) and future diagnoses of dyslexia (Vanderauwera et al., 2017), but also on their right

hemispheric homotopes. Higher FA in the right superior longitudinal fasciculus has

predicted future reading outcomes in children with dyslexia (Hoeft et al., 2011) and

longitudinal FA increases in this tract relates to positive reading development in children

with familial risk for dyslexia (Wang et al., 2017). These studies suggest the right

hemisphere may provide a compensatory mechanism in reading disabilities. This notion

is supported by a report suggesting that children with dyslexia show faster reading-related

volumetric changes in right, but not left, inferior frontal gyrus compared to typically reading

children (Phan et al., 2021). Longitudinal designs, as described above, have yielded

stronger results than analogous high-powered cross-sectional studies that have

suggested little-to-no significant relationship between DTI measures and individual

differences in reading skills (Meisler & Gabrieli, 2022a; Moreau et al., 2018; Roy et al.,

2022). Collectively, the extant literature implies that white matter infrastructure could have
a causal and dynamic relationship with reading outcomes, as opposed to being a static
genetically predisposed foundation that reflects individual differences in such outcomes.
96

While longitudinal studies of long-term reading development seem to have
converged on the importance of left-hemispheric reading circuitry in predicting and
tracking reading outcomes, neuroimaging studies focusing on short-term reading
instruction (on the order of days-to-weeks) have yielded few and mixed findings on rapid
anatomical correlates of reading remediation (as well as functional correlates – for
reviews, see Barquero et al., 2014; Braid & Richlan, 2022; Perdue et al., 2022). Huber et
al., (2018) found that decreases in MD across the brain, not limited to core reading
circuitry, were related to better reading intervention benefits across participants. A later
reanalysis of a subset of these participants found that this plasticity was likely not driven
by factors related to myelination, but rather extra-axonal characteristics (Huber et al.,
2021). One study reported better intervention responses were related to increases in FA
in the anterior left centrum semiovale (Keller & Just, 2009), a broad term for white matter
between the corpus callosum and cortical surface which is not considered a part of core
reading circuitry. Another study found decreased MD in several left hemispheric regions
after reading intervention, although right-hemispheric regions were not reported
(Richards et al., 2017). However, a different study concluded that white matter did not
change with reading intervention, but lower right-hemispheric dorsal white matter MD
prior to intervention predicted better intervention outcomes (Partanen et al., 2021). In
young pre-readers undergoing early literacy training, pre-to-post increases in FA in the
left AF and ILF were observed, but these were ultimately attributed to developmental, as
opposed to intervention-driven, processes (Economou et al., 2022). In summary, while
rapid white matter changes can be observed in a short period of time in the context of
intensive reading intervention, it is unclear whether these changes are reproducible,
domain-specific (that is, localized to tracts that typically support reading), or dissociable
from broader developmental trajectories.
122

Inconsistent findings could be driven by a variety of factors including publication
bias to report positive findings, small sample sizes, and variation in participant
characteristics, interventions, and neuroimaging acquisition and analysis protocols
(Perdue et al., 2022; Roy et al., 2022; Schilling, Rheault, et al., 2021; Schilling, Tax, et
al., 2021; Thornton & Lee, 2000). It is also a possibility that intervention-driven effects are
not robust or generalizable, reflecting unique properties of the intervention used or cohort
studied. Neuroimaging studies in general are prone to issues in replicability (Button et al.,
2013; Kharabian Masouleh et al., 2019; Marek et al., 2022; Turner et al., 2018). Notably,
the present study bears strong similarity to another reading intervention DWI study (Huber
et al., 2018), employing the same intervention curriculum among children within a similar
age-range and focusing on comparable DWI metrics. This resemblance offers a uniquely
well-suited opportunity to attempt to reproduce prior results.

In the present study, we examined changes in reading skill and white matter
properties before and after a six-week summer reading intervention among children with
reading disabilities (RD). We focused on 7 white matter tracts: the left AF and ILF as core
reading circuitry bundles, their right-sided homotopes as potential compensatory bundles,
bilateral corticospinal tracts (CST) as control bundles that are not thought to subserve
reading, and the splenium of the corpus callosum, which may support reading through
connecting bilateral visual cortices but has been shown to be microstructurally stable
during reading intervention (Huber et al., 2018). We hypothesized observing one of two
outcomes: (1) decreases in MD and/or increases in FA in all tracts (besides the splenium)
would be related to better intervention responses (e.g., Huber et al., 2018), or (2) this
effect would be localized to just the left AF, consistent with multiple studies tracking
reading development on longer time scales (Roy et al., 2022; Van Der Auwera et al.,
2021; Yeatman et al., 2012).

2. Materials and Methods

2.1. Ethics Statement

This project was approved by the Massachusetts Institute of Technology's
Committee on the Use of Humans as Experimental Subjects (protocol number:

1201004850). Informed written consent was obtained from parents or legal guardians,

while informed written assent was obtained from the participants, who were all minors.

2.2. Participants

Participants included in the present study were a subset of a larger sample, for

which reading (Christodoulou et al., 2017) and gray matter morphometric (Romeo et al.,

2018) findings have been previously reported. Forty-one participants passed all inclusion

and quality control criteria and were analyzed in the present study (see *Data Inclusion*

and Quality Control). All participants were between 7 and 9 years old and were entering

the summer having completed grades 1 or 2. Inclusion criteria included a history of

reading difficulty based on parental report and a manifestation of reading difficulty at study

enrollment. In particular, to be included in the study, participants had to have scored "At

Risk" or "Some Risk" on the Dynamic Indicators of Basic Early Literacy Skills test

(DIBELS; Good et al., 2002) and below the 25th percentile on at least 3 of the 5 following

measures: Elision and Nonword Repetition subtests from the Comprehensive Test of

Phonological Processing, 2nd Edition (CTOPP-2; Wagner et al., 1999), and the Objects,

Letters, and 2-set Letters and Numbers subtests of the Rapid Automatized Naming and

Rapid Alternating Stimulus Tests (RAN/RAS; Wolf & Denckla, 2005). Additionally,

participants had to score at or above the 16th percentile on the Matrices subtest of the

Kaufman Brief Intelligence Test, 2nd Edition (KBIT-2; Kaufman, 2004), which is a

measure of nonverbal cognitive ability. All the children were native English speakers.

Children were recruited from a local partner charter school and the Greater Boston area.

2.3. Reading Intervention

Participants were randomly assigned to either receive a reading intervention (*n* =

26) or be placed on a waiting-list (*n* = 15). Comprehensive details of the intervention have

been previously described (Christodoulou et al., 2017). Intervention participants

completed intensive reading instruction following *Seeing Stars: Symbol Imagery for*

Fluency, Orthography, Sight Words, and Spelling program (Bell, 1997). Instruction was

delivered by trained Lindamood-Bell teachers, who rotated classrooms hourly. The

program duration was 4 hours per day on 5 days per week for 6 weeks; intervention

duration totaled between 100 and 120 hours. Students received small group instruction

(3-to-5 students per group) to improve foundational reading skills including phonological

and orthographic processing. Children recruited from the local partner school received

the intervention on-site at their school (n = 9), while children recruited from the

community-at-large (n = 17) received the intervention at a dedicated space at the

Massachusetts Institute of Technology.

2.4. Outcome Measures

Standardized reading scores were collected from all participants before and after

the intervention period, regardless of whether they participated in the intervention. A

proximal reading measure of intervention response, the Symbol Imagery Test (SIT),

measured phonological and orthographic processing in reading (Bell, 2010). During the

SIT, participants briefly viewed cards with words or pseudowords for between 2 and 7

seconds and were then asked to report what they were shown. Cronbach's α values range

from .86 to .88, and the test–retest reliability is .95 (Bell, 2010). A relatively distal

composite reading index was calculated at each time point by averaging the following four

standardized reading scores: Sight Word Efficiency (SWE) and Phonemic Decoding

Efficiency (PDE) from the Test of Word Reading Efficiency, 2nd Edition (TOWRE-2;

Torgeson et al., 1999), and Word Identification (WID) and Word Attack (WA) from the

Woodcock Reading Mastery Tests, 3rd Edition (WRMT-3; Woodcock, 2011). Timed and

untimed single word reading skills were measured by SWE and WID, respectively, while

timed and untimed pseudoword reading skills were measured by PDE and WA,

respectively. For all four subtests, Form A was administered at the beginning of the study,

and Form B was administered at the end of the study to avoid practice or familiarity

effects. High alternate form reliability has been reported for standardized tests scores on

both the WRMT-3 subtests (Word ID: r = 0.93, Word Attack: r = 0.76; Woodcock, 2011)

and the TOWRE-2 subtests (SWE: r = 0.90, PDE: r = 0.92; Torgesen et al., 2012). Age-

normed scores for all tests were defined such that the population mean is 100, with a

standard deviation of 15.

2.5. Neuroimaging Acquisition

Participants were scanned at the Athinoula A. Martinos Imaging Center at the Massachusetts Institute of Technology using a 3 Tesla Siemens TimTrio scanner and standard 32 channel head coil. During each session, a T1-weighted (T1w) MPRAGE image was acquired with the following parameters: TR=2.53s, TE=1.64ms, Flip Angle=7°, 1mm isotropic voxels. A diffusion-weighted image (DWI) was acquired with the following parameters: TR=9.3s, TE=84ms, Flip Angle=90°, 2mm isotropic voxels, and 10 b0 volumes followed by 30 non-collinear directions at b=700 s/mm^2. Age-appropriate movies were shown during these scans to increase scan engagement and reduce head motion (Greene et al., 2018). Functional MRI tasks were also collected but are not discussed here. Before the first MRI session, participants were introduced to the MRI by visiting the center's pediatric mock scanner, which allows children to get acclimated with MRI noise and lying still in the machine, which improves scan compliance (de Bie et al., 2010; Gao et al., 2023).

2.6. MRI Preprocessing and Tract Segmentation

MRI preprocessing and tract segmentation were performed according to the longitudinal TRActs Constrained by UnderLying Anatomy (TRACULA) pipeline (Maffei et al., 2021; Yendiki et al., 2011, 2016), as part of *FreeSurfer* version 7.2 (Fischl, 2012; Fischl et al., 2002; Reuter et al., 2012). This method uses longitudinal anatomical priors to produce more plausible tracts compared to creating independent segmentations at each time point (Yendiki et al., 2016), as well as leverages high-quality training data to help inform tract shapes on routine-quality DWI data (Maffei et al., 2021). To achieve this, *FreeSurfer*'s longitudinal processing pipeline (Reuter et al., 2012) was run on each participant's pre and post T1w images to create an unbiased subject template image (Reuter & Fischl, 2011) using inverse consistent registration (Reuter et al., 2010). Information from this template was used to initialize several steps of the `recon-all` pipeline, such as skull-stripping and anatomical segmentation (Reuter et al., 2012).

DWI volumes from each image were aligned to the first b0 image in that scan. The b-matrix was rotated accordingly (Leemans & Jones, 2009). DWI images were corrected

for motion and eddy currents with FSL's `eddy` command (Andersson & Sotiropoulos,

2016). Information from this process was used to generate four measures of head motion

and image quality that inform the "total motion index" (Yendiki et al., 2014): mean volume-

by-volume head rotation, mean volume-by-volume head translation, proportion of slices

with signal dropout, and severity of signal dropout. The diffusion tensor was fitted using

FSL's `dtifit`. Mean diffusivity (MD) and fractional anisotropy (FA) were derived from

the tensor. A GPU-accelerated ball-and-stick model was fit for each DWI image (Behrens

et al., 2007; Hernández et al., 2013; Jbabdi et al., 2012). At each time point, a registration

was computed between the diffusion-weighted image and T1w image (native space)

using an affine boundary-based registration algorithm (Greve & Fischl, 2009). This

transformation was used to bring anatomical segmentations into DWI space. The DWI-

to-T1w and T1w-to-template registrations were multiplied to get a DWI-to-template

transformation. Information from high-resolution 7T training data (Maffei et al., 2021) was

used to estimate endpoint ROIs and pathways for white matter tracts in each participant's

template space images. These data were then brought back into the native DWI space of

each time point. The DWI ball-and-stick model and tract anatomical priors were used to

calculate the probability density of each pathway. From these, we collected the average

MD and FA from the cores of our tracts of interest (`[MD|FA]_Avg_Center`) to mitigate

concerns of noise and partial volume effects from fiber branching towards the exterior

and extremities of the bundles. These tracts included the bilateral AF, ILF, and CST, as

well as the splenium of the corpus callosum (Figure 1).

267   **Saggital**   **Axial**   **Coronal**

Figure 1: Tracts produced by TRACULA analyzed in the present study, overlaid on top of a fractional anisotropy image. Only the left hemispheric bundles are visualized for bilateral tracts. Pictured data come from a single representative participant.

2.7. Statistics and Analysis

Analyses were prepared, run, and visualized using Python packages Pandas 1.3.2 (McKinney, 2011), Statsmodels 0.13.5 (Seabold & Perktold, 2010), and Seaborn 0.12.1 (Waskom, 2021), respectively. We made a dataframe that contained the following phenotypic fields for each subject: ages at each scan (in months), sex (binary categorical factor), and reading measures at pre and post timepoints, as well as their longitudinal pre-to-post differences. For each tract, the pre and post MD and FA metrics were added to the dataframe, along with their longitudinal pre-to-post microstructural differences. The total motion indices, calculated separately for each time point, were also added to the dataframe (see *Data Inclusion and Quality Control*). We used ordinary least squares linear models to run multiple regressions, allowing us to relate reading measures to white matter microstructure while controlling for confounds.

For the primary analyses, we created models to predict pre-to-post changes in a given tract microstructural measure by the pre-to-post change in a reading measure across all participants, with nuisance regressors for sex, age at first scan, and motion indices at both timepoints. We also ran similar models using just participants in the reading intervention. As exploratory analyses to contextualize our main focus of longitudinal differences in tract microstructure, we also examined whether reading skill was related to tract FA and MD cross-sectionally at each time point across all participants. For these additional models, at a given time point, the tract metric was predicted based on the given standardized reading measure, controlling for sex, age, and motion for the given time point. Across all models, effect sizes (ΔR^2_{adj}) were calculated as the difference in adjusted R^2 coefficients between the full model and a reduced model without the reading score term of interest.

2.8. Data Inclusion and Quality Control

A total of 153 children were recruited as part of the larger overarching study. 52

participants had anatomical and DWI scans at both time points and were able to complete

the neuroimaging processing pipeline without errors. 44 of the remaining participants had

the necessary phenotypic data. As a quality assurance metric, we computed the total

motion index (TMI) as described in Yendiki et al., (2014). TMI is related to four measures:

rotation, translation, signal dropout prevalence, and signal dropout severity. For each

scan, we calculated each motion metric's difference from the study population mean for

the given time point, divided by the interquartile range of the metric. The TMI for each

scan is the cumulative sum of these calculations across the four motion metrics. 3

subjects had outliers in TMI at either time point and were excluded. For further quality

assurance, we confirmed that no remaining participant had any tract-averaged FA lower

than 0.3, which could indicate some combination of white matter disorganization and

partial volume effects from a tract branching into significant amounts of gray matter or

CSF. Thus, a total of 41 subjects (26 who received an intervention, and 15 in the non-

intervention group) were analyzed in the present study.

3. Results

3.1. Cognitive and Phenotypic Data

Phenotypic summary statistics for the participant cohort are provided in Table 1.

Of note, the intervention and non-intervention groups were matched in sex (χ^2 test, $p >$

0.7), but not handedness (χ^2 test, $p < 0.05$). However, we did not include handedness as

a regressor in our models due to a lack of evidence of handedness-related asymmetry in

white matter microstructure (López-Vicente et al., 2021). The two groups were also

matched in age, socioeconomic status, and reading performance across all subtests

cross-sectionally at each time point (two-sample *t*-test, $p > 0.1$ across all tests), with the

exception of the intervention group having a significantly higher SIT score post-

intervention (two-sample *t*-test, $p < 0.005$). Demonstrating the efficacy of the reading

intervention, the intervention group showed higher longitudinal pre-to-post differences in

the SIT and composite reading index (two-sample *t*-test, $p < 0.002$ across both tests),

driven by the non-intervention group regressing in both measures and the intervention group improving on the SIT and maintaining scores on the composite reading index (Figure 2). These results are consistent with what was observed in the larger cohort from which the present subset was derived (Christodoulou et al., 2017).

Participants	All (*n* = 41)	Intervention (*n* = 26)	Non-Intervention (*n* = 15)	Effect Size
Age at first scan (months)	94.85 (1.16)	95.00 (1.32)	94.60 (2.29)	*d* = 0.053
Sex (M/F)	26 / 15	16 / 10	10 / 5	*Φ* = 0.051
Handedness (L/R)	7 / 34	7 / 19*	0 / 15*	*Φ* = 0.345*
SES (years of parental education)	17.89 (0.41)	18.00 (0.50)	17.70 (0.75)	*d* = 0.112
SIT (Pre/Post/Diff)	89.76 (1.77) / 93.20 (1.69) / 3.439 (2.06)	88.62 (2.15) / 97.07 (1.87)* / 8.462 (2.15)*	91.73 (3.11) / 86.47 (2.53)* / -5.267 (3.21)*	*d* = 0.275 / *d* = 1.102* / *d* = 1.191*
Composite Reading Index (Pre/Post/Diff)	83.21 (1.28) / 81.11 (1.34) / -2.101 (0.957)	82.37 (1.51) / 82.34 (1.57) / -0.029 (1.27)*	84.67 (2.34) / 78.97 (2.42) / -5.694 (0.859)*	*d* = 0.280 / *d* = 0.395 / *d* = 1.102*
TOWRE SWE (Pre/Post)	83.49 (1.78) / 80.46 (1.99)	83.46 (2.23) / 80.81 (2.61)	83.53 (3.06) / 79.87 (3.11)	*d* = 0.006 / *d* = 0.073
TOWRE PDE (Pre/Post)	79.88 (1.40) / 77.63 (1.51)	78.46 (1.64) / 79.42 (1.74)	82.33 (2.48) / 74.29 (2.70)	*d* = 0.439 / *d* = 0.552
WJ Word ID (Pre/Post)	83.68 (1.45) / 82.68 (1.46)	83.50 (1.79) / 83.69 (1.89)	84.00 (2.57) / 80.93 (2.27)	*d* = 0.053 / *d* = 0.295
WJ Word Attack (Pre/Post)	86.00 (1.70) / 83.80 (1.43)	84.32 (2.06) / 85.42 (1.52)	88.80 (2.89) / 81.00 (2.84)	*d* = 0.421 / *d* = 0.489

Table 1: Cognitive and phenotypic summary statistics. Values are provided as Mean (SEM). All reading measures are age-standardized. Two-sample t-tests were used to test for differences between groups, with the exception of χ^2-tests for sex and handedness distributions. * denotes $p < 0.05$. Abbreviations: SES - Socioeconomic Status; SIT -

 Symbol Imagery Test; TOWRE - Test of Word Reading Efficiency; SWE - Sight Word

Efficiency; PDE - Phonemic Decoding Efficiency; WJ - Woodcock Johnson.

Figure 2: Changes in Composite Reading Index (left) and SIT (right) scores for

intervention (purple) and non-intervention (yellow) participants. Paired t-tests were used

to compare pre and post scores within groups, and two-sample t-tests were used to

compare scores at a given time point across groups. Significant tests (p < 0.05) are

annotated in the figure. Abbreviations: SIT - Symbol Imagery Test.

3.2. Cross-Sectional Relationships Between White Matter Microstructure and Reading

Scores

Across all participants at the beginning of the summer, lower MD ($p < 0.1$, ΔR^2_{adj} =

0.058) and higher FA ($p < 0.1$, ΔR^2_{adj} = 0.064) in the left ILF were marginally associated

with better SIT scores. Higher FA in the right ILF ($p < 0.05$, ΔR^2_{adj} = 0.106) and right CST

($p < 0.05$, ΔR^2_{adj} = 0.069), as well as lower MD in the right CST ($p < 0.05$, ΔR^2_{adj} = 0.079),

were related to better initial composite reading index scores. At the end of the summer,

better composite reading index scores were associated with higher FA ($p < 0.05$, ΔR^2_{adj}

= 0.084) and lower MD ($p < 0.05$, ΔR^2_{adj} = 0.086), in the right CST. No tract microstructure

values were associated with SIT scores at the end of the summer. No tests remained

significant at α = 0.05 with a Bonferroni factor of 28 (at each time point, 7 tracts X 2

microstructural metrics X 2 reading scores).

3.3. Relationship Between Changes in White Matter Microstructure and Reading Scores

Across all participants, pre-to-post decreases of MD in the left AF and left ILF were related to improvements in SIT scores over the summer, with MD changes in the left AF accounting for ~9% of variance in reading scores trajectories, and left ILF accounting for ~16% (Table 2, Figure 3). This effect was not present when considering the composite reading index. Decreasing splenium MD was marginally correlated with improvements in both reading measures ($p < 0.1$), accounting for ~5% of variance in each. Increasing FA in the left CST ($p < 0.05$) and to a lesser extent the left AF ($p < 0.1$) were related to improvements in SIT scores, but not the composite reading index (Figure 3, Table 3). No tests remained significant at $\alpha = 0.05$ after correction for multiple hypotheses with a Bonferroni factor of 28.

When running the same models on only the 26 participants who completed the intervention, the relationships between pre-to-post decreases in MD in the left ILF and improvement in SIT scores (Figure 3, $p < 0.05$, $\Delta R^2_{adj} = 0.238$) and between pre-to-post decreases in MD in the splenium and improvement in composite reading index scores ($p < 0.1$, $\Delta R^2_{adj} = 0.093$) remained. Additionally, there was a modest relationship between pre-to-post decreases of FA in the right ILF and improvement in SIT scores ($p < 0.1$, $\Delta R^2_{adj} = 0.118$). None of these additional models remained significant at $\alpha = 0.05$ after correction for multiple hypotheses with a Bonferroni factor of 28.

Tract	SIT Score			Composite Reading Index		
	β [95% CI]	ΔR^2_{adj}	*p*-value	β [95% CI]	ΔR^2_{adj}	*p*-value
Left AF	-0.385 [-0.746, -0.024]	0.092	0.037*	-0.219 [-0.541, 0.104]	0.024	0.177
Right AF	0.049 [-0.334, 0.431]	-0.026	0.798	0.080 [-0.249, 0.408]	-0.021	0.625
Left ILF	-0.480 [-0.839, -0.121]	0.157	0.010*	-0.079 [-0.417, 0.260]	-0.023	0.641
Right ILF	0.050 [-0.330, 0.430]	-0.026	0.792	0.104 [-0.222, 0.430]	-0.016	0.520

Left CST	-0.197 [-0.558, 0.165]	0.005	0.277	-0.073 [-0.388, 0.243]	-0.201	0.643
Right CST	-0.193 [-0.539, 0.154]	0.006	0.267	-0.156 [-0.454, 0.143]	0.002	0.298
Splenium	-0.292 [-0.624, 0.039]	0.046	0.082†	-0.265 [-0.549, 0.019]	0.054	0.066†

*Table 2: Multiple regression outcomes relating changes in age-standardized reading scores and tract MD across all participants. * denotes p < 0.05, † denotes p < 0.1. p-values are uncorrected for multiple comparisons. Abbreviations: AF - Arcuate Fasciculus; ILF - Inferior Longitudinal Fasciculus; CST - Corticospinal Tract; SIT - Symbol Imagery Test.*

Tract	SIT Score			Composite Reading Index		
	β [95% CI]	ΔR^2_{adj}	*p*-value	β [95% CI]	ΔR^2_{adj}	*p*-value
Left AF	0.292 [-0.052, 0.635]	0.045	0.094†	0.031 [-0.277, 0.339]	-0.024	0.839
Right AF	-0.063 [-0.449, 0.323]	-0.254	0.743	0.135 [-0.194, 0.464]	-0.009	0.411
Left ILF	0.180 [-0.210, 0.569]	-0.004	0.356	0.101 [-0.237, 0.439]	-0.019	0.548
Right ILF	-0.172 [-0.532, 0.188]	-0.001	0.338	-0.065 [-0.378, 0.248]	-0.021	0.677
Left CST	0.3929 [0.031, 0.755]	0.097	0.034*	0.089 [-0.242, 0.421]	-0.020	0.587
Right CST	0.155 [-0.191, 0.501]	-0.004	0.369	0.215 [-0.077, 0.507]	0.027	0.144
Splenium	0.036 [-0.340, 0.412]	-0.026	0.847	0.079 [-0.244, 0.402]	-0.020	0.621

*Table 3: Multiple regression outcomes relating changes in age-standardized reading scores and tract FA across all participants. * denotes p < 0.05, † denotes p < 0.1. p-values are uncorrected for multiple comparisons. Abbreviations: AF - Arcuate Fasciculus; ILF - Inferior Longitudinal Fasciculus; CST - Corticospinal Tract; SIT - Symbol Imagery Test.*

Figure 3: Partial regression plots relating changes in tract microstructure to changes in
standardized SIT scores. Confounds included age at first scan, sex, and motion indices
at each time point. Values on axes are residuals after accounting for nuisance regressors
in the model. Only tracts with an uncorrected p < 0.05 (denoted by *) across all
participants are shown. These include the left AF (left), ILF (middle), and CST (right). No
test reached this threshold with the composite reading index. Purple dots represent
intervention participants, and yellow does represent non-intervention participants. The
black solid line represents the best fit across all participants, and the purple dashed line
represents the best fit when considering only intervention participants. Abbreviations: AF
- Arcuate Fasciculus; ILF - Inferior Longitudinal Fasciculus; CST - Corticospinal Tract.

4. Discussion

In the present study, we investigated whether changes in white matter
microstructure were related to changes in reading skill during the summer among 41
children with reading disabilities. Reading ability trajectories varied on a wide spectrum,
including score regression (the "summer slump") and intervention-driven improvement.
We focused on 7 tracts within and outside of core reading circuitry, using two
microstructural measures (FA and MD), and two reading measures. One reading
measure, the SIT, was closely related to the intervention, while a separate composite
reading index was more distal to the intervention. We found that longitudinal decreases
in MD were related to improved SIT scores in core reading circuitry (the left AF and left
ILF). Longitudinal increases in FA in the left CST were also related to improved SIT
scores. Notably, none of these associations were present when considering the
composite reading index. We originally hypothesized seeing a relationship between white

matter microstructure and reading score changes in either just the left AF or more globally.

While neither of these hypothesized outcomes were explicitly met, the pattern of results,

particularly relating to the use of MD as done in other studies of short-term reading

intervention (Huber et al., 2018; Partanen et al., 2021), suggests that intervention effects

were strongest (and in the hypothesized direction) within core reading circuitry when

considering the reading measure most proximal to the intervention.

The design of our study most closely resembles that of Huber et al., (2018), but

with some important differences. The same intervention curriculum (*Seeing Stars*) was

used in both studies, with similar instruction hours per week, but the present study had a

6-week intervention period, as opposed to the 8 weeks in Huber et al., (2018). Notably,

students in the present study were taught in small group settings, while a 1-on-1 approach

was used in Huber et al., (2018). The longer intervention duration and more intense

individualized instruction may have been factors that led to children in that study

improving on their composite reading measure (composed of the same reading tests as

in the present study), as opposed to only maintaining scores as found in the present study.

Both studies had similar cohort sizes (41 and 43 children). However, the present study

had participants within a narrower age-range of 7-9 years compared to 7-12 years old in

Huber et al., (2018). Additionally, all participants in the present study were diagnosed with

a reading disability, while 10 children in Huber et al., (2018) were typical readers. Both

studies found that changes in MD in the left AF and ILF were related to reading outcomes

over the course of the summer. However, Huber et al., (2018) found more widespread

plasticity that did not include the splenium, while our cohort exhibited more domain-

specific plasticity that included the splenium ($p < 0.1$). Beyond the contrasts between

studies explained above, additional variation in data acquisition, processing, and

statistical techniques could have contributed to differences in results between studies

(Schilling, Rheault, et al., 2021; Schilling, Tax, et al., 2021).

It is encouraging that both the present study and Huber et al., (2018) found

intervention effects in the left AF and ILF, which provides converging lines of evidence

suggesting that intense educational instruction targets reading-relevant white matter

tracts, albeit with different findings about plasticity occurring in a broader range of tracts.
Future studies ought to address similar questions in different contexts to evaluate the
reproducibility and generalizability of these findings. Presently, there are not enough
extant studies on longitudinal neuroanatomical correlates of reading intervention (in either
gray or white matter) to perform meaningful meta-analyses. In functional MRI, a meta-
analysis of 8 studies with longitudinal neuroimaging and cognitive scores concluded that
there were no consistent locations where longitudinal changes in reading-invoked BOLD
signal and intervention response covaried (Perdue et al., 2022). However, individual
studies, including those that may have only contained one session of neuroimaging either
prior or after intervention, have found functional correlates of intervention response both
in putative reading regions as well as more globally (reviewed in Barquero et al., 2014;
Braid & Richlan, 2022; Perdue et al., 2022). One of the few studies of gray matter
morphometric correlates of intervention response was conducted on the same participant
pool as in the present study (Romeo et al., 2018). This study concluded that children who
improved exhibited significant cortical thickening in brain regions including the left middle
temporal gyrus, right superior temporal gyrus, and bilateral middle-inferior temporal
cortex, inferior parietal lobule, precentral cortex, and posterior cingulate cortex. While
some of these regions comprise left-lateralized core reading areas, others extend globally
beyond the reading network. Considering these spatially distinct patterns of results for
gray and white matter, it is unclear yet how to relate different anatomical measures of
plasticity in response to reading intervention.

In exploratory analyses, we created cross-sectional models to investigate whether
white matter microstructure and reading skills were associated at each time point. Lower
MD and higher FA in the left ILF were associated with better reading scores at the
beginning of the summer, consistent with the left ILF's critical role in supporting reading.
We also found that the right ILF and right CST microstructure had significant associations
with reading scores. Notably, this is not consistent with a previous report showing an
inverse relationship between right ILF FA and reading outcomes among children with
reading disabilities (Banfi et al., 2018), and in the present study, longitudinal
microstructural trajectories in these right-sided homotopes were not linked with reading

score changes. This might suggest that right-lateralized white matter serves as a static

compensatory agent that reflects early reading outcomes in reading disabilities (e.g.,

(Zuk, Dunstan, et al., 2021), but does not dynamically change with reading instruction.

However, given the small sample size and limited power of cross-sectional designs, this

should be interpreted with caution.

The significance of the models including the CST, both cross-sectionally and

longitudinally, were unexpected given its seeming lack of a role in reading as a primary

motor tract. Although effect sizes for these models were appreciably lower than those for

the left ILF and AF, this suggests that intervention effects could still be detected to some

extent outside of reading circuitry. The CST is not often focused in studies of reading

abilities, but one study found that volumes of bilateral CST were informative in predicting

dyslexia diagnoses in children (Cui et al., 2016). Additionally, other studies have found

that FA in the left CST predicted future phonological skills (a critical pre-reading ability) in

kindergarten (Zuk, Yu, et al., 2021), correlated with phonological processing in

preschoolers (Walton et al., 2018), and related to phonological encoding abilities in adults

with brain damage (Han et al., 2016). This suggests that the left CST could be co-opted

into reading circuitry in populations with deficient or not-fully developed language abilities.

This may deviate from the more frequent focus of compensation from right-sided reading

circuitry homotopes such as the right ILF and AF. However, this theory is consistent with

the left CST's importance in speech, evidenced by microstructural deficiencies in pre-

term children with poor oromotor outcomes (Northam et al., 2012) and stuttering

populations (Connally et al., 2014; Kronfeld-Duenias et al., 2016), and the relationship

between speech and reading disabilities (Catts, 1993; Hayiou et al., 2010). It is also a

possibility that the CST reconstructions largely intersected with the nearby corticobulbar

projections, as tractography is prone to overlaps (Schilling et al., 2022). Corticobulbar

projections innervate cranial nerves that support head and neck muscles, and properties

of corticobulbar tracts have also been related to speech and language outcomes in

preterm adolescents (Northam et al., 2019).

Associations between white matter microstructure and reading outcomes favored

proximal, but not distal, measures. The proximal outcome measure (SIT) was directly

aligned with the intervention program and is the metric that one would predict to be most

impacted by the intervention. We expected but did not find statistically significant

associations between white matter microstructure and the distal reading composite that

included four single word reading measures (timed and untimed, real and pseudo-words).

Previous research on this intervention reported the largest effect sizes (Cohen's *d*) for the

SIT measure (1.32), but more modest effect sizes for distal single word reading measures

that create the reading composite score (Untimed word reading (WRMT WID): 0.96;

untimed pseudoword reading (WRMT WA): 0.87; timed word reading (TOWRE SWE):

0.19; timed pseudoword reading (TOWRE PDE): 1.08; Christodoulou et al., 2017). The

association between plasticity in white matter microstructure and change in SIT scores

may reflect this more substantial change, perhaps reflecting a minimal threshold of

intervention impact that ties to structural plasticity.

The pattern of results in the present study favoring stronger effects in MD than FA

imply that the microstructural changes accompanying reading development are related to

extra-axonal factors, such as neurite density and CSF volume (Beaulieu, 2002; Genc et

al., 2017), as opposed to axonal factors such as myelination, orientation coherence, and

axonal density (Friedrich et al., 2020). This is consistent with other DWI studies of reading

intervention (Huber et al., 2018, 2021). However, multimodal research at various spatial

and temporal resolutions will need to be reconciled to perform the nontrivial task of

ascribing such changes to biophysical mechanisms (Jelescu et al., 2020). While higher

FA and lower MD are often thought to reflect more "healthy" white matter, these metrics

are biologically unspecific due to the variety of factors that can influence diffusion of water

in voxel-sized regions and the confounding impact of crossing fibers (De Santis et al.,

2014; Jones et al., 2013), which can impact as many as 90% of white matter voxels

(Behrens et al., 2007; Jeurissen et al., 2013). FIber-specific measures such as

quantitative anisotropy (Yeh et al., 2013) and fixel-based metrics (Raffelt et al., 2017),

and multicompartmental models such as NODDI (Zhang et al., 2012), can provide higher

biological specificity and have shown promise in better resolving brain-behavior

relationships in studies of reading abilities (Koirala et al., 2021; Meisler & Gabrieli, 2022b;

Sihvonen et al., 2021). Unfortunately, the low angular resolution and weak single-shelled

diffusion weighting of the present DWI acquisition scheme were not well-suited for these

more novel approaches (Genc et al., 2020), effectively limiting us to using DTI metrics.

However, future work should design data acquisition protocols to use these techniques in

longitudinal study designs.

Our results should be considered in the context of additional limitations. The small

amount of within-participant data (two timepoints) precluded us from running more

numerically sophisticated models, such as linear mixed-effect models as in Huber et al.,

(2018). This also limited our ability to infer when microstructural changes occurred over

the course of the intervention and characterize the temporal relationship between

changes in reading and tract properties (in other words, whether tract microstructural

changes preceded changes in reading scores, or vice-versa). To address these points,

future longitudinal studies of reading intervention should strive to contain at least 3

sessions of data collection (King et al., 2018). Our relatively small sample size of 41

reflects the challenges of collecting acceptable quality cognitive and multi-modal MRI data

in children with learning disabilities, which are particularly amplified for longitudinal

studies (Davis et al., 2022). Even with these limitations, we found that white matter

microstructural plasticity, predominantly in core reading circuitry, was related to changes

in reading abilities over the summer in the context of short-term intensive educational

intervention.

Data and Code Availability

Due to language used in the consenting process, we are not permitted to publicly

share subject MRI images. Images may be privately distributed upon reasonable request.

We share a CSV containing all necessary data to replicate the present results, as well as

the code to recreate the analyses and figures. All instructions and code for processing

data   and   running   the   statistical   analyses   can   be   found   at

https://github.com/smeisler/Meisler_ReadingInt_DWI. To   execute   the   *FreeSurfer*

workflows, we ran a Docker container containing *FreeSurfer* 7.2 and *FSL* 6.0.4 with

Singularity (3.9.5) (Kurtzer et al., 2017). The container can be collected with either `docker pull amirro/tracula:latest` or `singularity build tracula_container.img docker://amirro/tracula:latest`. Development of these software may introduce improvements and bug fixes that should be used in future research, so we encourage using the latest stable releases.

Author Contributions

Steven L. Meisler: Conceptualization, Formal analysis, Funding acquisition, Methodology, Software, Visualization, Writing - original draft, Writing - review & editing. **John D.E. Gabrieli:** Conceptualization, Funding acquisition, Supervision, Writing - review & editing. **Joanna A. Christodoulou:** Conceptualization, Funding acquisition, Investigation, Supervision, Writing - review & editing.

Funding

This work was supported by the National Institutes of Health (SLM: 5T32DC000038 and 1F31HD111139; JDEG and JAC: 1R01HD106122; JAC: 1R15HD102881), the Halis Family Foundation, and Reach Every Reader, a grant supported by the Chan Zuckerberg Foundation.

Declaration of Competing Interest

The authors declare no competing interests.

Acknowledgments

We thank all the participants and their families for volunteering their time to participate in the study. We also thank Atsushi Takahashi, Sheeba Arnold Anteraper, and Steven Shannon at the Athinoula A. Martinos Imaging Center at the McGovern Institute for Brain Research, Massachusetts Institute of Technology for technical assistance. We thank our colleagues and team members Rachel Romeo, Patricia Chang, Abigail Cyr, Pamela Hook, Jiayi Lin, Jack Murtagh, and Carly Schimmel.

References

Andersson, J. L. R., & Sotiropoulos, S. N. (2016). An integrated approach to correction for off-resonance effects and subject movement in diffusion MR imaging. *NeuroImage*, *125*, 1063–1078. https://doi.org/10.1016/j.neuroimage.2015.10.019

Banfi, C., Koschutnig, K., Moll, K., Schulte-Körne, G., Fink, A., & Landerl, K. (2018). White matter alterations and tract lateralization in children with dyslexia and isolated spelling deficits. *Human Brain Mapping*, *40*(3), 765–776. https://doi.org/10.1002/hbm.24410

Barquero, L. A., Davis, N., & Cutting, L. E. (2014). Neuroimaging of reading intervention: A systematic review and activation likelihood estimate meta-analysis. *PloS One*, *9*(1), e83668.

Basser, P. J., Mattiello, J., & LeBihan, D. (1994). MR diffusion tensor spectroscopy and imaging. *Biophysical Journal*, *66*(1), 259–267. https://doi.org/10.1016/S0006-3495(94)80775-1

Beaulieu, C. (2002). The basis of anisotropic water diffusion in the nervous system – a technical review. *NMR in Biomedicine*, *15*(7–8), 435–455. https://doi.org/10.1002/nbm.782

Behler, A., Kassubek, J., & Müller, H.-P. (2021). Age-Related Alterations in DTI Metrics in the Human Brain—Consequences for Age Correction. *Frontiers in Aging Neuroscience*, *13*. https://www.frontiersin.org/articles/10.3389/fnagi.2021.682109

Behrens, T. E. J., Berg, H. J., Jbabdi, S., Rushworth, M. F. S., & Woolrich, M. W. (2007). Probabilistic diffusion tractography with multiple fibre orientations: What can we gain? *NeuroImage*, *34*(1), 144–155. https://doi.org/10.1016/j.neuroimage.2006.09.018

Bell, N. (1997). *Seeing stars: Symbol imagery for phonemic awareness, sight words and spelling*. Gander Publishing Avila Beach, CA.

Bell, N. (2010). *Symbol imagery test*. Gander Publishing.

Ben-Shachar, M., Dougherty, R. F., & Wandell, B. A. (2007). White matter pathways in reading.

Current Opinion in Neurobiology, *17*(2), 258–270.

https://doi.org/10.1016/j.conb.2007.03.006

Blumenfeld-Katzir, T., Pasternak, O., Dagan, M., & Assaf, Y. (2011). Diffusion MRI of Structural

Brain Plasticity Induced by a Learning and Memory Task. *PLOS ONE*, *6*(6), e20678.

https://doi.org/10.1371/journal.pone.0020678

Borchers, L. R., Bruckert, L., Dodson, C. K., Travis, K. E., Marchman, V. A., Ben-Shachar, M., &

Feldman, H. M. (2019). Microstructural properties of white matter pathways in relation

to subsequent reading abilities in children: A longitudinal analysis. *Brain Structure &*

Function, *224*(2), 891–905. https://doi.org/10.1007/s00429-018-1813-z

Braid, J., & Richlan, F. (2022). The Functional Neuroanatomy of Reading Intervention. *Frontiers*

in Neuroscience, *16*, 921931. https://doi.org/10.3389/fnins.2022.921931

Button, K. S., Ioannidis, J. P. A., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S. J., &

Munafò, M. R. (2013). Power failure: Why small sample size undermines the reliability of

neuroscience. *Nature Reviews Neuroscience*, *14*(5), Article 5.

https://doi.org/10.1038/nrn3475

Catani, M., Jones, D. K., & Ffytche, D. H. (2005). Perisylvian language networks of the human

brain. *Annals of Neurology*, *57*(1), 8–16. https://doi.org/10.1002/ana.20319

Catani, M., & Mesulam, M. (2008). The arcuate fasciculus and the disconnection theme in

language and aphasia: History and current state. *Cortex*, *44*(8), 953–961.

https://doi.org/10.1016/j.cortex.2008.04.002

Catts, H. W. (1993). The Relationship Between Speech-Language Impairments and Reading

Disabilities. *Journal of Speech, Language, and Hearing Research*, *36*(5), 948–958.

https://doi.org/10.1044/jshr.3605.948

Christodoulou, J. A., Cyr, A., Murtagh, J., Chang, P., Lin, J., Guarino, A. J., Hook, P., & Gabrieli,

J. D. E. (2017). Impact of Intensive Summer Reading Intervention for Children With

Reading Disabilities and Difficulties in Early Elementary School. *Journal of Learning Disabilities*, *50*(2), 115–127. https://doi.org/10.1177/0022219415617163

Connally, E. L., Ward, D., Howell, P., & Watkins, K. E. (2014). Disrupted white matter in language and motor tracts in developmental stuttering. *Brain and Language*, *131*, 25–35. https://doi.org/10.1016/j.bandl.2013.05.013

Cooper, H., Nye, B., Charlton, K., Lindsay, J., & Greathouse, S. (1996). The Effects of Summer Vacation on Achievement Test Scores: A Narrative and Meta-Analytic Review. *Review of Educational Research*, *66*(3), 227–268. https://doi.org/10.3102/00346543066003227

Cui, Z., Xia, Z., Su, M., Shu, H., & Gong, G. (2016). Disrupted white matter connectivity underlying developmental dyslexia: A machine learning approach. *Human Brain Mapping*, *37*(4), 1443–1458. https://doi.org/10.1002/hbm.23112

Davis, B. R., Garza, A., & Church, J. A. (2022). Key considerations for child and adolescent MRI data collection. *Frontiers in Neuroimaging*, *1*, 981947. https://doi.org/10.3389/fnimg.2022.981947

Davison, K. E., Zuk, J., Mullin, L. J., Ozernov-Palchik, O., Norton, E., Gabrieli, J. D. E., Yu, X., & Gaab, N. (2022). Examining Shared Reading and White Matter Organization in Kindergarten in Relation to Subsequent Language and Reading Abilities: A Longitudinal Investigation. *Journal of Cognitive Neuroscience*, 1–17. https://doi.org/10.1162/jocn_a_01944

de Bie, H. M. A., Boersma, M., Wattjes, M. P., Adriaanse, S., Vermeulen, R. J., Oostrom, K. J., Huisman, J., Veltman, D. J., & Delemarre-Van de Waal, H. A. (2010). Preparing children with a mock scanner training protocol results in high quality structural and functional MRI scans. *European Journal of Pediatrics*, *169*(9), 1079–1085. https://doi.org/10.1007/s00431-010-1181-z

De Santis, S., Drakesmith, M., Bells, S., Assaf, Y., & Jones, D. K. (2014). Why diffusion tensor MRI does well only some of the time: Variance and covariance of white matter tissue

microstructure attributes in the living human brain. *NeuroImage*, *89*, 35–44.

https://doi.org/10.1016/j.neuroimage.2013.12.003

Dehaene, S., & Cohen, L. (2011). The unique role of the visual word form area in reading.

Trends in Cognitive Sciences, *15*(6), 254–262. https://doi.org/10.1016/j.tics.2011.04.003

Donnelly, P. M., Huber, E., & Yeatman, J. D. (2019). Intensive Summer Intervention Drives

Linear Growth of Reading Skill in Struggling Readers. *Frontiers in Psychology*, *10*, 1900.

https://doi.org/10.3389/fpsyg.2019.01900

Duffau, H., Herbet, G., & Moritz-Gasser, S. (2013). Toward a pluri-component, multimodal, and

dynamic organization of the ventral semantic stream in humans: Lessons from

stimulation mapping in awake patients. *Frontiers in Systems Neuroscience*, *7*.

https://www.frontiersin.org/articles/10.3389/fnsys.2013.00044

Economou, M., Van Herck, S., Vanden Bempt, F., Glatz, T., Wouters, J., Ghesquière, P.,

Vanderauwera, J., & Vandermosten, M. (2022). Investigating the impact of early literacy

training on white matter structure in prereaders at risk for dyslexia. *Cerebral Cortex*,

32(21), 4684–4697. https://doi.org/10.1093/cercor/bhab510

Enge, A., Friederici, A. D., & Skeide, M. A. (2020). A meta-analysis of fMRI studies of language

comprehension in children. *NeuroImage*, *215*, 116858.

https://doi.org/10.1016/j.neuroimage.2020.116858

Entwisle, D. R., Alexander, K. L., & Olson, L. S. (1997). *Children, schools, and inequality*.

Routledge.

Epelbaum, S., Pinel, P., Gaillard, R., Delmaire, C., Perrin, M., Dupont, S., Dehaene, S., &

Cohen, L. (2008). Pure alexia as a disconnection syndrome: New diffusion imaging

evidence for an old concept. *Cortex; a Journal Devoted to the Study of the Nervous*

System and Behavior, *44*(8), 962–974. https://doi.org/10.1016/j.cortex.2008.05.003

Fedorenko, E., Behr, M. K., & Kanwisher, N. (2011). Functional specificity for high-level

linguistic processing in the human brain. *Proceedings of the National Academy of*

Sciences, *108*(39), 16428–16433. https://doi.org/10.1073/pnas.1112937108

Fields, R. D. (2015). A new mechanism of nervous system plasticity: Activity-dependent

myelination. *Nature Reviews Neuroscience*, *16*(12), Article 12.

https://doi.org/10.1038/nrn4023

Fields, R. D., Araque, A., Johansen-Berg, H., Lim, S.-S., Lynch, G., Nave, K.-A., Nedergaard,

M., Perez, R., Sejnowski, T., & Wake, H. (2014). Glial Biology in Learning and Cognition.

The Neuroscientist, *20*(5), 426–431. https://doi.org/10.1177/1073858413504465

Fischl, B. (2012). FreeSurfer. *NeuroImage*, *62*(2), 774–781.

https://doi.org/10.1016/j.neuroimage.2012.01.021

Fischl, B., Salat, D. H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., van der Kouwe, A.,

Killiany, R., Kennedy, D., Klaveness, S., Montillo, A., Makris, N., Rosen, B., & Dale, A.

M. (2002). Whole Brain Segmentation: Automated Labeling of Neuroanatomical

Structures in the Human Brain. *Neuron*, *33*(3), 341–355. https://doi.org/10.1016/S0896-

6273(02)00569-X

Friedrich, P., Fraenz, C., Schlüter, C., Ocklenburg, S., Mädler, B., Güntürkün, O., & Genç, E.

(2020). The Relationship Between Axon Density, Myelination, and Fractional Anisotropy

in the Human Corpus Callosum. *Cerebral Cortex (New York, N.Y.: 1991)*, *30*(4), 2042–

2056. https://doi.org/10.1093/cercor/bhz221

Gao, P., Wang, Y.-S., Lu, Q.-Y., Rong, M.-J., Fan, X.-R., Holmes, A. J., Dong, H.-M., Li, H.-F., &

Zuo, X.-N. (2023). Brief mock-scan training reduces head motion during real scanning

for children: A growth curve study. *Developmental Cognitive Neuroscience*, *61*, 101244.

https://doi.org/10.1016/j.dcn.2023.101244

Genc, S., Malpas, C. B., Holland, S. K., Beare, R., & Silk, T. J. (2017). Neurite density index is

sensitive to age related differences in the developing brain. *NeuroImage*, *148*, 373–380.

https://doi.org/10.1016/j.neuroimage.2017.01.023

Genc, S., Tax, C. M. W., Raven, E. P., Chamberland, M., Parker, G. D., & Jones, D. K. (2020).

Impact of b-value on estimates of apparent fibre density. *Human Brain Mapping*, *41*(10),

2583–2595. https://doi.org/10.1002/hbm.24964

Good, R. H., Gruba, J., & Kaminski, R. A. (2002). Best Practices in Using Dynamic Indicators of

Basic Early Literacy Skills (DIBELS) in an Outcomes-Driven Model. In *Best practices in*

school psychology IV, Vols. 1-2 (pp. 699–720). National Association of School

Psychologists.

Greene, D. J., Koller, J. M., Hampton, J. M., Wesevich, V., Van, A. N., Nguyen, A. L., Hoyt, C.

R., McIntyre, L., Earl, E. A., & Klein, R. L. (2018). Behavioral interventions for reducing

head motion during MRI scans in children. *Neuroimage*, *171*, 234–245.

Greve, D. N., & Fischl, B. (2009). Accurate and robust brain image alignment using boundary-

based registration. *Neuroimage*, *48*(1), 63–72.

Gullick, M. M., & Booth, J. R. (2015). The direct segment of the arcuate fasciculus is predictive

of longitudinal reading change. *Developmental Cognitive Neuroscience*, *13*, 68–74.

https://doi.org/10.1016/j.dcn.2015.05.002

Han, Z., Ma, Y., Gong, G., Huang, R., Song, L., & Bi, Y. (2016). White matter pathway

supporting phonological encoding in speech production: A multi-modal imaging study of

brain damage patients. *Brain Structure and Function*, *221*(1), 577–589.

https://doi.org/10.1007/s00429-014-0926-2

Hayiou, -Thomas Marianna E., Harlaar, N., Dale, P. S., & Plomin, R. (2010). Preschool Speech,

Language Skills, and Reading at 7, 9, and 10 Years: Etiology of the Relationship.

Journal of Speech, Language, and Hearing Research, *53*(2), 311–332.

https://doi.org/10.1044/1092-4388(2009/07-0145)

Herbet, G., Zemmoura, I., & Duffau, H. (2018). Functional Anatomy of the Inferior Longitudinal

Fasciculus: From Historical Reports to Current Hypotheses. *Frontiers in Neuroanatomy*,

12. https://www.frontiersin.org/articles/10.3389/fnana.2018.00077

Hernández, M., Guerrero, G. D., Cecilia, J. M., García, J. M., Inuggi, A., Jbabdi, S., Behrens, T.

E. J., & Sotiropoulos, S. N. (2013). Accelerating Fibre Orientation Estimation from

Diffusion Weighted Magnetic Resonance Imaging Using GPUs. *PLOS ONE*, *8*(4),

e61892. https://doi.org/10.1371/journal.pone.0061892

Hoeft, F., McCandliss, B. D., Black, J. M., Gantman, A., Zakerani, N., Hulme, C., Lyytinen, H.,

Whitfield-Gabrieli, S., Glover, G. H., & Reiss, A. L. (2011). Neural systems predicting

long-term outcome in dyslexia. *Proceedings of the National Academy of Sciences*,

108(1), 361–366.

Huber, E., Donnelly, P. M., Rokem, A., & Yeatman, J. D. (2018). Rapid and widespread white

matter plasticity during an intensive reading intervention. *Nature Communications*, *9*(1),

1–13.

Huber, E., Mezer, A., & Yeatman, J. D. (2021). Neurobiological underpinnings of rapid white

matter plasticity during intensive reading instruction. *NeuroImage*, *243*, 118453.

https://doi.org/10.1016/j.neuroimage.2021.118453

Jbabdi, S., Sotiropoulos, S. N., Savio, A. M., Graña, M., & Behrens, T. E. J. (2012). Model-

based analysis of multishell diffusion MR data for tractography: How to get over fitting

problems. *Magnetic Resonance in Medicine*, *68*(6), 1846–1855.

https://doi.org/10.1002/mrm.24204

Jelescu, I. O., Palombo, M., Bagnato, F., & Schilling, K. G. (2020). Challenges for biophysical

modeling of microstructure. *Journal of Neuroscience Methods*, *344*, 108861.

https://doi.org/10.1016/j.jneumeth.2020.108861

Jeurissen, B., Leemans, A., Tournier, J.-D., Jones, D. K., & Sijbers, J. (2013). Investigating the

prevalence of complex fiber configurations in white matter tissue with diffusion magnetic

resonance imaging. *Human Brain Mapping*, *34*(11), 2747–2766.

Jones, D. K., Knösche, T. R., & Turner, R. (2013). White matter integrity, fiber count, and other

fallacies: The do's and don'ts of diffusion MRI. *NeuroImage*, *73*, 239–254.

https://doi.org/10.1016/j.neuroimage.2012.06.081

Kaufman, A. S., Kaufman, Nadeen L. ,. (2004). *Kaufman brief intelligence test KBIT 2 ; manual*.

/z-wcorg/.

Keller, T. A., & Just, M. A. (2009). Altering Cortical Connectivity: Remediation-Induced Changes

in the White Matter of Poor Readers. *Neuron*, *64*(5), 624–631.

https://doi.org/10.1016/j.neuron.2009.10.018

Kharabian Masouleh, S., Eickhoff, S. B., Hoffstaedter, F., Genon, S., & Alzheimer's Disease

Neuroimaging Initiative. (2019). Empirical examination of the replicability of associations

between brain structure and psychological variables. *ELife*, *8*, e43464.

https://doi.org/10.7554/eLife.43464

King, K. M., Littlefield, A. K., McCabe, C. J., Mills, K. L., Flournoy, J., & Chassin, L. (2018).

Longitudinal modeling in developmental neuroimaging research: Common challenges,

and solutions from developmental psychology. *Developmental Cognitive Neuroscience*,

33, 54–72. https://doi.org/10.1016/j.dcn.2017.11.009

Koirala, N., Perdue, M. V., Su, X., Grigorenko, E. L., & Landi, N. (2021). Neurite density and

arborization is associated with reading skill and phonological processing in children.

NeuroImage, *241*, 118426. https://doi.org/10.1016/j.neuroimage.2021.118426

Kronfeld-Duenias, V., Amir, O., Ezrati-Vinacour, R., Civier, O., & Ben-Shachar, M. (2016). The

frontal aslant tract underlies speech fluency in persistent developmental stuttering. *Brain*

Structure and Function, *221*(1), 365–381. https://doi.org/10.1007/s00429-014-0912-8

Kuhfeld, M., Lewis, K., & Peltier, T. (2023). Reading achievement declines during the COVID-19

pandemic: Evidence from 5 million U.S. students in grades 3–8. *Reading and Writing*,

36(2), 245–261. https://doi.org/10.1007/s11145-022-10345-8

Kurtzer, G. M., Sochat, V., & Bauer, M. W. (2017). Singularity: Scientific containers for mobility

of compute. *PLOS ONE*, *12*(5), e0177459. https://doi.org/10.1371/journal.pone.0177459

Langer, N., Peysakhovich, B., Zuk, J., Drottar, M., Sliva, D. D., Smith, S., Becker, B. L. C.,

Grant, P. E., & Gaab, N. (2017). White Matter Alterations in Infants at Risk for

Developmental Dyslexia. *Cerebral Cortex*, *27*(2), 1027–1036.

https://doi.org/10.1093/cercor/bhv281

Leemans, A., & Jones, D. K. (2009). The B-matrix must be rotated when correcting for subject

motion in DTI data. *Magnetic Resonance in Medicine*, *61*(6), 1336–1349.

https://doi.org/10.1002/mrm.21890

Lerma-Usabiaga, G., Carreiras, M., & Paz-Alonso, P. M. (2018). Converging evidence for

functional and structural segregation within the left ventral occipitotemporal cortex in

reading. *Proceedings of the National Academy of Sciences*, *115*(42), E9981–E9990.

https://doi.org/10.1073/pnas.1803003115

López-Vicente, M., Lamballais, S., Louwen, S., Hillegers, M., Tiemeier, H., Muetzel, R. L., &

White, T. (2021). White matter microstructure correlates of age, sex, handedness and

motor ability in a population-based sample of 3031 school-age children. *NeuroImage*,

227, 117643. https://doi.org/10.1016/j.neuroimage.2020.117643

Maffei, C., Lee, C., Planich, M., Ramprasad, M., Ravi, N., Trainor, D., Urban, Z., Kim, M., Jones,

R. J., Henin, A., Hofmann, S. G., Pizzagalli, D. A., Auerbach, R. P., Gabrieli, J. D. E.,

Whitfield-Gabrieli, S., Greve, D. N., Haber, S. N., & Yendiki, A. (2021). Using diffusion

MRI data acquired with ultra-high gradient strength to improve tractography in routine-

quality data. *NeuroImage*, *245*, 118706.

https://doi.org/10.1016/j.neuroimage.2021.118706

Marek, S., Tervo-Clemmens, B., Calabro, F. J., Montez, D. F., Kay, B. P., Hatoum, A. S.,

Donohue, M. R., Foran, W., Miller, R. L., Hendrickson, T. J., Malone, S. M., Kandala, S.,

Feczko, E., Miranda-Dominguez, O., Graham, A. M., Earl, E. A., Perrone, A. J.,

Cordova, M., Doyle, O., … Dosenbach, N. U. F. (2022). Reproducible brain-wide

association studies require thousands of individuals. *Nature*, *603*(7902), Article 7902.

https://doi.org/10.1038/s41586-022-04492-9

McKinney, W. (2011). pandas: A foundational Python library for data analysis and statistics.

Python for High Performance and Scientific Computing, *14*(9), 1–9.

Meisler, S. L., & Gabrieli, J. D. (2022a). A Large-Scale Investigation of White Matter

Microstructural Associations with Reading Ability. *NeuroImage*, 118909.

Meisler, S. L., & Gabrieli, J. D. (2022b). Fiber-specific structural properties relate to reading

skills in children and adolescents. *ELife*, *11*, e82088. https://doi.org/10.7554/eLife.82088

Metzler-Baddeley, C., Foley, S., de Santis, S., Charron, C., Hampshire, A., Caeyenberghs, K., &

Jones, D. K. (2017). Dynamics of White Matter Plasticity Underlying Working Memory

Training: Multimodal Evidence from Diffusion MRI and Relaxometry. *Journal of Cognitive*

Neuroscience, *29*(9), 1509–1520. https://doi.org/10.1162/jocn_a_01127

Moreau, D., Stonyer, J. E., McKay, N. S., & Waldie, K. E. (2018). No evidence for systematic

white matter correlates of dyslexia: An Activation Likelihood Estimation meta-analysis.

Brain Research, *1683*, 36–47. https://doi.org/10.1016/j.brainres.2018.01.014

Murphy, K. A., Jogia, J., & Talcott, J. B. (2019). On the neural basis of word reading: A meta-

analysis of fMRI evidence using activation likelihood estimation. *Journal of*

Neurolinguistics, *49*, 71–83.

Myers, C. A., Vandermosten, M., Farris, E. A., Hancock, R., Gimenez, P., Black, J. M., Casto,

B., Drahos, M., Tumber, M., Hendren, R. L., Hulme, C., & Hoeft, F. (2014). White Matter

Morphometric Changes Uniquely Predict Children's Reading Acquisition. *Psychological*

Science, *25*(10), 1870–1883. https://doi.org/10.1177/0956797614544511

Ng, S., Moritz-Gasser, S., Lemaitre, A.-L., Duffau, H., & Herbet, G. (2021). White matter

disconnectivity fingerprints causally linked to dissociated forms of alexia.

Communications Biology, *4*(1), Article 1. https://doi.org/10.1038/s42003-021-02943-z

Northam, G. B., Liégeois, F., Chong, W. K., Baker, K., Tournier, J.-D., Wyatt, J. S., Baldeweg,

T., & Morgan, A. (2012). Speech and Oromotor Outcome in Adolescents Born Preterm:

Relationship to Motor Tract Integrity. *The Journal of Pediatrics*, *160*(3), 402-408.e1.

https://doi.org/10.1016/j.jpeds.2011.08.055

Northam, G. B., Morgan, A. T., Fitzsimmons, S., Baldeweg, T., & Liégeois, F. J. (2019).

Corticobulbar Tract Injury, Oromotor Impairment and Language Plasticity in Adolescents

Born Preterm. *Frontiers in Human Neuroscience*, *13*.

https://www.frontiersin.org/articles/10.3389/fnhum.2019.00045

Partanen, M., Kim, D. H., Rauscher, A., Siegel, L. S., & Giaschi, D. E. (2021). White matter but

not grey matter predicts change in reading skills after intervention. *Dyslexia*, *27*(2), 224–

244.

Perdue, M. V., Mahaffy, K., Vlahcevic, K., Wolfman, E., Erbeli, F., Richlan, F., & Landi, N.

(2022). Reading intervention and neuroplasticity: A systematic review and meta-analysis

of brain changes associated with reading intervention. *Neuroscience & Biobehavioral*

Reviews, *132*, 465–494.

Phan, T. V., Sima, D., Smeets, D., Ghesquière, P., Wouters, J., & Vandermosten, M. (2021).

Structural brain dynamics across reading development: A longitudinal MRI study from

kindergarten to grade 5. *Human Brain Mapping*, *42*(14), 4497–4509.

https://doi.org/10.1002/hbm.25560

Raffelt, D. A., Tournier, J.-D., Smith, R. E., Vaughan, D. N., Jackson, G., Ridgway, G. R., &

Connelly, A. (2017). Investigating white matter fibre density and morphology using fixel-

based analysis. *NeuroImage*, *144*, 58–73.

https://doi.org/10.1016/j.neuroimage.2016.09.029

Reuter, M., & Fischl, B. (2011). Avoiding asymmetry-induced bias in longitudinal image

870   processing. *NeuroImage*, *57*(1), 19–21.

871   https://doi.org/10.1016/j.neuroimage.2011.02.076

872   Reuter, M., Rosas, H. D., & Fischl, B. (2010). Highly accurate inverse consistent registration: A

873   robust approach. *NeuroImage*, *53*(4), 1181–1196.

874   https://doi.org/10.1016/j.neuroimage.2010.07.020

875   Reuter, M., Schmansky, N. J., Rosas, H. D., & Fischl, B. (2012). Within-subject template

876   estimation for unbiased longitudinal image analysis. *NeuroImage*, *61*(4), 1402–1418.

877   https://doi.org/10.1016/j.neuroimage.2012.02.084

878   Richards, T. L., Berninger, V. W., Yagle, K. J., Abbott, R. D., & Peterson, D. J. (2017). Changes

879   in DTI Diffusivity and fMRI Connectivity Cluster Coefficients for Students with and

880   without Specific Learning Disabilities In Written Language: Brain's Response to Writing

881   Instruction. *Journal of Nature and Science*, *3*(4), e350.

882   Romeo, R. R., Christodoulou, J. A., Halverson, K. K., Murtagh, J., Cyr, A. B., Schimmel, C.,

883   Chang, P., Hook, P. E., & Gabrieli, J. D. E. (2018). Socioeconomic Status and Reading

884   Disability: Neuroanatomy and Plasticity in Response to Intervention. *Cerebral Cortex*,

885   *28*(7), 2297–2312. https://doi.org/10.1093/cercor/bhx131

886   Roy, E., Richie-Halford, A., Kruper, J., Narayan, M., Bloom, D., Brown, T. T., Jernigan, T. L.,

887   McCandliss, B. D., Rokem, A., & Yeatman, J. D. (2022). *White matter and literacy: A*

888   *dynamic system in flux* (p. 2022.06.21.497048). bioRxiv.

889   https://doi.org/10.1101/2022.06.21.497048

890   Sampaio-Baptista, C., & Johansen-Berg, H. (2017). White Matter Plasticity in the Adult Brain.

891   *Neuron*, *96*(6), 1239–1251. https://doi.org/10.1016/j.neuron.2017.11.026

892   Sampaio-Baptista, C., Khrapitchev, A. A., Foxley, S., Schlagheck, T., Scholz, J., Jbabdi, S.,

893   DeLuca, G. C., Miller, K. L., Taylor, A., Thomas, N., Kleim, J., Sibson, N. R.,

894   Bannerman, D., & Johansen-Berg, H. (2013). Motor Skill Learning Induces Changes in

White Matter Microstructure and Myelination. *Journal of Neuroscience*, *33*(50), 19499–

19503. https://doi.org/10.1523/JNEUROSCI.3048-13.2013

Saur, D., Kreher, B. W., Schnell, S., Kümmerer, D., Kellmeyer, P., Vry, M.-S., Umarova, R.,

Musso, M., Glauche, V., Abel, S., Huber, W., Rijntjes, M., Hennig, J., & Weiller, C.

(2008). Ventral and dorsal pathways for language. *Proceedings of the National Academy*

of Sciences, *105*(46), 18035–18040. https://doi.org/10.1073/pnas.0805234105

Saygin, Z. M., Norton, E. S., Osher, D. E., Beach, S. D., Cyr, A. B., Ozernov-Palchik, O.,

Yendiki, A., Fischl, B., Gaab, N., & Gabrieli, J. D. E. (2013). Tracking the Roots of

Reading Ability: White Matter Volume and Integrity Correlate with Phonological

Awareness in Prereading and Early-Reading Kindergarten Children. *Journal of*

Neuroscience, *33*(33), 13251–13258. https://doi.org/10.1523/JNEUROSCI.4383-12.2013

Schilling, K. G., Rheault, F., Petit, L., Hansen, C. B., Nath, V., Yeh, F.-C., Girard, G., Barakovic,

M., Rafael-Patino, J., Yu, T., Fischi-Gomez, E., Pizzolato, M., Ocampo-Pineda, M.,

Schiavi, S., Canales-Rodríguez, E. J., Daducci, A., Granziera, C., Innocenti, G., Thiran,

J.-P., … Descoteaux, M. (2021). Tractography dissection variability: What happens

when 42 groups dissect 14 white matter bundles on the same dataset? *NeuroImage*,

243, 118502. https://doi.org/10.1016/j.neuroimage.2021.118502

Schilling, K. G., Tax, C. M. W., Rheault, F., Hansen, C., Yang, Q., Yeh, F.-C., Cai, L., Anderson,

A. W., & Landman, B. A. (2021). Fiber tractography bundle segmentation depends on

scanner effects, vendor effects, acquisition resolution, diffusion sampling scheme,

diffusion sensitization, and bundle segmentation workflow. *NeuroImage*, *242*, 118451.

https://doi.org/10.1016/j.neuroimage.2021.118451

Schilling, K. G., Tax, C. M. W., Rheault, F., Landman, B. A., Anderson, A. W., Descoteaux, M.,

& Petit, L. (2022). Prevalence of white matter pathways coming into a single white matter

voxel orientation: The bottleneck issue in tractography. *Human Brain Mapping*, *43*(4),

1196–1213. https://doi.org/10.1002/hbm.25697

Scholz, J., Klein, M. C., Behrens, T. E. J., & Johansen-Berg, H. (2009). Training induces

changes in white matter architecture. *Nature Neuroscience*, *12*(11), 1370–1371.

https://doi.org/10.1038/nn.2412

Seabold, S., & Perktold, J. (2010). Statsmodels: Econometric and statistical modeling with

python. *Proceedings of the 9th Python in Science Conference*, *57*(61), 10.25080.

Shaywitz, S. E. (1998). Dyslexia. *New England Journal of Medicine*, *338*(5), 307–312.

https://doi.org/10.1056/NEJM199801293380507

Shin, J., Rowley, J., Chowdhury, R., Jolicoeur, P., Klein, D., Grova, C., Rosa-Neto, P., &

Kobayashi, E. (2019). Inferior Longitudinal Fasciculus' Role in Visual Processing and

Language Comprehension: A Combined MEG-DTI Study. *Frontiers in Neuroscience*, *13*,

875. https://doi.org/10.3389/fnins.2019.00875

Sihvonen, A. J., Virtala, P., Thiede, A., Laasonen, M., & Kujala, T. (2021). Structural white

matter connectometry of reading and dyslexia. *NeuroImage*, *241*, 118411.

Thiebaut de Schotten, M., Cohen, L., Amemiya, E., Braga, L. W., & Dehaene, S. (2014).

Learning to Read Improves the Structure of the Arcuate Fasciculus. *Cerebral Cortex*,

24(4), 989–995. https://doi.org/10.1093/cercor/bhs383

Thornton, A., & Lee, P. (2000). Publication bias in meta-analysis: Its causes and consequences.

Journal of Clinical Epidemiology, *53*(2), 207–216. https://doi.org/10.1016/S0895-

4356(99)00161-4

Torgesen, J. K., Wagner, R. K., & Rashotte, C. A. (2012). Test of word reading efficiency–

second edition (TOWRE-2). *Austin, TX: Pro-Ed*.

Torgeson, J. K., Wagner, R. K., & Rashotte, C. A. (1999). *Test Review: Test of Word Reading

Efficiency (TOWRE)*. Pro-ed.

Turner, B. O., Paul, E. J., Miller, M. B., & Barbey, A. K. (2018). Small sample sizes reduce the

replicability of task-based fMRI studies. *Communications Biology*, *1*(1), Article 1.

https://doi.org/10.1038/s42003-018-0073-z

Van Der Auwera, S., Vandermosten, M., Wouters, J., Ghesquière, P., & Vanderauwera, J.

(2021). A three-time point longitudinal investigation of the arcuate fasciculus throughout

reading acquisition in children developing dyslexia. *NeuroImage*, *237*, 118087.

https://doi.org/10.1016/j.neuroimage.2021.118087

Vanderauwera, J., Wouters, J., Vandermosten, M., & Ghesquière, P. (2017). Early dynamics of

white matter deficits in children developing dyslexia. *Developmental Cognitive*

Neuroscience, *27*, 69–77. https://doi.org/10.1016/j.dcn.2017.08.003

Vandermosten, M., Vanderauwera, J., Theys, C., De Vos, A., Vanvooren, S., Sunaert, S.,

Wouters, J., & Ghesquière, P. (2015). A DTI tractography study in pre-readers at risk for

dyslexia. *Developmental Cognitive Neuroscience*, *14*, 8–15.

https://doi.org/10.1016/j.dcn.2015.05.006

Vigneau, M., Beaucousin, V., Hervé, P. Y., Duffau, H., Crivello, F., Houdé, O., Mazoyer, B., &

Tzourio-Mazoyer, N. (2006). Meta-analyzing left hemisphere language areas:

Phonology, semantics, and sentence processing. *NeuroImage*, *30*(4), 1414–1432.

https://doi.org/10.1016/j.neuroimage.2005.11.002

von Hippel, P. T., Workman, J., & Downey, D. B. (2018). Inequality in Reading and Math Skills

Forms Mainly before Kindergarten: A Replication, and Partial Correction, of "Are Schools

the Great Equalizer?" *Sociology of Education*, *91*(4), 323–357.

https://doi.org/10.1177/0038040718801760

Wagner, R. K., Torgesen, J. K., Rashotte, C. A., & Pearson, N. A. (1999). *Comprehensive test*

of phonological processing: CTOPP. Pro-ed Austin, TX.

Wagner, R. K., Zirps, F. A., Edwards, A. A., Wood, S. G., Joyner, R. E., Becker, B. J., Liu, G., &

Beal, B. (2020). The prevalence of Dyslexia: A new approach to its estimation. *Journal of*

Learning Disabilities, *53*(5), 354–365.

Walton, M., Dewey, D., & Lebel, C. (2018). Brain white matter structure and language ability in

preschool-aged children. *Brain and Language*, *176*, 19–25.

https://doi.org/10.1016/j.bandl.2017.10.008

Wang, Y., Mauer, M. V., Raney, T., Peysakhovich, B., Becker, B. L., Sliva, D. D., & Gaab, N.

(2017). Development of tract-specific white matter pathways during early reading

development in at-risk children and typical controls. *Cerebral Cortex*, *27*(4), 2469–2485.

Waskom, M. L. (2021). Seaborn: Statistical data visualization. *Journal of Open Source*

Software, *6*(60), 3021.

Weiner, K. S., Yeatman, J. D., & Wandell, B. A. (2017). The posterior arcuate fasciculus and the

vertical occipital fasciculus. *Cortex; a Journal Devoted to the Study of the Nervous*

System and Behavior, *97*, 274–276. https://doi.org/10.1016/j.cortex.2016.03.012

Wolf, M., & Denckla, M. B. (2005). *RAN/RAS: Rapid automatized naming and rapid alternating*

stimulus tests. Pro-ed Austin, TX.

Woodcock, R. W. (2011). *Woodcock reading mastery tests: WRMT-III*. Pearson.

Wu, L.-Y., Xu, Y., Chen, L.-L., Yang, W.-R., Li, Y., Shang, S.-A., Luo, X.-F., Xia, W., Xia, J., &

Zhang, H.-Y. (2022). Test-retest reliability of diffusion kurtosis imaging metrics in the

healthy adult brain. *Neuroimage: Reports*, *2*(3), 100098.

https://doi.org/10.1016/j.ynirp.2022.100098

Xin, W., & Chan, J. R. (2020). Myelin plasticity: Sculpting circuits in learning and memory.

Nature Reviews Neuroscience, *21*(12), Article 12. https://doi.org/10.1038/s41583-020-

00379-8

Yeatman, J. D., Dougherty, R. F., Ben-Shachar, M., & Wandell, B. A. (2012). Development of

white matter and reading skills. *Proceedings of the National Academy of Sciences*,

109(44), E3045–E3053.

Yeatman, J. D., Rauschecker, A. M., & Wandell, B. A. (2013). Anatomy of the visual word form

area: Adjacent cortical circuits and long-range white matter connections. *Brain and*

Language, *125*(2), 146–155. https://doi.org/10.1016/j.bandl.2012.04.010

Yeatman, J. D., & White, A. L. (2021). Reading: The Confluence of Vision and Language.

Annual Review of Vision Science, *7*(1), 487–517. https://doi.org/10.1146/annurev-vision-

093019-113509

Yeh, F.-C., Verstynen, T. D., Wang, Y., Fernández-Miranda, J. C., & Tseng, W.-Y. I. (2013).

Deterministic Diffusion Fiber Tracking Improved by Quantitative Anisotropy. *PLOS ONE*,

8(11), e80713. https://doi.org/10.1371/journal.pone.0080713

Yendiki, A., Koldewyn, K., Kakunoori, S., Kanwisher, N., & Fischl, B. (2014). Spurious group

differences due to head motion in a diffusion MRI study. *NeuroImage*, *88*, 79–90.

https://doi.org/10.1016/j.neuroimage.2013.11.027

Yendiki, A., Panneck, P., Srinivasan, P., Stevens, A., Zöllei, L., Augustinack, J., Wang, R.,

Salat, D., Ehrlich, S., Behrens, T., Jbabdi, S., Gollub, R., & Fischl, B. (2011). Automated

Probabilistic Reconstruction of White-Matter Pathways in Health and Disease Using an

Atlas of the Underlying Anatomy. *Frontiers in Neuroinformatics*, *5*.

https://www.frontiersin.org/articles/10.3389/fninf.2011.00023

Yendiki, A., Reuter, M., Wilkens, P., Rosas, H. D., & Fischl, B. (2016). Joint reconstruction of

white-matter pathways from longitudinal diffusion MRI data with anatomical priors.

NeuroImage, *127*, 277–286. https://doi.org/10.1016/j.neuroimage.2015.12.003

Yu, Q., Peng, Y., Kang, H., Peng, Q., Ouyang, M., Slinger, M., Hu, D., Shou, H., Fang, F., &

Huang, H. (2020). Differential White Matter Maturation from Birth to 8 Years of Age.

Cerebral Cortex, *30*(4), 2674–2690. https://doi.org/10.1093/cercor/bhz268

Zemmoura, I., Herbet, G., Moritz-Gasser, S., & Duffau, H. (2015). New insights into the neural

network mediating reading processes provided by cortico-subcortical electrical mapping.

Human Brain Mapping, *36*(6), 2215–2230. https://doi.org/10.1002/hbm.22766

Zhang, H., Schneider, T., Wheeler-Kingshott, C. A., & Alexander, D. C. (2012). NODDI:

Practical in vivo neurite orientation dispersion and density imaging of the human brain.

NeuroImage, *61*(4), 1000–1016. https://doi.org/10.1016/j.neuroimage.2012.03.072

Zuk, J., Dunstan, J., Norton, E., Yu, X., Ozernov-Palchik, O., Wang, Y., Hogan, T. P., Gabrieli,

J. D. E., & Gaab, N. (2021). Multifactorial pathways facilitate resilience among

kindergarteners at risk for dyslexia: A longitudinal behavioral and neuroimaging study.

Developmental Science, *24*(1), e12983. https://doi.org/10.1111/desc.12983

Zuk, J., Yu, X., Sanfilippo, J., Figuccio, M. J., Dunstan, J., Carruthers, C., Sideridis, G., Turesky,

T. K., Gagoski, B., Grant, P. E., & Gaab, N. (2021). White matter in infancy is

prospectively associated with language outcomes in kindergarten. *Developmental*

Cognitive Neuroscience, *50*, 100973. https://doi.org/10.1016/j.dcn.2021.100973

1032

1033