Gut microbes contribute to variation in foraging intensity in the honey

bee, *Apis mellifera*.

Cassondra L. Vernier[1], Thi Lan Anh Nguyen[1,a], Tim Gernat[1], Amy Cash Ahmed[1], Zhenqing Chen[1],

Gene E. Robinson[1,2,3]

[1]Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana,

Illinois, USA.

[2] Department of Entomology, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA.

[3] Neuroscience Program, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA.

[a]Current address: Biology Department, Franciscan Missionaries of Our Lady University,

Louisiana, USA.

Corresponding Authors: cvernier@illinois.edu (CLV), generobi@illinois.edu (GER)

Competing Interests

The authors declare no competing interests.

Abstract

Gut microbiomes are increasingly recognized for mediating diverse biological aspects of their hosts, including complex behavioral phenotypes. Many studies have reported that experimental disruptions to the gut microbiome result in atypical host behavior. However, studies that address the role that natural variation in gut microbial communities plays in naturally occurring host behavioral variation are sparse. Here, we show that variation in the gut microbiome influences variation in foraging intensity in honey bees. Gut microbial community structure consistently differed between hive-based nurse bees and bees that leave the hive to forage for floral resources. These differences were associated with variation in the abundance of individual microbes, including *Lactobacillus mellis, Lactobacillus melliventris,* and *Bifidobacterium asteroides*. Manipulations of colony demography and individual foraging experience suggested that differences in microbiome composition were associated with task experience. Moreover, single microbe inoculations with *L. mellis, L. melliventris*, or *B. asteroides* caused changes in foraging intensity. These results demonstrate that gut microbiomes contribute to naturally occurring host behavioral variation, and support a model in which positive feedback interactions cause adaptive changes in both gut microbiome composition and host behavior.

Introduction

Gut microbiomes are emerging as important drivers and modulators of host phenotype [1], with evidence supporting the role of microbiomes in host digestion and nutrition, immune health, development, and, more recently, behavior [2–7]. Over the past decade, the "microbiota-gut-brain axis" [7], which describes bidirectional interactions between the gut microbiome and host brain and behavior, has developed as an increasingly important field of study with implications in understanding the ecology and evolution of host-microbe interactions and animal behaviors. However, current studies of the microbiota-gut-brain axis across animal taxa are largely

correlational, and causal studies exploring the relationship between gut microbes and host

behavior are sparse [7].

The functional relationship between hosts and microbiomes varies across host traits. In

many cases, this relationship is obligatory, where the microbiome is necessary for normal

functioning of the host [7–10]. In other cases, there appears to be a facultative relationship

between host and microbiome, such that differences in microbiome cause differences in

phenotype between individuals [11]. However, many of these host phenotypes result from

dysbiosis or fecal transplants from other species [11, 12], and are either not behavioral [11], or

represent behaviors associated with aging and senescence [13–18], severity of

neurodevelopmental disorders [19–21], or performance in artificial laboratory-based behavioral

tasks [22, 23]. Therefore, studies indicating that natural variation in the gut microbiome can induce

variation in adaptive, naturally occurring host behavior are lacking.

We use the European honey bees, *Apis mellifera,* to investigate the causal relationship

between microbiome and naturally occurring behavior. Honey bees represent a powerful model

to learn more about this relationship due to their complex, yet tractable social behaviors and well-

defined gut microbiome. In typical honey bee colonies, worker bees exhibit age-related division

of labor, which is based on a pattern of individual behavioral maturation. Typically, adult worker

bees perform brood care ("nursing") and other activities in the hive during the first 1-3 weeks of

adult life and then transition to foraging behaviors outside of the hive for the final 1-2 weeks of

their life [24, 25]. However, honey bee division of labor is flexible and responsive to changing

colony needs, and thus does not exclusively rely on the typical age patterns of behavioral

maturation [26–30]. In addition, honey bee foragers exhibit considerable individual variation in

preference for nectar or pollen [31, 32], exploratory behavior [33], and general foraging intensity

[34]. Therefore, honey bee division of labor involves individual variation in both behavioral

maturation and task performance [35].

The honey bee gut microbial community is well-characterized and relatively simple,

composed of 9 taxonomic clusters, including ~14 different species of facultatively anaerobic and

microaerophilic host-adapted bacteria that are consistently present, but may differ in abundance,

across different individuals and populations of honey bees [36]. Of particular interest, the

composition of honey bee gut microbial communities can be experimentally manipulated. All

species can be cultured [37] and used to inoculate young bees [38–40], who must acquire their

microbiome from older bees or hive materials [41]. Furthermore, previous studies indicate an

association between the gut microbiome and various aspects of division of labor in the honey bee

[42–45]. We took advantage of these behavioral and microbiome features to determine whether

gut microbes play a causal role in honey bee division of labor, with a specific focus on behavioral

maturation and variation in foraging intensity.

Materials and Methods

Animal Husbandry, treatments, and collections

Honey bee colonies were managed using standard beekeeping techniques at the University of

Illinois Bee Research Facility in Urbana, IL. Honey bees in this area are a mixture of subspecies,

primarily *Apis mellifera ligustica* and *carnica* subspecies. For collection of nurses and foragers,

nurses were identified as those actively feeding brood on a brood frame, and foragers were

identified as those returning to the hive with pollen loads on their hind legs or having a distended

abdomen due to nectar loads [28, 46]. To reduce genetic variation between workers in the

colonies used for the single-cohort colony, big-back colony and single microbe inoculation studies

(described below), we used bees derived from queens that were each instrumentally inseminated

with sperm from a single drone (SDI). Sample size for 16S sequencing studies was 10 bees per

group as in [47]. All bees used in 16S sequencing analyses were washed once with 12.5% bleach

in water and twice with double deionized water and flash frozen. All samples were stored at -80°C

until further analysis.

Single- and double-cohort colonies

Single-cohort colonies (SCC) were established as previously described [27, 28, 46] using ~1000

newly eclosed bees (<24 hours old). SCCs were kept outdoors, and bees were collected as

typical-age nurses and precocious foragers at approximately one week of age (7-9 days of age),

and as over-age nurses and typical-age foragers at three weeks of age. To obtain over-age

nurses, frames containing brood were removed and replaced with empty comb two weeks after

queen introduction to ensure that a new cohort of bees did not eclose before collections. For the

SCC depicted in Fig. 2 and Supplementary Fig. 1, bees from four SDI source colonies were used

to construct a single SCC. For each collection (Day 9 and Day 21), 10 bees of each source colony

were collected for each behavioral task group. On Day 9, nurses were only able to be collected

for two source colonies. For the SCC depicted in Supplementary Fig. 2, bees from a single SDI

source colony were used, and groups of 10 bees were collected at each collection (Day 7 and

Day 21) for each behavioral task group.

Double-cohort colonies (DCC) were established similarly to SCCs with a new, mated

queen, 1 plastic honeycomb frame provisioned with honey and pollen paste, 1 empty plastic

honeycomb frame, ~200 7-day old barcoded treatment bees (100 per treatment) and ~800 newly

emerged "background" bees (mixed colony backgrounds), whose addition contributed to

accelerated behavioral maturation of the older cohort [46, 48].

Big-back colonies

Big-back colonies were established as previously described [27, 46, 48] and kept outdoors. For

each colony, ~800-900 newly eclosed bees from a single SDI source colony were used; half with

a paint mark on the thorax, and half with a plastic tag attached to the thorax (~3 mm diameter, ~1

mm thick; "big-back" bees). Four days later, ~400 newly eclosed bees from ~10 mixed source

colonies (headed by naturally mated queens) were introduced in order to increase the proportion

of precocious foragers in the first cohort [27, 46, 48]. The entrance to the hive was blocked by a

piece of Plexiglas with holes in it to prevent the big-back bees from leaving the hive, but to allow

paint-marked bees to come and go freely.

Bees were collected at 10 days of age: returning active foragers and nurses collected as

described above, and big-back/inactive forager bees collected as they were attempting to leave

the hive via the holes in the plastic, indicating they too were in the behavioral state of foraging. 10

bees per group were collected from all three colonies, with the exception that only seven nurses

from the focal group in colony 2 were found and collected.

Single microbe inoculations

To identify effects of individual microbes on behavior, we treated groups of bees with either

an inoculum of a microbe of interest or sterile food in order to produce bees whose microbiomes

were composed of a single honey bee-associated microbe (single microbe-inoculated) or no

honey bee-associated microbes (microbiome-depleted control), respectively.

To obtain bees that lack the dominant honey bee bacterial species in the gut, modified

methods from [39, 49] were used. Specifically, tan-colored pupae with dark eyes were gently

removed from brood frames from three or four source colonies per trial using ethanol sterilized

forceps, placed dorsal side down in sterilized 3D-printed dental grade resin pieces fitted into

sterilized Plexiglas cages, and kept in a 34°C incubator with 60-65% relative humidity for 2 days

[50]. Eclosed bees were moved in groups of 30-40 to sterilized Plexiglas boxes (10 x 10 x 7 cm)

[51] in a 32°C incubator at 60-65% relative humidity. Each box was provisioned with a ~1 g

sterilized pollen patty and 1.7 mL sterilized 25% sucrose with or without inoculum. An equal

number of bees from each source colony were added to each box, such that each treatment group

(microbiome-depleted or single microbe inoculated) had an equal number of bees of each colony

background.

For single microbe inoculations, single colonies of bacteria (*B. asteroides* DSM 20089, *L.*

mellis DSM 26255, *L. melliventris* DSM 26256, DSMZ Leibniz-Institut, Germany) were cultured in

de Mann, Rogosa, Sharpe (MRS) (*B. asteroides*) or MRS + 20g/L fructose (*Lactobacillus* species)

broth shaking under anaerobic conditions at 35°C for two nights, spun down (3000 rpm for 5 min)

and resuspended in 1x PBS to an OD of 1. Every day for 5 days, 50 μl of a fresh culture solution

was added to 1.7 mL sterile 25% sucrose water in a new inverted microtube for each treatment

box. For microbiome-depleted bees, 50 μl of sterile 1x PBS was added to 1.7 mL 25% sucrose

water in a new inverted microtube for each treatment box. Bees were then fed sterile 50% sucrose

water for 2 days following the 5-day inoculation period. Bees were kept in treatment boxes until 7

days old, when they were randomly chosen for 16S analysis or behavioral assays. Microbe

survival in 25% sucrose water was ensured by placing 50 μl of each microbial culture (OD ~1) in

1.5 mL 25% sucrose water overnight and then plating this solution using species specific culturing

conditions. 16S rRNA sequencing and qPCR validations of microbiome-depleted bees and single

microbe inoculated bees after the 7-day treatment period are found in Supplementary Fig. 3A-B.

Gut microbiome DNA extraction, 16S rRNA sequencing and analysis

Frozen honey bee guts were dissected under sterile conditions on dry ice. Combined mid- and

hind- guts of individual bees were homogenized by maceration with a disposable sterile pestle

(VWR, Radnor, PA, USA). The homogenate was added to a PowerSoil Pro Bead Solution tube

(Qiagen, Germantown, MD, USA), and DNA was extracted using a DNeasy PowerSoil Pro DNA

isolation kit (Qiagen), following manufacturer's instructions. The hypervariable V4 region of the

16S rRNA gene was amplified by PCR in triplicates, with a negative control (no DNA), using

Platinum Hot Start PCR Master Mix (Invitrogen, Waltham, MA, USA), primers and barcodes

designed in [52] with a final concentration 0.25 μM, and the following cycling conditions: 94° 3

min, 35x[94° 45s/50° 60s/72° 90s], 72° 10 min. Before sequencing, PCR products were visualized

on agarose gels to confirm negative controls did not have amplification, while all samples had

expected amplification. All samples met these criteria, indicating no contamination. Samples were

pooled based upon concentrations and sequenced on an Illumina MiSeq with 2x250bp paired-

end reads. The samples were split among three sequencing runs: #1: SCCs (Dec 2020), #2: Big-

back colonies (July 2021), #3: Typical colony nurses and foragers, single microbe inoculated bees

(Dec 2021). Reads from these sequencing runs can be found in NCBI's Sequence Read Archive

PRJNA958253. Sequences for typical colonies from naturally mated queens were retained from

[43].

Samples were demultiplexed using QIIME2, and paired end reads were truncated at the

first base with a quality score of <Q3 using DADA2 [53]. Paired-end reads were merged and

amplicon sequence variants (ASV) were identified using DADA2 [54]. Chimeric ASVs were

removed and the remaining ASVs were taxonomically classified using the BEExact database [55].

ASVs that were taxonomically identified as a bee specific genus by the BEExact database, but

were unclassified at the species level, were subsequently classified to species level if possible,

using NCBI megaBLAST. To do this, representative 16S V4 sequences for each ASV obtained

from the QIIME2 rep-seqs.qza object were BLASTed against the entire NCBI nucleotide

database, and unclassified ASVs were called as the first identified full species with the lowest E-

score, if query cover > 80% and percent identity > 92%.

For sequencing run #1, 3,220,034 (1,610,017 pairs) total sequence reads were obtained,

1,431,503 pairs were filtered, merged and identified as non-chimeric (90%), and 627 ASVs were

identified. For sequencing run #2, 945,310 (472,655 pairs) total sequence reads were obtained,

424,785 pairs were filtered, denoised, merged and identified as non-chimeric (90%), and 110

ASVs were identified. For sequencing run #3, 2,538,722 (1,269,361 pairs) total sequence reads

were obtained, 1,141,046 pairs were filtered, denoised, merged and identified as non-chimeric
(90%), and 832 ASVs were identified.

To estimate the abundance of individual honey bee-associated microbial species in each
sample, the read counts for all ASVs that matched the same species were combined. Honey bee-
associated species (*Bartonella apis*, *Bifidobacterium asteroides*, *Bifidobacterium coryneforme*
[syn. *Bifidobacterium indicum*), *Bombella apis* [previously *Parasaccharibacter apium*), *Frischella*
perrara, *Gilliamella apicola*, *Lactobacillus apis*, *Lactobacillus helsingborgensis*, *Lactobacillus*
kullabergensis, *Lactobacillus* [syn. *Bombilactobacillus*) *mellifer*, *Lactobacillus* [syn.
Bombilactobacillus) *mellis*, *Lactobacillus melliventris*, and *Snodgrassella alvi*) [36, 56, 57] were
retained. All other ASV reads, representing non-honey bee-associated microbes and each ASV
comprising 0-1% of total reads per sample, were combined and labeled as "other." The variation
between samples in the abundance of "other" microbes is similar to previously published datasets
[38, 44]. To calculate the proportion of each species in each sample, that species' read count was
divided by the total read count for each sample. Raw read counts and/or proportion data were
used in analyses as described below.

For beta-diversity analyses, we used clr-transformations of ASV raw read counts in
statistical tests and Aitchison distance for visualization [58, 59]. To estimate the differential relative
abundance of individual microbial species between samples ("relative abundance"), we used
Analysis of Composition of Microbiomes with Bias Correction (ANCOM-BC) [60] on raw read
counts and visualized using proportion data. To estimate absolute bacterial species abundances
in individual samples ("absolute abundance"), we quantified the bacterial load in each sample
using quantitative PCR (qPCR), as in [38, 61]. We calculated primer efficiencies ($E = 10^{(-1/slope)}$)
and normalized each sample's raw 16S gene copy numbers ($n = E^{(intercept-Ct)}$x(DNA extraction
elution volume/3)) to the *A. mellifera Actin* gene, as in [38, 61]. We then multiplied the normalized
number of 16S rRNA gene copies per sample by the relative proportion of each microbe in that
sample (as determined by 16S rRNA sequencing data) to obtain number of genome equivalents

of each microbe, as in [38]. Log$_{10}$ values of genome equivalents, after replacing 0s with 1s, were

used in statistical analyses.

PCRs were performed in triplicate using 10 μl reactions of 0.5 μM primers targeting the

16S rRNA gene (F: AGGATTAGATACCCTGGTAGTCC, R: YCGTACTCCCCAGGCGG) or the

A. mellifera Actin gene (F: TGCCAACACTGTCCTTTCTG, R: AGAATTGACCCACCAATCCA)

[39] in 1X *Power*SYBR Green PCR Master Mix (Applied Biosystems, Waltham, MA, USA) with 3

μl of DNA. Each primer set included a no template control. Reactions were performed in 384 well

plates using a QuantStudio 6 Flex (Applied Biosystems) with the following cycling conditions: 50°

2 min, 95° 10 min, 40x[95° 15 sec/60° 1 min], melt curve: 95° 15 sec, 60° 1 min, 95° 15 sec.

Standard curves using serial dilutions of plasmids (TOPO pCR2.1 (Invitrogen)) containing the

target sequence (10^8-10^3 copies per 3 μl) were calculated from the molecular weight of the

plasmid and the DNA concentration of the plasmid.

We found an effect of colony replicate on gut microbial community structure across all

experiments, indicating that bees from different colonies have different gut microbial communities

and that early environment and/or genetics influences the composition of the gut microbiome.

This was the case when comparing bees from different typical colonies (Supplementary Fig. 1A-

B), among SCC bees originating from different colony sources (Supplementary Fig. 1C-D), and

across bees from different big-back colonies (Supplementary Fig. 1E). Colony differences in

honey bee microbiome composition have been previously reported [47].

Foraging assays and analysis using barcoded bees

To obtain comprehensive records of foraging activity, we used an automated behavioral tracking

system ("bCode") that uses a custom matrix barcode, enabling the unique identification of

individual bees [62]. After single microbe inoculations, 7-day old inoculated and microbiome-

depleted bees were cold-anaesthetized, a unique barcode was affixed to their thorax using super

glue, and a spot of colored paint associated with their treatment group was placed on their

abdomen. An equal number (~100) of bees from a single microbe inoculated group and a

corresponding microbiome-depleted group (same mix of colony backgrounds, same age) were

given barcodes from different sets, and were placed together in a DCC. Four colony replicates

per microbe treatment were performed. Two experimental days, corresponding with *L. mellis*

colony replicate 3 days 1-2 and *L. melliventris* colony replicate 2 days 1-2, experienced bad

weather, and thus no foraging occurred on those days.

Colonies were kept in a dark, temperature (32°C) and humidity (~50%) controlled building,

with access to the outside environment through a plastic tube. To monitor flight activity, an

entrance monitor with a video camera (Supplementary Methods) was attached to the hive

entrance, on the outside-facing end of this tube (Supplementary Fig. 4). The camera recorded

videos of bees entering and leaving the hive from 05:00 to 21:00 daily for a total of six days.

Barcodes in these videos were detected as in [62], and detections were filtered to remove tracking

errors and misidentifications. A mean of 97.82%, with a range between 96.24%-99.02%, of

detections were retained after these filtering steps. We then applied a flight activity detector [63,

64] (Supplementary Methods) to the remaining barcode detections to identify passes through the

entrance   monitor.   Computer   code   for   this   detector   can   be   found   at:

https://github.com/gernat/btools.

Passes through the entrance monitor determined by the flight activity detector were filtered

to remove unused barcodes, whose detections either occurred through detection error or due to

bee drift between experimental colonies. A mean of 99.19%, with a range of 95.4%-100%, of

events were retained after this filtering. Passes were then classified as "incoming," "outgoing," or

"other." Since incoming passes had a lower error rate than outgoing passes (Supplementary

Table 8), incoming passes alone were used to denote a foraging trip. Individual foraging trips that

occurred within 5 min of each other were condensed into a single event. A bee's first day of

foraging was defined as the first day on which it performed at least 4 foraging trips, with at least

50% of these trips occurring during peak foraging hours (11:00-15:00 CST), as per previous

studies [34, 65]. These criteria yield similar automated thresholds corresponding to human

observations of foraging behaviors [34]. All incoming passes from this first day of foraging were

counted as foraging trips, as were all subsequent incoming passes [34]. Likewise, an individual

bee was considered a "forager" from this day forward.

To quantify the degree of skew in foraging intensity among all workers, we calculated the

Gini coefficient [34, 66] for each experimental colony across all days and each experimental

colony on each day, since colony dynamics change with changes in weather and resource

availability, and thus likely changed throughout our studies. The values we obtained (Table 2) are

similar to those in previous studies [34, 66], indicating differences in foraging intensity between

individuals in each colony. To determine which bees performed the majority of the foraging for

each experimental colony on each day, we ranked individuals based on the proportion of total

foraging trips they performed for the colony on each day. A group of "elite foragers" was defined

as the subset of bees (variation, 5-40% of the foragers) performing $\geq$50% of the foraging trips for

each colony on each day [34, 66].

Although we monitored our experimental colonies for six days, we reasoned that any

behavioral effects seen after the first several days of monitoring may be confounded by changes

in colony dynamics due to forager mortality [67], "homogenization" of the microbiome among

colony members due to trophallaxis (sharing of liquids containing nutrients and signaling

molecules), or acquisition of honey bee-specific microbes from the environment. While we did not

test how long it takes for possible changes in microbiome due to these factors to occur, we chose

to focus on results for the first half (three days) of the experimental period below. Summary data

and statistics for the entire six-day period can be found in Supplementary Table 6.

Statistical Analysis

All statistical analyses were performed in R (v 4.2.0) [68]. For all analyses, assumptions (e.g.,

normality, homogeneity of variances) were checked before statistical analysis. The following

functions were used in the base statistics package: "shapiro.test" (Shapiro-Wilk Normality Test),

"fligner.test" (Fligner-Killeen Test of Homogeneity of Variances), "t.test" (Student's t-test),

"wilcox.test" (Mann-Whitney Rank Sum Test), "aov" (ANOVA), "TukeyHSD" (Tukey's HSD post

hoc test), "Kruskal.test" (Kruskal-Wallis test). Kruskal-Wallis post hoc was performed using

"kwAllPairsDunnTest", PMCMRplus package [69]. Microbiome beta diversity was analyzed using

Permutation MANOVAs with 999 permutations ("adonis2", vegan package [70]) on clr-

transformed ("transform", microbiome package [71]) raw read counts, followed by Pairwise

Permutation MANOVAs with 999 permutations and FDR p-value adjustment ("pairwise.adonis",

pairwiseAdonis package [72]), and visualized using Principal Components Analysis ("ordinate",

phyloseq package [73]) with Aitchison distance ("distance", phyloseq package [73]). The relative

abundance of individual microbes was analyzed using raw read counts and ANCOM-BC with FDR

adjustment ("ancombc2", ANCOMBC package [60]) with task as a fixed effect and colony as

random effect. Relative abundances were visualized using stacked barplots ("ggplot", ggplot2

package [74]). To compare the absolute abundance of individual microbes, Permutation ANOVAs

were used with 999 permutations ("perm.anova," RVAideMemoire package [75]), and p-values

were adjusted for multiple comparisons ("p.adjust" with FDR adjustment). Absolute abundance

data were visualized using dotplots ("ggplot," ggplot2 package [74]). Age at onset of foraging data

(using each individual bee's first day of foraging) were analyzed using a Cox Proportional Hazards

model ("coxph," survival package [76]), stratified by replicate [26, 65], and depicted as survival

plots using the cumulative proportion of bees from each treatment group that were identified as

foragers on each day. Proportions of foraging trips per individual were analyzed as generalized

linear mixed-effects models ("glmer," lme4 package [77]) with log-normal distributions and

nAGQ=0, inoculation treatment and day as main factors, and replicate and individual as random

factors, followed by "Anova" (car package [78]) to determine main factor significance, and

"emmeans" (emmeans package [79]) with Tukey's p-value adjustment for pairwise comparisons.

Two-way mixed ANOVAs with inoculation treatment as a between-subject factor and day as a

within-subjects factor, followed by pairwise one-way ANOVAs, were used in the remaining

behavioral analyses ("anova_test," rstatix package [80]). Gini coefficients were calculated using

"Gini" (DescTools package [81]). All behavioral data, except age at onset of foraging, were

visualized using boxplots ("ggplot," ggplot2 package [74]).

Results

Nurses and foragers differ in gut microbial community composition

To begin to investigate whether gut microbes influence honey bee division of labor, we compared

gut microbial communities in nurse and forager bees, which represent two of the canonical

behavioral task groups in honey bee division of labor, as well as two discrete time points in

behavioral maturation [25]. Nurses and foragers differ markedly in physiology [28, 82–87],

neuroanatomy [88], neurochemistry [33, 89, 90], gene expression [29, 91] and gene regulation

[65, 92–94]. Therefore, we reasoned that testing for consistent differences in gut microbial

community between these two groups would be a powerful way to test for associations between

gut microbiome variation and division of labor.

We performed 16S rRNA sequencing on gut samples from nurses and foragers from three

unrelated honey bee colonies and reanalyzed, using updated methods [53, 54], a similar dataset

from a previously published study that used a different 16S sequencing method [43]. We found

that nurses and foragers differed significantly in gut microbial community structure in both the

previously published dataset (Fig. 1A) and our new dataset (Fig. 1B), indicating robust differences

associated with division of labor and independent of sequencing method.

We next sought to identify individual microbes that differed in abundance between nurse

and forager bees, in order to identify candidate microbes specifically associated with division of

labor. Since microbes can vary between individuals in relative and/or absolute abundance, two

measures that are not necessarily linked and whose relative effects on host behavior are currently

unknown [61], we used both of these abundance measures to get a well-rounded sense of

individual microbes that are associated with honey bee division of labor. We found that individual

gut microbes significantly differed in both relative and absolute abundance between nurses and

foragers (Fig. 1C-D, Supplementary Table 1, Table 1). These results indicate that nurse and

forager bees differ significantly in their gut microbial communities. They also identify candidate

microbes that may be associated with division of labor.

357 Differences in gut microbial community composition between nurses and foragers

358 are dependent upon differences in behavioral state, not age

Under typical colony conditions, nurses and foragers differ in chronological age. However, honey

bee division of labor is flexible and responsive to changing colony needs. For example, if a colony

experiences a shortage of older bees, some younger individuals will begin to forage at an early

age, while a shortage of younger bees will cause bees to continue to act as nurse bees despite

advancing chronological age [26–30]. To determine whether the gut microbial differences

between nurses and foragers depend on differences in behavioral state and/or age, we exploited

this adaptive plasticity to create SCCs, a well-established experimental approach that separates

these two factors [27, 28, 46]. In two SCC replicates, we compared gut microbiomes between

age-matched nurse and forager bees when they were about one week of age (representing

typical-age nurses and precocious foragers) and three weeks of age (representing over-age

nurses and typical-age foragers).

Age-matched typical-age nurses and precocious foragers did not significantly differ in gut

microbial community structure (Fig. 2A, Supplementary Fig. 2A), while age-matched over-age

nurses and typical-age foragers did (Fig. 2B, Supplementary Fig. 2B). Similar results were

obtained for the abundance of individual microbial species: no microbes consistently differed in

abundance between typical-age nurses and precocious foragers across the two SCC replicates

(Fig. 2C, Fig. 2E, Supplementary Fig. 2C, Supplementary Fig. 2E, Supplementary Table 2,

Supplementary Table 3, Table 1), while two microbes consistently differed in relative abundance

and three microbes consistently differed in absolute abundance between over-age nurses and

typical-age foragers across the two SCC replicates (Fig. 2D, Fig. 2F, Supplementary Fig. 2D,

Supplementary Fig. 2F, Supplementary Table 2, Supplementary Table 3, Table 1). These findings

indicate that age-dependent differences in behavioral state are associated with differences in gut

microbial communities between nurses and foragers.

Together, our typical colony and three-week old SCC data indicate that *Lactobacillus*

mellis and *Lactobacillus melliventris* were consistently (i.e. identified in three or more analyses

across the two abundance measures) more abundant in nurses relative to foragers (Table 1),

while *Bartonella apis* and *Snodgrassella alvi* were consistently less abundant in nurses relative to

foragers (Table 1). In SCCs alone, *Bifidobacterium asteroides* showed consistent lower relative

abundance in over-age nurses compared to typical-age foragers (Table 1) and *Gilliamella apicola*

(Table 1) showed consistent higher absolute abundance in over-age nurses compared to typical-

age foragers. Such robust associations highlight these six microbes as candidate microbes whose

abundances have causal effects on division of labor in honey bees.

The observed age-dependent microbiome differences between nurses and foragers

suggest that associations between gut microbiome and behavioral state depend on worker

experience. Since SCCs are composed of bees of a single age cohort, three-week old SCCs are

composed of bees that have been performing their respective tasks for prolonged periods (1-2

weeks longer) compared to one-week old SCCs. Furthermore, the lack of gut microbial community

differences between one-week-old SCC nurses and foragers suggests that differences in

behavioral maturation rate are not associated with differences in gut microbial community

composition. Therefore, we hypothesized that task experience plays an important role in gut

microbial community composition. In the next section, we tested this by focusing on foraging

experience since it is known to have strong effects on behavior, brain chemistry, brain structure,

and brain gene expression [66, 89, 95, 96].

Foraging experience influences gut microbial community structure, while

behavioral state influences the abundance of individual microbes

In order to determine the causal effects of behavioral state and foraging experience on honey bee

gut microbiome composition, we controlled for behavioral state while manipulating foraging

experience. To do this, we created "big-back colonies" in which some foragers were freely able

to leave the colony ("active foragers"), while other bees, who appeared at the entrance and

showed an inclination to forage, were prevented from leaving due to the presence of a thick plastic

tag on their backs ("inactive foragers") [46, 48]. We were thus able to compare bees of the same

age and behavioral state that differed in foraging experience.

We found that nurses and inactive foragers did not differ in gut microbial community

structure, while active foragers and inactive foragers did (Fig. 3A). This finding indicates that

foraging experience plays an important role in defining differences in gut microbial community

structure. In contrast to these community structure results, which take below-species-level

diversity into account [54], we did not find any microbial species that differed in relative or absolute

abundance between age-matched active and inactive foragers (Fig. 3B-C, Supplementary Table

4). This suggests that the abundance of individual microbes is defined by behavioral state, and

not experience foraging outside of the hive. Although these results are seemingly contradictory to

our SCC data, which indicate that differences in the abundance of individual gut microbes may

be dependent upon worker experience (Fig. 2D,F, S2 Fig. D,F), it is possible that honey bee gut

microbes and foraging behaviors interact through a feedback model. For example, gut microbes

may change in abundance due to foraging experience, possibly incrementally (and thus

indiscernibly) at first and building with more experience, and this change in microbiome may then

feed back to influence host foraging behaviors. In the next section, we test one part of this model

by assessing the effects of inoculation with individual microbes on host foraging behavior.

Inoculation with individual microbes affects group- and individual-level foraging

intensity

To determine whether variations in gut microbiome composition play a causal role in

honey bee division of labor, we performed single microbe inoculations with three out of the six

individual microbes robustly associated with division of labor as described above: *B. asteroides*,

L. mellis and *L. melliventris*. Although it is a simplification of the natural microbiome, we chose to

test the effects of each microbe in isolation in order to increase experimental power. In addition

to the strong patterns of association reported above, we chose these three microbes due to *B.*

asteroides' known effects on host physiology with links to behavior and the *Lactobacillus* species'

high metabolic output [39], since gut microbial metabolites likely contribute to the microbiota-gut-

brain axis [7]. Using an automated tracking system [62], we determined the effect of inoculation

with these three microbes on two measures of foraging behavior important in division of labor in

honey bees: rate of behavioral maturation (age at onset of foraging) and foraging intensity

(amount of foraging trips performed by a group or individual) [35].

B. asteroides inoculated bees did not differ from microbiome-depleted bees in behavioral

maturation rate (Fig. 4A) across experimental replicates, but did differ in foraging intensity.

Overall, *B. asteroides* inoculated bees performed the majority of the foraging trips for the colony

on the first day after colony formation (Fig. 4B, Supplementary Table 5). These group-level effects

were not due to a difference in number of foragers between inoculated and microbiome-depleted

bees (Fig. 4C, Supplementary Table 5). Rather they were due to a difference in individual-level

foraging intensity, as individual *B. asteroides* inoculated foragers performed a majority of the

foraging trips for the colony on the first day after colony formation (Fig. 4D, Supplementary Table

5). Likewise, *B. asteroides* experimental colonies displayed a degree of skew in foraging intensity

among individuals (Table 2) and *B. asteroides* inoculated bees represented a higher proportion

of "elite forager" bees, i.e., a small subset of foragers that performed $\geq$50% of the colony's foraging

trips [34, 66], relative to microbiome-depleted bees on the first two days after colony formation

(Fig. 4E, Supplementary Table 5).

Similar to *B. asteroides* inoculated bees, *L. mellis* inoculated bees did not differ from

microbiome-depleted bees in behavioral maturation rate (Fig. 5A), but did differ in foraging

intensity. However, *L. mellis* inoculation caused an opposite effect on foraging intensity compared

to *B. asteroides* inoculation. Overall, *L. mellis* inoculated bees performed the minority of foraging

trips for the colony on the third day after colony formation (Fig. 5B, Supplementary Table 5).

Again, these group-level effects were not due to a difference in number of foragers between

inoculated and microbiome-depleted bees (Fig. 5C, Supplementary Table 5). Rather they were

due to a difference in individual-level foraging intensity, as *L. mellis* inoculated bees performed a

minority of foraging trips for the colony on the first day after colony formation (Fig. 5D,

Supplementary Table 5). Additionally, they could also be due to skew in foraging intensity

between individual foragers (Table 2), as *L. mellis* inoculated bees displayed a trend (with

marginal significance on the second day, Supplementary Table 5) of representing a lower

proportion of elite foragers than microbiome-depleted bees on all three days after colony

formation (Fig. 5E, Supplementary Table 5).

L. melliventris inoculated bees showed similar results to *L. mellis* inoculated bees. *L.*

melliventris inoculated bees did not differ from microbiome-depleted bees in behavioral

maturation rate (Fig. 6A), but did differ in foraging intensity. Overall, *L. melliventris* inoculated

bees performed the minority of foraging trips for the colony on the second day after colony

formation (Fig. 6B, Supplementary Table 5). Yet again, this group-level effect was not due to a

difference in number of foragers between inoculated and microbiome-depleted bees (Fig. 6C,

Supplementary Table 5). Rather, it was due to a difference in individual-level foraging intensity,

as individual *L. melliventris* inoculated foragers performed a minority of the foraging trips for the

colony on the second day after colony formation (Fig. 6D, Supplementary Table 5). Likewise, *L.*

melliventris colonies displayed a degree of skew in foraging intensity between individual foragers

(Table 2), and *L. melliventris* bees represented a lower proportion of elite foragers than

microbiome-depleted bees on the second day after colony formation (Fig. 6E, Supplementary

Table 5).

Overall, these results indicate that inoculation with individual gut microbes is sufficient to

cause increases or decreases in foraging intensity between worker bees at both the group- and

individual- level.

Discussion

Here, we show that natural variation in the gut microbiome can induce variation in a

naturally occurring host behavior. Specifically, our results indicate that differences in honey bee

gut microbial communities influence honey bee division of labor by affecting foraging intensity,

but not rate of behavioral maturation. This conclusion is supported by both microbe abundance

data and behavioral data: *B. asteroides* is found in higher abundance in foragers compared to

nurses, and inoculation with this species caused bees to have increased foraging intensity.

Similarly, *L. mellis* and *L. melliventris* are found in lower in abundance in foragers compared to

nurses, and inoculation with either of these species caused bees to have decreased foraging

intensity.

From a technical perspective, the observation that single microbe inoculations can cause

an increase or a decrease in foraging intensity dependent upon the microbe indicates that it is

unlikely that the results reported here are due to an artifact of the microbiological or behavioral

methods we used. In addition, the congruence of our microbe abundance and behavioral data

indicate that single microbe inoculations, which do not represent natural microbiome variations,

provide a useful method for functional analysis. Together, our results suggest that naturally

occurring increases in *B. asteroides*, which occur simultaneously with naturally occurring

decreases in *L. mellis* and *L. melliventris,* act synergistically to increase the intensity of foraging

behavior in individual bees. Honey bees, with a complex eusocial life history, live in large perennial

colonies and therefore collect large amounts of nectar and pollen to ensure colony survival during

periods when floral resources are not available. Thus, the influence of these three microbes on

honey bee foraging intensity indicates that host-microbe interactions likely help sustain the

perennial lifestyle of this eusocial insect.

While previous studies report correlational differences in gut microbiome associated with

honey bee behavioral task [42, 44, 61] and behavioral maturation [45], here, we provide evidence

that gut microbiomes play a causal role in defining behavioral differences between honey bees.

Given the age-related nature of honey bee division of labor, this research also shows that gut

microbiomes can drive adaptive variations in host behavioral phenotypes over the lifetime of

individual animals.

Our findings suggest that positive feedback interactions cause adaptive changes in both

gut microbiome composition and host behavior, such that changes in gut microbe abundance and

host behavior interact to strengthen one another and increase foraging intensity. Modeling

interactions in this way can provide the basis for future studies that probe the neural and metabolic

mechanisms by which *B. asteroides*, *L. mellis* and *L. melliventris* affect host behavior. Overall,

our findings highlight the importance of feedback mechanisms in microbiota-gut-brain

interactions, provide evidence that variation in gut microbiomes can drive variation in naturally

occurring behaviors, and support gut microbiomes as important drivers of host phenotypic

variation and plasticity.

Previous studies have shown that gut microbes may influence host behavior through a

variety of mechanisms including production of substances that cause changes in host brain gene

expression or host production of neurotransmitters and hormones [7]; these general mechanisms

are also associated with the regulation of honey bee division of labor [33, 35, 65, 87–90, 93, 95,

97, 98]. Likewise, previous studies indicate that some host factors associated with honey bee

division of labor, such as social interactions and diet [99, 100], may also modulate gut microbiome

composition [7]. Therefore, future research will work to identify specific mechanisms by which

honey bees and their gut microbiomes interact. Overall, studies using the honey bee as a model

have the potential to further elucidate the microbiota-gut-brain axis, and achieve a more

comprehensive understanding of interkingdom interactions.

535 Acknowledgements

We thank Nathan Beach, Sarah Magdalena Murphree, Sarai Stuart, Laura Kilikevicius, Rashmi

Bajaj, Maha Syed and Anthony Cantu for field assistance, Sue Cobey and Dr. Osman Kaftanoglu

for SDI queen rearing and inseminations, Ahmed Burki for assistance making qPCR standard

plasmids, Alvaro Hernandez, Chris Wright and staff at the Carver Biotechnology Center for

sequencing services, members of the Computer Network Resource Group of the Carl R. Woese

Institute for Genomic Biology (UIUC) for computational support, Konrad Wilk for annotating

missed barcodes, Richard B. Devine for contributions to video annotation software development,

and the workers of Turker Nation for annotating honey bee trajectories. Funding: US National

Science Foundation Division of Integrative Organismal Systems award number 2120378, PI:

GER.

Competing Interests

The authors declare no competing interests.

Data Availability Statement: 16S rRNA sequencing data are available from the NCBI Sequence

Read Archive (PRJNA958253) and all other data are available from Mendeley Data (DOI:

10.17632/f2s47y3nhn). Computer code for the barcode detector is available at

https://github.com/gernat/btools.

References

1. Zilber-Rosenberg I, Rosenberg E. Role of microorganisms in the evolution of animals and
plants: the hologenome theory of evolution. *FEMS Microbiol Rev* 2008; **32**: 723–735.
2. Cryan JF, Dinan TG. Mind-altering microorganisms: the impact of the gut microbiota on
brain and behaviour. *Nat Rev Neurosci* 2012; **13**: 701–712.
3. Flint HJ, Karen P, Louis P, Duncan SH. The role of the gut microbiota in nutrition and
health. *Nat Rev Gastroenterol Hepatol* 2012; **9**: 577–589.
4. Hooper L V., Littman DR, Macpherson AJ. Interactions Between the Microbiota and the
Immune System. *Science (1979)* 2012; 1268–1274.
5. Münger E, Montiel-Castro AJ, Langhans W, Pacheco-López G. Reciprocal Interactions
Between Gut Microbiota and Host Social Behavior. *Front Integr Neurosci* 2018; **12**: 21.
6. Ezenwa VO, Gerardo NM, Inouye DW, Medina M, Xavier JB. Animal Behavior and the
Microbiome. *Science (1979)* 2012; **338**: 198–199.
7. Sherwin E, Bordenstein SR, Quinn JL, Dinan TG, Cryan JF. Microbiota and the social brain.
Science (1979) 2019; **366**: 587.
8. Diaz Heijtz R, Wang S, Anuar F, Qian Y, Björkholm B, Samuelsson A, et al. Normal gut
microbiota modulates brain development and behavior. *Proceedings of the National
Academy of Sciences* 2011; **108**: 3047–3052.
9. Neufeld KM, Kang N, Bienenstock J. Reduced anxiety-like behavior and central
neurochemical change in germ-free mice. *Neurogastroenterology & Motility* 2011; **23**:
255–265.
10. Degroote S, Hunting DJ, Baccarelli AA, Takser L. Maternal gut and fetal brain connection:
Increased anxiety and reduced social interactions in Wistar rat offspring following peri-
conceptional antibiotic exposure. *Prog Neuropsychopharmacol Biol Psychiatry* 2016; **71**:
76–82.
11. Henry LP, Bruijning M, Forsberg SKG, Ayroles JF. The microbiome extends host
evolutionary potential. *Nat Commun* 2021; **12**: 5141.
12. Trevelline BK, Kohl KD. The gut microbiome influences host diet selection behavior.
Proceedings of the National Academy of Sciences 2022; **119**.
13. Shoji H, Takao K, Hattori S, Miyakawa T. Age-related changes in behavior in C57BL/6J
mice from young adulthood to middle age. *Mol Brain* 2016; **9**: 11.
14. Claesson MJ, Jeffery IB, Conde S, Power SE, O'Connor EM, Cusack S, et al. Gut microbiota
composition correlates with diet and health in the elderly. *Nature* 2012; **488**: 178–184.
15. Smith P, Willemsen D, Popkes M, Metge F, Gandiwa E, Reichard M, et al. Regulation of
life span by the gut microbiota in the short-lived African turquoise killifish. *Elife* 2017; **6**:
1–26.

16. Liu H, Wang X, Wang HD, Wu J, Ren J, Meng L, et al. Escherichia coli noncoding RNAs can
affect gene expression and physiology of Caenorhabditis elegans. *Nat Commun* 2012; **3**:
1–11.
17. Erny D, Prinz M. How microbiota shape microglial phenotypes and epigenetics. *Glia* 2020;
68: 1655–1672.
18. Canipe LG, Sioda M, Cheatham CL. Diversity of the gut-microbiome related to cognitive
behavioral outcomes in healthy older adults. *Arch Gerontol Geriatr* 2021; **96**: 104464.
19. Taniya MA, Chung H-J, Al Mamun A, Alam S, Aziz MdA, Emon NU, et al. Role of Gut
Microbiome in Autism Spectrum Disorder and Its Therapeutic Regulation. *Front Cell
Infect Microbiol* 2022; **12**.
20. Kelly JR, Minuto C, Cryan JF, Clarke G, Dinan TG. Cross Talk: The Microbiota and
Neurodevelopmental Disorders. *Front Neurosci* 2017; **11**.
21. Checa-Ros A, Jeréz-Calero A, Molina-Carballo A, Campoy C, Muñoz-Hoyos A. Current
Evidence on the Role of the Gut Microbiome in ADHD Pathophysiology and Therapeutic
Implications. *Nutrients* 2021; **13**: 249.
22. Li L, Solvi C, Zhang F, Qi Z, Chittka L, Zhao W. Gut microbiome drives individual memory
variation in bumblebees. *Nat Commun* 2021; **12**: 6588.
23. Davidson GL, Wiley N, Cooke AC, Johnson CN, Fouhy F, Reichert MS, et al. Diet induces
parallel changes to the gut microbiota and problem solving performance in a wild bird.
Sci Rep 2020; **10**: 20783.
24. Smith CR, Toth AL, Suarez A V., Robinson GE. Genetic and genomic analyses of the
division of labour in insect societies. *Nat Rev Genet* 2008; **9**: 735–748.
25. Robinson GE. Regulation of Division Of Labor In Insect Societies. *Annu Rev Entomol* 1992;
37: 637–665.
26. Ben-Shahar Y, Robichon A, Sokolowski MB, Robinson GE. Influence of Gene Action Across
Different Time Scales on Behavior. *Science (1979)* 2002; **296**: 741–744.
27. Vernier CL, Krupp JJ, Marcus K, Hefetz A, Levine JD, Ben-Shahar Y. The cuticular
hydrocarbon profiles of honey bee workers develop via a socially-modulated innate
process. *Elife* 2019; **8**: e41855.
28. Robinson GE, Page RE, Strambi C, Strambi A. Hormonal and Genetic Control of Behavioral
Integration in Honey Bee Colonies. *Science (1979)* 1989; **246**: 109–112.
29. Whitfield CW, Cziko A-M, Robinson GE. Gene Expression Profiles in the Brain Predict
Behavior in Individual Honey Bees. *Science (1979)* 2003; **302**: 296–299.
30. Ben-Shahar Y, Dudek NL, Robinson GE. Phenotypic deconstruction reveals involvement of
manganese transporter malvolio in honey bee division of labor. *Journal of Experimental
Biology* 2004; **207**: 3281–3288.
31. Calderone NW, Page RE. Effects of interactions among genotypically diverse nestmates
on task specialization by foraging honey bees (Apis mellifera). *Behav Ecol Sociobiol* 1992;
30: 219–226.
32. Pankiw T, Page Jr RE. Response thresholds to sucrose predict foraging division of labor in
honeybees. *Behav Ecol Sociobiol* 2000; **47**: 265–267.
33. Cook CN, Mosqueiro T, Brent CS, Ozturk C, Gadau J, Pinter-Wollman N, et al. Individual
differences in learning and biogenic amine levels influence the behavioural division

between foraging honeybee scouts and recruits. *Journal of Animal Ecology* 2019; **88**:
236–246.

34.   Tenczar P, Lutz CC, Rao VD, Goldenfeld N, Robinson GE. Automated monitoring reveals
extreme interindividual variation and plasticity in honeybee foraging activity levels. *Anim*
Behav 2014; **95**: 41–48.

35.   Robinson GE. Effects of a juvenile hormone analogue on honey bee foraging behaviour
and alarm pheromone production. *J Insect Physiol* 1985; **31**: 277–282.

36.   Kwong WK, Moran NA. Gut microbial communities of social bees. *Nat Rev Microbiol*
2016; **14**: 374–384.

37.   Engel P, James RR, Koga R, Kwong WK, McFrederick QS, Moran NA. Standard methods for
research on Apis mellifera gut symbionts. *J Apic Res* 2013; **52**: 1–24.

38.   Liberti J, Kay T, Quinn A, Kesner L, Frank ET, Cabirol A, et al. The gut microbiota affects
the social network of honeybees. *Nat Ecol Evol* 2022; **6**: 1471–1479.

39.   Kešnerová L, Mars RAT, Ellegaard KM, Troilo M, Sauer U, Engel P. Disentangling
metabolic functions of bacteria in the honey bee gut. *PLoS Biol* 2017; **15**: e2003467.

40.   Leonard SP, Powell JE, Perutka J, Geng P, Heckmann LC, Horak RD, et al. Engineered
symbionts activate honey bee immunity and limit pathogens. *Science (1979)* 2020; **367**:
573–576.

41.   Powell JE, Martinson VG, Urban-Mead K, Moran NA. Routes of Acquisition of the Gut
Microbiota of the Honey Bee Apis mellifera. *Appl Environ Microbiol* 2014; **80**: 7378–7387.

42.   Jones JC, Fruciano C, Marchant J, Hildebrand F, Forslund S, Bork P, et al. The gut
microbiome is associated with behavioural task in honey bees. *Insectes Soc* 2018; **65**:
419–429.

43.   Kapheim KM, Rao VD, Yeoman CJ, Wilson BA, White BA, Goldenfeld N, et al. Caste-
Specific Differences in Hindgut Microbial Communities of Honey Bees (Apis mellifera).
PLoS One 2015; **10**: e0123911.

44.   Copeland DC, Maes PW, Mott BM, Anderson KE. Changes in gut microbiota and
metabolism associated with phenotypic plasticity in the honey bee Apis mellifera. *Front*
Microbiol 2022; **13**.

45.   Ortiz-Alvarado Y, Clark DR, Vega-Melendez CJ, Flores-Cruz Z, Domingez-Bello MG, Giray T.
Antibiotics in hives and their effects on honey bee physiology and behavioral
development. *Biol Open* 2020; **9**.

46.   Ben-Shahar Y, Thompson CK, Hartz SM, Smith BH, Robinson GE. Differences in
performance on a reversal learning test and division of labor in honey bee colonies. *Anim*
Cogn 2000; **3**: 119–125.

47.   Vernier CL, Chin IM, Adu-Oppong B, Krupp JJ, Levine J, Dantas G, et al. The gut
microbiome defines social group membership in honey bee colonies. *Sci Adv* 2020; **6**:
eabd3431.

48.   Withers GS, Fahrbach SE, Robinson GE. Effects of experience and juvenile hormone on
the organization of the mushroom bodies of honey bees. *J Neurobiol* 1995; **26**: 130–144.

49.   Emery O, Schmidt K, Engel P. Immune system stimulation by the gut symbiont Frischella
perrara in the honey bee (Apis mellifera). *Mol Ecol* 2017; **26**: 2576–2590.

50. Fine JD, Shpigler HY, Ray AM, Beach NJ, Sankey AL, Cash-Ahmed A, et al. Quantifying the
effects of pollen nutrition on honey bee queen egg laying with a new laboratory system.
PLoS One 2018; **13**: 1–16.
51. Galbraith DA, Yang X, Niño EL, Yi S, Grozinger C. Parallel Epigenomic and Transcriptomic
Responses to Viral Infection in Honey Bees (Apis mellifera). *PLoS Pathog* 2015; **11**:
e1004713.
52. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-high-
throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms.
ISME J 2012; **6**: 1621–1624.
53. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. *Nat Methods* 2016; **13**: 581–
583.
54. Callahan BJ, McMurdie PJ, Holmes SP. Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. *ISME J* 2017; **11**: 2639–2643.
55. Daisley BA, Reid G. BEExact: a Metataxonomic Database Tool for High-Resolution
Inference of Bee-Associated Microbial Communities. *mSystems* 2021; **6**.
56. Smith EA, Anderson KE, Corby-Harris V, McFrederick QS, Parish AJ, Rice DW, et al.
Reclassification of seven honey bee symbiont strains as Bombella apis. *Int J Syst Evol*
Microbiol 2021; **71**.
57. Olofsson TC, Alsterfjord M, Nilson B, Butler È, Vásquez A. Lactobacillus apinorum sp. nov.,
Lactobacillus mellifer sp. nov., Lactobacillus mellis sp. nov., Lactobacillus melliventris sp.
nov., Lactobacillus kimbladii sp. nov., Lactobacillus helsingborgensis sp. nov. and
Lactobacillus kullabergensis sp. nov., isolated from the honey stomach of the honeybee
Apis mellifera. *Int J Syst Evol Microbiol* 2014; **64**: 3109–3119.
58. Aitchison J. The Statistical Analysis of Compositional Data. 1986. Springer Netherlands,
Dordrecht.
59. Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. Microbiome Datasets Are
Compositional: And This Is Not Optional. *Front Microbiol* 2017; **8**.
60. Lin H, Peddada S Das. Analysis of compositions of microbiomes with bias correction. *Nat*
Commun 2020; **11**: 3514.
61. Kešnerová L, Emery O, Troilo M, Liberti J, Erkosar B, Engel P. Gut microbiota structure
differs between honeybees in winter and summer. *ISME J* 2020; **14**: 801–814.
62. Gernat T, Rao VD, Middendorf M, Dankowicz H, Goldenfeld N, Robinson GE. Automated
monitoring of behavior reveals bursty interaction patterns and rapid spreading dynamics
in honeybee social networks. *Proceedings of the National Academy of Sciences* 2018;
115: 1433–1438.
63. Geffre AC, Gernat T, Harwood GP, Jones BM, Morselli Gysi D, Hamilton AR, et al. Honey
bee virus causes context-dependent changes in host social behavior. *Proceedings of the*
National Academy of Sciences 2020; **117**: 10406–10413.
64. Liaw A, Wiener M. Classification and Regression by randomForest. *R News* 2002; **2**: 18–
22.
65. Hamilton AR, Traniello IM, Ray AM, Caldwell AS, Wickline SA, Robinson GE. Division of
labor in honey bees is associated with transcriptional regulatory plasticity in the brain.
Journal of Experimental Biology 2019; **222**.

66.   Klein S, Pasquaretta C, He XJ, Perry C, Søvik E, Devaud J-M, et al. Honey bees increase
their foraging performance and frequency of pollen trips through experience. *Sci Rep*
2019; **9**: 6778.
67.   Rueppell O, Bachelier C, Fondrk MK, Page RE. Regulation of life history determines
lifespan of worker honey bees (Apis mellifera L.). *Exp Gerontol* 2007; **42**: 1020–1032.
68.   R Core Team. R: A language and environment for statistical computing. 2022. R
Foundation for Statistical Computing, Vienna, Austria.
69.   Pohlert T. The Pairwise Multiple Comparison of Mean Ranks Package (PMCMR). 2014. R
package.
70.   Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan:
Community Ecology Package. 2017. R package version 2.4-5.
71.   Lahti L, Shetty S. microbiome R package. 2019.
72.   Martinez Arbizu P. pairwiseAdonis: Pairwise multilevel comparison using adonis. 2020. R
package version 0.4.
73.   McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis
and Graphics of Microbiome Census Data. *PLoS One* 2013; **8**: e61217.
74.   Wickham H. Elegant graphics for data analysis. 2016. Springer-Verlag New York.
75.   Hervé M. RVAideMemoire: Testing and plotting procedures for biostatistics. 2022. R
package.
76.   Therneau TM. A package for survival analysis in R. 2022. R package.
77.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. *J*
Stat Softw 2015; **67**.
78.   Fox J, Weisberg S. An R Companion to Applied Regression, Third Edition. 2018. Sage
Publications, Thousand Oaks, CA.
79.   Lenth R. emmeans: Estimated marginal means, aka least-squares means. 2023. R package
version 1.8.5.
80.   Kassambara A. rstatix: Pipe-friendly framework for basic statistical test. 2023. R package
version 0.7.2.
81.   Signorell A. DescTools: Tools for descriptive statistics. 2023. R package version 0.99.48.
82.   Toth AL, Robinson GE. Worker nutrition and division of labour in honeybees. *Anim Behav*
2005; **69**: 427–435.
83.   Toth AL, Kantarovich S, Meisel AF, Robinson GE. Nutritional status influences socially
regulated foraging ontogeny in honey bees. *Journal of Experimental Biology* 2005; **208**:
4641–4649.
84.   Kubo T, Sasaki M, Nakamura J, Sasagawa H, Ohashi K, Takeuchi H, et al. Change in the
Expression of Hypopharyngeal-Gland Proteins of the Worker Honeybees (Apis
melliferaL.) with Age and/or Role. *J Biochem* 1996; **119**: 291–295.
85.   Ueno T, Takeuchi H, Kawasaki K, Kubo T. Changes in the Gene Expression Profiles of the
Hypopharyngeal Gland of Worker Honeybees in Association with Worker Behavior and
Hormonal Factors. *PLoS One* 2015; **10**: e0130206.
86.   Sigg D, Thompson CM, Mercer AR. Activity-Dependent Changes to the Brain and Behavior
of the Honey Bee, *Apis mellifera* (L.). *The Journal of Neuroscience* 1997; **17**: 7148–7156.
87.   Fahrbach SE, Robinson GE. Juvenile Hormone, Behavioral Maturation, and Brain
Structure in the Honey Bee. *Dev Neurosci* 1996; **18**: 102–114.

88.   Withers GS, Fahrbach SE, Robinson GE. Selective neuroanatomical plasticity and division
765   of labour in the honeybee. *Nature* 1993; **364**: 238–240.
766   89.   Finkelstein AB, Brent CS, Giurfa M, Amdam G V. Foraging Experiences Durably Modulate
767   Honey Bees' Sucrose Responsiveness and Antennal Lobe Biogenic Amine Levels. *Sci Rep*
768   2019; **9**: 5393.
769   90.   Schulz DJ, Robinson GE. Biogenic amines and division of labor in honey bee colonies:
770   behaviorally related changes in the antennal lobes and age-related changes in the
771   mushroom bodies. *J Comp Physiol A* 1999; **184**: 481–488.
772   91.   Greenberg JK, Xia J, Zhou X, Thatcher SR, Gu X, Ament SA, et al. Behavioral plasticity in
773   honey bees is associated with differences in brain microRNA transcriptome. *Genes Brain*
774   *Behav* 2012; **11**: 660–670.
775   92.   Sinha S, Jones BM, Traniello IM, Bukhari SA, Halfon MS, Hofmann HA, et al. Behavior-
776   related gene regulatory networks: A new level of organization in the brain. *Proceedings*
777   *of the National Academy of Sciences* 2020; **117**: 23270–23279.
778   93.   Chandrasekaran S, Ament SA, Eddy JA, Rodriguez-Zas SL, Schatz BR, Price ND, et al.
779   Behavior-specific changes in transcriptional modules lead to distinct and predictable
780   neurogenomic states. *Proceedings of the National Academy of Sciences* 2011; **108**:
781   18020–18025.
782   94.   Ament SA, Wang Y, Chen C-C, Blatti CA, Hong F, Liang ZS, et al. The Transcription Factor
783   Ultraspiracle Influences Honey Bee Social Behavior and Behavior-Related Gene
784   Expression. *PLoS Genet* 2012; **8**: e1002596.
785   95.   Lutz CC, Rodriguez-Zas SL, Fahrbach SE, Robinson GE. Transcriptional response to
786   foraging experience in the honey bee mushroom bodies. *Dev Neurobiol* 2012; **72**: 153–
787   166.
788   96.   Lutz CC, Robinson GE. Activity-dependent gene expression in honey bee mushroom
789   bodies in response to orientation flight. *Journal of Experimental Biology* 2013; **216**:
790   2031–2038.
791   97.   Ament SA, Blatti CA, Alaux C, Wheeler MM, Toth AL, Le Conte Y, et al. New meta-analysis
792   tools reveal common transcriptional regulatory basis for multiple determinants of
793   behavior. *Proceedings of the National Academy of Sciences* 2012; **109**: E1801–E1810.
794   98.   Khamis AM, Hamilton AR, Medvedeva YA, Alam T, Alam I, Essack M, et al. Insights into
795   the Transcriptional Architecture of Behavioral Plasticity in the Honey Bee Apis mellifera.
796   *Sci Rep* 2015; **5**: 11136.
797   99.   Brodschneider R, Crailsheim K. Nutrition and health in honey bees. *Apidologie* 2010; **41**:
798   278–294.
799   100.   Wild B, Dormagen DM, Zachariae A, Smith ML, Traynor KS, Brockmann D, et al. Social
800   networks predict the life and death of honey bees. *Nat Commun* 2021; **12**.
801
802

 Figures

Figure 1. Nurses and foragers from typical honey bee colonies differ in gut microbial

community. (A-B) Nurses and foragers differed in overall gut microbial community structure. (A)

Reanalyzed data from Kapheim et al 2015. Two-way Permutation MANOVA using Aitchison

Distance, Task: $F_{1,38}$ = 1.51, R^2 = 0.036, p = 0.018; Colony: $F_{4,38}$ = 1.72, R^2 = 0.165, p = 0.001,

Task*Colony: $F_{2,38}$ = 1.17, R^2 = 0.056, p = 0.105. n = 5 colonies. **(B)** New data. Two-way

Permutation MANOVA using Aitchison Distance, Task: $F_{1,59}$ = 4.52, R^2 = 0.069, p = 0.001; Colony:

$F_{2,59}$ = 2.35, R^2 = 0.072, p = 0.001, Task*Colony: $F_{2,59}$ = 1.22, R^2 = 0.037, p = 0.114. n = 10

bees/colony, 3 colonies. Depicted as Principal Components Analysis (PCA) plots. Lowercase

letters in legends denote statistically significant groups. **(C)** Nurses and foragers differed in

relative abundance of four individual microbial species (new data only). Depicted as stacked bar

plots, with each bar representing a single bee's microbiome. Asterisks in legend: *, $p \leq 0.05$, **,

$p \leq 0.01$, ANCOM-BC between nurses and foragers. See Supplementary Table 1 for all p-values.

(D) Nurses and foragers differed in absolute abundance of four individual microbial species (new

data only). Depicted as dot plots with all data points plotted, line represents median, n = 10

bees/colony, 3 colonies. *, $p \leq 0.05$, **, $p \leq 0.01$, Permutation ANOVA Test between nurses and

foragers. See Supplementary Table 1 for all p-values.

Figure 2. Age-matched nurses and foragers differ in gut microbial community. (A-B) Age-matched typical-age nurses and precocious foragers from a single cohort colony did not differ in gut microbial community structure at about one week of age (A), but age-matched over-age nurses and typical-age foragers significantly differed in gut microbial community structure at three weeks of age (B). 1 week: Two-way Permutation MANOVA using Aitchison Distance, Task: $F_{1,39}$ = 0.93, R^2 = 0.023, p = 0.523; Source colony: $F_{3,39}$ = 2.66, R^2 = 0.065, p = 0.001, Task*Colony: $F_{1,39}$ = 1.17, R^2 = 0.028, p = 0.259. n = 10 bees/source colony, 2 source colonies. 3 weeks: Two-way Permutation MANOVA using Aitchison Distance, Task: $F_{1,79}$ = 6.48, R^2 = 0.073, p = 0.001; Source colony: $F_{3,79}$ = 1.89, R^2 = 0.065, p = 0.001, Task*Colony: $F_{3,79}$ = 1.22, R^2 = 0.042, p = 0.094. n = 10 bees/source colony, 4 source colonies. Depicted as PCA plots. Lowercase letters in legends denote statistically significant groups. **(C-D)** Age-matched nurses and foragers from a single cohort colony did not differ in relative abundance of individual microbial species at one week of age (C), but did differ in relative abundance of five individual microbial species at three weeks of age (D). Depicted as stacked bar plots, with each bar representing a single bee's microbiome. Asterisks in legend: *, p≤0.05, **, p≤0.01, ANCOM-BC between nurses and foragers. See Supplementary Table 2 for all p-values. **(E-F)** Age-matched nurses and foragers from a single cohort colony did not differ in absolute abundance of individual microbial species at one week of age (E), but did differ in absolute abundance of six microbial species at three weeks of age (F). Depicted as dot plots with all data points plotted, line represents median, n = 10 bees/source colony, 2 source colonies (1 week) or 4 source colonies (3 weeks). *, p≤0.05, **, p≤0.01, Permutation ANOVA Test between nurses and foragers. See Supplementary Table 2 for all p-values.

Figure 3. Age-matched inactive and active foragers differ in gut microbial community structure, but not the abundance of individual microbes. (A) Age-matched nurses and inactive foragers were similar in gut microbial community structure, while age-matched active and

inactive foragers differed in gut microbial community structure. Two-way Permutation MANOVA

using Aitchison Distance, Task: $F_{2,86}$ = 1.82, R^2 = 0.038, p = 0.011; Colony: $F_{2,86}$ = 5.90, R^2 =

0.122, p = 0.001, Task*Colony: $F_{4,86}$ = 0.798, R^2 = 0.033, p = 0.896. n = 10 bees/colony, 3

colonies. Depicted as PCA plot. Lowercase letters in legends denote statistically significant

groups as determined by Pairwise Permutation MANOVA. **(B)** Age-matched inactive and active

foragers did not differ in the relative abundance of individual microbial species (B). Depicted as

stacked bar plots, with each bar representing a single bee's microbiome. Asterisks in legend: *,

p$\leq$0.05, **, p$\leq$0.01, ANCOM-BC between inactive and active foragers. See Supplementary Table

4 for all p-values. **(C)** Age-matched inactive and active foragers did not differ in absolute

abundance of individual microbial species. Depicted as dot plots with all data points plotted, line

represents median, n = 10 bees/colony, 3 colonies. *, p$\leq$0.05, **, p$\leq$0.01, Permutation ANOVA

Test between inactive and active foragers. See Supplementary Table 4 for all p-values.

Figure 4. Bees inoculated with *B. asteroides* do not differ from microbiome-depleted bees in behavioral maturation (age at onset of foraging), but do differ in foraging intensity. (A) *B. asteroides* inoculated bees did not differ from microbiome-depleted bees in age at onset of foraging. Cox Proportional Hazards, z = -0.935, p = 0.350. **(B)** *B. asteroides* inoculated foragers, as a group, performed the majority of foraging trips for the colony the first day after experimental colony construction. Two-way Mixed ANOVA, Treatment: $F_{1,6}$ = 5.86, p = 0.050, Day: $F_{2,12}$ = 0, p = 1, Treatment*Day: $F_{2,12}$ = 2.64, p = 0.112. See Supplementary Table 5 for Pairwise comparisons. **(C)** *B. asteroides* inoculated bees represented a similar number of foragers as microbiome-depleted bees on all three days after experimental colony construction. Two-way Mixed ANOVA, Treatment: $F_{1,6}$ = 1.02, p = 0.351, Day: $F_{1.07,6.42}$ = 14.01, p = 0.008, Treatment*Day: $F_{1.07,6.42}$ = 0.15, p = 0.726. See Supplementary Table 5 for Pairwise comparisons. **(D)** Individual *B. asteroides* inoculated foragers performed a majority of foraging trips for the colony the first day after experimental colony construction. Generalized Linear Mixed Effects Model, Treatment: χ^2 =

3.44, p = 0.064, Day: χ^2 = 92.39, p < 0.001, Treatment*Day: χ^2 = 8.03, p = 0.018. See

Supplementary Table 5 for Pairwise comparisons. **(E)** *B. asteroides* inoculated bees represented

a higher proportion of elite foragers than microbiome-depleted bees on the first two days after

experimental colony construction. Two-way Mixed ANOVA, Treatment: $F_{1,6}$ = 9.597, p = 0.021,

Day: $F_{2,12}$ = 0, p = 1, Treatment*Day: $F_{2,12}$ = 1.78, p = 0.210. See Supplementary Table 5 for

Pairwise comparisons. (A) depicted as survival plot. All other data depicted as box plots with data

points plotted, line represents median, x represents mean, and whiskers represent the minimum

and maximum values, n = 4 colonies. Asterisks used to denote comparisons between treatment

groups on each day only: *, p≤0.05, **, p≤0.001.

Figure 5. Bees inoculated with *L. mellis* do not differ from microbiome-depleted bees in behavioral maturation (age at onset of foraging), but do differ in foraging intensity. (A) *L. mellis* inoculated bees did not differ from microbiome-depleted bees in age at onset of foraging. Cox Proportional Hazards, $z = 0.118$, $p = 0.906$. **(B)** *L. mellis* inoculated foragers, as a group, performed the minority of foraging trips for the colony the third day after experimental colony construction. Two-way Mixed ANOVA, Treatment: $F_{1,4} = 1.17$, $p = 0.340$, Day: $F_{2,8} = 0$, $p = 1$, Treatment*Day: $F_{2,8} = 0.30$, $p = 0.750$. See Supplementary Table 5 for Pairwise comparisons. **(C)** *L. mellis* inoculated bees represented a similar number of foragers as microbiome-depleted bees on all three days after experimental colony construction. Two-way Mixed ANOVA, Treatment: $F_{1,6} = 0.01$, $p = 0.946$, Day: $F_{2,12} = 24.20$, $p < 0.001$, Treatment*Day: $F_{2,12} = 0.01$, $p = 0.987$. See Supplementary Table 5 for Pairwise comparisons. **(D)** Individual *L. mellis* inoculated foragers performed a minority of foraging trips for the colony on the first day after experimental colony construction. Generalized Linear Mixed Effects Model, Treatment: $\chi^2 = 4.56$, $p = 0.032$, Day: $\chi^2 =$

126.64, p < 0.001, Treatment*Day: χ^2 = 1.926, p = 0.382, See Supplementary Table 5 for Pairwise

comparisons. **(E)** *L. mellis* inoculated bees represented a similar proportion, though with a trend

of lower proportion, of elite foragers as microbiome-depleted bees all three days after

experimental colony construction. Two-way Mixed ANOVA, Treatment: $F_{1,4}$ = 4.18 p = 0.111, Day:

$F_{2,8}$ = 0, p = 1, Treatment*Day: $F_{2,8}$ = 0.813, p = 0.477. See Supplementary Table 5 for Pairwise

comparisons. (A) depicted as survival plot. All other data depicted as box plots with data points

plotted, line represents median, x represents mean, and whiskers represent the minimum and

maximum values, n = 4 colonies. Asterisks used to denote comparisons between treatment

groups on each day only: *, p≤0.05, **, p≤0.001.

Figure 6. Bees inoculated with *L. melliventris* do not differ from microbiome-depleted bees in behavioral maturation (age at onset of foraging), but do differ in foraging intensity. (A) *L. melliventris* inoculated bees did not differ from microbiome-depleted bees in age at onset of foraging. Cox Proportional Hazards, z = -0.383, p = 0.702. **(B)** *L. melliventris* inoculated foragers, as a group, performed the minority of foraging trips for the colony the second day after experimental colony construction. Two-way Mixed ANOVA, Treatment: $F_{1,4}$ = 5.76, p = 0.074, Day: $F_{2,8}$ = 0, p = 1, Treatment*Day: $F_{2,8}$ = 3.09, p = 0.101. See Supplementary Table 5 for Pairwise comparisons. **(C)** *L. melliventris* inoculated bees represented a similar number of foragers as microbiome-depleted bees on all three days after experimental colony construction. Two-way Mixed ANOVA, Treatment: $F_{1,6}$ = 0.163, p = 0.700, Day: $F_{1.13,6.76}$ = 22.43, p = 0.002, Treatment*Day: $F_{1.13,6.76}$ = 0.04, p = 0.869. See Supplementary Table 5 for Pairwise comparisons. **(D)** Individual *L. melliventris* inoculated foragers performed a minority of foraging trips for the colony the second day after experimental colony construction. Generalized Linear Mixed Effects

Model, Treatment: χ^2 = 1.14, p = 0.286, Day: χ^2 = 86.85, p < 0.001, Treatment*Day: χ^2 = 5.79, p

= 0.055, See Supplementary Table 5 for Pairwise comparisons. **(E)** *L. melliventris* inoculated bees

represented a lower proportion of elite foragers than microbiome-depleted bees on the second

day after experimental colony construction, but a higher proportion on the third day after

experimental colony construction. Two-way Mixed ANOVA, Treatment: $F_{1,4}$ = 4.65, p = 0.097,

Day: $F_{2,8}$ = 0, p = 1, Treatment*Day: $F_{2,8}$ = 3.71, p = 0.072. See Supplementary Table 5 for

Pairwise comparisons. (A) depicted as survival plot. All other data depicted as box plots with data

points plotted, line represents median, x represents mean, and whiskers represent the minimum

and maximum values, n = 4 colonies. Asterisks used to denote comparisons between treatment

groups on each day only: *, p$\leq$0.05, **, p$\leq$0.001.

Microbe	Relative abundance			Absolute abundance		
	Typical colonies	SCC Replicate 1 (week 3)	SCC Replicate 2 (week 3)	Typical colonies	SCC Replicate 1 (week 3)	SCC Replicate 2 (week 3)
B. apis	lower abundance	lower abundance		lower abundance		
B. asteroides		lower abundance	lower abundance			
B. coryneforme					higher abundance	
Bom. apis						
F. perrara						
G. apicola					higher abundance	higher abundance
L. apis						higher abundance
L. helsingborgensis	higher abundance					
L. kullabergensis				lower abundance	higher abundance	higher abundance
L. mellifer		lower abundance				
L. mellis	higher abundance				higher abundance	higher abundance
L. melliventris	higher abundance	higher abundance		higher abundance	higher abundance	
S. alvi		lower abundance	lower abundance	lower abundance	lower abundance	

Table 1: Summary table of individual microbe abundance data. Microbes that significantly differed in relative (left) and absolute (right) abundance measures in nurses relative to foragers in typical colonies and individual single cohort colony replicates. Higher abundance, higher in abundance in nurses relative to foragers (pink). Lower abundance, lower in abundance in nurses relative to foragers (blue).

Test Microbe	Colony Replicate	Overall Colony Gini Coefficient	Day 1 Gini Coefficient	Day 2 Gini Coefficient	Day 3 Gini Coefficient
B. asteroides	1	0.376	0.056	0.116	0.425
B. asteroides	2	0.362	0.352	0.320	0.278
B. asteroides	3	0.415	0.133	0.445	0.368
B. asteroides	4	0.305	0.199	0.303	0.242
L. mellis	1	0.339	0.130	0.320	0.304
L. mellis	2	0.459	0.304	0.453	0.355
L. mellis	3	0.310	NA	NA	0.310
L. mellis	4	0.302	0.126	0.230	0.363
L. melliventris	1	0.398	0.186	0.271	0.365
L. melliventris	2	0.283	NA	NA	0.281
L. melliventris	3	0.419	0.167	0.386	0.415
L. melliventris	4	0.391	0.174	0.453	0.267

Table 2: Gini coefficients for each experimental colony depicted in Figures 4-6. Gini

coefficients, representing the degree of skew in foraging intensity between individual foraging

bees, where 0 represents complete equality in foraging intensity between individuals and 1

represents complete inequality in foraging intensity between individuals, measured for each

experimental colony across all days ("Overall Colony Gini Coefficient"), and for each experimental

colony on each individual day ("Day X Gini Coefficient"). NAs represent days with bad weather in

which foraging did not occur.