Imaging of existing and newly translated proteins elucidates the mechanisms of sarcomere turnover

Guy Douvdevany [1,2], Itai Erlich[1,2], Lilac Haimovich-Caspi[2], Maksymilian Prondzynski[3,4,5], Maria Rosaria Pricolo[6], Jorge Alegre-Cebollada[6], Wolfgang A. Linke[7], Lucie Carrier[3,4], Izhak Kehat*[2]

[1]These authors contributed equally

[2]The Rappaport Institute and the Bruce Rappaport Faculty of Medicine, Technion – Israel Institute of Technology, 1 Efron Street, P.O. Box 9697 Haifa 3109601, Israel.

[3]Institute of Experimental Pharmacology and Toxicology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

[4]DZHK (German Centre for Cardiovascular Research), partner site Hamburg/Kiel/Lübeck, Hamburg, Germany;

[5]Current address: department of Cardiology, Boston Children's Hospital, Harvard Medical School, Boston, MA 02115, USA

[6]Centro Nacional de Investigaciones Cardiovasculares (CNIC), Madrid, Spain

[7]Institute of Physiology II, University of Munster, Germany

*Correspondence: ikehat@technion.ac.il

Keywords:

Sarcomere

Protein turnover

HaloTag

Protein degradation

Abstract

How the sarcomeric complex is continuously turned-over in long-living cardiomyocytes is unclear. Here, we imaged the exchange of old and new Halo-tagged sarcomeric proteins. We disprove the prevailing 'protein pool' model and instead show an ordered mechanism in which only newly translated proteins enter the sarcomeric complex while older ones are removed and degraded. We also show that degradation is independent of protein age, and that proteolytic extraction is a rate limiting step in the turnover. We imaged and quantified the turnover of expressed and endogenous sarcomeric proteins, including the giant protein titin, in cardiomyocytes in culture and in vivo, at the single cell and at the single sarcomere level. We show that replacement occurs at a similar rate within cells and across the heart and is markedly slower in adult cells. Our findings imply a new model of sarcomere turnover and sarcomeric subunit replacement.

Introduction

Almost all biological processes depend on the assembly of proteins into functional complexes. The protein subunits of these functional complexes must be continuously replaced since mammalian cells often live much longer than their proteins[1,2]. Two models for the assembly of complexes were previously proposed. In the first model the complex is in a dynamic equilibrium with a pool of cytoplasmic protein and continuously recycles and exchanges subunits with the pool through de- and re- assembly[3]. This exchange was shown to maintain simple complexes such as actin fibers and microtubules[4,5]. An alternative model proposes a sequential assembly[6], where the complex is formed following an ordered series of steps. For example, the 26S proteasome shows a sequential assembly[7]. However, the mechanism by which such complexes are continuously maintained once they are fully assembled, and the role of subunit recycling in their maintenance remain largely unknown[8].

The sarcomere is the basic contractile protein complex of striated muscle, including cardiomyocytes. It is composed of the contractile myofilament proteins myosin and actin linked by titin (TTN), along with several other regulatory, structural, and accessory proteins[9]. Cardiomyocytes must continuously maintain their sarcomeres since sarcomeric proteins have a short half-life[10], and the decreased sarcomere and sarcomeric protein content in failing hearts suggest that failure to do so may lead to heart failure.[11][12] According to the prevailing model of sarcomere maintenance, sarcomeres are maintained by cytoplasmic soluble protein pools with

free recycling between pools and sarcomeres[13–15]. This model was more recently referred to as the "sarcostat"[14,16–18]. Support for this model came from fluorescence recovery after photobleaching (FRAP) experiments with fluorescently tagged sarcomeric proteins, that show recovery of fluorescent- actin[19,20], myotilin[21,22], α-actinin 2 (ACTN2), telethonin, tropomyosin 1 (TPM1)[23], and TTN[24] after bleaching within several hours. In the latter study it was even proposed that TTN molecules can move between adjacent sarcomeric complexes.[24]

Here we developed a pulse-chase approach based on covalent binding of different color fluorescent HaloTag ligands to image the replacement of older sarcomeric proteins by newly translated ones in cardiomyocytes in culture and in vivo. We imaged and measured the turnover of overexpressed and endogenous sarcomeric proteins, including the largest known protein TTN within the sarcomeric complex, in cultured cells and in vivo, at the single cell and the single sarcomere level. In contrast to the current 'protein pool' model of sarcomere maintenance, our findings provide evidence for an ordered process in which newly translated proteins are continuously added to the complex, while assembled proteins are continuously removed, degraded, and not recycled. We further show that once assembled, newly synthesized proteins are just as likely to be removed and degraded as older ones, and that proteolytic cleavage is needed for their extraction from the sarcomere. Our study thus establishes the 'unidirectional replacement' model for cardiac sarcomeres.

Results

Sarcomere maintenance is a unidirectional process

To image protein turnover in single living cardiomyocytes, we initially expressed Halo-tagged sarcomeric proteins in cultured neonatal rat ventricular cardiomyocytes (NRVMs) using adenoviral vectors. For the Z-line ACTN2 and the A-band myosin regulatory light chain 2 (MYL2) sarcomeric proteins we used full-length sequence (ACTN2-Halo and Halo-MYL2), and for the large M-line myomesin-1 (MYOM1) and the A-band cardiac myosin-binding protein C (MYBPC3) we used the N- and C-terminal fragments, respectively (MYOM1-Halo and Halo-MYBPC3). These fragments are small enough to be packaged in adenoviral vectors and sufficient for stable integration into sarcomeres[25,26]. After 48 hours, to allow expression of the Halo-tagged proteins, we pulsed the cells with the Oregon Green (OG) HaloTag-ligand. HaloTag is a modified haloalkane dehalogenase that covalently and irreversibly binds to HaloTag-ligand[27]. Therefore, once bound by the ligand the existing Halo-tagged proteins in the cell remain permanently

tagged with fluorescent green. To ensure that all the Halo-tagged proteins in the cell are bound by a ligand, we followed by pulsing with the non-fluorescent blocker ligand 7-bromoheptanol (7BRO) at a high concentration (65µM)[28]. We then extensively washed the cells from excess ligands, photobleached a small area in the sarcomere, added the far-red HaloTag ligand Janelia Fluor 635 (JF-635)[29], and imaged the cardiomyocytes for at least 72 hours (Fig. 1a). We limited the frequency of imaging to 12-24 hours to reduce phototoxicity. The JF-635 ligand shows almost no light absorption in aqueous solution, but upon binding to HaloTag, it increases its absorbance and fluorescence by more than 100-fold, and therefore can be used for continuous live cell imaging[30]. As expected, we did not observe far-red fluorescence signal immediately after the addition of the JF-635 ligand, confirming that all the Halo-tagged proteins in the cell were already bound, and that unbound JF-635 does not significantly fluoresce. However, once new Halo-tagged sarcomeric proteins were translated their available HaloTag could bind the JF-635 ligand and display far-red fluorescence. Our approach therefore allowed us to follow two main populations of sarcomeric proteins in the myocytes: older proteins labeled in green and newly translated sarcomeric proteins labeled in far-red. Over time, we observed an increase in the sarcomeric far-red fluorescence signal across the myocyte's sarcomeres along with a simultaneous decrease in green fluorescence, showing that newly synthesized proteins were added to the sarcomere and gradually replaced the older ones (Fig. 1b). We did not observe old or new Halo-tagged proteins outside the sarcomere, indicating that un-incorporated sarcomeric proteins are degraded.

We wondered whether subunit recycling is part of the maintenance process, and therefore we photobleached a small part of the green-labeled sarcomere at the beginning of the time lapse (Fig. 1a,b). According to the 'protein pool' model, sarcomeres would be maintained by a pool of cytoplasmic proteins containing a mixture of both older recycled and newly translated sarcomeric proteins. As a result, the photobleached sarcomeric proteins should be gradually replaced by both 'old' green labeled proteins and 'new' far-red labeled proteins from the pool (Fig. S1). Contrary to this prediction, we found that the photobleached area gradually filled with 'new' far-red labeled proteins, whereas the green signal failed to recover, showing sarcomeric proteins are not recycled (Fig. 1b). We initially used strong laser power to completely bleach the green signal. The green signal did not recover under these conditions, but the emergence of red florescence was somewhat retarded inside the bleached area, indicative of some local photodamage. We therefore opted for a lower level of bleaching. There was still no recovery of the green signal under these circumstances, and uniform emergence of red fluorescence was observed. To test whether we could identify diffusion we also expressed ACTN2-Halo and

performed FRAP in rat fibroblasts. Green-labeled ACTN2-Halo fluorescence recovered in fibroblasts within a minute of bleaching (Fig. S2), in contrast to cardiomyocytes where no recovery occurred.

Figure 1. Imaging of sarcomere turnover shows a unidirectional process

a, Pulse-chase experiment timeline: Cardiomyocytes expressing Halo-tagged sarcomeric proteins were incubated with Oregon Green ligand, followed by a non-fluorescent blocker ligand 7-bromoheptanol (7BRO), washed, and incubated with far-red Janelia Fluor-635 HaloTag ligand (JF-635). A small square part of the cell was photobleached, followed by at least 72 hours of time-lapsed imaging. **b,** Representative time-lapse images of cardiomyocytes expressing Halo-tagged ACTN2, MYOM1, MYBPC3 and MYL2 in the green (top), far-red (middle), and the magnified merged channel (bottom), before the photobleaching, immediately after (0), and at 24,48, and 72 hours, showing gradual replacement of existing proteins (green) with newly synthesized ones (far-red). The dashed white square is the photobleached field. It is gradually filling with new far-red labeled proteins but not with green labeled ones. Scale bar = 30μm. **c,** Normalized sarcomeric fluorescence amplitude is shown for the green and far-red channels with fitted exponential decay and accumulation curves (dotted lines) and calculated half-lives (bold dot and number; n=11-26 cells for each protein from N=2-3 independent experiments). **d,** Normalized sarcomeric fluorescence intensity is shown in the far-red channel for unbleached (light red) and bleached (dark red) myofibrils, showing similar rates of accumulation. **e,** Normalized sarcomeric fluorescence intensity is shown in the green channel showing exponential decay in the unbleached myofibril (light green) and no recovery in the bleached myofibril (dark green), except for some recovery in MYL2.

To quantify our findings and measure the intensity of the green and far-red signals, we stacked the time-lapse images, drew a linear region of interest along the myofibril and calculated the median amplitude of the signal intensity over time in these two channels along the line (Fig. S3). In addition to the sarcomeric signal we also observed a nuclear accumulation of signal for MYOM1, which likely represents an aggregate, and were therefore careful not to include the nucleus in our analyses. Assuming a first order kinetic, we fitted an exponential decay function to the degradation, or a function that gives the difference between 1 and the exponential decay to fit to synthesis and calculated the sarcomeric half-lives of proteins from the fitted curves (Fig. 1c). These data show that the sarcomeric half-lives calculated for ACTN2, MYOM1, MYBPC3, and MYL2 were similar. Our approach does not require the assumption of a steady state for quantification, and we could calculate the half-life from either the degradation rate of green labeled proteins or from the accumulation of far-red tagged proteins. While these calculations yielded values in the same range, the half-lives calculated from accumulation were longer than those calculated from degradation. Some correction factor is likely to be required, and bleaching from the timelapse imaging could have contributed to accelerating green fluorescence decline

and delaying far-red fluorescence accumulation. Together these analyses show that proteins are turned over in the sarcomere with a half-life of 20-42 hours in cultured cardiomyocytes.

We measured and compared the increase in incorporation of newly translated far-red labeled proteins outside and inside the bleached area and found similar rates, showing that sarcomeric proteins in different myofibrils are replaced simultaneously and at similar rates, and that our photobleaching did not damage the sarcomere (Fig. 1d). Similarly, we measured and compared the changes in the existing, green-labeled protein signal outside and inside the bleached area (Fig. 1e). All the protein studied showed exponential decline in intensity outside the bleached area and for ACTN2, MYOM1 and MYBPC3 we did not observe any recovery of the green signal in the bleached area. The exception was a small resurgence of green fluorescence immediately after photobleaching for MYL2 followed by an exponential decay (Fig. 1e), which may represent a small mobile unincorporated pool of MYL2. However, even if such a MYL2 pool existed it would be relatively small. Clearly, most of the existing MYL2 proteins in the sarcomere were replaced by newly synthesized ones and not through recycling. We could also quantify the turnover of single sarcomeres using super-resolution. To this end we tracked several single sarcomeres in individual cells and measured their signal amplitude in the stacked time lapse images. We fitted an exponential decay function to the measured data and calculated the half-life for each sarcomere individually. Although some variability existed, the half-lives of individual sarcomeres within the same cell were similar (Fig S4).

Collectively, our data refute the prevailing 'protein pool' hypothesis and prove that sarcomere maintenance is unidirectional: new proteins are incorporated into the sarcomere, but older proteins are extruded and degraded and cannot be recycled. Our measurements of the half-lives also imply that sarcomeric proteins are replaced with similar rates throughout the myocyte sarcomeres.

Imaging turnover of endogenous sarcomeric proteins

To confirm that our turn-over model also applies to endogenous cardiac proteins we used neonatal mice ventricular cardiomyocytes (NMVMs) derived from animals in which HaloTag was knocked-in to the I-band section of the largest known protein, the sarcomeric protein TTN[31]. We used the same protocol as used in NRVMs to image the turnover of endogenous TTN in mouse cardiomyocytes. Similar to NRVMs we used a minimal sufficient bleaching in our experiments because we found that harsh bleaching damaged the cells. As we showed for over-expressed sarcomeric proteins, the pre-existing TTN proteins were gradually and uniformly replaced by

newly translated ones across the cardiomyocyte (Fig. 2a). After photobleaching there was no reemergence of the green fluorescence showing that endogenous TTN cannot be recycled (Fig. 2a). Quantification of our data confirmed this observation and showed that TTN is turned over with a half-life of about 15-31 hrs. (Fig. 2b-d).

Next, we used an approach that allowed us to knock in the HaloTag to endogenous genes in cultured cardiomyocytes. The CRISPR/Cas9-based homology-independent targeted integration (HITI) strategy can generate precise and efficient knock-in in genes of non-dividing cells like cardiomyocytes by using non-homologous end-joining (NHEJ), with no insertions or deletions at most junction sites[32]. To knock in the HaloTag in the endogenous *Actn2* locus in cardiomyocytes using HITI we identified a unique Cas9 target site upstream of the C-terminal stop codon of the *Actn2* gene. We used the same Cas9 target site to flank the donor integration cassette, which contained the *Actn2* C-terminus without a stop codon, followed in-frame by the HaloTag, allowing the same single guide RNA (sgRNA) to guide Cas9 to cut both in the genomic and on both sides of the cassette (Fig. S5a). Since integration of the cassette would regenerate the sgRNA site and allow Cas9 to cut the cassette out again, the Cas9 cut sites flanking the cassette were placed inversely to the sgRNA orientation in the genome, creating a hybrid junction that the sgRNA can no longer recognize after integration. We isolated NMVMs from Cas9-expressing mice[33], transduced them with adeno associated viruses (AAVs) encoding for the donor cassette and for either the *Actn2* C-terminal sgRNA or for control non-targeting sgRNA. Strong Z-line staining was observed after the HaloTag ligand addition in ~15% of cardiomyocytes only when *Actn2* sgRNA was used, and not when control non-targeting sgRNA was used, indicating that HaloTag cassette was integrated properly in frame in these cardiomyocytes (Fig S5b-c). We executed the same imaging protocol we used for TTN and for expressed Halo tagged proteins and followed the turnover of endogenous ACTN2 with knocked-in HaloTag. The turnover of endogenous ACTN2 in NMVM was similar to what we observed with virally expressed ACTN2-Halo in rat cardiomyocytes: pre-existing ACTN2 proteins were gradually and uniformly replaced by newly translated ones across the cardiomyocyte, and after photobleaching there was no reemergence of the green fluorescence, showing that endogenous ACTN2 cannot be recycled (Fig. 2e). Quantification of the data showed that endogenous ACTN2 is turned over with a half-life of about 19-31 hrs. (Fig. 2f-h). Together our data on ACTN2 and TTN demonstrated that both overexpressed and endogenous proteins cannot be recycled.

We also transduced NMVM with adenoviral vectors encoding for ACTN2-Halo and compared the turnover rates of exogenously over-expressed ACTN2-Halo to endogenous HITI generated

knock-in ACTN2-Halo (Fig 2i-j). We found that these turnover rates were similar, implying that expression level does not determine the turnover of sarcomeric proteins in the complex.

Figure 2. Imaging turnover of the endogenous sarcomeric proteins TTN and ACTN2.

a, Representative time-lapse images of neonatal mouse ventricular cardiomyocytes from Halo-tagged TTN knock-in mice in the green (top), far-red (middle), and in magnified merged channel (bottom), before the photobleaching, immediately after (0), and at 24,48, and 72 hours, showing gradual replacement of existing TTN proteins (green) with newly translated ones (far-red). The dashed white square is the photobleached field. It is gradually filling with new far-red labeled proteins but not with green labeled ones. Scale bar = 30µm. **b**, Normalized sarcomeric fluorescence amplitude is shown for the green and far-red channels with fitted exponential decay and accumulation curves (dotted lines) and calculated half-lives (bold dot and number). n=41 cells from N=3 independent experiments. **c**, Normalized sarcomeric fluorescence intensity is shown in the far-red channel for unbleached (light red) and bleached (dark red) myofibrils, showing similar degree of accumulation. **d**, Normalized sarcomeric fluorescence intensity is shown in the green channel showing exponential decay in the unbleached myofibril (light green) and no recovery in the bleached myofibril (dark green). **e**, Representative time-lapse images of NMVM after HITI knock-in of HaloTag to the *Actn2* locus using the same protocol as in (a), showing gradual replacement of existing endogenous ACTN2 proteins (green) with newly translated ones (far-red). The photobleached field is gradually filling with new far-red labeled proteins but not with green labeled ones. **f-h**, Quantification and calculation of assembly and degradation half-lives for endogenous ACTN2 as in (b-d). n=34 cells from N=2 independent experiments. **i-j**, Comparison of turnover rates of endogenous knocked-in ACTN2-Halo to exogenous virally expressed ACTN2-Halo in NMVMs. Normalized sarcomeric fluorescence amplitude is shown for the green (i) and far-red (j) channels with fitted exponential decay and accumulation curves (dotted lines) and calculated half-lives (bold dot and number). The deep and light purple dots and lines in (i) show the degradation of endogenous and exogenous virally expressed ACTN2 from the green channel respectively. The deep and light cyan dots and lines in (j) show the accumulation of endogenous and exogenous virally expressed ACTN2 from the far-red channel. n=34 endogenous HITI cardiomyocytes and n=14 exogenous virally transduced cardiomyocytes, each from N=2 independent experiments.

Sarcomere protein degradation is independent of protein age and controls turnover rate

It has been hypothesized that old proteins are identified by quality control pathways and targeted for degradation[34]. We therefore wanted to determine whether young and old proteins in the sarcomere are turned over at the same rate. We used MYOM1-Halo that has a very strong signal and expressed it in NRVM using an adenoviral vector. After 48 hours, we incubated the cells with the OG ligand and then with the JF-635 ligand for an additional 48 hours. To block all HaloTags we pulsed with the non-fluorescent blocker 7BRO at a high concentration (65μM) for 15 minutes after each fluorescent ligand, followed by extensive washing. This labeling protocol allowed us to follow the degradation of two main populations of sarcomeric proteins in the myocytes: older proteins labeled green and newer proteins labeled far-red over the next 72 hours (Fig. 3a). As before, we were careful not to include the MYOM1 nuclear aggregate in our analyses. We observed a gradual decline in the sarcomeric signal intensity for both the old and the new proteins (Fig. 3b). We fitted the signal data to an exponential decay function and found similar rates of decay and a similar half-life for both the old and new MYOM1 proteins (Fig. 3c,d). These results support a stochastic model where all proteins have the same likelihood of being removed and degraded, regardless of how old they are.

Both calpain and proteasomal proteases are responsible for the degradation of sarcomeric proteins[34]. However, it is not known if the proteolytic activity of these enzymes is required for the removal of proteins from the sarcomeric complex, or whether sarcomeric proteins can detach intact from the complex and are only degraded later. To differentiate between these two hypotheses, we used the peptide aldehyde MG-132, which is capable of inhibiting both calpains and the proteasome[35]. We expressed ACTN2-Halo in NRVMs using adenoviral vector and repeated the imaging protocol we used to study the turnover and measured the half-life of ACTN2-Halo with or without incubation with MG-132. The inhibition of proteases by MG-132 resulted in a marked delay in the removal of existing green-labeled ACTN2-Halo proteins from the sarcomeric structure with a significant increase in ACTN2 sarcomeric half-life (Fig. 3e-g). We performed the same analysis with endogenous TTN in NMVMs of the Halo-TTN knock-in mice, and similarly observed a marked delay in the removal of existing TTN and a significant increase in TTN sarcomeric half-life (Fig. 3h-j). This delayed turnover indicates that proteolytic activity is required for the removal of sarcomeric proteins such as ACTN2 and TTN from the sarcomeric structure.

Figure 3. Degradation is independent of protein age and requires proteases.

a, Degradation experiment timeline - Cardiomyocytes expressing Halo-MYOM1 for 48 hours were incubated with the OG and then with the JF-635 Halo-ligands for an additional 48 hours and imaged in time-lapse 72 hours. **b,** Representative images of cardiomyocytes at 0 ,24, 48, and 72 hours show a gradual decline of the sarcomeric signal in both the green and far-red channels. Scale bar = 10.25μm. **c-d,** Quantification of normalized sarcomeric fluorescence intensity in the green (OG) and far-red (JF) channels (c)**,** showing the half-lives of newer and older proteins in individual cardiomyocytes did not differ (d) (n=15 cells, N=2 independent experiments, two tailed paired Student's t-test p=0.3). **e,** Representative time-lapse images of NRVMs expressing ACTN2-Halo incubated with (bottom) or without (top) MG-132 at 0, 24, 48, and 72 hours in the green channel show that inhibition of proteases by MG-132 resulted in a marked delay in the removal of existing OG labeled proteins from the sarcomere. Scale bar=10μm **f-g**, Quantification of normalized sarcomeric fluorescence intensity for control (light green, Ctl) and MG-132 treated cardiomyocytes (dark green) (f) show much longer half-lives of ACTN2-Halo in MG-132-treated than in control cardiomyocytes (g) (n=9-41 cells, N=3 independent experiments, two tailed Student's t-test p<0.001). **h-j** similar analyses as in (e-g) in NMVM of Halo-TTN knock-in mice show much longer half-lives of endogenous Halo-TTN in MG-132-treated cardiomyocytes relative to controls. Scale bar = 10μm (n=24-33 cells, N=2 independent experiments, two tailed Student's t-test p<0.0001). **k**, Representative images of cardiomyocytes expressing ACTN2-Halo treated with control or MG-132 after 72 hours in the merged green and far-red channel with inset (dashed square) showing accumulation of newly translated far-red labeled proteins outside the Z-line and multiple aggregates of newly translated (yellow dotted circle) proteins and some aggregates of old proteins (white dotted circle). Scale bar = 10μm. **l**, Representative images of cardiomyocytes with knock-in Halo-TTN treated with control or MG-132 after 72 hours in the merged green and far-red channels with inset (dashed square) showing aggregates composed of newly translated far-red labeled proteins (yellow dotted circle) that displace the myofibrils. Scale bar = 10μm.

Earlier, we posited that sarcomeric proteins are translated abundantly, accompanied by swift proteasomal degradation of any unincorporated proteins[1]. In agreement with this proposal, we found that newly translated ACTN2-Halo proteins accumulated outside the Z-line structure following proteolytic blockade. By 72 hours multiple aggregates composed predominantly of newly translated far-red- labeled proteins were observed (Fig. 3k). Interestingly we also detected several discreet large aggregates predominantly made of older green labeled ACTN2-Halo

proteins (Fig. 3k). These aggregates could contain proteins derived from the disintegrating sarcomeric structure observed at this stage. Unlike ACTN2-Halo which was over-expressed in these experiments, the Halo-TTN is an endogenous protein. Large aggregates composed of newly translated far-red labeled Halo-TTN were observed between the myofibrils after proteolytic blockade, but these accumulations were not as dense as those of ACTN2-Halo (Fig. 3l). Together these data show that when proteolysis is blocked the removal of sarcomeric proteins such as ACTN2 and TTN from the sarcomere is delayed and the half-life of these proteins in the structure is increased. Newly translated proteins cannot push the existing intact proteins out of the structure without proteolysis, and if newly translated proteins cannot be integrated or degraded, they accumulate outside the sarcomere. These findings suggest that proteolysis is the rate limiting step in sarcomere turnover.

Imaging protein turnover in vivo using viral vectors

To image and quantify the turnover of sarcomeric proteins in vivo in the adult mouse heart we transduced neonatal mice with AAV9 vectors encoding TPM1-Halo or ACTN2-Halo full length proteins. After waiting for the mice to reach the age of 4 months we injected the non-fluorescent blocker 7BRO intravenously and intraperitoneally to ensure saturation of all the available HaloTags. We then waited either 3 hours or 3 days and intravenously delivered the fluorescent Janelia Fluor-525 HaloTag ligand (JF-525). To measure the maximal possible labeling intensity, we also delivered Halo-ligand JF-525 to mice that did not receive the 7BRO blocker. In addition to the '3 hours' group we included another negative control group of mice that received the ligands but not the AAVs encoding for Halo-tagged proteins (Fig 4a). We harvested the hearts 40 minutes after the injection of the JF-525 Halo-ligand and sectioned. Low and high magnification images of cardiac sections showed background signal in the 'no AAV' and '3 hours' negative control groups (Fig. 4b-e); the sarcomeric fluorescence signal was increased by 3 days and was highest in the 'no 7BRO' cardiomyocytes. In the latter, it was evident that the transduction rates are significant, albeit with some cell-to-cell variability. As we did in the cultured cardiomyocytes, we drew a region of interest along the myofibril in individual cardiomyocytes in the sections (Fig. 4 d-g) and calculated the median amplitude of the sarcomeric signal (Fig. 4 h-i). We fitted the data to a 1 minus exponential decay function and estimated the half-life of ACTN2 and TPM1 to be 312 and 91 hours, respectively (Fig. 4 j-k). Although we could not follow the same cardiomyocytes over time in vivo as we could in culture experiments, we could estimate the half-life from the average of multiple cells. We found that the

measured half-life of ACTN2 in vivo was ~10-fold longer than in cultured neonatal cardiomyocytes, indicating a slowing of the turn-over as cardiomyocytes matured.

Figure 4. Imaging of sarcomeric protein turnover in vivo

a, Neonatal mice were divided into four groups: In the 'No AAV' group no AAV injection was administered, while in the other three groups mice received AAV9 encoding for either ACTN2-Halo or TPM1-Halo. In adulthood all groups received an intravenous injection of the JF-525 Halo-ligand, and their hearts were harvested for analysis after 40 minutes. Mice in the 'No AAV' and the 'No 7BRO' groups did not receive intravenous injection of the non-fluorescent blocker

Imaging endogenous TTN turnover in vivo

While AAVs are efficient at expressing Halo-tagged sarcomeric proteins, their limited cloning capacity and variability in transduction efficiency may restrict their use. We therefore again used the mice in which HaloTag was knocked-in the I-band section of the giant sarcomeric protein TTN[31] to study the turnover of TTN in vivo. We employed a similar labeling approach to the one we used in the AAV transduced mice with injection of the non-fluorescent blocker 7BRO, followed by an injection of JF-525. We used similar negative controls with wild type mice that do not have Halo-tagged TTN and Halo-tagged TTN mice in which the fluorescent ligand was injected 3 hours after injection of the blocker, to confirm complete blocking of existing Halo-tagged TTN. We also injected Halo-ligand JF-525 to Halo-tagged TTN mice that did not receive the 7BRO blocker to measure the maximal possible labeling intensity. Here we included more labeling time points, with injection of Halo-ligand JF-525 at 3, 7, and 14 days after the injection of the blocker (Fig. 5a). We also added Halo-ligand JF-525 to the cardiac sections to ensure complete labeling. Representative images of the sections and of cells at low and high magnification showed no sarcomeric signal with injection after 3 hours, indicating blockade was complete, and increasing sarcomeric signal at 3, 7, and 14 days (Fig. 5b-d). The group that did not receive the non-fluorescent blocker showed maximal signal intensity. We used the same analysis approach we used for the AAV transduced hearts. We drew a region of interest along the myofibril axis on individual cardiomyocytes (Fig. 5d) and used an automated tool to calculate

the median amplitude of the sarcomeric signal along the line (Fig. 5e). We fitted the data to a 1 minus exponential decay function and estimated the half-life of endogenous TTN to be 191 hours (Fig. 5f). This number is more than 6-fold longer than the turnover of TTN in cultured NMVMs, indicating a marked slowdown in TTN turnover after maturation.

We next aimed to analyze the Halo-tagged TTN mice after induction of cardiac hypertrophy. Administration of phenylephrine (PE) activates α1-adrenergic receptors and is known to stimulate cardiomyocyte protein synthesis in mice[36]. We used a PE injection protocol in the Halo-tagged TTN mice and verified that it indeed resulted in a significant increase in normalized heart weight, indicative of hypertrophy (Fig S6a). We asked whether hypertrophy and an increased protein synthesis would also result in increased turnover rate of TTN. We therefore included groups of mice in which we injected PE, in parallel with the Halo-ligand pulse chase protocol (Fig. 5b-f and Fig S6b). The average maximal labeling intensity observed in the group that did not receive the 7BRO blocker was higher in the PE-treated mice, but the difference was not statistically significant. Accordingly, the half-life calculated for the PE group was slightly longer than in the control group (225 vs 191 hours), but the difference was not statistically significant. These data suggest that the turnover of titin is not significantly accelerated in hypertrophy.

Finally, we analyzed the turnover rates in different regions of the heart. We chose the left ventricular free wall, the interventricular septum and the right ventricular free wall as representative regions and measured the half-life of TTN in each of them. Our analysis showed that TTN is turned over in different regions of the heart at similar rates (Fig. S7).

Figure 5. In-vivo turnover of endogenous TTN in the normal heart and in hypertrophy
a, Experiment design. Adult Halo-TTN knock-in mice received a non-fluorescent Halo-ligand (7BRO). After a chase period of 3 hours, 3 days, 1 week, or 2 weeks, mice received the JF-525 ligand, and the hearts were harvested. To measure the maximal labeling, one group of HaloTag-TTN knock-in mice received the fluorescent ligand without the 7BRO blocker. A negative control group of wild-type mice also received the fluorescent ligand without the 7BRO blocker. **b-d,** Representative images of cardiac sections from the different groups for control and phenylephrine (PE) treated mice at low (b, scale bar = 100μm), and high magnification (c, scale bar = 30μm). The yellow box in (c) indicates a region of interest of ~10 μm length used for quantification and is shown at high magnification in (d). **e,** Quantification of signals from (d). The local maxima and minima were identified automatically and marked in red and blue dots, respectively. **f,** Quantification of sarcomere labeling intensity from control mice (red) and mice that received PE (green). Each dot represents the mean sarcomere intensity of 10 cells from 10 different fields, acquired from a section of a single mouse heart. The error bars represent the standard error. The no blocker control group data is indicated in infinity. The horizontal dashed line is the mean intensity in the no blocker positive control group and was used as total protein reference in the model. The curved dark dashed line is the fitting curve of the data to 1 minus exponential decay model. The vertical solid lines indicate the half-life of the TTN protein calculated from the model. Time is indicated in hours, and intensity in arbitrary units (AU). (n = 10 cells/mouse, from n = 3-8 mice/group, from N=4 independent experiments, ANOVA between control and PE p=0.205)

Discussion

Here we developed a pulse-chase approach based on the ability of Halo-tagged proteins to covalently bind fluorescent ligands. This approach allowed us to visualize pre-existing and newly translated proteins, measure their assembly and degradation rates, and uncover the mechanisms of sarcomeric maintenance. We showed that different sarcomeric proteins are uniformly turned over at similar rates, that turnover is slower in vivo than in cultured neonatal cardiomyocytes, and that the turnover of the giant protein TTN is uniform throughout the heart. We challenged the prevailing 'protein pool' hypothesis for sarcomeric maintenance and found that assembly is unidirectional - newly translated proteins continuously and uniformly enter the structure, whereas already assembled proteins can only be removed and degraded. Further, we demonstrated that older and newer sarcomeric proteins are not distinguished in the degradation process, and that proteolytic extraction from the complex is a rate limiting step in the turnover.

Since expressed sarcomeric proteins stoichiometrically replace endogenous ones[37] we could use virally expressed Halo-tagged sarcomeric proteins, and we confirmed that this applied to endogenous proteins as well. We relied on the TTN Halo-tag knock-in mice to image the turnover of the giant protein TTN. Since such tagged mice were not available for other proteins,

we developed a simple HITI approach that can be used to knock-in HaloTag to endogenous proteins in cultured cardiomyocytes by expressing Cas9, the donor HaloTag cassette, and in-vitro transcribed sgRNAs. We used this simple approach to follow the replacement and turn-over of endogenous ACTN2, and we believe that this approach can be adapted to almost any protein.

Proteomic approaches were used to measure the half-lives of cardiac proteins in vivo[38]. Such analyses are powerful and allow the study of thousands of proteins in parallel. However, proteomics averages the half-life of proteins over the entire heart tissue and cannot differentiate between cardiomyocytes and non-cardiomyocytes. We did not measure the half-life of proteins per se, but rather calculated the turnover inside the sarcomeric structure in cardiomyocytes. Nevertheless, for the giant protein TTN the half-life of turnover measured by us, ~8 days, was similar to the average 11.8 days half-life calculated by proteomics. Likewise, the half-lives we measured for ACTN2 and TPM1 of 13, and 3.8 days were comparable to the proteomics calculated half-lives of 11.4 and 7.15 days respectively [38]. A hypertrophic stimulation did not alter the half-life of TTN, as was also observed with proteomics[38]. The main advantage of our approach, however, is that it allowed us to study and image the turnover dynamics of proteins in living cultured cardiomyocytes and in vivo at the single cell and even at the single sarcomere level. We used a first-order kinetics model to analyze the assembly and degradation of the protein independently, without assuming a steady state, and found that the half-life for assembly was longer than that calculated for the degradation. This difference could reflect a true decrease in the total content of the sarcomeric protein in the cardiomyocytes throughout the measurement time, but we believe that it is more likely a result of bleaching during time lapse imaging which accelerated the decrease in green fluorescence and delayed the accumulation of far-red fluorescence. Nevertheless, the relative comparisons between sarcomeres, cells, or conditions would not be affected, even if some corrections are needed to calculate the exact half-lives.

With our approach, we were able to understand the key principles underlying protein complex maintenance, specifically for sarcomeres. Although it has been widely reported that sarcomeric proteins could be recycled[13–15], we showed that the maintenance is unidirectional - new proteins can get into the complex, while older ones can only exit, and cannot be recycled. Previous studies of sarcomere turnover used FRAP with genetically encoded fluorophores, typically eGFP or YFP, and extended the analysis time to many hours to show recovery. An inherent assumption of FRAP is that the transition between bright and dark states is irreversible, and that fluorescence recovery is only caused by fluorescent molecules diffusing into the bleached area. It is also possible, however, for fluorescent molecules to convert into a reversible dark state, in a

process called photoswitching. Several fluorescent molecules, including GFP, were shown to exhibit photoswitching[39], and such reversible photoswitching may appear as mimicking diffusion. Indeed, one study that used fluorescently tagged TTN found extensive reversible photobleaching that accounted for most of the TTN fluorescence recovery[40]. Our experiments did not reveal any recovery of OG Halo-tagged sarcomeric proteins, even when we bleached at a submaximal level. This fluor is likely less prone to photoswitching and unlike genetically encoded fluorophores do not need to fold or mature. We performed experiments in fibroblasts, where expressed sarcomeric proteins can freely diffuse to show that recovery following FRAP can be detected in our imaging setting. We did observe some quick recovery for MYL2, suggesting there may be a small diffusible pool for this protein.

In contrast with assembly that is ordered, degradation appeared to be stochastic. Cardiomyocytes degrade old and new proteins at similar rates, and likely cannot distinguish between them. We further showed that pharmacologic blockade of both calpains and proteasomal proteases delays the removal of existing proteins from the sarcomeric structure, indicating that proteolytic cleavage of sarcomeric proteins is required for their extraction. The removal of existing proteins, however, was slowed, but not completely stopped. We assume that the incomplete response to MG-132 is due to incomplete protease blockade, as we were limited in the dose of MG-132 due to toxicity, but we cannot exclude that a certain portion of sarcomeric proteins can exit the structure without being cleaved first. These data, together with the observation that the turnover of over-expressed ACTN2 occurs at the same rate as that of the endogenous protein, implies that the proteolytic extraction from the sarcomere is a rate limiting step in the turnover of sarcomeric proteins. These findings further suggest that the degradation of sarcomeric proteins may be regulated.

There are multiple sarcomeric proteins, many with different isoforms. The turnover principles we identified were similar for all the sarcomeric proteins we studied. Due to the complexity of our studies, such as the need for long and stable imaging of 72 hours for each cell, we were unable to extend the study to additional sarcomeric proteins. As these proteins all belong to the same complex it is highly likely that they share the same turnover mechanisms; however, we cannot rule out the possibility that some exceptions exist.

In conclusion, we discovered mechanisms of sarcomere maintenance that could be useful for developing future treatments for genetic cardiomyopathies as well as for heart failure and hypertrophy. In particular, the requirements for enzymatic cleavage as a rate limiting step in the process of replacement may be exploited to increase or delay the turnover of proteins in the

sarcomeric structure. Finally, while we demonstrated how sarcomeres are maintained in cardiomyocytes, our findings likely apply also to other heteromeric complexes in other cells.

Methods

Tissue culture

Neonatal rat ventricular myocytes (NRVMs) were isolated using the Worthington Biochemical Neonatal Cardiomyocyte Isolation system (cat#LK003300), according to the manufacturer's instructions, followed by Percoll gradient centrifugation, as previously described[41].

Neonatal mouse ventricular cardiomyocytes (NMVMs) were isolated according to previously published protocol[42]. Briefly, hearts were harvested and washed in 1xPBS supplemented with 20 mM BDM. Hearts were then minced and incubated in isolation medium (20 mM BDM, 0.0125% trypsin in HBSS (without Ca2+, Mg2+) overnight at 4°C. The following day the predigested hearts were transferred to a digestion medium (20 mM BDM, Worthington collagenase (cat#LK003240) in L15 medium) and were incubated at 37°C for 25 min with gentle shaking. Tissue fragments were triturated 15 times, and the cell suspension was strained. Cells were centrifuged at 300xg for 5 min. The supernatant was removed, and cells were resuspended in 10 ml of plating medium (65% DMEM high glucose, 19% M-199, 15% fetal calf serum, 1% penicillin/streptomycin).

Adenovirus vector production and transduction

The segments containing sarcomeric proteins or truncated protein coding sequence were amplified from a library of genomic NRVM DNA by PCR and cloned into Gateway cloning system (Thermo Fisher) 'Entry' plasmid. The coding sequence was verified by Sanger sequencing and cloned into pAd/CMV/V5-DEST adenovirus expression plasmid (Thermo Fisher). HEK293 cells were transfected with the adenoviral plasmids as previously described[41], for the generation and propagation of adenoviral particles. Cultured cardiomyocytes were transduced one day after seeding by incubation with the adenovirus for 4 hours, followed by washing.

The constructs used:

- Homo sapiens full size Actinin alpha 2 (ACTN2) – 894 amino acids, with HaloTag in the C terminus. GenBank: KR712245.1

- Rattus norvegicus Myomesin 1 (Myom1) 507 last amino acids, with HaloTag in the C terminus. GenBank: NM_001191584.1

- Rattus norvegicus full size Myosin light chain 2 (Mlc2) – 166 amino acids, with HaloTag in the N terminus. GenBank: NM_001035252.2.

- Rattus norvegicus Myosin binding protein C3 (Mybpc3) 416 last amino acids with HaloTag in the N terminus. GenBank: NM_001106490.1.

Pulse-chase in cardiomyocyte cultures

NRVMs or NMVMs were cultured on μ-Dish 35-mm Quad (Ibidi, 80416) coated with Matrigel Matrix (Corning, 354234), in 15% serum media. Cells were transduced with adenoviral vectors encoding the Halo tagged sarcomeric proteins. After 48 hours, 1 μM Oregon Green HaloTag Ligand (Promega G2801) was added to the cells for 15 min. The cells were then incubated with 65μM of 7-bromoheptanol (7BRO, Sigma-Aldrich 310913) medium for additional 15 min, washed 3 times. Janelia Fluor 635 ligand, 0.2 μM, (a kind gift from Luke Lavis) was added to the cells.

The MG-132 experiments were performed with adenovirus mediated ACTN2-Halo transduced NRVMs, or with NMVM isolated from Halo-TTN knock-in mice. We added 5 μM MG-132 (Apexbio, A2585) or vehicle and imaged over 72 hours.

Protein degradation experiment

For degradation experiments NRVM were transduced with adenoviral vector encoding for MYOM1-Halo. After 48 hours cells were incubated with 1 μM Oregon Green HaloTag Ligand for 15 min followed by 65μM 7BRO for 15 min and three washes. Janelia Fluor 635 ligand, 0.2 μM, was added, and cells were incubated for 48 hours to allow synthesis of new proteins and binding of the red ligand. The cells were incubated again with 65μM 7BRO medium, washed 3 times, and imaged over 72 hours.

Homology-independent targeted integration

Neonatal cardiomyocytes primary cultures were prepared from Rosa26-Cas9 knockin mice on C57BL/6J background (Jackson laboratory Strain #:024858)[33]. Cardiomyocytes were transduced with two AAV9 vectors (1 X 10^11 viral genomes each), one encoding the *Actn2*-HaloTag donor cassette and another one encoding for either control non-targeting sgRNA (GATGTTCAAGATTCAGGAGG) or the *Actn2* c-terminus sgRNA (CAGTACTGGGCCTGATCCGG) under the control of the U6 promoter.

Live cell imaging

Cells were imaged using Zeiss LSM 900 with Airyscan2 super-resolution system, attached to Axio Observer 7 inverted microscope, with PECON incubation insert, allowing regulation of temperature, CO_2, and humidity. The acquisition was done with Plan-Apochromat 63x/1.4 Oil DIC M27 (#420782-9900-799) Zeiss objective, with solid state laser with wavelengths of 488 & 640 nm. The photobleach was performed with the 488 nm laser in a small square area at the center of the field.

Image analysis

Time series images were aligned using the SIFT-algorithm, in ImageJ software[43]. For each cell linear regions of interest (ROIs) of ~10 μm were marked, inside and outside the photobleached area, and were used for all the time-lapse images of that cell. In the degradation and in vivo experiments, in which no photobleach was preformed, only one ROI was marked and analyzed. The intensity values of the ROI were analyzed using R (R: A language and environment for statistical computing (R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/). We determined the local maxima and minima points along the ROI using a moving window of 20 pixels (~1 μM). The amplitudes along the ROI were calculated from the local maxima and minima, and the median of all amplitude values was used as the representative value for the myofibril or cell. To calculate the half-life the median amplitude values from each cell were normalized between 0 and 1 to standardize the values and allow analysis of all cells together. Nonlinear Least Squares (NLS) function was used to fit all the representative values according to equation 1 for protein degradation, in exponential decay model, and equation 2 for protein synthesis, in 1 minus exponential decay model[44].

$$Eq.1: \qquad intensity = K * e^{(-t*\lambda)}$$

$$Eq.2: \quad intensity = 1 - K * e^{(-t*\lambda)}$$

K and λ are parameters estimated by the NLS function, *t* is the time and *intensity* is the calculated representative value.

The half-life calculated for each fitting model was calculated according to equation 3.

$$Eq.3: \qquad t\frac{1}{2} = \frac{\ln(2)}{\lambda}$$

Adeno-associated virus vector production and injection

The AAVs were produced according to a published protocol[45], as we previously described[46]. For transduction 50µl of virus containing a total of ~3 X 10^11 viral genomes was delivered intraperitoneally to 4-7-days old mouse pups.

Mouse pups were injected with AAV viruses, encoding for the following constructs or control (no virus):

- Homo sapiens full size Actinin alpha 2 (ACTN2) – 894 amino acids, with HaloTag in the C terminus. GenBank: KR712245.1

- Rattus norvegicus full size Tropomyosin 1 (TPM1) - 285 amino acids, with HaloTag in the C terminus. GenBank: NM_001301336.1

Animal experiments

All pups were born at the SPF unit of the pre-clinical research authority at the Technion. Health monitoring was carried out in accordance with FELASA recommendations. Each litter was housed with the dam weened at 4 weeks and separated by sex. Mice were maintained under climate-controlled conditions of 12:12 – hours light/dark cycle temperature range $21\pm2^{o}C$, relative humidity of 30-70% and fed ad libitum a commercial food pellet diet. Studies were conducted at the Technion, (IIT), Faculty of Medicine, Haifa, Israel, after obtaining approval from the institute's IACUC. All proceedings complied with the Animal Welfare Act of 1966 (P.L. 89-544), as amended by the Animal Welfare Act of 1970 (P.L.91-579) and 1976 (P.L. 94-279).

7-bromoheptanol (Sigma-Aldrich, 310913) (7BRO) 325 mM in DMSO (Sigma-Aldrich, D2650) was injected to adult mice retro-orbitally (15 µl) and intraperitoneally (50 µl) to achieve complete blocking. (Sigma-Aldrich, D2650)

Janelia Fluor-525 HaloTag ligand was a kind gift from Luke D. Lavis. One µl of 1µM stock solution was diluted in 15µl DMSO and injected retro-orbitally. The retro-orbital injections were done under sedation with isoflurane (Piramal, Isoflurane, USP Terrell).

In-vivo AAV overexpression

Adult mice (4 month old) recieved 7BRO (or control no 7BRO). After 3 or 72 hours the mice received Janelia Fluor-525 HaloTag ligand retro-orbitally. Mice were sacrificed, and hearts were harvested 40 minutes after the ligand injection.

In-vivo experiments in HaloTag-TTN knock-in mice

The HaloTag-TTN knock-in mice were previously described[31]. Adult mice (4-month-old) received 7BRO (or control no 7BRO). After either 3 hours, 3, 7, or 14 days the mice received the Janelia Fluor-525 HaloTag ligand. Mice were sacrificed, and hearts were harvested 40 minutes after the ligand delivery. For hypertrophy, 9mM PE (Sigma-Aldrich, P6126) and 0.2% ascorbic-acid (Sigma-Aldrich, 95209) in PBS were filtered through a 0.22 µm filter. Mice received three subcutaneous injections of 10 mg/kg bodyweight PE on days -7, -5, and -3, with day 0 being the day of the sacrifice.

Heart sections preparation

After harvest, hearts were incubated in 4% formaldehyde for 1-2 hours on ice, and then moved to overnight incubation with gentle tilting in 50% sucrose in PBS solution. Next the hearts were submerged in Tissue Freezing Medium (Leica, 7876270), and frozen in isopentane cooled in liquid nitrogen. Cryosection was carried out by Leica CM1860 UV Cryostat, 5µm sections were performed, and mounted on a slide. Sections were then fixed with 4% formaldehyde for 10 min at room temperature, permeabilized with 1% Triton X-100 (Sigma-Aldrich) in PBS for 10 min, nuclear counterstained with DAPI (Sigma-Aldrich, D9542) at 1µg/ml in PBS, mounted with 15µl of Fluoromount G (Enco, 0100-01) and sealed with optic round cover slides. For the TTN knock-in mice sections were washed twice with PBS containing 0.3% Triton X-100 (Sigma-Aldrich), and incubated with Janelia Fluor-525 HaloTag ligand for 30 minutes at 37° Celsius. Then, sections were washed once again with PBS containing 0.3% Triton X-100 (Sigma-Aldrich), nuclear counterstained with DAPI (Sigma-Aldrich, D9542) at 1µg/ml in PBS, mounted with 15µl of Fluoromount G (Enco, 0100-01) and sealed with optic round cover slides.

Heart sections imaging

Heart sections were imaged using Zeiss LSM 880 with Airyscan super-resolution system, attached to Axio Examiner Z1 microscope. The acquisition was done with Plan-Apochromat 63x/1.4 Oil DIC M27 (#420782-9900-799) Zeiss objective, with diode laser 405nm wavelength and argon laser 514 nm wavelength. Images were then Airyscan processed with ZEN software (Zeiss) using the standard parameters.

In-vivo image analysis

A linear ROI of several sarcomeres' length was marked on every cell, and the signal intensity values were extracted. The representative value was calculated the same way as in the live cells

imaging analysis. We did not include the 0- and 3-hours controls (background) data in the fitting because of the low signal to noise ratio at the early time points. We fitted the data with a 1-exponential decay model, limited to a maximal value of the average of the No 7BRO control group (c), as this control represents the steady-state protein level (Eq.4)

$$Eq.\,4: \quad intensity = c - K * e^{(-t*\lambda)}$$

Statistics

Data are shown as mean ± SE as indicated. A two tailed student's t-test was used for comparison.

Funding

Funding for this work was provided by the Israel Science Foundation (grant # 1385/20) to IK and by the Fondation Leducq Research grant no. 20CVD01 to IK and LC. JAC acknowledges funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No. [101002927]). The CNIC is supported by the Instituto de Salud Carlos III (ISCIII), the Ministerio de Ciencia e Innovación (MCIN, MCIN/AEI/10.13039/501100011033) and the Pro CNIC Foundation and is a Severo Ochoa Center of Excellence (grant CEX2020-001041-S funded by MCIN).

Acknowledgements

The authors wish to thank the Biomedical Core Facility at the Faculty of Medicine, Technion, the Technion Genomic Center, and the Pre – Clinical Research Authority at the Technion. The authors wish to thank **Luke D Lavis,** Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, Virginia, USA for his generosity in providing the Janelia Fluor HaloTag ligands.

1. Mathieson, T. *et al.* Systematic analysis of protein turnover in primary cells. *Nature Communications 2018 9:1* **9**, 1–10 (2018).

2. Spalding, K. L., Bhardwaj, R. D., Buchholz, B. A., Druid, H. & Frisén, J. Retrospective birth dating of cells in humans. *Cell* **122**, 133–143 (2005).

3. Goehring, N. W. & Hyman, A. A. Organelle growth control through limiting pools of cytoplasmic components. *Current Biology* vol. 22 R330–R339 Preprint at https://doi.org/10.1016/j.cub.2012.03.046 (2012).

4. Kortazar, D. *et al.* Role of cofactors B (TBCB) and E (TBCE) in tubulin heterodimer dissociation. *Exp Cell Res* **313**, 425–436 (2007).

5. Kotila, T. *et al.* Mechanism of synergistic actin filament pointed end depolymerization by cyclase-associated protein and cofilin. *Nat Commun* **10**, (2019).

6. Marsh, J. A. & Teichmann, S. A. Structure, dynamics, assembly, and evolution of protein complexes. *Annual Review of Biochemistry* vol. 84 551–575 Preprint at https://doi.org/10.1146/annurev-biochem-060614-034142 (2015).

7. Gallastegui, N. & Groll, M. The 26S proteasome: assembly and function of a destructive machine. *Trends Biochem Sci* **35**, 634–642 (2010).

8. Taggart, J. C., Zauber, H., Selbach, M., Li, G. W. & McShane, E. Keeping the Proportions of Protein Complex Components in Check. *Cell Syst* **10**, 125–132 (2020).

9. Yin, Z., Ren, J. & Guo, W. Sarcomeric protein isoform transitions in cardiac muscle: A journey to heart failure. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1852**, 47–52 (2015).

10. Martin, A. Turnover of cardiac troponin subunits. Kinetic evidence for a precursor pool of troponin-I. *Journal of Biological Chemistry* **256**, 964–8 (1981).

11. Hamdani, N. *et al.* More severe cellular phenotype in human idiopathic dilated cardiomyopathy compared to ischemic heart disease. *J Muscle Res Cell Motil* **31**, 289–301 (2010).

12. Chen, C. Y. *et al.* Suppression of detyrosinated microtubules improves cardiomyocyte function in human heart failure. *Nat Med* **24**, 1225–1233 (2018).

13. Sanger, J. M. & Sanger, J. W. The dynamic Z bands of striated muscle cells. *Sci Signal* **1**, (2008).

14. Islam, M., Diwan, A. & Mani, K. Come Together: Protein Assemblies, Aggregates and the Sarcostat at the Heart of Cardiac Myocyte Homeostasis. *Frontiers in Physiology* vol. 11 586 Preprint at https://doi.org/10.3389/fphys.2020.00586 (2020).

15. Rudolph, F. *et al.* Resolving titin's lifecycle and the spatial organization of protein turnover in mouse cardiomyocytes. *Proc Natl Acad Sci U S A* **116**, 25126–25136 (2019).

16. Hassoun, R. *et al.* Do they come together? Protein quality control, stress-activated signaling, and "sarcostat" in hypertrophic cardiomyopathy progression. *International Journal of Cardiology* vol. 347 44–45 Preprint at https://doi.org/10.1016/j.ijcard.2021.11.002 (2022).

17. Liu, W. & Bu, H. Sarcostat mechanism: A promising therapeutic strategy for misfolded proteins in cardiomyopathy. *European Journal of Clinical Investigation* vol. 52 Preprint at https://doi.org/10.1111/eci.13725 (2022).

18. Luo, F. & Bu, H. The sarcostat: A proposed framework to understand sarcomeric protein quality control system. *International Journal of Cardiology* vol. 344 140 Preprint at https://doi.org/10.1016/j.ijcard.2021.10.018 (2021).

19. Skwarek-Maruszewska, A., Hotulainen, P., Mattila, P. K. & Lappalainen, P. Contractility-dependent actin dynamics in cardiomyocyte sarcomeres. *J Cell Sci* **122**, 2119–2126 (2009).

20. Suzuki, H., Komiyama, M., Konno, A. & Shimada, Y. Exchangeability of actin in cardiac myocytes and fibroblasts as determined by fluorescence photobleaching recovery. *Tissue Cell* **30**, 274–280 (1998).

21. Kostan, J. *et al.* Molecular basis of F-actin regulation and sarcomere assembly via myotilin. *PLoS Biol* **19**, e3001148 (2021).

22. Wang, J. *et al.* Dynamics of Z-band based proteins in developing skeletal muscle cells. *Cell Motil Cytoskeleton* **61**, 34–48 (2005).

23. Wang, J. *et al.* Ectopic expression and dynamics of TPM1α and TPM1κ in myofibrils of avian myotubes. *Cell Motil Cytoskeleton* **64**, 767–776 (2007).

24. da Silva Lopes, K., Pietas, A., Radke, M. H. & Gotthardt, M. Titin visualization in real time reveals an unexpected level of mobility within and between sarcomeres. *J Cell Biol* **193**, 785–798 (2011).

25. Auerbach, D. *et al.* Different domains of the M-band protein myomesin are involved in myosin binding and M-band targeting. *Mol Biol Cell* **10**, 1297–1308 (1999).

26. Welikson, R. E. & Fischman, D. A. The C-terminal IgI domains of myosin-binding proteins C and H (MyBP-C and MyBP-H) are both necessary and sufficient for the intracellular crosslinking of sarcomeric myosin in transfected non-muscle cells. *J Cell Sci* **115**, 3517–3526 (2002).

27. Los, G. V. *et al.* HaloTag: A novel protein labeling technology for cell imaging and protein analysis. *ACS Chem Biol* **3**, 373–382 (2008).

28. Merrill, R. A. *et al.* A robust and economical pulse-chase protocol to measure the turnover of HaloTag fusion proteins. *Journal of Biological Chemistry* **294**, 16164–16171 (2019).

29. Grimm, J. B., Brown, T. A., English, B. P., Lionnet, T. & Lavis, L. D. Synthesis of Janelia Fluor HaloTag and SNAP-Tag Ligands and Their Use in Cellular Imaging Experiments BT - Super-Resolution Microscopy: Methods and Protocols. *Methods in Molecular Biology* **1663**, 179–188 (2017).

30. Deo, C. *et al.* The HaloTag as a general scaffold for far-red tunable chemigenetic indicators. *Nat Chem Biol* **17**, 718–723 (2021).

31. Rivas-Pardo, J. A. *et al.* A HaloTag-TEV genetic cassette for mechanical phenotyping of proteins from tissues. *Nat Commun* **11**, (2020).

32. Suzuki, K. *et al.* In vivo genome editing via CRISPR/Cas9 mediated homology-independent targeted integration. *Nature* **540**, 144–149 (2016).

33. Platt, R. J. *et al.* CRISPR-Cas9 knockin mice for genome editing and cancer modeling. *Cell* **159**, 440–455 (2014).

34. Martin, T. G. & Kirk, J. A. Under construction: The dynamic assembly, maintenance, and degradation of the cardiac sarcomere. *Journal of Molecular and Cellular Cardiology* vol. 148 89–102 Preprint at https://doi.org/10.1016/j.yjmcc.2020.08.018 (2020).

35. Kitagawa, S. Experimental manipulation of calpain activity in vitro. in *Methods in Molecular Biology* vol. 1915 209–218 (Humana Press Inc., 2019).

36. Scarborough, E. A. *et al.* Microtubules orchestrate local translation to enable cardiac growth. *Nat Commun* **12**, (2021).

37. Palermo, J., Gulick, J., Colbert, M., Fewell, J. & Robbins, J. Transgenic remodeling of the contractile apparatus in the mammalian heart. *Circ Res* **78**, 504–509 (1996).

38. Lau, E. *et al.* A large dataset of protein dynamics in the mammalian heart proteome. *Sci Data* **3**, 160015 (2016).

39. Mueller, F., Morisaki, T., Mazza, D. & McNally, J. G. Minimizing the impact of photoswitching of fluorescent proteins on FRAP analysis. *Biophys J* **102**, 1656–1665 (2012).

40. Cadar, A. G. *et al.* Real-time visualization of titin dynamics reveals extensive reversible photobleaching in human induced pluripotent stem cell-derived cardiomyocytes. *Am J Physiol Cell Physiol* **318**, C163–C173 (2020).

41. Golan-Lagziel, T. *et al.* Analysis of rat cardiac myocytes and fibroblasts identifies combinatorial enhancer organization and transcription factor families. *J Mol Cell Cardiol* **116**, 91–105 (2018).

42. Ehler, E., Moore-Morris, T. & Lange, S. Isolation and culture of neonatal mouse cardiomyocytes. *J Vis Exp* (2013) doi:10.3791/50154.

43. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature Methods 2012 9:7* **9**, 676–682 (2012).

44. Fornasiero, E. F. & Savas, J. N. Determining and interpreting protein lifetimes in mammalian tissues. *Trends Biochem Sci* **48**, 106–118 (2023).

45. Challis, R. C. *et al.* Systemic AAV vectors for widespread and targeted gene delivery in rodents. *Nat Protoc* **14**, 379–414 (2019).

46. Schlesinger-Laufer, M. *et al.* A simple adeno-associated virus-based approach for the generation of cardiac genetic models in rats. *F1000Res* **9**, (2020).

Figure S1. Models of sarcomere maintenance.

Schematic illustration of the photobleach pulse-chase experiment and the possible outcomes according to two sarcomere maintenance models. Halo tagged proteins are expressed using adenoviral vector and labeled with green fluorescent ligand, that is bleached in a small square area. The cardiomyocytes are then incubated with a far-red ligand. **a**, The protein pool model predicts the existence of a cytoplasmic pool of sarcomeric proteins composed of both recycled (green labeled) proteins and newly translated (far red labeled) proteins. Since the photobleached proteins are expected to recycle proteins with this pool, the bleached area is expected to fill with both green and red labeled proteins. **b**, The ordered assembly model on the other hand predicts that only new proteins can enter the structure and therefore the bleached proteins will only be replace by far-red labeled ones.

Figure S2. Recovery of Oregon Green labeled ACTN2-Halo fluorescence in fibroblasts after bleaching

a, Representative images from a FRAP experiment performed in fibroblasts transduced with ACTN2-Halo and labelled with Oregon green. Images show the same fibroblast before the bleach (pre-bleach), and at 0,1, and 18 minutes after the bleach, with almost complete recovery of the green signal within 1 minute. The dashed white square is the photobleached field. The dotted yellow rectangle is the region of interest used for quantification. Scale bar = 10 μm. **b**, Quantification of the green florescence recovery after photobleaching. We calculated the photobleaching rate (r) by comparing the fluorescence of the non-frapped region of the cell before (Fc0) and after photobleaching (Fc).

$$r = \frac{Fc}{Fc0}$$

Then, we normalized the fluorescence intensity of the photobleached area by dividing its raw intensity (Fs) by the photobleaching rate (r).

$$F = \frac{Fs}{r}$$

Figure S3. Semi-automated image analysis of sarcomere turnover pipeline

a, representative image of pulse-chase experiment of cardiomyocytes transduced with Halo-tagged MYOM1, pre- and after photobleach, over 72 hours. The image is a merge of Oregon Green signal of pre-exiting proteins, and JF-635 ligand far-red signal of newly translated proteins. The numbers above indicate the time in hours after the photobleach. The dashed white square indicates the bleached area. The yellow rectangles labeled b & c are the region of interest (ROI) chosen for the quantification. Scale bar = 20 µm. **b-c,** Fluorescent intensity of the ROI outside and inside the photobleached area (b & c respectively) is shown in green for the pre-exiting proteins, and in red for the newly translated proteins. Each graph represents one timepoint according to the representative image above. The distance is shown in µm, and the signal intensity is shown in arbitrary units (AU). The peaks and the nadir points were identified using a moving window of 1 µm, and labeled by blue and turquoise dots respectively. **d,** Example of analysis of a single timepoint signal. The grey rectangles represent the difference between a nadir and the adjacent peak. The yellow rectangles represent the difference between a peak and the adjacent nadir point. The median of these differences was used as the representative value for the specific timepoint and channel.

Figure S4. Turnover rates in single sarcomeres are similar.
Single sarcomeric units from one myofibril in the unbleached region were individually analyzed using time lapse imaging to measure the decay of their green fluorescence intensity and to calculate their individual half-lives. The measured sarcomeric fluorescent intensity is shown on the left as dots, with fitted exponential decay (dotted lines) and calculated half-lives (bold dot) for each sarcomeric unit. Numbers show the range of calculated half-lives for the different sarcomeric units. Time lapse images are shown on the right, scale bar = 2µm. Two individual myofibrils, each containing several sarcomeric units are shown for each Halo-tagged sarcomeric protein: **a-b** ACTN2-Halo, **c-d** MYOM1-Halo, **e-f** Halo-MYL2, **g-h** Halo-MYBPC3. The analysis was extended beyond 72 hours in myofibrils where there was no significant movement. The images were taken every 12 hours instead of every 24 hours in some cells where the initial signal intensity was very strong.

Figure S5. HITI to knock-in HaloTag into the C-terminus endogenous *Actn2* gene.
a, Illustration of the HITI targeting approach: a Cas9 site (sgRNA target) that is followed by a PAM (NGG) site was identified in the C-terminal end of *Actn2* before the stop codon. The scissors show the predicted cut site (top). A donor cassette encoding the C-terminal end of *Actn2* with no stop codon, followed by in-frame HaloTag, and flanked by the same Cas9 sites

(sgRNA target) motifs was cloned into an AAV vector (middle). The direction of the sgRNA target sites flanking the donor cassette are inversed relative to the direction of the site in the genome. Integration of the cassette in the correct orientation to the genomic locus results in a chimeric junction that is no longer recognized by the sgRNA (bottom). Insertion in the inverse orientation on the other hand would re-create the sgRNA site and will be cut out again (not shown). **b**, Representative images from Cas9 expressing mouse cardiomyocytes transduced with Actn2-HaloTag donor cassette virus and with either control non-targeting sgRNA (left) or *Actn2* c-terminus sgRNA (right) encoding virus after addition of the Oregon Green Halo-Ligand (time = 0 hours) showing cross-striated pattern only in cardiomyocytes transduced with both donor and *Actn2* sgRNA. A 10 µm linear region of interest drawn along the myofibril is marked by a yellow rectangle. **c**, The green -fluorescence intensity along the linear region of interest drawn in b.

Figure S6. Phenylephrine induced hypertrophy model.

a, Boxplot of heart weight normalized to body weight ratio (HW/BW in mg/g) of titin (TTN)-HaloTag knock-in mice, which did or did not receive phenylephrine (PE) 10 mg/kg in three doses over 7 days. The whiskers indicate the results range of the group, and the box is the interquartile range (n=8-10 mice per group, Student's t-test p-value =0.000043). **b**, The PE injection protocol we used for the TTN-HaloTag knock-in mice, where mice received 3 injections in the last 7 days of experiment.

Figure S7. Titin is turned over at similar rates in different regions of the heart.

a, We separately calculated the turnover of titin (TTN) in three cardiac regions: the left ventricular free wall, the interventricular septum and right ventricular free wall. Representative cardiac sections from these regions are shown, left scale bar = 805 µm, right scale bar = 32 µm. **b**, The sarcomeric signal intensity is plotted as a function of time from the 7BRO blocker injection, and each dot represents the mean of the sarcomere signal intensity from one mouse. A fitted 1-exponential decay curve is shown as a dashed line, and the calculated half-life as a solid vertical line. The dots, horizontal dashed lines, solid lines, and turnover-half lives are colored by cardiac region. Half-life of TTN was similar in all regions. (n = 7-25 cells/mouse/region, from n = 1-7 mice/timepoint).