"Dark taxonomy": a new protocol for overcoming the taxonomic impediments

for dark taxa and broadening the taxon base for biodiversity assessment

Rudolf Meier[1,2]

Amrita Srivathsan[1]

Sarah Siqueira Oliveira[2]

Maria Isabel P.A. Balbi[3]

Yuchen Ang[4]

Darren Yeo[5]

Jostein Kjærandsen[6]

Dalton de Souza Amorim[3]

[1]Center for Integrative Biodiversity Discovery, Leibniz Institute for Evolution and Biodiversity

Science, Museum für Naturkunde Berlin

[2]Institute for Biology, Humboldt University Berlin

[3]Departamento de Ecologia, Instituto de Ciências Biológicas, Universidade Federal de Goiás

[4]Departamento de Biologia, FFCLRP, Universidade de São Paulo

[5]Lee Kong Chian Natural History Museum, Faculty of Science, National University of Singapore

[6]Centre for Wildlife Forensics, National Parks Board Singapore

[7]Tromsø University Museum, UiT The Arctic University of Norway

Corresponding author: Rudolf Meier (Rudolf.Meier@hu-berlin.de)

Abstract.

We are entering the 6[th] mass extinction event on the planet with scarcely any data for "dark taxa"

that comprise most animal species. These taxa are neglected, because conventional taxonomic

methods are not well-equipped to process tens of thousands of specimens belonging to thousands of

species. We here test a new protocol for tackling these taxa by simultaneously addressing (1)

taxonomic impediments, (2) lack of biodiversity baseline data, (3) and low impact of revisionary

research. We propose to overcome the taxonomic impediments by carrying out revisions at small

geographic scales to keep the number of specimens and species manageable. To lower the risk of

taxonomic error, species delimitation and description are based on multiple data sources

("integrative taxonomy"). Secondly, we propose to create baseline data for dark taxa by initially

revising them only based on fresh specimens obtained with the same sampling methods that are also

used for biodiversity monitoring. This ensures that the data from the revision can be used as a

biodiversity baseline and the species most relevant for biomonitoring are revised first. Thirdly, we

propose to improve the impact of taxonomic revisions by publishing the results in two manuscripts

addressing different readerships (general and specialists). We illustrate our proposals by carrying

out a taxonomic revision of the fungus gnats (Diptera: Mycetophilidae) of Singapore based on

specimens obtained with Malaise traps placed at 107 sites across different habitats. We show that a

first batch of specimens (N=1,454) contains 120 species, of which 115 are new to science and

described in a separate taxonomic monograph. Species delimitation started with obtaining NGS

barcodes that were used for estimating the number of MOTUs (Molecular Operational Taxonomic

Units). This revealed 115-128 MOTUs depending on whether PTP, ABGD (P=0.001-0.060), or

objective clustering (2-5%) was used. MOTU boundaries were then revised with "LIT" (Large-

scale Integrative Taxonomy) which optimizes the integration of morphological and molecular data

by using the molecular data. LIT revealed a match ratio of 91% for 3% objective clustering

MOTUs, but an even higher match ratio of 97% was found for MOTUs obtained with 5% objective

clustering and ABGD (P=0.060). Using these parameters, only two MOTUs had to be fused for

perfect congruence between molecular and morphological data for the 120 species. To test the

completeness of our monograph, we then analyzed a second batch of 1,493 specimens. All clustering algorithms and settings revealed only 18 additional MOTUs; i.e., the first batch of specimens already covered 87% of the diversity, although 22% and 9% of the species are singletons or doubletons, respectively. Remarkably, >97% of all specimens in the second batch belonged to species described based on the first batch. The study thus demonstrates that revisions of dark taxa at moderate geographic scales with a moderate number of specimens from standardized traps can move dark taxa from being unavailable for biomonitoring to being well suited. In addition, the study increases the number of described species of Mycetophilidae in the Oriental Region by >25% indicating the size of taxonomic impediments for dark taxa in Southeast Asia.

Key words: Integrative Taxonomy, Dark Taxa, Reverse Taxonomic Workflow, Oriental Insect Fauna, NGS barcodes, Mycetophilidae

Introduction

Biodiversity assessment and monitoring are among the biggest and most urgent challenges of modern biology, given that many natural environments are disappearing fast and biodiversity loss is destabilizing whole ecosystems. However, due to taxonomic impediments, biodiversity assessment is difficult for many arthropod clades (e.g., Wheeler et al. 2004; Evenhuis 2007; de Carvalho et al. 2007). The problem is particularly severe in the tropics and for taxa characterized by high species diversity and abundance. For such taxa, conventional approaches to taxonomy involving morphospecies sorting tend to be ineffective, because they were not designed with hyperabundant and hyperdiverse taxa in mind (Hartop et al. 2022). Taxonomists have long been aware of these constraints and, thus, focused on projects covering only moderate numbers of species and dealt with a small number of specimens pertaining to taxa that are overall already well-known (i.e., with prior revisions, identification keys, illustrations etc.). Projects that satisfied these criteria have been readily available and selecting them has been good career advice for young taxonomists. However,

this also meant that many truly species-rich and abundant taxa remained neglected (Hartop et al.

2022; Srivathsan et al. 2023). Arguably, this has created a special kind of taxonomic impediment

("dark taxon impediment": Meier et al. 2022). Addressing this impediment has become high

priority because taxonomic chauvinism (Bonnet et al. 2002; Troudet et al. 2017) needs to be

overcome in an era where we need quantitative data on all of biodiversity.

What is needed are new taxonomic techniques that are scalable. Some examples illustrate how

scalable they have to be: a recent study barcoded 7,059 specimens of scuttle flies (Diptera:

Phoridae) collected by one Malaise trap over only eight weeks in a Ugandan National Park and

found evidence for >650 species (Srivathsan et al. 2019). Yet, only 462 scuttle fly species have

been described for the entire Afrotropical region (Phorid Catalog, 2023: accessed 13 August 2023).

Similarly, the diversity of the gall midges (Diptera: Cecidomyiidae) has been estimated to be 800+

species at a single site of Costa Rica (Brown et al. 2018, Borkent et al. 2018) and a species diversity

of gall midges close to 8,500 was estimated for Canada based on barcodes (Hebert et al. 2016),

although fewer than 250 Nearctic species have been described for this family (Savage et al. 2019).

These cases illustrate why truly hyperdiverse taxa are sometimes called "open-ended." They will

frustrate attempts at carrying out traditional comprehensive taxonomic revisions at reasonable

geographic scales (Bickel 2009). An alternative term that is nowadays also frequently used for such

clades is "dark taxa" (Hausmann et al. 2020, Hartop et al. 2022; Page 2011, 2016), which Hartop et

al. (2022) suggested should be restricted to clades where the undescribed fauna is estimated to

exceed the described fauna by at least one order of magnitude and the total diversity exceeds 1,000

species worldwide. These taxa need a different set of taxonomic protocols, which we here propose

to call "dark taxonomy" to highlight that these techniques are fundamentally different from what is

practiced for less diverse/abundant taxa.

The first key principle of "dark taxonomy" is embracing "faunistic sampling", i.e., the

geographic scope of a taxonomic revision is determined by the number of specimens and species

that can be reasonably handled and not by the putative range of species. Implicitly, almost all

"turbo-taxonomy" studies have embraced such sampling without much discussion (e.g., Fernandez-

Triana et al. 2014, 2022; Marsh et al., 2013; Riedel et al., 2013a,b; Riedel and Narakusumo 2019,

but see Dijkstra et al. 2015). The lack of open discussion is presumably because it goes against the

traditional recommendation that taxonomic revisions should cover all known species/specimens of a

clade at a geographic scale appropriate for the species included in the clade. For example, if many

species in a clade are found throughout the Palaearctic, one would carry out the revision at this

geographic scale in order to properly assess species diversity and intraspecific variability. Such

"completism" is desirable, but also contributes to taxonomic chauvinism because it cannot be

applied to dark taxa. It often delays embracing solutions needed for overcoming our taxonomic

ignorance in key taxa. Two ways to responsibly relax the requirements is to limit taxonomic

treatments to a small geographic area ("faunistics") and/or only work on a subset of the available

specimens. Our revision of Mycetophilidae (Diptera; fungus gnats) is an extreme example for both.

We only revise the fauna of a small island (Singapore, 730 km^2) and only used recently collected

specimens obtained with Malaise traps.

Faunistic sampling carries obvious risks. The most important one is incorrect species

delimitation because intraspecific genetic variability is poorly assessed for wide-spread species

(Bergsten et al. 2012). However, we predict that this error is manageable by applying the principles

of integrative taxonomy; i.e., species delimitation based on multiple data sources (Dayrat 2005). It

is likely that containment will be particularly effective if the characters used for species delimitation

were generated by different evolutionary processes. With regard to morphology, taxonomists

usually collect information from a broad array of structures including genitalia, legs, wings etc.

Genitalia are likely to be under strong sexual selection (e.g., Eberhard 1985, Rowe and Arnqvist

2012), while many other body parts are presumably more shaped by viability selection. It thus

appears probable that most speciation events will leave traces in such a diverse array of structures,

regardless of whether speciation was fast or slow. Of course, there are exceptions as is well known

from the literature on "cryptic species" (Bickford et al. 2007). Therefore, morphology should be

paired with a type of data that evolves largely via drift and thus reflects time of divergence. This is

the case for DNA barcodes, given that closely related species mostly differ with regard to

synonymous sites (Kwong et al. 2012; Pentinsaari et al. 2016). Large DNA barcode distances can

therefore flag cases where morphological differentiation is lower than expected based on time of

divergence. This can then trigger more detailed work to resolve species limits for taxa that have

high intra- or low interspecific variability ("reciprocal illumination"). We have repeatedly found

this to be very effective for resolving taxonomic problems for a range of fly families (Ang et al.

2013, Tan et al. 2010, Rohner et al. 2014, Laamanen et al. 2003).

The second element for tackling large numbers of specimens is what we call the "reverse

workflow.". Traditionally, species discovery has started with sorting specimens to species using

morphology. Subsequently, the morphospecies were validated with few barcodes from individuals

representing the morphospecies. This workflow is ineffective for dark taxa, where even samples

collected at a small geographic scales routinely contain too many specimens and species for

morphospecies sorting. Imagine having to sort the aforementioned >7,000 specimens of Ugandan

phorids into 650 species in this manner, or the 18,000 phorids in Hartop et al. (2022) into >350

species. As argued before, such samples should thus first be sorted to Molecular Operational

Taxonomic Units (MOTUs) with barcodes (Wang et al. 2018; Hartop et al. 2022). This is now

feasible because barcodes can be acquired semi-automatically and at low cost by staff who only

specialize in molecular methods (Meier et al. 2016; Srivathsan et al. 2019, 2021). Presorting large

numbers of specimens to putative species means that taxonomic experts can concentrate on revising

MOTU boundaries using other sources of data (Wang et al. 2018; Hartop et al. 2022).

Delimiting large numbers of species based on two types of data also requires explicit rules for

assessing and resolving conflict between data types. We recently proposed "Large-scale Integrative

Taxonomy" (LIT) for this purpose (Hartop et al. 2022). The LIT protocol addresses the "integrative

taxonomy conundrum" that collecting two different types of data for all specimens will slow down

taxonomy when faster speed is needed to address taxonomic impediments. Fortunately, the

conundrum can be overcome by speeding up collection of one type of data (e.g., semiautomatic

acquisition of barcodes) and then only testing MOTUs delimited with these data by subsampling

specimens for a second type of data (Hartop et al. 2022). This is what LIT implements. In the first

stage, barcodes are used to generate a set of MOTUs based on one criterion (e.g., clustering at 3%

pairwise distance). In the second step they are classified as "stable" or "instable" depending on

whether different algorithms and clustering thresholds change MOTU composition. In the third

stage, the MOTUs are tested by studying the morphology of specimens representing the main

and/or most divergent haplotypes— instable MOTUs are tested more extensively than stable ones

(see Hartop et al. 2022). In the fourth stage, those MOTUs that are congruent with morphological

evidence are accepted as species. The fifth stage is reserved for resolving conflict between

morphological and molecular evidence. As argued in Hartop et al. (2022), such conflict is expected

for recently diverged species which are expected to be incorrectly lumped into one MOTU (e.g.,

also due to introgression) and for "old" species with diverged allopatric populations which will be

split into multiple MOTUs. Conflict can thus usually be resolved by testing whether MOTU

boundaries obtained with different algorithms or clustering parameters are congruent with

morphological evidence. If so, these morphospecies can also be described because the molecular

and morphological data are not genuinely in conflict. If not, the MOTUs/morphospecies are left

unresolved until a third data source can be used to determine species boundaries. Hartop et al.

(2022) showed that only 5-10% of all specimens had to be checked to find congruent species

hypotheses for the ca. 350 MOTUs that were analyzed by studying 18,000 specimens.

Sorting specimens into putative species and revising the MOTU boundaries using LIT yields

species-level units. The next step is establishing which species are already described and which

need description. As illustrated in our companion monograph of mycetophilids of Singapore

(Amorim et al., 2023), the task of sorting through old descriptions is manageable for dark taxa in

the tropics because so few species have been described. The "superficial description impediment"

of Meier et al. (2022) may thus be somewhat less onerous for the taxa and regions of the planet that

are most in need for dark taxonomy. In other parts of the world, revisions are more likely to get

stuck after completing LIT because too many scientific names cannot be resolved. This is

frustrating, but it is already important progress when all species have been delimited based on two

types of data. Museomics is evolving quickly and accelerate the matching of delimited species

without names with barcodes from reliably identified or type material (e.g., Santos et al. 2023).

Developing high-throughput taxonomic techniques for dark taxa is important, but we will also

need more taxonomists to take on dark taxa. In proposing "dark taxonomy", we suggest two ways

for increasing impact. The first is that initial revisions of dark taxa at small geographic scales

should be based on specimens collected with the kind of sampling techniques that are also used for

biomonitoring biodiversity. For many groups of insects, this would be Malaise-, pitfall-, light-,

and/or flight intercept traps. Revising taxa based on such samples will simultaneously yield

biodiversity baselines and species descriptions that are immediately relevant for biomonitoring.

Furthermore, the DNA barcode database generated becomes a valuable resource for metabarcoding

and metagenomics projects. Fresh specimen samples are also easier to process at scale, because the

specimens are suitable for rapid barcoding and locality information is usually available in a digital

format.

The second strategy is to publish the results of such revisions in pairs of companion papers. The

first presents the results of interest to a broad readership, while the second with species descriptions

can be presented in a second monograph targeting taxonomic experts. To ensure that the

information in both manuscripts is available at the same time, we recommend joint publication as a

preprint. It should be duly noted, however, that this requires the registration of all scientific names

in Zoobank and the inclusion of a disclaimer according to section 8.2. of the ICZN code that the

preprint is not issued for the purpose of zoological nomenclature and should be considered not

published within the meaning of the code.

In this paper, we propose and test "dark taxonomy" by revising the fauna of Mycetophilidae of

Singapore based on an initial batch of 1,454 specimens. We then test the efficiency of the protocol

by using a second batch of 1,493 specimens to measure the proportion of species already described

based on the first batch. In addition, we illustrate the size of the taxonomic impediment for dark

taxa in the Oriental region by demonstrating that revising the fauna of a small island increases the

number of described species by >25%.

Material and Methods

The project proceeded through five phases, but the first would not have to be repeated for many

other dark taxa.

Phase 1: Collecting. Malaise traps were deployed for varying periods (2-6 months) between April

2012 and June 2019 at 107 sites in Singapore (Figs. 1A–B). The sites represent the following

habitat types in Singapore: degraded urban secondary forest, old and maturing secondary forest,

primary forest, coastal forest, swamp forest, freshwater swamp (lacking mature trees), and old-

growth and replanted mangrove forests (Fig. 2A–E). The samples were collected weekly and

preserved in 70% ethanol. Subsequently, they were sorted to order/family by parataxonomists and

entomologists. Mycetophilids were present in most sites, though not particularly abundant in some

habitats (e.g., mangroves). Details on the sampling campaign are in Yeo et al. (2021).

Phase 2: Estimating the number of MOTUs. We processed two batches of specimens. The first had

1,454 and the second, 1,493 specimens. The latter was sequenced after the cut-off date for the

descriptive monograph (Amorim et al. 2023). All specimens were sequenced for a 313-bp fragment

of the *cytochrome oxidase I* gene (*COI*) since such minibarcodes amplify better than the full-length

barcode (Srivathsan et al. 2021) and nevertheless yield similar species-level signal (Yeo et al.

2020). The minibarcode was amplified for each specimen using a protocol in Meier et al. (2016). In

the early phases of the study, DirectPCR (Wong et al. 2014) was employed; i.e., 1 or 2 legs of each

specimen were used as the DNA template for amplification. For specimens collected later, DNA

extraction was performed by immersing the whole specimen in Lucigen's QuickExtract solution or

HotSHOT buffer (Montero⬜Pau et al. 2008). PCR was performed with the following primer pair

mlCO1intF: 5'-GGWACWGGWTGAACWGTWTAYCCYCC-3' (Leray et al. 2013) and

jgHCO2198: 5'-TANACYTCNGGRTGNCCRAARAAYCA-3' (Geller et al. 2013). These primers

were labelled at the 5' end with 9-bp tags that differed from each other by at least three base pairs.

Every specimen in each sequencing library was assigned a unique combination of forward and

reverse primer tags. This allows for reads to be assigned to their specimen of origin. A negative

control was used for each 96-well PCR plate to detect contamination. Amplification success was

estimated via gel electrophoresis of a subsample of eight wells from each plate.

Figure 1. Singapore sites sampled with Malaise traps (modified from www.freeworldmaps.net). A.

Southeast Asia, with relative position of Singapore. B. Collecting sites in different environments.

Figure 2. Environments sampled in Singapore. A: Mangrove, B: Rainforest, C: Urban forest, D:

Swamp forest, E: Freshwater swamp.

Amplicons were pooled at equal volume within each plate. Equimolar pooling across plates was

performed by approximating concentration based on the presence and intensity of the gel bands.

The pooled samples were cleaned with Bioline SureClean Plus before being sent for library

preparation at AITbiotech using TruSeq Nano DNA Library Preparation Kits (Illumina) or the

Genome Institute of Singapore (GIS) using NEBNext DNA Library Preparation Kits (NEB).

Paired-end sequencing was performed on Illumina Miseq (2x300-bp or 2x250-bp) or Hiseq 2500

platforms (2x250-bp). The sequences were obtained across multiple runs, which allowed for

troubleshooting and re-sequencing specimens which failed to yield a sufficiently high number of

reads during the first sequencing attempt. Some of the specimens were also sequenced with

MinION (Oxford Nanopore) using primers labelled with slightly longer tags (13-bp; see Srivathsan

et al. 2019). The raw Illumina reads were processed with the bioinformatics and quality-control

pipeline described in Meier et al. (2016). A BLAST search to NCBI GenBank's nucleotide (nt)

database was also conducted to identify contaminant sequences by parsing the BLAST output with

readsidentifier (Srivathsan et al. 2015). Barcodes that matched non-target taxa at >97% identity

were discarded.

Barcodes lacking stop codons were aligned with MAFFT v7 (Katoh and Standley 2013) using

default parameters and analysed using three different species delimitation algorithms: objective

clustering (Meier et al. 2006), Automatic Barcoding Gap Discovery (ABGD: Puillandre et al. 2012)

and Poisson Tree Process (PTP: Zhang et al. 2013). Objective clustering was implemented via a

python script that implements the objective clustering algorithm in Meier et al. (2006) and we

obtained the results for four p-distance thresholds (2–5%) encompassing thresholds commonly used

in the literature for species delimitation (Ratnasingham and Hebert 2013). ABGD was performed

using uncorrected p-distances and the minimum slope parameter (-X) 0.1, with the default range of

priors (P = 0.001 – 0.1). Lastly, a maximum likelihood phylogeny was generated from the barcode

sequences in RAxML v.8 (Stamatakis 2014) with the GTRGAMMA model and with 20

independent tree searches. The barcode tree was then used for the PTP analysis using the

implementation provided in mPTP (--ml --single) (Kapli et al. 2017; Zhang et al. 2013) algorithms.

As described in Wang et al. (2019), the vials with the specimens were then physically sorted into

bags that corresponded to 3% p-distance clusters. Specimens from these bags were used for

morphological verification of species boundaries according to the rules of LIT.

Phase 3: Integrative Taxonomy

We started the morphological study before LIT was published by slide mounting at least one

male and one female (if available) for each 3% MOTU. The specimens were cleared with KOH,

dehydrated in ethanol, dissected to separate wings, abdomens with terminalia, and head/thorax. The

body parts were mounted in Euparal (modified from Walker and Crosby 1988; Huber and Reis

2011) under three separate coverslips. Particularly, the specimens that underwent DNA extraction

with Proteinase K overall worked well because the proteins were already digested, while the

exoskeleton was well preserved for slide mounting (Santos et al. 2018). However, some soft

specimens collapsed, affecting the quality of slide mounts.

This initially screen was then enhanced to also test the LIT rules outlined in Hartop et al. (2022).

All non-singleton barcode clusters obtained at 3% were classified as stable or instable based on a

stability index that quantifies whether cluster content changes when the clustering threshold is

increased from 1% to 3%. The clusters where the maximum pairwise distance between any two

specimens was >1.5% were also flagged as instable. All stable clusters were then tested by studying

the morphology of a pair of specimens representing the most distant haplotypes, while for all

instable clusters the morphology of one specimen representing the main haplotypes was also

studied. For most genera, the diagnoses include male terminalia features, such that only clusters

with males could provide sufficient information on conspecificity. For each cluster we then assessed

whether (1) the examined specimens from the same MOTU belonged to the same morphospecies

and (2) differed from the morphospecies in the "neighboring" clusters. If (1) was violated, the

cluster was split to test whether morphospecies and DNA barcode information were congruent at a

lower clustering threshold. If (2) was violated, it was tested whether fusing "neighboring" clusters

restored congruence with morphospecies. Eventually, we accepted those grouping decisions based

on DNA barcodes that were congruent with morphological information. No species delimitation

was performed when morphospecies were incongruent with all grouping statements based on

barcodes.

Phase 4: Species identification and description

To decide whether a specimen belonged to a described species or one that had to be newly

described, we extensively reviewed the literature on the Oriental fauna of Mycetophilidae, which is

here also cited because it forms the foundation of all taxonomic work on the group (Meier 2017).

For the Oriental species described by Edwards (1925, 1926, 1927, 1928, 1929, 1931, 1933, 1935,

1940), we studied and photographed primary types deposited at the Natural History Museum,

London. For more recently described Oriental species of Mycetophilidae, we were able to resolve

species identity based on published careful descriptions with good illustrations (Colless 1966;

Zaitzev 1982; Sivec and Plassmann 1982; Wu and Yang 1986; Bechev 1995; Søli 1996, 2002;

Kallweit 1998; Matile 1999; Xu and Wu 2002; Wu et al. 2003; Papp 2004; Hippa et al. 2005; Hippa

2006, 2007, 2008, 2009, 2011; Ševčík 2001; Ševčík and Hippa 2010; Hippa and Ševčík 2010, 2013;

Ševčík et al. 2011, 2012; Borkent and Wheeler 2012; Ševčík & Kjærandsen 2012; Kurina and

Hippa 2015; Hippa and Saigusa 2016; Hippa and Kurina 2018; Magnussen et al. 2019; Fitzgerald

2017; Kaspřák et al. 2017; Kjærandsen et al. 2023). Some Oriental species, especially those from

Sri Lanka, cannot be resolved because the descriptions are insufficiently detailed and the types were

lost or are unavailable. We here assumed that the species from Singapore are allospecific based on

geography, which is consistent with high endemism for Mycetophilidae species in South America

(Amorim and Santos 2018), although our revision (Amorim et al. 2023) also includes a species

whose distribution ranges from Japan through Singapore and Sumatra to Thailand. Future study will

have to reveal if any of the Sri Lankan species are conspecific with the material studied in

Singapore.

Phase 5: Completeness of the faunal assessment

The barcode data for the second batch of specimens (N=1,493) was used to test how

complete the revision based on the first batch of specimens was (N=1,454). For this purpose, the

genetic data were analyzed as described earlier for the first batch of sequences. Subsequently, we

assessed how complete the initial revision was in the light of the species diversity represented in the

second set of specimens. We also determined the number of new haplotypes, clusters, and how

many specimens from batch 2 belonged to species described based on batch 1. The new species are

currently being described. We then compared these observed values to those obtained by

randomized assignment of specimens to batch 1 and batch 2, assuming the same sample sizes (batch

1, N=1,454 and batch 2, N=1,493) in order to account for any effect of differences in sampling.

This was done using a custom script (https://github.com/asrivathsan/darktaxa). Lastly, we estimated

how many specimens would have had to be studied if LIT was performed on both batches of

specimens (N=2,947).

Results

Phase 1: Collecting

The material for this study was obtained during 3,526 Malaise trapping weeks distributed across

five different habitats in Singapore: mangroves (74 sites, 2,162 trap weeks), tropical forest (nine

sites, 567 weeks), urban forests (15 sites, 280 weeks), swamp forest (four sites, 262 weeks), coastal

forest (10 sites, 156 weeks), and freshwater swamp (seven sites, 99 weeks). A total of 69 traps

collected 3,030 mycetophilid specimens that were successfully sequenced. The overall number of

mycetophilid specimens collected is somewhat lower than expected for such a large number of

Malaise trap samples due to fact that almost two thirds of the samples (61%) were obtained from

mangroves that are comparatively poor for Mycetophilidae.

Phase 2: Molecular processing

The 1,454 COI barcodes from the first batch clustered into 115–128 MOTUs via objective

clustering at 2–5% uncorrected p-distances (Table 1). With ABGD, the number of MOTUs range

from 115–128. PTP clustered the barcodes into 127 MOTUs respectively. The 1,493 COI barcodes

from the second batch yielded 95 MOTUs from 5% p-distance clustering.

Phase 3: Integrative Taxonomy

Congruence between morphospecies and MOTUs was assessed with match ratios as described in

Ahrens et al. (2016). It was overall high (0.87-0.97), with the highest match ratio observed for 5%

clusters and MOTUs obtained with ABGD at P=0.06 (0.97, 113 morphospecies congruent).

Table 1. Integrative species delimitation using different algorithms for MOTU estimation. The

number of morphospecies with molecular data was 117.

Algorithm	PTP	Objective Clustering				ABGD				
		2%	3%	4%	5%	P=0.001 -0.005	P=0.008	P=0.013 -0.022	P=0.036	P=0.06 0
No. MOTUs	127	128	124	121	115	126	125	124	122	115
No. congruent morphospecies	108	107	110	113	113	109	110	110	112	113
Split morphospecies	9	10	7	4	0	8	7	7	5	0
Fused morphospecies	0	0	0	0	2	0	0	0	0	2
Match ratio	89%	87%	91%	95%	97%	90%	91%	91%	94%	97%

367   *Phase 4: Species identification, description and naming*

368  The application of LIT to the 124 MOTUs (3%) required morphological study of 224 specimens.

369 Of these 124 MOTUs, 27 were singletons, 73 MOTUs were stable, and 24 were instable (530

370 specimens). One challenge with carrying out the LIT analysis is that for approx. 40% of haplotypes,

371 only female specimens were available. This weakens the test for congruence with MOTU

372 boundaries. Overall, congruence with morphospecies required fusing closely related MOTUs in

373 seven cases (Figure 3; 7x2 MOTUs). Subsequently, 117 species were delimited and described

374 belonging to 21 known and one new genus (note that the monograph also describes three additional

375 species based on morphology only: Amorim et al. 2023). Only five species were already described

376 prior to this revision: *Chalastonepsia hokkaidensis* Kallweit, *Eumanota racola* Søli, *Metanepsia*

377 *malaysiana* Kallweit, *Parempheriella defectiva* Edwards, *Neoempheria dizonalis* EdwardsThe

378 remaining 112 new species with molecular and morphological data are described and illustrated in

379 the companion taxonomic monograph (Amorim et al. 2023). Images are also available from digital

380 reference collection "Biodiversity of Singapore" (https://singapore.biodiversity.online/taxon/A-

381 [Arth-Hexa-Dipt-Mycetophilidae](https://singapore.biodiversity.online/taxon/A-Arth-Hexa-Dipt-Mycetophilidae)).

383 *Phase 5: Completeness of the faunal assessment*

384  The second batch of specimens yielded 94 MOTUs at 5% in a combined analysis of all the

385 2,947 barcodes from both batches. 5% MOTUs were here used for clustering because it maximized

386 congruence between morphology and molecular data for batch 1. Of these, 76 belonged to MOTUs

387 already examined for species descriptions based on the specimens in the first batch (76/94 or 80%

388 of species in batch 2). All 18 newly discovered MOTUs were stable (no compositional changes with

389 all delimitation algorithms and settings used here) and overall so rare that they only represented 31

390 of the 1,493 specimens in the second batch. Overall, 97.9% of all specimens in batch 2 belonged to

391 MOTUs that were also found in batch 1 (Figure 4). In order to assess if the observed coverage of

392 species and specimens for batch 2 was influenced by sampling, we compared these values to those

393 obtained by randomized assignments of specimens to batch 1 and batch 2 (100 randomizations).

Randomized assignments of specimens to batch 2 revealed coverage that closely reflected observed

values: 82 ± 2.7% of MOTUs and 98.1 ± 0.4% of specimens would belong to MOTUs found in

batch 1 (Figure 4). Thus, if batch 1 is well studied in terms of integrative taxonomy, very few

specimens in batch 2 would belong to new 5% MOTUs and require additional taxonomic work.

Figure 3. Most MOTUs are stable and congruent with morphospecies (green; N=100). Remaining

species have splitting (red) and lumping (yellow) conflict with molecular data depending on species

delimitation algorithm.

At 5% objective clustering, the proportion of singletons was 22% (N=29) for all specimens, 17%

(N=25) for dataset 1, and 26% (N=24) for dataset 2. The corresponding numbers for doubletons

were 11% (N=14), 9% (N=11), and 16% (N=15). Carrying out a LIT analysis for all 2,947

specimens would have required checking 143 (3%) MOTUs of which 30 were singletons, 79 stable,

and 34 were instable (1,741 specimens). Despite the large number of specimens in the instable

clusters, only 272 specimens of the combined dataset would have to be checked to sample

haplotypes according to LIT.

Table 2. MOTU (3%) and specimen abundance for the second batch of specimens and all data.

Dataset	Batch 1	Batches 1 + 2
3% MOTUs (No. of specimens)	124 (1454)	143 (2947)
Singleton MOTUs	27	30
Stable MOTUs (No. of specimens)	73 (897)	79 (1176)
Instable MOTUs (No. of specimens)	24 (530)	34 (1741)
LIT: No. of specimens to be examined (singleton/stable/instable)	224 (27/130/67)	272 (30/136/106)

Figure 4. An overwhelming number of specimens and most species in specimen batch 2 belong to

species described based on specimens in batch 1 (~80% of species and >97% of specimens).

Dashed lines represent the observed values while the solid lines represent the results of 100

randomizations.

Discussion

Biodiversity loss is now considered one of the top five threats to planetary health (World

Economic Forum 2023). Yet, most animal species are undescribed, unidentifiable, and lack baseline

data for biomonitoring. Much of this diversity apparently belongs to dark taxa such as the top 20

species-rich families of flying insects that were identified in Srivathsan et al. (2023) based on

samples from five biogeographical regions. For many of these taxa, we know how to obtain

standardized quantitative samples (e.g., Malaise, pitfall, flight intercept traps), but most samples are

currently only evaluated with metabarcoding (Yu et al., 2012). However, bulk sequencing is unable

to overcome taxonomic impediments because it disrupts the association between specimen and

genetic data. In addition, metabarcoding only provides approximate abundance/biomass data. Note,

however, that even the barcoded specimens with vouchers in BOLD largely remain

morphologically unidentified beyond family level. For example, in 2021 95% of all Sciaroidea and

60% of all Mycetophilidae specimens were found to lack species-level identifications (Kjærandsen

2022). To overcome such problems, we propose that some samples for each site should be analyzed

with "dark taxonomy" to delimit species, obtain species-specific barcode libraries, facilitate

taxonomic documentation through description and naming, and yet generate a first set of

biodiversity baseline data also useful for subsequent biomonitoring with bulk sequencing

approaches.

Our study shows how this is a feasible goal even for dark taxa. We here show that dark

taxonomy can fairly quickly yield a good estimate of species diversity and provide species names

and barcodes for most common species belonging to a dark taxon. Our initial set of specimens

consisted only of 1,454 specimens. They are shown in the companion monograph to belong to 117

species delimitated with morphological and molecular data. We then tested the completeness of the

revision by using the data for an additional 1,493 specimens. Fortunately, the observed number of

MOTUs increased by only 16% (N=18); i.e., the first richness estimate was more than 80%

complete. Furthermore, 76 of the 117 species in the first batch were also represented in the second

batch. Not surprisingly, this involved mostly common species, so that the proportion of specimens

from the second batch that belong to species described in the first batch is very high (>97%). This

means that one moderately scaled revision of a dark taxon was sufficient to ready a country like

Singapore for the inclusion of this taxon (Mycetophilidae) in biomonitoring. This is impressive

given that the number of known and described species of Mycetophilidae alone in Singapore now

exceeds the number of species in major vertebrate groups (mammals: 83 spp.; amphibians: 26 spp.;

"reptiles": 109 spp; source: Singapore National Parks Board [Red Data Book] Species List, 2023).

Moreover, all the new species can be identified based either on barcodes or morphology and

sampled minimally invasively with Malaise traps.

Currently, very little is known about the natural history of tropical fungus gnats so it is difficult

to assess whether they could be used as indicator taxa for habitat quality or fungus richness.

However, this seems likely, given the dependency of Mycetophilidae on fungi and studies such as

the ones by Økland (1994, 1996) demonstrating that mycetophilids are good indicators of temperate

forest quality. However, more environmental correlates need to be collected for the tropics and it

will be important to associate larvae and adults (see Yeo et al. 2018) to understand to what extent

mycetophilid species are specialized on specific species of fungi and how mycetophilid species-

richness and abundance could be used as an indirect evidence of tropical forest health.

To fully appreciate the potential of dark taxonomy, it is important to consider that Srivathsan et

al. (2023) showed that more than half of the specimens and species in Malaise trap samples belong

to approximately 20 family-level dark taxa (see also Brown 2005; Karlsson et al. 2020). This means

that applying a technique like "dark taxonomy" to a few complete Malaise trap samples could

simultaneously yield biodiversity baseline data for many numerically dominant families and

hundreds of species simultaneously. This is realistic, because barcoding is now so easy and cheap

(Srivathsan et al. 2021) that dark taxonomy is frugal and thus suitable for biodiverse countries with

limited science funding (Brydegaard et al. 2024). PCR costs can be as low as 0.05 USD per

specimen, and sequencing with Illumina's NovaSeq costs only 0.01-0.02 USD /barcode (Srivathsan

et al. 2021). A self-contained barcoding lab covering all steps from specimens to barcodes can be

installed for <USD 10,000 if it utilizes Oxford Nanopore Technologies (ONT) sequencers (e.g.,

MinION; Srivathsan et al. 2024). The use of 3^{rd} generation sequencing technologies raises the

sequencing cost for DNA barcodes, but it still means that the consumable cost for barcoding the

~3,000 specimens in our study costs USD 150 (Illumina) to USD 375 (ONT). Specimen handling

costs are also dropping with the use of DIY microscopes (Wührl et al. 2024) and robots for sample

sorting (see Wührl et al. 2022). Carrying out taxonomic revisions for dark taxa will inevitably

remain challenging but "dark taxonomy" can help with simultaneously addressing taxonomic

impediments and the need for biodiversity baseline data.

<u>Objections to dark taxonomy</u>

The proposal of "dark taxonomy" is likely to be controversial for a variety of reasons. One is that

specimens/types from dry collections will initially only be used if they are important for resolving

species names, i.e., only barcodes for types or reliably identified specimens are important.

However, many rare and most extinct species are only found in dry collections. They are rarely or

never found in the kind of traps used for biomonitoring and thus not covered by "dark taxonomy"

revisions. However, we would argue that the advantages of dark taxonomy outweigh these

disadvantages. Concentrating on fresh material from standardized traps means that specimens are

suitable for barcoding. Secondly, standardized samples come with complete metadata and there is

no need to digitize thousands of specimen labels individually. None of this can be said for

taxonomic revisions that deal with a mixture of wet and dry specimens. This does not mean that dry

material will become irrelevant. Dark taxonomy is designed to rapidly improve taxonomic

knowledge for a poorly known taxon to reach the level where conventional taxonomic approaches

can be applied. At this stage, pinned museum specimens will become critical for testing uncertain

species boundaries and estimating the full species-level diversity of a clade.

We also suspect that some systematists may be concerned about working at small geographic

scales. However, there are two reasons why we are less worried. Firstly, we would like to point to

the success of those turbo-taxonomic studies that seem to have found few cases, if any, where

species boundaries had to be revisited later when the fauna of neighboring areas was covered. The

best example is the weevil genus *Trigonopterus*, where several hundred species have been described

in regional treatments (Narakusumo et al. 2019, 2020; Riedel 2022; Riedel and Narakusumo 2019;

Riedel and Tänzler 2016; Riedel et al. 2013b, 2014; Van Dam et al. 2016). Of course, only time will

tell as to which proportion of species proposed based on poor geographic sampling will have to be

revised as geographic sampling increases, but it appears unlikely that the proportion will be high

when delimitation is based on molecular and morphological data. This prediction is based on the

current observation that barcodes alone tend to successfully delimit the majority of species. Note,

however, that the precise proportion remains unknown given that most species still lack barcodes

and comparatively few species have been sampled across their full geographic range. Denser

sampling tends to obscure the signal in barcode data as can also be seen in our study given that the

proportion of specimens in instable MOTUs increased from 530 in dataset 1 (36%) to 1,741 in the

combined dataset (59%). However, overall, we are optimistic that the vast majority of species

delimited with dark taxonomy will be stable, because the species boundaries are determined by

morphological and molecular data. So, both data would have to be misleading to generate incorrect

species limits.

Those systematists who are still reluctant to accept that dark taxa should be revised at small

geographic scales may want to consider the alternative. By describing more than 100 species based

on 1,454 specimens collected in a small country such as Singapore (730 km^2), we are increasing the

number of known Oriental species by >25%. This indicates the alarming magnitude of taxonomic

impediment and neglect, given that Singapore should have reduced diversity because it has lost

most of its natural habitats, and many species belonging to charismatic taxa are known to have gone

extinct (Chisholm et al. 2023; Theng et al. 2020). Expanding the geographic scale of a mycetophilid

revision to the Malay Peninsula would mean revising the fauna of an area that is 300 times larger

(242,363 km^2), with more pristine habitats and elevational gradients that are likely to increase the

species diversity. It would surely be impossible to handle the mycetophilid fauna at this scale using

conventional means. One may propose to tackle the fauna one genus at a time, but this would not be

feasible for three species-rich genera that contribute over 60% of the new species described in the

companion monograph (*Neoempheria* Osten-Sacken: N=31; *Epicypta* Winnertz: N=29; *Manota*

Williston: N=14).

Working at small geographic scales does requires regular combining of information from

multiple faunistic revisions covering neighboring areas. For example, our taxonomic revision of

Singapore's Mycetophilidae will become essential background information for a taxonomic revision

of the same group in additional studies on the fauna of the Malay Peninsula. Fortunately, combining

and analyzing barcode data from different studies is straightforward (Meier et al. 2021; Vences et

al. 2021) as long as the same barcoding region is sequenced. Such analysis then yields information

on the stability and intraspecific variability of each MOTU. This information furthermore guides an

expanded LIT analysis involving morphological and/or nuclear data. It is important that this follow-

up work should not require the loan of type material to avoid delays and the cost and risks of

specimen shipping. This is one major reason why the morphology of new species should be well

documented (Ang et al. 2013). But such documentation is also important for other reasons. It is the

only way to provide comparison data for species that were described based on morphology over the

last 250 years. This is almost all, given that even now only 10% of all new species descriptions in

entomology contain molecular data (Miralles et al. 2021). Integrative species descriptions are also

critical for the analysis of specimens not suitable for sequencing and for allowing researchers

without access to molecular labs to participate in science. This is a large number of biologists all

over the world (Zamani et al. 2022a,b). Neglecting morphology in species description thus leads to

the build-up of a new "superficial description impediment" (Meier et al. 2022) in taxonomy because

eventually morphological data will have to be collected anyway. Neglect of morphology also seems

very shortsighted, given that it appears likely that specimen identification with molecular tools will

eventually be complemented with identification based on image-based AI tools (Høye et al. 2021,

Wührl et al. 2022, 2024). These tools need to be trained with images and arguably the best images

come from taxonomic revisions because the specimens have been competently identified (Meier

and Dikow 2004).

<u>Establishing integrative species limits with LIT</u>

An important aspect of our study was testing the performance of Hartop et al.´s (2022) Large-Scale-

Integrative Taxonomy (LIT). We find that it performs well for Mycetophilidae. Although the initial

clustering of barcodes/specimens was at a lower threshold (3%) than was eventually found to

maximize congruence between barcode and molecular data, the number of specimens that had to be

studied with morphology was moderate (N=224). The proportion of specimens was higher than in

the previous application of LIT (Mycetophilidae: 15.4%; Phoridae: 5.1% in Hartop et al. 2022), but

this was largely due to the high species/specimen ratio of 8% for Mycetophilidae. Carrying out a

LIT analysis for all data (N=2,947) would reduce the proportion of specimens to be studied to 9%

(Table 2). In the case of Singapore´s Mycetophilidae, the level of congruence between MOTUs

used for specimen sorting (3% clusters) and morphological data was very high (110 congruent

MOTUs). This means that only seven morphospecies were in conflict with 3% MOTUs. Note also

that the term "conflict" is hardly appropriate from an evolutionary point of view. Measurements

such as the match ratio only quantify which proportion of morphospecies are in perfect agreement

(i.e., form identical sets) with MOTUs obtained using one particular clustering threshold or

algorithm. Yet, the use of a single threshold for clustering barcodes has long been criticized as

inappropriate on theoretical grounds (Will and Rubinoff 2004). Indeed, we would expect that

closely related species should form several proper subsets within a single MOTU if the latter was

delimited using a threshold appropriate for "average species". Conversely, we would expect old

species to form union sets composed of several MOTUs. These expectations are in line with the

results of a recent empirical study on optimal sequence similarity thresholds (e.g., Bonin et al.

2023). LIT therefore not only establishes whether morphospecies are "in conflict" with MOTUs,

but also tests whether there are clustering thresholds that yield congruent MOTUs. Only if this is

not the case, LIT would consider barcode and morphological data to be in genuine conflict and the

species would remain undescribed until additional data become available (e.g., nuclear markers,

ecology). However, it is comforting that we did not find such a case in our current study or our

recent study on Swedish phorids (Hartop et al. 2022).

Publishing dark taxonomy revisions

Many taxonomic monographs consist of two main sections. The introduction and discussion

sections contain information that is of interest to a broad readership (e.g., information on biology,

species diversity, taxonomic history of a taxon) while the taxonomic section consists of species

descriptions that are mostly of interest to taxonomic specialists. Unfortunately, the information that

is of general interest is often overlooked by biologists and thus also not cited. We here suggest that

a dark taxonomy revision should be published in two separate publications. The taxonomic

monograph could only consist of an introduction/method section that presents the background

information needed to understand how the species hypotheses were derived, as well as the

taxonomic descriptions. Everything else could be in a separate companion paper intended for a

broader audience. Such a companion manuscript can, for example, present a rigorous quantitative

analysis of the habitat preferences and phenology of species if the revision was based on fresh

samples obtained from standardized samples. We believe that it is desirable to bring both

manuscripts together in the form of a preprint and this approach is also pursued here. In our general

paper, we introduce the concept of dark taxonomy, document that a single, moderately sized

revision can increase the number of described species in a dark taxon by >25% for the Oriental

region, and present evidence that the species discovered in a first revision cover a very large

proportion of specimens collected at different times. In the companion monograph, we describe and

illustrate the species and discuss the taxonomic affinities of the newly described fauna (Amorim et

al. 2023).

Conclusions

Dark taxa are everywhere. They are abundant, species-rich, and a major obstacle to holistic

biomonitoring. What is needed is a robust "dark taxonomy" protocol designed to tackle the

taxonomic impediments for this dark diversity. We would argue that whatever protocol may

eventually be used should satisfy two criteria: it should be efficient enough to deal with thousands

of specimens and species and it should also yield biodiversity baseline data, so that changes in

abundance and richness of dark taxa can be monitored in a not-too-distant future and utilized for

ecosystem management or rehabilitation purposes. We believe that our proposal of "dark

taxonomy" can be a major step in this direction. Robots and cost-effective sequencing enable

presorting of species to putative species. LIT reduces the amount of morphological work needed to

obtain integrative species boundaries. Lastly, barcoding of type specimens will accelerate the

process of distinguishing between species that need a scientific name and those that are already

described.

Data availability statement

Data files and/or online-only appendices can be found in the Dryad data repository:

https://datadryad.org/stash/share/xFTVi0kZueBDEU4AXU-xaNeNa9o-y_jYvSRFlQdjt7s. The

custom script used for randomization of the data set partitions is available from

https://github.com/asrivathsan/darktaxa.

Acknowledgements

All work described here was carried out as part of a comprehensive insect survey of Singapore

which was carried out in collaboration and with support from the National Parks Board of

Singapore (NParks). Special thanks go to Dr. Patrick Grootaert and the team from the National

Biodiversity Centre of NParks for their assistance in fieldwork (Permits: NP/RP12-022-4,

NP/RP12-022-5, NP/RP12-022-6). We would also like to thank the research staff, lab technicians,

undergraduate students, and interns of the Evolutionary Biology Laboratory for their help and

assistance. This project would have been impossible without their hard work. Financial support was

provided by Tier 2 funding from the Ministry of Education (Singapore).

References

Ahrens D., Fujisawa T., Krammer H.J., Eberle J., Fabrizi S., Vogler, A.P. 2016. Rarity and

incomplete sampling in DNA-based species delimitation. *Syst. Biol.* 65(3):478-494.

Amorim D.S., dos Santos C.M.D. 2018. Flies, endemicity, and the Atlantic Forest: a

biogeographical study using topographic units of analysis. *Aust. Syst. Bot.* 30:439–469.

Amorim D.S., Oliveira S. S., Balbi M. I. P. A., Ang Y., Yeo D., Srivathsan R., Meier R. 2023. An

integrative taxonomic treatment of the Mycetophilidae (Diptera: Bibionomorpha) from

Singapore reveals 115 new species on 730km2. BioRxiv DOI:

10.1101/2023.09.02.555672.

Ang Y., Wong L.J., Meier R. 2013. Using seemingly unnecessary illustrations to improve the

diagnostic usefulness of descriptions in taxonomy–a case study on *Perochaeta orientalis*

(Diptera, Sepsidae). *ZooKeys* 355:9–27.

Bechev D. 1995. *Allactoneura papuensis* spec. nov. from New Guinea (Insecta: Diptera:

Mycetophilidae). *Reichenbachia* 31:81–82.

Bergsten J., Bilton D.T., Fujisawa T., Elliott M., Monaghan M.T., Balke M., Hendrich L., Geijer J.,

Herrmann J., Foster G.N., Ribera, I. 2012. The effect of geographical scale of sampling on

DNA barcoding. *Syst. Biol.* 61(5): 851–869.

Bickel D.J. 1999. What museum collections reveal about species accumulation, richness, and rarity:

an example from the Diptera. In: Ponder W., Lunney, D., editors. *The Other 99%: The*

Conservation and Biodiversity of Invertebrates. Royal Zoological Society of New South

Wales. p.174–181.

Bickford D., Lohman D.J., Sodhi N.S., Ng P.K.L., Meier R., Winker K., Ingram K.K., Das I. 2007.

Cryptic species as a window on diversity and conservation. *Trends Ecol. Ecol.* 22(3):148–

155.

Bonin A., Guerrieri A., Ficetola G.F. 2023. Optimal sequence similarity thresholds for clustering of

molecular operational taxonomic units in DNA metabarcoding studies. *Molecular Ecology*

Resources 23(2):368–381.

Bonnet X., Shine R., Lourdais O. 2002. Taxonomic chauvinism. *Trends Ecol. Ecol.* 17(1):1–3.

Borkent C.J., Wheeler T.A. 2012. Systematics and Phylogeny of *Leptomorphus* Curtis (Diptera:

Mycetophilidae). *Zootaxa* 3549:1–117.

Borkent A., Brown B.V., Adler P.H., Amorim D.S., Barber K., Bickel D., Boucher S., Brooks S.E.,

Burger J., Burington Z.L., Capellari R.S., Costa D.N.R., Cumming J.M., Curler G., Dick

C.W., Epler J.H., Fisher E., Gaimari S.D., Gelhaus J., Grimaldi D.A., Hash J., Hauser M.,

Hippa H., Ibáñez-Bernal S., Jaschhof M., Kameneva E.P., Kerr P.H., Korneyev V.,

Korytkowski C.A., Kung G.-A., Kvifte G.M., Lonsdale O., Marshall S.A., Mathis W.,

Michelsen V., Naglis S., Norrbom A.A., Paiero S., Pape T., Pereira-Colavite A., Pollet M.,

Rochefort S., Rung A., Runyon J.B., Savage J., Silva V.C., Sinclair B.J., Skevington J.H.,

Stireman J.O.III, Swann J., Vilkamaa P., Wheeler T., Withworth T., Wong M., Wood

D.M., Woodley N., Yau T., Zavortink T.J., Zumbado M.A. 2018. Remarkable fly (Diptera)

diversity in a patch of Costa Rican cloud forest: why inventory is a vital science. *Zootaxa*

4402(1):53–90.

Brown B.V. 2005. Malaise trap catches and the crisis in Neotropical dipterology. *Am. Entomol.*

51(3):180–183.

Brown B.V., Borkent A., Adler P.H., Amorim D.S., Barber K., Bickel D., Boucher S., Brooks S.E.,

Burger J., Burington Z.L., Capellari R.S., Costa D.N.R., Cumming J.M., Curler G., Dick

C.W., Epler J.H., Fisher E., Gaimari S.D., Gelhaus J., Grimaldi D.A., Hash J., Hauser M.,

Hippa H., Ibáñez-Bernal S., Jaschhof M., Kameneva E.P., Kerr P.H., Korneyev V.,

Korytkowski C.A., Kung G.-A., Kvifte G.M., Lonsdale O., Marshall S.A., Mathis W.,

Michelsen V., Naglis S., Norrbom A.A., Paiero S., Pape T., Pereira-Colavite A., Pollet M.,

Rochefort S., Rung A., Runyon J.B., Savage J., Silva V.C., Sinclair B.J., Skevington J.H.,

Stireman J.O.III., Swann J., Thompson F.C., Vilkamaa P., Wheeler T.A., Witworth T.,

Wong M., Wood D.M., Woodley N., Yau T., Zavortink T.J., Zumbado M.A. 2018.

Comprehensive inventory of true flies (Diptera) at a tropical site. *Commun. Biol.* 1(21): 1–

8.

Brydegaard M., Pedales R. D., Feng V., Yamoa A. S.-D., Kouakou B., Månefjord H., Wührl L.,

Pylatiuk C., Amorim D. D. S., Meier R. 2024. Towards global insect biomonitoring with

frugal methods. Phil. Trans. R. Soc. B 379: DOI: 10.1098/rstb.2023.0103.

Chisholm R.A., Kristensen N.P., Rheindt F.E., Chong K.Y., Ascher J.S., Lim K.K.P., Ng P.K.L.,

Yeo D.C.J., Meier R., Tan H.H., Giam X., Yeoh Y.S., Seah W.W., Berman L.M., Tan

H.Z., Sadanandan K.R., Theng M., Jusoh W.F.A., Jain A., Huertas B., Tan D.J.X., Ng

A.C.R., Teo A., Yiwen Z., Cho T.J.Y., Sin Y.C.K.2023. Two centuries of biodiversity

discovery and loss in Singapore. PNAS 120(51): e2309034120.

Dayrat B. 2005. Towards integrative taxonomy. *Biol. J. Linn. Soc.* 85(3):407–417.

De Carvalho M.R., Bockmann F.A., Amorim D.S., Brandão C.R.F., de Vivo M., de Figueiredo J.L.,

Britski H.A., de Pinna M.C.C., Menezes N.A., Marques F.P.L., Papavero N., Cancello

E.M., Crisci J.V., McEachran J.D., Schelly R.C., Lundberg J.G., Gill A.C., Britz R.,

Wheeler Q.D., Stiassny M.L.J., Parenti L.R., Page L.M., Wheeler W.C., Faivovich J., Vari

R.P., Grande L., Humphries C.J., DeSalle R., Ebach M.C., Nelson G.J. 2007. Taxonomic

Impediment or Impediment to Taxonomy? A Commentary on Systematics and the

Cybertaxonomic–Automation Paradigm. *Evol. Biol.* 34:140–143.

Colless D.H. 1966. Diptera: Mycetophilidae. *Insects of Micronesia, Bernice P. Bishop Museum*

12(6):637–666.

Dijkstra K.D.B., Kipping J., Mézière N. 2015. Sixty new dragonfly and damselfly species from

Africa (Odonata). *Odonatologica* 44(4):447–678.

Eberhard W.G. 1985. Sexual Selection and Animal Genitalia. Harvard University Press.

Edwards F.W. 1925. British fungus–gnats (Diptera, Mycetophilidae). With a revised generic

classification of the family. *Trans. R. Entomol. Soc. Lond.* 73:505–670.

Edwards F.W. 1926. XIV – Diptera Nematocera from the mountains of Borneo. Sarawak Museum

Journal III(10):243–278.

Edwards F.W. 1927. Diptera Nematocera from the Dutch East Indies. *Treubia* 9:352–370.

Edwards F.W. 1928. Diptera Nematocera from the Federated Malay States Museum. *Journal of the*
Federated Malay States Museums 14:1–139.

Edwards F.W. 1929. Philipinne Nematocerous Diptera III. *Notulae Entomologicae* 9: 70–81.

Edwards F.W. 1931. Fauna Sumatrensis. Mycetophilidae, Diptera. *Beijdrage* 69:262–278.

Edwards F.W. 1933. Diptera Nematocera from Mount Kinabalu. *Journal of the Federated Malay*
States Museums 17:223–296.

Edwards F.W. 1935. Two new *Delopsis* collected by Dr. Leefmans in Java. Diptera,
Mycetophilidae. *Tijdschrift Voor Entomologie* 78:186–187.

Edwards F.W. 1940. Neotropical *Neoempheria* (Diptera, Mycetophilidae). *Novitates Zoologicae*
42(1):107–129.

Evenhuis N.L. 2007. Helping solve the "other" taxonomic impediment: completing the eight steps
to total enlightenment and taxonomic Nirvana. *Zootaxa* 1407:3–12.

Fernandez-Triana J.L., Whitfield J., Rodriguez J., Smith M., Janzen D., Hajibabaei M., Burns J.,
Solis A., Brown J., Cardinal S., Goulet H., Hebert P. 2014. Review of *Apanteles* sensu
stricto (Hymenoptera, Braconidae, Microgastrinae) from Area de Conservación
Guanacaste, northwestern Costa Rica, with keys to all described species from
Mesoamerica. *ZooKeys* 383:1–565.

Fernandez-Triana J.L., 2022. Turbo taxonomy approaches: lessons from the past and
recommendations for the future based on the experience with Braconidae (Hymenoptera)
parasitoid wasps. *ZooKeys*, 1087:199.

Fitzgerald S.J. 2017. *Phthinia* Winnertz (Diptera: Mycetophilidae): new species and records
from the Neotropical and Oriental regions. *Zootaxa* 4231:107–113.

Geller J., Meyer C., Parker M., Hawk H. 2013. Redesign of PCR primers for mitochondrial
cytochrome c oxidase subunit I for marine invertebrates and application in all☐taxa biotic
surveys. *Mol. Ecol. Resour.* 13(5):851 –861.

Hartop E., Srivathsan A., Ronquist F., Meier, R. 2022. Towards large-scale integrative taxonomy

(LIT): resolving the data conundrum for dark taxa. *Syst. Biol.* 71(6):1404–1422.

Hausmann A., Krogmann L., Peters R., Rduch V., Schmidt S. 2020. GBOL III: Dark Taxa.

Available from https://ibol.org/barcodebulletin/research/gbol-iii-dark-taxa/.

Hebert P.D., Ratnasingham S., Zakharov E.V., Telfer A.C., Levesque-Beaudin V., Milton M.A.,

Pedersen S., Jannetta P., de Waard J.R. 2016. Counting animal species with DNA

barcodes: Canadian insects. *Philos. Trans. R. Soc. Lond. B* 371:20150333.

Hippa H. 2006. Diversity of *Manota* Williston (Diptera: Mycetophilidae) in a Malaysian rainforest:

description of twenty-seven new sympatric species. *Zootaxa* 1161:1–49.

Hippa H. 2007. The genus *Manota* Williston (Diptera: Mycetophilidae) in Melanesia and Oceania.

Zootaxa 1502:1–44.

Hippa H. 2008. New species and new records of *Manota* Williston (Diptera, Mycetophilidae) from

the Oriental region. *Zootaxa* 1723:1–41.

Hippa H. 2009. New species and new records of *Manota* Williston (Diptera, Mycetophilidae) from

Thailand. *Zootaxa* 2017:1–33.

Hippa H. 2011. New species and new records of *Manota* Williston (Diptera, Mycetophilidae) from

Thailand, with a key to the Oriental and Palaearctic species. *Zootaxa* 2763:39–60.

Hippa H., Jaschhof M., Vilkamaa. 2005. Phylogeny of the Manotinae, with a review of *Eumanota*

Edwards, *Paramanota* Tuomikoski and *Promanota* Tuomikoski (Diptera, Mycetophilidae).

Studia Dipterologica 11:405–428.

Hippa H., Kurina O. 2018. Four new species and new records of *Manota* (Diptera: Mycetophilidae)

from Sulawesi, Indonesia. *Acta Entomol. Mus. Natl. Pragae* 58:249–256.

Hippa H., Saigusa T. 2016. Notes on Oriental and East Palaearctic *Manota* Williston (Diptera,

Mycetophilidae), with the description of seven new species. *Zootaxa* 4084:377–390.

Hippa H., Ševčík J. 2010. Notes on Oriental and Australasian Manotinae (Diptera, Mycetophilidae),

with the description of thirteen new species. *Zootaxa* 2333:1–25.

Hippa H., Ševčík J. 2013. Five new species and a new record of *Manota* (Diptera: Mycetophilidae)

from Sulawesi. *Acta Entomol. Mus. Natl. Pragae* 53: 763–775.

Høye T.T., Ärje J., Bjerge K., Hansen O.L., Iosifidis A., Leese F., Mann H.M., Meissner K.,

Melvad C., Raitoharju, J. 2021. Deep learning and computer vision will transform

entomology. *Proc. Natl. Acad. Sci. U.S.A.* 118(2):e2002545117.

Huber F., Reis F.H. 2011. Técnica alternativa para montagem de insetos em lâminas permanentes

para visualizaçao em microscopia óptica. *EntomoBrasilis* 4:13–19.

Kallweit U. 1998. Notes on the genus *Metanepsia* Edwards and its relatives from East Asia

(Insecta: Diptera: Mycetophlidae). *Reichenbachia* 53: 341–353.

Kapli P., Lutteropp S., Zhang J., Kobert K., Pavlidis P., Stamatakis A., Flouri T. 2017. Multi-rate

Poisson tree processes for single-locus species delimitation under maximum likelihood and

Markov chain Monte Carlo. *Bioinformatics* 33(11):1630–1638.

Karlsson D., Forshage M., Holston K., Ronquist F. 2020. The data of the Swedish Malaise Trap

Project, a countrywide inventory of Sweden's insect fauna. *Biodivers. Data J.* 8:e56286

Kaspřák D., Borkent C.J., Wahab R.A. 2017. *Leptomorphus sevciki* sp. nov., a remarkable new

wasp-mimicking fungus gnat from Brunei (Diptera: Mycetophilidae). *Acta Entomol. Mus.*

Natl. Pragae 57(1):195–203.

Katoh K., Standley D.M. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Molec. Biol. Evol.* 30(4):772–780.

Kjærandsen J. 2022. Current State of DNA Barcoding of Sciaroidea (Diptera)—Highlighting the

Need to Build the Reference Library Insects 13(2): 147. DOI: 10.3390/insects13020147.

Kjærandsen J., Kerr P., Lindemann J.P., Kurina O. 2023. When details matter: Integrative revision

of Holarctic *Coelophthinia* Edwards (Diptera, Mycetophilidae), including mapping of its

mitogenome, leads to the description of four new pseudocryptic species. *Biodivers. Data J.*

11: e98741.

Kurina O., Hippa H. 2015. A review of the South Pacific *Manota* Williston (Diptera,

Mycetophilidae), with description of thirteen new species. *Zootaxa* 4020:257–288.

Kwong S., Srivathsan A., Vaidya G., Meier R. 2012. Is the COI barcoding gene involved in

speciation through intergenomic conflict? *Mol. Phylogenet. Evol.* 62:1009–1012.

Laamanen T.R., Petersen F.T., Meier, R. 2003. Kelp flies and species concepts–the case of *Coelopa*

frigida (Fabricius, 1805) and *C. nebularum* Aldrich, 1929 (Diptera: Coelopidae). *J. Zool.*

Syst. Evol. Res. 41(2):127–136.

Leray M., Yang J.Y., Meyer C.P., Mills S.C., Agudelo N., Ranwez V., Boehm J.T., Machida, R.J.

2013. A new versatile primer set targeting a short fragment of the mitochondrial COI

region for metabarcoding metazoan diversity: application for characterizing coral reef fish

gut contents. *Front. Zool.* 10(1):34.

Magnussen T., Soli G.E.E., Kjaerandsen J. 2019. *Allodia* Winnertz from the Himalayas, with nine

species new to science (Diptera, Mycetophilidae). *ZooKeys* 820:119–138.

Marsh P., Wild A., Whitfield J. 2013. The Doryctinae (Braconidae) of Costa Rica: genera and

species of the tribe Heterospilini. *ZooKeys* 347:1–474.

Matile L. 1999. Première citation du genre *Parempheriella* Matile dans la Région Paléarctique

(Diptera, Mycetophilidae). *Rev. Fr. Entomol.* 21(3):115–118.

Meier R. 2017. Citation of taxonomic publications: the why, when, what and what not. *Syst. Ent.*,

42(2):301–304.

Meier R., Dikow T. 2004. Significance of specimen databases from taxonomic revisions for

estimating and mapping the global species diversity of invertebrates and repatriating

reliable specimen data. *Conser. Biol.* 18(2):478–488.

Meier R., Kwong S., Vaidya G., Ng P.K.L. 2006. DNA barcoding and taxonomy in Diptera: a tale

of high intraspecific variability and low identification success. *Syst. Biol.* 55(5):715–728.

Meier R., Wong W., Srivathsan A., Foo M. 2016. $1 DNA barcodes for reconstructing complex

phenomes and finding rare species in specimen□rich samples. *Cladistics* 32(1):100-110.

Meier R., Blaimer B.B., Buenaventura E., Hartop E., von Rintelen T., Srivathsan A., Yeo D. 2022.

A re-analysis of the data in Sharkey et al.'s (2021) minimalist revision reveals that BINs do

not deserve names, but BOLD Systems needs a stronger commitment to open science.

Cladistics 38(2):264–275. https://doi.org/10.1111/cla.12489

Meier R., Hartop E., Pylatiuk C., Srivathsan A. 2024. Towards holistic insect monitoring: species

discovery, description, identification and traits for all insects. Phil. Trans. R. Soc. B 379:

20230120. DOI: 10.1098/rstb.2023.0120.

Miralles A., Bruy T., Wolcott K., Scherz M.D., Begerow D., Beszteri B., Bonkowski M., Felden J.,

Gemeinholzer B., Glaw F., Glöckner F.O. 2020. Repositories for taxonomic data: where

we are and what is missing. *Syst. Biol.* 69(6):1231–1253.

Montero☐Pau J., Gómez A., Muñoz J. 2008. Application of an inexpensive and high☐throughput

genomic DNA extraction method for the molecular ecology of zooplanktonic diapausing

eggs. *Limnol. Oceanogr.: Methods* 6(6):218–222.

Narakusumo R.P., Balke M., Riedel, A. 2019. Seven New Species of *Trigonopterus* Fauvel

(Coleoptera, Curculionidae) from the Tanimbar Archipelago. *ZooKeys* 888: 75 –93.

Narakusumo R.P., Riedel A., Pons J. 2020. Mitochondrial genomes of twelve species of

hyperdiverse *Trigonopterus* weevils. *PeerJ* 8:e10017.

Økland B. 1994. Mycetophilidae (Diptera), an insect group vulnerable to forestry practices? A

comparison of clearcut, managed and semi-natural spruce forests in southern Norway.

Biodivers. Conserv. 3(): 68 –85.

Økland B. 1996. Unlogged forests: important sites for preserving the diversity of mycetophilids

(Diptera: Sciaroidea). *Biol. Conserv.* 76:297–310.

Page R.D. 2011. Dark taxa: GenBank in a post-taxonomic world. Available from

https://iphylo.blogspot.com/2011/04/dark-taxa-genbank-in-post-taxonomic.html

Page R.D. 2016. DNA barcoding and taxonomy: dark taxa and dark texts. *Philos. Trans. R. Soc.*

Lond. B 371:20150334.

Papp L. 2004. Seven new species of Manotinae (Diptera: Mycetophilidae) from Asia and Papua

New Guinea. *Acta Zool. Acad. Sci. Hung*. 50:227–244.

Pentinsaari M., Salmela H., Mutanen M., Roslin T. 2016. Molecular evolution of a widely-adopted

taxonomic marker (COI) across the animal tree of life. *Sci. Rep.* 6:35275.

Phorid Catalog (http//phorid.net/pcat/flist3.php). Accessed August 2023.

Puillandre N., Lambert A., Brouillet S., Achaz G. 2012. ABGD, Automatic Barcode Gap Discovery

for primary species delimitation. *Molec. Ecol.* 21(8):1864–1877.

Ratnasingham S., Hebert, P.D.N. 2013. A DNA-based registry for all animal species: the Barcode

Index Number (BIN) system. *PloS One*, 8(7):e66213.

Riedel A. 2022. Nine new species of *Trigonopterus* Fauvel (Coleoptera, Curculionidae) from

Sundaland. *ZooKeys* 1124:109–130.

Riedel A., Sagata K., Suhardjono Y.R., Tänzler R., Balke, M. 2013a. Integrative taxonomy on the

fast track-towards more sustainability in biodiversity research. *Front. Zool.* 10:1–9.

Riedel A., Sagata K., Surbakti S., Tänzler R., Balke M. 2013b. One hundred and one new species of

Trigonopterus weevils from New Guinea. *ZooKeys* 280:1–150.

Riedel A., Narakusumo R.P. 2019. One hundred and three new species of *Trigonopterus* weevils

from Sulawesi. *ZooKeys* 828:1–153.

Riedel A., Tänzler, R. 2016. Revision of the Australian species of the weevil genus *Trigonopterus*

Fauvel. *ZooKeys* 556:97–162.

Riedel A., Tänzler R., Balke M., Rahmadi C., Suhardjono Y.R. 2014. Ninety-eight new species of

Trigonopterus weevils from Sundaland and the Lesser Sunda Islands. *ZooKeys* 467:1–162

Rohner P.T., Ang Y., Lei Z., Puniamoorthy N., Blanckenhorn W.U., Meier, R. 2014. Genetic data

confirm the species status of *Sepsis nigripes* Meigen (Diptera: Sepsidae) and adds one

species to the Alpine fauna while questioning the synonymy of *Sepsis helvetica* Munari.

Invertebr. Syst. 28(5):555–563.

Rowe L., Arnqvist G. 2012. Sexual selection and the evolution of genital shape and complexity in

water striders. *Evolution* 66(1):40 –54.

Santos D., Ribeiro G.C., Cabral A.D., Sperança M.A. 2018. A non-destructive enzymatic method to

extract DNA from arthropod specimens: Implications for morphological and molecular

studies. *PLoS One* 13(2):e0192200.

Santos B., Miller M., Miklasevskaja M., McKeown J., Redmond N., Coddington J., Bird J., Miller

S., Smith A., Brady S., Buffington M. 2023. Enhancing DNA barcode reference libraries

by harvesting terrestrial arthropods at the Smithsonian's National Museum of Natural

History. *Biodivers. Data J.* 11:e100904

Savage J., Borkent A., Brodo F, Cumming J.M., Curler G., Currie D.C., deWaard J.R., Gibson J.F.,

Hauser M., Laplante L., Lonsdale O., Marshall S.A., O'Hara J.E., Sinclair B.J., Skevington

J.H., 2019. Diptera of Canada. In: Langor DW, Sheffield C.S., Editors. The Biota of

Canada – A Biodiversity Assessment. Part 1: The Terrestrial Arthropods. *ZooKeys* 819:

397–450. https://doi.org/10.3897/zookeys.819.27625

Ševčík J. 2001. First record of *Cordyla* Meigen and *Monoclona* Mik from the Oriental Region

(Diptera: Mycetophilidae). *Beitr. Ent.* 51(1):155–160.

Ševčík J., Bechev D., Hippa H. 2011. New Oriental species of Gnoristinae with pectinate antennae

(Diptera: Mycetophilidae). *Acta Entomol. Mus. Natl. Pragae* 51(2): 687–696.

Ševčík J., Hippa H. 2010. New species of *Chalastonepsia* and *Pectinepsia* gen. nov. (Diptera:

Mycetophilidae) from the Oriental and Australasian Regions. *Acta Entomol. Mus. Natl.*

Pragae 50(2): 595–608.

Sevcik J., Kjærandsen J. 2012. *Brachyradia*, a new genus of the tribe Exechiini (Diptera:

Mycetophilidae) from the Oriental and Australasian regions. *The Raffles Bulletin of*

Zoology 60(1): 117–127.

Ševčík J., Kjærandsen J, Marshall S.A. 2012. Revision of *Speolepta* (Diptera: Mycetophilidae),

with descriptions of new Nearctic and Oriental species. *The Canadian Entomologist*

144(1): 93-107. DOI:10.4039/tce.2012.10.

Singapore National Parks Board. 2023. Species List (Red Data Book List).

https://www.nparks.gov.sg/biodiversity/wildlife-in-singapore/species-list. Accessed

August 2023.

Sivec I., Plassmann E. 1982. Sechs neue Pilzmücken aus Sri Lanka (Diptera, Nematocera,

Mycetophilidae). *Spixiana* 5(1):7–13.

Søli G.E.E. 1996. *Chalastonepsia orientalis* gen.n., sp. n., a second genus in the tribe Metanepsiini

(Diptera, Mycetophilidae). *Tijdschr. Entomol.* 139:79–83.

Søli G.E.E. 2002. New species of *Eumanota* Edwards, 1933 (Diptera: Mycetophilidae). *Ann. Soc.*

Entomol. Fr. 38:45–53.

Srivathsan A., Sha J.C., Vogler A.P., Meier R. 2015. Comparing the effectiveness of metagenomics

and metabarcoding for diet analysis of a leaf☐feeding monkey (*Pygathrix nemaeus*).

Molec. Ecol. Resourc. 15(2):250–261.

Srivathsan A., Hartop E., Puniamoorthy J., Lee W.T., Kutty S.N., Kurina O., Meier R. 2019. Rapid,

large-scale species discovery in hyperdiverse taxa using 1D MinION sequencing. *BMC*

Biol. 17:96. https://doi.org/10.1186/s12915-019-0706-9

Srivathsan A., Lee L., Katoh K., Hartop E., Kutty S.N., Wong J., Yeo D., Meier, R. 2021.

ONTbarcoder and MinION barcodes aid biodiversity discovery and identification by

everyone, for everyone. *BMC Biol*. 19:217.

Srivathsan A., Ang Y., Heraty J., Hwang W.S., Wan J.F.A., Kutty S.N., Puniamoorthy J., Yeo D.,

Roslin T., Meier, R. 2023. Convergence of dominance and neglect in flying insect

diversity. *Nat. Ecol. Evol.* 7:1012–1021. https://doi.org/10.1038/s41559-023-02066-0.

Srivathsan, A., Feng, F., Suarez, D., Emerson, B., Meier R. 2024. ONTbarcoder 2.0: rapid species

discovery and identification with real-time barcoding facilitated by Oxford Nanopore

R10.4. *Cladistics.* 40(2):192-203.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* 30(9):1312–1313.

Tan D.S., Ang Y., Lim G.S., Ismail M.R.B., Meier, R. 2010. From 'cryptic species' to integrative

taxonomy: an iterative process involving DNA sequences, morphology, and behaviour

leads to the resurrection of *Sepsis pyrrhosoma* (Sepsidae: Diptera). *Zool. Scr.* 39(1):51–61.

Theng M., Wan J.F.A, Jain A., Huertas B., Tan D.J., Tan H.Z., Kristensen N.P., Meier R.,

Chisholm R.A. 2020. A comprehensive assessment of diversity loss in a well-documented

tropical insect fauna: Almost half of Singapore's butterfly species extirpated in 160 years.

Biol. Conserv. 242:108401.

Troudet J., Grandcolas P., Blin A., Vignes-Lebbe R., Legendre F. 2017. Taxonomic bias in

biodiversity data and societal preferences. *Sci. Rep.* 7(1):1-14.

Van Dam M.H., Laufa R., Riedel A. 2016. Four new species of *Trigonopterus* Fauvel from the

island of New Britain (Coleoptera, Curculionidae). *ZooKeys* 582:129 –141.

Vences M., Miralles A., Brouillet S., Ducasse J., Fedosov A., Kharchev V., Kumari S., Patmanidis

S., Puillandre N., Scherz M.D., Kostadinov I., Renner S.S. 2021. iTaxoTools 0.1:

Kickstarting a specimen-based software toolkit for taxonomists. *Megataxa* 6:77–92.

Walker A.K., Crosby T.K. 1988. The preparation and curation of insects. DSIR, information series

163. Wellington, New Zealand: Science Information Publishing Centre.

Wang W.Y., Srivathsan A., Foo M., Yamane S.K., Meier, R. 2018. Sorting specimen□rich

invertebrate samples with cost□effective NGS barcodes: Validating a reverse workflow for

specimen processing. *Molec. Ecol. Resourc.* 18(3):490–501.

Wheeler Q.D., Raven P.H., Wilson E.O. 2004. Taxonomy: Impediment or Expedient? *Science*

303(5656):285. DOI: 10.1126/science.303.5656.285.

Will K.W., Rubinoff D. 2004. Myth of the molecule: DNA barcodes for species cannot replace

morphology for identification and classification. *Cladistics* 20(1):47–55.

Wong W.H., Tay Y.C., Puniamoorthy J., Balke M., Cranston P.S., Meier R. 2014. 'Direct PCR'

optimization yields a rapid, cost effective, non-destructive and efficient method for

obtaining DNA barcodes without DNA extraction. *Molec. Ecol. Resourc.* 14(6):1271–

1280.

World Economic Forum, 2023. The Global Risks Report 2023 (18th ed.).

https://www.zurich.com/knowledge/topics/global-risks/the-global-risks-report-2023

Wu H., Yang C. 1986. A new genus and a new species of Mycomyiinae from Fujian province,

China (Diptera, Mycetophilidae). *Acta Entomol. Sin.* 39(1):86–89.

Wu H., Zheng L.-H., Xu H.-C. 2003. First record of *Allodia* Winnertz (Diptera, Mycetophilidae) in

China, with description of seven new species. *Acta Zool. Sin.* 28(2):349–355.

Wührl L., Pylatiuk C., Giersch M., Lapp F., von Rintelen T., Balke M., Schmidt S., Cerretti P.,

Meier R. 2022. DiversityScanner: Robotic handling of small invertebrates with machine

learning methods. *Molec. Ecol. Resourc.* 22(4):1626–1638.

Wührl L., Rettenberger L., Meier R., Hartop E., Graf J., Pylatiuk C. Entomoscope: An Open-Source

Photomicroscope for Biodiversity Discovery. IEEE Access 12: 11785-11794.

https://doi:10.1109/ACCESS.2024.3355272

Xu H.-C., Wu H. 2002. Two new species of the genus *Azana* from China (Diptera:

Mycetophilidae). *Acta Zootax. Sin.* 27:621–623.

Yeo D., Puniamoorthy J., Ngiam R.W.J., Meier R. 2018. Towards holomorphology in entomology:

rapid and cost effective adult–larva matching using NGS barcodes. *Syst. Ent.* 43(4):678–

691.

Yeo D., Srivathsan A., Meier R. 2020. Longer is not always better: Optimizing barcode length for

large-scale species discovery and identification. *Syst. Biol.* 69:999–1015.

Yeo D., Srivathsan A., Puniamoorthy J., Foo M., Grootaert P., Chan L., Guénard B., Damken C.,

Wahab R.A., Ang Y., Meier R. 2021. Mangroves are an overlooked hotspot of insect

diversity despite low plant diversity. *BMC Biol.* 19(1):1–17.

Yu D.W., Ji Y., Emerson B.C., Wang X., Ye C., Yang C., Ding Z. 2012. Biodiversity soup:

metabarcoding of arthropods for rapid biodiversity assessment and biomonitoring. *Methods*

Ecol. Evol.. 3(4):613–623.

Zaitzev A.I. 1982. Composition and systematic position of the genus *Allactoneura* de Meijere

(Diptera, Mycetophilidae). *Entomologicheskoye Obozreniye* 60: 901–913. [In Russian]

Zamani A., Pos D.D., Fric Z.F., Orfinger A.B., Scherz M.D., Bartoňová A.S., Gante H.F. 2022a.

The future of zoological taxonomy is integrative, not minimalist. *Syst. Biodivers*. 20(1):1–

14.

Zamani A., Fric Z.F., Gante H.F., Hopkins T., Orfinger A.B., Scherz M.D., Bartoňová A.S., Pos

D.D. 2022b. DNA barcodes on their own are not enough to describe a species. *Syst. Ent.*

47(3):385–389.

Zhang J., Kapli P., Pavlidis P., Stamatakis A. 2013. A general species delimitation method with

applications to phylogenetic placements. *Bioinformatics* 29(22):2869–2876.