SIDERITE: Unveiling Hidden Siderophore Diversity in the Chemical Space Through Digital Exploration

Authors: Ruolin He[1#], Shaohua Gu[1,2#], Jiazheng Xu[3], Xuejian Li[4], Haoran Chen[4], Zhengying Shao[3], Haofan Wang[1], Jiqi Shao[1], Wen-Bing Yin[5,6], Long Qian[1*], Zhong Wei[3*], Zhiyuan Li[1,2*]

[1] Center for Quantitative Biology, Academy for Advanced Interdisciplinary Studies, Peking University, Beijing, 100871, China

[2] Peking-Tsinghua Center for Life Sciences, Academy for Advanced Interdisciplinary Studies, Peking University, Beijing, 100871, China

[3] Jiangsu Provincial Key Lab for Organic Solid Waste Utilization, Jiangsu Collaborative Innovation Center for Solid Organic Waste Resource Utilization, National Engineering Research Center for Organic-based Fertilizers, Nanjing Agricultural University, Nanjing, PR China

[4] Beijing Addinghome Technology Co.,Ltd, Beijing, 100193, China

[5] State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing, PR China

[6] Savaid Medical School, University of Chinese Academy of Sciences, Beijing, PR China

[#] These authors contributed equally to this article.

[*] corresponding authors (email: long.qian@pku.edu.cn; weizhong@njau.edu.cn; zhiyuanli@pku.edu.cn)

Abstract

Siderophores, a highly diverse family of secondary metabolites, play a crucial role in facilitating the acquisition of the essential iron. However, the current discovery of siderophore relies largely on manual approaches. In this work, we introduced SIDERTE, a digitized siderophore information database containing 822 siderophore records with 649 unique structures. Leveraging this digitalized dataset, we gained a systematic overview of siderophores by their clustering patterns in the chemical space. Building upon this, we developed a ligand-based method for predicting new iron-binding molecules. Applying this method to a commercial library, we experimentally confirmed that 40 out of the 48 molecules predicted as siderophore candidates possessed iron-binding abilities. Expanding our approach to the COCONUT natural product database, we predicted a staggering 3,199 siderophore candidates, showcasing remarkable structure diversity that are largely unexplored. Our study provides a valuable resource for accelerating the discovery of novel iron-binding molecules and advancing our understanding towards siderophores.

Introduction

Siderophore is a diverse family of secondary metabolites that exhibit high affinities for binding and chelating iron, one of the essential elements for cellular processes including replication and respiration[1,2]. However, iron presents a limiting resource due to its low solubility in most environments[3]. To overcome this challenge, microbes synthesize small molecule iron chelators, known as siderophores, which are able to bind poorly soluble iron and transport it into the cell via specific transporters[3]. The significance of siderophores lies in their vital role in ensuring microbial survival and growth. Pathways associated with siderophore synthesis and uptake are widely present in bacteria, fungi, and archaea[4], constituting complex ecological games[5]. Additionally, as a special type of natural product, siderophores exhibit notable antibacterial and antifungal activities, making them promising candidates for the development of novel therapeutics[6]. Despite their importance, our current understanding of siderophores is still limited due to their high diversity.

The extensive range of microorganisms' lifestyles plays a pivotal role in shaping the remarkable diversity of siderophores. According to a review in 2014, over 500 different types of siderophores have been identified, with 270 having been structurally characterized[7]. Despite this wealth of structural information, the number of experimentally characterized siderophore biosynthetic pathways remains considerably lower[3,8]. Siderophore biosynthesis predominantly falls into two major classes of pathways: the non-ribosomal peptide synthetase (NRPS) pathway and the NRPS-independent siderophore synthetase (NIS) pathway[3,9]. Notably, the NRPS pathway has been recognized for its capacity to yield a diverse range of natural products[4]. With such a high diversity, a systematic overview of siderophores, encompassing their sources within producing organisms, biosynthetic pathways, chemical properties, and a quantitative

assessment of their diversity within the known chemical space of natural products, is still
to be developed.

Previous researchers have been working towards a systematic overview of siderophores.
In 2010, Robert C. Hider and Xiaole Kong provided a valuable source of information on
the chemistry and biology of siderophores in a seminal review, which included structural
features of 294 siderophores in appendix[3]. A siderophore database, known as the
"siderophore base" (http://bertrandsamuel.free.fr/siderophore_base/index.php), was
created by Samuel Bertrand in 2011, which contained 262 siderophores with structure
images, organism sources, and references. However, the database was last updated in
2013 and is no longer maintained. Moreover, new siderophore molecules [10-13], and even
new siderophore ligand types (diazeniumdiolate and 2-nitrosophenol)[14-17], are constantly
being discovered in various microorganisms. Information regarding these siderophores is
currently dispersed across various publications and needs to be systematically recorded.

Another significant challenge in achieving a systematic overview of siderophores is the
lack of digitalization, which hinders computational investigations. In the field of natural
products, large digitized databases such as COCONUT[18], LOTUS[19], and SuperNatural[20]
record their molecules in SMILES (Simplified Molecular Input Line Entry System) format.
SMILES is the commonly used format for storing and analyzing chemical molecules,
which translates a chemical's three-dimensional structure into a string of symbols that are
easily readable by computer softwares[21]. This format enables large-scale computational
investigations such as machine learning[22]. However, there is no systematically curated
digital dataset about siderophores. Digitalized natural product databases do not offer
publicly accessible and searchable instances of "siderophore" and only contain a fraction
of currently known siderophores. Previous case-by-case research on siderophores
reported their siderophore discoveries by images of the structural formula[10-14], as do
previous reviews and the siderophore base. Although tools are available for converting
chemical structure images to SMILES, their accuracy is limited[23-25]. Furthermore, the
images in previous reviews and Samuel Bertrand's siderophore base are not standardized
and have some errors, further complicating the conversion[3]. Therefore, in addition to
curating the comprehensive information set of siderophores, digitalization also constitutes
a significant challenge.

With advances in siderophore research and computational biology, the time is opportune
for a quest for answers. For instance, how shall we quantify the chemical similarity
between siderophores to understand whether the iron scavenging systems are under
convergent or divergent evolution? And practically, can we identify new compounds with
iron-binding abilities from the vast repository of natural products? Given the crucial role of
iron in microbial metabolism[1], identifying such iron-binding compounds would facilitate
targeted interventions in microbial communities[6,26] and guide therapeutic applications
(iron chelation therapy)[27]. Suppose a novel family of iron-chelating siderophores that can
be utilized by beneficial but not pathogenic microbes. These chemicals can serve as
targeted prebiotics[5,28,29]. Furthermore, iron homeostasis plays a critical role in human
health and is implicated in various conditions ranging from acute iron poisoning to cancer
and aging[30-32]. Unfortunately, medicines capable of modulating iron homeostasis remain

scarce[33]. Therefore, can we draw inspiration from siderophores and investigate the
iron-binding potential of the vast array of currently available small molecules? To address
these questions and many more, a comprehensive quantification of the currently known
siderophores repository is necessary.

Taken together, establishing a comprehensive siderophore database is crucial for gaining
a deeper understanding of siderophore synthesis, function, and application. To fulfill this
need, we have developed the Siderophore Information Database (SIDERTIE), a
user-friendly platform that includes 872 siderophore records with 649 unique structures in
the SMILES format, covering all known siderophores up to May 2023. Leveraging
SIDERTIE's digitalization capabilities, we presented the most comprehensive statistics of
siderophores to date, covering biosynthetic pathways, source of producers, and several
chemical characterizations. The dispersed distribution of known siderophores within the
chemical landscape of natural products hints at the vast, largely uncharted territory of

undiscovered diversity. Building upon this quantitative overview, we proposed a

ligand-based method to batch discover new siderophores. A total of 48 siderophore
candidates were identified by this method, 40 of which were experimentally confirmed to
possess iron-chelating properties . The creation of the siderophore database and the swift
discovery of new siderophores hold profound implications for advancing
siderophore-related scientific research.

Result

1. Overview of SIDERITE

The Siderophore Information Database (SIDERITE, http://siderite.bdainformatics.org)
contains 872 records covering all known siderophores up to May 2023. In addition to
siderophore records from previous databases and reviews[3,9,34-36], 224 records were
curated from single research articles for the first time (Table S1). In addition to the
expanded collection, SIDERITE records the siderophore structures in the SMILES format,
a string-based digitization of chemical structures that can be easily processed by various
algorithms. The digitized siderophore structures provide a more convenient way of
presenting the information compared to previous databases and reviews, which primarily
used the picture format. Overall, SIDERITE represents a valuable resource and facilitates
systematic computational investigations. Notably, in comparison to other siderophore
collections, SIDERITE boasts the largest collection of siderophores and stands out for
being freely accessible and digitized (Table 1).
138
139
140
141

 Table 1

143   **Table 1.** Comparison of databases related with siderophores

Database	Number of overlaps with SIDERITE***	Total number of compounds	Active	Digitalization	Free	Siderophore-specific	Search siderophores
NPASS[37]	30	96,235	Y	Y	Y	N	N
LOTUS[19]	83	276,518	Y	Y	Y	N	N
SuperNatural 3[20]	151	484,846	Y	Y	Y	N	N
NPAtlas 2[38]	308	33,372	Y	Y	Y	N	N
COCONUT[18]	330	407,270	Y	Y	Y	N	N
Dictionary of Natural Products*	373	> 230,000	Y	Y	N	N	Y##
siderophore base**	210#	262	N	N	Y	Y	Y
Robert C. Hider's Review[3]	294	294	Y	N	Y	Y	Y
SIDERITE	649	649	Y	Y	Y	Y	Y
Except SIDERITE	425	-	-	-	-	-	-

*https://dnp.chemnetbase.com/

**http://bertrandsamuel.free.fr/siderophore_base/index.php

***Compare canonical SMILES without stereoisomerism.

#Although siderophore base lists 262 siderophores, only 210 are left after removing incorrect records and merging siderophores with same structures.

##Although DNP is not a siderophore-specific database, it can be indexed with the keyword siderophore to 421 entries including 373 siderophores that overlap with SIDERITE. But it contains siderophores without structural records and molecules that are not siderophores.

Digitizing siderophores enables computational analysis, particularly in unifying
siderophores based on their chemical structures. By comparing the canonical SMILES of
siderophores, we identified 649 unique siderophore structures out of the 872 total records
(Table S2). During this process, we observed that many siderophores share identical
structures but have different names, such as acyl-desferrioxamine 1 and legonoxamine A,
bacillibactin and corynebactin, streptobactin and griseobactin, and brucebactin and
lysochelin. This observation indicates that the same siderophores were discovered in
different species or by different research groups[39-46]. Therefore, for each unique
siderophore structure, we merged corresponding records and designated one of their
names as the official "Siderophore name", while recording the other names as
"Siderophore other name".

The 649 unique siderophores in our database can be classified by their producer sources
(Fig 1A). At the kingdom level, the majority of 649 siderophores in the database are
produced by bacteria (85.90%), followed by fungi (12.40%), plants (1.54%), and animals
(0.15%). While most siderophores are specific to one producer source at the kingdom
level, there is one exception. Triacetylfusarinine can be produced by both bacteria
(*Paenibacillus triticisoli*) and fungi (*Penicillium* sp.). At the phyla level, 872 producers of

649 siderophores are spread across five major bacterial phyla (229 in Actinobacteria, 7 in
Cyanobacteria, 33 in Firmicutes, 412 in Proteobacteria, and 8 in Bacteroidetes) and three
major fungal phyla (124 in Ascomycota, 38 in Basidiomycota and, 8 in Mucoromycota,
Table S1).

Siderophores can also be classified by their biosynthetic pathways (Fig 1B). Previous
studies have suspected that the NPRS pathway is more dominant for siderophore
synthesis[3], yet no statistically concrete ratio between NRPS and NIS-derived siderophore
has been established. In SIDERITE, we found that related pathways can be classified into
NRPS (65.18%), NIS (21.73%), Hybrid NRPS/PKS (10.02%), Hybrid NRPS/NIS (2.77%),
and PKS (0.31%). Consistent with previous studies, NRPS was indeed the most abundant
biosynthetic pathway of siderophores, followed by NIS. PKS siderophores are rare, with
only two cases (proferrorosamine A and tetracycline).   In different kingdoms, the
composition of the biosynthetic pathway differs (Table S2). In plants and animals, all
siderophores are synthesized by NIS. In fungi, almost all siderophores are synthesized by
NRPS (90.12%, 73/81). In bacteria, the diversity of the siderophore biosynthetic pathways
is higher, with 62.90% (351/558) NRPS, 22.58% NIS (126/558), 11.29% hybrid
NRPS/PKS (63/558), and 2.87% hybrid NRPS/NIS (16/558).

Digitization also enables us to easily access the statistical properties of siderophores,
such as ligand distribution. Siderophores chelate iron by several common ligands[3], and a
single siderophore may use multiple types of ligands. In SIDERITE, the top five
combinations of ligands are present in 64.25% of the siderophores (Fig 1C). These top
five combinations are Hydroxamate (28.35%), Alpha-Hydroxycarboxylate+Hydroxamate
(12.02%), Catecholate+Hydroxamate (8.78%), Hydroxamate+Hydroxyphenyloxazoline
(8.63%) and Catecholate (6.47%). Hydroxamate and Catecholate are the most common
ligands found in 69.49% and 29.74% of siderophores, respectively (Fig 1D). The
Octahedral geometry, which allows six ligands to be arranged around the iron center with
minimum repulsion, is the most common coordination geometry observed during
siderophore-iron chelation[3]. As expected, most siderophores have a denticity number of
six (Fig 1E). However, there are exceptions, such as pacifibactin, desferrioxamine T1,
malleobactin D and pyoverdine 7.7, which contain eight denticities.

The molecular weight (MW) is another important property of siderophores, as it influences
the diffusion of siderophores[47]. Siderophores are typically small molecules, and the MW of
siderophores in our database ranges from 138.12 Da (salicylic acid) to 1766.86 Da
(pyoverdine IB3). Most siderophores (90.60%, 588/649) have a middle MW ranging from
300~1100 Da (Fig 1F). Of note, half of the smallest 12 siderophores (MW < 200 Da) are
monomers of other siderophores, such as salicylic acid, 2,3-dihydroxybenzoic acid, and
citrate. Most of the large siderophores (MW > 1200 Da) are pyoverdines (87.76%, 43/49),
as pyoverdines are frequently composed of more than ten amino acids. Further, we check
whether MW distribution is related to biosynthetic types. We found NRPS siderophores
exhibit a wide range of MW, from 206.20 Da (Spoxazomicin D) to 1766.86 Da (pyoverdine
IB3). NRPS siderophores are generally heavier (mean: 835.68 Da; median: 816.00 Da)

than NIS siderophores (mean: 506.93 Da; median: 516.63 Da).

Additionally, aqueous solubility and the diffusion coefficient also are important ecological

properties of siderophores. However, most siderophores have not been measured

experimentally for these two properties. Therefore, we employed SolTranNet[48] and

Stokes–Einstein Gierer-Wirtz Estimation (SEGWE)[49] to predict these properties using

SMILES representations (Fig S1). Out of the total 649 siderophores, 635 were predicted

to be water-soluble (predicted logS>-6). Among them, 551 siderophores exhibited a "good"

level of aqueous solubility (predicted logS>-4). Notably, 14 siderophores exhibit poor

aqueous solubility (predicted logS≤-6) due to the presence of long-chain fatty acids,

rendering them insoluble in water. The minimum and maximum predicted diffusion

coefficients are 2.66×10^{-10} m^2/s and 7.7×10^{-10} m^2/s respectively. The majority of

siderophores (611 out of 649) exhibit predicted diffusion coefficients within the range of

2.66 to 5.50×10^{-10} m^2/s, consistent with previous research[50].

Figure 1

Figure 1. The statistics of 649 unique siderophores in SIDERITE

A. Distribution of the siderophore producers by their kingdoms.

B. Distribution of the siderophore biosynthetic pathways.

C. Distribution of the ligand type combinations. For clarity, only the top ten combinations are shown, and the others are merged into "Others".

D. Distribution of the common ligand of siderophores. One siderophore could contribute to more than one ligand type if it contains many types of ligands.

E. Distribution of denticity numbers.

F. Distribution of the molecular weight.

2. **Clustering of siderophores by their structural similarities**

Siderophores have been known to exhibit remarkable structural diversity[3] (Fig S2). Converting siderophores into SMILES format enables us to quantify their chemical similarity more effectively, both within the SIDERITE database and between other natural products. To systematically assess the structural diversity of siderophores, we first locate all 649 SIDERITE structures to the vast chemical space of the COCONUT database by merging all molecules in these two databases, which encompasses over 4×10^5 natural products. By Tree MAP (TMAP) visualization of chemical similarity (Fig S3), we observed that the 649 siderophores could be grouped into 25 distinct clusters, which were separated from each other by natural products from the COCONUT. The clustering result shows siderophores have unevenly distributed structural diversity (Fig 2, Table S2). Most of these clusters (16 out of 25) only contain a few members (<5), while the largest four clusters account for 89.37% of the siderophore structures in total. We sorted the cluster indexes by their member counts in descending order; for instance, cluster 1 contains the most siderophore structures.

Within each cluster, there are common features of ligands or biosynthetic types. Cluster 1 (201, 30.97%) includes siderophores with benzene ring structures in the ligands such as catecholate, phenolate, hydroxyphenyloxazoline, and hydroxyphenylthiazoline. Siderophores in the cluster 1 are synthesized by both NRPS and NIS. Cluster 2 (197, 30.35%) only includes siderophores produced by NRPS pathways except Albomycins by the hybrid NRPS/NIS pathway. Albomycins are naturally occurring sideromycins (siderophore–antibiotic conjugates) produced by some streptomycetes, and their siderophore parts are synthesized by NRPS. From the perspective of ligands, most siderophores in the cluster 2 contain hydroxamate of (92.39%, 182/197), and many also contain alpha-hydroxycarboxylate (37.06%, 73/197). Cluster 3 (103, 15.87%) is all NIS siderophores. Like cluster 1, most of them contain hydroxamate (90.29%, 93/103), and many also contain alpha-hydroxycarboxylate (33.01%, 34/103). The sources of alpha-hydroxycarboxylate are mostly citrate. Cluster 4 (79, 12.17%) is NRPS siderophores with chromophores such as pyoverdines (93.67%, 74/79). Other small clusters all are located on the edge of four large clusters (Fig 2A). They consist of similar ligand composition (Fig S4-S16), which indicates the possibility of unusual siderophores evolving from common siderophores.

Molecule weight (MW), carbon-nitrogen ratio (C/N), and carbon-oxygen ratio (C/O) of siderophores pertain to their biosynthetic cost in various aspects: higher MW can indicate more building blocks; while producing products with lower C/N or C/O could influence growth under nitrogen-limited or hypoxic environments[51-53]. Therefore, we investigated these properties of siderophores in different clusters (Fig S17). Siderophores in cluster 4 have significantly higher MW than average, as most cluster members are pyoverdines.

275 Regarding C/N and C/O ratios, different clusters also exhibit different properties. For
276 example, siderophores in cluster 7 and cluster 10 have significantly higher-than-average
277 C/N because they only contain 1~2 nitrogen atoms, and siderophores in cluster 4 have a
278 significantly lower-than-average C/N (Fig S18).
279
Large clusters only provide an initial classification relative to other natural product
molecules. Further, we defined small groups within each of the 25 clusters by their
structure similarity coefficient (Dice coefficient, threshold 0.6, see method), and obtained
102 small groups in total. Each group is named by its cluster ID x and its group ID y within
this cluster. Accordingly, each siderophore is assigned a unique id $x.y.z$, where z stands
for the z-th record within the group. For example, enterobactin has the id 1.22.2. In this ID,
"1" means the 1^{st} large cluster. "22" means the 22^{nd} group within the 1^{st} large cluster. "2"
means the 2^{nd} record in the 1.22 cluster. In the future, newly discovered siderophores will
also be assigned a unique id when being incorporated into the SIDERITE database.
289
This cluster-group naming method provides an intuitive framework for comprehending the
siderophore clusters in the chemical space. The distribution of structural diversity of
siderophores in the chemical space is uneven. The three largest clusters contribute to
70.59% (72/102) of the groups, while the average member number of the remaining
groups is 1.36 (Table S2). All small groups encompass siderophores derived from unique
biosynthetic types, except groups 1.5 and 1.10, where siderophores are synthesized
through both hybrid NRPS/PKS and NRPS pathways. The siderophores in these two
groups have two biosynthetic types due to the optional modification from PKS. Also,
siderophores within the same group are typically produced by species from the same
kingdom. However, groups 2.1, 2.9, 3.7, and 9.1 contain producers from both bacteria and
fungi, and group (11.1) is produced by bacteria and animals.
301
For siderophores with unknown biosynthetic types, their biosynthetic types could be
inferred by other members with known biosynthetic types in the same small cluster
because almost all members in the same small cluster have the same biosynthetic types
(except for 1.5 and 1.10 clusters). It's useful for mining their biosynthetic genes in the
genome after discovering new siderophores, which would accelerate siderophore
research from structures to genes.
308

Figure 2

Figure 2. The visualization of large siderophore clusters in SIDERITE.

A. A network of siderophores connected by chemical similarities. Each node in the
network corresponds to a siderophore molecule, and the nodes are linked to their
most similar neighbors, forming a minimum spanning tree[54] (described in the method

section). Nodes are colored by their cluster IDs.

B. For the four largest clusters, example structures are provided, and the ligands of the

siderophores are colored according to their types. Siderophores are circled by

rounded rectangles to show their cluster IDs (same color codes as (**A**)).

3. Discovering new siderophores by the ligand-based method

The clustering analysis of known siderophores has unveiled the presence of common

ligands among these compounds. Drawing inspiration from this observation, we have

introduced an innovative rule-based approach, aimed at the discovery of novel

siderophores by their chemical structures (Figure 3A-3B). In this approach, we first

distilled 15 common ligands derived from the characteristics of known siderophores. Any

molecule containing at least one of these 15 ligands is identified as a potential

siderophore (Figure 3C and Table S3). Then, we exclude candidates containing any of the

8 modified siderophore ligands incapable of forming coordination bonds to chelate iron

(Figure 3D and Table S4). Besides, while the alpha-hydroxycarboxylate ligand is prevalent

within siderophores (Figure 3E), we have excluded it from our rule set due to its ubiquity in

non-siderophore molecules.

To verify this method, we used a commercial natural product library (the Natural Product

Library for high throughput screening, catalog number L6000, TargetMol, Shanghai, China,

June 2023) as a test set. After excluding six known siderophores, this dataset consisted a

total of 4,308 molecules (Figure 3B). Our ligand-based method identified 48 molecules as

potential siderophores, ranked by their highest Tanimoto similarity to known

siderophores[55] (Table S5). We checked the producer sources of these 48 molecules and

found that, except for chlorquinaldol (compound **29**) which is synthetic[56], the other 47 are

natural products derived from plant, insects, microbes, algae, or humans (Table S5).

Among these molecules, 22 are soluble in water, while the remaining 26 have poor

solubility in water (Table S5). To address this, we dissolved the poorly soluble molecules

by dimethyl sulfoxide (DMSO) instead of water. Subsequently, solutions of these 48

molecules were tested by chrome azurol S (CAS) assay, a universal colorimetric method

that detects iron-binding molecules[57].

The high positive rate from the CAS assay supports the effectiveness of our ligand-based

method. Among the tested molecules, 20 out of 22 (90.9%) water-soluble compounds and

20 out of 26 (76.9%) water-insoluble compounds exhibited iron-binding activity, as

evidenced by a noticeable change in the color of the CAS dye (Figure 3F-3G and Table

S6). Actually, most molecules with negative CAS results (compound **20**, **24**, **31**, **33**, **37**, **41,**

and **44**) exhibited unusual color patterns, which hinders accurate assessment. For

instance, their original solutions were significantly dark in color, or reacted with CAS

reagents and formed precipitates or turbidity. Actually, compound **24**, **37**, **41**, and **44** did

induce color change in the CAS assay solution, but their precipitation interfered with the

357 optical density measurement. Taken together, only one (compound **13**) out of the 48
358 molecules was confirmed to lack iron-chelating ability.

360 Of these 40 molecules, which we confirmed to exhibit iron-chelating capabilities (referred
361 to as 'potential siderophores' henceforth), the majority remains largely unexplored
362 regarding their iron-binding potential. Remarkably, only three (compounds **5**, **10**, and **48**,
363 see Figure 3H) with their seven derivatives have been previously associated with
364 iron-binding related research, but have not yet been regarded as siderophores: 1. Gallic
365 acid (compound **5**) has been systematically studied for complexation with iron[58] and has
366 been commonly used in the pharmaceutical industry. In these 40 molecules, there are six
367 gallic acid derivates (compounds **12**, **14**, **17**, **23**, **25**, and **27**) with iron-binding activity. 2.
368 Xanthurenic acid (compound **10**) is the precursor of *Pseudomonas* siderophore
369 quinolobactin[59]. 3. Gossypol (compound **48**) has been reported as dietary
370 supplementation with interactions with ferrous sulfate[60], implying iron-binding activity.
371 Gossypol derivate gossypol acetic acid (compound **47**) also shows iron-binding activity in
372 our assay. Taken collectively, at least 30 of the molecules we identified may represent
373 novel siderophores.

375 We then checked the chemical properties of these 40 potential siderophores. Out of the
376 40 molecules, the most frequent ligand type is catecholate (36/40). Among these, the
377 majority (19/36) are trihydroxy, such as gallic acid (compound 5, Figure 3H). The
378 remaining four molecules not possessing catecholate contain 8-hydroxyquinoline, which is
379 an uncommon ligand. It is worth noting that while it is known that 8-hydroxyquinoline can
380 chelate metal, only two siderophores (quinolobactin and thioquinolobactin) containing this
381 ligand have been reported so far[61,62]. Therefore, our discovery of the four molecules
382 significantly expands the list of 8-hydroxyquinoline siderophores. The four molecules are
383 robustine (compound **9**), xanthurenic acid (compound **10**, Figure 3H), jineol (compound
384 **16**, Figure 3H), and chlorquinaldol (compound **29**). Chlorquinaldol is a synthetic
385 compound. Robustine is produced by plant *Zanthoxylum integrifoliolum*[63] and
386 *Zanthoxylum avicennae*[64]. Jineol is produced by the insect *Scolopendra subspinipes*[65].
387 The biosynthetic pathways of robustine and jineol are currently unknown.

389 Regarding the functions of the 40 potential siderophores, most of them (excluding 14
390 molecules with unknown functions) are known to have antioxidant activities (14/40). This
391 percentage is significantly higher than the proportion of molecules with "antioxidant"
392 annotations in the commercial library (5.1%). The second most common function are
393 anticancer (7/40), followed by anti-inflammatory (4/40, Table S5). The high occurrence of
394 these pharmaceutical functions in potential siderophores motivates us to explore of the
395 large natural product repository for additional molecules with iron-binding capabilities.

397 Subsequently, we applied our ligand-based method to the large chemical database
398 COCONUT, excluding known siderophores from consideration. Remarkably, this analysis
399 led to the identification of 3,199 molecules exhibiting potential iron-binding activities from
400 a database containing over 0.4 million compounds (Table 1 and Table S7). By the

Tanimoto similarity clustering (Figure 3I), a large proportion of these potentially
iron-binding molecules is scattered in the chemical space, not close to any of the known
siderophores. Specifically, only 284 out of the 3,199 molecules exhibit significant
structural similarity (with a maximum Tanimoto similarity of > 60%) to the known 649
siderophores cataloged in SIDERITE. The remaining 2915 molecules are strong
candidates for novel siderophores with relatively unexplored chemical structures. This
analysis underscores the notion that the structural diversity of siderophores remains
largely concealed, inviting further in-depth exploration and investigation.

Figure 3

Figure 3. The rule-based siderophore discovery approach and the result of CAS test
experiments
A. Principle of rule-based siderophore discovery approach. The rules are summarized
from known siderophores based on the SIDERITE database. The predictions were
then tested by experiments.
B. Pipeline of the ligand-based siderophore discovery. Molecules containing at least one
of 15 ligands in (**C**) and not any of the 8 modified ligands in (**D**) are selected as
siderophore candidates.
C. The structures of siderophore ligands in the rules in (**B**). The rarity of ligands is noted
by the different background colors.
D. The structures of modified ligands in the rules in (**B**). Modifications that cause ligands
to lose iron-chelating abilities are marked in red.
E. The structure of common ligand alpha-hydroxycarboxylate.
F. The CAS result of the 25 water-soluble siderophore candidates. The solvent is water.

The mean and ± one standard deviation of OD (630nm) are shown in the bar graph.

The OD (630nm) of solvent is marked as a red line (mean value) and red dash line
(one standard deviation). The compounds with negative results in the CAS test are
marked in red.
G. The CAS result of 30 water-insoluble siderophore candidates. The solvent is DMSO.
Other legends are the same as (**F**).
H. The representative examples of compounds with iron-binding activities.
I. The structural diversity of new potential siderophores in the COCONUT database.
3,199 molecules with potential iron-binding activities and 649 known siderophores
were clustered with Tanimoto similarity. The source of molecules (COCONUT or
SIDERITE) is shown in the left bar by yellow and black colors, respectively.
436

4. The usage of SIDERITE database

438 SIDERITE is freely accessible at http://siderite.bdainformatics.org. The current version
439 serves as a platform for searching and displaying known siderophore chemical structures.
440 Users can access desired siderophores by the following three approaches (Fig 4A).
441

An indexing interface is provided so that users can quickly access the desired
siderophores. We visualize the SIDERITE database by TMAP[54], and label multiple
properties of siderophores (Figure 2A and Figure 4B,
http://siderite.bdainformatics.org/sidertmap). In the mapping, each siderophore is
represented as a distinct point that offers more detailed information when clicked.
Additionally, users can jump to the siderophore page by the interactive siderophore ID in
the information card (Figure 4B).
449

Users also can use the search function to access desired siderophores (Figure 4C-4D).
The platform offers fast fuzzy search functionality based on partial siderophore names, as
well as precise advanced search by multiple conditions. The advanced search feature

includes various conditions, including siderophore ID, siderophore name, siderophore
ligand, biosynthetic type, monomer name, number of monomers, monomer coverage,
theoretical denticity, molecular weight, microorganism, source (kingdom), source (phylum),
and class. Numeric conditions, such as the number of monomers and molecular weight,
can also be set within specified ranges.
458
Chemical similarity search is the third way to access the siderophore resource in
SIDERITE (Figure 4E). For any query molecule, this function helps users to find
structurally similar siderophores, and also predict whether this query molecule has
iron-binding activity. Users can input the SMILES notation of the query molecule or draw it
in the SMILES editor. If the molecule has been predicted to have potential iron-binding
activity by our ligand-based method (Figure 3) and is not already present in SIDERITE,
the SIDERITE platform will automatically indicate to the user that this is a potentially new
siderophore. Users can choose whether or not to upload this molecule to the SIDERITE
database without any obligations. Furthermore, this search will return siderophores in
SIDERITE that are chemically similar to the query molecule. The chemical similarity
metric supports Tanimoto, Dice, and Tversky, and the user can customize the cut-off
threshold.
471
472   We provide individual pages with detailed information for each siderophore structure (Fig
473   4F), which provide canonical SMILES, pictures, and related references. Alternative names
474   will be noted after the main name, separated by a forward slash. In the "Siderophore
475   Ligand" section, the number of specific ligands is specified within parentheses.

476 **Figure 4**

Figure 4. The SIDERITE database

A. Workflow of SIDERITE database usage.

B. Accessing siderophores by the indexing interface by the following steps: 1. Choose a type of siderophore legend. 2. Click the interested siderophore. 3. Jump to the siderophore page by the interactive siderophore ID in the information card.

C. Fast fuzzy siderophore search.

D. Advanced siderophore search by precise conditions.

E. Chemical similarity search, where users can type the SMILES string or draw in the SMILES editor.

F. The individual page with detailed information on each siderophore structure.

Discussion

In this work, we have constructed the most comprehensive siderophore database to date (http://siderite.bdainformatics.org/), with digitalized formats. Over the past few decades, a large number of siderophore structures have been identified. Some of them have been collected in Robert C. Hider and Xiaole Kong's review in 2010[3] and Samuel Bertrand's unmaintained siderophore database in 2011 (http://bertrandsamuel.free.fr/siderophore_base/index.php). As the field of siderophore research continues to advance, there is a growing need to establish a systematic and standardized classification system for siderophores. While previous dissemination of siderophore knowledge through pictorial formats has undoubtedly facilitated its spread and communication, these formats have not adequately supported systematic computational studies. Based on our digitalized dataset, we were able to obtain a systematic overview of siderophores, and derived an effective ligand-based method for discovering new iron-binding molecules. Our approaches have paved the way for more data-driven discoveries in the field.

Our work provided a statistical overview on siderophores, from the perspectives of biosynthetic type, producer source, and chemical properties. Particularly, the clustering patterns of siderophores in the chemical space hold implications for their eco-evolutionary histories[5], and the drives of their chemical diversity. Siderophores are considered public goods as they are predominantly secretive[66]. However, some species have evolved the ability to exploit siderophores without producing them, by synthesizing only the receptors for these siderophores[5,66]. To avoid this exploitation, producers can modify the structure of their siderophores[5,67,68]. This evolutionary pressure under such a Red Queen situation may generate large siderophore clusters with similar but diverse structures. For example, researchers recently found *Streptomyces venezuelae* produced the uncommon, alternative siderophore foroxymithine (cluster 2.7) to impose a more profound fitness when their main siderophore desferrioxamine (cluster 3.3) was exploited by the yeast *Saccharomyces cerevisiae* (cheater) in the co-culture[69]. Meanwhile, siderophores in clusters with few similar members may indicate other strategies, like utilizing multiple structurally distinctive siderophores to avoid intense competition. Across different

kingdoms, bacteria exhibit the most diverse synthetic types and the highest number of
siderophores, reflecting the intense iron competition observed among prokaryotes[5,70].

Dealing with such a high diversity, one advantages of digitalization is to facilitate the
annotation of previously unnoticed siderophores. During our construction of SIDERITE, 50
siderophore records were found by exploring chemically similar natural product molecules
around known siderophores on the COCONUT database mapping. Among them, 29
records are from references that do not contain "siderophore" in the title, making their
detection through literature searches more challenging. In fact, half of them were initially
reported as antibiotics or microbial inhibitors. Also, categorizing siderophores by the
SMILES format avoid rediscovering of the same structures from different organisms yet
annotating them as "new" siderophores[39-46].

Another advantage of digitalization is to provide resources for data-driven explorations,
such as in bio-retrosynthesis prediction. In recent years, deep learning approaches have
been employed for predicting the biosynthetic pathways of natural products[71]. In addition,
high-quality labeled siderophore data can be used to train supervised learning models,
leading to the design of novel artificial siderophores with enhanced biological activities.

In fact, our ligand-based method for searching iron-chelating structures serves as a
preliminary prototype for data-driven discoveries. Although simple in nature, it can be
seen as analogous to supervised learning, where features related to iron-binding (such as
rules for including or excluding certain ligands) are extracted from the existing SIDERITE
dataset and applied to chemicals with unknown iron-chelating abilities. With the high
successful rate validated by our CAS experiments, this ligand-based method is likely
applicable to the vast repository of natural products. In the COCONUT database, our
method predicted 3,199 molecules as iron-binding compounds. Particularly, siderophore
clusters only containing few members can suggest underexplored siderophore types and
highlights for future research. For example, pyridoxatin has only one member in the
cluster 18 of SIDERITE. Cordypyridone A and B with antimalarial activity are the nearest
natural products to pyridoxatin in the chemical space of COCONUT, and both compounds
were predicted as "iron-binding" by our method. However, cordypyridone A and B were not
previously considered to be siderophores[72], indicating underexplored siderophore types
that call for verification by subsequent experiments.

More inquiringly, most of the 3,199 structures are different from known siderophores,
suggesting that the true extent of siderophore diversity remains largely unexplored. Our
findings also imply a higher prevalence of iron-binding molecules in plants than
anticipated. These plant siderophores, known as phytosiderophores, are predominantly
produced by graminaceous plants[3]. Remarkably, our data demonstrates that 92.5% (37
out of 40) of our identified iron-binding molecules originate from plants (Table S5),
revealing substantial untapped potential for phytosiderophores.

Developing new methods to quickly discover new siderophores by systematically
summarizing the ligand diversity and structural features of siderophores holds immense
scientific and practical importance. Primarily, uncovering novel siderophores can open
avenues for innovative therapeutic strategies. Iron's critical role in various diseases,
including cancer and inflammation, underscores its significance[73-75]. It is known
siderophores have an antioxidant role by chelating iron because iron can cause reactive
oxygen species by reacting with H_2O_2[76-78]. And anticancer and anti-inflammatory activities
of siderophores are achieved by sequestering iron in the environment to limit the growth of
cancer cells[73] and microbes[74,75]. More than half of our newly identified iron-chelating
molecules have reported antioxidant, anticancer, and anti-inflammatory properties. These
molecules may also be used as environmental remediation of heavy metal pollution[79,80] .
575
576
Different from previous siderophore resources, which remain limited to single studies and
therefore do not provide a mechanism to perform cross-study and systematic analyses,
SIDERITE provides a curated community-wide platform. All siderophores are digitized,
which makes comparisons easy and thus avoids duplicate naming of the same molecule.
Each siderophore is assigned a three-level serial unique id based on the siderophore
structure similarity feature. The unique id assigned to each siderophore in this study also
provides a standardized nomenclature for future studies. With the advent of SIDERITE,
newly discovered siderophores can be easily identified and quickly known by other
researchers in this field. We invite all relevant researchers to join this community and
collaborate to promote the prosperity of the siderophore community.
587

Method

Siderophore information resource

A total of 872 siderophore information records were collected from various sources, including databases, reviews, and research articles. Among these, 355 records were obtained from Samuel Bertrand's Siderophore Base, while the remaining records were sourced from other databases, reviews, and research articles (Table S1). Specifically, we obtained 160 new records from the appendix of Robert C. Hider and Xiaole Kong's review, 37 new records from other reviews, 95 new records from the Dictionary of Natural Products (DNP) database (https://dnp.chemnetbase.com/), and 224 new records from research articles by searching with keywords such as "new siderophore," "novel siderophore," and "siderophore discovery". It is important to note that one new record which is a pyoverdine (a type of siderophore) was obtained from the LOTUS database with the keywords "pyoverdine". This was because this database only allowed searching siderophores by their specific names such as "pyoverdine", rather than by the general biological function such as "siderophore".

Each siderophore information record in our dataset comprises various details, including the siderophore name, the synonym name (if any), the type and number of ligands, the producing species, the biosynthetic type, and the structure (Table S1).

Biosynthetic type annotation

Information on 721 siderophore biosynthetic types is available in our database, which we documented based on references. For the remaining 151 siderophores with unknown biosynthetic types, we inferred their biosynthetic types by features of different types. The biosynthetic type of the siderophore would be annotated as "Putative NIS", if it contains monomers that only exist in the NIS siderophore such as citrate, diamine (e.g. 1,3-Diaminopropane, putrescine and cadaverine) or diamine derivatives (e.g. N-hydroxy cadaverine). The biosynthetic type of the siderophore would be annotated as "Putative NRPS" if the monomers in this siderophore all are amino acids.

SMILES conversion from siderophores

The Simplified Molecular-Input Line-Entry System (SMILES) is a notation to describe the chemical structure of molecules using a string format which is particularly useful for subsequent processing by computer. However, in our siderophore information resource, the SMILES format of siderophore structures and their annotations were only available in

the DNP database. In contrast, the literature and Siderophore Base only included
siderophore structures in the picture format which is more intuitive for readers but poses
difficulties for computational analysis. To address this limitation, we developed a
customized Python script that utilized the ChemSpider API and Chemical Identifier
Resolver (CIR) to convert siderophore names into the SMILES format. The SMILES
format structures of 34.93% (124/355) siderophore records in the Siderophore Base were
retrieved. For the remaining 748 siderophores that were not found or from other resources,
we manually drew their structures and obtained the SMILES format structures by SMILES
generator/checker   tool
(http://www.cheminfo.org/flavor/malaria/Utilities/SMILES_generator___checker/index.htm
l).
635
636   To ensure consistency and facilitate comparison, we then converted all SMILES strings
637   into canonical SMILES strings using another custom Python script. The RDKit package
638   was utilized for this conversion process, while simultaneously obtaining the molecular
639   formula and molecular weight of the siderophores.
640

Prediction of aqueous solubility and the diffusion coefficient for siderophores based on SMILES

643   To predict aqueous solubility, we used a machine learning tool SolTranNet with SMILES of
644   siderophores as input[48]. For the prediction of the diffusion coefficient, we used the
645   SEGWE calculator developed by Evans, R. *et al.*[49] with a temperature 298.15K and water
646   as solvent.
647

Visualization of SMILES

To compare and analyze siderophores intuitively, we visualize SIDERITE with other
chemical databases by TMAP[54]. The visualization mapping of SIDERITE with other
chemical databases consists of the vast chemical spaces for exploring new siderophores.
The TMAP mapping algorithm tends to produce a significant distance between a molecule
with and without specification of stereoisomersnon-stereoisomeric counterpart. To avoid
this bias, we removed stereoisomerism information based on the canonical SMILES. The
intersection of SIDERITE with other chemical databases is removed from the chemical
database in the visualization.
657

Clustering of siderophores

The identification of large clusters of siderophores was based on their arrangement within the vast chemical space. The chemical space is composed of numerous molecules, and in our study, we employed the COCONUT database[18] which comprises approximately 40.7w small molecules. Within this space, siderophores were found to be grouped into distinct large clusters, separated from other non-siderophore molecules.

To identify smaller clusters within each large siderophore cluster, we utilized a molecular structure similarity threshold of 60%. Specifically, siderophores were considered to be part of the same small cluster if their molecular similarity was greater than 60%, as determined using the Dice coefficient (Dc)[55].

Testing iron-binding activity by CAS test experiment

We initially screened 48 small molecules from the commercial natural product library for potential siderophore iron-chelating activity, alongside 8 small molecules that lacked such activity, using our developed method. Subsequently, we procured these small molecules individually (Table S5) The small molecules were then prepared by dissolving them in sterilized deionized water to achieve a final concentration of 1 mg/mL in an aqueous solution.

These aqueous solutions were employed to assess the iron-binding activity through the CAS test experiment[57]. Briefly, the liquid form of the CAS assay was employed, wherein 100 µl of the small molecule aqueous solution (four biological replicates for all small molecules) or sterilized deionized water as a control reference was added to 100 µl of the CAS assay solution in a 96-well plate. Following a static incubation of 2 hours at room temperature, the OD630 readings of the small molecule aqueous solution and sterilized deionized water and dimethyl sulfoxide (DMSO) controls were measured using a SpectraMax M5 plate reader at room temperature. Small molecules exhibiting siderophore iron-chelating activity would induce a color change in the CAS medium, leading to decreased OD630 measurements. Consequently, the ability of small molecules to chelate iron can be quantified using the OD630 measurements.

Funding

This work was supported by the National Natural Science Foundation of China (No. 42107140, No. 32071255, No. T2321001, No.42325704), the National Key Research and Development Program of China (No. 2021YFF1200500), National Postdoctoral Program for Innovative Talents (No. BX2021012).

Reference

1   Puig, S., Ramos-Alonso, L., Romero, A. M. & Martinez-Pastor, M. The elemental role

of iron in DNA synthesis and repair. *Metallomics* **9**, 1483-1500,

doi:10.1039/c7mt00116a (2017).

2   Read, A. D., Bentley, R. E. T., Archer, S. L. & Dunham-Snary, K. J. Mitochondrial

iron-sulfur clusters: Structure, function, and an emerging role in vascular biology.

Redox Biol **47**, doi:ARTN 102164

10.1016/j.redox.2021.102164 (2021).

3   Hider, R. C. & Kong, X. L. Chemistry and biology of siderophores. *Nat Prod Rep* **27**,

637-657, doi:10.1039/b906679a (2010).

4   He, R. *et al.* Knowledge-guided data mining on the standardized architecture of NRPS:

Subtypes, novel motifs, and sequence entanglements. *PLOS Computational Biology*

19, e1011100, doi:10.1371/journal.pcbi.1011100 (2023).

5   Kramer, J., Özkaya, Ö. & Kümmerli, R. Bacterial siderophores in community and host

interactions. *Nature Reviews Microbiology* **18**, 152–163,

doi:10.1038/s41579-019-0284-4 (2020).

6   Page, M. G. P. The Role of Iron and Siderophores in Infection, and the Development

of Siderophore Antibiotics. *Clinical Infectious Diseases* **69**, S529-S537,

doi:10.1093/cid/ciz825 (2019).

7   Ahmed, E. & Holmström, S. J. M. Siderophores in environmental research: roles and

applications. *Microbial Biotechnology* **7**, 196-208,

doi:https://doi.org/10.1111/1751-7915.12117 (2014).

8   Barry, S. M. & Challis, G. L. Recent advances in siderophore biosynthesis. *Current*

Opinion in Chemical Biology **13**, 205-215,

doi:https://doi.org/10.1016/j.cbpa.2009.03.008 (2009).

9   Carroll, C. S. & Moore, M. M. Ironing out siderophore biosynthesis: a review of

non-ribosomal peptide synthetase (NRPS)-independent siderophore synthetases. *Crit*

Rev Biochem Mol **53**, 356-381, doi:10.1080/10409238.2018.1476449 (2018).

10   Gauglitz, J. M. & Butler, A. Amino acid variability in the peptide composition of a suite

of amphiphilic peptide siderophores from an open ocean Vibrio species. *JBIC Journal*

of Biological Inorganic Chemistry **18**, 489-497, doi:10.1007/s00775-013-0995-3

(2013).

11   Gao, Y., Walt, C., Bader, C. D. & Müller, R. Genome-Guided Discovery of the

Myxobacterial Thiolactone-Containing Sorangibactins. *ACS Chemical Biology* **18**,

924-932, doi:10.1021/acschembio.3c00063 (2023).

12   Grosse, C., Brandt, N., Van Antwerpen, P., Wintjens, R. & Matthijs, S. Two new

siderophores produced by Pseudomonas sp. NCIMB 10586: The anti-oomycete

non-ribosomal peptide synthetase-dependent mupirochelin and the

NRPS-independent triabactin. *Frontiers in Microbiology* **14**,

doi:10.3389/fmicb.2023.1143861 (2023).

13   Rehm, K. *et al.* Chryseochelins—structural characterization of novel citrate-based

siderophores produced by plant protecting Chryseobacterium spp. *Metallomics* **15**,

doi:10.1093/mtomcs/mfad008 (2023).

14   Hermenau, R. *et al.* Gramibactin is a bacterial siderophore with a diazeniumdiolate

ligand system. *Nature Chemical Biology* **14**, 841-843, doi:10.1038/s41589-018-0101-9

(2018).

15   Hermenau, R. *et al.* Genomics-Driven Discovery of NO-Donating Diazeniumdiolate

Siderophores in Diverse Plant-Associated Bacteria. *Angewandte Chemie International*

Edition **58**, 13024-13029, doi:https://doi.org/10.1002/anie.201906326 (2019).

16   Martinet, L. *et al.* A Single Biosynthetic Gene Cluster Is Responsible for the

Production of Bagremycin Antibiotics and Ferroverdin Iron Chelators. *mBio* **10**,

e01230-01219, doi:doi:10.1128/mBio.01230-19 (2019).

17   Martinet, L., Baiwir, D., Mazzucchelli, G. & Rigali, S. Structure of New Ferroverdins

Recruiting Unconventional Ferrous Iron Chelating Agents. *Biomolecules* **12**, 752

(2022).

18   Sorokina, M., Merseburger, P., Rajan, K., Yirik, M. A. & Steinbeck, C. COCONUT

online: Collection of Open Natural Products database. *Journal of Cheminformatics* **13**,

2, doi:10.1186/s13321-020-00478-9 (2021).

19   Rutz, A. *et al.* The LOTUS initiative for open knowledge management in natural

products research. *eLife* **11**, e70780, doi:10.7554/eLife.70780 (2022).

20   Gallo, K. *et al.* SuperNatural 3.0—a database of natural products and natural

product-based derivatives. *Nucleic Acids Research* **51**, D654-D659,

doi:10.1093/nar/gkac1008 (2022).

21   Weininger, D. SMILES, a chemical language and information system. 1. Introduction

to methodology and encoding rules. *Journal of Chemical Information and Computer*

Sciences **28**, 31-36, doi:10.1021/ci00057a005 (1988).

22 Hirohara, M., Saito, Y., Koda, Y., Sato, K. & Sakakibara, Y. Convolutional neural

network based on SMILES representation of compounds for detecting chemical motif.

BMC Bioinformatics **19**, 526, doi:10.1186/s12859-018-2523-5 (2018).

23 Rajan, K., Brinkhaus, H. O., Zielesny, A. & Steinbeck, C. A review of optical chemical

structure recognition tools. *Journal of Cheminformatics* **12**, 60,

doi:10.1186/s13321-020-00465-0 (2020).

24 Clevert, D.-A., Le, T., Winter, R. & Montanari, F. Img2Mol – accurate SMILES

recognition from molecular graphical depictions. *Chemical Science* **12**, 14174-14181,

doi:10.1039/D1SC01839F (2021).

25 Khokhlov, I., Krasnov, L., Fedorov, M. V. & Sosnin, S. Image2SMILES:

Transformer-Based Molecular Optical Recognition Engine**. *Chemistry–Methods* **2**,

e202100069, doi:https://doi.org/10.1002/cmtd.202100069 (2022).

26 Figueiredo, A. R. T., Ozkaya, O., Kummerli, R. & Kramer, J. Siderophores drive

invasion dynamics in bacterial communities through their dual role as public good

versus public bad. *Ecol Lett* **25**, 138-150, doi:10.1111/ele.13912 (2022).

27 Poggiali, E., Cassinerio, E., Zanaboni, L. & Cappellini, M. D. An update on iron

chelation therapy. *Blood Transfus* **10**, 411-422, doi:10.2450/2012.0008-12 (2012).

28 Gu, S. H. *et al.* Competition for iron drives phytopathogen control by natural

rhizosphere microbiomes. *Nat Microbiol* **5**, 1002-+, doi:10.1038/s41564-020-0719-8

(2020).

29 Gu, S. *et al.* Siderophore-Mediated Interactions Determine the Disease

Suppressiveness of Microbial Consortia. *mSystems* **5**,

doi:10.1128/mSystems.00811-19 (2020).

30   Papanikolaou, G. & Pantopoulos, K. Iron metabolism and toxicity. *Toxicology and*

Applied Pharmacology **202**, 199-211, doi:https://doi.org/10.1016/j.taap.2004.06.021

(2005).

31   Hughes, C. E. *et al.* Cysteine Toxicity Drives Age-Related Mitochondrial Decline by

Altering Iron Homeostasis. *Cell* **180**, 296-+, doi:10.1016/j.cell.2019.12.035 (2020).

32   Sun, H. Y. *et al.* Metabolic Reprogramming in Cancer Is Induced to Increase Proton

Production. *Cancer Res* **80**, 1143-1155, doi:10.1158/0008-5472.Can-19-3392 (2020).

33   Roemhild, K. *et al.* Iron metabolism: pathophysiology and pharmacology. *Trends*

Pharmacol Sci **42**, 640-656, doi:10.1016/j.tips.2021.05.001 (2021).

34   Reitz, Z. L., Hardy, C. D., Suk, J., Bouvet, J. & Butler, A. Genomic analysis of

siderophore β-hydroxylases reveals divergent stereocontrol and expands the

condensation domain family. *Proceedings of the National Academy of Sciences* **116**,

19805-19814, doi:doi:10.1073/pnas.1903161116 (2019).

35   Ferreira, D., Seca, A. M. L., Pinto, D. C. G. A. & Silva, A. M. S. Targeting human

pathogenic bacteria by siderophores: A proteomics review. *J Proteomics* **145**, 153-166,

doi:10.1016/j.jprot.2016.04.006 (2016).

36   Ye, L. M. *et al.* A combinatorial approach to the structure elucidation of a pyoverdine

siderophore produced by a Pseudomonas putida isolate and the use of pyoverdine as

a taxonomic marker for typing P-putida subspecies. *Biometals* **26**, 561-575,

doi:10.1007/s10534-013-9653-z (2013).

37   Zhao, H. *et al.* NPASS database update 2023: quantitative natural product activity and

species source database for biomedical research. *Nucleic Acids Research*,

doi:10.1093/nar/gkac1069 (2022).

38	van Santen, J. A. *et al.* The Natural Products Atlas 2.0: a database of

microbially-derived natural products. *Nucleic Acids Research* **50**, D1317-D1323,

doi:10.1093/nar/gkab941 (2022).

39	D'Onofrio, A. *et al.* Siderophores from Neighboring Organisms Promote the Growth of

Uncultured Bacteria. *Chem Biol* **17**, 254-264, doi:10.1016/j.chembiol.2010.02.010

(2010).

40	Maglangit, F., Tong, M. H., Jaspars, M., Kyeremeh, K. & Deng, H. Legonoxamines

A-B, two new hydroxamate siderophores from the soil bacterium, Streptomyces sp.

MA37. *Tetrahedron Lett* **60**, 75-79, doi:10.1016/j.tetlet.2018.11.063 (2019).

41	Wilson, M. K., Abergel, R. J., Raymond, K. N., Arceneaux, J. E. L. & Byers, B. R.

Siderophores of Bacillus anthracis, Bacillus cereus, and Bacillus thuringiensis.

Biochem Bioph Res Co **348**, 320-325, doi:10.1016/j.bbrc.2006.07.055 (2006).

42	Zajdowicz, S. *et al.* Purification and Structural Characterization of Siderophore

(Corynebactin) from Corynebacterium diphtheriae. *Plos One* **7**, doi:ARTN e34591

10.1371/journal.pone.0034591 (2012).

43	Patzer, S. I. & Braun, V. Gene Cluster Involved in the Biosynthesis of Griseobactin, a

Catechol-Peptide Siderophore of Streptomyces sp ATCC 700974. *J Bacteriol* **192**,

426-435, doi:10.1128/Jb.01250-09 (2010).

44	Matsuo, Y. *et al.* Streptobactin, a Tricatechol-Type Siderophore from Marine-Derived

Streptomyces sp YM5-799. *J Nat Prod* **74**, 2371-2376, doi:10.1021/np200290j (2011).

827 45 Carrero, M. I. G., Sangari, F. J., Aguero, J. & Lobo, J. M. G. Brucella abortus strain

828   2308 produces brucebactin, a highly efficient catecholic siderophore. *Microbiol-Sgm*

829   **148**, 353-360, doi:Doi 10.1099/00221287-148-2-353 (2002).

830 46 Miller, A. L., Li, S. R., Eichhorn, C. D., Zheng, Y. B. & Du, L. C. Identification and

831   Biosynthetic Study of the Siderophore Lysochelin in the Biocontrol Agent Lysobacter

832   enzymogenes. *J Agr Food Chem*, doi:10.1021/acs.jafc.3c01250 (2023).

833 47 Kummerli, R., Schiessl, K. T., Waldvogel, T., McNeill, K. & Ackermann, M. Habitat

834   structure and the evolution of diffusible siderophores in bacteria. *Ecol Lett* **17**,

835   1536-1544, doi:10.1111/ele.12371 (2014).

836 48 Francoeur, P. G. & Koes, D. R. SolTranNet-A Machine Learning Tool for Fast

837   Aqueous Solubility Prediction. *J Chem Inf Model* **61**, 2530-2536,

838   doi:10.1021/acs.jcim.1c00331 (2021).

839 49 Evans, R., Dal Poggetto, G., Nilsson, M. & Morris, G. A. Improving the Interpretation of

840   Small Molecule Diffusion Coefficients. *Anal Chem* **90**, 3987-3994,

841   doi:10.1021/acs.analchem.7b05032 (2018).

842 50 Völker, C. & Wolf-Gladrow, D. A. Physical limits on iron uptake mediated by

843   siderophores or surface reductases. *Marine Chemistry* **65**, 227-244,

844   doi:https://doi.org/10.1016/S0304-4203(99)00004-3 (1999).

845 51 Demain, A. L. Microbial Secondary Metabolism - a New Theoretical Frontier for

846   Academia, a New Opportunity for Industry. *Ciba F Symp* **171**, 3-23 (1992).

847 52 Sinsabaugh, R. L., Hill, B. H. & Shah, J. J. F. Ecoenzymatic stoichiometry of microbial

848   organic nutrient acquisition in soil and sediment. *Nature* **462**, 795-U117,

849   doi:10.1038/nature08632 (2009).

850   53   Chen, Y. L. *et al.* Linking microbial C:N:P stoichiometry to microbial community and

851   abiotic factors along a 3500-km grassland transect on the Tibetan Plateau. *Global*

852   *Ecol Biogeogr* **25**, 1416-1427, doi:10.1111/geb.12500 (2016).

853   54   Probst, D. & Reymond, J. L. Visualization of very large high-dimensional data sets as

854   minimum spanning trees. *Journal of Cheminformatics* **12**, doi:ARTN 12

855   10.1186/s13321-020-0416-x (2020).

856   55   Maggiora, G., Vogt, M., Stumpfe, D. & Bajorath, J. Molecular similarity in medicinal

857   chemistry. *J Med Chem* **57**, 3186-3204, doi:10.1021/jm401411z (2014).

858   56   Vivanco, J. M., Bais, H. P., Stermitz, F. R., Thelen, G. C. & Callaway, R. M.

859   Biogeographical variation in community response to root allelochemistry: novel

860   weapons and exotic invasion. *Ecol Lett* **7**, 285-292,

861   doi:10.1111/j.1461-0248.2004.00576.x (2004).

862   57   Schwyn, B. & Neilands, J. B. Universal chemical assay for the detection and

863   determination of siderophores. *Anal Biochem* **160**, 47-56,

864   doi:10.1016/0003-2697(87)90612-9 (1987).

865   58   Fazary, A. E., Taha, M. & Ju, Y. H. Iron Complexation Studies of Gallic Acid. *J Chem*

866   *Eng Data* **54**, 35-42, doi:10.1021/je800441u (2009).

867   59   Matthijs, S. *et al.* The Pseudomonas siderophore quinolobactin is synthesized from

868   xanthurenic acid, an intermediate of the kynurenine pathway. *Molecular Microbiology*

869   **52**, 371-384, doi:10.1111/j.1365-2958.2004.03999.x (2004).

60   Barraza, M. L. *et al.* Iron Sulfate and Feed Pelleting to Detoxify Free Gossypol in

Cottonseed Diets for Dairy-Cattle. *J Dairy Sci* **74**, 3457-3467, doi:DOI

10.3168/jds.S0022-0302(91)78536-6 (1991).

61   Mossialos, D. *et al.* Quinolobactin, a New Siderophore of *Pseudomonas fluorescens*

ATCC 17400, the Production of Which Is Repressed by the Cognate Pyoverdine.

Applied and Environmental Microbiology **66**, 487-492,

doi:doi:10.1128/AEM.66.2.487-492.2000 (2000).

62   Matthijs, S. *et al.* Thioquinolobactin, a Pseudomonas siderophore with antifungal and

anti-Pythium activity. *Environ Microbiol* **9**, 425-434,

doi:https://doi.org/10.1111/j.1462-2920.2006.01154.x (2007).

63   石井, 永., 陳, 益., 赤池, 美., 石川, 勉. & 盧, 盛. ミカン科植物成分の研究(第 44 報)

台湾産ネワタノキ Xanthoxylum integrifoliolum(MERR.)MERR.(Fagara integrifoliola

MERR.) の成分検索 その 1 根木質部の成分. *藥學雜誌* **102**, 182-195,

doi:10.1248/yakushi1947.102.2_182 (1982).

64   Chen, J.-J., Wang, T.-Y. & Hwang, T.-L. Neolignans, a Coumarinolignan, Lignan

Derivatives, and a Chromene: Anti-inflammatory Constituents from Zanthoxylum

avicennae. *J Nat Prod* **71**, 212-217, doi:10.1021/np070594k (2008).

65   Moon, S.-S. *et al.* Jineol, a Cytotoxic Alkaloid from the Centipede Scolopendra

subspinipes. *J Nat Prod* **59**, 777-779, doi:10.1021/np960188t (1996).

66   Cordero, O. X., Ventouras, L. A., DeLong, E. F. & Polz, M. F. Public good dynamics

drive evolution of iron acquisition strategies in natural bacterioplankton populations. *P*

Natl Acad Sci USA **109**, 20059-20064, doi:10.1073/pnas.1213344109 (2012).

67   Butaite, E., Baumgartner, M., Wyder, S. & Kummerli, R. Siderophore cheating and

cheating resistance shape competition for iron in soil and freshwater Pseudomonas

communities. *Nature Communications* **8**, doi:ARTN 414

10.1038/s41467-017-00509-4 (2017).

68   Ozkaya, O., Balbontin, R., Gordo, I. & Xavier, K. B. Cheating on Cheaters Stabilizes

Cooperation in Pseudomonas aeruginosa. *Curr Biol* **28**, 2070-+,

doi:10.1016/j.cub.2018.04.093 (2018).

69   Shepherdson, E. M. F. & Elliot, M. A. Cryptic specialized metabolites drive

Streptomyces exploration and provide a competitive advantage during growth with

other microbes. *Proceedings of the National Academy of Sciences* **119**, e2211052119,

doi:doi:10.1073/pnas.2211052119 (2022).

70   Raines, D. J. *et al.* Bacteria in an intense competition for iron: Key component of the

Campylobacter jejuni iron uptake system scavenges enterobactin hydrolysis product.

Proceedings of the National Academy of Sciences **113**, 5850-5855,

doi:doi:10.1073/pnas.1520829113 (2016).

71   Zheng, S. J. *et al.* Deep learning driven biosynthetic pathways navigation for natural

products with BioNavi-NP. *Nature Communications* **13**, doi:ARTN 3342

10.1038/s41467-022-30970-9 (2022).

72   Isaka, M., Tanticharoen, M., Kongsaeree, P. & Thebtaranonth, Y. Structures of

cordypyridones A-D, antimalarial N-hydroxy- and N-methoxy-2-pyridones from the

insect pathogenic fungus Cordyceps nipponica. *J Org Chem* **66**, 4803-4808, doi:DOI

10.1021/jo0100906 (2001).

73   Pita-Grisanti, V., Chasser, K., Sobol, T. & Cruz-Monserrate, Z. Understanding the

Potential and Risk of Bacterial Siderophores in Cancer. *Front Oncol* **12**, doi:ARTN

867271

10.3389/fonc.2022.867271 (2022).

74   Ganz, T. & Nemeth, E. Iron homeostasis in host defence and inflammation. *Nat Rev*

Immunol **15**, 500-510, doi:10.1038/nri3863 (2015).

75   Nairz, M. & Weiss, G. Iron in infection and immunity. *Mol Aspects Med* **75**, doi:ARTN

100864

10.1016/j.mam.2020.100864 (2020).

76   Chuljerm, H. *et al.* Characterization of two siderophores produced by Bacillus

megaterium: A preliminary investigation into their potential as therapeutic agents.

Bba-Gen Subjects **1864**, doi:ARTN 129670

10.1016/j.bbagen.2020.129670 (2020).

77   Achard, M. E. S. *et al.* An antioxidant role for catecholate siderophores in Salmonella.

Biochem J **454**, 543-549, doi:10.1042/Bj20121771 (2013).

78   Peralta, D. R. *et al.* Enterobactin as Part of the Oxidative Stress Response Repertoire.

Plos One **11**, doi:ARTN e0157799

10.1371/journal.pone.0157799 (2016).

79   Hesse, E. *et al.* Ecological selection of siderophore-producing microbial taxa in

response to heavy metal contamination. *Ecol Lett* **21**, 117-127, doi:10.1111/ele.12878

(2018).

80   Roskova, Z., Skarohlid, R. & McGachy, L. Siderophores: an alternative bioremediation

strategy? *Sci Total Environ* **819**, 153144, doi:10.1016/j.scitotenv.2022.153144 (2022).

937