1   **SMAD4 promotes somatic-germline contact during murine oocyte growth**

2   Sofia Granados-Aparici[1,3,4,5], Qin Yang[1] and Hugh Clarke[1,2,3]

[1] Research Institute, McGill University Health Centre, 1001 Boul. Décarie, Montréal, QC, Canada H4A 3J1

[2] Departments of Obstetrics and Gynecology and Biology, Division of Experimental Medicine, McGill

University

[3] Corresponding authors (sograap@gmail.com; hugh.clarke@mcgill.ca)

[4] Cancer CIBER (CIBERONC), Madrid.

[5] Pathology Department, Medical School, University of Valencia-INCLIVA, Valencia.

Abstract

Development of the mammalian oocyte requires physical contact with the surrounding granulosa cells of

the follicle, which provide it with essential nutrients and regulatory signals. This contact is achieved

through specialized filopodia, termed transzonal projections (TZPs), that extend from the granulosa cells

to the oocyte surface. Transforming growth factor (TGFβ) family ligands produced by the oocyte increase

the number of TZPs, but how they do so is unknown. Using an inducible Cre recombinase strategy together

with expression of green fluorescent protein to verify Cre activity in individual granulosa cells, we

examined the effect of depleting the canonical TGFβ mediator, SMAD4. We observed a 20-50% decrease

in the total number of TZPs in SMAD4-depleted granulosa cell-oocyte complexes, and a 50% decrease in

the number of newly generated TZPs when the granulosa cells were reaggregated with granulosa cell-free

wild-type oocytes. Three-dimensional image analysis revealed that TZPs of SMAD4-depleted cells were

also longer than controls and more frequently oriented towards the oocyte. Strikingly, the

transmembrane proteins, N-cadherin and Notch2, were reduced by 50% in these cells. SMAD4 may thus

modulate a network of cell adhesion proteins that stabilize the attachment of TZPs to the oocyte, thereby

amplifying signalling between the two cell types.

Impact statement

Genetic deletion of SMAD4 in the granulosa cells of growing ovarian follicles reduces the number of

transzonal projections that contact the oocyte, implicating TGFβ-family signaling in maintaining soma-

germ line communication.

Key words

SMAD / oocyte / TZPs / filopodia / follicle / signalling

Introduction

The development of a healthy oocyte within the ovarian follicle requires continuing contact between the germ cell and the somatic granulosa cells that surround it (El-Hayek and Clarke, 2016, Clarke, 2018, Richani et al., 2021, Robert, 2021, Li and Albertini, 2013, Doherty et al., 2022, Xie et al., 2023, Marchais et al., 2022). Signals from the granulosa cells initiate the increase in volume of the oocyte when primordial (non-growing) follicles are activated to enter the growth phase (Zhang et al., 2014, Saatcioglu et al., 2016, Nagamatsu et al., 2019). Subsequently, the granulosa cells provide nucleotides, amino acids and pyruvate to the growing oocyte via gap junctions that connect the two cell types (Xie et al., 2023, Clarke, 2018, Marchais et al., 2022). When gap junctional communication between the two cell types is genetically disrupted, oocytes do not develop normally and cannot be fertilized (Gittens and Kidder, 2005, Simon et al., 1997, Winterhager and Kidder, 2015, Carabatsos et al., 2000). Once the oocyte reaches full-size, it acquires the ability to undergo the final stage of development, termed meiotic maturation, but is prevented from precociously doing so by cyclic GMP that is produced and transferred to it by the granulosa cells, also via gap junctions (Jaffe and Egbert, 2016, Wigglesworth et al., 2013, Norris et al., 2009, Zhang et al., 2010). In vitro, oocytes can develop normally when enclosed by granulosa cells but not when the granulosa cells are removed and provided in co-culture, further emphasizing the crucial importance of physical contact between the two cell types (O'Brien et al., 2003, Eppig, 1979, Herta et al., 2021).

In primordial follicles, a single layer of squamous granulosa cells surrounds the oocyte and contact it via adherens junctions (Mora et al., 2012, Jorgensen, 2013). Following activation of growth, a glycoprotein-rich coat termed the zona pellucida becomes established around the oocyte (Litscher and Wassarman, 2020, Wassarman and Litscher, 2018). The zona pellucida steadily thickens as growth continues, reaching a final thickness of about 7 µm in mice and 15 µm in humans. Despite the barrier imposed by the zona

pellucida, the granulosa cells and oocyte maintain direct contact by a dense network of thin filopodia-like

projections, termed transzonal projections (TZPs), that grow out from the granulosa cells through the zona

pellucida (Clarke, 2022, Macaulay et al., 2014, Baena and Terasaki, 2019, Motta et al., 1994, Zhang et al.,

2021, Albertini and Rider, 1994). The tips of the TZPs establish contact with the oocyte plasma membrane

through heterotypic adherens junctions comprising N-cadherin (granulosa cell) and E-cadherin (oocyte)

and harbour the gap junctions that link the two cell types (Mora et al., 2012) and likely also through focal

adhesions (McGinnis and Kinsey, 2015). New TZPs are generated throughout follicular growth, thereby

increasing intercellular molecular trafficking (El-Hayek et al., 2018a). A genetically imposed decrease in

the number of TZPs leads to defects in oocyte gene expression and to abnormal meiotic maturation and

embryonic development (Crozet et al., 2023). Despite the essential role of TZPs during oocyte

development in mammals, as well as the presence of analogous structures in non-mammalian species

(Kessel et al., 1985, Schroeder, 1981), the molecular mechanisms that govern their generation and

stability are not yet understood.

The TGFβ signalling pathway regulates the functions and interactions of a wide range of cell types under

various physiological and pathological conditions (Derynck and Budi, 2019, Peng et al., 2022). Binding of

the ligand to a complex of type I and type II receptors located at the cell surface activates signalling within

the cell, most commonly through the members of the SMAD family of transcription factors, although

SMAD-independent signaling can also occur (Aashaq et al., 2022, Derynck and Budi, 2019). Upon

phosphorylation of the tetrameric type II and type I receptors and subsequent phosphorylation of

receptor-activated SMAD proteins (R-SMADs; SMAD1, 2, 3, 5 and 8), a heterotrimeric complex with

SMAD4 is formed which enters the nucleus and regulates gene expression by binding to specific SMAD-

binding DNA elements. Although R-SMADs can signal in the absence of SMAD4, evidence indicates that

the presence of SMAD4 in the heterotrimer amplifies TGFβ cell signalling by stabilizing the complex and

by favouring the binding of other transcriptional co-repressors or co-activators of target genes (Aragón et

al., 2019, Guglielmi et al., 2021).

Genetic, pharmacological and in vitro studies have revealed multiple roles for TGFβ/SMAD signaling

during follicular growth and development (Patton et al., 2021, Pangas and Rajkovic, 2015). TGFβ signalling

via SMAD3 in granulosa cells acts initially in primordial follicles to maintain their quiescence (Granados-

Aparici *et al*, 2019; Wang *et al*, 2014) and later, following their activation, becomes excluded from the

nucleus in association with rapid granulosa cell growth and proliferation (Hardy *et al*, 2018). The TGFβ

superfamily members, growth-differentiation factor (GDF) 9, which signals through SMAD2/3, and bone

morphogenetic protein (BMP) 15, which signals through SMAD1/5/8, are produced by the oocyte (Dong

et al., 1996, Yan et al., 2001). Female mice lacking GDF9 rarely generate more than a single layer of

granulosa cells (Elvin et al., 1999a, Elvin et al., 1999b, Dong et al., 1996). This defect can be rescued by

preventing the production of inhibin, and the doubly mutant follicles can develop at least to the pre-antral

stage (Wu et al., 2004). Genetic inactivation of SMAD4 in the granulosa cells from growing follicles leads

to a reduction in the number of antral follicles and ovulated eggs, as well as the appearance of luteinized

follicles containing entrapped oocytes (Pangas et al., 2006). Similar phenotypes are observed when both

SMAD2 and SMAD3 are inactivated in the granulosa cells (Li et al., 2008). These effects are likely due at

least in part to a failure of the cumulus layer to expand in response to the ovulatory surge of LH (Pangas

et al., 2006, Li et al., 2008), as expansion is known to require SMAD signaling and to be required for

ovulation (Dragovic et al., 2007, Hao et al., 2022, Mottershead et al., 2015, Peng et al., 2013, Yu et al.,

2013).

Two lines of evidence point to a potential role for SMAD signaling in the generation or maintenance of

TZPs. First, deletion of GDF9 leads to both a decrease in the number of TZPs and a failure of a proportion

of the TZPs to correctly orient towards the oocyte (Carabatsos et al., 1998). Second, TZP number is

reduced in follicles grown in vitro following RNAi-mediated depletion of GDF9 in the oocyte (El-Hayek et

al., 2018a). To test the role of SMAD4 specifically, we used a Cre-lox strategy to inactivate *Smad4* in

granulosa cells, through expression of the recombinase under the control of the *Amhr2* (anti-Mullerian

hormone receptor 2) promoter, as previously described (Jorgez et al., 2004, Pangas et al., 2006). Using

image analysis tools and three-dimensional reconstructions, we show that depletion of SMAD4 in the

granulosa cells decreases the number of TZPs that project from individual granulosa cells in both intact

granulosa cell-oocyte complexes (GOCs) and in reaggregated complexes. Morphometric parameters

including length and orientation are also altered. The decrease in TZP number in intact GOCs is associated

with a reduction in the amount of N-cadherin and Notch2 in the granulosa cells, suggesting that the

composition of the granulosa cell surface is perturbed in the absence of SMAD4. We propose that SMAD4,

through its activity to increase expression of target genes, stabilizes TZPs by promoting their adhesion to

the oocyte surface, thereby helping to increase the number of TZPs and communication between the

growing oocyte and its follicular microenvironment.

Results

We began by verifying the expression of SMAD4 during folliculogenesis. As shown in Fig. 1A and consistent

with prior reports, SMAD4 was detectable in primordial and growing follicles as well as in the granulosa

cells collected from pre-antral follicles. SMAD2 and SMAD3 were also present, with SMAD2 being the

major species detected in oocytes. Two strategies were tested to deplete SMAD4 from the granulosa cells

of growing follicles. First, mice were generated carrying two floxed *Smad4* alleles and a transgene

encoding Cre recombinase under the control of the *Amhr2* promoter (Fig. 1B). These mice also carried a

transgene (*mTmG*, see Methods) constructed such that Tomato fluorescent protein is expressed in the

absence of Cre activity, whereas Cre-mediated recombination switches expression to green fluorescent

protein (GFP), thus allowing us to identify cells where Cre had been active. Upon immunostaining of

ovarian sections or whole-mount intact follicles, only a small number of cells expressed GFP (Fig. 1C, left).

Consistent with this observation, less than 10% of FACS-sorted granulosa cells were GFP-positive (Fig. 1C,

right). Thus, in our system, *Amhr2-Cre* did not mediate efficient recombination of floxed DNA in granulosa

cells. This is consistent with results described in previous reports (Pangas et al., 2006, Li et al., 2008).

We therefore turned to mice that expressed Cre under the control of a tamoxifen-inducible form of the

estrogen receptor (ER) gene promoter (*ER-Cre*). Because ER is expressed in pituitary gonadotropes and

SMAD4 regulates the transcription of the *Fshb* gene encoding the β-subunit of follicle-stimulating

hormone (Li et al., 2018), this model could not be used in vivo. Granulosa cell-oocyte complexes (GOCs)

from mid-growth stage follicles (diameter 100-120 μm) were therefore isolated, exposed to tamoxifen for

one day and then incubated in tamoxifen-free medium for another five days. Under these conditions, a

substantial fraction of the granulosa cells underwent Cre-mediated recombination, as indicated by the

expression of GFP in immunostained GOCs and by FACS analysis (Fig. 1E). This in vitro system was used

for the experiments described here.

To examine the efficiency of Cre-mediated *Smad4* recombination, the mRNA level was assessed in intact

GOCs using the RNAscope technique and in purified granulosa cells using qRT-PCR. Following tamoxifen

treatment and culture, there was an approximately 60% reduction in the number of foci in the granulosa

cells from 5.78 +/- 0.36 (mean, s.e.m., n=21) in *ER-Cre$^-$*; *Smad4$^{fl/fl}$* mice to 2.16 +/- 0.26 (n=26) in *ER-Cre$^+$*;

Smad4$^{fl/fl}$ mice (p<0.0001), and a 70% reduction in the amount of PCR product from granulosa cells (n=5,

p=0.0001) (Fig. 1F). Immunoblotting of granulosa cells revealed a corresponding 70% reduction in SMAD4

protein in the granulosa cells of *ER-Cre$^+$*; *Smad4$^{fl/fl}$* mice compared to *ER-Cre$^-$*; *Smad4$^{fl/fl}$* controls (n=11,

p<0.0001) (Fig. 1G). Thus, the in vitro ER-Cre strategy substantially reduced although did not eliminate

SMAD4 from the granulosa cells.

To examine the effect of SMAD4 depletion on the TZPs, GOCs were isolated and cultured as above, fixed

and stained using anti-GFP to label cells where Cre had been active and phalloidin to label the TZPs, and

imaged them using confocal microscopy. First, the TZPs were analyzed as a population. Because TZPs arise

from both the innermost and more distal granulosa cell layers (Baena and Terasaki, 2019), it was not

possible to reliably determine how many granulosa cells contributed the GFP-positive TZPs in a GOC. The

total number of phalloidin-stained TZPs – irrespective of whether they were GFP-positive – that projected

to the oocyte in equatorial optical sections of GOCs from control ($Smad4^{wt/wt}$) and experimental ($Smad4^{fl/fl}$)

mice were therefore counted (Fig. 2A, B). TZP-number was normalized to the diameter of the

corresponding oocyte, although the mean oocyte diameter did not differ between the groups (Fig. 2B),

We observed a 20.4 +/- 0.05% reduction in the number of TZPs in the GOCs from the $Smad4^{fl/fl}$ mice (n=63

[control], 59 [experimental], p=0.002).

A limitation of quantifying TZPs in intact GOCs is that the granulosa cells, depending on their number and

spatial distribution, can variably impair their resolution and discrimination even using confocal

microscopy. Therefore, the number of TZPs was also counted after mechanically removing the granulosa

cells from the GOCs. This analysis revealed a striking 51.1 +/- 0.04% reduction in the normalized number

of TZPs in the GOCs from the $Smad4^{fl/fl}$ mice (n=44 [control], 47 [experimental], p<0.0001) (Fig. 2C, D). As

observed using intact GOCs, the mean oocyte diameter did not differ (Fig. 2D). To verify the difference in

TZP-number, we also compared $ER-Cre^+$; $Smad4^{fl/fl}$ to a different control genotype, $ER-Cre^-$; $Smad4^{fl/fl}$ mice.

A quantitatively similar 43.6 +/- 0.04% decrease in TZPs within the $ER-Cre^+$ population (n=164 [control],

172 [experimental], p<0.0001) (Fig. 2D) was observed following tamoxifen treatment and culture.

Together, these results indicate that depletion of SMAD4 in the granulosa cells is associated with a

decrease in the number of TZPs.

TZPs provide the physical platform that enables essential gap junctional communication between the

granulosa cells and the oocyte. To test whether the reduction in TZP number in SMAD4-depleted

granulosa cells impaired this communication, we used a fluorescence recovery after photobleaching

(FRAP) assay. Control and experimental GOCs were treated with tamoxifen and incubated as above. After

five days, they were exposed to calcein-AM, a fluorescent dye that can pass through gap junctions. The

fluorescence in the oocyte of each GOC was bleached using the laser of a confocal microscope and its

recovery was then recorded. As shown in Fig. 2E, which plots the fluorescence in the oocyte as a function

of time after photobleaching and at the end of the recording period normalized to the initial fluorescence,

we observed no difference between groups in the rate of transfer of fluorescence from the granulosa cells

to the oocyte. Thus, despite the decreased number of TZPs, depletion of SMAD4 did not detectably impair

gap junctional communication between the granulosa cells and the oocyte.

As TZPs project not only from the cells immediately adjacent to the zona pellucida but also from those in

more distal layers, it was possible that the reduced number of TZPs in *ER-Cre$^+$*; *Smad4$^{fl/fl}$* complexes

reflected an impaired ability of the granulosa cells to proliferate. *ER-Cre$^+$; Smad4$^{+/+}$* and *ER-Cre$^+$; Smad4$^{fl/fl}$*

GOCs were incubated for 5 hr in the presence of the thymidine analogue, EdU, added five days after the

tamoxifen exposure. No difference between the groups in the fraction of EdU-positive cells was observed

(Fig. 2F), however, implying that the reduced number of TZPs was not due to impaired proliferation of the

granulosa cells.

We then focused our analysis on individual granulosa cells. GOCs where individual GFP-positive cells could

be readily identified and the TZPs were easily attributable to a particular cell (Fig. 3A) were selected, serial

optical sections were obtained using confocal microscopy, and 3-dimensional images were reconstructed

using Imaris software. Fig. 3B shows one such reconstruction, where GFP-positive granulosa cells are

pseudo-coloured in green. We observed a 25% decrease in the mean volume of *Smad4$^{fl/fl}$* granulosa cells

compared to wild-type cells (323.2 +/- 11.2 vs 242.6 +/- 10.0; n=293, 322; *p*<0.0001), but no difference

between genotypes in the sphericity of the cells (Fig. 3C). The TZPs in a subset of the GFP-positive

granulosa cells were then analyzed (Fig. 3D). In this subset, there was a 15% difference in mean cell volume

(392.2 +/- 20.8 vs 334.0 +/- 21.2; n=55, 63) which did not reach statistical significance (*p*=0.054). However,

whereas wild-type cells harbored a mean of 15.8 +/- 0.87 (n=55) TZPs per cell, this number was reduced

by 35% in *Smad4^{fl/fl}* cells to 10.5 +/- 0.71 (n=65, *p*<0.0001). As compared to those of wild-type cells, the

TZPs of the *Smad4^{fl/fl}* cells were about 10% longer (4.64 +/- 0.08 μm vs 5.19 +/- 0.09 μm; n=893, 720;

p<0.0001) and very slightly narrower (0.40 +/- 0.003 μm vs 0.38 +/- 0.003 μm; n=870, 719; *p*<0.0001)(Fig.

3E). We did not detect other differences between the genotypes in TZP morphology. These results

obtained using reconstructed 3-dimensional images of GFP-positive cells establish that the number of

TZPs is reduced in individual SMAD4-depleted granulosa cells.

A limitation of analyzing TZPs in GFP-positive cells within GOCs is that that this strategy cannot distinguish

TZPs that were generated in a low-SMAD4 environment (i.e., after the Cre recombinase was expressed)

from those generated earlier in a normal-SMAD4 environment. To ensure that we examined only TZPs

that were generated after Cre had been expressed, we used our previously published reaggregation assay

(El-Hayek et al., 2018a) (Fig. 4A). GOCs obtained from mid-growth stage follicles of *ER-Cre; mTmG* mice

carrying the appropriate *Smad4* alleles were exposed to tamoxifen for one day and then incubated for an

additional four days. The granulosa cells were then separated from the oocyte and combined with an

oocyte that did not carry the *mTmG* gene. These reaggregated GOCs were incubated for an additional 6

days, then fixed and stained using anti-GFP (Fig. 4B).

When the entire population of granulosa cells in the reaggregated GOCs was analyzed, there was a ~30%

reduction in mean volume in the *Smad4^{fl/fl}* cells as compared to *Smad4^{+/+}* cells (394.5 +/- 18.8 vs 279.8 +/-

16.6; n=185, 159; p<0.0001) and no difference in cell sphericity (Fig. 4C), as observed using intact GOCs.

There was no difference between genotypes, however, in mean volume among the cells whose TZPs were

characterized (Fig. 4D). Within this group, the mean number of TZPs per cell was reduced by ~50%, from

14.2 +/- 2.0 (n=16) in the *Smad4*$^{+/+}$ cells to 6.84 +/- 1.1 (n=25) in the *Smad4*$^{fl/fl}$ cells (Fig. 4D, p=0.001). A

~20% increase in the mean length of the TZPs of *Smad4*$^{fl/fl}$ cells (3.71 +/- 0.17 vs 4.40 +/- 0.28; n=174, 154;

p=0.03) was also detected, as in intact GOCs, but no difference in the mean width (Fig. 4E).

The increased mean length of the TZPs of *Smad4*$^{fl/fl}$ cells compared to *Smad4*$^{+/+}$ cells prompted us to

analyze the trajectory of individual TZPs. We traced their path from their point of origin to their tip, then

measured the distance between the point of origin and the oocyte and subtracted from this the distance

between the tip and the oocyte. A large positive difference indicates that the tip of the TZP is relatively

close to the oocyte; conversely, a small positive or a negative difference indicates that that tip is relatively

far from the oocyte (Fig. 4F). This analysis revealed that the mean difference was 50% greater in the

Smad4$^{fl/fl}$ cells compared to *Smad4*$^{+/+}$ cells (1.61 +/- 0.16 vs 2.40 +/- 0.21; n=174, 157; *p*=0.003) (Fig. 4G).

This indicates that the tips of the TZPs of the *Smad4*$^{fl/fl}$ cells were, on average, closer to the oocyte than

those of the *Smad4*$^{+/+}$ cells. We also found that the proportion of TZPs that were oriented away from the

oocyte trended towards being greater in the *Smad4*$^{+/+}$ cells, although the difference did not reach

statistical significance (Fig. 4G). Taken together, the trajectory analyses provide additional evidence that

the TZPs of *Smad4*$^{fl/fl}$ cells differ from those of *Smad4*$^{+/+}$ cells.

The observation that TZP-number was modestly reduced when intact GOCs were analyzed, yet markedly

reduced when oocytes from which the granulosa cells had been stripped were analyzed, suggested that

the attachment between the TZPs and oocyte might be weakened when SMAD4 was depleted. To

evaluate this, the amount of N-cadherin, which is the principal cell adhesion molecule that has been

identified in granulosa cells, was compared between *Smad4*$^{+/+}$ and *Smad4*$^{fl/fl}$ granulosa cells. Consistent

with the previous results (see Fig. 1E), SMAD4 was reduced by 73.5 +/- 0.05% in the *Smad4*$^{fl/fl}$ granulosa

cells (Fig. 5A, n=6, *p*<0.0001). Strikingly, N-cadherin was reduced by 52.7 +/- 0.04 % (n = 6, *p*=0.0004) in

the *Smad4[fl/fl]* granulosa cells (Fig. 5A). Moreover, Notch2, a transmembrane receptor expressed in

granulosa cells that associates with membrane-bound Jagged1 expressed on the oocyte, was reduced by

52.6 +/- 0.05% (n=5; *p*=0.0003) in these cells (Fig. 5B). In contrast to the protein, however, there was no

detectable reduction in the amount of encoding mRNA in either case as assayed using qRT-PCR, although

Smad4 was reduced in the same samples (Fig. 5C). Thus, a reduction in SMAD4 in granulosa cells is

associated with a reduction in certain transmembrane proteins, including those responsible for cell

adhesion.

As we previously reported that depletion of GDF9, which is secreted by the oocyte and signals in the

granulosa cells through SMAD4, was associated with a reduction in mRNAs encoding structural

components of TZPs – *Myo10*, *Fscn1*, and *Daam1* (El-Hayek et al., 2018a), we measured the quantity of

these mRNAs using qRT-PCR. We found that they were not depleted in the *Smad4[fl/fl]* granulosa cells

compared to the *Smad4[+/+]* cells (Fig. 5D). This implies either that the SMAD4 which remained was

sufficient to induce expression of these genes or the GDF9 regulates their expression through a SMAD4-

independent pathway.

Discussion

We have examined the role of SMAD4 establishing the TZPs that connect the granulosa cells to the oocyte.

Using a tamoxifen-inducible Cre recombinase strategy, we were able to reduce SMAD4 in the granulosa

cells to about 30% of its wild-type level. This reduction was associated with a decrease ranging from 20-

50% in the number of TZPs in the granulosa cells assessed as a population (i) in intact GOCs and (ii) after

removal of the granulosa cell bodies to increase optical resolution, and as individual cells (iii) in intact

GOCs and (iv) in reconstructed complexes. Previous studies where SMAD4, or both SMAD2 and SMAD3,

was depleted from granulosa cells described a loose or loss association of the cumulus cells with the

oocyte in some follicles (Pangas et al., 2006, Li et al., 2008). Moreover, as described in the Introduction,

blocking production of GDF9 by the oocyte leads to a decrease in the number of TZPs (Carabatsos et al.,

1998, El-Hayek et al., 2018a). Together these observations suggest that SMAD4 regulates the size of the

population of TZPs in growing follicles.

The population of TZPs presumably reflects the net result of the generation of new TZPs and the retraction

of existing TZPs, as is the case for filopodia (Gallop, 2020, Heckman and Plummer, 2013, Mattila and

Lappalainen, 2008). SMAD4 could thus promote either or both processes. The observation that GFP-

expressing granulosa cells in reaggregated complexes generated new TZPs implies that SMAD4 is not

required for this process, although it does not rule out a role in increasing the rate of generation. A clue

to the function of SMAD4 may lie in the observation that the decrease in the number of TZPs in SMAD4-

depleted cells was quantitatively greater when the granulosa cell bodies were removed prior to analysis.

This suggests that the process of removing the cell bodies also removed the TZPs more easily in the

SMAD4-depleted cells than in wild-type cells.

In this context, it is intriguing that N-cadherin, which is thought to stabilize attachment of the TZP tips to

the oocyte surface, was depleted in the mutant cells. We propose that, by increasing the quantity of N-

cadherin in the granulosa cells, SMAD4 increases the strength of adhesion of the TZPs to the oocyte

surface (Fig. 6). This in turn would decrease the probability of retraction, thereby increasing the steady-

state number of TZPs. This model is consistent with the observation that the non-attached (free-ended)

TZPs of SMAD4-depleted cells were on average longer and closer to the oocyte than those of wild-type

cells. Specifically, if the ability of the TZPs to establish a stable contact with the oocyte were impaired in

SMAD4-depleted cells, a relatively large proportion might continue to 'search' for an attachment and to

lengthen during the searching process.

Mechanistically, some reports have linked SMAD4 to transcription of the *Cdh2*, which encodes N-cadherin

(Kang et al., 2014, Yang et al., 2015), but we did not detect a difference in mRNA quantity between normal

and SMAD4-depleted granulosa cells. In view of the reduced quantity in the latter of Notch2, also a

membrane-associated protein, it may be speculated that SMAD4 might act more broadly to regulate the

expression of genes whose products maintain the normal configuration of the granulosa cell plasma

membrane. An alternative explanation is raised by observations that granulosa-cell depletion of SMAD4

is associated with the presence of follicles containing luteinized granulosa cells enclosing an entrapped

oocyte and increased expression of markers of luteinization, including *Cyp11a1*, *Sfrp4*, *Star* and *Ptgfr*

(Pangas et al., 2006). This suggests that the SMAD4-depleted granulosa cells in the GOCs studied here

might have become luteinized, and that luteinization in turn caused a reduction in the number of TZPs.

We did not detect increased expression of the luteinization markers (unpublished data), suggesting that

luteinization did not occur under the conditions of our experiments. This difference may arise because we

examined GOCs that had been maintained in vitro, whereas the previous study examined follicles in intact

ovaries of prepuberal mice following injection of an FSH analogue.

Regardless of the mechanism linking SMAD4 to TZP number, it is noteworthy that the number of TZPs that

project from an individual granulosa cells – about 15 in this study and about 40 in a previous study where

more advanced (antral) follicles were examined (Baena and Terasaki, 2019) – considerably exceeds the

number of filopodia that are present on individual blastomeres of mouse embryos (5-6) (Fierro-Gonzalez

et al., 2013) or on chick embryonic cells (~4) (Sagar et al., 2015). SMAD4 may therefore act in granulosa

cells to amplify an innate ability to generate filopodia, which become TZPs and increase the efficiency of

communication between the growing germ cell and its somatic microenvironment.

309 **Methods**

Key Resources Table				
Reagent type	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
gene (mouse)	Smad4	GenBank	Gene ID: 17128	
Strain (mouse)	*Smad4[tm2.1Cxd]*/J	Jackson Laboratories	strain 017462	Supplied by Dr. Daniel. J. Bernard, McGill University
Strain (mouse)	*Amhr2[tm3(cre)Bhr]*	MMRRC	strain 014245-UNC	Supplied by Dr. Makoto Nagano, McGill University
Strain (mouse)	B6.129-*Gt(ROSA)26Sor[tm1(cre/ERT2)Tyj]*/J	Jackson Laboratories	strain 008463	
Strain (mouse)	B6.129(Cg)-*Gt(ROSA)26Sor[tm4(ACTB-tdTomato,-EGFP)Luo]*/J	Jackson Laboratories	strain 007676	
RNA in situ hybridization probe	*Smad4*, exon 8	Advanced Cell Diagnostics	857571	
RNA in situ hybridization probe	*Bacillus subtilis DapB*	Advanced Cell Diagnostics	310043	
Primary antibody	SMAD2/3	Cell Signaling Technology	8685	

Primary antibody	SMAD4	Sigma	HPA019154	
Primary antibody	N-cad	BD Biosciences	B610920	
Primary antibody	Notch	Abcam	Ab8926	
Primary antibody	GFP	Invitrogen	A11122	
Secondary antibody	Goat anti-rabbit IgG-HRP	Promega	W4011	
Secondary antibody	Goat anti-mouse IgG-HRP	Promega	W4021	
Secondary antibody	Donkey anti-rabbit IgG- Alexa 488	Invitrogen	A21206	
Reagent	Tamoxifen	Sigma	T176	
Reagent	Phalloidin	Thermo Fisher	A12380	
Reagent	EdU	Thermo Fisher	C10337	

Primers				
Target	application	5-prime	3-prime	Predicted size
Smad4 -wt	Genotype	GGGCAGCGTAGCATATAAGA	GACCCAAACGTCACCTTCAG	390
Smad4 - flox	Genotype	GGGCAGCGTAGCATATAAGA	GACCCAAACGTCACCTTCAG	450
Smad4 - recombined	Genotype	TAAGAGCCACAGGGTCAAGC	GACCCAAACGTCACCTTCAG	450
Amhr2-Cre	Genotype	CTCTGGTGTAGCTGATGATC	TAATCGCCATCTTCCAGCAG	380
ER-Cre	Genotype	CGTGATCTGCAACTCAGTC	AGGCAAATTTTGGTGTACGG	150

Smad4	mRNA	TCACAATGAGCTTGCATTCC	CCATCCACAGTCACAACAGG	143
Cdh2 (N-Cad)	mRNA	CCAGGAAAAGTGGCAGGTAG	CACTGTGCTTGGCAAGTTGT	121
Notch2	mRNA	GACCCTATCCTACCCTCTAGTG	AGCAGGATGAAGAACAGGATG	103
Myo10	mRNA	TCCAGACAGACTATGGGCAGG	GGAAGCCATGTCGTCCACG	109
Fscn1	mRNA	AGAACGCCAGCTGCTACTTT	CGAGGAATCACTACCCACCG	331
Daam1	mRNA	GCGGCTGCTCAGAGTATAGAAA	AAACATGGCTTCCCTGTGTTTG	273
RPL-19	mRNA	GAAATCGCCAATGCCAACTC	CTTCCCTATGCCCATATGCC	147

Mouse strains

All experiments were performed in compliance with the regulations and policies of the Canadian Council

on Animal Care and were approved by the Animal Care Committee of the Research Institute of the McGill

University Health Centre (RI-MUHC, protocol 7783). CD-1 mice were obtained from Charles River (St-

Constant, QC). The following transgenic strains were obtained from commercial suppliers: B6.129-

Gt(ROSA)26Sor^{tm1(cre/ERT2)Tyj}/J (hereafter *ER-Cre*) carrying a tamoxifen-inducible sequence encoding Cre

recombinase under the control of the promoter of the estrogen receptor 2 gene (Jackson Laboratories,

Bar Harbor, ME, strain 008463); B6.129(Cg)-*Gt(ROSA)26Sor^{tm4(ACTB-tdTomato,-EGFP)Luo}*/J (Jackson, strain 007676;

hereafter *mTmG*) carrying sequences encoding a membrane-targeted tdTomato and membrane-targeted

EGFP; Cre-mediated recombination of *loxP* sites switches expression from RFP to EGFP. *Smad4^{tm2.1Cxd}*/J

(hereafter *Smad4^{fl}*) carrying *loxP* sites flanking exon 8 (Jackson, strain 017462) was kindly supplied by Dr.

Daniel Bernard (McGill University). *Amhr2^{tm3(cre)Bhr}* (hereafter *Amhr2-Cre*) carrying a sequence encoding

Cre recombinase under the control of the promoter of the Anti-Mullerian hormone receptor 2 gene was

kindly supplied by Dr. Makoto Nagano (McGill University).

To generate *Amhr2-Cre^+; Smad4^{fl/fl}* females carrying an *mTmG* allele, *Smad4^{fl/wt}; mTmG^{+/+}* females were

crossed with *Amhr2-Cre^+; Smad4^{fl/wt}* males. To generate *ER-Cre^+; Smad4^{fl/fl}* females carrying an *mTmG*

allele, *Smad4$^{fl/wt}$; mTmG$^{+/+}$* females were crossed with *ER-Cre^{+}; Smad4$^{fl/wt}$* males. *ER-Cre^{-}; mTmG^{+};*

Smad4$^{fl/fl}$ females and *ER-Cre^{+}; mTmG^{+}; Smad4$^{wt/wt}$* females served as controls. Genotyping of the offspring

was done using the EZ Tissue/Tail DNA Isolation Plus PCR kit (EZ Bioresearch). After DNA extraction from

tail snips, PCR was used to detect wild type, floxed and recombined *Smad4* and the presence of *ER-Cre*

and *Amhr2-Cre* alleles. Primer sequences are given in Supplemental Table 1.

Tissue embedding and immunohistochemistry

Sections of paraffin-embedded ovaries were deparaffinized using xylene, rehydrated in a decreasing

ethanol series (100, 90, 70%) for 3 min each, and then placed in Milli-Q water for 5 min. Slides were heated

intermittently in a microwave during 20 min in 0.1 M citrate buffer solution at pH 6.0, washed in PBS for

10 min, and blocked with CAS universal blocking solution (Thermo Fisher, 008120) for 30 min at room

temperature to reduce nonspecific binding. After overnight incubation with primary antibody, slides were

washed three times in PBS at room temperature and then incubated for 1 h at room temperature with

secondary antibody with DAPI. Slides were washed twice in PBS for 5 min and mounted on a drop of

VECTASHIELD solution (Vector Laboratories, H-1200-10). Images were acquired using a Zeiss LSM 880

confocal microscope.

Collection, culture and tamoxifen treatment of granulosa cell-oocyte complexes (GOCs)

GOCs were obtained as previously described (El-Hayek et al., 2018b, O'Brien et al., 2003). Briefly, ovaries

of 10- to 12-day-old mice were dissected from the ovarian bursa. Follicles were mechanically isolated by

carefully tearing the ovaries apart and using fine needles to separate the follicles from each other in

HEPES-buffered minimal essential medium (MEM, pH 7.2, Life Technologies, Burlington, ON) (MEM-H). A

Pipetman P1000 was used to carefully pick up and transfer the individual follicles to dishes containing

fresh medium. GOCs were obtained by incubating follicles in the presence of collagenase (10 µg/ml;

Cedarlane, Burlington, ON) and DNase I (10 µg/ml; Sigma) for 5 min at 37°C in air. GOCs were collected

and briefly washed in NaHCO$_3$-buffered MEM (MEM-C) and cultured in 4-well plates containing 500 µl of

MEM-C supplemented with ITS (10 µg/mL insulin, 5.5 µg/mL transferrin and 5 ng/mL selenium; Sigma),

FSH (10 mIU/ml; EMD Serono, Mississauga, ON), and estradiol (10 nM, Sigma, E2758) in an atmosphere

of 5% CO$_2$ in air. To induce Cre-mediated recombination of the *Smad4* allele, GOCs were incubated with

4-hydroxytamoxifen (1 µg/ml, Sigma, T176) in the above medium for 24 hours. They were then transferred

to medium without tamoxifen.

Immunoblotting

Cell samples were lysed in 10 µl of Laemmli buffer. After denaturation at 95∘C for 10 min, proteins were

separated using 4–15% precast polyacrylamide gels (BioRad, 4568085) and transferred onto a

polyvinylidene fluoride membrane (Amersham, Oakville, ON, Canada) under constant voltage. The

membrane was subsequently blocked using 5% BSA, in 0.1% Tween-PBS. The membrane was washed

three times in Tween-PBS and incubated with primary antibody overnight at 4∘C. After washing for 45

min, the membrane was incubated in secondary antibody conjugated to horseradish peroxidase

(Promega) at a dilution of 1:5000 for 1h at room temperature. After washing for 30 min, bound antibody

was revealed using ECL+ (Amersham).

RNA purification and quantitative real-time PCR

RNA was extracted using a Picopure RNA isolation kit (Life Technologies) following the manufacturer's

instructions and eluted in 10 µL of the provided elution buffer, as described (El-Hayek *et al*, 2018a). Briefly,

SuperScript II Reverse Transcription kit (Life Technologies) was used to generate cDNA. PCR amplification

was performed using a Corbett Rotorgene 6000 (Montreal Biotech, Montreal, QC).   Each reaction

contained 4 µL of EvaGreen Mix (Montreal Biotech), 13 µL of UltraPure DNase/RNase-free distilled water

(Life Technologies), 1 µL of 10 mM primers and 2 µL of cDNA (diluted 1:20 from original stock). Primers

were designed using Primer-BLAST (National Institutes of Health) and obtained from Sigma. For each

primer pair, a standard curve was generated using serial dilutions of cDNA prepared from ovarian RNA

and used to determine the efficiency of amplification. Melt-curve analysis and electrophoresis of

amplified products confirmed that only a single product of the expected size was generated. Data was

analyzed using software provided by the manufacturer. Relative quantities of amplified product were

calculated according to 2-ΔΔCT method, using *Rpl19* for normalization.

Immunofluorescence

GOCs and denuded oocytes were fixed for 15 min in freshly prepared 2% (w/v) para-formaldehyde (Fisher

Scientific 04042) in PBS (pH 7.2) containing 0.1% Triton X-100 (ACROS 9002-93-1) (PBST), then washed in

PBST. The specimens were incubated overnight in primary antibody in PBST at 4 °C with gentle agitation,

washed twice in PBST, then incubated for 1 h at room temperature in the secondary antibody as well as

phalloidin and DAPI in PBST, then washed in PBST. GOCs or denuded oocytes were placed in a 2 µl drop

of PBS containing 0.3% polyvinylpyrrolidone (Sigma, PVP360) on a glass bottom dish covered with mineral

oil. Images were acquired using a Zeiss LSM880 confocal microscope. Images were obtained as 16-bit RGB

stack files. GOCs were imaged with a Plan Apo 63x/1.4 oil immersion objective and the following lasers

and filters (in parentheses): 488 nm (491–553 nm), 561 nm (571–642) and 405 nm (410–484). Laser

strength ranged between 1.5 and 2% and digital gain was adjusted until the first saturated pixels were

observed for each channel.

EdU incorporation and detection

EdU (5-ethynyl-2'-deoxyuridine) detection was performed on GOCs according to the Click-iT EdU Alexa

Fluor 488 Imaging Kit manufacturer's instructions (Molecular Probes, C10337). Briefly, cells were

incubated in MEM containing 100 μM EdU for 5 hrs to allow for incorporation into newly synthesized DNA,

then fixed, permeabilized, and incubated in Click-iT cocktail reaction for 30 minutes. EdU labeling was

visualized by confocal microscopy and quantified with ImageJ (NIH).

In situ hybridization

GOCs were fixed in freshly prepared 2% paraformaldehyde in phosphate-buffered saline (PBS, pH 7) for

15 min at room temperature, dehydrated by passing through a gradient of ice-cold methanol (50%, 75%,

100%) for 3 minutes each, then sealed and stored at 4°. The next day, the GOCs were rehydrated using

the reverse methanol gradient (75%, 50%, PBS) for 3 minutes each. Up to 10 GOCs were placed in the

middle of a glass microscope slide, encircled by a hydrophobic barrier created using an Immedge pen, and

incubated with protease solution (ACD, 322337) for 20 min. Slides were rinsed 3 times with PBS and

processed as indicated by the manufacturer for the BaseScope Assay (Advanced Cell Diagnostics (ACD),

323971). Briefly, slides were incubated with the *Smad4* probe specific for the exon8 allele (ACD, 857571)

or negative control probe against *Bacillus subtilis DapB* (ACD, 310043) at 40◦C for 2 h in a HybEZ oven.

Subsequent amplification and detection steps were performed as recommended by the manufacturer and

slides were taken for image acquisition using a Zeiss LSM 880 confocal microscope.

Fluorescence-activated cell sorting

GOCs were dissected from the ovaries of 12-15 d-old *Amhr2-Cre$^+$; mTmG$^+$; Smad4$^{fl/fl}$* mice or were

collected after tamoxifen treatment and culture in the case of *ER-Cre+; mTmG+ Smad4$^{fl/fl}$* mice. GOCs

were dissected from *Amhr2-Cre$^-$* and *ER-Cre-* littermates were used as negative controls of EGFP

expression**.** They were transferred to a 4-well dish containing Ca-free MEM (Thermo Fisher, 21068028)

supplement with 0.25% trypsin (Thermo Fisher, 25200056), 1mM EDTA (Sigma) and DNAse I for several

minutes. After pipetting up and down for a few times to dissociate the cells, fetal bovine serum was added

(4% final concentration) to inactivate the trypsin. Cells were centrifuged at 1500xg for 5 minutes and

supernatant was carefully removed under the dissecting microscope to better visualise the pellet. FACs

buffer containing PBS, 1%FBS, 1mM EDTA and 1mg/ml of DAPI was used to resuspend the pellet. Cells

were resuspended and passed through a Flowmi® 40-μm strainer (Merck,BAH136800040) to remove

oocytes and cell clumps, and then kept on ice for sorting. Resulting granulosa cells were analyzed on a BD

FACSAria Fusion 8.0.2 sorter at the Immunophenotyping Facility of the RI-MUHC. All events were collected

for each sample. A 488-nm excitation laser was used for both Tomato and EGFP excitation since the

wavelength of 488 nm falls within the peak excitation of EGFP but only at 25% of peak excitation for

tdTomato. Data were analyzed using FlowJo FACS analysis software. Cells were gated for singlets (using

forward scatter vs. side scatter area), viability (DAPI scatter vs. side scatter area)], and fluorescence (FITC

and PE) without compensation.

Measurement of granulosa cell-oocyte communication using fluorescence recovery after

photobleaching (FRAP)

A protocol for FRAP was adapted from our previous study (El-Hayek & Clarke, 2015). GOCs were incubated

for 15 min in MEM supplemented with calcein-AM and then for 60 min in calcein-free MEM to allow

transfer of calcein from the granulosa cells into oocyte. A circular region of interest of fixed diameter (10

μm, to ensure the bleaching did not extend to the surrounding granulosa cells) was set in the center of

the enclosed oocyte and subjected to 60 repetitive bleaches of 50 iterations (1 min total duration) using

the argon laser at 50% transmission strength. The intensity of fluorescence in the oocyte and the layer of

granulosa cells immediately surrounding it was recorded before bleaching and after the 10th, 20th, 30th,

40th, 50th, and 60th bleaching. Fluorescence was calculated as intensity of fluorescence in granulosa cell

layer at each time point divided by the pre-bleach intensity of the layer. As oocytes of all experimental

groups were subjected to the same bleaching procedure within the fixed ROI, the loss in fluorescence was

further divided by initial pre-bleach intensity of the oocyte normalized to intensity of fluorescence in

oocyte at the given time point, in order to correct for differences in bleaching effect in oocytes of different

sizes.

Generation of granulosa cell-oocyte reaggregates

GOCs were obtained from 12-day *ER/Cre+; mTmG+; Smad4+/+* and *ER-Cre+; mTmG+; Smad4fl/fl* mice,

exposed to tamoxifen for 24 hr and incubated for an additional five days as described above. They were

then incubated in Ca-Mg-free PBS and drawn in and out of a mouth-controlled micropipette to generate

a suspension of denuded oocytes surrounded by the zona pellucida and individual or small clumps of

granulosa cells. Briefly, the granulosa cells were deposited together with oocytes from CD-1 mice into the

base of a 200-µl microfuge tube and PBS was added to a total volume of 100 µl. The tube was spun at top

speed in a microcentrifuge for one minute, then rotated 180° and centrifuged for one minute. This step

was repeated twice more. The tube was then cut near the base and the cell-pellet was carefully scooped

out and incubated for 2 days in 4-well plates in MEM supplemented with FSH, estradiol and ITS. At the

end of the incubation period, complexes of oocytes surrounded by granulosa cells were fixed and

processed for analysis.

Image analysis

To quantify the number of F-actin-stained-TZPs, a confocal optical section was obtained at the equatorial

plane of the oocyte. Using Fiji software (National Institutes of Health, Bethesda, MD), a segmented circle

was drawn around the oocyte in the middle of the zona pellucida, and the fluorescence intensity at each

point on the line was obtained. Each point whose value was above the background value of the oocyte

cytoplasm and higher than each of its immediately neighboring points was counted as a TZP.

To quantify EdU incorporation in granulosa cells, Fiji was used to calculate the ratio of Edu-labelled cells

per GOC by dividing the number of EdU-positive cells by the total number of cells counterstained with

DAPI.

To individually segment granulosa cells and their corresponding TZPs, Imaris 9.7.2 Software (Bitplane) was

used. First, z-stacks of 0.2 μm thickness covering the middle region of a follicle were imaged to obtain a

conventional 2D Maximum Intensity Projection (MIP). Then, a 3D rendered segmentation of GFP-positive

granulosa cells was created with the "Cell detection" tool where cell boundaries were detected based on

GFP membrane staining. Smallest diameter was set to 100μm, membrane detail of 0.2 μm and the

threshold was adjusted based on local contrast. Segmented cells with a volume less than 100μm^2 were

deleted and fusing or fragmented cells were edited using the "merge" or "split" tools to match the original

image. The oocyte was also segmented using the phalloidin red channel. After edition, cells were

converted to surface objects with the tool "Convert to surfaces" to record their measurements. To render

individual TZPs, the "Filament tracer" tool and semi-automated tracing with the tool "Autopath" was used

to draw the path of each TZP from the granulosa cell body (start) to the oocyte surface (end)

(Supplemental Figure 1A). TZP diameter (thickness) was calculated automatically with a minimum

diameter of 0.1 μm. To determine the distance of the TZP tip to the oocyte surface, the segmented TZP

was automatically divided into discrete spots using the extension "Filament to Spots". This allowed the

calculation of the distance from each spot to the oocyte surface. All measurements were exported using

the "distance to oocyte surface" from each TZP spot. To determine the TZP orientation (away or towards

the oocyte), the distance of the spot located at the TZP start was subtracted from the distance of the spot

located at the TZP tip. A positive difference indicates that the TZP tip is closer to the oocyte and a negative

difference indicates that the TZP tip is farther from the oocyte (Supplemental Figure 1B).

Statistical analysis

Data are presented as mean ± standard error of mean and were analyzed and represented using GraphPad

Prism 8.4.2 software. Statistical significance was evaluated using one- or two-sample two-tailed t-tests.

Acknowledgements

We thank Drs. Shibo Feng and Min Fun (RI-MUHC Imaging Platform) and Marie-Hélène Lacombe, Hélène

Pagé-Veillette and Dr. Ekaterina Iourtchenko (RI-MUHC Immunophenotyping Platform) for their

invaluable assistance. We acknowledge the core facility CELLIM supported by MEYS CR (LM2023050

Czech-BioImaging) for their generous support and assistance in obtaining image analysis data using Imaris,

and thank our colleagues for advice and discussion. Supported by grants to H.J.C. from the Canadian

Institutes of Health Research and the Natural Sciences and Engineering Research Council. H.J.C. gratefully

acknowledges the support of the Richard Cruess Chair in Reproductive Biology at McGill University.

Author contributions

Conceptualization: S.G.A. and H.C.

Methodology: S.G.A., Q.Y. and H.C.

Investigation: S.G.A. and Q.Y.

Funding acquisition: H.C.

Supervision: S.G.A. and H.C.

Writing – original draft: S.G.A. and H.C.

Writing – review and editing: All authors have read and agreed to the final version of the manuscript.

Conflict of interest

The authors declare no conflicts of interest.

References

AASHAQ, S., BATOOL, A., MIR, S. A., BEIGH, M. A., ANDRABI, K. I. & SHAH, Z. A. 2022. TGF-β signaling: A
recap of SMAD-independent and SMAD-dependent pathways. *Journal of Cellular Physiology,*
237**,** 59-85.
ALBERTINI, D. F. & RIDER, V. 1994. Patterns of intercellular connectivity in the mammalian cumulus-
oocyte complex. *Microsc Res Techn,* 27**,** 125-133.
ARAGÓN, E., WANG, Q., ZOU, Y., MORGANI, S. M., RUIZ, L., KACZMARSKA, Z., SU, J., TORNER, C., TIAN, L.,
HU, J., SHU, W., AGRAWAL, S., GOMES, T., MÁRQUEZ, J. A., HADJANTONAKIS, A. K., MACIAS, M.
J. & MASSAGUÉ, J. 2019. Structural basis for distinct roles of SMAD2 and SMAD3 in FOXH1
pioneer-directed TGF-β signaling. *Genes Dev,* 33**,** 1506-1524.
BAENA, V. & TERASAKI, M. 2019. Three-dimensional organization of transzonal projections and other
cytoplasmic extensions in the mouse ovarian follicle. *Sci Rep,* 9**,** 1262.
CARABATSOS, M. J., ELVIN, J., MATZUK, M. M. & ALBERTINI, D. F. 1998. Characterization of oocyte and
follicle development in growth differentiation factor-9-deficient mice. *Dev Biol,* 204**,** 373-384.
CARABATSOS, M. J., SELLITTO, C., GOODENOUGH, D. A. & ALBERTINI, D. F. 2000. Oocyte-granulosa cell
heterologous gap junctions are required for the coordination of nuclear and cytoplasmic meiotic
competence. *Dev Biol,* 226**,** 167-179.
CLARKE, H. J. 2018. Regulation of germ cell development by intercellular signaling in the mammalian
ovarian follicle. *Wiley Interdiscip Rev Dev Biol,* 7**,** 10.1002/wdev.294.
CLARKE, H. J. 2022. Transzonal projections: Essential structures mediating intercellular communication
in the mammalian ovarian follicle. *Mol Reprod Dev,* 89**,** 509-525.
CROZET, F., LETORT, G., BULTEAU, R., DA SILVA, C., EICHMULLER, A., TORTORELLI, A. F., BLÉVINAL, J.,
BELLE, M., DUMONT, J., PIOLOT, T., DAUPHIN, A., COULPIER, F., CHÉDOTAL, A., MAÎTRE, J. L.,
VERLHAC, M. H., CLARKE, H. J. & TERRET, M. E. 2023. Filopodia-like protrusions of adjacent
somatic cells shape the developmental potential of oocytes. *Life Sci Alliance,* 6**,** e202301963.
DERYNCK, R. & BUDI, E. H. 2019. Specificity, versatility, and control of TGF-β family signaling. *Sci Signal,*
12**,** eaav5183.
DOHERTY, C. A., AMARGANT, F., SHVARTSMAN, S. Y., DUNCAN, F. E. & GAVIS, E. R. 2022. Bidirectional
communication in oogenesis: a dynamic conversation in mice and Drosophila. *Trends Cell Biol,*
32**,** 311-323.
DONG, J., ALBERTINI, D. F., NISHIMORI, K., KUMAR, R. M., LU, N. & MATZUK, M. 1996. Growth
differentiation factor-9 is required during early ovarian folliculogenesis. *Nature,* 383**,** 531-535.
DRAGOVIC, R. A., RITTER, L. J., SCHULZ, S. J., AMATO, F., THOMPSON, J. G., ARMSTRONG, D. T. &
GILCHRIST, R. B. 2007. Oocyte-secreted factor activation of SMAD 2/3 signaling enables initiation
of mouse cumulus cell expansion. *Biol Reprod,* 76**,** 848-857.
EL-HAYEK, S. & CLARKE, H. J. 2016. Control of oocyte growth and development by intercellular
communication within the follicular niche. *Res Probl Cell Differ,* 58**,** 191-224.
EL-HAYEK, S., YANG, Q., ABBASSI, L., FITZHARRIS, G. & CLARKE, H. J. 2018a. Mammalian oocytes locally
remodel follicular architecture to provide the foundation for germline-soma communication.
Current Biology, 28**,** 1124-1131.
EL-HAYEK, S., YANG, Q. & CLARKE, H. J. 2018b. Growth in vitro of granulosa cell-oocyte complexes of the
mouse. *Methods Mol Biol,* 1818**,** 1-11.

ELVIN, J., YAN, C., WANG, P., NISHIMORI, K. & MATZUK, M. 1999a. Molecular characterization of the
follicular defects in the growth differentiation factor 9-deficient ovary. *Mol Endocrinol,* 13**,** 1018-
1034.
ELVIN, J. A., CLARK, A. T., WANG, P., WOLFMAN, N. M. & MATZUK, M. M. 1999b. Paracrine actions of
growth differentiation factor-9 in the mammalian ovary. *Mol Endocrinol,* 13**,** 1035-48.
EPPIG, J. J. 1979. A comparison between oocyte growth in coculture with granulosa cells and oocytes
with granulosa cell-oocyte junctional contact maintained in vitro. *J Exp Zool,* 209**,** 345-53.
FIERRO-GONZALEZ, J. C., WHITE, M. D., SILVA, J. C. & PLACHTA, N. 2013. Cadherin-dependent filopodia
control preimplantation embryo compaction. *Nat Cell Biol,* 15**,** 1424-33.
GALLOP, J. L. 2020. Filopodia and their links with membrane traffic and cell adhesion. *Semin Cell Dev*
Biol, 102**,** 81-89.
GITTENS, J. E. & KIDDER, G. M. 2005. Differential contributions of connexin37 and connexin43 to
oogenesis revealed in chimeric reaggregated mouse ovaries. *J Cell Sci,* 118**,** 5071-8.
GUGLIELMI, L., HELIOT, C., KUMAR, S., ALEXANDROV, Y., GORI, I., PAPALEONIDOPOULOU, F.,
BARRINGTON, C., EAST, P., ECONOMOU, A. D., FRENCH, P. M. W., MCGINTY, J. & HILL, C. S. 2021.
Smad4 controls signaling robustness and morphogenesis by differentially contributing to the
Nodal and BMP pathways. *Nature Communications,* 12**,** 6374.
HAO, X., YUAN, F., CUI, Y. & ZHANG, M. 2022. Oocyte-secreted factor TGFB2 enables mouse cumulus cell
expansion in vitro. *Mol Reprod Dev,* 89**,** 554-562.
HECKMAN, C. A. & PLUMMER, H. K., 3RD 2013. Filopodia as sensors. *Cell Signal,* 25**,** 2298-311.
HERTA, A.-C., AKIN, N., BILLOOYE, K., SAUCEDO-CUEVAS, L., LOLICATO, F., SEGERS, I., ANCKAERT, E. &
SMITZ, J. 2021. Reversing complete mechanical transzonal projections disruption during mouse
in vitro follicle culture with unaltered oocyte competence†. *Biology of Reproduction,* 104**,** 1373-
1385.
JAFFE, L. A. & EGBERT, J. R. 2016. Regulation of mammalian oocyte meiosis by intercellular
communication within the ovarian follicle. *Ann Rev Physiol,* 79**,** 237-260.
JORGENSEN, J. S. 2013. Defining the neighborhoods that escort the oocyte through its early life events
and into a functional follicle. *Mol Reprod Dev,* 80**,** 960-76.
JORGEZ, C. J., KLYSIK, M., JAMIN, S. P., BEHRINGER, R. R. & MATZUK, M. M. 2004. Granulosa Cell-Specific
Inactivation of Follistatin Causes Female Fertility Defects. *Mol Endocrinol,* 18**,** 953-967.
KANG, Y., LING, J., SUZUKI, R., ROIFE, D., CHOPIN-LALY, X., TRUTY, M. J., CHATTERJEE, D., WANG, H.,
THOMAS, R. M., KATZ, M. H., CHIAO, P. J. & FLEMING, J. B. 2014. SMAD4 regulates cell motility
through transcription of N-cadherin in human pancreatic ductal epithelium. *PLoS One,* 9**,**
e107948.
KESSEL, R. G., TUNG, H. N., ROBERTS, R. & BEAMS, H. W. 1985. The presence and distribution of gap
junctions in the oocyte-follicle cell complex of the zebrafish, Brachydanio rerio. *J Submicrosc*
Cytol, 17**,** 239-53.
LI, Q., PANGAS, S. A., JORGEZ, C. J., GRAFF, J. M., WEINSTEIN, M. & MATZUK, M. M. 2008. Redundant
roles of SMAD2 and SMAD3 in ovarian granulosa cells in vivo. *Mol Cell Biol,* 28**,** 7001-7011.
LI, R. & ALBERTINI, D. F. 2013. The road to maturation: somatic cell interaction and self-organization of
the mammalian oocyte. *Nat Rev Mol Cell Biol,* 14**,** 141-52.
LI, Y., SCHANG, G., WANG, Y., ZHOU, X., LEVASSEUR, A., BOYER, A., DENG, C. X., TREIER, M., BOEHM, U.,
BOERBOOM, D. & BERNARD, D. J. 2018. Conditional deletion of FOXL2 and SMAD4 in
gonadotropes of adult mice causes isolated FSH deficiency. *Endocrinology,* 159**,** 2641-2655.
LITSCHER, E. S. & WASSARMAN, P. M. 2020. Zona pellucida proteins, fibrils, and matrix. *Annual Review*
of Biochemistry, 89**,** 695-715.

MACAULAY, A. D., GILBERT, I., CABALLERO, J., BARRETO, R., FOURNIER, E., TOSSOU, P., SIRARD, M. A.,
CLARKE, H. J., KHANDJIAN, E. W., RICHARD, F. J., HYTTEL, P. & ROBERT, C. 2014. The gametic
synapse: RNA transfer to the bovine oocyte. *Biol Reprod,* 91**,** 90.
MARCHAIS, M., GILBERT, I., BASTIEN, A., MACAULAY, A. & ROBERT, C. 2022. Mammalian cumulus-oocyte
complex communication: a dialog through long and short distance messaging. *J Assist Reprod
Genet,* 39**,** 1011-1025.
MATTILA, P. K. & LAPPALAINEN, P. 2008. Filopodia: molecular architecture and cellular functions. *Nat
Rev Mol Cell Biol,* 9**,** 446-54.
MCGINNIS, L. K. & KINSEY, W. H. 2015. Role of focal adhesion kinase in oocyte-follicle communication.
Mol Reprod Dev, 82**,** 90-102.
MORA, J. M., FENWICK, M. A., CASTLE, L., BAITHUN, M., RYDER, T. A., MOBBERLEY, M., CARZANIGA, R.,
FRANKS, S. & HARDY, K. 2012. Characterization and significance of adhesion and junction-related
proteins in mouse ovarian follicles. *Biol Reprod,* 86**,** 153.
MOTTA, P., MAKABE, S., NAGURO, T. & CORRER, S. 1994. Oocyte follicle cells association during
development of human ovarian follicle. A study by high resolution scanning and transmission
electron microscopy. *Arch Histol Cytol,* 57**,** 369 - 394.
MOTTERSHEAD, D. G., SUGIMURA, S., AL-MUSAWI, S. L., LI, J. J., RICHANI, D., WHITE, M. A., MARTIN, G.
A., TROTTA, A. P., RITTER, L. J., SHI, J., MUELLER, T. D., HARRISON, C. A. & GILCHRIST, R. B. 2015.
Cumulin, an oocyte-secreted heterodimer of the transforming growth factor-beta family, is a
potent activator of granulosa cells and improves oocyte quality. *J Biol Chem,* 290**,** 24007-24020.
NAGAMATSU, G., SHIMAMOTO, S., HAMAZAKI, N., NISHIMURA, Y. & HAYASHI, K. 2019. Mechanical
stress accompanied with nuclear rotation is involved in the dormant state of mouse oocytes. *Sci
Adv,* 5**,** eaav9960.
NORRIS, R. P., RATZAN, W. J., FREUDZON, M., MEHLMANN, L. M., KRALL, J., MOVSESIAN, M. A., WANG,
H., KE, H., NIKOLAEV, V. O. & JAFFE, L. A. 2009. Cyclic GMP from the surrounding somatic cells
regulates cyclic AMP and meiosis in the mouse oocyte. *Development,* 136**,** 1869-1878.
O'BRIEN, M. J., PENDOLA, J. K. & EPPIG, J. J. 2003. A revised protocol for in vitro development of mouse
oocytes from primordial follicles dramatically improves their developmental competence. *Biol
Reprod,* 68**,** 1682-1686.
PANGAS, S. A., LI, X., ROBERTSON, E. J. & MATZUK, M. M. 2006. Premature luteinization and cumulus
cell defects in ovarian-specific Smad4 knockout mice. *Mol Endocrinol,* 20**,** 1406-1422.
PANGAS, S. A. & RAJKOVIC, A. 2015. Follicular Development: Mouse, Sheep, and Human Models. *In:*
PLANT, T. M. & ZELEZNIK, A. J. (eds.) *Knobil and Neill's Physiology of Reproduction (Fourth
Edition).* San Diego: Academic Press.
PATTON, B. K., MADADI, S. & PANGAS, S. A. 2021. Control of ovarian follicle development by TGFβ family
signaling. *Curr Opin Endocr Metab Res,* 18**,** 102-110.
PENG, D., FU, M., WANG, M., WEI, Y. & WEI, X. 2022. Targeting TGF-β signal transduction for fibrosis and
cancer therapy. *Molecular Cancer,* 21**,** 104.
PENG, J., LI, Q., WIGGLESWORTH, K., RANGARAJAN, A., KATTAMURI, C., PETERSON, R. T., EPPIG, J. J.,
THOMPSON, T. B. & MATZUK, M. M. 2013. Growth differentiation factor 9:bone morphogenetic
protein 15 heterodimers are potent regulators of ovarian functions. *Proc Natl Acad Sci U S A,*
110**,** E776-85.
RICHANI, D., DUNNING, K. R., THOMPSON, J. G. & GILCHRIST, R. B. 2021. Metabolic co-dependence of
the oocyte and cumulus cells: essential role in determining oocyte developmental competence.
Human Reproduction Update, 27**,** 27-47.
ROBERT, C. 2021. Nurturing the egg: the essential connection between cumulus cells and the oocyte.
Reprod Fertil Dev, 34**,** 149-159.

SAATCIOGLU, H. D., CUEVAS, I. & CASTRILLON, D. H. 2016. Control of oocyte reawakening by Kit. *PLoS*
Genet, 12**,** e1006215.
SAGAR, PRÖLS, F., WIEGREFFE, C. & SCAAL, M. 2015. Communication between distant epithelial cells by
filopodia-like protrusions during embryonic development. *Development,* 142**,** 665-71.
SCHROEDER, T. E. 1981. Microfilament-mediated surface change in starfish oocytes in response to 1-
methyladenine: implications for identifying the pathway and receptor sites for maturation-
inducing hormones. *J Cell Biol,* 90**,** 362-71.
SIMON, A., GOODENOUGH, D., LI, E. & PAUL, D. 1997. Female infertility in mice lacking connexin 37.
Nature, 385**,** 525-529.
WASSARMAN, P. M. & LITSCHER, E. S. 2018. The mouse egg's zona pellucida. *Curr Top Dev Biol,* 130**,** 331-
356.
WIGGLESWORTH, K., LEE, K. B., O'BRIEN, M. J., PENG, J., MATZUK, M. M. & EPPIG, J. J. 2013.
Bidirectional communication between oocytes and ovarian follicular somatic cells is required for
meiotic arrest of mammalian oocytes. *Proc Natl Acad Sci U S A,* 110**,** E3723-9.
WINTERHAGER, E. & KIDDER, G. M. 2015. Gap junction connexins in female reproductive organs:
implications for women's reproductive health. *Hum Reprod Update,* 21**,** 340-52.
WU, X., CHEN, L., BROWN, C. A., YAN, C. & MATZUK, M. M. 2004. Interrelationship of growth
differentiation factor 9 and inhibin in early folliculogenesis and ovarian tumorigenesis in mice.
Mol Endocrinol, 18**,** 1509-19.
XIE, J., XU, X. & LIU, S. 2023. Intercellular communication in the cumulus–oocyte complex during
folliculogenesis: A review. *Frontiers in Cell and Developmental Biology,* 11.
YAN, C., WANG, P., DEMAYO, J., DEMAYO, F. J., ELVIN, J. A., CARINO, C., PRASAD, S. V., SKINNER, S. S.,
DUNBAR, B. S., DUBE, J. L., CELESTE, A. J. & MATZUK, M. M. 2001. Synergistic roles of bone
morphogenetic protein 15 and growth differentiation factor 9 in ovarian function. *Mol*
Endocrinol, 15**,** 854-66.
YANG, H., WANG, L., ZHAO, J., CHEN, Y., LEI, Z., LIU, X., XIA, W., GUO, L. & ZHANG, H. T. 2015. TGF-β-
activated SMAD3/4 complex transcriptionally upregulates N-cadherin expression in non-small
cell lung cancer. *Lung Cancer,* 87**,** 249-57.
YU, C., ZHANG, Y. L. & FAN, H. Y. 2013. Selective Smad4 knockout in ovarian preovulatory follicles results
in multiple defects in ovulation. *Mol Endocrinol,* 27**,** 966-78.
ZHANG, H., RISAL, S., GORRE, N., BUSAYAVALASA, K., LI, X., SHEN, Y., BOSBACH, B., BRÄNNSTRÖM, M. &
LIU, K. 2014. Somatic cells initiate primordial follicle activation and govern the development of
dormant oocytes in mice. *Curr Biol,* 24**,** 2501-2508.
ZHANG, M., SU, Y. Q., SUGIURA, K., XIA, G. & EPPIG, J. J. 2010. Granulosa cell ligand NPPC and its
receptor NPR2 maintain meiotic arrest in mouse oocytes. *Science,* 330**,** 366-9.
ZHANG, Y., WANG, Y., FENG, X., ZHANG, S., XU, X., LI, L., NIU, S., BO, Y., WANG, C., LI, Z., XIA, G. &
ZHANG, H. 2021. Oocyte-derived microvilli control female fertility by optimizing ovarian follicle
selection in mice. *Nat Commun,* 12**,** 2523.

Figure 1. SMAD4 is efficiently depleted in granulosa cells of GOCs in vitro using tamoxifen-induced Cre

recombinase

A. Immunoblot of primordial follicles, growing follicles, and granulosa cells and oocytes isolated from

antral follicles, probed using anti-SMAD4 and anti-SMAD2/3.

B. Breeding scheme to delete *Smad4* in the granulosa cells using *Amhr2-Cre*. The *mTmG* cassette

enables Cre recombinase activity to be verified through expression of EGFP.

C. Left: Confocal images of histological sections of paraffin-embedded 12-d ovary (i, ii) and an isolated

follicle (iii) stained using an anti-GFP antibody and DAPI to counterstain DNA. Relatively few

granulosa cells are GFP-positive (arrow in iii). Scale bars: 50 µm, 20 µm, 10 µm. Right: Histogram and

chart showing the percentage of FACS-sorted GFP+ and GFP- granulosa cells from growing follicles.

Granulosa cells expressing both td tomato and GFP and GFP alone were scored as GFP+.

D. Breeding scheme to delete *Smad4* in vitro using generate tamoxifen-inducible *ER-Cre*. The *mTmG*

cassette enables Cre recombinase activity to be verified through expression of EGFP.

E. Left: Schematic illustrating time-line of GOC exposure to tamoxifen and subsequent incubation.

Center: Bright-field image of GOCs following incubation and confocal image of a representative GOC

stained using anti-GFP antibody and DAPI to counterstain DNA. Scale bar = 10µm and 100µm,

respectively. Right: Bar chart showing the percentage of FACS-sorted GFP+ and GFP- granulosa cells

from GOCs. Granulosa cells expressing td tomato and GFP and GFP alone are counted as GFP+.

F. Left: Confocal image of representative GOCs stained with single-molecule RNA FISH probe for *Smad4*

exon 8 and DAPI to counterstain DNA. Right: Mean and SEM of the average number of *Smad4* mRNA

foci per cell (left) and relative mRNA amounts of *Smad4* quantified using qRT-PCR (right) in granulosa

cells five days after tamoxifen treatment.

G. Immunoblot and plot of mean and SEM of SMAD4 protein content in granulosa cells from GOCs five

days after tamoxifen treatment.

Statistical analysis using t-test (F, left) and one-sample t-test (F, right and G), ***P< 0.001; ****P <

0.0001.

Figure 2. Depletion of SMAD4 reduces TZP-number but does not affect granulosa cell communication

with the oocyte or proliferation.

A.   Confocal images of GOCs five days after tamoxifen treatment stained with anti-GFP, phalloidin to

visualize F-actin and DAPI to counterstain DNA. Arrow shows GFP-labelled TZPs. Scale bar = 20 μm.

B.   Mean and SEM of oocyte diameter and TZP number five days after tamoxifen treatment. TZP

number in each GOC is normalized to the oocyte diameter.

C.   Confocal image of a denuded GOC five days after tamoxifen treatment and removal of the

granulosa cells, stained with anti-GFP, phalloidin to visualize F-actin and DAPI to counterstain DNA.

Arrow shows phalloidin-labelled TZPs that remain embedded in the zona pellucida. Scale bar =

20μm.

D.   Mean and SEM of oocyte diameter and TZP number five days after tamoxifen treatment in denuded

GOCs. TZP number for each specimen is normalized to the oocyte diameter.

E.   Left: Time course of fluorescence recovery in oocyte of GOCs following bleaching. Each line

represents an individual GOC. Right: Mean and SEM of the ratio of the final to pre-bleach

fluorescence in each GOC.

F.   Left: Confocal image of a GOC after tamoxifen treatment followed by treatment with EdU and

visualization. DNA stained using DAPI. Scale bar = 20 μm. Right: Mean and SEM of the ratio of Edu-

labelled cells to the total number of cells.

Statistical analysis using t-test, *P< 0.05; ****P < 0.0001; ns, not significant.

Figure 3. Depletion of SMAD4 in GOCs reduces the number of TZPs per granulosa cell and alters their

morphometric parameters.

A.   Schematic illustrating the experimental protocol. GFP+ granulosa cells in clusters were used for

calculations in C and individual GFP+ granulosa cells were used for calculations in D.

B.   Three-dimensional reconstruction of a GOC, showing granulosa cell bodies, TZPs and oocyte cortex.

Serial optical confocal sections were processed using Imaris software. GOCs were stained using

anti-GFP (green), phalloidin (red) and DAPI (blue). Scale bar = 10 μm (upper, middle), 3 μm (lower).

C.   Mean and SEM of volumes and sphericities of individual granulosa cells from a cluster and those

selected for TZP analyses.

D.   Mean and SEM of volumes of individual granulosa cells selected for TZP analysis, number of TZPs

per cell, TZP length and width.

Statistical analysis using t-test, ****$P < 0.0001$; ns, non-significant.

Figure 4. Depletion of SMAD4 in reaggregates reduces the number of newly generated TZPs and alters

their morphometric parameters.

A. Schematic illustrating the reaggregation procedure.

B. Three-dimensional reconstruction of a reaggregate, showing granulosa cell bodies, TZPs and oocyte

cortex. Serial optical confocal sections were processed using Imaris software. GOCs were stained

using anti-GFP (green), phalloidin (red) and DAPI (blue). Scale bar =10 μm (upper), 3 μm (lower).

C. Mean and SEM of volumes and sphericities of individual granulosa cells.

D, E Mean and SEM of volumes of individual granulosa cells, number of TZPs per cell, TZP length and TZP

width.

F. Upper: Schematic illustrating how the distance from the origin and the tip, respectively, of TZPs

(blue) and non-TZP projections (red) to the oocyte surface was determined. Lower: TZP trajectories

of 3D-reconstructed TZPs, representing the distance from origin relative to the position on the TZP

axis.

G. Left: Mean and SEM of the difference between the distance from the origin and the tip of individual

TZPs to the oocyte surface. Blue symbols (>0) represent TZPs oriented towards the oocyte; red

symbols (<0) represent non-TZPs oriented away from the oocyte. Right: Mean and SEM of the

proportion of TZPs in individual cells that were oriented away from the oocyte.

Statistical analysis using t-test, *P < 0.05; **P < 0.01; ****P < 0.0001; ns, non-significant.

Figure 5. Expression of transmembrane N-cadherin and Notch proteins is reduced in SMAD4-depleted

granulosa cells.

A.   Immunoblot and graph showing mean and SEM of SMAD4 and N-cadherin in granulosa cells.

Protein quantities are normalized to amounts in ER-Cre⁻ cells.

B.   As in A, showing Notch2.

C.   Mean and SEM of relative mRNA amounts of *Smad4*, *Cdh2* (encoding N-cadherin), and *Notch2* in

granulosa cells purified from GOCs five days after tamoxifen treatment. mRNA quantities are

normalized to amounts in ER-Cre⁻ cells.

D.   As in C, showing *Smad4*, *Myo10, Fscn1* and *Daam1*.

Statistical analysis using one-sample t-test, ***P < 0.001; ****P < 0.0001; ns non-significant.

Figure 6. Model for regulation of the size of the TZP population.

Upper: Dynamics of an individual TZP. When growing TZP contacts the surface of the oocyte, it becomes

stably attached if there is sufficient N-cadherin, as in wild-type cells. In SMAD4-depleted cells, there is

insufficient N-cadherin to stabilize the attachment. The TZP may continue to elongate but eventually

becomes retracted. Lower: Dynamics of the population of TZPs. Owing to increased stability of

attachment of individual TZPs to the oocyte surface, the total number of TZPs becomes larger in wild-

type cells than in SMAD4-depleted cells.

Figure 1

Figure 2

Figure 3

A

B

ER-Cre⁺; Smad4⁺/⁺   *ER-Cre⁺; Smad4^{fl/fl}*

C

D

Figure 4

Figure 5

Figure 6