Immigration delays but does not prevent adaptation following

environmental change: experimental evidence

Authors:

Lily F. Durkee[1,2*], Laure Olazcuaga[2,3], Brett A. Melbourne[4], and Ruth A. Hufbauer[1,2*]

[1] Graduate Degree Program in Ecology, Colorado State University, Fort Collins, Colorado 80523,

USA

[2] Department of Agricultural Biology, Colorado State University, Fort Collins, Colorado 80523,

USA

[3] Centre National de la Recherche Scientifique (CNRS), Station d'Ecologie Théorique et

Expérimentale (UAR2029), Moulis, France

[4] Department of Ecology & Evolutionary Biology, University of Colorado, Boulder, Colorado

80309, USA

*Corresponding author: L.Durkee@colostate.edu; Ruth.Hufbauer@colostate.edu

Running head: **Evolutionary rescue with immigration**

Abstract

In today's rapidly changing world, it is critical to examine how animal populations will

respond to severe environmental change. Following events such as pollution or deforestation

that cause populations to decline, extinction will occur unless populations can adapt in

response to natural selection, a process called evolutionary rescue. Theory predicts that

immigration can delay extinction and provide novel genetic material that can prevent

inbreeding depression and facilitate adaptation. However, when potential source populations

have not experienced the new environment before (*i.e.,* are naive), immigration can counteract

selection and constrain adaptation. This study evaluated the effects of immigration of naive

individuals on evolutionary rescue using the red flour beetle, *Tribolium castaneum,* as a model

system. Small populations were exposed to a challenging environment, and three immigration

rates (zero, one, or five migrants per generation) were implemented with migrants from a

benign environment. Following an initial decline in population size across all treatments,

populations receiving no immigration gained a higher growth rate one generation earlier than

those with immigration, illustrating the constraining effects of immigration on adaptation. After

seven generations, a reciprocal transplant experiment found evidence for adaptation regardless

of immigration rate. Thus, while the immigration of naive individuals briefly delayed adaptation,

it did not increase extinction risk or prevent adaptation following environmental change.

Keywords: evolutionary rescue, dispersal, *Tribolium castaneum*, gene flow, translocation

1 | Introduction

A critical goal of conservation biology is to establish how to effectively manage

populations in human-altered habitats. Environmental change such as pollution (Loria *et al.*,

2019), deforestation (Foster *et al.*, 2021), and fragmentation (Cheptou *et al.*, 2017) can reduce

habitat quality and connectivity, which can decrease the average fitness of residing

populations. If fitness drops below the replacement rate, populations will decline in size and

will approach extinction if the downward trend persists. Immigration, either through

maintenance of dispersal corridors or translocations, is one approach readily available to

conservation practitioners to support declining populations, particularly in fragmented

landscapes (Frankham *et al.*, 2019).

Theoretical work within the conservation literature recommends an immigration rate of

one to ten migrants per generation for isolated populations (Mills & Allendorf, 1996; Frankham

et al., 2019). More specifically, theory suggests a rate of one migrant per generation to offset

population divergence and loss of heterozygosity that occurs through drift (Mills & Allendorf,

1996), and five migrants per generation to prevent damaging inbreeding depression and

maintain adaptive genetic variation in changing environments (Frankham *et al.*, 2019). Ideally,

immigration can occur consistently through dispersal corridors, and if translocations are

needed, consistent immigration should be sustained. However, these recommendations have

been tested only partially. Laboratory experiments have demonstrated the benefits of

immigration when comparing isolated populations to populations that received a single

immigration event under scenarios of inbreeding (Spielman & Frankham, 1992) or low diversity

populations facing environmental change (Hufbauer *et al.*, 2015). Only one study we know of,

Newman & Tallmon (2001), tested immigration that occurred every generation. They assessed

rates of one or 2.5 migrants per generation in recently isolated mustard plant populations, and

found that the rate of one migrant per generation was sufficient to maintain genetic variation

and decrease inbreeding when compared to isolated populations (Newman & Tallmon, 2001).

In this era of global change, it is important to also test these recommendations under more

severe environmental conditions.

A separate but related body of literature focuses on how populations declining due to

exposure to challenging environments can avoid extinction through adaptive evolution, a

process called evolutionary rescue (Gomulkiewicz & Holt, 1995; Carlson *et al.*, 2014). Examples

include adaptation in weed or insect populations to pesticides, antibiotic resistance in bacteria,

or invertebrate adaptation to metal pollution in streams (Bell, 2017). Experiments show that

immigration or admixture can enhance the probability of evolutionary rescue if populations are

exposed to a challenging environment (Hufbauer *et al.*, 2015; Durkee *et al.*, 2023).

Furthermore, theoretical studies show that the probability of rescue can increase with

immigration, primarily due to the added genetic variation, which increases the chance of

beneficial alleles being present (Tomasini & Peischl, 2020; Czuppon *et al.*, 2021). To contrast,

other theoretical studies show that gene flow from maladapted immigrants can limit

adaptation by swamping adaptive alleles in the recipient population, reducing fitness (*i.e.,*

outbreeding depression) (Slatkin, 1987; Lenormand, 2002; Brady *et al.*, 2019).

To reduce the likelihood of outbreeding depression, using a source population from a

habitat similar to the recipient population (*i.e.,* habitat matching) is recommended by models

(Edelaar & Bolnick, 2012) and in the context of conservation (Frankham *et al.*, 2019). However,

this is often not possible, particularly under scenarios of novel environmental change (Aitken &

Whitlock, 2013) or when the species of concern is rare (Frankham *et al.*, 2019). Even when

migrants are not adapted to their new habitat, for example when individuals from a

geographically distant population (*e.g.,* Hedrick & Fredrickson, 2010) or from captivity (*e.g.,*

Crone *et al.,* 2007) are used, every-generation immigration of one to five individuals is meant to

be low enough to minimize outbreeding depression while still maintaining genetic diversity and

adaptive potential of populations (Newman & Tallmon, 2001; Frankham *et al.*, 2019). However,

to our knowledge, no study has compared the recommended immigration rates of one and five

migrants per generation and their impacts on adaptation in the context of severe

environmental change.

Here, we evaluated how immigration rates recommended in the conservation literature

(Frankham *et al.*, 2019) impacted evolutionary rescue (Carlson *et al.*, 2014). To do this, we first

created experimental populations of red flour beetles (*Tribolium castaneum*) from a genetically

admixed stock population (Durkee *et al.,* 2023). We then introduced them to a challenging

environment to mimic a situation where a large, diverse population experiences sudden habitat

change that decreases population size and reduces habitat quality, for example when forest are

logged (Hillers *et al.,* 2008). Each experimental population received one of three immigration

treatments (zero, one, or five individuals every generation), and we observed the effects of

immigration on population persistence and growth. In habitats like these that have limited

space and degraded resources, population recovery may be constrained by density-dependent

processes like competition for food and space (Osmond & de Mazancourt, 2013; Nordstrom *et*

al., 2023). Thus, we evaluated both density-dependent and density-independent population

growth through time to tease apart the impacts of negative density dependence and

adaptation. Our experiment was designed to address the following question: In populations

experiencing severe environmental change, how do different immigration rates affect *(i)*

population size and growth rates over time, *(ii)* the timing of evolutionary rescue, and *(iii)*

adaptation to the challenging environment?

2 | Methods

To evaluate the question posed in the introduction, we carried out an experiment where

we subjected replicated, independent populations to a challenging environment that caused

them to decline in size. This environment, therefore, would have led to extinction unless the

populations were able to adapt sufficiently to the challenging conditions. Hereafter, we refer to

this as the rescue experiment (Figure 1a-b).

Rescue experiment initiation and propagation

We created a genetically diverse *Tribolium castaneum* stock population comprised of

lineages described in Durkee *et al.* (2023). The stock population had been reared on standard

media (19:1 wheat flour to brewer's yeast) and in standard environmental conditions (31°C and

50-80% humidity) in 4x4x6cm patches for nine, 35-day non-overlapping generations prior to the

start of the rescue experiment. To start, we created 25 independent populations from the stock

population of 300 individuals each (Figure 1a). These populations were used to initiate

replicated experimental populations and maintain populations to serve as a source of migrants

(hereafter, source populations). These populations were divided at the start of the rescue

experiment, and therefore represent newly isolated populations with no history of divergence,

similar to Newman & Tallmon (2001).

Experimental populations were each initiated with 50 adult beetles, 36 days old, and

population sizes were allowed to fluctuate naturally in subsequent generations. Each

population was placed onto a single patch of challenging environment created using media

containing the pyrethroid insecticide deltamethrin (DeltaDust, Bayer). This event represents a

sudden environmental change that reduced fitness and created replicated, isolated

populations. The experimental patches each contained a concentration of deltamethrin that a

dose-response curve created from preliminary experiments (Figure S1) indicated would

decrease the population growth rate to approximately 0.40 from a typical value of around 2.

Without adaptation, this reduced growth rate should have led to extinction in 10 generations

(based on simulations with the stochastic Ricker model developed for *T. castaneum* by

Melbourne & Hastings, 2008). The experimental populations (N=50) were initiated in 25 groups

of three, each group from one of the original 25 independent populations (Figure 1a).

Source populations were initiated with 150 adults, 36 days old, and each served as a

source of migrants for a group of three experimental populations (Figure 1a). Thus, 25 source

populations were initiated in total, each from one of the original 25 populations. We intended

for the sources to represent larger populations, and so we used the large population size

(N=150) from Hufbauer *et al.* (2015). Sources were maintained at 150 adults every generation

regardless of the number of offspring produced, to approximate populations with a carrying

capacity of 150 individuals. Source populations were each maintained on a single patch

containing standard dry media, a benign environment for *T. castaneum*. Thus, these individuals

were not exposed to the challenging environment present in the experimental patches.

Hereafter, we refer to these individuals as naive migrants, to refer to both evolutionary and

behavioral naivety.

The adult beetles in the experimental and source patches were given 24 hours to mate

and oviposit. Adults were then were discarded, and the eggs developed for an additional 34

days. The first census event (N_1, Figure 1b) occurred on day 35. Subsequently, one of three

immigration treatments was applied to each experimental population in every group: zero, one,

or five migrants. The migrants were moved from the source to each experimental patch

manually using a paintbrush. This occurred at the start of the following oviposition period. This

procedure (immigration of zero, one, or five migrants, 24-hour oviposition, removal of adults,

34-day development period, census) repeated each subsequent generation. The experiment

was terminated after seven generations because nearly all the experimental populations had

either gone extinct or their growth rates and population sizes had increased, indicative of

evolutionary rescue (see Figure 2).

In summary, there were 25 groups, each consisting of one source and three

experimental populations (one of each treatment). Thus, there were 25 replicates of each

treatment, which we split across three temporal blocks (eight groups in blocks 1 and 2, nine in

block 3; Figure 1a). Within each group, treatments were set up sequentially, and propagated for

the first four generations in the following order to minimize opportunities for experimental

error: 1-migrant, 5-migrants, and then the 0-migrant populations. As this was not random, we

tested if this order mattered in a follow-up experiment and found no significant effect of

treatment order on the growth rate the following generation (Figure S2). During generations 4

through 7, the experimental protocol was well established, and so we randomized the order of

set-up.

Source population dispersal arrays

The immigration recommendations we followed (one or five migrants per generation)

are based on either natural dispersal or translocation. Thus, in order to approximate natural

dispersal and make our results applicable to either case, we used migrants with higher

demonstrated dispersal ability to represent individuals that would be more likely to leave the

source patch and move to a new patch in the absence of translocation (Edelaar & Bolnick, 2012;

Cote *et al.*, 2022). Dispersal arrays were created to select dispersing individuals. On day 34

during generations 1-6, habitat patches housing source populations were connected to 3-patch

dispersal arrays connected with 2mm holes, creating a 4-patch dispersal array. Hereafter, we

refer to the original source patch as patch 1 and the subsequent patches, in order, as patches 2-

4, which contained media that had been previously used to rear beetles for 35 days to keep

resource availability consistent throughout the array. Adult beetles were given 24 hours to

disperse out of patch 1. On day 35, all adults were censused in each of the four patches of the

dispersal array (Figure S3). Individuals that dispersed to patches 2-4 were used as migrants

(hereafter called dispersing migrants) with preference given to individuals that dispersed to the

furthest patch (*i.e.,* individuals were first chosen from patch 4, then patch 3, then 2).

In order to maintain the population size of the source populations at 150 to start every

generation, growth rates needed to be above the replacement rate. However, growth rates

were sometimes lower than expected, which likely occurred due to effects of demographic

stochasticity, which strongly influences the *T. castaneum* system (Melbourne & Hastings, 2008).

When this happened, populations were supplemented with individuals from the stock

population at the start of the oviposition period, after the dispersal period and immigration to

the experimental populations occurred. After the first generation, two source populations were

initialized with 110 individuals due to limited availability of stock individuals.

Common environment and reciprocal transplant experiment

After seven generations, we terminated the rescue experiment, and all extant

populations (58 experimental and 25 source populations) were placed at a constant density of

40 individuals on standard media for a single generation (hereafter, "common environment"

generation; Figure 1c). All adults were kept within their experimental populations, *i.e.,* no

mixing occurred within or across treatments. In *T. castaneum,* maternal environmental effects

can strongly affect growth rate (Hufbauer *et al.,* 2015; Van Allen & Rudolf, 2013), and so this

generation where the environment and density were kept the same for all populations served

to standardize maternal effects for the subsequent reciprocal transplant experiment. With only

a single generation, we do not standardize grand-maternal effects. However, inference from

multigenerational studies suggests that grand maternal effects are relatively weak in *T.*

castaneum populations (Van Allen & Rudolf, 2013; Hufbauer *et al.,* 2015).

In the ninth generation, a reciprocal transplant experiment was completed to assess

adaptation to the challenging environment and potential costs to adaptation in the source

environment (Figure 1d). We divided individuals from each population into three groups and

randomly assigned them to one of three environments: standard media (0ppm, hereafter

"source environment"), media containing the concentration of deltamethrin present in the

challenging environment of the rescue experiment (2.968ppm, hereafter "experimental

environment"), and media containing a higher concentration of deltamethrin (5.641ppm,

hereafter "severe environment"). The dose-response curve (Figure S1) indicated that the severe

environment would reduce the growth rate of naive populations to near zero. Each replicate

patch of these three environmental treatments received 40 beetles, which were allowed 24

hours for oviposition, and 34 additional days for development prior to census. Replication

within population depended on the number of available individuals following the common

environment generation (see Figure S4).

Figure 1. Experimental design for the rescue and reciprocal transplant experiments. a) We

began with a large stock population ($N \approx 2500$), from which eight or nine independent

populations were initiated ($N=300$). Each of these populations was split into one source

population, initiated and then maintained at $N=150$ individuals, and three experimental

populations, initiated with $N_0=50$ individuals with population size allowed to fluctuate

thereafter. The source populations were maintained on a benign environment (tan shading),

and the experimental populations were maintained in a challenging environment containing

the insecticide deltamethrin (light gray shading). **b)** In the rescue experiment, census events

(gens 1 & 7 shown) are indicated with three cartoon beetles, and census occurred every

generation on day 35, following a 24-hour oviposition period and 34 days of development

(indicated with dotted arrows). Immigration from the source to the experimental populations

(dashed arrows) occurred after each census event at the start of the oviposition period, and

three treatments were implemented: 0 migrants (purple square outlines), 1 migrant (yellow

diamond outlines), and 5 migrants per generation (blue circle outlines). **c)** All extant

populations were placed on a benign, common environment for a single generation following

the census in generation seven at a density of N=40, with as many replicates created as

possible. **d)** In the reciprocal transplant experiment, all populations were split into groups of 40

and placed simultaneously and at constant densities on three different environments: benign,

challenging, and severe (dark gray shading). This whole procedure **(a-d)** was repeated across

three temporal blocks, with eight replicates in blocks 1 and 2 and nine replicates in block 3.

Data analysis

All statistical analyses were performed in R version 4.1.2 (R Core Team, 2021).

Population size at census, which occurred on day 35 during generations 1-7, was used to assess

population dynamics throughout the rescue experiment. Populations were considered rescued

if they grew to a population size greater than 50, and maintained a growth rate greater than 1

(the replacement rate) for at least two of the final four generations. This allowed for decreases

in population size due to negative density dependence. Extinction risk and survival times were

quantified using a survival analysis with the *ggsurvfit* function in the *survival* package (Therneau

2021). Extinct populations were defined as those with zero adults at census during the first six

generations or one adult by generation seven. Overall extinction rates were calculated as raw

percentages with confidence intervals around each percentage for each treatment.

Population growth rate was calculated as the mean number of surviving adult offspring

per individual in the population, *i.e.,* population size at census each generation (N_t) divided by

the number of parents that produced those individuals (parent population size, N_{t-1}), or N_t/N_{t-1}.

We evaluated both density-dependent and density-independent growth rates through time

using a linear mixed model with the *lme4* package (Bates *et al.*, 2015). We included immigration

treatment (0 migrants, 1 migrant, or 5 migrants per generation), time (t) in generations (from 1

to 7 generations, categorical), and parent population size (N_{t-1}, integers) as fixed effects, and

temporal block (block 1, block 2, block 3), source population (nested within block), and

population identity (nested within source population) as random effects. Growth rate data

were *log(x+1)* transformed to help satisfy the equal variance assumption. Diagnostic plots were

visualized for all models using the *DHARMa* package (Hartig 2022).

From this model, we generated estimates of both density-dependent and density-

independent population growth rates using the *predict* function with different values for the

parent population size (N_{t-1}) each generation. To estimate density-dependent growth rates, we

allowed the values for N_{t-1} to vary for each generation and treatment combination, matching

the observed mean population sizes at the start of each generation. This estimate was used to

describe the population growth rate data and quantify the effects of immigration treatment in

a finite habitat by considering the constraints of competition for resources and cannibalism that

affected the populations in our experiment (Pointer *et al.*, 2021). To estimate density-

independent growth rates from the fitted model, we set N_{t-1} to zero for every generation and

treatment, which is equivalent to estimating the y-intercepts when plotting model-estimated

growth rate over N_{t-1} (Figure S5). These density-independent estimates allowed us to track

adaptation to the environment by revealing the change in intrinsic fitness of individuals in the

population unfettered by the effects of density. We expected the density-independent growth

rates to be higher than the density-dependent estimates due to the known negative density

dependence in *T. castaneum* (Melbourne & Hastings 2008). Confidence intervals for both

density-dependent and density-independent growth rate estimates were estimated by

parametric bootstrap (percentile method) using the *bootMer* function with 5000 simulations in

the *lme4* package (Bates *et al.*, 2015). Comparisons between treatments were made by

estimating confidence intervals by parametric bootstrap (percentile method) for the difference

in means between treatments within each generation using the *simpleboot* and *boot* packages

with 5000 simulations (Peng, 2019; Canty & Ripley, 2022).

The timing of evolutionary rescue was assessed in each treatment group using a

Generalized Additive Mixed (GAM) model (Wood, 2017) similar to Olazcuaga *et al.* (2023). Only

rescued populations were considered in this analysis. We modeled the population size of

rescued populations using a negative binomial distribution to account for overdispersion using

the package *mgcv* (Wood, 2004, 2011). To allow for the relationship between the population

size and time to be nonlinear, we modeled population size as a smooth function of generation

with the number of knots set to 4, which was chosen to maximize the adjusted R-squared value.

Explanatory variables also included immigration treatment and the interaction of treatment

and generation as fixed effects, allowing for a smooth function for each treatment. The full

model included population identity and temporal blocks as random effects. The best-fit model

was chosen by comparing to reduced models using the *compareML* function from the *itsadug*

package (van Rij *et al.*, 2022).

We used data from the reciprocal transplant experiment to further evaluate adaptation

to the experimental environment, as well as potential fitness costs of adaptation in a benign

environment. Growth rate in the reciprocal transplant experiment was square-root

transformed and used as the response variable in a linear mixed effects model that included

population type (source, 0-migrant, 1-migrant and 5-migrant), test environment (source,

experimental, severe), and the interaction between population type and test environment as

fixed effects. Population identity was included as a random effect nested within temporal block.

Comparisons within each environment and among immigration treatments were made using

95% bootstrapped confidence intervals (percentile method) for the difference in means, which

were generated from 5000 iterations.

3 | Results

Population size and extinction

During the rescue experiment, a majority of populations experienced evolutionary

rescue (Figure 2, black lines), and extinction probability was similar for all three treatments

(Figure 2, red lines). The difference in the extinction risk was statistically similar across

treatments throughout the experiment (Figure S6). Additionally, there was no clear difference

in survival time among treatments ($\chi_2^2 = 0.1$, $p = 1.00$).

Figure 2. Population size through time in discrete generations for the populations receiving 0,

1, or 5 migrants per generation during the rescue experiment. Population sizes represent the

number of adult beetles present at census (day 35 each generation). Each black line represents

a rescued population, each orange line represents an extant population that was not

considered rescued, and each red line represents a population that went extinct or had a

population size of one individual in generation seven. Immigration events are indicated with

instantaneous increases in population size, which occurred at the start of the first six

generations. The percent of populations that went extinct and simple confidence intervals

around each percent are provided.

Growth rate through time

Density-dependent growth rates increased through generation four or five, then

decreased again as population density increased (Figure 3a, Table S1). Density-independent

growth rates were below or close to the replacement rate (growth rate of 1) in generations 1-3

(Figure 3b; confidence intervals overlap with dashed line). This changed in generation four,

when the 0-migrant treatment had a mean density-independent growth rate much larger than

the replacement rate, illustrating rapid adaptation within those populations (Figure 3b). In

generation four, density-independent growth rate in the 0-migrant populations was on average

0.808 (95% CI [0.207, 1.453]) higher than the 1-migrant populations and 0.969 [0.478, 1.573]

higher than the 5-migrant populations. By generation five, all treatments gained and then

maintained a mean density-independent growth rate greater than 1, demonstrating adaptation

to the challenging environment. The comparison between the density-independent and

density-dependent estimates of growth rates suggest that the decrease in the latter by

generation seven was likely due to the effects of negative density dependence, which increased

as populations grew.

Figure 3. Density-dependent growth rate (a) and density-independent growth rate (b) throughout the

rescue experiment. Each point outlined in black is the predicted mean for each treatment (purple

square = 0 migrants, yellow diamond = 1 migrant per generation, and blue circle = 5 migrants per

generation). Error bars are bootstrapped 95% confidence intervals. Points without black outlines are raw

growth rate values, and each point represents one population. The dotted line represents a growth rate

of 1, the replacement rate. Two estimates for growth rate are presented: a) the density-dependent

estimates are predictions from the linear mixed model for each generation and treatment combination,

used to incorporate the effects of negative density-dependence and provide estimates that describe the

raw data (shown), and b) the density-independent estimates (*i.e.,* predictions using N_{t-1} = 0), showing

adaptation over time.

Timing of evolutionary rescue

Among the rescued populations only (black lines in Figure 2), the analysis of population

size with a GAM model revealed a shallow U-shaped curve (Figure 4). The estimated curves

371   from the best-fit model, which included treatment and the interaction of treatment and

372   generation as fixed effects and population identity as a random effect, are shown in Figure 4.

373   The interaction between immigration treatment and generation was significant (

 , $p <$

0.0001), with the 0-migrant populations growing slightly faster than populations receiving

migrants (Figure 4). The 1-migrant and 5-migrant treatments maintained similar population

trajectories throughout the experiment.

Figure 4. Evolutionary rescue analysis. Population size over time (in generations) of rescued

populations receiving 0 migrants (purple line, squares), 1 migrant (yellow line, diamonds), or 5

migrants (blue line, circles) per generation estimated by the best-fit GAM model. Each point

represents one population, which are staggered to reduce overlap. The lines are estimates of

mean population size for the three immigration treatments, and the shaded region around each

line is the 95% confidence interval given by the GAM model.

Adaptation and associated costs

Growth rate in the reciprocal transplant experiment was influenced by an interaction

between environment and population type ($F_{6, 168.04} = 25.16$, $p < 0.0001$), in a pattern showing

that adaptation to the pesticide was linked to reduced fitness in the benign source environment

(Figure 5). In the benign environment, the experimental populations had lower growth rates on

average compared to the source populations (confidence intervals are negative and do not

overlap with zero: 95% $CI_{0\text{-migrant} - source}$ [-0.855, -1.628]; 95% $CI_{1\text{-migrant} - source}$ [-0.911, -1.748]; 95%

$CI_{5\text{-migrant} - source}$ [-0.641, -1.482]), and a higher growth rate in the severe habitat (confidence

intervals are positive and do not overlap with zero: 95% $CI_{0\text{-migrant} - source}$ [0.442, 0.171]; 95% $CI_{1\text{-}}$

$_{migrant - source}$ [0.424, 0.210]; 95% $CI_{5\text{-migrant} - source}$ [0.464, 0.216]). Additionally, the 5-migrant

populations performed marginally better than the 0-migrant populations in the experimental

environment (95% $CI_{0\text{-migrant} - 5\text{-migrant}}$ [-0.0675, 0.572]).

Unexpectedly, the experimental and source populations performed similarly in the

challenging experimental environment (Figure 5, all confidence intervals of the differences

overlap with zero: 95% $CI_{0\text{-migrant} - source}$ [-0.404, 0.209]; 95% $CI_{1\text{-migrant} - source}$ [-0.220, 0.403]; 95%

$CI_{5\text{-migrant} - source}$ [-0.153, 0.452]). The experimental populations did not perform as well as

expected based on their growth rates at the end of the rescue experiment (Figure 3). Our

expectation for growth rates were based on predictions from the linear mixed model created

for the rescue experiment with N_{t-1} = 40 (the number of parents used in the reciprocal

transplant) and gen = 7 (the time closest to the start of the reciprocal transplant). Those

predicted values were 1.80, 95% CI [1.32, 2.39], 1.52 [1.12, 2.00], and 1.75 [1.31, 2.26] for the

0-migrant, 1-migrant, and 5-migrant treatments, respectively. The average growth rates we

observed in the reciprocal transplant experiment in the experimental environment were on

average 40% lower than these estimates, with the means 0.85 [0.64, 1.09], 1.04 [0.80, 1.32],

1.10 [0.86, 1.35].

412   Furthermore, source populations did not perform as poorly as expected in the

experimental environment. We expected them to perform similarly to the first generation of

the experimental populations in the rescue experiment prior to adaptation. The prediction

given N_{t-1} = 40 was 0.63 [0.38, 0.83] for the 0-migrant populations at the start of the rescue

experiment, but the mean growth rate of source populations was about 46% higher at 0.92

[0.72, 1.15].

Figure 5. Reciprocal transplant experiment results. Growth rate was measured in three

environments: the source environment (standard dry media), the challenging experimental

environment, and a severe environment containing a higher concentration of insecticide. Each

point outlined in black is the model-estimated mean for each population type (purple squares =

0-migrant populations, yellow diamonds = 1-migrant populations, blue circles = 5-migrant

populations, and green triangles = source populations). Black error bars are bootstrapped 95%

confidence intervals; for the source populations in the severe environment (green triangles),

the black error bars are not visible because they are too small to be seen. Thick lines connecting

these points emphasize the treatment-environment interaction. Each point without the black

outline represents a single population, with the light error bars representing raw 95%

confidence intervals across replicates; points without error bars have no replicates. Points are

staggered and jittered to reduce overlap.

4 | Discussion

Our study evaluated the effects of different immigration rates on adaptation to a new,

challenging environment using experimental populations of *Tribolium castaneum*. We

evaluated differences between initially declining populations receiving two immigration rates

(one or five migrants per generation) with isolated populations that received no immigration.

The migrants were not adapted to the challenging environment, allowing us to examine the

degree to which immigration can limit adaptation (Savolainen *et al.*, 2013). While the

experimental populations declined in size initially, leading some populations to go extinct,

around 80% of populations were able to persist for the duration of the seven-generation

experiment. Most of these populations had growth rates above the replacement rate and had

large population sizes, characteristic of evolutionary rescue. We also found that extinction rate

was similar regardless of immigration. Notably, growth rates increased one generation sooner

in the isolated populations compared to those with immigration. This delay was brief, as the

following generation, growth rates were comparable across the different immigration

treatments. We also tested for adaptation using a reciprocal transplant experiment. While

some of the results from that experiment were unexpected, overall it provided evidence that

our experimental populations were adapted to the insecticide that created the challenging

environment. There was also a cost to that adaptation, evident from the experimental

populations having lower growth rates than the sources in the benign environment.

Observed delay in the increase in growth rate: possible mechanisms

Populations without immigration gained a much higher density-independent growth

rate (more than doubling on average) compared to the populations with immigration (near or

just above the replacement rate) in the fourth experimental generation. The following

generation, all populations grew at a similar rate regardless of immigration. The delay we

observed may have been due to *i)* differences in density dependence, *ii)* phenotypic plasticity,

or *iii)* outcrossing with naive migrants slowing adaptation to the challenging environment.

First, theory shows that reduced negative density dependence in a degraded habitat

may enhance rescue (Uecker *et al.*, 2014; Czuppon *et al.*, 2021). In our system, population

densities were low in the experimental populations in the initial generations, and thus negative

density-dependent processes were likely relatively weak. The added immigrants increased

density slightly relative to the treatment without migrants, and this could have contributed to

lower growth rates. However, if this were the case, the populations receiving five migrants each

generation should have experienced further reductions in growth than the populations

receiving one per generation, as their density was increasing by a greater amount (31% increase

for the 5-migrant populations on average compared to a 7% increase for the 1-migrant

populations). Thus, density dependence is an unlikely explanation for the one-generation delay

in increased growth rates.

Second, growth rate in the challenging environment could be influenced by phenotypic

plasticity, which could have led to the lag in increased growth rates. Recently, it has been

shown that plasticity in insects can be mediated by associated microbes (Kolodny &

Schulenburg, 2020). Variation in the microbial communities in host insects is high (Lange *et al.*,

2023), and the microbial community can evolve, which can facilitate persistence in new and

challenging environments (Kolodny & Schulenburg, 2020). Microbes, therefore, can serve as an

important source of adaptive phenotypic plasticity (Kolodny & Schulenburg, 2020; Ghalambor

et al., 2007). Insect-associated microbes can be transmitted from parent to offspring (Lange *et*

al., 2023), indicating that plasticity could be multi-generational. If the microbial community

influences fitness in the presence of deltamethrin, it is possible that plasticity and adaptation in

the microbiome helped facilitate the observed increase in growth rate during the rescue

experiment. The delay in increase for populations with immigration could be due to the

microbiome from the benign environment being continually re-introduced by the migrants.

Third, migration is well known as a constraint to adaptation (Slatkin, 1987). For example,

theory has shown that immigration between different habitat types can limit local adaptation

through genetic swamping of adaptive alleles (Lenormand, 2002; Bolnick & Nosil, 2007), and

reduce the probability (Schiffers *et al.*, 2013) or speed (Holt & Barfield, 2015) of evolutionary

rescue because of maladaptation in the new environment. It seems likely that migration of

naive individuals slowed adaptation to the novel environment, which may have been enhanced

by the mating behavior of *T. castaneum*. In this species, males reared in high-quality

environments have increased mating rate and insemination success compared to males reared

in poor environments (Lewis *et al.*, 2012). Thus, male migrants from the benign source

environment may have successfully mated with multiple females in the experimental patches,

enhancing the rate of gene flow of maladapted alleles. Female migrants would have also likely

spread maladapted alleles, as a single female *T. castaneum* can lay over 10 eggs per day (Pray

et al., 1949). Thus, regardless of sex, outcrossing with the naive migrants likely contributed to

the brief delay in growth rate increase. Given the strength of selection pressure, however,

adaptation likely enabled growth rates to increase in the following generation, which we

observed on average across all treatments.

We suggest that the increases in population size and growth rate, likely through

adaptation and possibly aided by adaptive plasticity, can be described as evolutionary rescue.

We further evaluate the effects of adaptation *vs.* plasticity when we discuss the reciprocal

transplant experiment below. The finding that rescue can occur with intermediate levels of

immigration of not adapted individuals is supported by several modeling studies that assess

evolutionary rescue in heterogenous, structured systems (Gomulkiewicz *et al.*, 1999; Uecker *et*

al., 2014; Tomasini & Peischl, 2020; Czuppon *et al.*, 2021). This result has also been shown in a

case study of Trinidadian guppies, where rescue was able to occur with gene flow from a

maladapted source population (Fitzpatrick *et al.,* 2014, 2020).

Observed delay in rescue: implications

Our finding of only a brief (one generation) delay of adaptation has practical

implications. Such delayed adaptation could be detrimental to populations at imminent risk of

extinction due to small size (Gomulkiewicz & Holt, 1995; Melbourne & Hastings, 2008).

However, for populations not at immediate risk of extinction, such a delay may be

inconsequential, and worth the potential increase in genetic diversity that occurs through

immigration. Such increased diversity may prevent inbreeding depression from developing, and

improve the ability of the population to adapt to future environmental changes (Frankham *et*

al., 2019).

Thus, these results suggest that naive migrants can be suitable to use in management

efforts. Furthermore, they are often more feasible to obtain, either from captive rearing

programs (Crone *et al.,* 2007) or from large source populations living in a different habitat (*e.g.,*

different host plants in Bisschop *et al.,* 2019) or geographic region (*e.g.,* isolated guppy

populations in Fitzpatrick *et al.,* 2020), when habitat matching is not possible.

Additionally, our findings are consistent with refuge strategies to delay evolution of pest

resistance to toxins, such as in crops genetically modified to express *Bacillus thuringiensis* (*Bt*).

The immigration of susceptible individuals from sections of fields planted in non-*Bt* crops, or

growing crops in areas with plentiful alternative host plants for the pest has proven to be highly

effective (Carrière *et al.,* 2012). In those situations, particularly with 50% or more of a pest

population being susceptible, resistance can be successfully managed for multiple generations

(Tabashnik *et al.,* 2008). Given our much smaller proportion of susceptible individuals, it makes

sense that the delay of the evolution of resistance was brief.

Reciprocal transplant experiment: interpretation of unexpected results

The two most straightforward takeaways from the reciprocal transplant experiment are

i) the experimental populations had higher growth rates than the source populations in the

severe environment, suggesting they had adapted to the insecticide used to create the

challenging environment in the rescue experiment and *ii)* the experimental populations had

lower growth rates in the benign environment than the source populations, suggesting that

there was a cost associated with adaptation to the insecticide. However, the source populations

had higher than expected growth rates in the experimental environment, and the experimental

populations had lower than expected growth rates in the experimental environment

inconsistent with the hypothesis of adaptation. We next discuss *i)* how experimental challenges

might have affected the source populations, *ii)* how the common environment generation could

have reduced growth rates by eliminating effects of plasticity, and *iii)* how multiple pathways of

adaptation in the experimental populations could explain their performance in the reciprocal

transplant experiment.

First, the high performance of the source populations in the experimental environment

could be due to challenges in implementing the experimental design. For instance, accidental

contamination of the source population with deltamethrin, or accidental gene flow from the

experimental to the source populations could have occurred during the rescue experiment. We

took great care to avoid both forms of contamination, but if contamination nevertheless

occurred, it could have allowed the source populations to adapt to low concentrations of the

insecticide. The source populations were not adapted to high concentrations of the pesticide,

as evidenced by their performance in the severe environment.

A second consideration is that the common environment generation could have masked

differences in growth rates, if they were initially caused by plasticity. As described above,

microbiome-mediated phenotypic plasticity could have contributed to the increase in growth

rate in the rescue experiment (Lange *et al.,* 2023). If microbiome-mediated plasticity drove

performance, the generation in a common environment prior to the reciprocal transplant

experiment would have alleviated selection pressure and reduced the ability of the

experimental populations to tolerate the insecticide. This could help explain the similar

performance of the experimental and source populations in the experimental environment.

However, there does not appear to be strong evidence for the masking of phenotypic plasticity

in the severe or benign environments given that the experimental populations had a much

higher growth rate than the sources in the severe environment and a lower growth rate in the

benign. We suggest that those differences can be explained by our last point, adaptation, which

is discussed below.

Finally, adaptation to deltamethrin likely explains the relative performance of the

populations in the three environments. More specifically, it is possible that the experimental

populations adapted through more than one mechanism during the rescue experiment. First,

adaptation to detoxify the insecticide compound likely occurred, as rapid evolutionary

responses to deltamethrin is well documented in pest populations of *T. castaneum* (Collins,

1998; Stuart *et al.*, 1998) through the increased expression of cytochrome P450 enzymes (Kalsi

& Palli, 2015). In other insect species, this mechanism of resistance to deltamethrin has a

confirmed cost (Tchouakui *et al.*, 2020), and such a cost is consistent with the reduced

performance of experimental populations in the benign environment. The second way that

populations could have adapted is by increasing the rate of egg cannibalism to supplement

their diets, as cannibalism propensity is a genetically based trait that can evolve (Stevens, 1989)

that has been shown to enhance fitness in challenging environments (Via, 1999; Pointer *et al.*,

2021). These potential pathways, evolution of increased detoxification and increased

cannibalism, are not mutually exclusive, and both may have facilitated increased population

growth in the experimental environment during the rescue experiment. Then, in the reciprocal

transplant experiment, the populations were initiated at lower density (40 parents) in the three

environments. In this lower density environment, the number of eggs available for cannibalizing

was reduced compared to what was available in the latter half of the rescue experiment (in

generations 4-7, N_{t-1} > 150 on average). Thus, the decreased density reduced the eggs available

for cannibalism, limiting one of the adaptive pathways and reducing overall growth rates of the

experimental populations in the experimental environment. In the severe environment, growth

rates of the experimental populations were low but still greater than the source populations,

which can be explained primarily by their evolved ability to detoxify the pesticide.

Considerations for future experiments

593   The bottleneck events that occurred during the rescue experiment, some as small as 10

individuals, would likely have longer-term consequences (Frankham *et al.*, 1999; Olazcuaga *et*

al., 2023). Census population size in a wide range of natural populations is almost always larger

than the effective population size (the median N_e/N_c ratio across plant and animal taxa is

approximately 0.1; for *T. castaneum,* it is estimated to be between 0.5-0.9) and bottlenecks of

N_e < 50 can result in reductions in genetic variation and increases in inbreeding (Wade, 1980;

Jamieson & Allendorf, 2012). For this reason, we suggest that future work examine the effects

of demographic bottlenecks of this severity by propagating experiments like this for longer (*i.e.,*

more than 8 generations). Longer-term experiments would be able to test the hypothesis that

the benefits of immigration become more apparent over longer time scales. Such experiments

would help shed light on the longer-term impacts of immigration for small and isolated

populations, an important question when considering whether to implement translocations in

declining natural populations (Frankham, 2015, 2016). Genomic methods would also provide

important insights for monitoring the genetic contributions of migrants, the genetic

differentiation between the experimental and source populations, and for confirming the genes

contributing to adaptation (Whiteley *et al.*, 2015). Genomics would also allow us to more

clearly differentiate between adaptation and plasticity (Koch & Guillaume, 2020).

Conclusions

This study explored the effects of immigration of naive individuals on large, outbred,

and recently diverged populations following population decline due to sudden environmental

degradation. To our knowledge, we are the first to experimentally test the effects of the

recommended immigration rates of one compared to five migrants per generation on

adaptation in a challenging environment. We found that though immigration slowed adaptation

in the short-term, extant populations with and without immigration were on average equally

well adapted to the challenging conditions after seven generations. Thus, we confirm the

findings of previous theoretical work (Tomasini & Peischl, 2020; Czuppon *et al.*, 2021) by

demonstrating experimentally that evolutionary rescue can occur both with and without

immigration following severe environmental change.

Data Availability Statement

Data and R scripts are publicly available on the Dryad Digital Repository

(https://doi.org/10.5061/dryad.79cnp5j3c).

Conflict of Interest

The authors declare no conflicts of interest.

References

Bates, D., Maechler, M., Bolker, B. & Walker, S. 2015. Fitting linear mixed-effects models using
lme4. *J Stat Softw* 67: 1–48.

Bell, G. 2017. Evolutionary rescue. *Annu Rev Ecol Evol Syst* 48: 447–71.

Bisschop, K., Mortier, F., Etienne, R.S. & Bonte, D. 2019. Transient local adaptation and source-
sink dynamics in experimental populations experiencing spatially heterogeneous environments.
Proceedings of the Royal Society B: Biological Sciences 286.

Bolnick, D.I. & Nosil, P. 2007. Natural selection in populations subject to a migration load.
Evolution 61: 2229–2243.

Brady, S.P., Bolnick, D.I., Angert, A.L., Gonzalez, A., Barrett, R.D.H., Crispo, E., *et al.* 2019. Causes
of maladaptation. *Evol Appl* 12: 1229–1242.

Canty, A. & Ripley, B. 2022. boot: Bootstrap R Functions.

Carlson, S.M., Cunningham, C.J. & Westley, P.A.H. 2014. Evolutionary rescue in a changing
world. *Trends in Ecology & Evolution* 29: 521-530.

Carrière, Y., Ellers-Kirk, C., Hartfield, K., Larocque, G., Degain, B., Dutilleul, P., *et al.* 2012. Large-
scale, spatially-explicit test of the refuge strategy for delaying insecticide resistance. *Proc Natl*
Acad Sci 109: 775–780.

Cheptou, P.O., Hargreaves, A.L., Bonte, D. & Jacquemyn, H. 2017. Adaptation to fragmentation:
Evolutionary dynamics driven by human influences. *Philosophical Transactions of the Royal*
Society B: Biological Sciences 372.

Collins, P.J. 1998. Inheritance of resistance to pyrethroid insecticides in *Tribolium castaneum*
(Herbst). *J Stored Prod Res* 34: 395–401.

Cote, J., Dahirel, M., Schtickzelle, N., Altermatt, F., Ansart, A., Blanchet, S., *et al.* 2022. Dispersal
syndromes in challenging environments: A cross-species experiment. *Ecology Letters* 25: 2675-
2687.

Crone, E.E., Pickering, D. & Schultz, C.B. 2007. Can captive rearing promote recovery of
endangered butterflies? An assessment in the face of uncertainty. *Biol Conserv* 139: 103–112.

Czuppon, P., Blanquart, F., Uecker, H. & Débarre, F. 2021. The effect of habitat choice on
evolutionary rescue in subdivided populations. *American Naturalist* 197: 625–643.

Durkee, L.F., Olazcuaga, L., Szymanski, R., Melbourne, B.A. & Hufbauer, R.A. 2023. Genetic
mixing facilitates adaptation to a novel environmental constraint. *Ecol Entomol* 48: 517–522.

Edelaar, P. & Bolnick, D.I. 2012. Non-random gene flow: An underappreciated force in evolution
and ecology. *Trends Ecol Evol* 27: 659–665.

Fitzpatrick, S. W., Gerberich, J. C., Kronenberger, J. A., Angeloni, L. M., & Funk, W. C. 2015.
Locally adapted traits maintained in the face of high gene flow. *Ecology Letters* 18: 37–47.

Fitzpatrick, S.W., Bradburd, G.S., Kremer, C.T., Salerno, P.E., Angeloni, L.M. & Funk, W.C. 2020.
Genomic and fitness consequences of genetic rescue in wild populations. *Current Biology* 30:
517-522.e5.

Foster, B.J., McCulloch, G.A., Vogel, M.F.S., Ingram, T. & Waters, J.M. 2021. Anthropogenic
evolution in an insect wing polymorphism following widespread deforestation. *Biol Lett* 17.

Frankham, R. 2016. Genetic rescue benefits persist to at least the F3 generation, based on a
meta-analysis. *Biol Conserv* 195: 33–36.

Frankham, R. 2015. Genetic rescue of small inbred populations: meta-analysis reveals large and
consistent benefits of gene flow. *Mol Ecol* 24: 2610–2618.

Frankham, R., Ballou, J.D., Ralls, K., Eldridge, M., Dubash, M.R., Fenster, C.B., *et al.* 2019. *A*
Practical Guide for Genetic Management of Fragmented Animal and Plant Populations, 1st ed.
Oxford University Press, Oxford, UK.

Frankham, R., Lees, K., Montgomery, M.E., England, P.R., Lowe, E.H. & Briscoe, D.A. 1999. Do
population size bottlenecks reduce evolutionary potential? *Anim Conserv* 2: 255–260.

Ghalambor, C.K., McKay, J.K., Carroll, S.P. & Reznick, D.N. 2007. Adaptive versus non-adaptive
phenotypic plasticity and the potential for contemporary adaptation in new environments.
Funct Ecol 21: 394–407.

Gomulkiewicz, R. & Holt, R.D. 1995. When does evolution by natural selection prevent
extinction? *Evolution* 49: 201.

Gomulkiewicz, R., Holt, R. D., & Barfield, M. (1999). The Effects of Density Dependence and
Immigration on Local Adaptation and Niche Evolution in a Black-Hole Sink Environment. In
Theoretical Population Biology (Vol. 55).

Florian Hartig (2022). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
Regression Models. R package version 0.4.6. https://CRAN.R-project.org/package=DHARMa

Hedrick, P.W. & Fredrickson, R. 2010. Genetic rescue guidelines with examples from Mexican
wolves and Florida panthers. *Conservation Genetics* 11: 615–626.

Hillers, A., Veith, M., & Rödel, M. O. (2008). Effects of forest fragmentation and habitat
degradation on West African leaf-litter frogs. *Conservation Biology* 22: 762–772.

Holt, R.D. & Barfield, M. 2015. The influence of imperfect matching habitat choice on evolution
in source–sink environments. *Evol Ecol* 29: 887–904.

Hufbauer, R.A., Szücs, M., Kasyon, E., Youngberg, C., Koontz, M.J., Richards, C., *et al.* 2015.
Three types of rescue can avert extinction in a changing environment. *PNAS* 112: 10557–10562.

Jamieson, I.G. & Allendorf, F.W. 2012. How does the 50/500 rule apply to MVPs? *Trends Ecol*
Evol 27: 578–584.

Kalsi, M. & Palli, S.R. 2015. Transcription factors, CncC and Maf, regulate expression of CYP6BQ
genes responsible for deltamethrin resistance in *Tribolium castaneum. Insect Biochem Mol Biol*
65: 47–56.

Kawecki, T.J. & Ebert, D. 2004. Conceptual issues in local adaptation. *Ecol Lett* 7: 1225–1241.

Koch, E.L. & Guillaume, F. 2020. Restoring ancestral phenotypes is a general pattern in gene
expression evolution during adaptation to new environments in *Tribolium castaneum. Mol Ecol*
29: 3938–3953.

Kolodny, O., & Schulenburg, H. (2020). Microbiome-mediated plasticity directs host evolution
along several distinct time scales. *Philosophical Transactions of the Royal Society B: Biological*
Sciences, 375(1808).

Lange, C., Boyer, S., Bezemer, T. M., Lefort, M. C., Dhami, M. K., Biggs, E., Groenteman, R.,
Fowler, S. V., Paynter, Q., Verdecia Mogena, A. M., & Kaltenpoth, M. (2023). Impact of
intraspecific variation in insect microbiomes on host phenotype and evolution. *ISME Journal* 17:
1798–1807.

Lenormand, T. 2002. Gene flow and the limits to natural selection. *Trends Ecol Evol* 17: 183–
189.

Lewis, S.M., Tigreros, N., Fedina, T. & Ming, Q.L. 2012. Genetic and nutritional effects on male
traits and reproductive performance in *Tribolium* flour beetles. *J Evol Biol* 25: 438–451.

Loria, A., Cristescu, M.E. & Gonzalez, A. 2019. Mixed evidence for adaptation to environmental
pollution. *Evol Appl* 12: 1259–1273.

Melbourne, B.A. & Hastings, A. 2008. Extinction risk depends strongly on factors contributing to
stochasticity. *Nature* 454: 100–103.

Mills, L.S. & Allendorf, F.W. 1996. The one-migrant-per-generation rule in conservation and
management. *Conservation Biology* 10: 1509–1518.

Newman, D. & Tallmon, D.A. 2001. Experimental evidence for beneficial fitness effects of gene
flow in recently isolated populations. *Conservation Biology* 15: 1054–1063.

Nordstrom, S.W., Hufbauer, R.A., Olazcuaga, L., Durkee, L.F., & Melbourne, B.A. (2023). How
density dependence, genetic erosion, and the extinction vortex impact evolutionary
rescue. *Proceedings of the Royal Society B.* 203: 1-12.

Olazcuaga, L., Lincke, B., Delacey, S., Durkee, L.F., Melbourne, B.A. & Hufbauer, R.A. 2023.
Population demographic history and evolutionary rescue: influence of a bottleneck event. *Evol*
Appl 16: 1483–1495.

Osmond, M.M. & de Mazancourt, C. 2013. How competition affects evolutionary rescue.
Philosophical Transactions of the Royal Society B: Biological Sciences 368.

Peng, R.D. 2019. simpleboot: Simple Bootstrap Routines.

Pointer, M.D., Gage, M.J.G. & Spurgin, L.G. 2021. *Tribolium* beetles as a model system in
evolution and ecology. *Heredity* 126: 869–883. Springer US.

Rinkevich, F.D., Leichter, C.A., Lazo, T.A., Hardstone, M.C. & Scott, J.G. 2013. Variable fitness
costs for pyrethroid resistance alleles in the house fly, *Musca domestica*, in the absence of
insecticide pressure. *Pestic Biochem Physiol* 105: 161–168.

Savolainen, O., Lascoux, M. & Merilä, J. 2013. Ecological genomics of local adaptation. *Nat Rev*
Genet 14: 807–820.

Schiffers, K., Bourne, E.C., Lavergne, S., Thuiller, W. & Travis, J.M.J. 2013. Limited evolutionary
rescue of locally adapted populations facing climate change. *Philosophical Transactions of the*
Royal Society B: Biological Sciences 368: 0–9.

Slatkin, M. 1987. Gene flow and the geographic structure of natural populations. *Science (1979)*
236: 787–792.

Spielman, D. & Frankham, R. 1992. Modeling problems in conservation genetics using captive
Drosophila populations: Improvement of reproductive fitness due to immigration of one
individual into small partially inbred populations. *Zoo Biol* 11: 343–351.

Stevens, L. 1989. The genetics and evolution of cannibalism in flour beetles (genus *Tribolium*).
Evolution 43: 169–179.

Stuart, J.J., Ray, S., Harrington, B.J., Neal, J.J. & Beeman, R.W. 1998. Genetic Mapping of a Major
Locus Controlling Pyrethroid Resistance in *Tribolium castaneum* (Coleoptera: Tenebrionidae). *J*
Econ Entomol 91: 1232–1238.

Tabashnik, B.E., Gassmann, A.J., Crowder, D.W. & Carrière, Y. 2008. Insect resistance to Bt
crops: Evidence versus theory. *Nat Biotechnol* 26: 199–202.

Tchouakui, M., Riveron Miranda, J., Mugenzi, L.M.J., Djonabaye, D., Wondji, M.J., Tchoupo, M.,
et al. 2020. Cytochrome P450 metabolic resistance (CYP6P9a) to pyrethroids imposes a fitness

cost in the major African malaria vector Anopheles funestus. *Heredity* 124: 621–632. Springer
US.

R Core Team 2021. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Tomasini, M. & Peischl, S. 2020. When does gene flow facilitate evolutionary rescue? *Evolution*
74: 1640–1653.

Uecker, H., Otto, S.P. & Hermisson, J. 2014. Evolutionary rescue in structured populations.
American Naturalist 183.

Van Allen, B.G. & Rudolf, V.H.W. 2013. Ghosts of habitats past: Environmental carry-over
effects drive population dynamics in novel habitat. *American Naturalist* 181: 596–608.

van Rij, J., Wieling, M., Baayen, R.H. & van Rijn, H. 2022. itsadug: Interpreting Time Series and
Autocorrelated Data Using GAMMs.

Via, S. 1999. Cannibalism facilitates the use of a novel environment in the flour beetle,
Tribolium castaneum. Heredity 82: 267–275.

Wade, M.J. 1980. Effective population size: the effects of sex, genotype, and density on the
mean and variance of offspring numbers in the flour beetle, *Tribolium castaneum. Genet Res*
36: 1–10.

Whiteley, A.R., Fitzpatrick, S.W., Funk, W.C. & Tallmon, D.A. 2015. Genetic rescue to the rescue.
Trends in Ecology & Evolution 30: 42-49.

Wood, S.N. 2011. Fast stable restricted maximum likelihood and marginal likelihood estimation
of semiparametric generalized linear models. *Journal of the Royal Statistical Society* 73: 3–36.

Wood, S.N. 2017. *Generalized Additive Models: An Introduction with R*, 2nd ed. Chapman and
Hall/CRC.

Wood, S.N. 2004. Stable and efficient multiple smoothing parameter estimation for generalized
additive models. *J Am Stat Assoc* 99: 673–686.