Immigration delays but does not prevent adaptation following

environmental change: experimental evidence

Authors:

Lily F. Durkee[1,2*], Laure Olazcuaga[2], Brett A. Melbourne[3], and Ruth A. Hufbauer[1,2*]

[1] Graduate Degree Program in Ecology, Colorado State University, Fort Collins, Colorado 80523,

USA

[2] Department of Agricultural Biology, Colorado State University, Fort Collins, Colorado 80523,

USA

[3] Department of Ecology & Evolutionary Biology, University of Colorado, Boulder, Colorado

80309, USA

*Corresponding author: lilyfdurkee@gmail.com; Ruth.Hufbauer@colostate.edu

Abstract

An important and pressing goal in conservation is to determine how to effectively manage populations experiencing environmental change. When populations begin to decline, extinction will occur unless populations can adapt in response to natural selection, a process called evolutionary rescue. Theory predicts that immigration can delay extinction and provide novel genetic material that may reduce inbreeding depression and facilitate adaptation. However, when potential source populations have not experienced the new environment before (*i.e.,* are naive), immigration can counteract selection and constrain adaptation. This study evaluated the effects of immigration of naive individuals on evolutionary rescue using the red flour beetle, *Tribolium castaneum,* as a model system. Small populations were exposed to a challenging environment, and three immigration rates (0, 1, or 5 migrants per generation) were implemented with migrants from a benign environment. Following an initial decline in population size across all treatments, populations receiving no immigration gained a positive growth rate one generation earlier than those with immigration, illustrating the constraining effects of immigration on adaptation. After seven generations, a reciprocal transplant experiment found evidence for local adaptation regardless of immigration rate. Thus, while the immigration of naive individuals briefly delayed adaptation, it did not increase extinction risk or prevent adaptation to severe and abrupt environmental change.

Keywords: evolutionary rescue, dispersal, *Tribolium castaneum*, gene flow, translocation

1 | Introduction

A critical goal of conservation biology is to establish how to effectively manage

populations in human-altered habitats. Environmental change such as pollution (Loria *et al.*,

2019), deforestation (Foster *et al.*, 2021), and fragmentation (Cheptou *et al.*, 2017) can reduce

habitat quality and connectivity, which can decrease the average fitness of residing

populations. If fitness drops below the replacement rate, populations will decline in size and

will approach extinction if the downward trend persists. Immigration, either through

maintenance of dispersal corridors or translocations, is one approach readily available to

conservation practitioners to support declining populations, particularly in fragmented

landscapes (Frankham *et al.*, 2019).

The conservation literature recommends an immigration rate of one to ten migrants per

generation for isolated populations (Mills & Allendorf, 1996). More specifically, one migrant per

generation aims to offset population divergence that occurs through drift, while five migrants

per generation aims to prevent inbreeding depression and maintain adaptive potential in

changing environments (Frankham *et al.*, 2019). Ideally, immigration can occur consistently

through dispersal corridors, and if translocations are needed, consistent immigration should be

sustained. However, these recommendations have been tested only partially. Laboratory

experiments have demonstrated the benefits of immigration when comparing isolated

populations to populations that received a single immigration event under scenarios of

inbreeding (Spielman & Frankham, 1992) or low diversity populations facing environmental

change (Hufbauer *et al.*, 2015). Only one study we know of, Newman & Tallmon (2001), tested

immigration that occurred every generation. They assessed rates of 1 or 2.5 migrants per

generation in newly-isolated mustard plant populations, and found that the rate of 1

migrant/gen was sufficient to maintain genetic variation and decrease inbreeding when

compared to isolated populations (Newman & Tallmon, 2001). In this era of global change, it is

important to also test these recommendations in more severe environmental conditions.

A separate but related body of literature focuses on how populations declining due to

exposure to challenging environments can avoid extinction through adaptive evolution, a

process called evolutionary rescue (Gomulkiewicz & Holt, 1995; Carlson *et al.*, 2014). Examples

include adaptation in weed or insect populations to pesticides, antibiotic resistance in bacteria,

or invertebrate adaptation to metal pollution in streams (Bell, 2017). Experiments show that

immigration or admixture can enhance the probability of evolutionary rescue if populations are

exposed to a challenging environment (Hufbauer *et al.*, 2015; Durkee *et al.*, 2023). However,

gene flow from maladapted immigrants is well known to limit adaptation, slowing the increase

in frequency of adaptive alleles and reducing fitness (*i.e.,* outbreeding depression) (Slatkin,

1987; Lenormand, 2002; Brady *et al.*, 2019).

To reduce the likelihood of outbreeding depression, habitat matching between source

and recipient populations is recommended (Edelaar & Bolnick, 2012; Frankham *et al.*, 2019).

However, this is often not possible, particularly under scenarios of novel environmental change

(Aitken & Whitlock, 2013) or when the species of concern is rare (Frankham *et al.*, 2019). Even

when migrants are not adapted to their new habitat, for example when individuals from a

geographically distant population (*e.g.,* Hedrick & Fredrickson, 2010) or from captivity (*e.g.,*

Crone et al., 2007) are used, every-generation immigration of 1 to 5 individuals is meant to be

low enough to maintain genetic diversity and adaptive potential of populations (Newman &

Tallmon, 2001; Frankham *et al.*, 2019). However, to our knowledge, no study has tested the

recommended immigration rates of one vs. five migrants per generation experimentally in the

context of severe environmental change.

Here, we evaluated how immigration recommendations from the conservation

literature (Frankham *et al.*, 2019) impacted evolutionary rescue (Carlson *et al.*, 2014) in a

challenging environment. To do this, we used red flour beetle (*Tribolium castaneum*)

populations without a history of inbreeding or bottlenecks to model a situation where large,

diverse populations experience sudden habitat change, causing them to become both small and

fragmented. In habitats like these that have limited space and degraded resources, population

recovery may be constrained by density-dependent processes like competition for food and

space (Osmond & de Mazancourt, 2013; Nordstrom *et al.*, 2023). We exposed these

populations to a new, challenging environment, and then implemented three different rates of

immigration: 0, 1, or 5 individuals each generation from a large population not adapted to the

new environment. We evaluated both density-dependent and density-independent population

growth to tease apart the impacts of negative density dependence and adaptation. Our

experiment was designed to address the following question: In populations experiencing severe

environmental change, how do different immigration rates affect *(i)* population size and growth

rate over time, *(ii)* the timing of evolutionary rescue, and *(iii)* adaptation and the associated

consequences of adapting to the challenging environment?

2 | Methods

To evaluate the question posed in the introduction, we carried out an experiment where

we subjected replicate populations to a challenging environment that caused populations to

decline in size, and therefore would have led to extinction unless populations adapted

sufficiently to the challenging conditions. Hereafter, we refer to this as the rescue experiment.

We created a genetically diverse *T. castaneum* stock population comprised of lineages

described in Durkee et al. (2023). The stock population had been reared on standard media

(19:1 wheat flour to brewer's yeast) and in standard environmental conditions (31°C and 50-

80% humidity) in 4x4x6cm patches for nine non-overlapping generations prior to the start of

the rescue experiment. Each generation was five weeks long, with adult oviposition occurring

on day 35. To start the experiment, we initiated 25 independent populations from the stock

population. These independent populations were used to initiate experimental populations and

maintain populations to serve as a source of migrants (hereafter, source populations).

Rescue experiment initiation and propagation

Experimental populations were each initiated with 50 adults that were placed onto a

single patch of challenging environment created using media containing the pyrethroid

insecticide deltamethrin (DeltaDust, Bayer). These two events together represent a sudden

environmental change causing both habitat degradation and population fragmentation

simultaneously. The experimental patches contained 2.968ppm of deltamethrin, a

concentration that a dose-response curve created from preliminary experiments indicated

would decrease population growth from a typical value of 2, or doubling every generation, to

0.40 (Figure S1). This growth rate guaranteed population extinction in 10 generations without

adaptation or immigration based on simulations with the stochastic Ricker model developed for

T. castaneum by Melbourne & Hastings (2008). Adults were 36 days old at population initiation,

and they were given 24 hours to mate and oviposit in the experimental patches. Adults were

discarded after oviposition, and offspring were then given 35 days to develop. Experimental

populations were propagated for 7 generations in this way, with populations censused

completely each generation on day 35. In total, 75 experimental populations were initiated

across three temporal blocks.

Source populations of migrants were initiated with 150 adults from the stock

population and then maintained at 150 adults each subsequent generation, regardless of the

number of offspring produced. This approximates a population with a carrying capacity of 150

individuals. Source populations were each maintained on a single patch containing standard dry

media, a benign environment for *T. castaneum*. Thus, these individuals were not exposed to the

challenging environment present in the experimental patches. Hereafter, we refer to these

individuals as naive migrants, to refer to both evolutionary and behavioral naivety. We

expected that the source populations would have a growth rate greater than the replacement

rate. However, when this did not occur, populations were supplemented with individuals from

the stock population to maintain the oviposition density at 150 adults per patch. This

supplementation occurred at the start of the oviposition period (day 35), after immigration to

the experimental populations occurred. After the first generation, two source populations were

initialized with 110 individuals due to limited availability of stock individuals. In total, 25 source

populations were maintained total across the three temporal blocks, one for each group of

immigration treatments.

Three immigration rates were used in the rescue experiment: zero, one, or five

migrants every generation. Each generation, experimental populations received migrants from

the source population on day 35 before the oviposition period. Immigration occurred by

moving individuals using a paintbrush from the source population to each experimental patch.

Across the 7 experimental generations, 6 migrants were added to the 1-migrant populations

and 30 migrants were added to the 5-migrant populations in total. Each immigration treatment

was replicated 25 times, in 25 separate groups comprised of three treatments each, split across

three temporal blocks. Within each group, treatments were set up sequentially using the same

pool of adult beetles, and propagated for the first four generations in the following order: 1-

migrant, 5-migrants, and 0-migrants. As this was not random, we tested if this order mattered

in a follow-up experiment and found no significant effect of patch order on the growth rate the

following generation (Figure S2).

The two immigration recommendations of one or five migrants per generation are

based on either natural dispersal or translocation. Thus, in order to approximate natural

dispersal and make our results applicable to either case, we used migrants with higher

demonstrated dispersal ability to represent individuals that would be more likely to leave the

source patch and move to a new patch in the absence of translocation (Edelaar & Bolnick, 2012;

Cote *et al.*, 2022). **Dispersal arrays** were created to select dispersing individuals. On day 34

during generations 1-6, habitat patches housing source populations were connected to 3-patch

dispersal arrays connected with 2mm holes, creating a 4-patch dispersal array. Hereafter, we

refer to the original source patch as patch 1 and the subsequent patches, in order, as patches 2-

4, which contained media that had been previously used to rear beetles for 35 days to keep

resource availability consistent throughout the array. Adult beetles were given 24hrs to

disperse out of patch 1. On day 35, all adults were censused in each of the four patches of the

dispersal array (see Figure S3). Individuals that dispersed to patches 2-4 were used as migrants

(hereafter called dispersing migrants) with preference given to individuals that dispersed to the

furthest patch (*i.e.,* individuals were first chosen from patch 4, then patch 3, then 2).

Reciprocal transplant experiment

After seven generations, we terminated the rescue experiment, and a reciprocal

transplant experiment was completed using the 58 extant experimental populations and the 25

source populations to assess adaptation to the challenging environment and potential costs to

adaptation in the source environment. We measured the growth rate of each population for

one generation in each of three environments: standard media (0ppm, hereafter "source

environment"), media containing the concentration of deltamethrin present in the challenging

environment of the rescue experiment (2.968ppm, hereafter "experimental environment"), and

media containing a higher concentration of deltamethrin (5.641ppm, hereafter "severe

environment"). Using the dose-response curve created in preliminary experiments (Figure S1),

this severe concentration was intended to reduce the growth rate of naive populations to zero.

To standardize maternal and environmental effects, all extant populations were placed

at a constant density of 40 individuals on standard media for one generation during generation

8 (hereafter, "common garden" generation). For three populations, a density of fewer than 40

individuals was used due to small population size ($N < 50$) at the conclusion of the rescue

experiment. All adults were kept within their experimental populations, *i.e.,* no mixing occurred

within or across treatments.

The reciprocal transplant experiment occurred in generation 9 using the adult offspring

from the common garden generation, and replication depended on the number of available

individuals (see Figure S6). The growth rate of each experimental population was measured in

each of the three test environments (source, experimental, and severe) with an initial

population size of 40. In total, there were 23-24 replicates of populations from the 0-migrant

populations (average replication of 1.1/population) across the three environments, 19-20

replicates (average replication of 1.2/population) from the 1-migrant populations, 22 replicates

for the 5-migrant treatment (average replication of 1.1/population), and 24-25 replicates for

the source populations (replication of 1/population). Additional replication occurred when

enough individuals (*i.e.,* 160 or more) were available (Figure S6).

Data analysis

All statistical analyses were performed in R version 4.1.2 (R Core Team, 2021).

Population size through time was used to assess population dynamics throughout the rescue

experiment. Populations were considered rescued if they grew to a population size greater than

50, and maintained a growth rate of greater than 1 (the replacement rate) for at least two of

the final four generations. This allowed for decreases in population size due to negative density

dependence. Extinction risk and survival times were quantified using a survival analysis with the

survival package (Therneau 2021). Extinct populations were defined as those with zero adults at

census during generations 1-6 or one adult by generation 7. Overall extinction rates were

calculated as raw percentages with confidence intervals around each percentage for each

treatment.

Population growth was calculated as the mean number of surviving adult offspring per

individual in the population, *i.e.,* census population size measured on day 35 (N_t) divided by the

initial population size (N_{t-1}), or $\frac{N_t}{N_{t-1}}$. We evaluated both density-dependent and density-

independent growth rates through time using a linear mixed model with the *lme4* package

(Bates *et al.*, 2015). We included immigration treatment (0 migrants, 1 migrant, or 5 migrants

per generation), time (t) in generations (from 1 to 7 generations, categorical), and initial

population size (N_{t-1}, continuous) as fixed effects, and temporal block (block 1, block 2, block

3), source population (nested within block), and population identity (nested within source

population) as random effects. Growth rate was square root transformed to accommodate zero

values and to ensure that the equal variance assumption was satisfied.

From this model, we generated estimates of both density-dependent and density-

independent population growth rates. To estimate density-dependent growth rates, we set the

values for the initial population size (an explanatory variable in the model) to vary for each

generation and treatment combination, matching the observed mean initial population sizes.

These estimates were used to describe the population growth data and quantify the effects of

immigration treatment in a finite habitat by considering the constraints of competition for

resources and cannibalism that affected the populations in our experiment (Pointer *et al.*,

2021). To estimate density-independent growth rates from the fitted model, we set the values

of initial population size to zero (*i.e.,* $N_{t-1} = 0$) for every generation and treatment, which is

equivalent to estimating the y-intercepts (see Figure S5). These density-independent estimates

allowed us to track adaptation to the environment by revealing the change in intrinsic fitness of

individuals in the population unfettered by the effects of density. We expected the density-

independent growth rates to be higher than the density-dependent estimates due to the

known negative density dependence in *T. castaneum* (Melbourne & Hastings 2008). Confidence

intervals for both density-dependent and density-independent growth rate estimates were

estimated by parametric bootstrap (percentile method) using the *bootMer* function with 5000

simulations in the *lme4* package (Bates *et al.*, 2015). Comparisons between treatments were

made by estimating confidence intervals by parametric bootstrap (percentile method) for the

difference in means between treatments within each generation using the *simpleboot* and *boot*

packages with 5000 simulations (Peng, 2019; Canty & Ripley, 2022).

The timing of evolutionary rescue was assessed in each treatment group using a

Generalized Additive Mixed (GAM) model (Wood, 2017) similar to Olazcuaga *et al.* (2023). Only

rescued populations were considered in this analysis. We modeled the population size of

rescued populations using a negative binomial distribution to account for overdispersion using

the package *mgcv* (Wood, 2004, 2011). To allow for the relationship between the population

size and time to be nonlinear, we modeled population size as a smooth function of generation

with the number of knots set to 4, which was chosen to maximize the R-squared value.

Explanatory variables also included immigration treatment and the interaction of treatment

and generation as fixed effects, allowing for a smooth function for each treatment. The full

model included population identity and temporal blocks as random effects. The best-fit model

was chosen by comparing to reduced models using the *compareML* function from the *itsadug*

package (van Rij *et al.*, 2022).

We used the reciprocal transplant data to further evaluate adaptation to the

experimental environment, as well as potential fitness costs of adaptation in a benign

environment. Growth rate in the reciprocal transplant experiment was used as the response

variable in a linear mixed effects model that included population type (source, 0-migrant, 1-

migrant and 5-migrant), test environment (source, experimental, severe), and the interaction

between population type and test environment as fixed effects. Population identity was

included as a random effect nested within temporal block. Comparisons within each

environment and among immigration treatments were made using 95% bootstrapped

confidence intervals (percentile method) for the difference in means, which were generated

from 5000 iterations.

3 | Results

Population size and extinction

During the rescue experiment, a majority of populations experienced evolutionary

rescue (Figure 1, black lines), and extinction probability was similar for all three treatments

(Figure 1, red lines). The difference in the extinction risk was statistically unclear across

treatments throughout the experiment (Figure S4). Additionally, there was no clear difference

in survival time among treatments ($\chi_2^2 = 0.1$, p = 1.00).

274
275

Figure 1. Population size through time in discrete generations for the 0-migrant, 1-migrant,

and 5-migrant per generation treatments during the 7-generation rescue experiment. Each

black line represents a rescued population, each orange line represents an extant population

that was not considered rescued, and each red line represents a population that went extinct or

had a population size of one individual in generation 7. Immigration events are indicated with

instantaneous increases in population size, which occurred at the start of generations 1-6.

Extinction rates (% of red lines) and simple confidence intervals around each percent are

provided.

3.2 | Growth rate through time

Density-dependent growth rates increased through generation four, then decreased

again as population density increased (Figure 2a, Table S1). Density-independent growth rates

were below or close to the replacement rate (growth rate of 1) in generations 1-3 (Figure 2b;

confidence intervals overlap with dashed line). This changed in generation 4, when the 0-

migrant treatment had a mean density-independent growth rate much larger than the

replacement rate, illustrating rapid adaptation within those populations (Figure 2). In

generation 4, density-independent growth rate in the 0-migrant populations was on average

0.808 (95% CI [0.207, 1.453]) higher than the 1-migrant populations and 0.969 (0.478, 1.573)

higher than the 5-migrant populations. By generation 5, all treatments gained and then

maintained a mean density-independent growth rate greater than 1, showing adaptation to the

environment. The comparison between the density-independent and density-dependent

estimates of growth rates suggest that decrease in the latter by generation 7 was likely due to

the effects of negative density dependence, which increased as populations grew (see Figures 1

& 3).

Figure 2. Density-dependent growth rate (a) and density-independent growth rate (b) throughout the

rescue experiment (generation 1 to 7). Each point outlined in black is the predicted mean for each

treatment (purple square = 0 migrants, yellow diamond = 1 migrant, and blue circle = 5 migrants). Error

bars are bootstrapped 95% confidence intervals. Points without black outlines are raw growth rate

values, and each point represents one population. The dotted line represents a growth rate of 1, the

replacement rate. Two estimates for growth rate are presented: **a)** the density-dependent estimates are

predictions from the linear mixed model using average initial population size for each generation and

treatment combination, used to incorporate the effects of negative density-dependence and provide

estimates that describe the raw data (shown), and **b)** the density-independent estimates (*i.e.,* using an

initial population size of 0), showing adaptation over time.

Timing of evolutionary rescue

Among the rescued populations only (black lines in Figure 1), the analysis of population

size with a GAM model revealed a shallow U-shaped curve (Figure 3). The estimated curves

from the best-fit model, which included treatment and the interaction of treatment and

generation as fixed effects and population identity as a random effect, are shown in Figure 4.

The interaction between immigration treatment and generation was significant ($\chi^2_3 > 330$, $p <$

0.0001), with the 0-migrant populations growing slightly faster than populations receiving

migrants (Figure 3). The 1-migrant and 5-migrant treatments maintained similar population

trajectories throughout the experiment.

Figure 3. Evolutioanry rescue analysis. Population size of rescued populations receiving no

migrants (purple line, squares), one migrant (yellow line, diamonds), or five migrants (blue line,

circles) through time in generations estimated by the best-fit GAM model. Each point

represents one population, which are staggered to reduce overlap. The lines are estimates of

mean population size for the three immigration treatments from the best-fit GAM model. The

shaded region around each line is the 95% confidence interval given by the GAM model.

Adaptation and associated costs

Growth rate in the reciprocal transplant experiment was influenced by an interaction

between environment and population type ($F_{6, 168.04} = 25.16$, $p < 0.0001$), in a pattern showing

that adaptation to the challenging environment resulted in reduced fitness in the benign source

environment (Figure 4). There, the experimental populations had lower growth rates on

average compared to the source populations (95% $CI_{0\text{-migrant} - source}$ [-0.855, -1.628]; 95% $CI_{1\text{-migrant}}$

$_{- source}$ [-0.911, -1.748]; 95% $CI_{5\text{-migrant} - source}$ [-0.641, -1.482]), and a higher growth rate in the

severe habitat (95% $CI_{0\text{-migrant} - source}$ [0.442, 0.171]; 95% $CI_{1\text{-migrant} - source}$ [0.424, 0.210]; 95% $CI_{5\text{-}}$

$_{migrant - source}$ [0.464, 0.216]). Additionally, the 5-migrant populations performed marginally

better than the 0-migrant populations in the experimental environment (95% $CI_{0\text{-migrant} - 5\text{-migrant}}$

[-0.0675, 0.572]).

Unexpectedly, the experimental and source populations performed similarly in the

challenging experimental environment (Figure 4, all confidence intervals of the differences

overlap with zero: 95% $CI_{0\text{-migrant} - source}$ [-0.404, 0.209]; 95% $CI_{1\text{-migrant} - source}$ [-0.220, 0.403]; 95%

$CI_{5\text{-migrant} - source}$ [-0.153, 0.452]). The experimental populations did not perform as well as

expected based on their growth rates at the end of the rescue experiment (Figure 2). Our

expectation for growth rates were based on predictions with initial population size of 40 (the

size used in the reciprocal transplant) from the linear mixed model created for the growth rate

data in the rescue experiment. Those predicted values were 1.73, 95% CI [1.19, 2.34], 1.46

[1.02, 1.98], and 1.68 [1.21, 2.23] for the 0-migrant, 1-migrant, and 5-migrant treatments,

respectively. The average growth rates we observed in the reciprocal transplant experiment in

the experimental environment were lower than these estimates, with the means 0.86 [0.64,

1.10], 1.04 [0.81, 1.31], 1.09 [0.87, 1.36] that fall at the edge or outside the predicted values

and CIs, and confidence intervals that overlap only slightly.

Furthermore, source populations did not perform as poorly as expected. We expected

them to perform similarly to the first generation of the experimental populations in the rescue

experiment prior to adaptation. The prediction given N_{t-1} = 40 was 0.62 [0.38, 0.92] for the 0-

migrant populations, while the mean growth rate of source populations was 0.92 [0.72, 1.15].

The CIs overlap partially, but the overlap does not include the mean, confirming that the source

growth rates were unexpectedly high.

Figure 4. Reciprocal transplant experiment results. Growth rate was measured in three

environments: the source environment (standard dry media), the challenging experimental

environment (2.968ppm deltamethrin, the concentration used during the rescue experiment),

and a severe environment containing a higher concentration (5.641ppm). Each point outlined in

black is the model-estimated mean for each treatment (purple square = no migrants, yellow

diamond = 1 migrant, blue circle = 5 migrants, and green triangle = source population). Black

error bars are bootstrapped 95% confidence intervals; for the source populations in the severe

environment (green), the black error bars are not visible because they are too small to be seen.

Thick lines connecting these points emphasize the treatment-environment interaction. Each

point without the black outline represents a single population, with the light error bars

representing raw 95% confidence intervals across replicates; points without error bars have no

replicates. Points are staggered and jittered to reduce overlap.

4 | Discussion

Our study evaluated the effects of different immigration rates on adaptation to a new,

challenging environment using experimental populations of *T. castaneum*. We evaluated

differences between populations receiving two immigration rates (1 or 5 migrants per

generation) with isolated populations that received no immigration. The migants were not

adapted to the challenging environment, allowing us to examine the degree to which

immigration can limit adaptation (Savolainen *et al.*, 2013). While the experimental populations

declined in size initially, leading some populations to go extinct, around 80% of populations

were able to persist for the duration of the 7-generation experiment. A majority of these

populations had growth rates above the replacement rate and large population sizes,

characteristic of evolutionary rescue. We also found that extinction rate was similar regardless

of immigration. However, we observed that growth rates increased one generation sooner in

the isolated populations compared to those with immigration, indicating that outcrossing with

migrants that were not adapted to the challenging environment likely delayed adaptation. This

delay was brief, as the following generation, growth rates were comparable across the different

immigration treatments. We also tested for adaptation using a reciprocal transplant

experiment, where we observed a pattern of local adaptation, with experimental populations

having higher growth rates compared to naive populations in a severe environment, and lower

growth rates than the naive populations in a benign environment, which suggests that

adaptation to the challenging environment was associated with a cost.

Observed delay in adaptation

Populations without immigration gained a density-independent growth rate greater

than the replacement rate one generation before the populations with immigration. The

following generation, all populations grew at a similar rate regardless of immigration. Taken

together, we can conclude immigration of individuals not adapted to a new, challenging

environment, even at a reasonably high level, did not prevent adaptation, only that it slowed

the process. Such delayed adaptation could be detrimental to populations experiencing urgent,

time-sensitive extinction risk due to small size (Melbourne & Hastings, 2008). However, for

populations not at immediate risk of extinction, such a delay may be inconsequential, and

worth the potential increase in genetic diversity that occurs through immigration, which may

improve the ability of the population to adapt to future environmental changes. This finding of

only a brief (one generation) delay of adaptation has practical implications. Naive migrants are

often more feasible to obtain, either from captive rearing programs (Crone *et al.,* 2007) or from

large source populations living in a different habitat (*e.g.,* different spider mite host plants in

Bisschop et al., 2019) or geographic region (*e.g.,* geographically isolated guppy populations in

Fitzpatrick et al., 2020), when habitat matching is not possible.

Our findings are consistent with refuge strategies to delay evolution of pest resistance

to toxins, such as in crops genetically modified to express *Bacillus thuringiensis* (*Bt*). The

immigration of susceptible individuals from sections of fields planted in non-*Bt* crops, or

growing crops in areas with plentiful alternative host plants for the pest has been proven to be

highly effective (Carrière *et al.*, 2012). In those situations, particularly with 50% or more of a

pest population being susceptible, resistance can be successfully managed for multiple

generations (Tabashnik *et al.*, 2008). Given our much smaller proportion of susceptible

individuals, it makes sense that the delay of the evolution of resistance was brief.

Reciprocal transplant experiment: thoughts on adaptation and genetic load

We observed consistent signals of adaptation in the experimental populations in the

second half of the rescue experiment and in the severe environment of the reciprocal

transplant experiment. However, we did not expect the experimental and source populations

to have similar growth rates in the experimental environment – the experimental populations

performed worse than expected, and the source populations performed better than expected,

as detailed in the results. We did observe that the populations receiving five migrants per

generation performed marginally better than the isolated populations. This suggests that there

may be a benefit of immigration in the longer term by preventing or alleviating genetic load

(Hedrick & Garcia-Dorado, 2016; Stewart *et al.*, 2017).

However, the performance of the experimental populations overall was lower than

expected, and this is challenging to explain. It does not seem likely to be a general effect of

genetic load, as just two generations prior, in generation 7 of the rescue experiment,

populations demonstrated density-independent growth rates well above the replacement rate.

The relatively high growth rate of the source populations in the challenging environment may

be related to adaptation to a different environmental challenge, high density, which affected

source populations. The source populations were maintained at a consistently higher density (N

> 150 on average for all 7 generations; Figure S3) than the experimental populations (N > 150 in

generations 4-7 only; Figure 1) in habitat patches of the same size. High density creates a

stressful environment for *T. castaneum,* which results in adults releasing stress hormone and

engaging in increased cannibalism (Hirashima *et al.,* 1993; Pointer *et al.,* 2021). This heightened

competition for twice as many generations may have resulted in adaptation to higher density,

perhaps through evolution of increased cannibalism (Stevens, 1989) along with the use of

cannibalism to persist in the challenging, high-density environment (Via, 1999). In our reciprocal

transplant experiment, increased cannibalism may have enabled the source populations to

perform comparably to the experimental populations in the challenging experimental

environment as well.

Reciprocal transplant experiment: costs to adaptation

The adaptation of the experimental populations to the challenging environment, while

clear when examining the density-independent growth rates, was only apparent in the

reciprocal transplant experiment in the severe environment. Adaptation to moderate-to-high

concentrations of deltamethrin is well-documented in pest populations of *T. castaneum*

(Collins, 1998; Stuart *et al.,* 1998; Kalsi & Palli, 2015), and in our study, adaptation enabled

tolerance of higher concentration of the insecticide. The low performance of insecticide-

adapted populations in a benign environment suggests a cost to adaptation, which has been

documented in the literature for other insect species. For example, detoxification of pyrethroid

insecticides through the over-expression of cytochrome P450 enzymes leads to reduced

fecundity and fertility in mosquitoes (Tchouakui *et al.*, 2020). While a direct cost to pyrethroid

resistance in *T. castaneum* has to our knowledge not been documented, it seems likely that

there would be a fitness cost given that the cytochrome P450 pathway is also involved in

resistance to deltamethrin in the species (Kalsi & Palli, 2015).

Considerations for future experiments

The bottleneck events that occurred during the rescue experiment, some as small as 10

individuals, will likely have longer-term consequences (Frankham *et al.*, 1999; Olazcuaga *et al.*,

2023). Census population size is almost always larger than the effective population size (the

median N_e/N_c ratio across plant and animal taxa is approximated to be 0.1; for *T. castaneum,* it

is estimated to be between 0.5-0.9) and bottlenecks of $N_e < 50$ can result in reductions in

genetic variation and increases in inbreeding (Wade, 1980; Jamieson & Allendorf, 2012). For

this reason, we suggest that future work examine the effects of demographic bottlenecks of

this severity by propagating experiments like this for longer (*i.e.,* more than 8 generations).

Longer-term experiments would be able to test the hypothesis that the benefits of immigration

become more apparent (*i.e.,* genetic load may be lower in populations with immigration as

opposed to those without) over longer time scales. Such experiments would help shed light on

the longer-term impacts of immigration for small and isolated populations, an important

question when considering whether to implement translocations in declining natural

populations (Frankham, 2015, 2016). Genomic methods would also provide important insights

for monitoring the genetic contributions of migrants and their effects on population fitness

(Whiteley *et al.*, 2015).

Conclusions

Overall, this study explored the effects of immigration of naive individuals on adaptation

of previously large and outbred populations following sudden environmental degradation and

habitat fragmentation. Of course, the ideal solution would be to restore the habitat, and this

study provides support for using immigration as a tool to sustain populations for multiple

generations until restoration can occur (Whiteley *et al.*, 2015; Frankham *et al.*, 2019). We found

that though immigration slowed adaptation in the short-term, extant populations with and

without immigration were on average adapted to the new and challenging environment after

seven generations. Broadly, the balance between migration and selection can determine the

magnitude of adaptation in natural populations as well as the amount of genetic variation

present among connected populations (Kawecki & Ebert, 2004; Yeaman & Jarvis, 2006;

Savolainen *et al.*, 2013; Reid *et al.*, 2021). Thus, we demonstrate that evolutionary rescue can

occur both with and without consistent immigration from a naive source population. Our

findings suggest the deleterious effects of naive migrants are short-lived, and should not

prevent managers from implementing consistent immigration through corridors or

translocations to maintain genetic diversity and increase adaptive potential in populations

exposed to novel environmental change.

500 Acknowledgements

501   This work was supported by the U.S. National Science Foundation Graduate Research

502 Fellowship No. 006784 to LFD and Grant Nos. 1930222 and 1930650 to BAM and RAH,

503 respectively. We thank Sarah DeLacey, Freddie Habarecht, Mayra Contreras-Hernandez, Desiree

504 Moore, and Rachel Szymanski for assistance maintaining laboratory *T. castaneum* colonies used

505 in this experiment. We thank XX anonymous reviewers for their constructive feedback.

507 Conflict of Interest

508 The authors declare no conflicts of interest.

510 References

Bates, D., Maechler, M., Bolker, B. & Walker, S. 2015. Fitting Linear Mixed-Effects Models Using
lme4. *J Stat Softw* **67**: 1–48.
Bell, G. 2017. Evolutionary Rescue. *Annu Rev Ecol Evol Syst* **48**: 447–71.
Bisschop, K., Mortier, F., Etienne, R.S. & Bonte, D. 2019. Transient local adaptation and source-
sink dynamics in experimental populations experiencing spatially heterogeneous
environments. *Proceedings of the Royal Society B: Biological Sciences* **286**.
Brady, S.P., Bolnick, D.I., Angert, A.L., Gonzalez, A., Barrett, R.D.H., Crispo, E., *et al.* 2019. Causes
of maladaptation. *Evol Appl* **12**: 1229–1242.
Canty, A. & Ripley, B. 2022. boot: Bootstrap R Functions.
Carlson, S.M., Cunningham, C.J. & Westley, P.A.H. 2014. Evolutionary rescue in a changing
world. *Trends in Ecology & Evolution* **29**: 521-530.
Carrière, Y., Ellers-Kirk, C., Hartfield, K., Larocque, G., Degain, B., Dutilleul, P., *et al.* 2012. Large-
scale, spatially-explicit test of the refuge strategy for delaying insecticide resistance. *Proc*
Natl Acad Sci U S A **109**: 775–780.
Cheptou, P.O., Hargreaves, A.L., Bonte, D. & Jacquemyn, H. 2017. Adaptation to fragmentation:
Evolutionary dynamics driven by human influences. *Philosophical Transactions of the Royal*
Society B: Biological Sciences **372**.
Collins, P.J. 1998. Inheritance of resistance to pyrethroid insecticides in Tribolium castaneum
(Herbst). *J Stored Prod Res* **34**: 395–401.

Cote, J., Dahirel, M., Schtickzelle, N., Altermatt, F., Ansart, A., Blanchet, S., *et al.* 2022. Dispersal
531   syndromes in challenging environments⬚: A cross- species experiment. 1–13.

532   Crone, E.E., Pickering, D. & Schultz, C.B. 2007. Can captive rearing promote recovery of
533   endangered butterflies? An assessment in the face of uncertainty. *Biol Conserv* **139**: 103–
534   112.

535   Durkee, L.F., Olazcuaga, L., Szymanski, R., Melbourne, B.A. & Hufbauer, R.A. 2023. Genetic
536   mixing facilitates adaptation to a novel environmental constraint. *Ecol Entomol* **48**: 517–
537   522.

538   Edelaar, P. & Bolnick, D.I. 2012. Non-random gene flow: An underappreciated force in evolution
539   and ecology. *Trends Ecol Evol* **27**: 659–665.

540   Fitzpatrick, S.W., Bradburd, G.S., Kremer, C.T., Salerno, P.E., Angeloni, L.M. & Funk, W.C. 2020.
541   Genomic and Fitness Consequences of Genetic Rescue in Wild Populations. *Current Biology*
542   **30**: 517-522.e5.

543   Foster, B.J., McCulloch, G.A., Vogel, M.F.S., Ingram, T. & Waters, J.M. 2021. Anthropogenic
544   evolution in an insect wing polymorphism following widespread deforestation. *Biol Lett* **17**.

545   Frankham, R. 2016. Genetic rescue benefits persist to at least the F3 generation, based on a
546   meta-analysis. *Biol Conserv* **195**: 33–36.

547   Frankham, R. 2015. Genetic rescue of small inbred populations: meta-analysis reveals large and
548   consistent benefits of gene flow. *Mol Ecol* **24**: 2610–2618.

549   Frankham, R., Ballou, J.D., Ralls, K., Eldridge, M., Dubash, M.R., Fenster, C.B., *et al.* 2019. *A*
550   *Practical Guide for Genetic Management of Fragmented Animal and Plant Populations*, 1st
551   ed. Oxford University Press, Oxford, UK.

552   Frankham, R., Lees, K., Montgomery, M.E., England, P.R., Lowe, E.H. & Briscoe, D.A. 1999. Do
553   population size bottlenecks reduce evolutionary potential? *Anim Conserv* **2**: 255–260.

554   Gomulkiewicz, R. & Holt, R.D. 1995. When does Evolution by Natural Selection Prevent
555   Extinction? *Evolution (N Y)* **49**: 201.

556   Hedrick, P.W. & Fredrickson, R. 2010. Genetic rescue guidelines with examples from Mexican
557   wolves and Florida panthers. *Conservation Genetics* **11**: 615–626.

558   Hedrick, P.W. & Garcia-Dorado, A. 2016. Understanding Inbreeding Depression, Purging, and
559   Genetic Rescue. *Trends Ecol Evol* **31**: 940–952.

560   Hirashima, A., Nagano, T., Takeya, R. & Eto, M. 1993. Effect of larval density on whole-body
561   biogenic amine levels of Tribolium freemani hinton. *Comparative Biochemistry and*
562   *Physiology. Part C: Comparative* **106**: 457–461.

563   Hufbauer, R.A., Szücs, M., Kasyon, E., Youngberg, C., Koontz, M.J., Richards, C., *et al.* 2015.
564   Three types of rescue can avert extinction in a changing environment. *PNAS* **112**: 10557–
565   10562.

566   Jamieson, I.G. & Allendorf, F.W. 2012. How does the 50/500 rule apply to MVPs? *Trends Ecol*
567   *Evol* **27**: 578–584.

568   Kalsi, M. & Palli, S.R. 2015. Transcription factors, CncC and Maf, regulate expression of CYP6BQ
569   genes responsible for deltamethrin resistance in Tribolium castaneum. *Insect Biochem Mol*
570   *Biol* **65**: 47–56.

571   Kawecki, T.J. & Ebert, D. 2004. Conceptual issues in local adaptation. *Ecol Lett* **7**: 1225–1241.

572   Lenormand, T. 2002. Gene flow and the limits to natural selection. *Trends Ecol Evol* **17**: 183–
573   189.

Loria, A., Cristescu, M.E. & Gonzalez, A. 2019. Mixed evidence for adaptation to environmental
pollution. *Evol Appl* **12**: 1259–1273.
Melbourne, B.A. & Hastings, A. 2008. Extinction risk depends strongly on factors contributing to
stochasticity. *Nature* **454**: 100–103.
Mills, L.S. & Allendorf, F.W. 1996. The one-Migrant-per-Generation Rule in Conservation and
Management. *Conservation Biology* **10**: 1509–1518.
Newman, D. & Tallmon, D.A. 2001. Experimental evidence for beneficial fitness effects of gene
flow in recently isolated populations. *Conservation Biology* **15**: 1054–1063.
Nordstrom, S.W., Hufbauer, R.A., Olazcuaga, L., Durkee, L.F. & Melbourne, B.A. 2023. How
density dependence, genetic erosion, and the extinction vortex impact evolutionary
rescue. *bioRxiv* 1–42.
Olazcuaga, L., Lincke, B., Delacey, S., Durkee, L.F., Melbourne, B.A. & Hufbauer, R.A. 2023.
Population demographic history and evolutionary rescue: influence of a bottleneck event.
Evol Appl 1483–1495.
Osmond, M.M. & de Mazancourt, C. 2013. How competition affects evolutionary rescue.
Philosophical Transactions of the Royal Society B: Biological Sciences **368**.
Peng, R.D. 2019. simpleboot: Simple Bootstrap Routines. R package version 1.1-7.
https://CRAN.R-project.org/package=simpleboot
Pointer, M.D., Gage, M.J.G. & Spurgin, L.G. 2021. Tribolium beetles as a model system in
evolution and ecology. *Heredity (Edinb)* **126**: 869–883. Springer US.
Reid, J.M., Arcese, P., Nietlisbach, P., Wolak, M.E., Muff, S., Dickel, L., *et al.* 2021. Immigration
counter-acts local micro-evolution of a major fitness component: Migration-selection
balance in free-living song sparrows. *Evol Lett* **5**: 48–60.
Savolainen, O., Lascoux, M. & Merilä, J. 2013. Ecological genomics of local adaptation. *Nat Rev*
Genet **14**: 807–820. Nature Publishing Group.
Slatkin, M. 1987. Gene flow and the geographic structure of natural populations. *Science (1979)*
236: 787–792.
Spielman, D. & Frankham, R. 1992. Modeling problems in conservation genetics using captive
Drosophila populations: Improvement of reproductive fitness due to immigration of one
individual into small partially inbred populations. *Zoo Biol* **11**: 343–351.
Stevens, L. 1989. The genetics and evolution of cannibalism in flour beetles (genus *Tribolium*).
Evolution (N Y) **43**: 169–179.
Stewart, G.S., Morris, M.R., Genis, A.B., Szűcs, M., Melbourne, B.A., Tavener, S.J., *et al.* 2017.
The power of evolutionary rescue is constrained by genetic load. *Evol Appl* **10**: 731–741.
Stuart, J.J., Ray, S., Harrington, B.J., Neal, J.J. & Beeman, R.W. 1998. Genetic Mapping of a Major
Locus Controlling Pyrethroid Resistance in Tribolium castaneum (Coleoptera:
Tenebrionidae). *J Econ Entomol* **91**: 1232–1238.
Tabashnik, B.E., Gassmann, A.J., Crowder, D.W. & Carrière, Y. 2008. Insect resistance to *Bt*
crops: Evidence versus theory. *Nat Biotechnol* **26**: 199–202.
Tchouakui, M., Riveron Miranda, J., Mugenzi, L.M.J., Djonabaye, D., Wondji, M.J., Tchoupo, M.,
et al. 2020. Cytochrome P450 metabolic resistance (CYP6P9a) to pyrethroids imposes a
fitness cost in the major African malaria vector *Anopheles funestus. Heredity (Edinb)* **124**:
621–632. Springer US.

Therneau, T. M. (2021). A Package for Survival Analysis in S. version 3.2-13, URL:
https://CRAN.R-project.org/package=survival
R Core Team 2021. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.
van Rij, J., Wieling, M., Baayen, R.H. & van Rijn, H. 2022. itsadug: Interpreting Time Series and
Autocorrelated Data Using GAMMs. R package version 2.4.1.
Via, S. 1999. Cannibalism facilitates the use of a novel environment in the flour beetle,
Tribolium castaneum. *Heredity (Edinb)* **82**: 267–275.
Wade, M.J. 1980. Effective population size: the effects of sex, genotype, and density on the
mean and variance of offspring numbers in the flour beetle, *Tribolium castaneum. Genet*
Res **36**: 1–10.
Whiteley, A.R., Fitzpatrick, S.W., Funk, W.C. & Tallmon, D.A. 2015. Genetic rescue to the rescue.
Trends in Ecology & Evolution **30**: 42-49.
Wood, S.N. 2011. Fast stable restricted maximum likelihood and marginal likelihood estimation
of semiparametric generalized linear models. *Journal of the Royal Statistical Society* **73**: 3–
36.
Wood, S.N. 2017. *Generalized Additive Models: An Introduction with R*, 2nd ed. Chapman and
Hall/CRC.
Wood, S.N. 2004. Stable and efficient multiple smoothing parameter estimation for generalized
additive models. *J Am Stat Assoc* **99**: 673–686.
Yeaman, S. & Jarvis, A. 2006. Regional heterogeneity and gene flow maintain variance in a
quantitative trait within populations of lodgepole pine. *Proceedings of the Royal Society B:*
Biological Sciences **273**: 1587–1593.
640