Global conservation prioritization approach provides credible results at a regional scale

Michael Roswell[1,2] mroswell@umd.edu 0000-0002-8479-9184

Anahí Espíndola[1] anahiesp@umd.edu 0000-0001-9128-8836

[1]Department of Entomology, University of Maryland, College Park, MD, USA

[2]Department of Biology, University of Maryland, College Park, MD, USA

Abstract

Aim 	Conservationists and managers must direct resources and enact measures to protect

species, despite uncertainty about their present statuses. One approach to covering the data gap is

borrowing information from data-rich species or populations to guide decisions about data-poor

ones, via machine learning. Recent efforts demonstrated proof-of-concept at the global scale, leaving

unclear whether similar approaches are feasible at the local and regional scales at which conservation

actions most typically occur. To address this gap, we tested a global-scale predictive approach at a

regional scale, using two groups of taxa.

Location 	Maryland, USA.

Taxa 	Vascular land plants and lepidopterans.

Methods 	Using publicly available occurrence and biogeographic data, we trained random forest

classifiers to predict the state-level conservation status of species in each of the two focal taxa. We

assessed model performance with cross-validation, and explored trends in the predictions.

Results 	Our models had strong discriminatory ability, accurately predicting status for species

with existing status assessments. They predict that the northwestern part of Maryland, USA, which

overlaps the Appalachian Mountains, harbors a higher concentration of unassessed, but threatened

plants and lepidopterans. Our predictions track known biogeographic patterns, and unassessed

species predicted as most likely threatened in Maryland were often recognized as also needing

conservation in nearby jurisdictions, providing external validation to our results.

Main Conclusions   We demonstrate that a modelling approach developed for global analysis can

be downscaled and credible when applied at a regional scale that is smaller than typical species

ranges. We identified select unassessed plant and lepidopteran species, and the western, montane

region of Maryland as priority targets for additional monitoring, assessment, and conservation. By

rapidly aggregating disparate data and integrating information across taxa, models like those we used

can complement traditional assessment tools and assist in prioritization for formal assessments, as

well as protection.

Key words

Predictive modelling, conservation status, random forest, lepidopterans, plants, downscaling,

prioritization, multi-taxon

Introduction

Conservationists and natural resource managers cannot wait until all the data are in, and must direct

resources and enact measures to assess – and ultimately protect – species and populations, despite

uncertainty about their present status. One stepping-stone approach to bridging the data gap is

borrowing information from data-rich species or populations to guide decision making about data-

poor ones (Callaghan, Nakagawa, & Cornwell, 2021; Free et al., 2020; Pelletier, Carstens, Tank,

Sullivan, & Espíndola, 2018; Zizka, Andermann, & Silvestro, 2022), using predictive modelling

frameworks. Increasingly, publicly available datasets that aggregate observations (Heberling, Miller,

Noesgaard, Weingart, & Schigel, 2021) enable formulating and fitting predictive models to guide

conservation and natural resource management (Bastin et al., 2019; Caetano et al., 2022; Callaghan et

al., 2021; Panter et al., 2020; Pelletier et al., 2018; Zizka et al., 2022). The predictions from such models

might be used to support conservation actions, and guide prioritization for resource-intensive status

assessments (Pelletier et al., 2018). On the latter, national and subnational jurisdictions generate

taxon-specific management plans triggered by formal status assessments (e.g., via National and

State-level Endangered Species Acts in the United States (Department of the Interior, 1973; Nongame

and endangered species conservation act, 2022)). In this context, status predictions from predictive

models such as those mentioned above are not sufficient to trigger protections, but might undergird

the listing process, by assisting with monitoring and assessment prioritization in a resource-limited

world.

Several knowledge gaps or "shortfalls" prevent strategic species protection (Cardoso, Erwin, Borges,

& New, 2011b; Mokany & Ferrier, 2011). Here, we focus on the "Rabinowitzian" shortfall, i.e., the gap

between knowing species' occurrence patterns and using that information directly for evaluating their

conservation risks (Cardoso, Erwin, Borges, & New, 2011a; Crisfield, Blanchet, & Gravel, 2022; Isbell et

al., 2023; Rabinowitz, 1981). Models that use existing data efficiently can identify under-recognized

patterns of threat and endangerment (Regan et al., 2008), and enable efficient allocation of

monitoring, assessment, and protection resources (Hochkirch et al., 2021). For these goals, statistical

learning tools that aggregate disparate data streams can recognize meaningful patterns that might

escape notice in more granular analyses. These types of tools were used by Pelletier et al. (2018), who

used only a handful of spatial and climatic covariates, and were able to predict the conservation

status of unassessed land plants. They identified taxa likely to meet IUCN red list criteria, as well as

previously unrecognized hotspot regions that likely harbor high concentrations of threatened but

unassessed taxa. The authors anticipated that their approach could easily serve conservation needs at

smaller spatial scales, which would make their method directly applicable at the scales at which most

conservation actions take place.

Although the approach Pelletier et al. (2018) used can be technically readily implemented at nearly

any scale, downscaling entails radical shifts in data availability, quality, and biogeographic

representation that could improve or erode the utility of the approach. At smaller, regional scales,

data quality may be more similar between assessed and un-assessed taxa, which should improve

predictions. Additionally, at smaller scales, complete datasets containing a wider variety of

potentially appropriate covariates (e.g., land use and its history) may be available. Nevertheless, at

smaller spatial scales, records of species' occurrences are unlikely to cover species' whole ranges, and

the number of species in a smaller region is also likely to be smaller, potentially limiting statistical

power. Additionally, it is unknown whether the approach is extensible to other taxa beyond plants

with little modification. If the same suites of geographic covariates can predict conservation status of

disparate taxa, this predictive approach has high promise to assist in a variety of conservation

contexts.

Here, we downscale the approach Pelletier et al. (2018) took to predict conservation statuses,

applying similar methods to vascular land plants (the focal higher taxon in Pelletier et al. (2018),

hereafter "plants") and lepidopterans (a new focal group) within a sub-national management region

(the state of Maryland, USA). We selected these two taxa because the state of Maryland maintains a

rare, threatened, and endangered list for each (Maryland Natural Heritage Program, 2021a, 2021b)and

the conservation biology of these two taxa differ (Isbell et al., 2023; Wagner, Fox, Salcido, & Dyer,

2021). We asked whether, despite differences in the quality and quantity of data available, as well as

the ecology of the taxa upon which predictions are made, these highly general predictive modelling

tools can help cover the Rabinowitzian shortfall, and guide conservation decisions at smaller, sub-

national scales. In discussing our results, we evaluate model performance and prediction credibility,

and then explore our predictions. We identify species that are likely threatened in the state of

Maryland, USA, and regions within the state that are likely to harbor a high density of threatened,

unassessed species. Finally, we use our case studies to discuss the promises and potential pitfalls of

using similar predictive modelling tools for conservation prioritization at regional and local scales.

Materials and Methods

Method Overview

To predict the conservation status of lepidopterans and plants in the state of Maryland, we obtained

publicly available species occurrence and spatial data. We used data for all assessed species to build

random forest classifiers, tuning hyperparameters and evaluating model discriminatory ability, and

expected generalization error, through nested cross-validation. Then, for each taxon group (i.e.,

lepidopterans and plants), we fit a final model that was trained on all assessed species, and uses these

to predict conservation status for all unassessed species known to occur within the state. We mapped

predictions to visually screen for spatial patterns, and then conducted a limited literature search to

contextualize status predictions with independent lines of evidence. In the following sections, we

provide details on the data we used, how we constructed and assessed models, the ways we

generated and visualized predictions, and how we used independent lines of evidence to corroborate

predictions.

Data

Plant and Lepidopteran Records

We obtained observations of plants and lepidopterans from the state of Maryland between 1989-2021

from GBIF (Appendix S2), irrespective of life stage. We filtered this dataset to include only records

associated with valid species observed in the wild. We assigned each species to a conservation status

category, based on Maryland's state-level rankings, as recorded in NatureServe (NatureServe, 2023).

In order to do this, we first harmonized the taxonomies used by NatureServe and GBIF with

programmatic tools from the R Packages *rgbif, natserv,* and *taxize* (Chamberlain, 2020; Chamberlain,

Oldoni, et al., 2022; Chamberlain, Szoecs, et al., 2022), and some manual curation (Appendix S1), and

then merged the NatureServe conservation status ranks with the occurrence data based on the

consensus species identification. Because relatively few plants, butterflies, or moths had state

conservation statuses in Maryland, we grouped their state conservation categories into three groups

as follows: "secure" taxa were all those categorized as S4 or S5, "threatened" taxa were all those

categorized as S1, S2, S3, SH, or SX; all other taxa were considered unranked. To restrict our analysis

to only species considered native to the state, we first filtered records based on the national and

subnational exotic/native fields from NatureServe, and subsequently, for plants, also filtered them

based on a recent Maryland flora (Knapp & Naczi, 2020). Six threatened, one secure, and 331 unranked

lepidopteran species, and 84 threatened, 20 secure, and 306 unranked plant species were dropped

from future analyses because they were represented by only a single occurrence location. Overall, our

filtered dataset contained records for, respectively, 29, 77 and 1269 threatened, secure and unranked

lepidopterans, and 261, 123 and 940 threatened, secure and unranked plant species. The 1375

lepidopteran species were represented by a total of 78,184 records; 1324 plant species were

represented by 110,232 records.

Environmental Variables

For each location record of a valid native species, we extracted latitude, longitude, and a suite of

associated spatial variables. We extracted current and historic landcover, and indices of landcover

change, from the National Land Cover Dataset (J. Wickham, Stehman, Sorenson, Gass, & Dewitz,

2021), which includes 30m x 30m landcover data at 5-year intervals from 2001 to 2016. We also

extracted interpolated climate data from the expanded CHELSA dataset, which includes 19 variables

that summarize estimated daily, monthly, and annual temperature and precipitation (bioclim), and an

additional 50 variables derived from physical data such irradiance, pressure, and humidity, as well as

their relationships to biological tolerance ranges (e.g., growing season length estimates) at the

resolution of 30 x 30 arc-seconds (Brun, Zimmermann, Hari, Pellissier, & Karger, 2022). Additionally,

for each occurrence record, we extracted the estimated slope (at roughly 100m x 100m resolution)

from the 1-m Statewide Slope layer (Eastern Shore Regional GIS Cooperative, 2021). In total, this

included 69 spatial variables.

Although we gathered spatial data at the occurrence level, we fit models at the species level. To create

the species-level datasets, we summarized each environmental variable across all occurrences of a

species to a pair of summary statistics: for continuous variables we recorded the mean and standard

deviation, and for categorical variables we recorded the modal value and the number of unique levels

observed for the given species. In addition, for each species, we recorded the minimum and maximum

latitude and longitude of their records. Variables with only a single value or missing values prevented

model fitting and were thus dropped from the datasets for both groups of organisms. To serve as a

sort of null model for variable importance, we also included a random continuous variable as a

predictor in each model. Models were built using these 133 summarized species-level predictors.

Model Construction and Evaluation

We divided the dataset into a training dataset, comprised of all species within the categories

"threatened" or "secure," and a prediction dataset, comprised of all unranked species. In order to

tune hyperparameters and evaluate model performance, while retaining as much training data as

possible, we used nested repeated 10 x 10-fold cross-validation. To do this, we created 10 x 10 outer

folds of the training dataset, and on each outer fold, trained a random forest classifier using 10 x 10-

fold cross-validation, using the R packages *caret* and *randomForest* (Kuhn et al., 2022; Liaw, Wiener,

Breiman, & Cutler, 2022). On the inner folds, we conducted a grid search over the hyperparameter

"mtry," which controls the number of features available to use for splitting the data at each node, and

has been shown among the random forest hyperparameters to substantively affect model

performance (Hastie, Tibshirani, & Friedman, 2017; Probst, Wright, & Boulesteix, 2019). We left the

other hyperparameters at their default values, as these are typically considered adequate for model

construction (Probst et al., 2019). We used the area under the receiver-operating characteristic curve

(AUROC) as our measure of performance in parameter tuning, as this measure emphasizes model

discriminatory ability (Ling, Huang, & Zhang, 2003). Although the Random Forest algorithm enables

less computationally intensive model assessment using out-of-bag errors, here we used traditional

cross-validation, which would be applicable to any model fitting algorithm (Kuhn et al., 2022).

For each of the 100 outer folds, we fit a "final model" with the entire dataset, save the held-out fold,

based on the selected hyperparameters, and used this model to make predictions on the held-out

fold. Using random forest votes as a measure of class probability, we computed AUROC, and to

evaluate model performance on unseen data, we computed the mean and standard deviation of the

100 AUROC values computed on these 100 outer folds. In addition, we recorded the mean and

standard deviation in AUROC from the inner, tuning folds, as large differences between AUROC from

the inner and outer folds could indicate a lack of stability in model performance, or a tendency to

overfit the training data.

After the extensive cross-validation described above, we re-trained each model (i.e., one for plants

and one for lepidopterans) on the entire training dataset, again tuning the hyper-parameter "mtry"

with 10x repeated 10-fold cross-validation. We used the "final model" for each taxon group from this

step to make all predictions on unclassified species. The performance estimates we obtained from

previous cross-validation provide the expected performance of these final models on unseen data,

with two major caveats. First, additional data were used to train the final models, which are therefore

not exactly equivalent to the models built on subsets of the same data. Second, we use these final

models for predicting conservation status for unassessed, rather than assessed species. For

unassessed species, we can only evaluate model performance indirectly, by comparing predictions

with non-definitive external data (see below).

Optimal Threshold

Depending on the application, greater precaution may be warranted for misclassification errors for

either threatened or relatively secure species, implying different ideal thresholds for the share of

random forest "votes" required to classify a species into a certain category. As the threshold for

classifying a species as threatened increases, fewer species will be classified as threatened, and the

chance of mis-classifying a species that is, in fact, relatively secure as "threatened" falls, while the

chance of mis-classifying a threatened species as "secure" rises. In evaluating model performance, we

focused on discriminatory ability (using AUROC, which effectively considers all possible dichotomous

thresholds (Hand, 2009)), and did not assess accuracy (i.e., the average rate of prediction errors) (Ling

et al., 2003). However, in order to visualize predictions, we selected a binary classification threshold,

acknowledging that across different applications, other thresholds might be preferable (Walker, Leão,

Bachman, Bolam, & Nic Lughadha, 2020). To set a threshold, we assumed equal costs of

misclassification for threatened and secure species. For each taxon group, we selected the threshold

that maximized summed sensitivity and specificity on out-of-fold training data using the function

performance from the R package *ROCR* (Tobias, Sander, & Beerenwinkel, 2022).

Visualizing Predictions

We created state maps using the R packages *raster, tigris, sf, rnaturalearth,* and *ggplot2* (Etten et al.,

2023; Massicotte, South, Hufkens, & Philippe, 2023; Pebesma et al., 2023; H. Wickham et al., 2022) to

visualize areas in which many records of putatively threatened species were found. We divided the

study region into 8km x 8km grid cells, and each cell value represented the number of species

classified as threatened according to the threshold for each higher taxon (see above). Such maps can

be used to prioritize locations for additional survey and assessment resources, or to prioritize

locations for conservation funding, as locations with a high concentration of putatively threatened

species may both yield valuable data for evaluating species' conservation statuses, and warrant

precautionary protection.

Assessing Predictors

Random forest models are poorly suited to inference on causal relationships, but it is possible to

diagnose which predictors most strongly drive model predictions. Towards this end, we measured

"MDA" using the R package r*andomForest* (Bénard, Da Veiga, & Scornet, 2022; Breiman, 2001;

Ishwaran, 2007; Liaw et al., 2022). MDA approximates the degree to which each variable improves the

predictive accuracy of trees within the random forest. We ranked all variables based on their MDA

scores within each cross-validation run such that the variable with the greatest MDA score in a

particular run would be ranked first. We then plotted the distribution of MDA ranks across cross-

validation runs for the top five predictors for each taxon group. Variables exhibiting consistent, low

MDA ranks had relatively strong and/or clear associations with conservation status.

Literature Search

Although we cannot directly test whether predictions on unassessed species are correct without

formal status assessments, we can use external data sources as additional lines of evidence to

corroborate or challenge our predictions. To do this, we gathered external data for the ten species for

each taxon group that received the most and fewest votes for the category "threatened". For each of

these species, we first asked if the species had been recognized as threatened in nearby jurisdictions

by querying the species name on NatureServe (NatureServe, 2023)(Appendix S2). If we found that a

conservation status had been assigned elsewhere in Eastern North America (i.e., the continental USA

east of the Mississippi, and the Canadian provinces of Ontario, Québec, Nova Scotia, New Brunswick,

Newfoundland and Labrador, Prince Edward's Island, and tribal/first nations lands within this region),

we attempted to recover documentation of a previous status assessment. A match between our status

prediction and the recorded statuses in neighboring locations supports our prediction. Second, we

also examined contemporary geographic range data for the same top species. To do this, we

visualized recent occurrence records for each species using iNaturalist (iNaturalist Network, 2022). For

species predicted as "threatened" (and the opposite for species predicted as "secure"), if it appeared

that species current range limits may be in or near Maryland, we considered our predictions further

supported.

Results

Model Construction and Performance

On average, the discriminatory abilities were considerable for models for both taxon groups (mean

AUROC plants = 0.720, sd = 0.091; lepidopterans = 0.846, sd = 0.134; Fig. 1), with both models generally

outputting higher "threatened" probabilities to threatened species than to secure ones (Fig. 2). We did

not detect large differences in AUROC between the inner and outer folds, which would have indicated

unstable models. Hyperparameter selection was highly variable for lepidopteran and more consistent

for plant models (mean mtry for plant models = 2.1, sd = 0.4; lepidopterans = 16.5, sd = 28). Assuming

equal cost to misclassification for both threatened and secure species, the optimal fraction of random

forest votes to consider a species "threatened" (vs. "secure") for binary classification differed between

the two taxon groups: 0.606 for plants and 0.348 for lepidopterans. At these thresholds, both models

accurately classify previously assessed species.

Predictions for unassessed species

At the selected vote thresholds, our classifiers predicted 53% (502/940) of unassessed plant species

and 63% (798/1269) of unassessed lepidopteran species as threatened in Maryland (Table S1). The

most predictive variables differed between plants and lepidopterans (Fig. 3). For lepidopterans,

variables related to coldest temperatures tended to place among the top variables. These included

the mean last and first days of year above critical temperatures (5º and 10º C), the mean daily

minimum temperature in the coldest month, and the latitude of the northernmost Maryland record.

For plants, the top variables were often associated with variability, such as that measured in daily

mean air temperature during the coldest month, and in daily and annual temperature ranges. The

mean maximum relative humidity and maximum Maryland latitude were also important predictors for

plant conservation status.

Model predictions highlighted the northwesternmost part of the state as a possible conservation

hotspot for both plants and lepidopterans (Fig. 4). Although unassessed species predicted as

"threatened" were found throughout the state, their occurrence records appear concentrated in the

northwestern half of Maryland (Fig. 4). Furthermore, the mean per-cell predicted threat probabilities

showed a strong east-west gradient, with the highest mean threat probabilities in the westernmost

parts of Maryland (Fig. S1). In the case of plants, a higher number, but not high proportion, of species

were predicted as "threatened" in the Piedmont region in the center of the state (Figs. 4, S1).

When corroborating predictions with external data, we found that species with high "threatened"

probabilities were either unassessed in Eastern North America (5/20) or had been identified as

Vulnerable or Imperiled in at least one nearby jurisdiction (15/20). Recent occurrence patterns for

many of the species strongly predicted as "threatened" suggest they have a range primarily north of

Maryland, often with southern range extensions along the Appalachian Mountains (Table 1).

Lepidopterans receiving the highest "secure" probabilities were rarely listed as Vulnerable or

Imperiled in nearby jurisdictions (1/10). The case was different for plants, where 9/10 species had

been assessed as Vulnerable or Imperiled elsewhere in Eastern North America; primarily to the west

and north of Maryland. We note that many of these plant species had also been assessed as relatively

secure in coastal states from Delaware southwards, where climates tend to be warmer and where

their ranges primarily extend. These patterns also support, to some degree, our predictions.

Discussion

Here, we showed that random forest models can help cover the Rabinowitzian shortfall, using the biogeographic information about species contained in their occurrence records to evaluate conservation status. We showed that this machine learning method, originally developed for global analysis, applies at regional scales (here, the state of Maryland) and to a variety of taxa. We identified unassessed plant and lepidopteran species that are likely to be threatened and thus can be considered as having high conservation priorities. Our prediction mapping highlighted the western, montane section of the state as a potential hotspot for species monitoring, assessment, and protection. In the following sections, we discuss the implications of our results, explore the biogeographic trends recognized by our models, and describe how our study supports the use of similar methods to cover Rabinowitzian shortfalls in additional contexts.

Publicly Available Data and Simple Predictive Models Identify Regional-Scale Conservation Priorities
Several studies have demonstrated proof-of-concept for predictive modelling approaches for monitoring or conservation prioritization for plant and animal groups, but at continental or global scales (Bastin et al., 2019; Parsons, Pelletier, Wieringa, Duckett, & Carstens, 2022; Pelletier et al., 2018; Wieringa, 2022; Zhang, Campomizzi, & Lebrun-Southcott, 2022; Zizka et al., 2022). Because decisions about monitoring, research, and protection are often made at smaller spatial scales, before recommending adoption of these tools for conservation practice, it is crucial to determine if they still provide useful predictions at those scales (Meyer & Pebesma, 2021; Wyborn & Evans, 2021), as we have done here.

We anticipated several obstacles to prediction would be larger at the smaller geographical scale considered here, but also recognized mitigating advantages. Formally similar models trained on global vs. regional datasets may reflect fundamentally different biogeographic processes (e.g., continental vs. local), and are built on data of different quality and quantity. Disadvantages at regional scales include the fact that occurrence records are unlikely to cover complete species' ranges, thereby obscuring their ecological preferences (A. Lee-Yaw, L. McCune, Pironon, & N. Sheth, 2022; Phillips et al., 2009). Another limitation is that the variation of environmental covariates used is likely smaller, theoretically limiting the strength of any ecological signals and increasing the chance that model

predictions would be based on noise (Garcia-Rosello, Gonzalez-Dacosta, Guisande, & Lobo, 2023;

Hughes et al., 2021). Finally, and most simply, the number of taxa available on which to train models

at regional scales is likely to be much smaller, further reducing the chance that predictive models

reflect generally applicable trends.

Mitigating some of these concerns, at smaller scales, geographic datasets may have higher resolution

and higher quality, or be more consistently available across species or space, which should improve

the performance of resulting models. In this vein, we used complete, publicly-accessible, high-

resolution covariates at the regional scale, such as land use, land cover, and their history, as well as

slope – information that may be less readily available at much larger scales (Gogol-Prokurat, 2011).

Though subject to many of the environmental and spatial gaps and biases typical among large

biodiversity data (Bachman, Nic Lughadha, & Rivers, 2018; Gallagher et al., 2023; Kitzes & Shirley,

2016; Newbold, 2010), occurrence datasets from our study region benefit from high densities of

monetary wealth, human population, and roads, as well as proximity to flagship natural history

collections and older research universities. These advantages may, in this study, have

counterbalanced the limitations of the smaller spatial scale discussed above.

Overall, even as our regional datasets differ from those used in larger scale predictive models, they

appear to contain sufficient information to generate credible predictions. First, we found evidence

that the models were using stable – and perhaps robust – signals to generate predictions. Across

cross-validation runs of models for the same higher taxon, the same top predictors were frequently

identified, implying sample sizes and signal strengths were adequate within our regional-scale data.

Furthermore, the resulting models discriminated well between previously assessed threatened and

relatively secure species. We found binary thresholds for each higher taxon at which all previously

assessed species were classified accurately (albeit at quite different vote counts; 39% for

lepidopterans and 61% for plants). Relatively low representation of threatened species among

assessed lepidopterans (29%) versus plants (68%) may partly explain differences in the selected

thresholds (Peirce, 1884).

Our models showed strong performance on unseen data via cross-validation. Nevertheless, because

we do not suspect that the training data (assessed species) are a representative sample of all species,

we remain cautious in our claims of success. We know *a priori* that species have historically been

prioritized for assessment based on a variety of social and ecological factors that, while potentially

shared among assessed species, may not predict threat across other species that have not been

historical priorities for monitoring or protection (Bachman et al., 2019; Brummitt et al., 2015; Di Marco

et al., 2017; Fraixedas, Roslin, Antão, Poyry, & Laine, 2022; Ricketts, Daily, & Ehrlich, 2002). A strong

test of our predictions would require performing formal status assessments across a large,

representative subsample of previously unassessed species, which is beyond the scope of the present

study. Here, we informally validated selected predictions using external data, which were largely

corroboratory and supports the use of this predictive approach at such a regional scale.

Predicted Conservation Needs Reflects Biogeographic Trends

Generally, our predictions received support from the external data sources, with some caveats.

Among the 10 species per group that received the strongest predictions of threat from our models, all

had either previously been recognized as threatened in nearby jurisdictions or were unassessed.

These findings are compatible with elevated threat in Maryland, as well. Without exception, plant and

lepidopteran species that we predicted with high probability as "secure" in Maryland that had also

been assessed nearby were found to be secure in at least one jurisdiction in Eastern North America as

well, supporting our predictions. We note that among the unassessed plants predicted as most likely

to be secure, most were assessed as threatened elsewhere in the wider region, which does not

support our predictions. However, apparent trends in recent occurrences of species predicted as

"threatened" and "secure" with high probabilities may partly explain the latter result, and provide

biogeographic context and support to our predictions.

Our models predicted the likely statuses of species, but their occurrences point to spatial patterns

relevant in the context of regional conservation and climate change. First, we found that the species

our models predicted as "threatened" were concentrated in the northwestern part of the state, which

includes a central portion of the Appalachian Mountains. Second, our models predicted threat in both

taxon groups for northern-ranged, likely cold-adapted species, and relative security for southern-

ranged, likely warmer-adapted species. Montane habitats harbor a high proportion of total, rare, and

threatened species (Körner, 2004) and the Appalachians are no exception (Jenkins, Van Houtan,

Pimm, & Sexton, 2015). The Appalachian Mountains are home to species considered relicts from

glacial periods (Pickering, Kays, Meier, Andrew, & Yatskievych, 2003) that face increasing threats from

climate warming (La Sorte & Jetz, 2010; Zhu, Papeş, Armsworth, & Giam, 2022). It is thus expected that

our predictions would also highlight montane regions of Maryland as potential conservation hotspots.

Furthermore, the northwestern half of Maryland contains many habitats recognized as key for wildlife

conservation (Maryland Department of Natural Resources, 2015b), as well as priority areas for

conservation and monitoring (Maryland Department of Natural Resources, 2015a), constituting a

compelling stage for long-range conservation (Lawler et al., 2015). Beyond the Appalachian region,

our results recovered a large number (but not proportion) of plant species predicted as threatened in

the Piedmont region of the state, which has seen rapid development over the past 50 years (Maryland

Department of Planning, 2010). Sheer densities of roads, institutions, money, and people in the center

of the state may explain this result. However, that section of Maryland is experiencing stark habitat

loss and fragmentation, and may also be an emerging priority for monitoring, status assessment, and

conservation.

Even as they are not designed for inference, our models recovered ecological signals both within (and

between) the plant and lepidopteran models in measures of variable importance. These signals point

towards mechanisms that drive threat for some species and relative security for others. The latitude of

the northernmost Maryland occurrence was a top predictor variable for both lepidopteran and plant

status, with species with occurrences at higher latitudes being predicted as "threatened" with higher

probabilities. Otherwise, top predictor variables differed between the taxon groups, and likely reflect

the ecology of each group. For plants, variables associated with niche breadth were important, along

with latitude and humidity. Niche breadth has long been associated with vulnerability (Slatyer, Hirst,

& Sexton, 2013), and plants restricted to microclimates or habitat types that are rare in Maryland are

likely under threat from anthropogenic change. For lepidopterans, the top variables reflected low-

temperature tolerance, which we interpret to mean that more cold-associated lepidopterans tend to

be more threatened in Maryland. Abundance and range shifts and declines in response to changing

climate are well-documented in lepidopterans (Crossley et al., 2022; Hill, Kawahara, Daniels,

Bateman, & Scheffers, 2021), and it is likely that some of Maryland's most threatened insect fauna are

retreating to relatively cool and high-elevation refugia.

Applications

Our methodology is not intended to replace species conservation assessments (Zizka et al., 2022) nor

to stand in for highly contextual, place-based ecological knowledge (Wyborn & Evans, 2021), but

rather as a tool that can inform prioritization in a resource-limited conservation world (Sinclair et al.,

2018). As resources for protection are already biased towards monitoring and research, approaches

like ours can use existing data to accelerate movement of conservation resources towards active

conservation work (Buxton et al., 2020). Maryland's 2015-2025 State Wildlife Action Plan calls for

greater coherence in monitoring efforts, especially for species that have yet to be recognized as "of

greatest conservation need" (Maryland Department of Natural Resources, 2015c). Towards these

ends, first, our models provide guidance about assessed and unassessed plant and lepidopteran

species (table S1) in a coherent, quantitative framework that can immediately integrate new data as

they become available. Second, beyond individual species predictions, we also produce regional

maps (Fig. 3, S1), that highlight priority areas where conservation needs likely intersect; these maps

can support proactive, place-based conservation (Cardillo & Meijaard, 2012). Third, we tentatively

identify ecological features of threatened and relatively secure plants and lepidopterans, which

warrant further investigation and may be applicable in other regions or to other taxa as well. Our

results echo calls to protect climate refugia and unique habitat types within Maryland, and perhaps to

assist climate-driven migration (Maryland Department of Natural Resources, 2015a). Finally, as the

data inputs for the approach we used are publicly available, and we have made scripts containing our

workflows open (CITE CODE REPOSITORY), any interested party can easily reproduce our analysis,

update our predictions with new data, or extend our method to additional taxa and locations.

Future applications of our approach would ideally involve stakeholders (Knight et al., 2008). In our

study, direct decision-maker input during model construction might have altered which data sources

were used or how species were grouped into individual models (here, we generated separate models

for plants and lepidopterans), and would also have affected how the models were trained on those

data. Decision-makers, uniquely positioned to name the problems associated with different types of

misclassification errors, could help translate these problems into loss functions and classifier training

criteria (Hand, 2009; Walker et al., 2020). These criteria help tune the model to a specific application

(e.g., by avoiding predicting that species are threatened when they are not, or the opposite), via

hyperparameter values and vote threshold selections. This approach is fit for stakeholder-driven

model construction: models were fit on a personal laptop; the decision trees at the heart of random

forests are concrete, easily apprehended, and thus accessible; the data we used are publicly available

and well-documented. Overall, this modelling approach is well-suited to use by local and regional

conservation planners and decision-makers.

Conclusion

Predictive methods cannot replace localized knowledge systems nor systematic species assessments

(Betts et al., 2020). Instead, in a fast-changing world in which most species, including threatened

species, remain unassessed and data-deficient, they extend the conservation utility of existing data.

Here, we showed that a predictive method for prioritizing unassessed species, originally developed for

global applications, can be flexibly applied at smaller, regional scales and to a variety of taxa. We

show how using occurrence records as input data enables spatially explicit predictions, which can

highlight priority locations, and could facilitate place-based or ecosystem-level conservation. We

highlighted Maryland (USA) unassessed plant and lepidopteran species likely to be threatened, and

identified the western part of the state and Appalachian Mountains as hotspots for such likely-

threatened species. Such emergent results can help resolve a central conservation mystery: which

taxa are most vulnerable to which drivers (Isbell et al., 2023; Sánchez-Bayo & Wyckhuys, 2019). The

random forests we used "learn" from existing, publicly accessible datasets using simple decision

trees, require relatively little computational resources or natural history expertise to implement, and

can be used "in house" by researchers, planners, or managers at regional scales. As a result of the

immediacy of our method, as well as the encouraging results from the plant and lepidopteran case

studies we present here, we advocate for regional applications of similar predictive tools to better

identify conservation priorities. Finally, future applications of these methods could more explicitly

integrate stakeholders, as well as consider other niche (e.g., species interactions or co-occurrences)

into both threat assessment and conservation prioritization.

References

A. Lee-Yaw, J., L. McCune, J., Pironon, S., & N. Sheth, S. (2022). Species distribution models rarely

predict the biology of real populations. *Ecography*, *2022*(6), 1–16.

https://doi.org/10.1111/ecog.05877

Bachman, S. P., Field, R., Reader, T., Raimondo, D., Donaldson, J., Schatz, G. E., & Lughadha, E. N.

(2019). Progress, challenges and opportunities for Red Listing. *Biological Conservation*,

234(November 2018), 45–55. https://doi.org/10.1016/j.biocon.2019.03.002

Bachman, S. P., Nic Lughadha, E. M., & Rivers, M. C. (2018). Quantifying progress toward a

conservation assessment for all plants. *Conservation Biology*, *32*(3), 516–524.

https://doi.org/10.1111/cobi.13071

Bastin, J.-F., Finegold, Y., Garcia, C., Mollicone, D., Rezende, M., Routh, D., … Crowther, T. W. (2019).

The global tree restoration potential. *Science*, *365*(6448), 76–79.

https://doi.org/10.1126/science.aax0848

Bénard, C., Da Veiga, S., & Scornet, E. (2022). Mean decrease accuracy for random forests:

inconsistency, and a practical solution via the Sobol-MDA. *Biometrika*, *109*(4), 881–900.

https://doi.org/10.1093/biomet/asac017

Betts, J., Young, R. P., Hilton-Taylor, C., Hoffmann, M., Rodríguez, J. P., Stuart, S. N., & Milner-Gulland,

E. J. (2020). A framework for evaluating the impact of the IUCN Red List of threatened species.

Conservation Biology, *34*(3), 632–643. https://doi.org/10.1111/cobi.13454

Breiman, L. (2001). Random Forest. *Machine Learning*, (45), 5032.

Brummitt, N. A., Bachman, S. P., Griffiths-Lee, J., Lutz, M., Moat, J. F., Farjon, A., … Lughadha, E. M. N.

(2015). Green plants in the red: A baseline global assessment for the IUCN Sampled Red List

Index for Plants. *PLoS ONE*, *10*(8), 1–22. https://doi.org/10.1371/journal.pone.0135152

Brun, P., Zimmermann, N. E., Hari, C., Pellissier, L., & Karger, D. N. (2022). Global climate-related

predictors at kilometre resolution for the past and future Earth System Science Data Discussions.

Earth System Science Data, *14*(June), 5573–5603. Retrieved from

https://doi.org/10.16904/envidat.332,

Buxton, R. T., Avery-Gomm, S., Lin, H. Y., Smith, P. A., Cooke, S. J., & Bennett, J. R. (2020). Half of

resources in threatened species conservation plans are allocated to research and monitoring.

Nature Communications, *11*(1), 1–8. https://doi.org/10.1038/s41467-020-18486-6

Caetano, G. H. O., Chapple, D. G., Grenyer, R., Raz, T., Rosenblatt, J., Tingley, R., … Roll, U. (2022).

Automated assessment reveals extinction risk of reptiles is widely underestimated across space

and phylogeny. *BioRxiv*. https://doi.org/10.1101/2022.01.19.477028

Callaghan, C. T., Nakagawa, S., & Cornwell, W. K. (2021). Global abundance estimates for 9,700 bird

species. *Proceedings of the National Academy of Sciences of the United States of America*, *118*(21),

1–10. https://doi.org/10.1073/pnas.2023170118

Cardillo, M., & Meijaard, E. (2012). Are comparative studies of extinction risk useful for conservation?

Trends in Ecology and Evolution, *27*(3), 167–171. https://doi.org/10.1016/j.tree.2011.09.013

Cardoso, P., Erwin, T. L., Borges, P. A. V., & New, T. R. (2011a). The seven impediments in invertebrate

conservation and how to overcome them. *Biological Conservation*, *144*(11), 2647–2655.

https://doi.org/10.1016/j.biocon.2011.07.024

Cardoso, P., Erwin, T. L., Borges, P. A. V, & New, T. R. (2011b). The seven impediments in invertebrate

conservation and how to overcome them. *Biological Conservation*, *144*(11), 2647–2655.

https://doi.org/10.1016/j.biocon.2011.07.024

Chamberlain, S. (2020). Package 'natserv.'

Chamberlain, S., Oldoni, D., Barve, V., Desmet, P., Geffert, L., McGlinn, D., … Waller, J. (2022). Package

'rgbif .' Retrieved from https://github.com/ropensci/rgbif

Chamberlain, S., Szoecs, E., Foster, Z., Arendsee, Z., Boettiger, C., Ram, K. ;, … Grenié, M. (2022). taxize:

Taxonomic Information from Around the Web. Retrieved from https://docs.ropensci.org/taxize/

(website),%0Ahttps://github.com/ropensci/taxize (devel), https://taxize.dev%0A(user manual)

Crisfield, V., Blanchet, F. G., & Gravel, D. (2022). How and why species are rare: A mechanistic

reappraisal of the Rabinowitz rarity framework.

https://doi.org/10.22541/au.166748076.66307375/v1

Crossley, M. S., Meehan, T. D., Moran, M. D., Glassberg, J., Snyder, W. E., & Davis, A. K. (2022). Opposing

global change drivers counterbalance trends in breeding North American monarch butterflies,

(May), 1–10. https://doi.org/10.1111/gcb.16282

Department of the Interior, U. S. F. & W. S. (1973). Endangered Species Act of 1973, As Amended

through the 108th Congress. *Endangered Species Act Of 1973*, 1–44. Retrieved from

papers://a25d78e7-d6e0-4b2b-9b83-8d9ae6ec0592/Paper/p134

Di Marco, M., Chapman, S., Althor, G., Kearney, S., Bensancon, C., Butt, N., … Watson, J. E. M. (2017).

Changing trends and persisting biases in three decades of conservation science. *Global Ecology*

and Conservation, *10*, 32–42. https://doi.org/10.1016/j.gecco.2017.01.008

Eastern Shore Regional GIS Cooperative. (2021). Statewide 1m slope map. Retrieved May 22, 2023,

from

https://lidar.geodata.md.gov/imap/rest/services/Statewide/MD_statewide_slope_m/ImageServ

er

Etten, J. Van, Sumner, M., Cheng, J., Baston, D., Bevan, A., Bivand, R., … Mosher, S. (2023). Package

'raster.' Retrieved from https://rspatial.org/raster BugReports

Fraixedas, S., Roslin, T., Antão, L. H., Poyry, J., & Laine, A. L. (2022). Nationally reported metrics can't

adequately guide transformative change in biodiversity policy. *Proceedings of the National*

Academy of Sciences of the United States of America, *119*(9), 1–4.

https://doi.org/10.1073/pnas.2117299119

Free, C. M., Jensen, O. P., Anderson, S. C., Gutierrez, N. L., Kleisner, K. M., Longo, C., … Walsh, J. C.

(2020). Blood from a stone: Performance of catch-only methods in estimating stock biomass

status. *Fisheries Research*, *223*(November 2019), 105452.

https://doi.org/10.1016/j.fishres.2019.105452

Gallagher, R. V., Allen, S. P., Govaerts, R., Rivers, M. C., Allen, A. P., Keith, D. A., … Adams, V. M. (2023).

Global shortfalls in threat assessments for endemic flora by country. *Plants People Planet*,

(February), 1–14. https://doi.org/10.1002/ppp3.10369

Garcia-Rosello, E., Gonzalez-Dacosta, J., Guisande, C., & Lobo, J. M. (2023). GBIF falls short of

providing a representative picture of the global distribution of insects. *Systematic Entomology*,

(May 2022), 1–9. https://doi.org/10.1111/syen.12589

Gogol-Prokurat, M. (2011). Predicting habitat suitability for rare plants at local spatial scales using a

species distribution model. *Ecological Applications*, *21*(1), 33–47. https://doi.org/10.1890/09-

1190.1

Hand, D. J. (2009). Measuring classifier performance: A coherent alternative to the area under the ROC

curve. *Machine Learning*, *77*(1), 103–123. https://doi.org/10.1007/s10994-009-5119-5

Hastie, T., Tibshirani, R., & Friedman, J. (2017). *The elements of statistical learning: data mining,*

inference, and prediction (2nd ed.). Springer. Retrieved from

http://www.springerlink.com/index/D7X7KX6772HQ2135.pdf

Heberling, J. M., Miller, J. T., Noesgaard, D., Weingart, S. B., & Schigel, D. (2021). Data integration

enables global biodiversity synthesis. *Proceedings of the National Academy of Sciences of the*

United States of America, *118*(6), 1–7. https://doi.org/10.1073/pnas.2018093118

Hill, G. M., Kawahara, A. Y., Daniels, J. C., Bateman, C. C., & Scheffers, B. R. (2021). Climate change

effects on animal ecology: butterflies and moths as a case study. *Biological Reviews*, *96*(5), 2113–

2126. https://doi.org/10.1111/brv.12746

Hochkirch, A., Samways, M. J., Gerlach, J., Böhm, M., Williams, P., Cardoso, P., … Dijkstra, K. D. B.

(2021). A strategy for the next decade to address data deficiency in neglected biodiversity.

Conservation Biology, *35*(2), 502–509. https://doi.org/10.1111/cobi.13589

Hughes, A. C., Orr, M. C., Ma, K., Costello, M. J., Waller, J., Provoost, P., … Qiao, H. (2021). Sampling

biases shape our view of the natural world. *Ecography*, *44*(9), 1259–1269.

https://doi.org/10.1111/ecog.05926

iNaturalist Network. (2022). Observations · iNaturalist. Retrieved May 17, 2023, from

https://www.inaturalist.org/observations

Isbell, F., Balvanera, P., Mori, A. S., He, J. S., Bullock, J. M., Regmi, G. R., … Palmer, M. S. (2023). Expert

perspectives on global biodiversity loss and its drivers and impacts on people. *Frontiers in*

Ecology and the Environment, *21*(2), 94–103. https://doi.org/10.1002/fee.2536

Ishwaran, H. (2007). Variable importance in binary regression trees and forests. *Electronic Journal of*

Statistics, *1*, 519–537. https://doi.org/10.1214/07-EJS039

Jenkins, C. N., Van Houtan, K. S., Pimm, S. L., & Sexton, J. O. (2015). US protected lands mismatch

biodiversity priorities. *Proceedings of the National Academy of Sciences of the United States of*

America, *112*(16), 5081–5086. https://doi.org/10.1073/pnas.1418034112

Kitzes, J., & Shirley, R. (2016). Estimating biodiversity impacts without field surveys: A case study in

northern Borneo. *Ambio*, *45*(1), 110–119. https://doi.org/10.1007/s13280-015-0683-3

Knapp, W. M., & Naczi, R. F. C. (2020). *Vascular plants of Maryland, USA: A comprehensive account of the*

state's botanical diversity. Washington, D. C.: Smithsonian Institution Scholarly Press.

Knight, A. T., Cowling, R. M., Rouget, M., Balmford, A., Lombard, A. T., & Campbell, B. M. (2008).

Knowing but not doing: Selecting priority conservation areas and the research-implementation

gap. *Conservation Biology*, *22*(3), 610–617. https://doi.org/10.1111/j.1523-1739.2008.00914.x

Körner, C. (2004). Mountain biodiversity, its causes and function. *Ambio*, *33*(SPEC. ISS. 13), 11–17.

https://doi.org/10.1007/0044-7447-33.sp13.11

Kuhn, M., Wing, J., Weston, S., Williams, A., Keefer, C., Engelhardt, A., … Hunt, T. (2022). Package

'caret.'

La Sorte, F. A., & Jetz, W. (2010). Projected range contractions of montane biodiversity under global

warming. *Proceedings of the Royal Society B: Biological Sciences*, *277*(1699), 3401–3410.

https://doi.org/10.1098/rspb.2010.0612

Lawler, J. J., Ackerly, D. D., Albano, C. M., Anderson, M. G., Dobrowski, S. Z., Gill, J. L., … Weiss, S. B.

(2015). The theory behind, and the challenges of, conserving nature's stage in a time of rapid

change. *Conservation Biology*, *29*(3), 618–629. https://doi.org/10.1111/cobi.12505

Liaw, A., Wiener, M., Breiman, L., & Cutler, A. (2022). Package "randomForest."

Ling, C. X., Huang, J., & Zhang, H. (2003). AUC: A statistically consistent and more discriminating

measure than accuracy. *IJCAI International Joint Conference on Artificial Intelligence*, 519–524.

Maryland Department of Natural Resources. (2015a). Conservation actions. In *2015-2025 State Wildlife*

Action Plan. Retrieved from

https://dnr.maryland.gov/wildlife/Documents/SWAP/SWAP_Chapter7.pdf

Maryland Department of Natural Resources. (2015b). Maryland's Key Wildlife Habitats. In *Maryland*

State Wildlife Action Plan 2015-2025. Retrieved from

https://dnr.maryland.gov/wildlife/Documents/SWAP/SWAP_Chapter4.pdf

Maryland Department of Natural Resources. (2015c). Monitoring and Effectiveness Measures. In

Maryland State Wildlife Action Plan 2015-2025.

Maryland Department of Planning. (2010). *A summary of land use trends in Maryland*. Retrieved from

https://planning.maryland.gov/Documents/OurProducts/landuse/MDP2010_LU_Summary.pdf

Maryland Natural Heritage Program. (2021a). *List of Rare, Threatened, and Endangered Animals*.

Maryland Natural Heritage Program. (2021b). *Rare, Threatened, and Endangered Plants of Maryland*.

Retrieved from http://dnr2.maryland.gov

Massicotte, P., South, A., Hufkens, K., & Philippe, M. (2023). Package 'rnaturalearth.' Retrieved from

https://docs.ropensci.org/rnaturalearth/

Meyer, H., & Pebesma, E. (2021). Predicting into unknown space? Estimating the area of applicability

of spatial prediction models. *Methods in Ecology and Evolution*, *12*(9), 1620–1633.

https://doi.org/10.1111/2041-210X.13650

Mokany, K., & Ferrier, S. (2011). Predicting impacts of climate change on biodiversity: A role for semi-

mechanistic community-level modelling. *Diversity and Distributions*, *17*(2), 374–380.

https://doi.org/10.1111/j.1472-4642.2010.00735.x

NatureServe. (2023). NatureServe Explorer. In *NatureServe*. Retrieved from

https://explorer.natureserve.org/

Newbold, T. (2010). Applications and limitations of museum data for conservation and ecology, with

particular attention to species distribution models. *Progress in Physical Geography*, *34*(1), 3–22.

https://doi.org/10.1177/0309133309355630

651 Nongame and endangered species conservation act, Code of Maryland § (2022).

652 Panter, C. T., Clegg, R. L., Moat, J., Bachman, S. P., Klitgård, B. B., & White, R. L. (2020). To clean or not

653  to clean: Cleaning open-source data improves extinction risk assessments for threatened plant

654  species. *Conservation Science and Practice*, *2*(12), 1–14. https://doi.org/10.1111/csp2.311

655 Parsons, D. J., Pelletier, T. A., Wieringa, J. G., Duckett, D. J., & Carstens, B. C. (2022). Analysis of

656  biodiversity data suggests that mammal species are hidden in predictable places. *Proceedings of*

657  *the National Academy of Sciences of the United States of America*, *119*(14).

658  https://doi.org/10.1073/pnas.2103400119

659 Pebesma, E., Bivand, R., Racine, E., Sumner, M., Cook, I., Keitt, T., … Dunnington, D. (2023). Package

660  'sf.' Retrieved from https://r-spatial.github.io/sf/

661 Peirce, C. S. (1884). The numerical measure of the success of predictions. *Science*, *4*(93), 453–454.

662  https://doi.org/10.1126/science.ns-4.93.453.b

663 Pelletier, T. A., Carstens, B. C., Tank, D. C., Sullivan, J., & Espíndola, A. (2018). Predicting plant

664  conservation priorities on a global scale. *Proceedings of the National Academy of Sciences*,

665  *115*(51), 13027–13032. https://doi.org/10.1073/pnas.1804098115

666 Phillips, S. J., Dudík, M., Elith, J., Graham, C. H., Lehmann, A., Leathwick, J., & Ferrier, S. (2009). Sample

667  selection bias and presence-only distribution models: Implications for background and pseudo-

668  absence data. *Ecological Applications*, *19*(1), 181–197. https://doi.org/10.1890/07-2153.1

669 Pickering, J., Kays, R., Meier, A., Andrew, S., & Yatskievych, K. (2003). The Appalachians. In P. R. Gil, R.

670  A. Mittermeier, C. G. Mittermeier, J. P. Pilgrim, G. Fonseca, W. R. Konstant, & T. Brooks (Eds.),

671  *Wilderness Earth's Last Wild Places* (pp. 458–467). Conservation International, Washington, DC.

672  Retrieved from http://www.discoverlife.org/co/

673 Probst, P., Wright, M. N., & Boulesteix, A. L. (2019). Hyperparameters and tuning strategies for random

674  forest. *Wiley Interdisciplinary Reviews: Data Mining and Knowledge Discovery*, *9*(3), 1–15.

675  https://doi.org/10.1002/widm.1301

676 Rabinowitz, D. (1981). Seven forms of rarity. *The Biological Aspects of Rare Plant Conservation*, 205–

677  217. https://doi.org/10.1177/004728702237415

678 Regan, H. M., Hierl, L. A., Franklin, J., Deutschman, D. H., Schmalbach, H. L., Winchell, C. S., & Johnson,

679  B. S. (2008). Species prioritization for monitoring and management in regional multiple species

680  conservation plans. *Diversity and Distributions*, *14*(3), 462–471. https://doi.org/10.1111/j.1472-

681  4642.2007.00447.x

Ricketts, T. H., Daily, G. C., & Ehrlich, P. R. (2002). Does butterfly diversity predict moth diversity?

Testing a popular indicator taxon at local scales. *Biological Conservation*, *103*(3), 361–370.

https://doi.org/10.1016/S0006-3207(01)00147-1

Sánchez-Bayo, F., & Wyckhuys, K. A. G. (2019). Worldwide decline of the entomofauna: A review of its

drivers. *Biological Conservation*, *232*(January), 8–27.

https://doi.org/10.1016/j.biocon.2019.01.020

Sinclair, S. P., Milner-Gulland, E. J., Smith, R. J., McIntosh, E. J., Possingham, H. P., Vercammen, A., &

Knight, A. T. (2018). The use, and usefulness, of spatial conservation prioritizations. *Conservation*

Letters, *11*(6), 1–7. https://doi.org/10.1111/conl.12459

Slatyer, R. A., Hirst, M., & Sexton, J. P. (2013). Niche breadth predicts geographical range size: A

general ecological pattern. *Ecology Letters*, *16*(8), 1104–1114. https://doi.org/10.1111/ele.12140

Tobias, A., Sander, O., & Beerenwinkel, N. (2022). Package ' ROCR .'

Wagner, D. L., Fox, R., Salcido, D. M., & Dyer, L. A. (2021). A window to the world of global insect

declines: Moth biodiversity trends are complex and heterogeneous. *Proceedings of the National*

Academy of Sciences of the United States of America, *118*(2), 1–8.

https://doi.org/10.1073/PNAS.2002549117

Walker, B. E., Leão, T. C. C., Bachman, S. P., Bolam, F. C., & Nic Lughadha, E. (2020). Caution Needed

When Predicting Species Threat Status for Conservation Prioritization on a Global Scale.

Frontiers in Plant Science, *11*(April), 1–4. https://doi.org/10.3389/fpls.2020.00520

Wickham, H., Chang, W., Henry, L., Pedersen, T. L., Takahashi, K., Wilke, C., … Dunnington, D. (2022).

ggplot2: Elegant Data Visualisations Using the Grammar of Graphics. R package version 3.4.0.

Wickham, J., Stehman, S. V., Sorenson, D. G., Gass, L., & Dewitz, J. A. (2021). Thematic accuracy

assessment of the NLCD 2016 land cover for the conterminous United States. *Remote Sensing of*

Environment, *257*, 112357. https://doi.org/10.1016/j.rse.2021.112357

Wieringa, J. G. (2022). Comparing predictions of IUCN Red List categories from machine learning and

other methods for bats. *Journal of Mammalogy*, *103*(3), 528–539.

https://doi.org/10.1093/jmammal/gyac005

Wyborn, C., & Evans, M. C. (2021). Conservation needs to break free from global priority mapping.

Nature Ecology and Evolution, *5*(10), 1322–1324. https://doi.org/10.1038/s41559-021-01540-x

Zhang, X., Campomizzi, A. J., & Lebrun-Southcott, Z. M. (2022). Predicting population trends of birds

worldwide with big data and machine learning. *Ibis*, *164*(3), 750–770.

713   https://doi.org/10.1111/ibi.13045

714 Zhu, G., Papeş, M., Armsworth, P. R., & Giam, X. (2022). Climate change vulnerability of terrestrial

715   vertebrates in a major refuge and dispersal corridor in North America. *Diversity and Distributions*,

716   *28*(6), 1227–1241. https://doi.org/10.1111/ddi.13528

717 Zizka, A., Andermann, T., & Silvestro, D. (2022). IUCNN – Deep learning approaches to approximate

718   species' extinction risk. *Diversity and Distributions*, *28*(2), 227–241.

719   https://doi.org/10.1111/ddi.13450

721 **Acknowledgements:**

722 This research was supported by University of Maryland startup funds awarded to AE.

Tables and figures

Figure 1. Distribution of area under the receiver operating curve (AUROC) values on unseen data in
10x repeated 10-fold cross-validation. For plants (a) mean AUROC was moderate (mean = 0.72; sd =
0.09). For lepidopterans (b), model performance was stronger but more variable (AUROC mean = 0.84,
sd = 0.13). In each panel, mean AUROC is shown with a red vertical line; dashed vertical line indicates
the expected performance of a model making random guesses (AUROC of 0.50).

Figure 2. Random forest predictions from final models on occurrence records of assessed (a-d) and unassessed (e, f) species. Models distinguished threatened and relatively secure species in the training dataset (lighter points in panels a and b, darker in c and d), and identify a concentration of putatively threatened, unassessed species in Western Maryland (trend towards darker points in western portion of maps in e, f).

Figure 3. Distribution of variable importance ranks for the five predictors with the lowest mean MDA

ranks (i.e., those with higher importance across cross-validation runs) for plants (green) and

lepidopterans (yellow). Maximum ranks exceed the total number of variables because each level of

categorical variables receives its own MDA measure.

a   plants

b   lepidopterans

745   **Figure 4.** Number of species classified as threatened by the final models using the taxon group-

746   specific probability thresholds (0.614 for plants, 0.39 for lepidoptera), per 8km x 8km grid cell across

747   the state of Maryland. Colors represent number of species (see figure scales).

Table 1. Top and bottom ten species per taxon group, based on the predicted probability of being threatened. Known conservation status from other jurisdictions in Eastern North America is shown, if available. Codes indicate state/province conservation status and the state/province in question. Row coloring indicates species range type. Orange: iNaturalist occurrence records appeared concentrated north of Maryland, with a southward extension along the Appalachian Mountains; blue: occurrences concentrated north of Maryland, without an Appalachian extension; green: occurrences neither concentrated to the north of Maryland nor clustered along the Appalachian Mountains.

taxon group	genus	species	predicted status	vote fraction (threatened)	Status elsewhere	notes
plants	Pyrola	elliptica	threatened	0.95	S1 VA, NC; S2 DE, S4 WV, NJ	
	Poa	autumnalis	threatened	0.94	SX MI, IL, SH NJ, S1 PA, S3 DE, S4 NC, S5 SC, VA, KY	
	Pyrola	americana	threatened	0.94	SX KY, IL, S2 NJ, NC, TN, IN; S4 NJ, VT; S5 VA, WV	
	Eragrostis	hypnoides	threatened	0.938	SH ME, NH; S2 NC, DE; S3 NJ, VT, S4 NY, VA, WV, KY	
	Bidens	connata	threatened	0.936	S1 RI; SH WV; S4 VA; QE; S5 NJ, NY, KY	
	Viola	macloskeyi	threatened	0.932	S1 DE; S3 GA, NC; S4 VA, WV; S5 NY, QE, ON	
	Dichanthelium	longiligulatum	threatened	0.928	S1 Nj, S3 NC; S4 VA, KY	
	Carex	gynandra	threatened	0.926	S1 OH; S2 De; S4 Nj, NC; S5 VA, WV, NY, ON, KY	
	Gentiana	clausa	threatened	0.926	S2 QE, RI; S3 NC, OH; S4VA, WV, NY, NJ	
	Selaginella	rupestris	threatened	0.926	S1 OH, NB, NS; S2 IA, NJ; S3 WV, VT, NC, IL, MA; S4 NY; S5 ON, QE, GA	
	Elymus	virginicus	secure	0.134	S4 WV, S5 ON, NC, IL, IN, KY	
	Juncus	dichotomus	secure	0.14	S1 PA, WV, OH; S2 NY, S4 DE, GA, S5 NC, VA, NJ	
	Carex	longii	secure	0.144	S1 OH, WV, S2 KY, PA; S3 NY; S4 NJ, DE; S5 VA	
	Cyperus	retrorsus	secure	0.15	S1 MA, NY; S2 KY; S4 NJ; S5 GA, NC, VA	
	Polygala	mariana	secure	0.154	S1 TN, S2 NJ, S4 DE, NC; S5 KY, VA	
	Eupatorium	album	secure	0.16	S1 CT, OH, S3 IN, WV; S4 DE; S5 NC, VA, KY	
	Lobelia	puberula	secure	0.162	S1 PA; S4 KY; S5 NC, WV, VA	

	Panicum	*virgatum*	secure	0.162	S2 QE; S3 VT; S4 ON, KY; S5 IL, NC, VA, NC
	Fimbristylis	*autumnalis*	secure	0.166	S1 VT; S2 ME; S3 IL; S4 NC, ON, QE; S5 KY, VA, WV, NJ, NY
	Dichanthelium	*scoparium*	secure	0.168	SH WV, MA; S1 IL, IN, OH, PA, NY; S4 NJ; S5 NC, VA, DE
lepidopterans	*Olethreutes*	*bipartitana*	threatened	0.936	
	Scoparia	*cinereomedia*	threatened	0.93	
	Euphyia	*intermediata*	threatened	0.928	
	Haploa	*confusa*	threatened	0.926	
	Catocala	*blandula*	threatened	0.916	S1 KY, IN; S4 PA
	Panthea	*acronyctoides*	threatened	0.916	S3 VA; S4 NB, ON
	Argyresthia	*calliphanes*	threatened	0.898	S1 CT; S2 MA; S4 NH, NB; S5 ON
	Cingilia	*catenaria*	threatened	0.898	
	Chrysanympha	*formosa*	threatened	0.896	S1 IN; S3 NC; S4 NY, PA, NB; S5 ON
	Idia	*diminuendis*	threatened	0.894	S2 QE; S3 NH; S4 PA
	Hyphantria	*cunea*	secure	0	S5 ON; ID, PA
	Marimatha	*nigrofimbria*	secure	0	S1 NY, S4 PA
	Nomophila	*nearctica*	secure	0	S5 PA
	Palthis	*asopialis*	secure	0	S3 ON, S4 NY
	Hemaris	*diffinis*	secure	0.002	S4 ON, S5 PA, KY
	Atteva	*punctella*	secure	0.004	S4 ON, S5 PA
	Hylephila	*phyleus*	secure	0.004	S4 KY, WV; S5 IN, MS, AL, FL, GA, SC, NC, VA, **MD (See appendix S1)** Syn. *Papilio phyleus*
	Malacosoma	*americana*	secure	0.004	S5 ON, PA
	Pyrrharctia	*isabella*	secure	0.004	S4 ID, S5 ON, PA
	Actias	*luna*	secure	0.006	S4 QE, S5 ON, PA, VT