Endonucleolytic RNA cleavage drives changes in gene expression during the innate immune response

Agnes Karasik[1], Hernan A. Lorenzi[2,] Andrew V. DePass[1] and Nicholas R. Guydosh[1]

[1]Laboratory of Biochemistry and Genetics, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD 20892

[2]TriLab Bioinformatics Core, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD 20892

Keywords: RNase L, endonuclease, Integrated Stress Response, ZAKα, JNK, ribosome collisions

Key summary:

- Activated RNase L acts together with other dsRNA-sensing pathways to induce a strong immune response via JNK and p38 signaling
- Generic RNA fragmentation plays a key role in inducing transcription, potentially through ribosome collisions
- Activation of RNase L inhibits the effects of eIF2α phosphorylation
- Translation of the antiviral IFIT mRNAs is inhibited by active RNase L

Funding

This research was supported by the Intramural Research Program of the NIH, The National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) (DK075132 to N.R.G.) and the Postdoctoral Research Associate Training Program (PRAT) at the National Institute of General Medical Sciences (NIGMS) (FI2GM128743 to A.K.).

Acknowledgements

We are thankful for the insightful discussions with Drs. Bret Hassel, Alan Hinnebusch, Jon Lorsch, Stacy Horner and Tom Dever. A549 cells were a kind gift from Dr. Bernie Moss. We also thank Dr. Behdad Afzali and Dr. Daniel Chauss for their assistance with electroporation experiments and the assistance of the genomic cores at the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) and the National Heart, Lung, and Blood Institute (NHLBI). A.K. is grateful to have been chosen for a MOSAIC K99/R00 award (K99 GM143484) and for associated mentoring support from the American Society of Cell Biology partnership activities.

Summary

During viral infection, several dsRNA sensors are activated in the cell and trigger signaling that leads to changes in gene expression. One of these sensors activates an endonuclease, RNase L, that cleaves many types of RNA in the cell. However, how the resultant widespread RNA decay affects gene expression is not fully understood. Here we found that gene expression changes caused by activating dsRNA sensing pathways are tuned by RNase L activation, pointing to intricacy in the innate immune response where multiple inputs are integrated to create an optimized output. We show that RNase-L-dependent RNA fragmentation induces the activation of the Ribotoxic Stress Response, potentially through ribosome collisions. The p38 and JNK pathways that are activated as part of this response promote outcomes that inhibit the virus, such as programmed cell death. We also show that RNase L appears to limit the translation of genes that are regulated by another dsRNA-induced pathway, the Integrated Stress Response. Intriguingly, we found the activity of the generic endonuclease, RNase A, recapitulates many of the same molecular phenotypes as activated RNase L, demonstrating how widespread RNA cleavage events can directly evoke an antiviral program.

Introduction

Viral infections are recognized by Pattern Recognition Receptors (PRRs) that trigger the innate immune response and ultimately promote elimination of pathogens. Foreign nucleic acid (RNA or DNA) sensing is one of the major ways viral pathogens are recognized (Hartmann, 2017). Viruses have either an RNA genome or produce RNA intermediates, and thus sensing of viral RNA in the cytoplasm is vital for recognizing and fighting against viruses. A subset of PRRs recognize double stranded RNA (dsRNA) (Hartmann, 2017). One group within these PRRs is the interferon-induced family of Oligo-Adenylate Synthetases (OAS1-3) that produce an oligomer of adenylates chained together through 2'-5' linkages (2-5A) upon dsRNA binding. 2-5A can selectively bind Ribonuclease L (RNase L) to induce dimerization that is required for its transition from a latent to an activated state. Activated RNase L then cleaves a variety of host and viral RNAs, triggering many downstream processes and physiological changes in the cell. These changes include the loss of most mRNAs (and therefore gene expression in the cell), production of chemokines and cytokines, activation of inflammasomes, inhibition of cell migration, autophagy, senescence, and cell death (Banerjee et al., 2015; Castelli et al., 1997; Donovan et al., 2017; Silverman, 2007). Intriguingly, the function of RNase L appears to extend beyond its role in protecting against viruses. RNase L mutations increase risk for multiple types of cancer, including prostate and breast cancer (Carpten et al., 2002; Casey et al., 2002; Lee et al., 2013; Long et al., 2013), and autoimmune diseases (Magg et al., 2021).

Besides RNase L, another interferon-regulated PRR, Protein Kinase R (PKR), can induce reduction in protein synthesis in the cell through phosphorylation of eukaryotic Initiation Factor 2α (eIF2α). Phospho-eIF2α inhibits initiation of translation of the majority of mRNAs, presumably to limit viral protein production, and triggers a number of changes in the cell that are collectively referred to as the Integrated Stress Response (ISR) (Costa-Mattioli and Walter, 2020; Hinnebusch et al., 2016; Young and Wek, 2016). Additionally, the dsRNA receptors RIG-I and MDA5 induce the interferon and inflammatory response by activating signaling that leads to transcription. Activation of the different dsRNA response pathways in the cytoplasm can have intersecting functions, wherby activated RIG-I induces gene expression, while activated RNase L and PKR reduce gene

expression. However, how the different dsRNA sensing pathways, particularly RNase L,

act together to achieve a strong antiviral response is largely unknown.

Activation of RNase L can induce and alter transcription (Andersen et al., 2009;

Burke et al., 2019; Malathi et al., 2005; Rath et al., 2019). During RNase L activation,

many proinflammatory genes and cytokines (e.g CXCL8) were found to be upregulated

in early microarray studies (Malathi *et al.*, 2005). Importantly, more recent RNA-

sequencing (RNA-seq) studies found that most transcripts are degraded (~60 to >99%)

by RNase L. Interestingly, immune transcripts, such as interferon-β (IFNβ), are able to

compensate for this degradation via transcriptional upregulation or resistance to cleavage

due to underrepresentation of favorable RNase L cleavage motifs (UU or UA) (Burke *et*

al., 2019; Rath *et al.*, 2019). Furthermore, many Interferon Stimulated Genes (ISGs) are

upregulated during the dsRNA response in both WT and *RNASEL* KO cells, suggesting

that they do not depend on RNase L, whereas upregulation of cytokines, such as CXCL8,

CXCL2, and other immune genes, including *IFNL1* and *2*, and *IFIT3*, were at least

partially dependent on RNase L (Burke *et al.*, 2019). These observations suggest that

active RNase L together with the other dsRNA pathways may amplify transcription of

antiviral genes. In some cell lines, RNase-L cleaved RNA fragments play a role in

activating RIG-I (Malathi et al., 2007; Manivannan et al., 2020) and the inflammasome

(Chakrabarti et al., 2015), however these were not observed in human lung carcinoma

cells (Burke *et al.*, 2019; Karasik et al., 2021; Rath *et al.*, 2019). One limitation of the

existing studies that used RNA-seq is that RNase L was activated by the dsRNA mimic

(poly I:C), an approach that simultaneously activates many other pathways (Burke *et al.*,

2019; Rath *et al.*, 2019). These pathways can mask RNase L-specific effects and make

it difficult to assess how RNase L activation and the other dsRNA pathways act together

to induce the antiviral immune response.

Importantly, activation of RNase L can also induce pathways related to the

Ribotoxic Stress Response (RSR) (Iordanov et al., 2000; Li et al., 2004), which triggers

apoptosis through transcription activated by JNK (c-JUN N-terminal Kinase) signaling,

which is a mitogen-activated protein kinase (MAPK) family member. While several

MAP2K family members can phosphorylate JNK (Iordanov *et al.*, 2000), the mechanism

of how they are activated by RNase L has remained unclear. The RSR was originally

shown to be activated by defective translation, so it was suggested that pathways related

to the RSR were induced when RNase L damaged ribosomes by cleaving rRNA (Iordanov

et al., 2000; Li *et al.*, 2004; Siddiqui and Malathi, 2012). However, more recently, it was

revealed that ribosomes with cleaved rRNAs are functional, suggesting that ribosome

damage may not be responsible for activating the RSR (Rath *et al.*, 2019). In addition,

the RSR was recently shown to include activation of the MAP3K family member ZAKα

that senses collisions between ribosomes (Wu et al., 2020). ZAKα phosphorylates the

MAP2Ks that, in turn, phosphorylate JNK, and another MAPK, p38 (Robinson et al., 2022;

Vind et al., 2020a; Vind et al., 2020b). These observations suggest that activation of ZAKα

by ribosome collisions could occur when RNase L induces the RSR. However, the source

of any potential ribosome collisions in RNase L activated cells remains unknown. One

possible source of ribosome collisions was suggested by our previous observation that

activation of RNase L leads to translation of fragmented mRNAs (Karasik *et al.*, 2021).

Ribosomes that translate mRNA fragments would have the possibility of reaching a 3'

end and stalling, potentially leading to the formation of ribosome collisions.

More broadly, the overall loss of translated mRNA in the cell upon RNase L

activation is expected to change the balance of free ribosomal subunits and mRNAs (Rath

et al., 2019). This altered balance could change the efficiency with which the remaining,

full-length transcripts are translated. Consistent with this, we previously observed an

increase in the relative proportion of ribosomes in alternative Open Reading Frames

(ORFs) or "altORFs" following RNase L activation (Karasik *et al.*, 2021). We proposed

that translation of these altORFs became more favorable as excess free ribosomes

promoted translation initiation at the 5' ends of the RNA fragments created by RNase L.

However, how RNase L and other dsRNA pathways lead to changes in translation on

mRNAs remained unclear.

Here we investigated how direct activation of RNase L, and attendant global mRNA

degradation, contributes to the activation of the innate immune response. We found that

RNase-L mediated RNA cleavage strengthened multiple parts of the antiviral

transcriptional response and, in particular, induced ZAKα activation, presumably via

ribosome collisions. In addition, activation of RNase L also changed how efficiently intact

mRNAs were translated, in some cases by inhibiting the effect of eIF2α phosphorylation.

Our results highlight the multifaceted role of RNase L and endonucleolytic cleavage in
promoting antiviral functions.

Results

Activation of RNase L induces transcription and phosphorylation of ZAKα

To measure changes in the transcriptome under activation of RNase L, but not
other components of the dsRNA-induced innate immune response, we performed RNA-
seq on wild type (WT) and *RNASEL* KO cells that were transfected with 2-5A (1 μM), the
direct and specific activator of RNase L (Figure 1A). Since RNase L cleaves rRNA, we
utilized a rRNA degradation assay to assess the level of RNase L activation (Floyd-Smith
et al., 1981). Across all our replicates, we found that rRNA degradation was similar (within
a 2-fold difference, Figure S1A). Based on previous reports that used spike-in oligos for
normalization (Burke *et al.*, 2019; Rath *et al.*, 2019), these levels of rRNA degradation
correspond to a loss of ~60% to >99% of individual mRNAs in the cell. We therefore
assume that the absolute abundance of most mRNAs in the treated cells under study
here substantially decreases ("baseline loss").

Then, we analyzed differential expression (DE) patterns in 2-5A treated vs
untreated cells by DEseq2 (Love et al., 2014). The DE changes in our experiments
represent differences in relative abundance of the small pool of mRNA that is not
degraded. We infer that mRNAs that appear upregulated (described throughout as
relative "upregulation") actually decrease in abundance, but to a lesser extent than the
general population, due to resistance to RNase L cleavage activity or compensation by
active transcription. The DE values therefore characterize how the global loss of mRNA
varies across the transcriptome. We found hundreds of differentially expressed genes
("DE genes," $p_{adjusted}<0.05$) in 2-5A treated WT cells, revealing that the level of these
mRNAs significantly differs from the baseline loss that generally affects all transcripts
(Figure 1B). 2-5A treatment did not induce many changes in differential gene expression
in *RNASEL* KO cells (Table S1), as expected, since 2-5A is a specific activator of RNase
L. DE genes that appeared to be upregulated by RNase L activation (Figure 1B and S1B,
defined as log2 fold changes >1) included transcripts that encode proinflammatory

proteins and cytokines (e.g. *CXCL2* and *CCL20*), such as interleukins (e.g. *CXCL8* and

IL6), suggesting that direct RNase L activation stimulates an inflammatory response. In

contrast, we did not detect an interferon (IFN) response, such as increased transcript

levels for *IFNB1* (Burke *et al.*, 2019; Karasik *et al.*, 2021; Rath *et al.*, 2019). These findings

are largely consistent with previous RNA-seq data from human cells (WT vs *RNASEL*

KO) that were transfected with poly I:C (Burke *et al.*, 2019; Rath *et al.*, 2019).

We noted that some of the genes that appear to be upregulated in 2-5A treated

cells encode transcription factors or activators, such as *JUN*, *FOSB*, *FOXA2*, *GDF15* and

ATF3 (Figure 1B). This is consistent with previous findings for some of these genes, such

as *FOXA2* and *GDF15* (Malathi *et al.*, 2005; Rath et al., 2015). Gene Ontology (GO)

analysis of molecular functions offered additional support since the top categories of DE

genes were related to transcription regulation (Figure 1C). This finding suggests that the

apparent increase in some transcript levels in our DE analysis could be due to new

transcription that is induced by these factors. Since RNase L reduces global mRNA levels

in the cell, it is unclear whether the observed relative upregulation of these transcription

factor genes in our RNA-seq data is sufficient to increase production of protein. Thus, we

probed selected transcription factor protein levels by western blotting. Interestingly, we

found that protein levels for FOSB and JUN increased in 2-5A treated WT, but not in

RNASEL KO cells (Figure 1D and S1C). This finding suggests that activation of RNase L

can actually raise the abundance of these proteins, leading to downstream transcriptional

activation.

Next, we assessed which pathways are triggered in 2-5A treated cells that could

explain the observed transcriptional upregulation. We found that several target genes in

the JNK, p38, ERK, and NF-κB pathways appeared to be increased in our data,

suggesting that at least some of these pathways (which are known to have overlapping

targets) are activated. For instance, the transcription factor GDF15 controls inflammation,

cell repair, and growth, and is known to be upregulated by the activity of the JNK and

ERK kinases when prostate cancer cells are treated with 2-5A (Malathi *et al.*, 2005). To

dissect which of these pathways are specifically responsible for the observed

transcriptional changes in our dataset, we used western blotting to assess activation

through phosphorylation of these pathways. We found that RNase L can activate JNK

and p38 (Figure 1E and S1D), but we did not detect activation of pathways related to ERK (Figure S1E, ERK phosphorylation) or NF-κB (Figure S2, p65 nuclear fraction). These results suggest that transcriptional changes observed in the cells here are driven, in part, by the activation of JNK and p38. We also noticed that multiple "DUSP" mRNAs, such as *DUSP1* and *8*, that encode negative regulators of the JNK pathway were upregulated (Figure 1B and S1B), suggesting a need for modulation of JNK when RNase L is activated. Interestingly, JNK activation can be apoptotic or induce autophagy. Here we observed the induction of proapoptotic genes, such as JUN, but not genes expressed during autophagy, such as ATF5 (Dhanasekaran and Reddy, 2017).

It has been suggested that activation of RNase L can trigger the RSR, which can lead to downstream activation of the JNK and p38 pathway (Iordanov *et al.*, 2000; Wu *et al.*, 2020). Since it was recently demonstrated that this can involve the kinase ZAKα, we tested whether ZAKα is activated in 2-5A treated cells by examining its phosphorylation state by western blotting. We found that ZAKα is strongly activated due to 2-5A treatment in WT but not in *RNASEL* KO cells, indicating that phosphorylation of JNK and p38 under RNase L might occur through ribosome collisions during RNase L activation (Figure 1E-F and S1D).

The OAS-RNase L pathway acts together with other dsRNA sensing pathways to achieve a strong activation of innate immune signaling

We next asked whether activation of RNase L modulates the effects of other pathways that are typically activated during infection by other dsRNA sensors (Malathi *et al.*, 2007; Manivannan *et al.*, 2020). To simultaneously activate multiple pathways and test RNase L's role, we transfected WT and *RNASEL* KO A549 cells with poly I:C, a dsRNA mimic (0.25 µg/ml) and performed RNA-seq (Figure 2A). Poly I:C is known to strongly activate OAS and, in turn, RNase L as well as other dsRNA sensing pathways, including PKR, and RIG-I/MDA5. First, we assessed the levels of RNase L activation by measuring the level of rRNA degradation and comparing it to that in cells treated by the direct activator, 2-5A. We found that in all replicates of poly I:C and 2-5A treated cells, the level of rRNA degradation was comparable (within 2-fold), suggesting a similar level of RNase L activation (Figure S1A).

Next, we determined the DE genes in WT cells upon poly I:C treatment with
DEseq2. We found that poly I:C treatment resulted in a higher number of relatively
upregulated DE genes (~1200, log2 fold changes >1, $p_{adjusted}$ <0.05) as compared to 2-
5A treatment (~700 genes) and that additional pathways were activated, as expected
given the broader effects of poly I:C (Figure 2B). In particular, in both WT and *RNASEL*
KO cells, PKR and RIG-I/MDA5 pathways were activated, as judged by the levels of
eIF2α phosphorylation (shown in (Karasik *et al.*, 2021)) and the increase in ISGs (Figure
2C and Table S1), respectively. Consistent with this, we observed that *PPP1R15A*
(*GADD34*), a gene involved in the ISR (see below) and known to be transcribed as a
result of both interferon production via the IRF3/7 pathway and PKR activation (Dalet et
al., 2017), also exhibited a relative increase compared to other transcripts (~32-fold) in
poly I:C treated WT cells (Table S1 and Figure 2C).

Notably, we also found that most genes that are differentially expressed upon 2-
5A treatment were also differentially expressed in poly I:C treated cells (Figure 2B, note
overlap between yellow- and magenta-colored circles). This suggests that these changes
are largely unimpaired in cells treated with poly I:C and are not masked by the effects of
the other activated dsRNA pathways. We noted that many genes that did not meet the
criteria for significant differential expression in WT cells treated with 2-5A changed in
ways that closely mirrored changes observed in cells treated with poly I:C (Figure 2C,
Pearson's R = 0.87). This correlation indicates that the small relative gene expression
changes in 2-5A treated cells that do not meet the DE threshold may indicate partial
activation of pathways that get amplified when other dsRNA sensors are triggered in poly
I:C treated cells.

To more closely investigate how simultaneous activation of RNase L and other
pathways triggered by dsRNA affects the JNK and p38 pathways, we examined the
mRNA levels of target genes and phosphorylation of the transcription factors themselves.
Based on our finding above that 2-5A treatment (RNase L activation alone) activates the
JNK and p38 pathways (Figure 1E, Figure 2C, blue dots), we defined a subset of 54
transcripts that are commonly associated with JNK and p38 and are activated in 2-5A
treated samples (see Methods section for more details). Since ~50% of the genes
downstream of JNK and p38 overlap, we analyzed these MAPK pathways together. We

assessed transcriptional induction of JNK/p38 downstream targets under activation of
RNase L only (WT + 2-5A), other dsRNA pathways only (RNASEL KO cells + poly I:C),
or both pathways (WT + poly I:C) (Figure 2D). As expected based on the selection of
targets, we found that these JNK/p38-related mRNAs were highly induced in WT + 2-5A
treated cells. We also noted some activation due to other dsRNA pathways alone (KO +
poly I:C). Interestingly, when combined, the RNase L and other dsRNA pathways together
suggested the effects may be additive since the level of induction was highest in WT cells
treated with poly I:C (Figure 2D). We therefore asked whether these trends were
supported by the level of JNK and p38 phosphorylation, as measured by western blotting
(Figure 2E). In support of additivity between the RNase L and the other dsRNA pathways,
we observed the strongest phosphorylation of JNK and p38 when both RNase L and the
other dsRNA pathways were triggered compared with either one alone (Figure 2E and F).
In addition, we found that ZAKα, which we showed to be phosphorylated by 2-5A
(Figure 1E) was also phosphorylated in poly I:C treated WT cells to a similar extent,
suggesting that activation of ZAKα by RNase L is not affected by the other dsRNA
pathways (Figure 2E). Importantly, ZAKα phosphorylation was not detectable in RNase
L KO cells that were treated with poly I:C, suggesting its activation (presumably via
ribosome collisions) requires RNase L activity. This model therefore implies that JNK and
p38 phosphorylation in the absence of RNase L (Figure 2E) must use an MAP3K other
than ZAKα and that, together with ZAKα, affords the strongest induction of p38/JNK
targets in cells treated with poly I:C (Figure 1E).
Next, we assessed if activation of RNase L affects the interferon response that is
triggered by dsRNA binding to MDA5 and RIG-I (Figure 2C, red dots). We defined a
subset of ISGs (N = 45, see Methods for details) that were upregulated in WT cells treated
with poly I:C. We then looked at the activation of these genes across our experimental
conditions (Figure 2D, right panel). We found no activation of these genes with 2-5A
treatment, consistent with previous RNA-seq findings in A549 cells (Burke *et al.*, 2019;
Karasik *et al.*, 2021; Rath *et al.*, 2019), but contrasting with models where RNase L
cleaved RNA fragments stimulate other dsRNA sensors, potentially in other cell lines
(Malathi *et al.*, 2007; Manivannan *et al.*, 2020). In cells treated with poly I:C, we observed
some decrease in activation for this limited subset of genes in *RNASEL* KO cells

compared to WT cells (Figure 2D). This finding suggests that RNase L may have a minor

role in augmenting activation of other dsRNA sensors in these cells (Malathi *et al.*, 2007;

Manivannan *et al.*, 2020). We further investigated these trends by western blotting (Figure

2E and F). In these experiments, we visualized the phosphorylation of IRF3, one of the

main transcription factors that induces ISGs in response to interferon. We were unable to

detect activation of IRF3 in cells treated with 2-5A but observed activation in cells treated

with poly I:C, in agreement with our RNA-seq analysis. We also observed a small

decrease (~20%) in activation of IRF3 in *RNASEL* KO cells treated with poly I:C, in line

with our observations from RNA-seq, and a model where RNase L may have some role

in affecting the response to interferon.

We also asked whether RNase L can contribute to the other pathways that play a

role in the dsRNA-induced innate immune response, such as NF-κB and ERK. Activation

of NF-κB (p65) was only detectable in poly I:C treated cells and the levels of activation

appeared to be similar in both WT and *RNASEL* KO cells, indicating that RNase L has no

further effects on this pathway (Figure S2). We also found that 2-5A or poly I:C treatments

had no effect on ERK activation in A549 cells, suggesting that the observed effects in our

experiments are not due to activation of ERK (Figure S2).

Taken together, our analysis shows that activated RNase L can act together with

other dsRNA pathways to promote p38 and JNK signaling, but has only minor effects on

the induction of ISGs (Figure 2G).

A generic endonuclease, RNase A, induces transcriptional changes similar to

those triggered by RNase L

Next, we wanted to assess what changes directly result from the global RNA

cleavage that occurs after RNase L activation. RNase L is known to interact with several

proteins, including those that make up the cytoskeleton and the nucleopore complex

(Bisbal et al., 1995; Burke et al., 2021; Le Roy et al., 2005; Nogimori et al., 2019; Nogimori

et al., 2020; Tnani et al., 1998). Thus, it is possible that some of the observed changes

upon RNase L activation and dimerization arise from functional protein-protein

interactions and not RNA cleavage. To further investigate how global RNA cleavage

changes gene expression, we electroporated a generic RNase, RNase A, in *RNASEL* KO

A549 cells and performed RNA-seq after 4.5 hours (Figure 3A). RNase A (bovine

pancreatic ribonuclease) is a small, generic endonuclease that primarily cleaves at

pyrimidine bases (U and C) in ssRNAs (Nogués et al., 1995). As a control, we

electroporated similar amounts of Bovine Serum Albumin (BSA), that is similar in size,

but has no RNase activity. We anticipated that RNase A would cause widespread RNA

decay but would lack RNase L specific interactions. During our experiments, we followed

RNase A activity by monitoring rRNA cleavage, similar to our above experiments with

RNase L. We observed that rRNAs are degraded in RNase A electroporated cells, but

with different RNA cleavage patterns (Figure S3A). Based on RNase A's specificity

(cleavage at Us and Cs), we expect that most RNA species (including mRNAs) would be

targeted and thus fragmented by RNase A.

We found >1000 differentially expressed genes in RNase A electroporated cells

(Table S1). As noted for RNase L activation above, gene expression is reported in

absolute terms from the average global loss of mRNA; therefore, genes that appear to be

upregulated may simply be less downregulated than those in the same treatment group.

Interestingly, most upregulated DE genes found in 2-5A treated cells overlap with those

identified here for RNase A activation (~65 % of those in the RNase L dataset, Figure 3B

and C, Figure S3B) and the expression levels were correlated (R = 0.62, Figure 3C). This

suggests that most RNase L-dependent DE changes are the direct result of RNA

cleavage rather than potential indirect non-enzymatic activities.

Next, we assessed if targets of the JNK signaling pathway were upregulated in

RNase A electroporated cells as they were in 2-5A treated ones. We found that the JNK

and p38 pathway-related mRNAs were upregulated in RNase A electroporated samples

(Figure 3D). JNK activation was also confirmed by western blotting (Figure 3E). Strikingly,

we found that p38 and ZAKα are also phosphorylated (Figure 3E). Our results therefore

indicate that transcriptional changes, particularly through the JNK and p38 pathways, can

be induced by generic RNA cleavage, regardless of the effector endonuclease.

RNA fragmentation by generic RNases induces altORF translation

Previously, we examined the translation that takes place in RNase L activated cells

by using ribosome profiling to find where ribosomes translate the reduced pool of mRNAs

(Karasik *et al.*, 2021). One of the most striking findings from our previous study is that the
relative abundance of ribosome footprints increased in non-coding regions of transcripts,
including the 5' and 3' UTR regions, as well as out-of-frame parts of coding sequences.
This phenomenon likely results when ribosomes initiate translation at the first available
open reading frame on the 3' mRNA fragments that are generated by RNase L, a
phenomenon that we termed "altORF translation" (Karasik *et al.*, 2021). In particular, we
observed strong altORF translation in the 5' and 3' UTRs. The model predicts that the
activity of any generic endonuclease should lead to the same outcome. To test this
hypothesis, we electroporated RNase A into *RNASEL* KO A549 cells, as described
above, and performed ribosome profiling on those same samples after 4.5 hours (Figure
4A).

The major hallmarks of altORF translation that we previously observed in RNase
L activated cells were recapitulated in the cells where RNase A was electroporated
(Figure 4B-D). The most striking signature of altORF translation is an increase in
ribosome footprints in the 3'UTR. This region of the mRNA is rarely translated under
normal conditions, but we found ribosomes in this region when RNase L was active
(Karasik *et al.*, 2021). Cells where RNase A was active exhibited similar ribosome profiling
patterns in the 3'UTRs, consistent with the altORF translation model (Figure 4B, *ACTG1*
as an example gene model). We also investigated whether this signature was present in
RNase-A treated samples globally. To assess characteristics of altORF translation in a
global manner, we computed the ratios of ribosome profiling footprint levels in 3' UTR
regions and the main ORFs of mRNAs (3'UTR:ORF ratios) as we did before for active
RNase L. We found that in all RNase treated samples, 3'UTR:ORF ratios increased
compared to controls (Figure 4C) (Karasik *et al.*, 2021). Increased 5'UTR translation was
also observed in RNase L activated cells (Karasik *et al.*, 2021) and we found similar
patterns when computing 5'UTR:ORF ratios for RNase A treated cells, further confirming
altORF translation in these samples (Figure S4A). Increased reads in the 5' and 3'UTRs
were also observed in metagene averages at the start and stop codon of main ORFs for
RNase A electroporated samples as compared to the control (Figure 4D and S4B), further
confirming ribosome occupancy and potential altORF translation in the UTR regions.

The data from cells that were electroporated with RNase A suggest the
phenomenon of altORF translation takes place whenever any generic endonuclease, not
only RNase L, is active in the cytoplasm. However, electroporation of RNase A is artificial,
and we wondered whether another naturally activated endonuclease would trigger the
translation of altORFs. The endonuclease Ire1 is homologous to RNase L and plays a
pivotal role in the Unfolded Protein Response (UPR) since it is activated by protein
misfolding in the ER (Walter and Ron, 2011). Ire1 is required for cytoplasmic splicing of
the *XBP1* mRNA through recognition of specific sequences and structural elements.
However, Ire1 can also cleave more generically at UGC sequence motifs. This broader
activity is referred to as Regulated Ire1 Dependent Decay (RIDD) and could potentially
lead to altORF translation (Guydosh et al., 2017; Hollien et al., 2009). To test this
hypothesis, we analyzed a publicly available ribosome profiling dataset where human
cells were treated with thapsigargin, a potent inducer of the UPR and Ire1 (Woo et al.,
2018) (GEO ID: GSE103719). We found that thapsigargin-treated cells also exhibited
characteristics of altORF translation, including increased 3'UTR:ORF ratios (Figure 4E).
These results show that RNase L is not the only naturally activated endonuclease that
can lead to the translation of altORFs.
414

RNA fragmentation can modulate translational efficiency
The major effect on gene expression of active RNase L is that the majority of
mRNAs are lost, resulting in reduced protein synthesis for most genes. However,
transcriptional upregulation counteracts this and, in some cases, overrides (Figure 1D)
this effect. In addition, it is conceivable that gene expression from the remaining mRNAs
could be tuned at the translational level and thus further refine the amount of protein
synthesized from the remaining transcriptome. As an example of this, an increase in the
translation of some host mRNAs was reported during vaccinia virus infection, which is a
potent activator of RNase L (Dai et al., 2017). In this way, mRNA degradation could have
a role in regulating translation.
To investigate whether RNase L activity alters which mRNAs are translated, we
took advantage of our ribosome profiling data (reported previously (Karasik *et al.*, 2021))
and with its matched RNA-seq data (reported here) assessed translational efficiency (TE)

of coding sequences. The TE metric is defined as the ribosome footprint level in a coding

region, normalized to the corresponding RNA-seq level, and serves as a proxy for the

relative loading of a transcript by ribosomes. We therefore plotted the change in ribosome

footprint levels against the corresponding change in RNA-seq level (Figure 5A) for genes

where either of these values were found to change significantly by DESeq2 (see

Methods). Those genes with large changes in TE will appear furthest from the diagonal

(slope >1 indicates TE increase; <1 TE decrease). We noted several cases where dots

were shifted from the diagonal, suggesting functionally important regulation of translation.

Some of the largest changes we observed affected a group of transcripts that were

expressed at higher relative levels when RNase L was active at the mRNA level (via 2-

5A or poly I:C in WT, but not in *RNASEL* KO, cells) but were counterbalanced by

downregulation of expression at the translational level (lower TE, Figure 5A, dots to the

right of the origin and below the dotted diagonal). This category includes the extracellular

matrix proteins *MUC5AC* and *MUC5B* (Figure 5A). We directly computed the TE for these

genes for each replicate and found the effect of reduced TE to be consistent across

replicates (Figure 5B). We also noticed an enrichment for genes encoding proteins

involved in transcription or related to transcription in this category (e.g. *KLF6*, *ZNF296*),

suggesting that RNase L tunes the production of proteins involved in transcription at the

translational level.

In another class of genes, we observed that relative mRNA levels were increased

upon RNase L activation (via 2-5A or poly I:C in WT, but not in *RNASEL* KO, cells) and

that the level of ribosome footprints also increased to a larger extent and would therefore

be expected to augment gene expression (Figure 5A, dots to the right of the origin and

above the dotted diagonal). This category included *CXCL1-3* and *8*, genes that encode

chemokines and cytokines. We also directly computed the TE for these genes for each

replicate and found the effects to be consistent (Figure 5B). Interestingly, we found that

mRNAs encoded by the mitochondrial genome also fell into this category (Figure 5A,

orange dots). This observation is consistent with a study of TEs during vaccinia infection

where higher levels of translation of mitochondrially-encoded transcripts in conjuction of

higher oxidative phosphorylation activity was reported. This was proposed to enhance

viral replication due to the high energy demand of viral replication (Dai *et al.*, 2017). Our

data therefore suggest that this phenomenon is driven by RNase L.

Intriguingly, we also observed a trend among mRNAs that are induced by dsRNA

pathways apart from RNase L, including *IFIT1-3* and *IFNB1* (induced by interferon) and

the gene *GADD34* (induced by interferon and the ISR as noted above). These genes

exhibited ~2 fold higher TEs in poly I:C treated *RNASEL* KO cells as compared to WT

cells (Figure 5A, dots shift above diagonal in third vs second panel), suggesting that

RNase L may play some role in reducing their expression at the translational level as a

way to tune their transcriptional induction.

RNase L activation modulates the Integrated Stress Response

Given the role of *GADD34* in the ISR, we further examined the ribosome profiling

data for clues as to how its translation is regulated by RNase L. The ISR is activated when

PKR binds dsRNA and phosphorylates the translation initiation factor eIF2α, ultimately

resulting in a loss of its availability (Wek et al., 2006). The downstream effects of eIF2α

phosphorylation are collectively known as the integrated stress response (ISR)

(Hinnebusch *et al.*, 2016; Young and Wek, 2016). Phosphorylation of eIF2α generally

limits the loading of ribosomes on mRNAs in the cell, reducing their TE. However, there

is a small class of mRNAs, including *ATF4*, *GADD34* (*PPP1R15A*), and *CHOP* (*DDIT3*),

that undergo the opposite response and are translationally activated (Young and Wek,

2016). ATF4 and CHOP are transcription factors that induce expression of stress-

responsive genes, while GADD34 is a phosphatase of eIF2α and thus serves as a

counterbalance to ISR activation (Fusakio et al., 2016; Novoa et al., 2001). We found that

relative transcript levels increased in poly I:C treated WT and *RNASEL* KO cells as

compared to respective controls for *GADD34* (as noted above, 5-32-fold) and *CHOP* (5-

9-fold). This is consistent with work showing both genes are transcriptionally induced by

ATF4 (Ma et al., 2002; Ma and Hendershot, 2003). *GADD34* is likely further induced by

the interferon-induction IRF3 pathway (Dalet *et al.*, 2017). The *GADD34* mRNA was

identified as a strong target of RNase L driven degradation (Burke *et al.*, 2019) and thus

absolute levels of *GADD34* are likely lower in WT cells as compared to *RNASEL* KO cells.

The transcripts for *ATF4*, *CHOP*, and *GADD34* contain one or more open reading frames

in their 5'UTR (uORFs) with unique features that allow them to suppress translation of the

main ORF in non-stressed cells, but activate it when eIF2α becomes phosphorylated

(Dever et al., 2023; Young and Wek, 2016). As noted above, the TE for *GADD34* was

observed to increase 2-4-fold in in poly(I:C) treated cells (WT or *RNASEL* KO) (Figure

6A). We observed a ~2-fold change in TE for *ATF4* in WT and *RNASEL* KO cells and a

more modest increase for *CHOP* (Figure S5A). These data are consistent with ISR

induction in our WT and *RNASEL* KO cells.

Several explanations could account for our observation that the change in

GADD34 TE was less in WT vs *RNASEL* KO cells (Figure 6A). One possibility is that

levels of Phospho-eIF2α are decreased in *RNASEL* WT vs KO cells, leading to reduced

translational activation via uORF regulation in the WT cells. Alternatively, RNase L could

directly affect TE by altering the shift in translation from the uORFs to the main ORF. To

determine whether RNase L had an effect on uORF-based regulation, we quantified the

ratio of ribosome footprints in the 5'UTR and compared it to footprints that mapped to the

main ORF (Figure 6B-C, see Methods). We noted a substantial decrease in uORF

translation relative to main ORF translation in poly I:C treated WT or *RNASEL* KO cells

due to eIF2α phosphorylation, as expected (Dever *et al.*, 2023). However, this decrease

was less in the case of the WT cells compared to *RNASEL* KO cells (Figure 6B-C),

suggesting that RNase L activation may have some role in affecting this shift. Similar

RNase L dependent shifts in 5'UTR vs main ORF ribosome footprints for *ATF4* and *CHOP*

suggest this regulation may also extend to those genes (Figure S5B-C). It has been

shown that the *GADD34* mRNA is heavily targeted by RNase L, making the mRNA and

protein product in poly I:C treated *RNASEL* KO cells much more detectable than in WT

cells (Burke *et al.*, 2019). Consistent with this, we also found that GADD34 protein levels

increased in poly I:C-treated *RNASEL* KO cells (Figure 6D). Thus, this mechanism where

RNase L suppresses main ORF translation augments the dominant underlying trend in

expression that is driven by mRNA decay.

Taken together, these results show a role of RNase L in inhibiting the translation

of *GADD34*, thus modulating the ISR during the innate immune response, and suggests

the RNase L may have a broader role in regulating translational control.

Translation of IFIT mRNAs is inhibited by activation of RNase L

Since we observed that poly I:C induced higher translation efficiency in *RNASEL* KO vs WT cells for the genes *IFIT1*, *2* and *3*, (Figure 5A, right panel), we asked whether this trend could be explained by uORF-mediated translational control as in the case of the ISR-related mRNAs. IFIT proteins are one of the key antiviral proteins that are stimulated by interferon and inhibit viral translation (Fensterl and Sen, 2015). Since these genes have low expression under conditions without poly I:C stimulation, we focused on data from cells treated with poly I:C.

To determine whether uORFs were translated on these mRNAs, we examined ribosome densities in their 5'UTR in ribosome profiling data. We found that 5'UTRs of *IFIT1*, *2* and *3* contained ribosome peaks at start codons of potential uORFs (Figure 7A and Figure S6A, green lines). The uORFs used near-cognate start codons (CUG or GUG instead of AUG, Figure 7A and Figure S6A). These data suggest a model for translational control where the uORFs inhibit main ORF translation and it is conceivable that this inhibition could be regulated by eIF2α phosphorylation. As in the case of *GADD34*, this apparent uORF inhibition may be regulated by RNase L since there was lower 5'UTR translation relative to main ORF translation in poly I:C treated *RNASEL* KO vs WT cells that was correlated with higher TE (Figure 7B and C). These data show that expression of these genes may be controlled by inhibitory uORFs and that this mechanism may depend on RNase L (Figure 7D).

Discussion

In this study, we examined how direct RNase L activation by 2-5A, and attendant global RNA degradation in the cell, affects gene expression. Our work shows the global loss in mRNA is compensated by increases in transcription and tuned by changes in translation.

We showed that one way mRNA cleavage leads to new transcription is via activation of the RSR. RNase L activation alone (via 2-5A) turns on the p38 and JNK MAPK pathways independently of other dsRNA pathways that are activated by poly I:C (Figure 1). While JNK activation was previously documented in several cell lines, simultaneous p38 activation was not observed when RNase L alone was activated

(Iordanov *et al.*, 2000; Malathi *et al.*, 2005). Since poly I:C treatment strongly induces p38,

it is possible that weak activation of p38 was masked in previous studies, or the extent of

activation could be cell line dependent. The activation of JNK is known to be associated

with the RSR (Iordanov et al., 1997; Wang et al., 2005) and it was initially suggested to

be involved in innate immunity due to damage to rRNA by RNase L (Iordanov *et al.*, 2000;

Li *et al.*, 2004). However, more recent studies showed that even with their rRNA damaged

by RNase L, ribosomes can still translate (Rath *et al.*, 2019), suggesting that rRNA

cleavage may not lead to ribosome disassembly and therefore sensing of damaged

rRNAs. Recently, it was found that the RSR can be activated by the ribosome collision

sensor and MAP3K, ZAKα (Vind *et al.*, 2020b; Wu *et al.*, 2020). We established that ZAKα

can also be triggered directly by RNase L activity, but not by the other dsRNA pathways.

Moreover, it is likely that other mechanisms, beyond the RSR, exist to activate

transcription in response to global RNA degradation. For instance, global RNA loss in the

cytoplasm was proposed to free up RNA binding proteins that, in turn, modulate

transcription due to their relocation to the nucleus (Hartenian and Glaunsinger, 2019).

Our observation raises the question of why ribosome collisions might be occurring

in these cells. Ribosome collisions are known to form when a ribosome stalls during

elongation and upstream ribosomes bump into it (Wolin and Walter, 1988). One possibility

is that mRNAs undergoing active translation are cleaved by RNase L, resulting in the

ribosomes stalling whenever they reached a 3' end of a 5' cleavage fragment. Prior work

using ribosome profiling demonstrated this kind of 3'-end stalling and disome formation

when the homologous nuclease, Ire1, cleaved mRNAs during the unfolded protein

response in fission yeast (Guydosh *et al.*, 2017). Consistent with the idea of an increase

in ribosome collisions during viral infection, it was shown that vaccinia virus infection leads

to an increase in ubiquitination of the ribosomal protein us10 by the ubiquitin ligase

ZNF598, which is known to be directly activated by ribosome collisions (Sundaramoorthy

et al., 2021). Alternatively, ribosome collisions could form for other reasons, including the

widespread translation of alternative (non-canonical) ORFs that we previously reported

(Karasik *et al.*, 2021), since such ORFs may be less evolutionarily tuned to minimize slow

translation (Müller et al., 2023). We also note that ZAKα has been reported to be

somewhat activated by individual stalled ribosomes, implying ribosome collisions are not

strictly required for the effects observed here (Snieckute et al., 2022). It is also possible

that ZAKα could be activated in other ways that involve the cleaved RNA fragments

through a yet unknown mechanism. The effect of RNase L-induced activation of the RSR

was previously shown to be JNK-mediated programmed cell death in response to viral

infections (Li *et al.*, 2004). Interestingly, both ZAKα and RNase L activation were shown

to trigger the inflammasome independently (Chakrabarti *et al.*, 2015; Robinson *et al.*,

2022). Considering our data, it is plausible that RNase L also triggers inflammasome

activation through ZAKα, and contributes to cell death.

Previous RNA-seq studies hinted that RNase L activation may act together with

the other dsRNA pathways to amplify the innate immune response (Burke *et al.*, 2019).

Our findings here confirm this synergy by showing that activation of JNK and p38 occur

by both RNase L dependent and independent routes. Since impairment of RNase L

activation can have serious health consequences linked to a reduced ability to fight viral

infections (Banday et al., 2022; Wickenhagen et al., 2021), it is possible that the loss of

this two-pronged activation weakens other branches of the immune response that are

independently triggered by dsRNA. In addition, our observation that ERK is not activated

and p38 is activated by 2-5A has not been consistently observed in previous studies

(Iordanov *et al.*, 2000; Malathi *et al.*, 2005), further suggesting that the activation of

kinases by RNase activity is regulated by other factors, possibly in a cell-line dependent

way.

Beyond control of new transcription, we also examined roles for RNase L in

changing which transcripts are translated. We noted several prominent cases (Figure 5)

where mRNA TEs were changed when RNase L was activated by 2-5A. The mechanism

underlying these translational changes remains unclear, however, several possible

explanations exist. Certain transcripts, such as *IFNB1*, are retained in the nucleus via

direct interactions with nucleopores when RNase L is activated (Burke *et al.*, 2021) or a

viral nuclease is activated (Fisher et al., 2022). This mechanism would explain the

apparent decrease in translational efficiency of this gene (decreased TE in poly I:C

treated *RNASEL KO* as compared to WT, Figure 5) and possibly other transcripts are

selectively retained in the nucleus. In addition, TE changes could arise because the

availability of ribosomal subunits and the corresponding translation machinery, such as

initiation factors, likely increase because the overall level of mRNA is greatly reduced

upon RNase L activation (Rath *et al.*, 2019; Thoms et al., 2020). Such a dramatic change

in this "ribosome homeostasis" (as defined, (Lodish, 1974; Mills and Green, 2017)) or

balance between mRNA and the translational machinery have been shown to result in

changes in which transcripts are favored for translation (Gaikwad et al., 2021; Mills et al.,

2016). Such a result is thought to occur because mRNAs that are generally less efficient

at recruiting ribosomes can be more heavily translated due to an abundance of ribosomal

subunits promoting translation via mass action.

We also focused on changes in translation efficiency related to the ISR since it is

a well-established regulatory pathway known to rely on control of translation via

phosphorylation of eIF2α by the dsRNA sensor PKR (Dever *et al.*, 2023; Young and Wek,

2016). One consequence of eIF2α phosphorylation is the translational activation of a set

of genes, such as *GADD34*, due to a shift in translation initiation from uORFs to the main

ORF (Young and Wek, 2016). Intriguingly, we found that activation of RNase L overrides

this effect of eIF2α phosphorylation and limits the translational activation of *GADD34*.

This translational suppression would serve to further enhance the inhibition of this gene

due to its known susceptibility to RNase L mediated degradation (Figure 6) (Burke *et al.*,

2019). One possible explanation for the inhibitory effect of RNase L on the translation of

this gene is that the availability of eIF2α may increase when RNase L is active due to the

general reduction in demand for translation, as noted above. As a result, some

phosphorylation of eIF2α population may be insufficient to influence translation initiation,

thus preventing the shift from uORF to main ORF translation on these transcripts.

Alternatively, since RNase L cleaves rRNAs, it is plausible that this damage could alter

the choice of start codon (uORF vs main ORF) by scanning 40S subunits. Another

possibility is that we previously showed (Karasik *et al.*, 2021) that RNase L activation

leads to an increase in the proportion of ribosome footprints in 5'UTRs globally. This effect

may arise from direct initiation on mRNA cleavage fragments generated when RNase L

cleaves in the 5'UTR. This change in initiation could therefore lead to the reduced

translational activation of *GADD34*.

Additionally, we found that RNase L-dependent changes in translation efficiency

of *IFIT1*, *IFIT2*, and *IFIT3* (as well as other interferon-related transcripts) is correlated with

an apparent shift in uORF to main ORF translation. One potential explanation for this
effect is that these *IFIT* genes, much like *GADD34*, could be activated by the ISR. In this
scenario, RNase L would similarly serve to limit their activation. While the expression of
these genes is likely already downregulated at the mRNA level by RNase L mediated
degradation, this mechanism offers additional control and suppression. Such dampening
could be important for limiting autoimmune conditions. For example, impairments in the
OAS-RNase L pathway is one of the contributing factors for MIS-C (Multisystem
Inflammatory Syndrome in Children) (Lee et al., 2023). In the case of MIS-C, the disease
appears to manifest due to the lack of RNase L's control of inflammation. Thus, active
RNase L may serve a dual role in innate immunity, activating key pathways while
counterbalancing others.

Finally, we found that the generic endonuclease RNase A evokes a transcriptional
response, including that mediated by ZAKα, that is very similar to that of RNase L (Figure
3). In addition, RNase A, or the ISR-mediated endonuclease IRE1, also triggers the
translation of altORFs, much like RNase L (Figure 4). This evidence suggests other
endonucleases, such as the viral endonuclease Nsp1 or the apoptotic host endonuclease
EndoU (Deng and Baker, 2018; Fisher *et al.*, 2022; Poe et al., 2014), could potentially
cause similar outcomes to RNase L. It is intriguing that virally-encoded endonucleases
may bring about outcomes that are similar to host endonucleases and suggests the
effects of generic endonuclease activation can be tuned to benefit either the virus or the
host. For example, in the case of Nsp1, nuclear mRNA retention and cytoplasmic mRNA
degradation were shown to be important for limiting the interferon response and
promoting viral replication (Burke *et al.*, 2021; Fisher *et al.*, 2022). On the other hand,
more limited dampening of these pathways, particularly for *IFNB1* (Burke *et al.*, 2019)
and *IFIT* genes shown here (Figure 7) could be important for preventing autoimmune
conditions or cytokine storms, such as noted above. Further studies are needed to
elucidate these possibilities.

Overall, our data offer a model where activation of RNase L induces ribosome
stalling and likely collisions, leading eventually to cell death via key transcriptional
pathways (Figure 1) related to the kinase ZAKα. This suggests that ZAKα could be a
potential target for therapeutics aimed at modulating RNase L's effects. In addition,

RNase L may have other roles in modulating the ISR via translation of ISR target genes.
The effects of RNase L on transcription and translation may be important for reinforcing
or dampening effects evoked by other dsRNA sensors, pointing to linkages between
pathways in the innate immune response. Further study is needed to dissect these
connections that bring together the entire spectrum of mRNA activities, including
synthesis, translation, and degradation.

Materials and Methods

Cell culture. WT and RNase L KO A549 lung carcinoma cells were cultured in RPMI
(Gibco Cat # 60870127) complemented with 10% Fetal Bovine Serum (Gibco FBS). A549
cells were tested and negative for Mycoplasma contamination throughout the study.
Mycoplasma testing was performed using eMyco Valid Mycoplasma PCR detection kit
(BioLink) or ATCC Universal Mycoplasma Detection Kit. Cells were incubated at 37°C in
the presence of 5% CO_2. Cell lines used in this study were a kind gift of Dr. Bernie Moss
(NIH) and generated as described in (Liu and Moss, 2016).

rRNA cleavage assay. Total RNA was extracted from ~3×10^5 cells using Direct-zol mini
kit (Zymo) or Trizol reagent (Invitrogen) according to the manufacturer's protocol. The
amount of total RNA was computed by absorbance at 260 nm measured by NanoDrop
(Thermo Fischer Scientific) and then diluted to 50-500 ng/µl. Then RNA samples were
run on a Bioanalyzer 2100 (Agilent) using the RNA 6000 Nano kit (Agilent) or TapeStation
with the Agilent RNA ScreenTape assay. Minute run-to-run shifts in band sizes are due
to inherent limitations of the instrument.

Sample preparation for RNA-seq and ribosome profiling. Cells used in experiments
were grown until they reached near confluency (70-90% confluent) in a T75 flask. Then
cells were transfected with the pre-incubated mixture of 1 µM 2-5A or 0.25 µg/ml poly I:C
(Invitrogen, low molecular weight) and Lipofectamine 3000 (7.5 µl/6-well) in Opti-Mem
serum free media as recommended by the manufacturer for 4.5 hours. Control cells

(untreated) were treated with lipofectamine only. 2-5A was synthesized as described
before (Karasik *et al.*, 2021). For electroporation of RNase A or BSA, confluent T75s (~6-
8x10^6 cells) were trypsinized and resuspended in MaxCyte electroporation buffer and
electroporated in 3XOC25 cuvettes (MaxCyte) in the presence of 160 ng/ml RNase A or
BSA with standard settings for A549 in the MaxCyte ATX electroporator. Cells were plated
on T75 flasks and collected after 4.5 hours incubation. Then cell lysates were prepared
and collected as for ribosome profiling (Karasik *et al.*, 2021). The same samples were
split for RNA-seq and ribosome profiling.

Ribosome profiling. Ribosome profiling of RNase A and BSA electroporated cells was
carried out as previously reported (Karasik *et al.*, 2021), except that rRNA depletion was
carried out by siTools riboPool rRNA depletion kit designed for human ribosome profiling.
The ribosome profiling data was processed and signatures of altORF translation were
assessed as before (Karasik *et al.*, 2021). Sequencing was performed on HiSeq 3000
(single end 50) at the NHLBI or NIDDK DNA Sequencing and Genomics Core. These
ribosome profiling data and previously published data were used for analysis of
translation efficiency (see below).

Total RNA-seq. Total RNA was extracted from 100 µl of lysates prepared for ribosome
profiling using Direct-zol mini kit (Zymo) or Trizol reagent (Invitrogen) according to the
manufacturer's protocol. The amount of total RNA was computed by absorbance at 260
nm measured by NanoDrop (Thermo Fischer Scientific) and then diluted to 50-500 ng/µl.
Then RNA samples were run on Bioanalyzer 2100 (Agilent) using the RNA 6000 Nano kit
(Agilent). Samples for sequencing were subject to rRNA depletion and library preparation
by Illumina Truseq kit. Sequencing was performed on HiSeq 3000 or Novaseq 6000
(paired end 50 or 75 cycle) at the NHLBI DNA Sequencing and Genomics Core.

Computational analysis.
Processing and alignment of RNA-seq reads.
After removing residual adapters from sequencing reads with the bbduk.sh tool from
bbtools (v37.62), sequencing data were mapped to a database of highly abundant rRNA

and tRNA sequences with bowtie2 (v 2.4.5) (Langmead et al., 2009). Unmapped reads
were then mapped to the human reference genome (GRCh38) with STAR (v2.7.10a)
using default parameters except for alignSJDBoverhangMin=1,
outFilterMismatchNmax=2, alignEndsType=EndToEnd, and leaving multimaper reads
mapping to less than 11 loci (Dobin et al., 2013). Duplicated reads were removed with
picard MarkDuplicates (v2.26.10) (Picard toolkit) and quantification of fragments per gene
was carried out with the featureCounts tool from Subread software (v2.0.1) (Liao et al.,
2014) using Ensembl gene annotations (Ensembl version hg38) by counting read-pairs
falling within coding regions (CDS) as fragments and adding fractional counts for
multimapper reads. BigWig files were generated with bamCoverage tool from DeepTools
(v3.5.1) (Ramírez et al., 2014).

Analysis of RNA-seq data.
Processed bulk RNaseq data was analyzed in R using the following workflow:
https://github.com/TriLab-bioinf/TK_65_RNAseL_manuscript.   Differential   gene
expression analysis was performed with the R package DESeq2 (v1.34.0) (Love *et al.*,
2014). GO enrichment analysis were carried out with the gseGO function from the R
package ClusterProfiler (v4.2.2) (Yu et al., 2012).
To create violin plots in Figure 2 and Figure 3 we utilized gene collections from
Harmonizome 3.0 (Rouillard et al., 2016) that were based on Gene References Into
Function (GeneRIFs, JNK, p38 datasets). Within these datasets we used a subgroup of
genes that were upregulated with 2-5A treatment (JNK/p38 plots, *N*=54, Table S3). For
interferon-related genes we used the hallmark datataset from the Molecular Signature
Database (Liberzon et al., 2015) ("interfron-gamma-response") for a subgroup of genes
that were upregulated in poly I:C treated WT cells (interferon-related plots, *N*=45, Table
S3). We performed paired t-test to determine significant differences between the groups
within each plot (see legends for more details).

Processing of ribosome profiling footprint data.
Ribosome profiling data was processed as described before (Karasik *et al.*, 2021). After
processing, reads were 5' aligned with STAR (v2.8.2a) to the human genome (hg38,

UCSC). Multimapping was allowed up to 200 times (--outFilterMultimapNmax option) and
the maximum mismatch/read was set to 1 (--outFilterMismatchNmax option). For both the
RNA-seq and ribosome profiling data CDS read counts were generated by featureCounts
tool from Subread software (v2.0.1) including multimappers as fractional counts and
reads on exon-exon junctions (-M --fraction and -J options, respectively).

Analysis of translational efficiencies.
We used our DEseq2 pipeline as described above for RNA-seq to obtain differential
expression values (log2 fold changes) for the ribosome profiling experiments. These
changes (RNA-seq and ribosome profiling log2 fold change values) were plotted against
each other in Figure 5A to visualize where translation efficiency (TE) was changing. We
used a filter for log2 fold changes > 2 and p adjusted value < 0.01 to evaluate the most
significant TE changes. The output of the DEseq2 analysis for the RNA-seq and ribosome
profiling are provided as Supplemental Table 1 and 2. For further analysis of individual
genes, we computed TEs directly by dividing reads per million (rpm) values of ribosome
profiling and RNA-seq by previously established pipelines in our lab (see more details in
(Guydosh, 2021; Karasik *et al.*, 2021)). This allowed us to directly compare replicates
within a treatment group.

Analysis of translation for altORF analysis and individual genes.
For more detailed analysis of ribosome profiling data, we implemented a reduced-
transcriptome alignment and analysis pipeline as described elsewhere (Karasik *et al.*,
2021) (Karasik 2021, (Ribofootprinter paper biorxiv 2021). Using these tools, we
computed the global ratio of UTR to main ORF reads and metagene plots. We also used
these tools to output reads for individual gene tracks. When uORFs overlapped the main
ORF (i.e. for *ATF4*), the overlap region was assigned to the UTR.

Western blotting. Protein levels were assessed by SDS-PAGE electrophoresis and
immunoblotting during 2-5A and poly I:C treatment. For western blotting cells were grown
on a 6-well plate up to ~70% confluency then transfected with 2-5A or poly I:C as
described above. Cells were trypsinized after 4.5 hours (Gibco TrypleExpress) and

collected by centrifugation at 200 g for 5 minutes. The cell pellet then was resuspended in 90 µl of 10 mM Tris-HCL pH=8.0, 150 mM NaCl and 1 % triton-X and incubated on ice for 5 minutes. Samples were then centrifuged for 1500 rpm, 4 minutes at 4°C. The supernatant was then mixed with equal amount of 2X SDS-PAGE buffer (Quality Biological) and samples were sonicated with a Brenson sonicator for 10-15 seconds until the sample was no longer viscous. 5-10 µl of sample was run on a 4-20% gradient Mini-PROTEAN Tris-HCl gel (BioRad) or 7% PhosTag (Wako) gel prepared according to manufacturer's instruction and as described before (Wu *et al.*, 2020). Proteins were transferred to a 0.2 µm PVDF membrane using the Trans-Blot Turbo System (BioRad, 1.3 A, 7 minutes for 4-20% gradient Mini-PROTEAN Tris-HCl gel or 1.3 A for 14 minutes for PhosTag gel). Membranes were blocked in EveryBlot Blocking Buffer (BioRad) for 10 minutes at room temperature. This was followed by incubation with primary antibodies overnight in TBST (Tris-buffered saline plus 0.1% tween) and 1% milk. The following antibodies were used: anti-p38 MAPK (Cell signaling Technologies #9212, 1:2000 dilution), anti-P-p38 MAPK (Cell signaling Technologies, 1:2000 dilution, #9211), anti-IRF3 (Cell signaling Technologies #11904), anti-P-IRF3 Ser396 (Cell signaling Technologies #4947), anti-ZAK (Bethyl A301-993A), anti-P-JNK (Thr183/Tyr185) (Cell signaling Technologies #4668), anti-JNK (Cell signaling Technologies #9252), anti-p65 (Cell signaling Technologies #8242), anti-ERK1/2 (Cell signaling Technologies #4695), anti P-ERK1/2 (Cell signaling Technologies #4370), anti-IFIT1 (abcam #236256), anti-GADD34 (abcam #236256). After washing 3 times for 5 minutes in TBST, secondary antibodies were incubated with the PVDF membrane for 1 hour at room temperature (Goat anti-rabbit antibody, BioRad, 1:3000). Then, after additional washes (3 times, 5 minutes each) in TBST, the PVDF membrane were incubated with Clarity Western ECL substrate (BioRad) for 5 minutes and proteins were visualized by Amersham Imager 600. Western blots were repeated for 2-4 times using biological replicates and blots were quantified using Fiji (Schindelin et al., 2012). Replicate images will be provided on Mendeley Data.

Figure Legends:

Figure 1. *RNase L activation leads to transcriptional changes through ZAK, JNK and p38.*

A Schematics of RNA-seq experiments. A549 cells were transfected with 2-5A and RNA-

seq was performed as described in Materials and Methods. **B** Volcano-plot showing

upregulation of example proinflammatory cytokines and transcription factors during 2-5A

treatment. p_{adj} value cutoff=0.5, log_2fold change cutoff=1. **C** Gene Ontology analysis of

molecular functions shows that upregulated genes are enriched in genes related to

transcription activities. The first five categories with the highest Normalized Enrichment

Scores are shown where number of genes/category > 100 genes and $p_{adjusted}$ <0.05. **D**

Western blots show that transcription factors FOSB and JUN are increased due to 2-5A

activation. Different FOSB isoforms are marked with black arrows. **E** Western blots show

JNK and p38 are activated in A549 cells after 2-5A treatment. Different JNK isoforms are

marked with black arrows. ZAKα is hyperphosphorylated during RNase L activation.

Upper bar shows phosphorylated ZAKα in a Phos-Tag gel (shifted from its expected

molecular weight). Lower line points to non-phosphorylated ZAKα. **F** Working model of

JNK/p38 pathway activation by RNase L. RNase L cleaves mRNAs that are being

translated. At the end of the resulting fragments, ribosomes can collide and activate ZAKα

and subsequently JNK and p38.

Figure 2. *OAS-RNase L and other dsRNA pathways act together to induce the*

proinflammatory response. **A** Schematics of RNA-seq experiments. A549 cells were

transfected with poly I:C (leading to activation of MDA5, RIG-I, PKR, and OAS, besides

RNase L) and RNA-seq was performed as described in Materials and Methods. **B** Venn-

diagram showing the overlap of DE genes in cells treated with poly I:C in WT (magenta),

2-5A in WT (yellow) and poly I:C in *RNASEL* KO cells (green, abbreviated to KO). **C**

Correlation between differential expression changes in 2-5A (activation of RNase L) vs

poly I:C (activation of all dsRNA pathways including RNase L) treated cells. Blue and red

dots represent JNK/p38 MAPK activation-related or IFN stimulated genes (as defined in

Materials and Methods), differentially expressed in 2-5A or poly I:C treated cells,

respectively. Black dot represents *GADD34*. **D** Violin plot representation of genes related

to JNK and p38 MAPK activation and interferon stimulation. Statistical testing (paired t-

tests) showed significant differences between data sets as indicated (p value < 0.001,

marked by ***). **E** Synergistic effects of the OAS-RNase L pathway on activation of JNK

(upper panels), p38 (middle panels) and IRF3 (lower panels) assayed by Western

blotting. ZAKα is hyperphosphorylated during dsRNA response. Upper bar shows

phosphorylated ZAKα in a Phos-Tag gel (shifted from its expected molecular weight).

Lower line points to non-phosphorylated ZAKα. **F** Quantification of western blot for poly

I:C treated cells WT and *RNASEL* KO cells to assess level of synergy in activation

between RNase L and other dsRNA pathways. Activation levels are normalized to that of

poly I:C treated WT cells. Data reveal strong RNase L dependence for JNK and little

dependence for p38 and IRF3. **G** Model for RNase L acting synergistically with the other

dsRNA pathways to induce a strong antiviral effect, particularly for JNK.

Figure 3. *Global RNA cleavage in the cell leads to transcriptional changes.* **A** Schematics

of RNA-seq experiments. *RNASEL* KO A549 cells were electroporated with RNase A or

BSA (control) and RNA-seq experiments were performed as described in Materials and

Methods. **B** Venn-diagram showing the number of upregulated genes (defined as p

adjusted value < 0.05 and log2fold changes >1) in RNase A electroporated samples

(purple) and 2-5A treated cells (yellow). **C** Comparison of RNA log2 fold changes between

RNase L activated vs RNase A electroporated samples. Blue dots are representing

transcripts related to JNK and p38 activation ($N = 53$, as defined in Materials and

Methods). **D** Violin plot representation of those genes related to JNK and p38 activation

in RNase A electroporated cells. *RNASEL* KO is abbreviated to KO. Statistical analysis

(paired t-test) showed that the log2 fold changes between RNase A electroporated and

2-5A activated samples were not significantly different (p value > 0.05, marked by n.s.).

E JNK, p38 and ZAKα are activated due to RNase A treatment, as shown by western

blots. Different JNK isoforms are marked with black arrows. Upper bar shows

phosphorylated ZAKα in a Phos-Tag gel (shifted from its expected molecular weight).

Lower line points to non-phosphorylated ZAKα.

Figure 4. *RNA fragmentation leads to alternative ORF (altORF) translation.* **A** Schematics

of ribosome profiling experiments. *RNASEL* KO A549 cells were electroporated with

RNase A or BSA (Control) and ribosome profiling experiments were performed as

described in Materials and Methods. **B** Ribosome profiling reads in the 3'UTR of the
ACTG1 gene model shows the signature of altORF translation in RNase A electroporated
and RNase L activated A549 cells, but not in the controls (-2-5A and BSA). **C** Increased
3'UTR:ORF ratios indicate higher translation of 3'UTRs when active RNases are present
in the cell, but not in controls (-2-5A and BSA). **D** Normalized average ribosome footprint
occupancy (metagene plot) around the stop codon of main ORFs (left panel) reveals
increased ribosome footprint levels in the 3' UTRs (right panel, zoom in). **E** Increased
3'UTR:ORF ratios indicate altORF translation during thapsigargin treatment and
activation of the UPR, but does not occur in untreated cells.

In all panels data shown for RNase L activation (+2-5A) was obtained from (Karasik *et*
al., 2021).

Figure 5. *RNase L activation influences translational control and effects of poly I:C*
treatment. **A**. Comparison of changes in RNA-seq vs ribosome profiling under conditions
of 2-5A treated WT cells (left), poly I:C treated WT cells (center), and poly I:C treated
RNASEL KO A549 cells (right). To emphasize the highest and most significant changes,
genes were filtered for the following criteria: log2 fold changes >2 (either RNA-seq or
ribosome profiling), $p_{adjusted}$ <0.01, normalized mean counts across the samples > 50 (see
Materials and Methods for details). Orange circles represent genes encoded by the
mitochondria. **B.** TEs from individual replicates of representative genes with
downregulated (MUC5AC and MUC5B) or upregulated (CXCL3 or CXCL8) translation
were computed. TEs calculated as described in Materials and Methods. *RNASEL* KO is
abbreviated to KO in all panels.

Figure 6. *RNase L modulates elements of the Integrated Stress Response.* **A** Bar plots
representing TEs of *GADD34* mRNA during poly I:C treatment in WT and *RNASEL KO*
cells. These data show that TEs changes are greater in *RNASEL* KO cells during the ISR
as compared to respective controls. **B** Ribosome profiling tracks of gene models for
GADD34 (*PPP1R15A*) in WT and *RNASEL* KO cells during poly I:C treatment suggest a
change in 5'UTR vs main ORF translation ratio in *RNASEL* KO cells during the ISR as
compared to WT cells and their respective controls. **C** 5'UTR:main ORF ratios of ribosome

profiling data in WT and *RNASEL* KO cells during poly I:C treatment were computed for

individual replicates and show a reduction in cells where RNase L is absent. **D.** Western

blot for GADD34 shows increased levels of protein in poly I:C treated *RNASEL* KO cells,

reflecting both the lack of mRNA degradation and enhanced TE in cells where RNase L

is absent.

Figure 7. *IFIT mRNAs undergo changes in translation that depend on RNase L.* **A** uORFs

were identified on *IFIT1* mRNAs based on the increase in ribosomes density at alternative

start codons (CUG, green markings) in poly I:C treated cells. Red marking shows the start

codon of the main ORF. **B** TE of *IFIT1-3* mRNAs are decreased in RNase L deficient cells

(comparing WT and *RNASEL* KO poly I:C treated cells) consistently across replicates. **C**

IFIT1-3 mRNAs have less 5'UTR and increased main ORF translation in poly I:C treated

RNASEL KO (abbreviated to KO) cells as compared to WT cells. **D** Schematics of how

eIF2α phosphorylation and activation of RNase L affects gene expression in the cell.

While both mechanisms lead to reduced gene expression, eIF2α phosphorylation

increases translation on ISR-related mRNAs but RNase L reduces their translation.

Results suggest that and *IFIT* mRNA translation might be also controlled by uORF and

eIF2α phosphorylation (marked by asterisk since further confirmation is needed).

S1 Related to Figure 1. A Cleavage patterns of total RNA extracts show activation of

RNase L in 2-5A or poly I:C treated WT, but not *RNASEL* KO, A549 cells. rRNA cleavage

assay performed on a BioAnalyzer. Black arrows indicate the 28 and 18S rRNAs and red

arrows show RNase L degradation products. **B** Heatmap representation of the most

highly expressed genes during RNase L activation. Trends are consistent across

replicates. **C** Western blots show that transcription factors FOSB and JUN do not increase

in *RNASEL* KO cell due to 2-5A activation. **D** Western blots show JNK, p38 and ZAKα

are not activated in *RNASEL* KO cells after 2-5A treatment. **E** Western blots show ERK

is not activated in WT cells after 2-5A treatment.

S2 Related to Figure 2. Western blots show no detectable effect of RNase L activation on NF-κB activation by poly I:C and no detectable activation by 2-5A. ERK activation was not observed during 2-5A or poly I:C treatment.

S3 Related to Figure 3. A Cleavage of RNAs in the cell occurs when RNase A is electroporated into cells, but not when BSA is electroporated, as observed by rRNA cleavage assays performed on BioAnalyzer. Arrows indicate the 28S and 18S rRNAs. **B** Volcano-plot showing upregulation of example proinflammatory cytokines and transcription factors during RNase A electroporation. p_{adj} value cutoff=0.5, log_2fold change cutoff=1.

S4 Related to Figure 4. A Increased 5'UTR:ORF ratios indicating higher uORF translation when active RNases are present in the cell, but not in controls (-2-5A and BSA electroporated). **B** Normalized average ribosome footprint occupancy (metagene plot) around the start codon of main ORFs reveals increased ribosome footprint levels in the 5' UTRs when an active RNase is present vs the respective control.

In all panels data shown for RNase L activation (+2-5A) was obtained from (Karasik *et al.*, 2021).

S5 Related to Figure 6. A Bar plots representing TEs of *ATF4* and *CHOP* mRNA during poly I:C treatment in WT and *RNASEL KO* cells across replicates. **B** Ribosome profiling tracks for gene model of *ATF4* and *CHOP* in WT and *RNASEL* KO cells during 2-5A or poly I:C treatment. Data show the poly I:C dependent shift toward main ORF vs 5'UTR translation is greater when RNase L is absent. Asterisks show RNase L dependent ribosome profiling peaks in 2-5A and poly I:C treated cells that likely correspond to altORF translation initiation events. **C** 5'UTR:main ORF ratios of ribosome profiling peaks in WT and *RNASEL* KO cells during poly I:C treatment for *ATF4* and *CHOP*.

S6 Related to Figure 7. A uORFs were identified on *IFIT2* and *3* mRNAs based on the increase in ribosomes densities on alternative start codons (CUG or GUG, green markings) in poly I:C treated WT or *RNASEL* KO cells. Red marking shows the start codon

for the main ORF. **B** Western blot for *IFIT1* shows increased protein levels in poly I:C

treated RNase L KO cells, suggesting control of TE or RNA degradation reduce levels in

WT cells.

Table S1. Results tables from DEseq2 analysis for RNA-seq (WT and RNASEL KO, +/-

2-5A, +/- poly I:C, BSA vs RNase A electroporated)

Table S2. Results tables from DEseq2 analysis for ribosome profiling (WT and RNASEL

KO, +/-2-5A, +/- poly I:C, BSA vs RNase A electroporated)

Table S3. List of genes that are upregulated by JNK/p38 and interferon response in WT

2-5A and poly I:C treated cells, respectively. This list was used to create violin plots in

Figure 2D and 3D.

References

Andersen, J.B., Mazan-Mamczarz, K., Zhan, M., Gorospe, M., and Hassel, B.A. (2009). Ribosomal protein mRNAs are primary targets of regulation in RNase-L-induced senescence. RNA biology *6*, 305-315. 10.4161/rna.6.3.8526.

Banday, A.R., Stanifer, M.L., Florez-Vargas, O., Onabajo, O.O., Papenberg, B.W., Zahoor, M.A., Mirabello, L., Ring, T.J., Lee, C.H., Albert, P.S., et al. (2022). Genetic regulation of OAS1 nonsense-mediated decay underlies association with COVID-19 hospitalization in patients of European and African ancestries. Nature genetics *54*, 1103-1116. 10.1038/s41588-022-01113-z.

Banerjee, S., Li, G., Li, Y., Gaughan, C., Baskar, D., Parker, Y., Lindner, D.J., Weiss, S.R., and Silverman, R.H. (2015). RNase L is a negative regulator of cell migration. Oncotarget *6*, 44360-44372. 10.18632/oncotarget.6246.

Bisbal, C., Martinand, C., Silhol, M., Lebleu, B., and Salehzada, T. (1995). Cloning and characterization of a RNAse L inhibitor. A new component of the interferon-regulated 2-5A pathway. The Journal of biological chemistry *270*, 13308-13317.

Burke, J.M., Gilchrist, A.R., Sawyer, S.L., and Parker, R. (2021). RNase L limits host and viral protein synthesis via inhibition of mRNA export. Sci Adv *7*. 10.1126/sciadv.abh2479.

Burke, J.M., Moon, S.L., Matheny, T., and Parker, R. (2019). RNase L Reprograms Translation by Widespread mRNA Turnover Escaped by Antiviral mRNAs. Molecular cell *75*, 1203-1217.e1205. 10.1016/j.molcel.2019.07.029.

Carpten, J., Nupponen, N., Isaacs, S., Sood, R., Robbins, C., Xu, J., Faruque, M., Moses, T., Ewing, C., Gillanders, E., et al. (2002). Germline mutations in the ribonuclease L gene in families showing linkage with HPC1. Nature genetics *30*, 181-184. 10.1038/ng823.

Casey, G., Neville, P.J., Plummer, S.J., Xiang, Y., Krumroy, L.M., Klein, E.A., Catalona, W.J., Nupponen, N., Carpten, J.D., Trent, J.M., et al. (2002). RNASEL Arg462Gln variant is implicated in up to 13% of prostate cancer cases. Nature genetics *32*, 581-583. 10.1038/ng1021.

Castelli, J.C., Hassel, B.A., Wood, K.A., Li, X.L., Amemiya, K., Dalakas, M.C., Torrence, P.F., and Youle, R.J. (1997). A study of the interferon antiviral mechanism: apoptosis activation by the 2-5A system. The Journal of experimental medicine *186*, 967-972. 10.1084/jem.186.6.967.

Chakrabarti, A., Banerjee, S., Franchi, L., Loo, Y.M., Gale, M., Jr., Núñez, G., and Silverman, R.H. (2015). RNase L activates the NLRP3 inflammasome during viral infections. Cell Host Microbe *17*, 466-477. 10.1016/j.chom.2015.02.010.

Costa-Mattioli, M., and Walter, P. (2020). The integrated stress response: From mechanism to disease. Science (New York, N.Y.) *368*. 10.1126/science.aat5314.

Dai, A., Cao, S., Dhungel, P., Luan, Y., Liu, Y., Xie, Z., and Yang, Z. (2017). Ribosome Profiling Reveals Translational Upregulation of Cellular Oxidative Phosphorylation mRNAs during Vaccinia Virus-Induced Host Shutoff. Journal of virology *91*. 10.1128/jvi.01858-16.

Dalet, A., Argüello, R.J., Combes, A., Spinelli, L., Jaeger, S., Fallet, M., Vu Manh, T.P.,
Mendes, A., Perego, J., Reverendo, M., et al. (2017). Protein synthesis inhibition and
GADD34 control IFN-β heterogeneous expression in response to dsRNA. Embo j *36*, 761-
782. 10.15252/embj.201695000.

Deng, X., and Baker, S.C. (2018). An "Old" protein with a new story: Coronavirus
endoribonuclease is important for evading host antiviral defenses. Virology *517*, 157-163.
10.1016/j.virol.2017.12.024.

Dever, T.E., Ivanov, I.P., and Hinnebusch, A.G. (2023). Translational regulation by
uORFs and start codon selection stringency. Genes Dev *37*, 474-489.
10.1101/gad.350752.123.

Dhanasekaran, D.N., and Reddy, E.P. (2017). JNK-signaling: A multiplexing hub in
programmed cell death. Genes Cancer *8*, 682-694. 10.18632/genesandcancer.155.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*, 15-21. 10.1093/bioinformatics/bts635.

Donovan, J., Rath, S., Kolet-Mandrikov, D., and Korennykh, A. (2017). Rapid RNase L-
driven arrest of protein synthesis in the dsRNA response without degradation of
translation machinery. RNA (New York, N.Y.) *23*, 1660-1671. 10.1261/rna.062000.117.

Fensterl, V., and Sen, G.C. (2015). Interferon-induced Ifit proteins: their role in viral
pathogenesis. J Virol *89*, 2462-2468. 10.1128/jvi.02744-14.

Fisher, T., Gluck, A., Narayanan, K., Kuroda, M., Nachshon, A., Hsu, J.C., Halfmann,
P.J., Yahalom-Ronen, Y., Tamir, H., Finkel, Y., et al. (2022). Parsing the role of NSP1 in
SARS-CoV-2 infection. Cell Rep *39*, 110954. 10.1016/j.celrep.2022.110954.

Floyd-Smith, G., Slattery, E., and Lengyel, P. (1981). Interferon action: RNA cleavage
pattern of a (2'-5')oligoadenylate--dependent endonuclease. Science (New York, N.Y.)
212, 1030-1032. 10.1126/science.6165080.

Fusakio, M.E., Willy, J.A., Wang, Y., Mirek, E.T., Al Baghdadi, R.J., Adams, C.M.,
Anthony, T.G., and Wek, R.C. (2016). Transcription factor ATF4 directs basal and stress-
induced gene expression in the unfolded protein response and cholesterol metabolism in
the liver. Mol Biol Cell *27*, 1536-1551. 10.1091/mbc.E16-01-0039.

Gaikwad, S., Ghobakhlou, F., Young, D.J., Visweswaraiah, J., Zhang, H., and
Hinnebusch, A.G. (2021). Reprogramming of translation in yeast cells impaired for
ribosome recycling favors short, efficiently translated mRNAs. Elife *10*.
10.7554/eLife.64283.

Guydosh, N.R. (2021). ribofootPrinter: A precision python toolbox for analysis of ribosome
profiling data. bioRxiv.

Guydosh, N.R., Kimmig, P., Walter, P., and Green, R. (2017). Regulated Ire1-dependent
mRNA decay requires no-go mRNA degradation to maintain endoplasmic reticulum
homeostasis in S. pombe. Elife *6*. 10.7554/eLife.29216.

Hartenian, E., and Glaunsinger, B.A. (2019). Feedback to the central dogma: cytoplasmic
mRNA decay and transcription are interdependent processes. Crit Rev Biochem Mol Biol
54, 385-398. 10.1080/10409238.2019.1679083.

Hartmann, G. (2017). Nucleic Acid Immunity. Advances in immunology *133*, 121-169.
10.1016/bs.ai.2016.11.001.

Hinnebusch, A.G., Ivanov, I.P., and Sonenberg, N. (2016). Translational control by 5'-
untranslated regions of eukaryotic mRNAs. Science (New York, N.Y.) *352*, 1413-1416.
10.1126/science.aad9868.

Hollien, J., Lin, J.H., Li, H., Stevens, N., Walter, P., and Weissman, J.S. (2009). Regulated
Ire1-dependent decay of messenger RNAs in mammalian cells. J Cell Biol *186*, 323-331.
10.1083/jcb.200903014.

Iordanov, M.S., Paranjape, J.M., Zhou, A., Wong, J., Williams, B.R., Meurs, E.F.,
Silverman, R.H., and Magun, B.E. (2000). Activation of p38 mitogen-activated protein
kinase and c-Jun NH(2)-terminal kinase by double-stranded RNA and
encephalomyocarditis virus: involvement of RNase L, protein kinase R, and alternative
pathways. Mol Cell Biol *20*, 617-627. 10.1128/mcb.20.2.617-627.2000.

Iordanov, M.S., Pribnow, D., Magun, J.L., Dinh, T.H., Pearson, J.A., Chen, S.L., and
Magun, B.E. (1997). Ribotoxic stress response: activation of the stress-activated protein
kinase JNK1 by inhibitors of the peptidyl transferase reaction and by sequence-specific
RNA damage to the alpha-sarcin/ricin loop in the 28S rRNA. Mol Cell Biol *17*, 3373-3381.
10.1128/mcb.17.6.3373.

Karasik, A., Jones, G.D., DePass, A.V., and Guydosh, N.R. (2021). Activation of the
antiviral factor RNase L triggers translation of non-coding mRNA sequences. Nucleic
Acids Res *49*, 6007-6026. 10.1093/nar/gkab036.

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-
efficient alignment of short DNA sequences to the human genome. Genome Biol *10*, R25.
10.1186/gb-2009-10-3-r25.

Le Roy, F., Salehzada, T., Bisbal, C., Dougherty, J.P., and Peltz, S.W. (2005). A newly
discovered function for RNase L in regulating translation termination. Nat Struct Mol Biol
12, 505-512. 10.1038/nsmb944.

Lee, D., Le Pen, J., Yatim, A., Dong, B., Aquino, Y., Ogishi, M., Pescarmona, R.,
Talouarn, E., Rinchai, D., Zhang, P., et al. (2023). Inborn errors of OAS–RNase
L in SARS-CoV-2–related multisystem inflammatory syndrome in children.
Science (New York, N.Y.) *0*, eabo3627. doi:10.1126/science.abo3627.

Lee, T.Y., Ezelle, H.J., Venkataraman, T., Lapidus, R.G., Scheibner, K.A., and Hassel,
B.A. (2013). Regulation of human RNase-L by the miR-29 family reveals a novel
oncogenic role in chronic myelogenous leukemia. J Interferon Cytokine Res *33*, 34-42.
10.1089/jir.2012.0062.

Li, G., Xiang, Y., Sabapathy, K., and Silverman, R.H. (2004). An apoptotic signaling
pathway in the interferon antiviral response mediated by RNase L and c-Jun NH2-terminal
kinase. The Journal of biological chemistry *279*, 1123-1131. 10.1074/jbc.M305893200.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics *30*, 923-930. 10.1093/bioinformatics/btt656.

Liberzon, A., Birger, C., Thorvaldsdóttir, H., Ghandi, M., Mesirov, J.P., and Tamayo, P. (2015). The Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell Syst *1*, 417-425. 10.1016/j.cels.2015.12.004.

Liu, R., and Moss, B. (2016). Opposing Roles of Double-Stranded RNA Effector Pathways and Viral Defense Proteins Revealed with CRISPR-Cas9 Knockout Cell Lines and Vaccinia Virus Mutants. J Virol *90*, 7864-7879. 10.1128/jvi.00869-16.

Lodish, H.F. (1974). Model for the regulation of mRNA translation applied to haemoglobin synthesis. Nature *251*, 385-388. 10.1038/251385a0.

Long, T.M., Chakrabarti, A., Ezelle, H.J., Brennan-Laun, S.E., Raufman, J.P., Polyakova, I., Silverman, R.H., and Hassel, B.A. (2013). RNase-L deficiency exacerbates experimental colitis and colitis-associated cancer. Inflammatory bowel diseases *19*, 1295-1305. 10.1097/MIB.0b013e318281f2fd.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550. 10.1186/s13059-014-0550-8.

Ma, Y., Brewer, J.W., Diehl, J.A., and Hendershot, L.M. (2002). Two distinct stress signaling pathways converge upon the CHOP promoter during the mammalian unfolded protein response. J Mol Biol *318*, 1351-1365. 10.1016/s0022-2836(02)00234-6.

Ma, Y., and Hendershot, L.M. (2003). Delineation of a negative feedback regulatory loop that controls protein translation during endoplasmic reticulum stress. The Journal of biological chemistry *278*, 34864-34873. 10.1074/jbc.M301107200.

Magg, T., Okano, T., Koenig, L.M., Boehmer, D.F.R., Schwartz, S.L., Inoue, K., Heimall, J., Licciardi, F., Ley-Zaporozhan, J., Ferdman, R.M., et al. (2021). Heterozygous OAS1 gain-of-function variants cause an autoinflammatory immunodeficiency. Sci Immunol *6*. 10.1126/sciimmunol.abf9564.

Malathi, K., Dong, B., Gale, M., Jr., and Silverman, R.H. (2007). Small self-RNA generated by RNase L amplifies antiviral innate immunity. Nature *448*, 816-819. 10.1038/nature06042.

Malathi, K., Paranjape, J.M., Bulanova, E., Shim, M., Guenther-Johnson, J.M., Faber, P.W., Eling, T.E., Williams, B.R., and Silverman, R.H. (2005). A transcriptional signaling pathway in the IFN system mediated by 2'-5'-oligoadenylate activation of RNase L. Proc Natl Acad Sci U S A *102*, 14533-14538. 10.1073/pnas.0507551102.

Manivannan, P., Siddiqui, M.A., and Malathi, K. (2020). RNase L Amplifies Interferon Signaling by Inducing Protein Kinase R-Mediated Antiviral Stress Granules. Journal of Virology *94*, e00205-00220. 10.1128/jvi.00205-20.

Mills, E.W., and Green, R. (2017). Ribosomopathies: There's strength in numbers. Science (New York, N.Y.) *358*. 10.1126/science.aan2755.

Mills, E.W., Wangen, J., Green, R., and Ingolia, N.T. (2016). Dynamic Regulation of a Ribosome Rescue Pathway in Erythroid Cells and Platelets. Cell Rep *17*, 1-10. 10.1016/j.celrep.2016.08.088.

Müller, M.B.D., Kasturi, P., Jayaraj, G.G., and Hartl, F.U. (2023). Mechanisms of readthrough mitigation reveal principles of GCN1-mediated translational quality control. Cell *186*, 3227-3244.e3220. 10.1016/j.cell.2023.05.035.

Nogimori, T., Nishiura, K., Kawashima, S., Nagai, T., Oishi, Y., Hosoda, N., Imataka, H., Kitamura, Y., Kitade, Y., and Hoshino, S.I. (2019). Dom34 mediates targeting of exogenous RNA in the antiviral OAS/RNase L pathway. Nucleic Acids Res *47*, 432-449. 10.1093/nar/gky1087.

Nogimori, T., Ogami, K., Oishi, Y., Goda, R., Hosoda, N., Kitamura, Y., Kitade, Y., and Hoshino, S.I. (2020). ABCE1 Acts as a Positive Regulator of Exogenous RNA Decay. Viruses *12*. 10.3390/v12020174.

Nogués, M.V., Vilanova, M., and Cuchillo, C.M. (1995). Bovine pancreatic ribonuclease A as a model of an enzyme with multiple substrate binding sites. Biochim Biophys Acta *1253*, 16-24. 10.1016/0167-4838(95)00138-k.

Novoa, I., Zeng, H., Harding, H.P., and Ron, D. (2001). Feedback inhibition of the unfolded protein response by GADD34-mediated dephosphorylation of eIF2alpha. J Cell Biol *153*, 1011-1022. 10.1083/jcb.153.5.1011.

Picard toolkit. Broad Institute, GitHub repository *https://broadinstitute.github.io/picard/*.

Poe, J.C., Kountikov, E.I., Lykken, J.M., Natarajan, A., Marchuk, D.A., and Tedder, T.F. (2014). EndoU is a novel regulator of AICD during peripheral B cell selection. The Journal of experimental medicine *211*, 57-69. 10.1084/jem.20130648.

Ramírez, F., Dündar, F., Diehl, S., Grüning, B.A., and Manke, T. (2014). deepTools: a flexible platform for exploring deep-sequencing data. Nucleic Acids Res *42*, W187-191. 10.1093/nar/gku365.

Rath, S., Donovan, J., Whitney, G., Chitrakar, A., Wang, W., and Korennykh, A. (2015). Human RNase L tunes gene expression by selectively destabilizing the microRNA-regulated transcriptome. Proc Natl Acad Sci U S A *112*, 15916-15921. 10.1073/pnas.1513034112.

Rath, S., Prangley, E., Donovan, J., Demarest, K., Wingreen, N.S., Meir, Y., and Korennykh, A. (2019). Concerted 2-5A-Mediated mRNA Decay and Transcription Reprogram Protein Synthesis in the dsRNA Response. Molecular cell *75*, 1218-1228.e1216. 10.1016/j.molcel.2019.07.027.

Robinson, K.S., Toh, G.A., Rozario, P., Chua, R., Bauernfried, S., Sun, Z., Firdaus, M.J., Bayat, S., Nadkarni, R., Poh, Z.S., et al. (2022). ZAKα-driven ribotoxic stress response activates the human NLRP1 inflammasome. Science (New York, N.Y.) *377*, 328-335. 10.1126/science.abl6324.

Rouillard, A.D., Gundersen, G.W., Fernandez, N.F., Wang, Z., Monteiro, C.D., McDermott, M.G., and Ma'ayan, A. (2016). The harmonizome: a collection of processed

datasets gathered to serve and mine knowledge about genes and proteins. Database
(Oxford) *2016*. 10.1093/database/baw100.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat Methods *9*, 676-682. 10.1038/nmeth.2019.

Siddiqui, M.A., and Malathi, K. (2012). RNase L induces autophagy via c-Jun N-terminal
kinase and double-stranded RNA-dependent protein kinase signaling pathways. The
Journal of biological chemistry *287*, 43651-43664. 10.1074/jbc.M112.399964.

Silverman, R.H. (2007). Viral encounters with 2',5'-oligoadenylate synthetase and RNase
L during the interferon antiviral response. J Virol *81*, 12720-12729. 10.1128/jvi.01471-07.

Snieckute, G., Genzor, A.V., Vind, A.C., Ryder, L., Stoneley, M., Chamois, S., Dreos, R.,
Nordgaard, C., Sass, F., Blasius, M., et al. (2022). Ribosome stalling is a signal for
metabolic regulation by the ribotoxic stress response. Cell Metab *34*, 2036-2046.e2038.
10.1016/j.cmet.2022.10.011.

Sundaramoorthy, E., Ryan, A.P., Fulzele, A., Leonard, M., Daugherty, M.D., and Bennett,
E.J. (2021). Ribosome quality control activity potentiates vaccinia virus protein synthesis
during infection. J Cell Sci *134*. 10.1242/jcs.257188.

Thoms, M., Buschauer, R., Ameismeier, M., Koepke, L., Denk, T., Hirschenberger, M.,
Kratzat, H., Hayn, M., Mackens-Kiani, T., Cheng, J., et al. (2020). Structural basis for
translational shutdown and immune evasion by the Nsp1 protein of SARS-CoV-2.
Science (New York, N.Y.) *369*, 1249-1255. 10.1126/science.abc8665.

Tnani, M., Aliau, S., and Bayard, B. (1998). Localization of a molecular form of interferon-
regulated RNase L in the cytoskeleton. J Interferon Cytokine Res *18*, 361-368.
10.1089/jir.1998.18.361.

Vind, A.C., Genzor, A.V., and Bekker-Jensen, S. (2020a). Ribosomal stress-surveillance:
three pathways is a magic number. Nucleic Acids Res *48*, 10648-10661.
10.1093/nar/gkaa757.

Vind, A.C., Snieckute, G., Blasius, M., Tiedje, C., Krogh, N., Bekker-Jensen, D.B.,
Andersen, K.L., Nordgaard, C., Tollenaere, M.A.X., Lund, A.H., et al. (2020b). ZAKα
Recognizes Stalled Ribosomes through Partially Redundant Sensor Domains. Molecular
cell *78*, 700-713.e707. 10.1016/j.molcel.2020.03.021.

Walter, P., and Ron, D. (2011). The unfolded protein response: from stress pathway to
homeostatic regulation. Science (New York, N.Y.) *334*, 1081-1086.
10.1126/science.1209038.

Wang, X., Mader, M.M., Toth, J.E., Yu, X., Jin, N., Campbell, R.M., Smallwood, J.K.,
Christe, M.E., Chatterjee, A., Goodson, T., Jr., et al. (2005). Complete inhibition of
anisomycin and UV radiation but not cytokine induced JNK and p38 activation by an aryl-
substituted dihydropyrrolopyrazole quinoline and mixed lineage kinase 7 small interfering
RNA. The Journal of biological chemistry *280*, 19298-19305. 10.1074/jbc.M413059200.

Wek, R.C., Jiang, H.Y., and Anthony, T.G. (2006). Coping with stress: eIF2 kinases and
translational control. Biochem Soc Trans *34*, 7-11. 10.1042/bst20060007.

Wickenhagen, A., Sugrue, E., Lytras, S., Kuchi, S., Noerenberg, M., Turnbull, M.L.,
Loney, C., Herder, V., Allan, J., Jarmson, I., et al. (2021). A prenylated dsRNA sensor
protects against severe COVID-19. Science (New York, N.Y.) *374*, eabj3624.
10.1126/science.abj3624.

Wolin, S.L., and Walter, P. (1988). Ribosome pausing and stacking during translation of
a eukaryotic mRNA. Embo j *7*, 3559-3569. 10.1002/j.1460-2075.1988.tb03233.x.

Woo, Y.M., Kwak, Y., Namkoong, S., Kristjánsdóttir, K., Lee, S.H., Lee, J.H., and Kwak,
H. (2018). TED-Seq Identifies the Dynamics of Poly(A) Length during ER Stress. Cell Rep
24, 3630-3641.e3637. 10.1016/j.celrep.2018.08.084.

Wu, C.C., Peterson, A., Zinshteyn, B., Regot, S., and Green, R. (2020). Ribosome
Collisions Trigger General Stress Responses to Regulate Cell Fate. Cell *182*, 404-
416.e414. 10.1016/j.cell.2020.06.006.

Young, S.K., and Wek, R.C. (2016). Upstream Open Reading Frames Differentially
Regulate Gene-specific Translation in the Integrated Stress Response. The Journal of
biological chemistry *291*, 16927-16935. 10.1074/jbc.R116.733899.

Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). clusterProfiler: an R package for
comparing biological themes among gene clusters. Omics *16*, 284-287.
10.1089/omi.2011.0118.

1257

Figure 1.

A A549 WT or *RNASEL* KO → +2-5A → Activation of RNase L only → Total RNA-seq

B Volcano plot: Log2 fold changes (x-axis) vs -Log p adjusted (y-axis). Labeled genes: CXCL2, DUSP8, CXCL8, FOSB, JUN, DUSP1, CCL20, ATF3, IL6, GDF 5, FOXA2, ATF5

C Normalized Enrichment Score (NES):
- Signaling receptor activator activity
- Receptor ligand activity
- Cytokine activity
- Transcription repressor activity
- Transcription activator activity

D JUN, FOSB, H3 — Control, 2-5A 1 µM, 5 µM

E JNK, P-JNK; p38, P-p38; P-ZAKα, ZAKα — Control, 2-5A 1 µM, 5 µM

F Active RNase L → RNA cleavage → Ribosome collision (ZAKα → p38, JNK) → Apoptosis

Figure 2.

Figure 3.

A — A549 RNASEL KO cells → electroporation of RNase A → Total RNA-seq

B — RNase L / RNase A — 1670 | 475 | 277

C — JNK/p38
+2-5A vs -2-5A log2 fold change (y-axis)
RNase A vs BSA log2 fold change (x-axis)

D — JNK/p38
Log2 fold change
KO | WT (+ 2-5A) | RNase A
n.s.

E —
BSA | RNase A
JNK
P-JNK

BSA | RNase A
p38
P-p38

BSA | RNase A
P-ZAKα
ZAKα

Figure 4.

Figure 5.

A

B

Figure 6.

A

C

B

GADD34

WT Control

WT + poly I:C

KO Control

KO + poly I:C

uORF1 uORF2
5'UTR   Main ORF   3'UTR

D

Figure 7.

S1.

A

WT   *RNASEL* KO   WT   *RNASEL* KO   WT

1 µM 2-5A   -   +   -   -   +   -   -   +   -   -   +   -   +
poly I:C   -   -   +   -   -   +   -   -   +   -   -   +   -

28S

18S

4000
nt

2000

1000

Replicate 1   Replicate 2   Replicate 3

B

-2-5A   +2-5A

Normalized expression

KLF6
INO80D
CHERP
CCN2
ATF3
DUSP8
CXCL8
CXCL1
PRDM1
LCAT
CXCL3
BICRA
NR4A3
ARL14
FOSB
P2RY11
PYGM
IL6
CCL20
ALOXE3
ZNF296
ARID5B
WNK4
DNAH17
SRRM2
DUSP1
LAMA5
AMOTL2
PPRC1
KMT2B
SPEN
SRCAP
NCOR2
KMT2D
PPP1R15A
MUC5B
NUAK2
POLR2A
FOSL2
LENG8
ZNF217
CHD2
C15orf39
CCN1
KDM6B
CXCL2
NFKBIZ
ANKRD1
SETD1B
SETD1A
EP400
TSC22D2
LATS2
LIF
MAP3K14
JUN
IRS2
ABL2
GADD45B
SOCS3

C

RNASEL KO

Control   1 µM   5 µM

2-5A

JUN

FOSB

H3

D

RNASEL KO

Control   1 µM   5 µM

2-5A

JNK

P-JNK

p38

P-p38

P-ZAKα

ZAKα

E

WT

Control   1 µM   5 µM

2-5A

ERK

P-ERK

S2.

S3.

A

B

S4.

A

B

S5.

S6.

A

B