ORIGINAL ARTICLE

Repeat turnover meets stable chromosomes: repetitive DNA sequences mark speciation and gene pool boundaries in sugar beet and wild beets

Nicola Schmidt[1], Katharina Sielemann[2,3], Sarah Breitenbach[1], Jörg Fuchs[4], Boas Pucker[5], Bern Weisshaar[2], Daniela Holtgräwe*[2], and Tony Heitkam*[1]

[1]Faculty of Biology, Technische Universität Dresden, 01069 Dresden, Germany

[2]Genetics and Genomics of Plants, Center for Biotechnology (CeBiTec) & Faculty of Biology, Bielefeld University, 33615 Bielefeld, Germany

[3]Graduate School DILS, Bielefeld Institute for Bioinformatics Infrastructure (BIBI), Bielefeld University, 33615 Bielefeld, Germany

[4]Leibniz Institute of Plant Genetics and Crop Plant Research (IPK) Gatersleben, 06466 Stadt Seeland, Germany

[5]Plant Biotechnology and Bioinformatics, Institute of Plant Biology & Braunschweig Integrated Centre of Systems Biology (BRICS), TU Braunschweig, 38106 Braunschweig, Germany

*Manuscript corresponding authors:

Tony Heitkam
Faculty of Biology
Technische Universität Dresden
01069 Dresden
Germany
Phone: (+49) 351 463 39593
E-Mail: tony.heitkam@tu-dresden.de

Daniela Holtgräwe
CeBiTec & Faculty of Biology
Bielefeld University
33615 Bielefeld
Germany
(+49) 521 106 8724
dholtgra@cebitec.uni-bielefeld.de

Running title: Repeats in Beets

ABSTRACT

• *Background*

Sugar beet (*Beta vulgaris* subsp. *vulgaris*) and its crop wild relatives share a base

chromosome number of nine and similar chromosome morphologies. Yet, interspecific

breeding is impeded by chromosome and sequence divergence that is still not fully

understood. Since repetitive DNA sequences represent the fastest evolving parts of the

genome, they likely impact genomic variability and contribute to the separation of beet

gene pools. Hence, we investigated if innovations and losses in the repeatome can be

linked to chromosomal differentiation and speciation.

• *Results*

We traced genome- and chromosome-wide evolution across sugar beet and twelve wild

beets comprising all sections of the beet genera *Beta* and *Patellifolia*. For this, we

combined data from short and long read sequencing, flow cytometry, and cytogenetics

to build a comprehensive data framework for our beet panel that spans the complete

scale from DNA sequence to chromosome up to the genome.

Genome sizes and repeat profiles reflect the separation of the beet species into three

gene pools. These gene pools harbor repeats with contrasting evolutionary patterns: We

identified section- and species-specific repeat emergences and losses, e.g. of the

retrotransposons causal for genome expansions in the section *Corollinae/Nanae*. Since

most genomic variability was found in the satellite DNAs, we focused on tracing the 19

beetSat families across the three beet sections/genera. These taxa harbor evidence for

contrasting strategies in repeat evolution, leading to contrasting satellite DNA profiles

and fundamentally different centromere architectures, ranging from chromosomal

uniformity in *Beta* and *Patellifolia* species to the formation of patchwork chromosomes

in *Corollinae/Nanae* species.

• *Conclusions*

We show that repetitive DNA sequences are causal for genome size expansion and

contraction across the beet genera, providing insights into the genomic underpinnings of

beet speciation. Satellite DNAs in particular vary considerably among beet taxa, leading

to the evolution of distinct chromosomal setups. These differences likely contribute to

the barriers in beet breeding between the three gene pools. Thus, with their

isokaryotypic chromosome sets, beet genomes present an ideal system for studying the

link between repeats, genome variability, and chromosomal differentiation/evolution

and provide a theoretical basis for understanding barriers in crop breeding.

KEY WORDS

genome divergence; chromosomes; speciation; repetitive DNA; transposable elements;

satellite DNAs; crop wild relatives; sugar beet; *Beta*; *Patellifolia*

Background

Interplay between chromosomal stability and genome evolution

Large evolutionary leaps in terms of speciation and genome divergence are occasionally

accompanied by chromosome number changes caused by polyploidisation events and/or

structural rearrangements (*Arabidopsis*: Yogeeswaran *et al.*, 2005; *Carex*: Escudero *et*

al., 2012 & 2015; *Cylicomorpha*: Rockinger *et al.*, 2016). In contrast, some genera

display genomic divergence despite harboring karyotypes with stable base chromosome

numbers across all known species (McCann *et al.*, 2020; Vitales *et al.*, 2020a; Pellicer

et al., 2021). That even extends to large intraspecific differences in genome size despite

a consistent chromosome number (*Euphrasia*: Becher *et al.*, 2021). Therefore, the

question arises how inter- and intraspecific genomic divergence accumulates while the

chromosomal setup is maintained.

One of the links between the genome and its partitioning into chromosomes are

repetitive DNA sequences, which provide structure to eukaryotic karyotypes. Despite

their fast evolution, repetitive DNA sequences have a conserved function by providing

sequence material to the main structural regions, such as centromeres and telomeres.

Nevertheless, how the evolution of the repeatome itself relates to the global

chromosome and genome evolution is still a matter of debate (e.g. centromere paradox;

Henikoff *et al.*, 2001; Presting, 2018).

Species of the Beta *and* Patellifolia *genera are marked by dynamic genomes, but stable*

chromosome numbers

We wondered how genomic diversity despite stability of chromosomal number is

reflected in the repeat composition. To test this, we used species of the beet genera *Beta*

and *Patellifolia* as examples – plant taxa that have been used to study repetitive DNA

evolution for over thirty years (Schmidt *et al.*, 1990; and later publications from the lab

of late Thomas Schmidt). The members of these two genera belong to chromosomally

stable taxa, all marked by a base chromosome number of x=9 (despite some higher

ploidies) and containing roughly equally-sized, metacentric chromosomes.

The genera *Beta* and *Patellifolia* comprise at least eleven species that are separated by

up to 38.4 million years of evolution (Hohmann *et al.*, 2006). Species from the genus

Beta are divided into two or three different sections (*Beta*, *Corollinae/Nanae*)

depending on whether the endemic species *B. nana* is considered a separate section

(Kadereit *et al.*, 2006; Frese and Ford-Lloyd, 2020; Sielemann *et al.*, 2022). Cultivated

beets such as sugar beet are varieties of *B. vulgaris* subsp. *vulgaris* within the section

Beta. The presumed progenitor of all cultivated beets, *B. vulgaris* subsp. *maritima*, also

belongs to this section (Frese and Ford-Lloyd, 2020). For better readability, *B. vulgaris*

subsp. *vulgaris*, *B. vulgaris* subsp. *adanensis*, and *B. vulgaris* subsp. *maritima* will be

hereafter referred to as *B. vulgaris*, *B. adanensis*, and *B. maritima*, respectively.

Sugar beet is a relatively young crop that went through an exceptionally narrow

bottleneck during its 200 years of domestication (Fischer, 1989). This has led to very

low genetic diversity and a loss of several valuable traits such as pathogen resistances

and tolerance to adverse environmental conditions (e.g. drought, salinated soil; Panella

et al., 2020). Therefore, it is necessary to harvest genetic variation in the beet

germplasm to improve cultivated beet varieties. However, crossing experiments of

sugar beet with many of its crop wild relatives (CWRs) resulted in reproduction-

defective offspring such as sterile and semi-fertile, as well as aneuploid and

anorthoploid plants indicating that postzygotic isolation mechanisms, e.g. the lack of

chromosome homology due to larger genomic differences, are causal for the limited

gain of genetically improved seeds rather than prezygotic isolation mechanisms (Frese

and Ford-Lloyd, 2020).

Based on the crossability with *B. vulgaris*, the wild beets are grouped into three

different gene pools. These correspond to the three beet main taxa with section *Beta*

species representing the primary gene pool, *Corollinae/Nanae* species representing the

secondary gene pool, and *Patellifolia* species representing the tertiary gene pool. Given

the fact that di- and polyploid species are present in all of these gene pools (Frese and

Ford-Lloyd, 2020) and due to the chromosome uniformity across all beet clades,

polyploidisation and restructuring of chromosomes seem to play a rather subordinate

role in the emergence of genomic variety in *Beta* and *Patellifolia* species. Instead,

differences at the DNA sequence level (Sielemann *et al.*, 2023a) may be the cause of

flawed chromosome pairing, thus resulting in the observed crossing barriers.

The rapid evolution of repetitive elements can be a major factor in reduced pairing

between homologous and homoeologous chromosomes (Dvorak, 1983) as it leads to

varying repeat compositions in the genomes of even closely related species. Repetitive

DNA sequences include transposable elements (TEs) and tandem repeats (TRs) that

differ in their origin, amplification mode, and sequence characteristics (Bennetzen,
2005; Bennetzen and Wang, 2014).

There are two classes of TEs: class I or retrotransposons with a 'copy-and-paste'
transposition mechanism and an RNA intermediate, and class II or DNA transposons
with a 'cut-and-paste' mechanism, thus DNA being the intermediate in replication.
Class I is divided into two subclasses, those flanked by long terminal repeats (LTR
retrotransposons) and those without (non-LTR retrotransposons; Finnegan, 1989; Wells
and Feschotte, 2020). The classes and subclasses are further divided hierarchically into
order, superfamily, family, subfamily, and lineage (as reviewed in Wicker *et al.*, 2007;
Piégu *et al.*, 2015; Neumann *et al.*, 2019).

TRs on the other hand include ribosomal genes, telomeres and satellite DNAs. Satellite
DNAs (satDNAs) are highly repetitive non-coding sequences that are arranged in large
tandem arrays and contribute to structurally important chromosomal regions such as the
centromeres (Lower *et al.*, 2018; Garrido-Ramos, 2021).

Beet genomes, in particular the sugar beet genome, are well characterized regarding
retrotransposons (Wollrab *et al.*, 2012; Weber *et al.*, 2013; Heitkam *et al.*, 2014;
Schwichtenberg *et al.*, 2016; Maiwald *et al.*, 2021), DNA transposons (Menzel *et al.*,
2006; Menzel *et al.*, 2012), and even related viral relics (Schmidt *et al.*, 2021).
Additionally, at least 13 satDNA families are characterized (e.g. Dechyeva and
Schmidt, 2006; Zakrzewski *et al.*, 2010). In beets, the most iconic repeat identified so
far is the centromeric satDNA pBV (Schmidt and Metzlaff, 1991), which differs from
the centromeric satDNA in more distant wild beets (pTS5, pTS4.1; Schmidt and
Heslop-Harrison, 1996; Gindullis *et al.*, 2001). However, there is not yet a full picture
of the repeat landscape across beet genomes that is needed to identify the evolutionary
relationships across time and across species. Expanding the knowledge about the repeat
landscapes may help to explain the different crossabilities between the beet gene pools.

To understand the impact of repeats on crossing barriers, beet genomic resources are
needed. So far there are three published reference genome sequence assemblies for
sugar beet at our disposal (consecutive RefBeet versions: Dohm *et al.*, 2014;
consecutive EL10 versions: Funk *et al.*, 2018 and McGrath *et al.*, 2020;
KWS2320ONT_v1.0: Sielemann *et al.*, 2023b) and the number of studies comparing
the wild beet germplasm to cultivated varieties is continuously increasing (Sielemann *et*
al., 2022, 2023a; Wascher *et al.*, 2022). Hence, sugar beet and its CWRs are well-suited

to serve as a reference for maintaining stable karyotypes despite global changes across
the repetitive genome.

To illuminate how the karyotypically stable wild beet genomes differ from another, we
focused on repetitive DNA sequences as one of the most impactful contributors to
genomic variability. We asked how repeat evolution across wild beets can be linked to
speciation and, hence, to crossing barriers. We further investigated if the wild beet
germplasm's splits into species, sections, and genera (i.e. primary, secondary, and
tertiary gene pools) are mirrored by broad genomic innovations in the repeatome. For
this, we chose 17 beet accessions, including all major beet taxa, measured their genome
sizes, and determined their chromosome numbers and karyotypes. To add a (pan-
)genomic layer, we generated low-pass whole genome shotgun data for all accessions as
well as long reads for selected beet genomes. After assessing the phylogenetic
placements of these accessions (Sielemann *et al.*, 2022, 2023a), we estimated their
repetitive DNA content and finely classified their repetitive DNA sequences in the
respective TE and TR hierarchies. This not only allows following repetitive DNA
evolution comprehensively across the beet genera, determining repeat gains, losses and
replacements, but also linking it back to chromosomal location and karyotypic stability.

RESULTS

Corollinae *species have the largest monoploid genome sizes among sugar beet and wild*
beets

To provide the foundation for later analyses of genome and chromosome evolution, we
first determined the chromosome configuration and ploidy levels of all 17 cultivated
and wild beet accessions, and then estimated their genome sizes by flow cytometry
(Additional file 1: Fig. S1; Table 1). Building onto the plastome-based phylogenetics
framework (Sielemann *et al.*, 2022), our 17-beet-species-panel allows comprehensive
investigation of the *Beta* and *Patellifolia* germplasm.

The beet genome sizes are in accordance with the respective ploidy level (Table 1;
Additional file 1: Fig. S1). Diploid members of the section *Beta* have 1C genomes sizes
of approx. 700 Mbp, whereas the genome of the sole tetraploid species within this
section, *B. macrocarpa*, is nearly twice as large. In contrast, diploid members of the
section *Corollinae* achieve considerably higher values with an average 1C genome size
of almost 940 Mbp. Also, the polyploid species *B. corolliflora* and *B. intermedia* have

genome sizes slightly larger than mathematically expected (2027 Mbp instead of
1880 Mbp, and approx. 2470 Mbp instead of 2350 Mbp). With a value of approx.
760 Mbp, the genome size of the diploid *B. nana* is more similar to that of the diploids
from the section *Beta* than from the section *Corollinae*. Regarding the sister genus,
diploid *Patellifolia* species have genome sizes similar to that of the diploids from the
section *Beta*, whereas the genome of the tetraploid *P. patellaris* is again slightly larger
than mathematically expected (1502 Mbp instead of 1435 Mbp).

Repeat abundance and genome size correlate positively for all major repeat types,
except for tandem repeats
We estimated repeat proportions in the genomes of all species through individual as
well as comparative clustering using the RepeatExplorer2 pipeline. Combined, the
repeats identified for each species represent between 53% (Bptu) and 68% (Bint) of the
total genome (Additional file 1: Table S1). For better comparability and to minimize the
dependence of the genome size from the ploidy level, polyploid species were also
analyzed with downsampled read sets so that they correspond to a diploid chromosome
set ('ploidy corrected genome size'). In general, we observe a high correlation between
repeat proportion (absolute values) and genome size, with $r2 = 0.996$ ($p < 8.94e{-}17$) for
the ploidy corrected genome sizes (Fig. 1A). Further, we observe a clustering of species
with larger genome sizes and higher repeat contents (*Corollinae* members) vs. species
with smaller genome sizes and lower repeat contents (section *Beta*, *B. nana*, and the
genus *Patellifolia*; Fig. 1A).
When individual repeat classes (i.e. LTR retrotransposons, DNA transposons, and
satDNAs) are considered, the distinct taxonomic groups can be resolved according to
their genome sizes and respective repeat contents (absolute values; Fig. 1B-D,
shadings). Again, a positive correlation can be observed when the overall proportion of
LTR retrotransposons ($r2 = 0.948$, $p = 7.32e{-}9$) and DNA transposons ($r2 = 0.798$, $p =$
$1.22e{-}4$) is plotted against the ploidy corrected genome sizes, respectively (Fig. 1B, C).
However, when specifically focusing on the proportion of satDNAs, an extraordinarily
high amount (up to roughly 14%, see Additional file 1: Table S1) is found in the beet
species with small genome sizes, in particular *B. vulgaris* and *B. adanensis*, whereas the
beet species with the largest genome sizes from the section *Corollinae/Nanae* (with the
exception of *B. macrorhiza*) show the lowest contents of satDNAs (less than 5%). This

results in a negative trend regarding the relation between the satDNA proportion and the

genome size (r2 = -0.166, p > 0.52; Fig. 1D). The same observations apply for the

overall and individual repeat fractions calculated as proportions (Additional file 1:

Fig. S2).

Genomic TEs differences across sugar beet and wild beets reflect the separation into

beet sections and genera

The read clustering results based on a $0.1\times$ coverage of the ploidy corrected genome

size for each species are analyzed. The repetitive fraction of all analyzed genomes is

composed mainly of LTR retrotransposons (14-30% of the genome) with twice the

relative content in *B. corolliflora* and *B. intermedia* (*Corollinae* section) compared to

species of the section *Beta* (see Additional file 1: Table S1). With the exception of

B. vulgaris, in which Ty1-*copia* and Ty3-*gypsy* retrotransposons show a rather balanced

relation to each other, in most other genomes, the Ty3-*gypsy* elements dominate

(Fig. 2A-C, see Additional file 1: Table S1). This observation is most pronounced in

P. webbiana, where the proportion of Ty3-*gypsy* retrotransposons is four times bigger

than that of Ty1-*copia* retrotransposons.

As a substantial difference between species from the section *Beta* and the remaining

taxa (section *Corollinae/Nanae* and genus *Patellifolia*), their most abundant LTR

retrotransposons belong to different superfamilies: In section *Beta*, SIRE/Maximus

retrotransposons from the Ty1-*copia* superfamily contribute the highest TE share (3-

4%), whereas the genomes of the other species consist largely of Athila

retrotransposons from the Ty3-*gypsy* superfamily (4-6% in *Corollinae* species; 8-9% in

Patellifolia species). However, *B. nana* stands out as the proportion of Athila elements

(2%) within its genome is clearly surpassed by the proportion of Ogre (3%) and Tekay

elements (5%). In this respect, *B. nana* resembles the species from the section *Beta* in

which Tekay elements are the most abundant Ty3-*gypsy* retrotransposons as well (2-4%

of the genome, with approximately as many Tekay elements as SIRE/Maximus

elements in the tetraploid *B. macrocarpa*).

Another notable observation is that beets from the section *Corollinae/Nanae* are

characterized by a quite high amount of non-autonomous LTR retrotransposons (3-5%,

see Additional file 1: Table S1) compared to the other species. The investigation of the

corresponding clusters revealed that this increased share can be traced back to one
single, so far uncharacterized element.

Taking a look at the DNA transposon fraction, species from the genus *Beta* are
dominated by EnSpm/CACTA terminal inverted repeat (TIR) transposons, whereas the
most abundant DNA transposons in *Patellifolia* species belong to the hAT superfamily
(Fig. 2D). We find a higher genome proportion of DNA transposons in species with
larger genome sizes, with *Patellifolia* species generally having fewer DNA transposons
compared to beets of the genus *Beta*.

TEs are generally conserved across the beet genomes and show section-specific
abundances

An all-to-all comparison across the beets (accession 1 of every species; see Table 1)
revealed that the overall most abundant repeat, a so far unknown LTR retrotransposon
of the Athila/Errantivirus lineage, occurs in all analyzed genomes, representing a
genome proportion of 0.89% (Bnan1) up to 8.08% (Pweb1; Additional file 1: Fig. S3).
However, the high Athila/Errantivirus proportion in *Corollinae* species is caused by
another LTR retrotransposon originally described as five distinct dispersed repeats by
Gao *et al.* (2000; pBC1054, pBC227, pBC305, pBC507/169, and pBC537). In general,
about half of the repeat families are found in all analyzed beet genomes. The other half
consists of species-, section-, and genus-specific repeats. There are fewer *Patellifolia*-
specific repeats (33 out of 395 clusters) than *Beta*-specific repeats (172 out of 395
clusters), resulting in a lower overall repeat diversity in the genus *Patellifolia*. However,
the *Patellifolia*-specific repeats are highly abundant, whereas all repeats in the *Beta*
genomes (with the exception of some satDNAs, see paragraph below) are rather
moderately abundant. Yet, we also find section-specific repeats with regard to the
sections *Beta* (8 out of 172 genus Beta-specific clusters) and *Corollinae/Nanae* (11 out
of 172 genus *Beta*-specific clusters). Several LTR retrotransposons (Athila, Tekay, Tat,
and Ale/Retrofit elements) seem to be re-amplified in the beet genomes of the
Corollinae/Nanae section, whereas an increased abundance of SIRE/Maximus elements
was observed in the beet genomes of the *Beta* section. Furthermore, a so far unknown
Ogre LTR retrotransposon, as well as the non-autonomous LTR retrotransposon
mentioned in the paragraph above, set *Corollinae/Nanae* apart from all other analyzed
beets (see Fig. 2C and Additional file 1: Fig. S3).

Beet satDNAs break ranks (1): Beta *and* Patellifolia *satDNAs are few, but highly*

amplified, whereas Corollinae/Nanae *satDNAs are diverse and lowly abundant*

Comparing *Beta* and *Patellifolia* genomes, the greatest repeatome difference is found in

the satDNAs (Fig. 3). They contribute roughly 6% to the genomes of *Patellifolia*

species and only 4% to the genomes of the *Corollinae/Nanae* species (with the

exception of *B. macrorhiza*; see Additional file 1: Table S1). In contrast, in genomes of

the section *Beta*, satDNAs contribute up to 14%, which is about as much as the LTR

retrotransposon fraction.

The reason for this divergence is the extraordinarily high abundance of two iconic sugar

beet satDNAs (see Additional file 1: Fig. S3), the centromeric beetSat01-pBV and the

intercalary beetSat02-pEV (Schmidt and Metzlaff, 1991; Schmidt *et al.*, 1991).

BeetSat01-pBV and beetSat02-pEV are highly abundant in all species of the section

Beta contributing up to 8% and 5% of the genomes, respectively (Fig. 3; see Additional

file 1: Table S2). However, beetSat02-pEV also occurs in beets of the section

Corollinae/Nanae and within the sister genus *Patellifolia* with low or moderate

abundance (Fig. 3). In contrast, canonical beetSat01-pBV satDNA arrays (as described

by Schmidt and Metzlaff, 1991; Zakrzewski *et al.*, 2011) are strictly limited to the beets

of the section *Beta*.

Although genomes of the *Patellifolia* genus contain less satDNAs than those of the

section *Beta*, they are also dominated by abundant satDNA families (Fig. 3; see

Additional file 1: Table S2): Again, the most prominent satDNAs in the *Patellifolia*

species are those that constitute the centromeres: the centromeric beetSat03-pTS5 and

the pericentromeric beetSat04-pTS4.1 (Schmidt & Heslop-Harrison, 1996). However, in

sequence and monomer length they are substantially different from beetSat01-pBV. In

addition, we identified another highly abundant and genus-specific satDNA (beetSat05).

The satDNA designated as beetSat06 is the only satDNA that is distributed quite

equally among all beet genomes (Fig. 3; Additional file 1: Table S2). This subtelomeric

satDNA ubiquitously occurs on all chromosomes and was previously described under

different names depending on the respective plant species (e.g. in *B. vulgaris* it is

known as pAv34; Dechyeva & Schmidt, 2006). However, Dechyeva and Schmidt

(2006) found that, in the wild beet *B. nana*, this satDNA is restricted to one single pair

of chromosomes which is consistent with the reduced abundance of beetSat06 that we

observed in *B. nana* using the comparative repeat analysis (Fig. 3).

Generally, within the *Corollinae/Nanae*, the specific satDNA quantities set *B. nana*

slightly apart from the other species within this section (Fig. 3; Additional file 1: Table

S2). This applies for other known satDNAs such as beetSat07-pHC8 (Gindullis *et al.*,

2001; Fig. 4G, H) or beetSat10-pRN (Kubis *et al.*, 1997; Fig. 4I, J), as well as for the

satDNAs newly identified during the comparative repeat analysis (beetSat15–18;

Additional file 1: Data S1 and Data S2). These satDNAs are most pronounced in beet

genomes of the section *Corollinae/Nanae* without the conspicuously high abundances

observed in the section *Beta* and the sister genus *Patellifolia*. Among the satDNAs that

define the *Corollinae/Nanae* genomes, only beetSat13 is prominent, especially in

B. macrorhiza (Fig. 3). This satDNA is also known as pBC1447 in *B. corolliflora* and

ChenSat-1a in the related *Chenopodium quinoa* (Gao *et al.*, 2000; Heitkam *et al.*, 2020).

Overall, regarding the interplay of satDNA diversity and abundance, we conclude that

Beta and *Patellifolia* contain relatively few satDNA families that can reach high copy

numbers. In contrast, species within the *Corollinae/Nanae* accumulate a wide satDNA

variety with only low amplification levels (Fig. 3).

Beet satDNAs break ranks (2): Species- and section-specific array expansions lead to

the emergence of unique satDNA landscapes across the beet genomes

As presented above, individual satDNAs can occur in high abundance in one beet

species and in low abundance in another. To understand, if beet satDNA organization is

retained across species and abundance patterns, we investigated their tandem

arrangement across the three main beet taxa. By using long reads from *B. vulgaris*

(section *Beta*), *B. corolliflora* (section *Corollinae*), and *P. procumbens* (genus

Patellifolia) we showed that nearly all identified satDNAs occur in long tandem arrays

in at least one of the three beet taxa (see dotplot visualization in Additional file 1:

Fig. S4). Only beetSat18 does not show any long arrays in neither of the three species.

This is easily explained as this satDNA is specific for *B. nana*, with head-to-tail

arrangements identified on *B. nana* short reads.

We noted variation in monomer length and sequence (Additional file 1: Fig. S5), often

with section specificity. Sugar beet's main satDNA beetSat01-pBV is especially

intriguing: It occurs only scarcely in *Patellifolia* genomes, forming no tandem arrays.

However, in *B. corolliflora*, the beetSat01-pBV monomer is part of a different,

tandemly arranged repeat, resulting in a much longer monomer size of approx. 1800 bp

(Additional file 1: Fig. S4). This larger tandem repeat shows no further sequence

similarity to publicly available nucleotide or protein database entries. Indeed, it may

serve as a starting point to understand formation of one of the largest satDNA families

in any plant genome. Similar potential start points of satDNA emergence were detected

for beetSat10, beetSat11, beetSat12, beetSat17, and beetSat18 (Additional file 1:

Fig. S4; arrows), suggesting a general pattern.

Further, monomer length variations were detected, indicating section specificity, but

none as striking as for beetSat1-pBV. For instance, beetSat02-pEV variants diverge

slightly in monomer length and sequence among the three beet taxa (Additional file 1:

Fig. S5), whereas beetSat06 shows variation in higher order arrangement.

Overall, regarding satDNA amplification and array formation, we conclude that

different satDNAs amplify in different beet species. Whereas large arrays can form in

one wild beet genome, the same satDNA can occur only as a relic in the next. Similarly,

emergence of species-/section-specific satDNA variants occurs. We observe that the

respective beet satDNA landscapes are characterized by different evolutionary

mechanisms: Whereas few *Beta* and *Patellifolia* satDNAs underwent amplification and

homogenization, the satDNAs landscape of *Corollinae/Nanae* species is still rather

dynamic, mirrored by the emergence of several new satDNAs.

Beet satDNAs break ranks (3): Beta *and* Patellifolia *satDNAs constitute major parts of*

357   *all chromosomes, whereas* Corollinae/Nanae *satDNAs are restricted to*

358   *chromosome subsets*

To understand the structural chromosome makeup across the beets, we again focused on

the three species *B. vulgaris*, *B. corolliflora* and *P. procumbens* to represent the three

main beet taxa. First, we investigated the five main satDNAs: beetSat01-pBV to

beetSat05.

Our two-color FISH onto sugar beet chromosomes shows that beetSat01-pBV (green)

constitutes the centromeres, whereas beetSat02-pEV (red) builds large intercalary

blocks along all *B. vulgaris* chromosomes (Fig. 4A, B; Kubis *et al.*, 1998). In contrast,

beetSat02-pEV is restricted to only a subset of chromosomes and/or comparatively

small intercalary regions along *B. corolliflora* and *P. procumbens* chromosomes

368 (Fig. 4C, D). This indicates that beetSat02-pEV is not only specific in its monomer

369 sequence (see above), but also in its chromosomal localization among the three beet

370 taxa.

371 In *P. procumbens*, beetSat03-pTS5 marks most, but not all centromeres (Fig. 4E, green

372 signals; see also Gindullis *et al.*, 2001). BeetSat05 resides at as many chromosomes (10-

373 14 centromeres; Fig. 4E, blue signals). Thus, each centromere contains either beetSat03-

374 pTS5 or beetSat05, or both. All centromeres are flanked by beetSat04-pTS4.1 that

375 accompanies the centromeric beetSat03-pTS5 and beetSat05 signals, but also occupies

376 some distal locations (Fig. 4E, red signals). All five main satDNAs are restricted to the

377 DAPI-positive heterochromatin in interphase nuclei (Fig. 4B, F). Concluding, we note

378 that chromosomes of *B. vulgaris* and *P. procumbens* – despite their phylogenetic

379 distance – are organized similarly, with main satDNAs building the centromeres and

380 intercalary regions.

381 This is in sharp contrast to the chromosome organization in *B. corolliflora*, a

382 representative of the *Corollinae/Nanae* section: Out of the many satDNAs with varying

383 degrees of amplification and homogenization (see above), we selected six for

384 hybridizations (beetSat07-pHC8, beetsSat08, beetSat10-pRN, beetSat13, beetSat15, and

385 beetSat17; Figure 4G-N). None of them is exclusively localized at the centromeres.

386 Instead, intercalary signals are common as well and (peri-)centromeric signals are

387 restricted to only some chromosomes, often being less prominent than centromeric

388 *Beta/Patellifolia* signals. The comparison of two of these satDNAs at *B. vulgaris* and

389 *B. corolliflora* chromosomes reveals that the signal patterns are more (beetSat07-pHC8)

390 or less (beetSat10-pRN) similar in both genomes (Fig. 4G-J):

391 BeetSat07-pHC8 resides on all *B. vulgaris* chromosomes in intercalary and distal

392 regions, but also close to at least two centromeres (Fig. 4G). In *B. corolliflora*,

393 beetSat07-pHC8 resides in two centromeric and several intercalary regions as well.

394 Major signals are pronounced on six chromosomes (presumably three chromosome

395 pairs; Fig. 4H), including the two centromeric signals.

396 The second satDNA, beetSaat10-pRN, is lowly abundant in *B. vulgaris* (see Fig. 3),

397 thus producing only few, faint and scattered signals (Fig. 4I). In contrast, beetSat10-

398 pRN hybridized strongly to 6-8 chromosomes of *B. corolliflora*, predominantly in the

399 centromeres (Fig. 4J).

To better understand how the patchy satDNA emergence/amplification patterns in the

Corollinae/Nanae affect the chromosomes, we also localized beetSat08, beetSat13,

beetSat15, and beetSat17 (Figure 4K-N):

BeetSat08 resides on nearly all *B. corolliflora* chromosomes (Fig. 4K). Two major and

two minor signals mark the centromeres of four chromosomes, whereas the majority of

signals was rather weak and distributed over the intercalary and distal regions.

BeetSat13, which is known to be associated with the centromeres of *C. quinoa*

chromosomes (ChenSat-1a; Heitkam *et al.*, 2020), can also be found at the centromeres

of at least ten *B. corolliflora* chromosomes (Fig. 4L). BeetSat13 signals near the

centromere were detected on seven additional chromosomes. The remaining

chromosomes did not hybridize with the beetSat13 probe or showed intercalary

beetSat13 signals.

All six major sites of beetSat15 hybridization are associated with *B. corolliflora*

centromeres as well (Fig. 4M).

The comparatively low abundance of beetSat17 (see Fig. 3) resulted in only few

hybridization signals in *B. corolliflora* (Fig. 4N): Signals were detected on twelve

chromosomes with at least two of them showing weak centromeric signals.

Overall, regarding the chromosomal impacts of the vastly different satDNA landscapes

in *Beta* and *Patellifolia* versus *Corollinae/Nanae*, we note that the different

evolutionary patterns affect chromosome structure, especially the centromeres. *Beta* and

Patellifolia centromeres are made up of few, highly abundant, homogenized satDNAs.

Instead, the *Corollinae/Nanae* harbor 'patchwork centromeres': We identified at least

28 (peri-)centromeric signals with six different satDNAs (beetSat07-pHC8, beetSat08,

beetSat10-pRN, beetSat13, beetSat15, beetSat17) on *B. corolliflora* chromosomes.

From this, we conclude that *Corollinae/Nanae* centromeres are constituted by a

multitude of different satDNAs, whereas *Beta* and *Patellifolia* centromeres are

constituted by one or two main satDNAs, respectively.

DISCUSSION

A beet species panel to understand the evolving genome under karyotypic stability

We leverage a comprehensive repeatome study across 17 accessions of cultivated and

wild beet species, spanning two sister genera, *Beta* and *Patellifolia*. We target all major

species across all sections, including higher polyploids. Building onto the plastome-

based phylogenetics framework (Sielemann *et al.*, 2022, 2023a), our 17-beet-species-

panel represents a well-characterized sugar beet and CWR panel with regard to tracing

genome evolution, chromosome stability, pangenomics, and taxonomy.

Beet and wild beet genome size variation results from repeat content fluctuations

Considering the respective ploidy level, the beet genome sizes show relatively moderate

variation: 1.4-fold variation among the diploids, 1.5-fold variation among the

tetraploids, and also 1.5-fold variation among the ploidy corrected genome sizes.

Meanwhile, intraspecific differences are minimal, with almost no variation. For taxa

with high chromosomal variability, e.g. across *Euphrasia* individuals (1.3-fold

variation), structural changes such as the loss or gain of chromosome fragments are

likely responsible for genome size variations (Becher *et al.*, 2021). In contrast, we

hypothesize that genome size variation within plant taxa with stable chromosome

setups, such as the analyzed beet species, depends strongly on genomic repeats. This

can be demonstrated by a clear correlation between genome size and repeat fraction. For

the beet species, we have found such a correlation not only between the overall repeat

content, but especially between the LTR retrotransposons and the respective genome

size, indicating that the amplification and elimination of LTR retrotransposons in

particular is causal for genome size differences between beets of the same ploidy. In

comparison to other angiosperms, the determined correlation is at least as high (e.g.

compared to *Fabeae* sp.; Macas *et al.*, 2015), if not higher (e.g. compared to *Eleocharis*

sp., *Solanum* sp., *Hesperis*-clade sp.; Zedek *et al.*, 2010; Michael, 2014; Gaiero *et al.*,

2019; Hloušková *et al.*, 2019; Gantuz *et al.*, 2021), pointing to a particularly strong

impact of repeats (i.e. LTR retrotransposons) on beet genome sizes.

Specific amplification and elimination of repeats may explain why the genome size of

B. nana rather resembles those of the section *Beta*, even though this species is

repeatedly considered a member of the section *Corollinae* (Kadereit *et al.*, 2006; Frese

and Ford-Lloyd, 2020; Sielemann *et al.*, 2022). However, genome sizes of beets from

the genus *Patellifolia* are most similar to those of the section *Beta*, suggesting that

repeat amplification and/or acquisition caused a genome upsizing in *Corollinae/Nanae*

species.

Repeat abundances mirror the phylogenies of the beet and wild beet sections

With 53-68%, the repetitive fraction of beet genomes represents typical values

compared to other Amaranthaceae (approx. 76% in quinoa: Heitkam *et al.*, 2020;

approx. 51% in spinach: Li *et al.*, 2021). Discrepancies in repeat abundances to a

previous RepeatExplorer2 analysis in *B. vulgaris* (Kowar *et al.*, 2016) are moderate and

result from the exclusion of organellar DNA from our read set as well as the

improvement of the repeat annotation by including a more comprehensive, custom

repeat database. In general, the repeat content results not only from the number of

repetitive sequences, but also from their type: A repeat with a longer element structure

accounts for a larger share of the genome than a shorter repeat. Within the beet

genomes, the most frequent repeats belong to the LTR retrotransposons, namely the

Ty3-*gypsy* retrotransposons, which include the longest known TEs in plants (i.e. Ogre

elements with >23 kb in length; Orozco-Arias *et al.*, 2019). The predominance of Ty3-

gypsy retrotransposons within the repeat fraction was observed for other angiosperms as

well (Kelly *et al.*, 2015; Macas *et al.*, 2015; Gaiero *et al.*, 2019; Hloušková *et al.*, 2019;

Dodsworth *et al.*, 2020). However, the rather balanced share of Ty3-*gypsy* and Ty1-

copia retrotransposons in *B. vulgaris* may reflect different TE dynamics in the

domesticated beet cultivars, whereas the independent amplification and/or acquisition of

Ty3-*gypsy* retrotransposons may have increased their abundance within the CWRs.

We found that most TE families are distributed across all analyzed beet species,

indicating that the present beet repeat set can be traced back to the last common

ancestor of both genera *Beta* and *Patellifolia*. After speciation from this ancestor, a few

new TE families emerged, though rather sparsely. Instead, specific TE

amplification/elimination has led to genus- and section-specific TE abundances. This

global trend in beet genomes is mirrored in the ups and downs of individual TE

families/lineages (e.g. SIRE/Maximus elements: Weber *et al.*, 2010; chromoviruses:

Weber *et al.*, 2013; LINEs: Heitkam *et al.*, 2014; non-autonomous LTR

retrotransposons: Maiwald *et al.*, 2021). Differences in the repeat profiles support the

relationships of the current beet phylogenies (Frese and Ford-Lloyd, 2020; Sielemann *et*

al., 2022) with characteristic repeatomes for species from the sections *Beta*,

Corollinae/Nanae, and from the genus *Patellifolia*, respectively. However, whereas the

amplification of TE families/lineages (in particular Ogre and non-autonomous LTR

retrotransposons) has probably led to an increase in genome size of *Corollinae/Nanae*

species, accumulation of specific TEs in the genomes of *Patellifolia* or section *Beta*

species seem to have taken place at a comparable level, so that no striking differences in

genome size are apparent between those beet taxa. To be more precise, there has been a

strong amplification of few repeats in *Patellifolia* genomes, while many repeats have

been amplified rather moderately in genomes of the section *Beta*.

Since the genomic shock of polyploidisation may stimulate rapid and dynamic genomic

changes such as the activation of retrotransposons and viral elements (McClintock,

1984; Lopez-Gomollon *et al.*, 2021), genome size may subsequently increase. The

overall repeat content is indeed greater within the polyploid beet genomes compared to

their closest diploid relatives, mainly due to an increased abundance of Ty3-*gypsy*

retrotransposons. For most polyploid beet species, we measured larger genome sizes

than mathematically expected, which can thus be attributed to a general accumulation of

all kinds of present Ty3-*gypsy* retrotransposons instead of the targeted amplification of

distinct TEs. During the 'cycle of polyploidy', polyploid plants usually undergo genome

downsizing (Wendel, 2015), which may be the case for *B. macrocarpa* (since its

genome size is smaller than mathematically expected) but not for the other polyploid

beet species, indicating that these are relatively young polyploids (less than 0.9-1.4

million years; Romeiras *et al.*, 2016).

It is often observed that genomic repeat profiles contain a phylogenetic signal

(Dodsworth *et al.*, 2015; McCann *et al.*, 2020; Vitales *et al.*, 2020b; Herklotz *et al.*,

2021). In the case of *Beta* and *Patellifolia* species, this is only partly true: Overall, the

repeatomes enable to separate the beet species into genera and sections. However,

repeatome differences at the subspecies level do not reflect the currently proposed

relationships of the subspecies taxa: As an example, *B. vulgaris* is considered to be

more closely related to *B. maritima* accessions, regardless of their origin, than to

B. adanensis (Wascher *et al.*, 2022). According to kmer-based genomic distance,

B. adanensis may even be considered a distinct species rather than a subspecies

(Wascher *et al.*, 2022). In contradiction with this report, focusing on the three

subspecies *B. adanensis*, *B. vulgaris*, and *B. maritima*, the repeat profiles of the first two

are most similar to each other. This may be explained by the *B. maritima* accessions

used here: both Bmar1 and Bmar2 are from the Atlantic coast and are therefore less

closely related to *B. vulgaris* than its presumed progenitor *B. maritima* from the

Mediterranean area. From this, we conclude that the repeat profiles are not suited to

address still debated phylogenetic questions regarding the (sub-)species level of beets.

This also concerns the non-uniform treatment of *P. procumbens* and *P. webbiana* as

distinct or as the same species.

The abundance of a distinct repeat depends on several factors that either cause an

increase, a reduction, or that keep the current copy number stable. Those factors include

an enhanced amplification (as mentioned above), the targeted or non-targeted

elimination from the genome (e.g. by defense mechanisms and/or the loss of whole

chromosomes or chromosome fragments), and selective pressure for the maintenance of

specific genomic regions. An example in which some of these factors come into play is

the section-specific abundance of endogenous pararetroviruses (EPRVs) found among

the beets. There are three to five times more EPRVs within the genomes of wild beets

from the section *Corollinae* compared to beets from the sister genus *Patellifolia*, the

section *Beta*, and even *B. nana*. Such differences were also found between wild and

cultivated potato species (Gaiero *et al.*, 2019) and it was assumed that the EPRVs

underwent either an increased amplification in wild potatoes or a selective bias in

cultivated potatoes. As for sugar beet, we know that there is a trade-off between the

targeted inactivation and a simultaneous preservation of EPRV sequences (Schmidt *et*

al., 2021). Therefore, we believe that *Corollinae* species accumulated pararetroviral

sequences by the gain of further EPRVs in comparison to the remaining beet species.

Similar section-specific emergence and loss occurred also for other TE

families/lineages, especially among the LTR retrotransposons.

SatDNAs emerge, amplify and vanish without impacting the global genome size

The only repeat type for which no correlation with the beet genome size was found are

the satDNAs. Strikingly, of all repeats, the satDNAs show the most pronounced

specificity in abundance and distribution among the beet genomes, pointing to the fact

that satDNAs are the most dynamic repeats (Garrido-Ramos, 2021). With this study, we

present the first all-encompassing account of every present *Beta/Patellifolia* satDNA,

involving all known as well as so far unpublished satDNAs. By comparison to publicly

available databases, we determined that, with the exception of beetSat02-pEV (also

found in quinoa: Schmidt *et al.*, 2014), beetSat06-pAv34 (pRs34; also found in spinach:

Dechyeva and Schmidt, 2006), and beetSat13 (ChenSat-1a; also found in quinoa:

Heitkam *et al.*, 2020), the satDNAs are specific for and restricted to beet species from

the genera *Beta* and *Patellifolia*.

Usually, the repeatome of plants is constituted mostly by LTR retrotransposons,

accounting for up to 80% of the plant genome size (Orozco-Arias *et al.*, 2019). Only

few plants are known in which the proportion of satDNAs is comparable to that of TEs

(e.g. olive: Barghini *et al.*, 2014; radish: He *et al.*, 2015; *Fritillaria affinis*: Kelly *et al.*,

2015). Here, we present sugar beet (*B. vulgaris*) as another plant species with peculiar

high amounts of satDNAs. However, with diminishing phylogenetic relationship to

sugar beet, the proportion of satDNAs decreases in its CWRs. In addition, each beet

section/genus has its own distinct repeat profile, which sets it apart from the other

(Fig. 5). Thus, we observed satDNAs specific for the genus *Patellifolia*, as well as for

the sections *Corollinae/Nanae* (three different satDNAs, each) and *Beta* (one satDNA).

Even the sole two satDNAs that are present in all analyzed beet species (i.e. beetSat02-

pEV, beetSat06-pAV34) show variability in abundance as well as sequence and

monomer length, resulting in genus- and section-specific variants that may initiate

homogenization processes and sequence shifts in the future.

As beetSat02-pEV and beetSat06-pAV34 are distributed among all analyzed beets

(Fig. 5) and as variants of them are also present in more distantly related plant species

(beetSat02-pEV in quinoa: Schmidt *et al.*, 2014; beetSat06-pAV34 in spinach:

Dechyeva and Schmidt, 2006), we assume that these satDNAs were already present in

the common ancestor of both genera *Beta* and *Patellifolia*. During the subsequent beet

speciation, a great variety of new satDNAs emerged and some of them accumulated

section-specifically by remarkable re-amplification. Such a restriction to individual

sections or species is sometimes known for TEs, but is particularly pronounced for the

fast evolving satDNAs (Orozco-Arias *et al.*, 2019, Garrido-Ramos, 2021). For example,

beetSat13's specificity in abundance is visible even at the level of beet accessions

(Bmrh1 vs. Bmrh2). Moreover, this satDNA can be found in all *Corollinae/Nanae*

species as well as in the more distantly related quinoa (ChenSat-1a: Heitkam *et al.*,

2020), but not in beet genomes from the genus *Patellifolia* and the section *Beta* (Fig. 5).

The patchy distribution of beetSat13 among Amaranthaceae members may be either

explained by its loss in most of the beet genomes except those from the section

Corollinae/Nanae, or by an independent acquisition into the genome of the

Corollinae/Nanae ancestor. It was argued that in quinoa this satDNA emerged from a

CACTA-like TE (Belyayev *et al.*, 2020). Thus, an independent acquisition/generation

of beetSat13 into/within the genome of the *Corollinae/Nanae* ancestor may have

occurred similarly by spawning from a TE or even by the transmission of

extrachromosomal circular molecules consisting of this satDNA (Navrátilová *et al.*,

2008).

SatDNAs impact beet chromosome architecture in a gene pool-dependent manner

Apart from their role as intercalary heterochromatin (beetSat02-pEV) and (sub-

)telomere contributors (beetSat06-pAV34), satDNAs often provide the DNA backbone

onto which the centromeres are formed (Melters *et al.*, 2013; Garrido-Ramos, 2021).

Strikingly, this iconic location is not occupied by the same satDNA among beets, but by

a peculiarly high number of different satDNAs (Fig. 5). It is not unusual for the

centromeric satDNAs to differ even in very closely related species (Henikoff *et al.*,

2001; Melters *et al.*, 2013). However, whereas the centromeric satDNA in beets from

the section *Beta* is well defined and conserved across the whole karyotype (Schmidt and

Metzlaff, 1991; Zakrzewski *et al.*, 2011), beets from the section *Corollinae/Nanae* and

the genus *Patellifolia* seem to differ in their satDNA composition even from

chromosome to chromosome. This phenomenon was previously observed in potatoes

(Gong *et al.*, 2012) and especially in legumes (Avila Robledillo *et al.*, 2020), as well as

in animal species (chicken: Shang *et al.*, 2010). Usually, the most abundant satDNA of

a species is the one constituting the centromeres (Melters *et al.*, 2013). Yet, while this is

true for beets from the section *Beta*, there is no clearly predominant satDNA in beets

from the section *Corollinae/Nanae* and the genus *Patellifolia*, going along with the

observation that the variety of similarly abundant satDNAs is spread over the (peri-

)centromeric regions of subsets of chromosomes. This is particularly remarkable as

these satDNAs (i.e. beetSat07-pHC8 to beetSat11, beetSat13, beetSat15, and beetSat17

in *Corollinae/Nanae* species; beetSat03-pTS5 and beetSat05 in *Patellifolia* species)

differ considerably in their monomer lengths and/or sequences and yet chromosomes

maintain an error-free cell division. Furthermore, centromeric satDNA families of other

closely related plant taxa (e.g. *Arabidopsis* species), albeit species-specific, nevertheless

derived from a common ancestor (summarized by Garrido-Ramos, 2021), which does

not seem to be true for the beets: Monomer sequence and length differences are too

large to be explained by high mutation rates, even though some satDNAs (e.g.

beetSat01-pBV) indeed evolve towards quite considerable changes in monomer

sequence and length (see Additional file 1: Fig. S4). The centromere drive model

(Henikoff *et al.*, 2001) does not seem to apply either, as the presence of multiple

centromeric satDNAs with different sequences precludes any sequence-dependent

coevolution with the kinetochore complex (Avila Robledillo *et al.*, 2020). Considering

the evolutionary mode of satDNAs, which are thought to typically evolve in a concerted

manner (Šatović-Vukšić and Plohl, 2023), it may be that *Corollinae/Nanae* and

Patellifolia species are only at an intermediate stage on the way to centromere and

satDNA homogenization. This is supported by the fact that beet species from the section

Corollinae/Nanae in particular belong to a highly variable hybrid complex (Frese and

Ford-Lloyd, 2020). Hence, hybridization events between genotypes with different

satDNA profiles may frequently disrupt any ongoing satDNA homogenization

processes. This may have led to the observed variety of lowly amplified satDNAs,

which were then selectively retained to form unique satDNA mixes at each centromere

within one nucleus. This would then result in the observed combination of different

chromosome sets with different satDNA architectures. To account for their unique

chromosome make-up, we refer to the *Corollinae/Nanae* chromosomes as patchwork

chromosomes (Fig. 5; middle). Alternatively, keeping a set of different centromeres

may hold an advantage for these beets, especially in chromosome recognition during

meiotic pairing. For the *Corollinae/Nanae*, we cannot yet determine if there is

chromosome-complementarity among individual satDNAs. Nevertheless, multicolor

FISH indicates a possible chromosome-complementarity of beetSat03-pTS and

beetSat05 in the genus *Patellifolia*. Taking together, we detect vastly different satDNA

landscapes that vary in satDNA family number, abundance and location. These provide

the DNA backbone for the formation of overall different chromosome architectures

within a nucleus, leading from overall similar chromosome/centromere structures

(section *Beta*), over conceptually similar chromosomes (*Patellifolia*) towards

completely different chromosomal/centromeric setups (*Corollinae/Nanae*).

Conclusion

In the use of wild germplasm for breeding, crossing barriers often impede the

sustainable introgression of target traits from wild species into crops. This is also the

case for cultivated and wild beet species, which are taxa with isokaryotypic

chromosomes, but which nevertheless exhibit strong crossing borders between gene

pools. We found that global differences between beet genomes can be attributed

primarily to their repeatomes, especially to the specific composition of satDNAs,

whereas the base number and morphology of chromosomes is rather consistent. After

the divergence from the common beet ancestor, genus- and section-specific repeat

variants propagated by independent transposition and centromeres evolved distinctly,

corresponding to the beet phylogeny and the categorization into gene pools. Based on

their repeatome, beet species divide into three different groups (section *Beta* as primary

gene pool, section *Corollinae/Nanae* as secondary gene pool, and genus *Patellifolia* as

tertiary gene pool), supporting the repeated handling of *B. nana* as a member of the

section *Corollinae*. In comparison to the other beet species, this section is characterized

by a peculiarly high number of different satDNAs with comparable abundances. We

found that these satDNAs contribute to the *Corollinae/Nanae* centromeres in a

composite manner, resulting in a highly variable chromosome composition (patchwork

chromosomes). These insights are valuable for future beet breeding as they help to face

the challenge of overcoming postzygotic crossing barriers and provide unique insights

into genome, repeatome, and chromosome evolution in karyotypically stable taxa.

METHODS

Plant material, genome size measurement, and DNA extraction

The 17 investigated beet accessions cover the whole spectrum of species within the

genera *Beta* and *Patellifolia* (see Table 1). Seeds of the *B. vulgaris* subsp. *vulgaris*

genotype KWS2320 were obtained from KWS Saat SE, Einbeck, Germany. The seeds

of all other accessions were obtained from the Leibniz Institute of Plant Genetics and

Crop Plant Research Gatersleben, Germany. The material of the KWS SAAT SE & Co.

KGaA, Einbeck and IPK Gatersleben was transferred under the regulations of the

standard material transfer agreement (SMTA) of the International Treaty. The plants

were grown under long day conditions in a greenhouse.

Nuclei extraction and staining were performed using the CyStain PI OxProtect reagent

kit from Sysmex. Nuclear DNA was measured by flow cytometry, using *Raphanus*

sativus (2C = 1.11 pg DNA) or *Pisum sativum* (2C = 9.07 pg DNA; Doležel *et al.*,

1992) as internal standard. Four DNA estimations were carried out for each plant (5000

nuclei per analysis) on at least two different days. Nuclear DNA content (2C value in

[pg]) was calculated as: sample peak mean/standard peak mean × 2C DNA content of

the standard. DNA amounts in picograms were converted to the number of base pairs

using the conversion factor 1 pg DNA = 0.978 × 109 bp (Doležel *et al.*, 2003). The

mean nuclear DNA content was then calculated for each plant as 1C value (in [Mbp];

Table 1).

Genomic DNA was extracted from approximately 100 mg of fresh leaf tissue samples

using the NucleoSpin® Plant II protocols from Macherey-Nagel. Libraries were

prepared using the TruSeq Nano DNA library preparation kit (Fa. Illumina) and were

sequenced on an Illumina HiSeq1500 sequencer at the CeBiTec Sequencing Core

Facility, Bielefeld University, Germany (250 bp paired-end reads).

Repeat classification and quantification

The Illumina reads were trimmed to a length of 100 bp using Trimmomatic (v0.39;

Bolger *et al.*, 2014). High quality of the trimmed reads was ensured by FastQC

examination (v0.11.5; Andrews, 2020). Bowtie2 (v2.2.6; Langmead and Salzberg,

2012) was applied to identify and subsequently remove reads representing organellar

DNA from the sequence data by mapping them to a database containing publicly

available chloroplast and mitochondrial DNAs of *Beta* species

(https://ncbi.nlm.nih.gov/nuccore). The sequence reads were then subsampled to reduce

the genome coverage to 0.1× for all species, and different numbers of paired-end reads

sampled depending on the genome size. To standardize all read pre-treatments, we have

used the mentioned tools embedded in the preparation module of the ECCsplorer

pipeline (Mann *et al.*, 2022). For the identification and quantification of repetitive

sequences, the similarity-based read clustering method was applied as described by

Novák *et al.* (2010) and implemented in the RepeatExplorer2 pipeline (Novák *et al.*,

2013, 2020). We used the pipeline default settings (90% similarity over 55% of the read

length) and included our custom database of Betoideae repeats

(https://zenodo.org/record/8255813) for individual analyses (each beet accession

distinctly) as well as an all-vs-all comparison across all species (accession 1 of every

species; see Table 1). When comparing different accessions of the same species

(Bmar1/Bmar2, Bmrh1/Bmrh2, Bint1/Bint2, and Bnan1/Bnan2), the RepeatExplorer2

pipeline produced highly similar results for the respective repeat compositions (see

Additional file 1: Table S1). Any discrepancies resulted from the read sampling as we

confirmed by using re-sampled read datasets as input. Hence, only one accession per
species was included in the comparative RepeatExplorer2 analysis.

For this all-vs-all comparison, each read set was downsampled to represent 4% of the
respective genome (coverage of 0.04×) based on 'ploidy corrected genome sizes' (1C
values [Table 1] divided by 2 [for the tetraploid species *B. macrocarpa*, *B. corolliflora*,
and *P. patellaris*] and 2.5 [for the pentaploid species *B. intermedia*], respectively).
Nevertheless, the pipeline reached a limit, retrieving 3,985,090 reads. Although
automatic annotation of read clusters was improved by the inclusion of our custom
Betoideae repeat database, not all clusters were classified. The cluster graph shapes of
all unassigned clusters that represented at least 0.001% of the investigated genomes,
were examined manually and consensus sequences, if provided, were searched using
publicly available databases (https://blast.ncbi.nlm.nih.gov/Blast.cgi). Finally, the
relative abundance of each repeat was calculated based on the number of reads in the
respective cluster, excluding remaining organellar reads.

Sequence comparison
Multiple sequence alignments were generated with the MAFFT (v7.017; Katoh and
Standley, 2013) and MUSCLE (v3.8.31; Edgar, 2004) local alignment tools. They have
been manually refined and used for the calculation of pairwise sequence identities. We
explored and visualized sequences with the multipurpose software Geneious 6.1.8
(Kearse *et al.*, 2012). Dotplots were generated with FlexiDot (Seibt *et al.*, 2018) with a
word size of 20 bp using long reads of *B. vulgaris* (PacBio; accession number
SRX3402137; Funk *et al.*, 2018), *B. corolliflora* (ONT; accession number
ERS13530775; Sielemann *et al.*, 2023a), and *P. procumbens* (ONT; accession number
ERS13530778; Sielemann *et al.*, 2023a).

Generation of satDNA probes
The probe for the *Patellifolia*-specific satDNA beetSat05 was ordered as EXTREmer
oligonucleotides synthesized by *Eurofins Genomics* based on the RepeatExplorer2
consensus sequence (Additional file 1: Data S1).
Standard PCR reactions of genomic *B. corolliflora* (BETA 408) and *B. nana*
(BETA 546) DNA were performed using primer pairs designed for two further beet-
specific satDNAs (beetSat15 and beetSat17; Additional file 1: Table S3). The PCR

conditions were 94 °C for 3 min followed by 35 cycles of 94 °C for 30 s, primer-

specific annealing temperature for 30 s, 72 °C for 25 sec (beetSat15) and 45 s

(beetSat17), respectively, and a final incubation at 72 °C for 5 min. PCR fragments

were purified, cloned and commercially sequenced. Sequenced inserts spanning two

(beetSat17; 279 bp) and six (betSat15; 303 bp) monomers of the respective satDNA

were used as probes for the following hybridization experiments. Their identity to the

respective reference sequence (RepeatExplorer2 consensus sequence) was 96.4%

(beetSat15) and 93.7% (beetSat17), respectively.

Preparation of chromosome spreads and fluorescent in situ hybridization (FISH)

The meristem of young leaves was used for the preparation of mitotic chromosomes.

The plant tissues were treated as described by Schmidt *et al.* (2023). Accumulation of

metaphases was achieved by a combination of an incubation with 2 mM 8-

hydroxyquinoline for 3 h and nitrous oxide for 30 min. Fixed plant material was

digested using the 'leaf enzyme solution II' (Schmidt *et al.*, 2023).

The probes for the chromosomal localization of several satDNAs using FISH were

labeled directly as well as indirectly: The probes 'pBV I' (Schmidt and Metzlaff, 1991;

Kubis *et al.*, 1998) and 'beetSat05' (this manuscript; Additional file 1: Data S1) for

specific centromeric satDNA families were directly labeled with DY647-dUTP

(Dyomics). The probe 'pEV I' (similar to Schmidt *et al.*, 1991; accession number

OY726583) marking an intercalary satDNA family (Kubis *et al.*, 1998) was directly

labeled with DY415-dUTP (Dyomics). The probe 'pTS4.1' for the pericentromeric

satDNA in *Patellifolia* species (Schmidt *et al.*, 1990; Schmidt and Heslop-Harrison,

1996) was indirectly labeled by PCR in the presence of digoxigenin-11-dUTP detected

by antidigoxigenin-fluorescein isothiocyanate (FITC; both from Roche Diagnostics).

The remaining satDNAs ('pTS5': Schmidt *et al.*, 1990, Schmidt and Heslop-Harrison,

1996; 'pHC8': Gindullis *et al.*, 2001; 'pRN1': Kubis *et al.*, 1997; beetSat08: accession

number OY726584 [corresponding to 'BlSat1' from Ha, 2018]; beetSat13

[corresponding to 'ChenSat-1a']: Heitkam *et al.*, 2020; beetSat15 and beetSat17: this

manuscript, accession numbers OY726585 and OY726586) were indirectly labeled by

PCR in the presence of biotin-16-dUTP (Roche Diagnostics) detected by streptavidin-

Cy3 (Sigma–Aldrich). The hybridization procedure was performed as described

previously (Schmidt *et al*., 1994) with a stringency of 82%. Chromosomes were

counterstained with DAPI (4',6'-diamidino-2-phenylindole; Böhringer, Mannheim) and

mounted in an antifade solution (CitiFluor). Slides were examined with a fluorescence

microscope (Zeiss Axioplan 2 imaging) equipped with appropriate filters. Images were

acquired directly with the Applied Spectral Imaging v. 3.3 software coupled to the high-

resolution CCD camera ASI BV300-20A. After separate capture for each fluorochrome,

the individual images were combined computationally and processed using Adobe

Photoshop CS5 software (Adobe Systems, San Jose, CA, USA). We used only contrast

optimization, Gaussian and channel overlay functions affecting all pixels of the image

equally.

DECLARATIONS

Ethics approval and consent to participate

The material of the KWS SAAT SE & Co. KGaA, Einbeck and IPK Gatersleben was

transferred under the regulations of the standard material transfer agreement (SMTA) of

the International Treaty. Plants were grown in accordance with German legislation.

Consent for publication

Not applicable.

Availability of data and materials

Illumina whole genome sequence data are available at EBI under accession numbers as

indicated in Table 1. ONT reads for *B. corolliflora* and *P. procumbens* are available at

EBI under the accession numbers ERS13530775 and ERS13530778. RepeatExplorer2

outputs have been made available at ZENODO (doi: 10.5281/zenodo.7821055;

https://zenodo.org/record/7821055). Satellite DNA consensus sequences and the

sequences used as FISH probes are available in Additional file 1: Data S1 and Data S2.

Furthermore, the cloned sequences of the satDNAs beetSat02-pEV and beetSat08, as

well as the newly identified satDNAs beetSat15 and beetSat17 were submitted to ENA

(accession numbers OY726583-OY726586).

Competing interests

The authors declare no competing interests.

Funding

This work was supported by the German Federal Ministry of Education and Research

(call „Epigenetics: Opportunities for Plant Research", grant 031B1221A). KS was

funded by Bielefeld University.

Authors' contributions

NS, BW, DH, and TH designed the study. NS selected and cultivated the plants. NS,

KS, and BP performed DNA extraction. NS and JF performed flow cytometry. BP

performed sequencing. NS and SB performed FISH. NS implemented the bioinformatic

methodology, analyzed the data and prepared the figures and tables. NS, KS, and TH

wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We acknowledge the KWS SAAT SE & Co. KGaA, Einbeck and the Genbank

Gatersleben for providing seeds and data for the investigated accessions. Computational

resources for RepeatExplorer2 analysis were provided by the ELIXIR-CZ project

(LM2015047), part of the international ELIXIR infrastructure.

SUPPLEMENTARY DATA

Additional supporting information may be found in the online version of this article and

consists of the following. Fig. S1: DAPI-stained chromosomes of 17 different beet

accessions. Fig. S2: Correlation between repeat proportion and genome size in *Beta* and

Patellifolia species. Fig. S3: Comparative repeat composition among *Beta* and

Patellifolia species. Fig. S4: Arrangement of beet satDNA monomers on long reads of

B. vulgaris (Bvul), *B. corolliflora* (Bcor), and *P. procumbens* (Ppro). Fig. S5:

Comparative nucleotide alignment of the three section-specific beetSat02-pEV variants.

Table S1: Genome proportion [%] of different repeat classes and superfamilies (repeat

proportion) in *Beta* and *Patellifolia* species. Table S2: Genome proportion [%] of

different satDNAs (and tandem repeats) across *Beta* and *Patellifolia* species. Table S3:

Primer sequences for the amplification of beet-specific satellite DNAs. Data S1: beetSat

nucleotide sequences derived as RepeatExplorer2 consensuses from the comparative

analysis. Data S2: beetSat sequences used as FISH probes.

LITERATURE CITED

Andrews S. FastQC: a quality control tool for high throughput sequence data. 2020.

http://www.bioinformatics.babraham.ac.uk/projects/fastqc.

Avila Robledillo L, Neumann P, Koblizkova A, Novak P, Vrbova I, Macas J.

Extraordinary sequence diversity and promiscuity of centromeric satellites in the

legume tribe *Fabeae*. Mol Biol Evol. 2020;37:2341–2356.

Barghini E, Natali L, Cossu RM, Giordani T, Pindo M, Cattonaro F, et al. The peculiar

landscape of repetitive sequences in the olive (*Olea europaea* L.) genome. Genome Biol

Evol. 2014;6:776–791.

Becher H, Powell RF, Brown MR, Metherell C, Pellicer J, Leitch IJ, Twyford AD. The

nature of intraspecific and interspecific genome size variation in taxonomically complex

eyebrights. Ann Bot. 2021;128:639–651.

Belyayev A, Josefiová J, Jandová M, Mahelka V, Krak K, Mandák B. Transposons and

satellite DNA: on the origin of the major satellite DNA family in the *Chenopodium*

genome. Mob DNA. 2020;11:20.

Bennetzen JL. Mechanisms of recent genome size variation in flowering plants. Ann

Bot. 2005;95:127–132.

Bennetzen JL, Wang H. The contributions of transposable elements to the structure,

function, and evolution of plant genomes. Annu Rev Plant Biol. 2014;65:505–530.

Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics. 2014;30:2114–2120.

Dechyeva D, Gindullis F, Schmidt T. Divergence of satellite DNA and interspersion of

dispersed repeats in the genome of the wild beet *Beta procumbens*. Chromosome Res.

2003;11:3–21.

Dechyeva D, Schmidt T. Molecular organization of terminal repetitive DNA in *Beta*

species. Chromosome Res. 2006;14:881–897.

Dodsworth S, Chase MW, Kelly LJ, Leitch IJ, Macas J, Novák P, Piednoël M, Weiss-

Schneeweiss H, Leitch AR. Genomic repeat abundances contain phylogenetic signal.

Syst Biol. 2015;64:112–126.

Dodsworth S, Kovarik A, Grandbastien MA, Leitch IJ, Leitch AR. Repetitive DNA

dynamics and polyploidization in the genus *Nicotiana* (Solanaceae). In: Ivanov NV,

Sierro N, Peitsch MC, editors. The tobacco plant genome. Compendium of Plant

Genomes. Springer, Cham; 2020. p. 85–99.

Dohm JC, Minoche AE, Holtgräwe D, Capella-Gutiérrez S, Zakrzewski F, Tafer H, et
al. The genome of the recently domesticated crop plant sugar beet (*Beta vulgaris*).
Nature 2014;505:546–549.

Doležel J, Sgorbati S, Lucretti S. Comparison of three DNA fluorochromes for flow
cytometric estimation of nuclear DNA content in plants. Physiol Plant. 1992;85:625–
631.

Doležel J, Bartoš J, Voglmayr H, Greilhuber J. Nuclear DNA content and genome size
of trout and human. Cytom Part A. 2003;51:127–128.

Dvorak J. Evidence of genetic suppression of heterogenetic chromosome pairing in
polyploid species of *Solanum*, sect. *Petota*. Can J Genet Cytol. 1983;25:530–539.

Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and
space complexity. BMC Bioinformatics 2004;5:113.

Escudero M, Hipp AL, Waterway MJ, Valente LM. Diversification rates and
chromosome evolution in the most diverse angiosperm genus of the temperate zone
(*Carex*, Cyperaceae). Mol Phylogenet Evol. 2012;63:650–655.

Escudero M, Maguilla E, Loureiro J, Castro M, Castro S, Modesto L. Genome size
stability despite high chromosome number variation in *Carex* gr. *laevigata*. Am J Bot.
2015;102:233–238.

Finnegan DJ. Eukaryotic transposable elements and genome evolution. Trends Genet.
1989;5:103–107.

Fischer HE. Origin of the 'Weisse Schlesische Rübe' (white Silesian beet) and
resynthesis of sugar beet. Euphytica. 1989;41:75–80.

Frese L, Ford-Lloyd B. Taxonomy, phylogeny, and the genepool. In: Biancardi E,
Panella L, McGrath J, editors. *Beta maritima*. Springer, Cham; 2020. p. 121–151.

Funk A, Galewski P, McGrath JM. Nucleotide-binding resistance gene signatures in
sugar beet, insights from a new reference genome. Plant J. 2018;95:659–671.

Gaiero P, Vaio M, Peters SA, Schranz ME, de Jong H, Speranza PR. Comparative
analysis of repetitive sequences among species from the potato and the tomato clades.
Ann Bot. 2019;123:521–532.

Gantuz M, Marfil CF, Masuelli RW. Transposable elements and genome expansion in
cultivated and wild potato and tomato species. In: Carputo D, Aversano R, Ercolano
MR, editors. The wild *Solanums* genomes. Compendium of Plant Genomes. Springer,
Cham, 2021. p. 201–214.

Gao D, Schmidt T, Jung C. Molecular characterization and chromosomal distribution of
species-specific repetitive DNA sequences from *Beta corolliflora*, a wild relative of
sugar beet. Genome. 2000;43:1073–1080.

Garrido-Ramos MA. The genomics of plant satellite DNA. In: Ugarković Đ, editor.
Satellite DNAs in physiology and evolution. Progress in Molecular and Subcellular
Biology, vol 60. Springer, Cham. 2021. p. 103–143.

Ghaffari R, Cannon EK, Kanizay LB, Lawrence CJ, Dawe RK. Maize chromosomal
knobs are located in gene-dense areas and suppress local recombination. Chromosoma.
2013;122:67–75.

Gindullis F, Desel C, Galasso I, Schmidt T. The large-scale organization of the
centromeric region in *Beta* species. Genome Res. 2001;11:253–265.

Gong Z, Wu Y, Koblízková A, Torres GA, Wang K, Iovene M, Neumann P, Zhang W,
Novák P, Buell CR, Macas J, Jiang J. Repeatless and repeat-based centromeres in
potato: implications for centromere evolution. Plant Cell. 2012;24:3559–3574.

Ha BH. Structure, organization, and evolution of satellite DNAs in species of the genera
Beta and *Patellifolia*. Doctoral dissertation, Technische Universität Dresden. 2018.
https://nbn-resolving.org/urn:nbn:de:bsz:14-qucosa-238083.

He Q, Cai Z, Hu T, Liu H, Bao C, Mao W, Jin W. Repetitive sequence analysis and
karyotyping reveals centromere-associated DNA sequences in radish (*Raphanus sativus*
L.). BMC Plant Biol. 2015;15:105.

Heitkam T, Holtgräwe D, Dohm JC, Minoche AE, Himmelbauer H, Weisshaar B,
Schmidt T. Profiling of extensively diversified plant LINEs reveals distinct plant-
specific subclades. Plant J. 2014;79:385–397.

Heitkam T, Weber B, Walter I, Liedtke S, Ost C, Schmidt T. Satellite DNA landscapes
after allotetraploidization of quinoa (*Chenopodium quinoa*) reveal unique a and B
subgenomes. Plant J. 2020;103:32–52.

Henikoff S, Ahmad K, Malik HS. The centromere paradox: stable inheritance with
rapidly evolving DNA. Science. 2001;293:1098–1102.

Herklotz V, Kovařík A, Wissemann V, Lunerová J, Vozárová R, Buschmann S, et al.
Power and weakness of repetition – Evaluating the phylogenetic signal from repeatomes
in the family Rosaceae with two case studies from genera prone to polyploidy and
hybridization (*Rosa* and *Fragaria*). Front Plant Sci. 2021;12:738119.

Hloušková P, Mandáková T, Pouch M, Trávníček P, Lysak MA. The large genome size
variation in the *Hesperis* clade was shaped by the prevalent proliferation of DNA
repeats and rarer genome downsizing. Ann Bot. 2019;124:103–120.

Hohmann S, Kadereit JW, Kadereit G. Understanding Mediterranean-Californian
disjunctions: molecular evidence from Chenopodiaceae-Betoideae. Taxon. 2006;55:67–
78.

Kadereit G, Hohmann S, Kadereit JW. A Synopsis of Chenopodiaceae Subfam.
Betoideae and Notes on the Taxonomy of *Beta*. Willdenowia. 2006;Bd. 36, H. 1(Special
Issue: Festschrift Werner Greuter):9–19.

Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol. 2013;30:772–780.

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious
Basic: an integrated and extendable desktop software platform for the organization and
analysis of sequence data. Bioinformatics. 2012;28:1647–1649.

Kelly LJ, Renny-Byfield S, Pellicer J, Macas J, Novák P, Neumann P, et al. Analysis of
the giant genomes of *Fritillaria* (Liliaceae) indicates that a lack of DNA removal
characterizes extreme expansions in genome size. New Phytol. 2015;208:596–607.

Kowar T, Zakrzewski F, Macas J, Kobližková A, Viehoever P, Weisshaar B, Schmidt
T. Repeat composition of CenH3-chromatin and H3K9me2-marked heterochromatin in
sugar beet (*Beta vulgaris*). BMC Plant Biol. 2016;16:120.

Kubis S, Heslop-Harrison J, Schmidt T. A family of differentially amplified repetitive
DNA sequences in the genus *Beta* reveals genetic variation in *Beta vulgaris* subspecies
and cultivars. J Mol Evol. 1997;44:310–320.

Kubis S, Schmidt T, Heslop-Harrison JS. Repetitive DNA elements as a major
component of plant genomes. Ann Bot. 1998;82:45–55.

Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012;9:357–359.

Li N, Li X, Zhou J, Yu L, Li S, Zhang Y, Qin R, Gao W and Deng C. Genome-wide
analysis of transposable elements and satellite DNAs in *Spinacia* species to shed light
on their roles in sex chromosome evolution. Front Plant Sci. 2021;11:575462.

Lopez-Gomollon S, Müller SY, Baulcombe DC. Endogenous virus SRNA regulates
gene expression following genome shock in tomato hybrids. bioRxiv. 2021.
https://doi.org/10.1101/2021.09.20.461014.

Lower SS, McGurk MP, Clark AG, Barbash DA. Satellite DNA evolution: old ideas,
new approaches. Curr Opin Genet Dev. 2018;49:70–78.

Macas J, Novak P, Pellicer J, Čížková J, Koblížková A, Neumann P. In depth
characterization of repetitive DNA in 23 plant genomes reveals sources of genome size
variation in the legume tribe Fabeae. PLoS ONE 2015;10:1–23.

Maiwald S, Weber B, Seibt KM, Schmidt T, Heitkam T. The Cassandra retrotransposon
landscape in sugar beet (*Beta vulgaris*) and related Amaranthaceae: recombination and
re-shuffling lead to a high structural variability. Ann Bot. 2021;127:91–109.

Mann L, Seibt KM, Weber B, Heitkam T. ECCsplorer: a pipeline to detect
extrachromosomal circular DNA (eccDNA) from next-generation sequencing data.
BMC bioinform. 2022;23:1–15.

McCann J, Macas J, Novák P, Stuessy TF, Villaseñor JL, Weiss-Schneeweiss H.
Differential genome size and repetitive DNA evolution in diploid species of
Melampodium sect. *Melampodium* (Asteraceae). Front Plant Sci. 2020;11:362.

McClintock B. The significance of responses of the genome to challenge. Science.
1984;226:792–801.

McGrath JM, Funk A, Galewski P, Ou S, Townsend B, Davenport K, et al. A
contiguous *de novo* genome assembly of sugar beet EL10 (*Beta vulgaris* L.). DNA Res.
2023;30:dsac033.

Melters DP, Bradnam KR, Young HA, Telis N, May MR, Ruby JG, et al. Comparative
analysis of tandem repeats from hundreds of species reveals unique insights into
centromere evolution. Genome Biol. 2013;14:1–20.

Menzel G, Dechyeva D, Keller H, Lange C, Himmelbauer H, Schmidt T. Mobilization
and evolutionary history of miniature inverted-repeat transposable elements (MITEs) in
Beta vulgaris L.. Chromosome Res. 2006;14:831–844.

Menzel G, Krebs C, Diez M, Holtgräwe D, Weisshaar B, Minoche AE. Survey of sugar
beet (*Beta vulgaris* L.) *hAT* transposons and MITE-like *hATpin* derivatives. Plant Mol
Biol. 2012;78:393–405.

Michael TP. Plant genome size variation: bloating and purging DNA. Brief Funct
Genomics. 2014;13:308–317.

Navrátilová A, Koblizkova A, Macas J. Survey of extrachromosomal circular DNA
derived from plant satellite repeats. BMC Plant Biol. 2008;8:1–13.

Neumann P, Novák P, Hoštáková N, Macas J. Systematic survey of plant LTR-retrotransposons elucidates phylogenetic relationships of their polyprotein domains and provides a reference for element classification. Mob DNA. 2019;10:1–17.

Novák P, Neumann P, Macas J. Graph-based clustering and characterization of repetitive sequences in next-generation sequencing data. BMC Bioinform. 2010;11:1–12.

Novák P, Neumann P, Pech J, Steinhaisl J, Macas J. RepeatExplorer: a Galaxy-based web server for genome-wide characterization of eukaryotic repetitive elements from next generation sequence reads. Bioinformatics. 2013;29:792–793.

Novák P, Neumann P, Macas J. Global analysis of repetitive DNA from unassembled sequence reads using RepeatExplorer2. Nat Protoc. 2020;15:3745–3776.

Orozco-Arias S, Isaza G, Guyot R. Retrotransposons in plant genomes: structure, identification, and classification through bioinformatics and machine learning. Int J Mol Sci. 2019;20:3837.

Panella LW, Stevanato P, Pavli O, Skaracis G. Source of useful traits. In: Biancardi E, Panella L, McGrath J, editors. *Beta maritima*. Springer, Cham; 2020. p. 167–218.

Pellicer J, Fernández P, Fay MF, Michálková E, Leitch IJ. Genome size doubling arises from the differential repetitive DNA dynamics in the genus *Heloniopsis* (Melanthiaceae). Front Genet. 2021;12:726211.

Piégu B, Bire S, Arensburger P, Bigot Y. A survey of transposable element classification systems - A call for a fundamental update to meet the challenge of their diversity and complexity. Mol Phylogenet Evol. 2015;86:90–109.

Presting GG. Centromeric retrotransposons and centromere function. Curr Opin Genet Dev. 2018;49:79–84.

Rockinger A, Carvalho FA, Renner SS. Chromosome number reduction in the sister clade of *Carica papaya* with concomitant genome size doubling. Am J Bot. 2016;103:1082–1088.

Romeiras MM, Vieira A, Silva DN, Moura M, Santos-Guerra A, Batista D, et al. Evolutionary and biogeographic insights on the Macaronesian *Beta-Patellifolia* species (Amaranthaceae) from a time-scaled molecular phylogeny. PLoS ONE. 2016;11:e0152456.

Šatović-Vukšić E, Plohl M. Satellite DNAs – From localized to highly dispersed genome components. Genes. 2023;14:742.

Schmidt N, Seibt KM, Weber B, Schwarzacher T, Schmidt T, Heitkam T. Broken,
silent, and in hiding: tamed endogenous pararetroviruses escape elimination from the
genome of sugar beet (*Beta vulgaris*). Ann Bot. 2021;128:281–299.

Schmidt N, Weber B, Klekar J, Liedtke S, Breitenbach S, Heitkam T. Preparation of
mitotic chromosomes using the dropping technique. In: Heitkam T, Garcia S, editors.
Walker J, series editor. Methods in Molecular Biology: Plant Cytogenetics and
Cytogenomics. Humana New York, US. 2023. p. 151–162.

Schmidt M, Hense S, Minoche AE, Dohm JC, Himmelbauer H, Schmidt T, Zakrzewski
F. Cytosine methylation of an ancient satellite family in the wild beet *Beta procumbens*.
Cytogenet Genome Res. 2014;143:157–167.

Schmidt T, Junghans H, Metzlaff M. Construction of *Beta procumbens*-specific DNA
probes and their application for the screening of *B. vulgaris* x *B. procumbens* (2n = 19)
addition lines. Theoret Appl Genetics. 1990;79:177–181.

Schmidt T, Jung C, Metzlaff M. Distribution and evolution of two satellite DNAs in the
genus *Beta*. Theoret Appl Genetics. 1991;82:793–799.

Schmidt T, Metzlaff M. Cloning and characterization of a *Beta vulgaris* satellite DNA
family. Gene. 1991;101:247–250.

Schmidt T, Heslop-Harrison JS. Variability and evolution of highly repeated DNA
sequences in the genus *Beta*. Genome. 1993;36:1074–1079.

Schmidt T, Schwarzacher T, Heslop-Harrison JS. Physical mapping of rRNA genes by
fluorescent *in-situ* hybridization and structural analysis of 5S rRNA genes and
intergenic spacer sequences in sugar beet (*Beta vulgaris*). Theoret Appl Genetics.
1994;88:629–636.

Schmidt T, Heslop-Harrison JS. High-resolution mapping of repetitive DNA by *in situ*
hybridization: molecular and chromosomal features of prominent dispersed and
discretely localized DNA families from the wild beet species *Beta procumbens*. Plant
Mol Biol. 1996;30:1099–1113.

Schwichtenberg K, Wenke T, Zakrzewski F, Seibt KM, Minoche A, Dohm JC, et al.
Diversification, evolution and methylation of short interspersed nuclear element
families in sugar beet and related Amaranthaceae species. Plant J. 2016;85:229–244.

Seibt KM, Schmidt T, Heitkam T. FlexiDot: highly customizable, ambiguity-aware
dotplots for visual sequence analyses. Bioinformatics. 2018;34:3575–3577.

Shang WH, Hori T, Toyoda A, Kato J, Popendorf K, Sakakibara Y, et al. Chickens
possess centromeres with both extended tandem repeats and short non-tandem-repetitive
sequences. Genome Res. 2010;20:1219–28.

Sielemann K, Pucker B, Schmidt N, Viehöver P, Weisshaar B, Heitkam T, Holtgräwe
D. Complete pan-plastome sequences enable high resolution phylogenetic classification
of sugar beet and closely related crop wild relatives. BMC Genomics 2022;23:113.

Sielemann K, Schmidt N, Guzik J, Kalina N, Pucker B, Viehöver P, et al. Pangenome of
cultivated beet and crop wild relatives reveals parental relationships of a tetraploid wild
beet. bioRxiv. 2023a; https://doi.org/10.1101/2023.06.28.546919.

Sielemann K, Pucker B, Orsini E, Elashry A, Schulte L, Viehoever P, et al. Genomic
characterization of a nematode tolerance locus in sugar beet. bioRxiv. 2023b;
https://doi.org/10.1101/2023.06.22.546034.

Vitales D, Álvarez I, Garcia S, Hidalgo O, Feliner GN, Pellicer J, et al. Genome size
variation at constant chromosome number is not correlated with repetitive DNA
dynamism in *Anacyclus* (Asteraceae). Ann Bot. 2020a;125:611–623.

Vitales D, Garcia S, Dodsworth S. Reconstructing phylogenetic relationships based on
repeat sequence similarities. Mol Phylogenet Evol. 2020b;147:106766.

Wascher FL, Stralis-Pavese N, McGrath JM, Schulz B, Himmelbauer H, Dohm JC.
Genomic distances reveal relationships of wild and cultivated beets. Nat Commun.
2022;13:2021.

Weber B, Schmidt T. Nested Ty3-*gypsy* retrotransposons of a single *Beta procumbens*
centromere contain a putative chromodomain. Chromosome Res. 2009;17:379–396.

Weber B, Wenke T, Frömmel U, Schmidt T, Heitkam T. The Ty1-*copia* families
SALIRE and Cotzilla populating the *Beta vulgaris* genome show remarkable
differences in abundance, chromosomal distribution, and age. Chromosome Res.
2010;18:247–263.

Weber B, Heitkam T, Holtgräwe D, Weisshaar B, Minoche AE, Dohm JC, et al. Highly
diverse chromoviruses of *Beta vulgaris* are classified by chromodomains and
chromosomal integration. Mobile DNA. 2013;4:1–16.

Wells JN, Feschotte C. A field guide to eukaryotic transposable elements. Annu Rev
Genet. 2020;54:539–561.

Wendel JF. The wondrous cycles of polyploidy in plants. Am J Bot. 2015;102:1753–
1756.

Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, et al. A unified
classification system for eukaryotic transposable elements. Nat Rev Genet. 2007;8:973–
982.
Wollrab C, Heitkam T, Holtgräwe D, Weisshaar B, Minoche AE, Dohm JC, et al.
Evolutionary reshuffling in the Errantivirus lineage Elbe within the *Beta vulgaris*
genome. Plant J. 2012;72:636–651.
Yogeeswaran K, Frary A, York TL, Amenta A, Lesser AH, Nasrallah JB. Comparative
genome analyses of *Arabidopsis* spp.: inferring chromosomal rearrangement events in
the evolutionary history of *A. thaliana*. Genome Res. 2005;15:505–515.
Zakrzewski F, Wenke T, Holtgräwe D, Weisshaar B, Schmidt T. Analysis of a *c₀t-1*
library enables the targeted identification of minisatellite and satellite families in *Beta*
vulgaris. BMC Plant Biol. 2010;10:1–14.
Zakrzewski F, Weisshaar B, Fuchs J, Bannack E, Minoche AE, Dohm JC, et al.
Epigenetic profiling of heterochromatic satellite DNA. Chromosoma. 2011;120:409–
422.
Zakrzewski F, Weber B, Schmidt T. A molecular cytogenetic analysis of the structure,
evolution, and epigenetic modifications of major DNA sequences in centromeres of
Beta species. In: Jiang J, Birchler JA, editors. Plant centromere biology. Oxford: John
Wiley & Sons. 2013. p. 39–55.
Zakrzewski F, Schubert V, Viehoever P, Minoche AE, Dohm JC, Himmelbauer H, et al.
The CHH motif in sugar beet satellite DNA: a modulator for cytosine methylation. Plant
J. 2014;78:937–950.
Zedek F, Šmerda J, Šmarda P, Bureš P. Correlated evolution of LTR retrotransposons
and genome size in the genus *Eleocharis*. BMC Plant Biol. 2010;10:1–10.
1145
1146

1147 **TABLES**

1148 **Table 1:** Sampled beet and wild beet species, including five-letter code, accession
1149 details, ploidy level / chromosome number, and genome size. Gene pools are marked
1150 according to Frese and Ford-Lloyd, 2020.

beet gene pools	species	code	accession	ENA accession no.	chromosome configuration	genome size 1C [Mbp]
primary	*Beta vulgaris* subsp. *vulgaris*	Bvul1	KWS2320	ERR6110425	2n=2x=18	712
	Beta vulgaris subsp. *adanensis*	Bada1	BETA 1233	ERR6110427	2n=2x=18	709
	Beta vulgaris subsp. *maritima*	Bmar1	BETA 1101	ERR6110429	2n=2x=18	703
	Beta vulgaris subsp. *maritima*	Bmar2	BETA 2322	ERR6110431	2n=2x=18	694
	Beta patula	Bptu1	BETA 548	ERR6110433	2n=2x=18	706
	Beta macrocarpa	Bmca1	BETA 881	ERR6110435	2n=4x=36	1370
secondary	*Beta lomatogona*	Blom1	BETA 674	ERR6110449	2n=2x=18	929
	Beta macrorhiza	Bmrh1	BETA 830	ERR6110451	2n=2x=18	936
	Beta macrorhiza	Bmrh2	BETA 576	ERR6110453	2n=2x=18	946
	Beta corolliflora	Bcor1	BETA 408	ERR6110441	2n=4x=36	2027
	Beta intermedia	Bint1	BETA 431	ERR6110455	2n=5x=45	2468
	Beta intermedia	Bint2	BETA 923	ERR6110457	2n=5x=45	2471
	Beta nana	Bnan1	BETA 546	ERR6110459	2n=2x=18	766
	Beta nana	Bnan2	BETA 570	ERR6110461	2n=2x=18	754
tertiary	*Patellifolia procumbens*	Ppro1	BETA 951	ERR6110463	2n=2x=18	712
	Patellifolia webbiana	Pweb1	BETA 526	ERR6110465	2n=2x=18	723
	Patellifolia patellaris	Ppat1	BETA 892	ERR6110467	2n=4x=36	1502

1151

Fig. 1: Correlation between repeat fraction and genome size in *Beta* and *Patellifolia* species. For better comparability with natural diploids (green), polyploid species were downsampled as artificial diploids (orange). Shapes and shades indicate the different beet sections and genera, respectively. Samples are abbreviated according to Table 1. (A) Positive correlation between the total repeat fraction and genome size. (B) Positive correlation between LTR retrotransposon fraction and genome size. (C) Positive correlation between DNA transposon fraction and genome size. (D) The satellite DNA fraction and genome size tend to relate negatively.

Fig. 2: Repeat composition of several *Beta* and *Patellifolia* species. The analyzed samples and abbreviations accord to accession 1 of the respective species (see Table 1) as the repeat compositions of different accessions from the same species are highly similar (see Additional file 1: Table S1). (A) Proportion of major repeat classes shown in relation to the total genome size. (B) Composition of the Ty1-*copia* LTR retrotransposon fraction. (C) Composition of the Ty3-*gypsy* LTR retrotransposon fraction. (D) Composition of the DNA transposon fraction.

Fig. 3: Quantification of satDNAs within different *Beta* and *Patellifolia* genomes. The relative satDNA abundance is indicated by the size of the circles, which corresponds to the number of short reads within the respective satDNA cluster. The monomer sizes as well as relevant references are provided. Further previously described 'satDNAs' and minisatellites (i.e. Dione/*Fok*I-satellite and Niobe/*Alu*I-satellite: Zakrzewski et al. 2010, 2014; BvMSats: Zakrzewski et al. 2010; BvuSats: Li et al. 2021) were not included as we found that these repeats are not arranged in long arrays and therefore we consider them tandem repeats rather than satDNA. The monomer sequences of newly identified satDNAs can be found in Additional file 1: Data S1.

Fig. 4: Multicolor fluorescent *in situ* hybridization to chromosome spreads of *B. vulgaris*, *B. corolliflora*, and *P. procumbens*. DAPI-stained mitotic chromosomes and interphase nuclei are shown in gray. (A) The centromeric and intercalary localization of the satDNAs beetSat01-pBV (green signals) and beetSat02-pEV (red signals), respectively, become visible on prometaphase chromosomes

of *B. vulgaris*. (B) The interphase nucleus of *B. vulgaris* shows that beetSat01-pBV (green signals) and beetSat02-pEV (red signals) are located in heterochromatic regions. (C, D) The satDNA beetSat02-pEV can also be found on almost all chromosomes of *B. corolliflora* (C) and *P. procumbens* (D) in intercalary as well as distal regions, forming large as well as small arrays. Whereas the signals are mostly restricted on one chromosome arm each in *B. corolliflora* (C), they can be often found on both chromosome arms in *P. procumbens* (D). (E) The centromeric and pericentromeric localization of the satDNAs beetSat03-pTS5 (green signals) and beetSat04-pTS4.1 (red signals), respectively, become visible on prometaphase chromosomes of *P. procumbens*. Most centromeres (14 out of 18) also show large beetSat05 arrays (blue signals); in particular those that are not constituted by beetSat03-pTS5. (F) The interphase nucleus of *P. procumbens* shows that beetSat03-pTS5 (green signals), beetSat04-pTS4.1 (red signals), and beetSat05 (blue signals) are located in heterochromatic regions. (G, H) The satDNA beetSat07-pHC8 was detected on almost all arms of all chromosomes of *B. vulgaris* (G) and *B. corolliflora* (H): Signals were present in intercalary as well as distal regions, forming large as well as small arrays. Moreover, two centromeric beetSat07-pHC8 signals were detected in both species. (I, J) Hybridization sites for beetSat10-pRN are few in *B. vulgaris* (I), whereas large signal clusters were detected on most chromosomes of *B. corolliflora* (J). (I) The low beetSat10-pRN abundance in *B. vulgaris* is revealed by faint signals mostly on both arms of the respective chromosome. (J) The major hybridization sites of the beetSat10-pRN probe on *B. corolliflora* chromosomes are associated with the centromeres. Intercalary hybridization sites can be found mostly on one arm of further chromosomes. (K) The centromere of four *B. corolliflora* chromosomes showed signals (two major, two minor) after hybridization with the beetSat08 probe, whereas almost all other chromosomes showed intercalary and distal signals. (L) BeetSat13 can also be found at the centromeres of at least seventeen *B. corolliflora* chromosomes (ten major and seven minor signals). Eight further chromosomes hybridized in intercalary regions. (M) The beetSat15 hybridization resulted in at least six *B. corolliflora* chromosomes with signals at the centromeres and again eight chromosomes with intercalary signals. (N) BeetSat17 signals were detected on twelve *B. corolliflora* chromosomes with at least two of them being near the centromere. Information on probe labeling and detection can be found in the methods section. Scale bars = 5 µm.

Fig. 5: Scenario for the evolution of satDNAs during the history of beet speciation. As the satDNAs beetSat02-pEV and beetSat06 are also present in the spinach/quinoa outgroup (Dechyeva and Schmidt, 2006; Schmidt *et al.*, 2014), we assume an emergence of these satDNAs even before the split of the beet crown. All other identified satDNAs seem to have appeared earliest in the direct beet ancestor with the subsequent emergence of genus- and section-specific satDNAs and different amplification and differentiation patterns. The age of the respective beet taxa has been estimated in million years ago (mya) by Hohmann *et al.* (2006).