Non-autonomous Vnd acts with autonomous Appl during mushroom body axon growth in *Drosophila*

Claire Marquilly[1], Ana Boulanger[1], Germain U. Busto[1#], Robert P Zinzen[2], Bassem A Hassan[3], Lee G Fradkin[4], and Jean-Maurice Dura[1]

[1] IGH, Univ Montpellier, CNRS, Montpellier, France
[2] Systems Biology Imaging Platform, Max Delbrueck Centrum in der Helmholtz Gemeinschaft, Berlin Institute for Medical Systems Biology, Berlin, Germany
[3] Institut du Cerveau-Paris Brain Institute (ICM), Sorbonne Université, Inserm, CNRS, Hôpital Pitié-Salpêtrière, Paris, France
[4] Institute for Stem Cell Biology and Regenerative Medicine, Stanford University, Stanford, CA USA

[#]Present address: INM, Univ Montpellier, INSERM, Montpellier, France

Abstract

The Amyloid Precursor Protein (APP) is associated with Alzheimer's disease. *Appl* is the single *Drosophila* APP ortholog and is expressed in all neurons throughout development. Appl was previously shown to cell-autonomously modulate axon outgrowth in the mushroom bodies (MBs), the fly olfactory memory center. However, we found that *Appl^d*, the only reported null allele, affects the normal function of *vnd*, the gene just proximal to *Appl*. To decipher developmental defects specifically due to a loss of only *Appl* function, we generated a precise *Appl* null allele (*Appl^{C2.1}*) by CRISPR/Cas9 genomic engineering. With *Appl^{C2.1}*, we confirmed the partial requirement for *Appl* in MB axon outgrowth. We also produced new CRISPR *vnd* alleles removing either *vnd-B* or *vnd-A* function. We report here that *vnd* is also required for MB β-branch axon outgrowth and to a much greater extent than *Appl* itself. Moreover, *vnd* is expressed in neurons close to, but not within, the MB during development and is required non-cell-autonomously for MB axon outgrowth

Introduction

Alzheimer's disease (AD) is associated with extracellular accumulation of amyloid fibrils derived from the Amyloid Precursor Protein (APP). APPs have therefore been intensely investigated, however their physiological function in the brain remains unclear and controversial (Nicolas and Hassan, 2014; Selkoe and Hardy, 2016; Soldano and Hassan, 2014). Although there are three paralogues in mammals (APP, APLP1 and APLP2), *Drosophila* encodes a single APP homologue, called *Appl*, that is expressed in all neurons throughout development. It has been shown that the type I transmembrane protein coding gene *Appl* is a conserved neuronal modulator of a Wnt planar cell polarity (Wnt/PCP) pathway, a regulator of cellular orientation within the plane of an epithelium (Soldano et al., 2013). It has been proposed that Appl is part of the membrane complex

formed by the core PCP receptors Fz (*frizzled*) and Vang (*Van Gogh*) required for axon growth (Liu
et al., 2021; Soldano et al., 2013).
The mushroom bodies (MBs) are two bilaterally symmetric structures in the central brain
that are required for learning and memory (Lin, 2023). Each MB is comprised of 2000 neurons
that arise from 4 neuroblasts. Three types of neurons appear sequentially during development:
the embryonic/early larval γ, the larval α'β', and the late larval/pupal αβ types. Each αβ neuron
projects an axon that branches to send an α branch dorsally, which contributes to the formation
of the α lobe, and a β branch medially, which contributes to the formation of the β lobe (Lee et
al., 1999). Appl's expression continues in adult flies, notably in the MB αβ neurons (Leyssen et al.,
2005; Rabah et al., 2025). So far, *Appl^d* is the only reported null *Appl* allele and it results from a
synthetic genomic deletion removing the *Appl* locus (Luo et al., 1992). The *Appl^d*-bearing
chromosome was selected, after γ irradiation, as a translocation of a partial duplication of the X
chromosome on the Y chromosome to a X chromosome terminal deficiency (S1 Fig). *Appl^d* flies
are viable, fertile and display no gross structural defects in the brain (Luo et al., 1992). However,
the *Appl* signaling pathway is required for proper axon outgrowth in the MBs since *Appl^d* flies
display modestly penetrant axonal defects in αβ neurons. Importantly, Appl is required cell-
autonomously for β-branch axon outgrowth (Cassar and Kretzschmar, 2016; Soldano et al., 2013).
The *ventral nervous system defective* (*vnd*) gene, which is immediately proximal to *Appl*,
encodes a Nk2-class homeodomain transcription factor, that acts in a context-dependent manner
as an activator or repressor and is essential for the development of the nervous system. The *vnd*
gene encodes two proteins: Vnd-A and Vnd-B whose mRNAs arise from two different promoters.
These two proteins differ in their aminoterminal domains and are identical in the remainder of
their sequences. While Vnd-A is a transcription repressor for promoters containing Nk-2 binding
sites, Vnd-B directly activates transcription, also likely via binding to the same binding site.
(Stepchenko et al., 2011). Flies bearing a null allele of vnd in hemi- or homozygous condition are
embryonic lethal (Jimenez et al., 1995).
Interestingly, we found that the *Appl^d* chromosome also genetically affects *vnd* function.
To genetically dissect *Appl* and *vnd* functions, we generated CRISPR alleles that precisely delete
the *Appl* gene without affecting *vnd* function (*Appl$^{C2.1}$*) and that precisely delete each one of the
two *vnd* transcripts without affecting *Appl* (*vnd^{CΔA}* and *vnd^{CΔB}*). To analyze the exact
developmental defects due to the specific loss of the *Appl* function only, we used *Appl$^{C2.1}$* and first
confirmed the partial requirement of *Appl* in MB β-branch axon outgrowth. We showed here that
vnd-A, but not *vnd-B,* is also required for MB β-branch axon outgrowth. Unexpectedly, *vnd* is
expressed in neurons close to, but not within, the MB during development and is required non-
cell-autonomously, likely by promoting the production of a secreted factor(s) involved in the MB
axon outgrowth.
77
78 **Results**
79
Generating new *Appl* CRISPR alleles
The complex chromosome structure resulting in the *Appl^d* allele complicates its subsequent
genetic manipulation and might be a reason for the sensitivity of its phenotype to genetic
background (S1 Fig). Therefore, we produced new null *Appl* alleles by removing the entire *Appl*
transcriptional unit via CRISPR/Cas9-mediated deletion (Doudna and Charpentier, 2014). We

recovered $Appl^{C1.4}$ and $Appl^{C2.1}$, two CRISPR $Appl$ null alleles (Fig 1 and S2 Fig), following an
adaptation of a published protocol to produce ~50kb deletions (Port et al., 2014). $Appl^d$ MBs
display a modestly penetrant cell-autonomous axon growth defect of the β-branches which
manifests as an absence of the β lobe (Liu et al., 2021; Soldano et al., 2013).
Surprisingly, we found a strong difference in the penetrance of the absence of the β lobe
phenotype in the two $Appl$ CRISPR alleles. Although 9% of $Appl^{C2.1}$ MBs lacked the β lobe, slightly
lower than the 14.5% described for $Appl^d$ (Marquilly et al., 2021), $Appl^{C1.4}$ MBs displayed a much
higher penetrance (66%) of this phenotype (Fig 1). We determined the extents of the precise
deletions present in the two alleles via sequencing of PCR amplicons and found that while the
$Appl^{C2.1}$ deletion precisely removed the $Appl$ transcriptional unit, the $Appl^{C1.4}$ allele also removed
a part of the vnd transcriptional unit (Fig 1). We then similarly mapped the $Appl^d$ deletion and
found that, unlike $Appl^{C2.1}$, it removes most of the intergenic region between $Appl$ and vnd which
possibly influences vnd function (Fig 1). We then assessed, by genetic complementation tests, if
the different $Appl$ null alleles possibly affected vnd function. We used vnd^A, a molecularly
characterized lethal allele which impacts both the vnd-A and vnd-B transcripts (Fig 2) (Haelterman
et al., 2014). Although $Appl^{C2.1}/vnd^A$ transheterozygous MBs displayed no anatomical MB
phenotypes, the $Appl^d/vnd^A$ and $Appl^{C1.4}/vnd^A$ MBs displayed 14% and 69% of β lobe absence,
respectively (Fig 2). Flies heterozygous for any of the four mutations individually displayed no MB
phenotypes: vnd^A/+ (n = 72), $Appl^{C1.4}$/+ (n = 60), $Appl^{C2.1}$/+ (n = 42) and $Appl^d$/+ previously
described (Soldano et al., 2013). Therefore, $Appl^d$ and $Appl^{C1.4}$, but not $Appl^{C2.1}$ affect vnd
functions in MB development. We could write $Appl^d = Appl^{null}\ vnd^{weak}$, $Appl^{C1.4} = Appl^{null}\ vnd^{strong}$
and only $Appl^{C2.1} = Appl^{null}\ vnd^+$. Taken together, these data strongly suggest that vnd is also
involved in the MB β-branch axon growth and likely to a much greater extent than $Appl$ itself.
108
Generation of new vnd CRISPR alleles deleted either for vnd-B or vnd-A function.
The two vnd transcripts, vnd-A and vnd-B, produce two different Vnd proteins, respectively Vnd-
A and Vnd-B (Fig 2). It was proposed that, while Vnd-A has its main role during embryogenesis,
Vnd-B acts during metamorphosis (Stepchenko et al., 2011). To determine which Vnd isoform is
required for MB β-branch axon growth, we produced the new vnd alleles $vnd^{CΔA}$ and $vnd^{CΔB}$ by
CRISPR/Cas9 genomic engineering that eliminate either the vnd-A or the vnd-B isoform,
respectively (Fig 3 and S3 Fig). Males bearing a $vnd^{CΔB}$ mutant allele ($vnd^{CΔB}$/Y), as well as $vnd^{CΔB}$/
$Appl^{C1.4}$ females, are viable and have essentially wild-type MBs (Fig 3). Males bearing a $vnd^{CΔA}$
mutant allele ($vnd^{CΔA}$/Y) are embryonic lethal similarly to other vnd null alleles which affect both
isoforms. $Appl^{C1.4}/vnd^{CΔA}$ female MBs displayed a strong phenotype of β lobe absence with a
penetrance (64%) like those of $Appl^{C1.4}/vnd^A$ female MBs (Fig 4). We sought to determine the
respective contribution of the lack of $Appl^+$ and the lack of vnd^+ for the strong β lobe absence in
$Appl^{C1.4}/vnd^{CΔA}$ females. For this purpose, we added either a duplication of $Appl^+$ or a duplication
of vnd^+ to this genotype. We found that $Appl^{C1.4}/vnd^{CΔA}$; Dp-$Appl^+$/+ females displayed 69% of β
lobe absence, a penetrance not distinguishable from that observed without the duplication of
wildtype $Appl$ (p = 0.42). Therefore, in these two types of females ($Appl^{C1.4}/vnd^{CΔA}$; +/+ and
$Appl^{C1.4}/vnd^{CΔA}$; Dp-$Appl^+$/+), the phenotype of β lobe absence seemed entirely due only to vnd
lack-of-function. Indeed, $Appl^{C1.4}/vnd^{CΔA}$; +/+ females could be considered as $Appl^{null}$
$vnd^{strong}/Appl^+\ vnd^{null}$ being homozygous mutant for vnd but heterozygous null for $Appl$. Similarly,
$Appl^{C1.4}/vnd^{CΔA}$; Dp-$Appl^+$/+ females could be considered as $Appl^{null}\ vnd^{strong}/Appl^+\ vnd^{null}$; Dp-

Appl⁺/+ and are homozygous mutant for *vnd* but wild-type for *Appl*. Supporting the hypothesis of
Vnd role in β axon growth, we observed a complete rescue when an extra dose of *vnd⁺* is supplied
as in *Appl^{C1.4}*/*vnd^{CΔA}*; *Dp-vnd⁺*/+ females (Fig 4). *Appl^{C1.4}*/Y males have no *Appl⁺* function (null
allele) and display reduced *vnd⁺* function. Moreover, *Appl^{C1.4}*/Y male MB phenotype is moderately
rescued by the presence of an extra dose of *Appl⁺* but strongly rescued by an extra dose of *vnd⁺*
(Fig 4). Taken together, these results strongly indicate that the *vnd-A* transcript but not the *vnd-*
B transcript is specifically required for MB β-branch axon growth.
136
Vnd is expressed around the MBs and is not required within the MBs.
In order to know where *vnd* is expressed, we employed a *vnd-T2A-GAL4* line where GAL4 is under
the control of endogenous *vnd* regulatory sequences via CRISPR gene targeting so that GAL4 and
vnd are translated from a single mRNA transcript (Chen et al., 2015; Lee et al., 2020). UAS-GFP
labelling revealed a pattern in the ventral ganglion of the third larval instar central nervous system
(L3 CNS) similar to that described with immunostaining with antibodies against Vnd (Stepchenko
et al., 2011) thus validating the use of the *vnd-T2A-GAL4* line as a *bona fide vnd* reporter (S4 Fig).
GFP was detected, in the developing brain, close to the MBs visualized by anti-Fas2 staining, from
L3 to 24 h after puparium formation (APF) (Fig 5A, A'- 5D, D'). Staining reveals a structure in the
brain whose cell bodies are organized in a honeycomb pattern. To determine the identity of the
cell bodies, we performed DAPI staining at the L3 stage. We observed a significant DAPI staining
in these structures indicating that they correspond to cell nuclei (Fig 5E, E'). Moreover, these cell
nuclei are Vnd positive and Repo negative (Fig 5F, F'-5G, G') indicating that these cells correspond
to neurons, rather than glia. The neurites emanating from these cell nuclei are very close to the
developing MB medial lobe from which the adult β lobe develops (Fig 5). Noticeably, we did not
observe any GFP labeling within the MBs themselves. This unexpected expression pattern
suggests a non-cell-autonomous role for Vnd in the MB β-branch axon growth. We tested this
hypothesis by MARCM mosaic analysis which allowed the generation of homozygous *vnd* loss-of-
function MB clones in an otherwise heterozygous genetic background and overcame the
homozygous lethality phenotype (Lee et al., 1999; Lee and Luo, 1999). Mitotic recombination was
induced in late-stage embryos/early first instar larvae and the clones were analyzed at the adult
stage. We obtained 20 MB *vnd^{A/A}* clones that include the αβ neurons. All the 20 *vnd* mutant clones
displayed a β-branch axon growth that looked identical to that in wild-type clones (S5 Fig).
Moreover, we obtained similar results, namely a normal β-branch axon growth, with two
additional different null alleles, *vnd^{Δ38}* (8 clones) and *vnd⁶* (4 clones) (S5 Fig). Taken together,
these results demonstrate that, although *vnd* function is strongly required for MB β-branch axon
growth, *vnd* is neither required nor expressed in the MBs themselves. Therefore, it is most likely
that Vnd regulates MB axon growth by a non-cell-autonomous mechanism.
165
166
167   **Discussion**
168
It has been proposed that Appl is part of the membrane complex formed by the core PCP proteins
Fz and Vang (Soldano et al., 2013). *Appl^d* was, before this report, the only *Appl* null allele
described. *Appl^d* mutant flies are completely lacking Appl function and exhibit 14.5 % MB β-lobe
loss (Marquilly et al., 2021) in accordance with what has been reported (Liu et al., 2021; Soldano

et al., 2013). This modestly penetrant axonal phenotype due to the lack of Appl could be due to
partially redundant function provided by the other transmembrane receptors of the complex.
Flies homozygous for the loss of function allele *Vang*[stbm-6] exhibit 50% β-lobe loss (Liu et al., 2021).
As the *Appl*[d] allele involves complex chromosomic rearrangements (Luo et al., 1992), we
leveraged CRISPR/Cas9 chromosomal engineering to produce a defined deletion of around 50 kb
to specifically eliminate the *Appl* transcriptional unit. We recovered two *Appl* complete deletions
(Fig 1): *Appl*[C1.4] and *Appl*[C2.1] removing around 52 kb and 47 kb and exhibiting 66 % and 9 % β-lobe
loss respectively. DNA sequences analysis revealed that, while the *Appl*[C2.1] deletion removes only
Appl sequences, the *Appl*[C1.4] deletion additionally removes a part of the proximally located
transcriptional unit corresponding to the *vnd* gene (Fig 1). This indicates that in addition to Appl,
vnd is likely involved in the β-branch axon outgrowth. Moreover, as can be expected from the
extent of the deletions and based on genetic complementation tests, the *Appl*[C1.4] deletion, but
not the *Appl*[C2.1] deletion, affects *vnd* function. Most importantly, the *Appl*[d] deletion, which
removes most of the Appl coding sequence and the intergenic region between *Appl* and *vnd*, also
affects *vnd* function (Fig 1 and 2).
The role of *vnd* in β-branch axon outgrowth is complicated by the fact that the viable *vnd*
mutant alleles (*Appl*[d] *and* the *Appl*[C1.4] deletion), do not encode functional Appl. Nevertheless,
employing a small (less than 100 kb) genomic *Appl*[+] duplication, we found that flies lacking only
vnd function still exhibit 69% β-lobe loss and that this mutant phenotype is completely rescued
by a small genomic *vnd*[+] duplication (Fig 4). This demonstrates a requirement of *vnd* function in
MB β-branch axon outgrowth. To better understand the roles of both Vnd isoforms in the MB β-
branch axon outgrowth, we produced two mutations, *vnd*[CΔB] and *vnd*[CΔA] , which remove the *vnd-*
B and the *vnd-A* transcript respectively (Fig 3 and S3 Fig). *vnd-A* mRNA is expressed strongly in
embryos and much less in larvae and adults contrarily to *vnd-B* mRNA (Stepchenko et al., 2011).
It was therefore expected that the phenotype seen in adult MBs of pupal-born neurons should be
due to *vnd-B* rather than *vnd-A*. Surprisingly, although Vnd-A has a significant role in MB axon
growth, apparently Vnd-B plays very little to no role in this process. Noticeably, the MB phenotype
of *Appl*[C1.4]/*vnd*[CΔA] flies is similar in penetrance to that of *Appl*[C1.4]/*vnd*[A] where *vnd*[A] impacts both
the *vnd-A* and *vnd-B* transcripts. This indicates that a lack of *vnd-A* function, and not that of vnd-
B, underlies the poor β-branch axon outgrowth phenotype.
If Vnd interacts with Appl within the MBs, we hypothesized that it might be expressed
within the MBs, and act cell-autonomously like Appl. However, endogenous *vnd* transcription
monitored by GAL4 expression that is translated from the same mRNA transcript and was
validated by anti-Vnd staining did not reveal Vnd expression within the MBs. Rather, *vnd* is
transcribed in cells within a neuronal brain structure near the developing MB medial lobe from
which the adult β lobe develops (Fig 5). Using MARCM mosaic analysis, we show that β axons
extended from MB clones null for *vnd* function exhibit wild-type growth patterns (S5 Fig)
demonstrating a non-cell-autonomous requirement for Vnd. Therefore, we favor the hypothesis
that *vnd* expressed in a MB surrounding brain structure causes the expression of one or more
secreted factors regulating the axon growth of the developing β medial lobe.
213
214
215   **Materials and methods**
216

Drosophila stocks

Drosophila stocks
All crosses were performed on standard culture medium at 25° C. Except where otherwise stated,
all alleles and transgenes have been described previously (http://flystocks.bio.indiana.edu/). The
following alleles were used: $Appl^{C1.4}$, $Appl^{C2.1}$, $vnd^{C\Delta A}$ and $vnd^{C\Delta B}$ were generated in this study. $Appl$-
y^+ (BL #56073), $Appl^d$ (BL #43632), vnd^A (BL #57139), $vnd^{\Delta 38}$ (Chu et al., 1998), vnd^6 (Jimenez et
al., 1995). The following transgenes were used: *UAS-mCD8::GFP* (BL #5137) and the duplications
resulting from genomic transgenes *Dp(1;3) DC430* (BL #32288) and *Dp(1.3) DC009* (BL #30219)
which are both inserted into the same attP site at 65B2. For simplicity, we refer to *Dp(1;3) DC430*
as $Dp\text{-}Appl^+$ and *Dp(1.3) DC009* as $Dp\text{-}vnd^+$. $Dp\text{-}Appl^+$ is ~ 97 kb and contains 11 genes including
the *Appl* gene but not the *vnd* gene. $Dp\text{-}vnd^+$ is also ~ 97 kb and contains 15 genes including the
vnd gene but not the *Appl* gene. We used two GAL4 lines: *vnd-T2A-GAL4* (Lee et al., 2020) which
reveals *vnd* transcription and *c739-GAL4* (BL #7362) expressed in adult αβ MB neurons (Aso et al.,
2009). Recombinant chromosomes were obtained by standard genetic procedures. In the case of
the recombination between y^+ and $Appl^{C2.1}$ (~ 170 kb apart) we screened for [y^+] viable males
coming from $y^+ vnd^{\Delta 38}/y Appl^{C2.1}$ females. Note that $vnd^{\Delta 38}$ is an embryonic lethal mutation. Four
[y^+] viable males out of 11320 males were obtained. One male was sterile, but three $y^+ Appl^{C2.1}$ /Y
males gave progeny (percentage of recombination circa 0.027%) and were confirmed by genomic
PCR. MARCM stocks: *tubP-GAL80, hsFLP122, hsFLP1, FRT19A* (all on the X chromosome).
235
CRISPR-Cas9 strategy
All guide RNA sequences (single guide RNA (sgRNA)) were selected using the algorithm
targetfinder.flycrispr.neuro.brown.edu/ containing 20 nucleotides each (PAM excluded) and are
predicted to have zero off-targets. We selected one pair of sgRNA for the *Appl* gene and two pairs
of sgRNA for the *vnd* gene. For *Appl*, the pair is targeting the entire transcriptional unit (S2 Fig).
For *vnd*, each pair is targeting either the A specific region of *vnd* or the B specific region of *vnd* (S3
Fig). We used the following oligonucleotide pairs: CRISPR-1 Appl fwd and CRISPR-1 Appl rev to
target the all *Appl* region, CRISPR-1 vnd A fwd and CRISPR-1 vnd A rev to target the A region of
vnd, CRISPR-1 vnd B fwd and CRISPR-1 vnd B rev to target the B region of *vnd* (see the
corresponding oligonucleotide sequences in S2 and S3 Fig.). We introduced two sgRNA sequences
into pCFD4 (Port et al., 2014), a gift from Simon Bullock (Addgene plasmid # 49411) by Gibson
Assembly (New England Biolabs) following the detailed protocol at crisprflydesign.org. For PCR
amplification, we used the protocol described on that website. Construct injection was performed
by Bestgene (Chino Hills, CA) and all the transgenes were inserted into the same attP site
(VK00027 at 89E11). Being aware that the DNA excision of nearly 50 kb might be rare, we have
decided to use a positive screen before validation by genomic PCR. We have used a stock where
the *Appl* gene is marked by a y^+ construct. The deletion of the *Appl* gene should therefore be
associated to y^- phenotype. Transgenic males expressing the *Appl* sgRNAs and bearing an
isogenized X chromosome ($y Appl\text{-}y^+ Act\text{-}Cas9 w^*$) were crossed to $FM7c/ph^0 w$ females. 180
crosses were set up and only two crosses gave rise to $y Appl^{deletion?} Act\text{-}Cas9 w^*/FM7c$ females
indicating 1% of CRISPR efficacy. From each of these two crosses, a single $y Appl^{deletion} Act\text{-}Cas9$
$w^* /FM7c$ female was crossed with *FM7c* males to make a stock ($Appl^{C1.4}$ and $Appl^{C2.1}$ where Cas9
was removed). The deletion was then validated by genomic PCR using two pairs of primers:
$Appl^{C1.4}$ fwd (GAGCCAGATACACAAGCACA) / $Appl^{C1.4}$ rev (GGCTTTGTTTACTTCCTGGC) and $Appl^{C2.1}$
fwd (TCCTACTACGTTCCACAATC) / $Appl^{C2.1}$ rev (TAATGCCCAACATATCCAAC). The precise

endpoints of the deletion were determined by sequencing (Genewiz, France). Transgenic males
expressing the different *vnd* sgRNAs were crossed to *y nos-Cas9 w** females bearing an isogenized
X chromosome. 100 crosses were set up for each sgRNA pair, with up to 5 males (because of poor
viability) containing both the sgRNAs and *nos-Cas9,* and 5 *FM7c/ph⁰ w* females. From each
selected cross, a single *y vnd^{deletion?} nos-Cas9 w* /FM7c* female was crossed with *FM7c* males to
make a stock which was validated for the presence of an indel by genomic PCR with primers
flanking the anticipated deletion and the precise endpoints of the deletion were determined by
sequencing (Genewiz, France) using *vnd*-specific primers: *vnd^{CΔA}* fwd (ccaacaaagccgagagtctct) /
vnd^{CΔA} rev (cgggaatttctaagccagggt) and *vnd^{CΔB}* fwd (cgatttggggcgttgtgagta) / *vnd^{CΔB}* rev
(gttgggctttaatccgggagt). A CRISPR efficacy of at least 5% was obtained in both cases. We kept a
single stock from each set of CRISPR experiment: *vnd^{CΔA}* and *vnd^{CΔB}* where Cas9 was removed.

Brain dissection, immunostaining and MARCM mosaic analysis
Adult fly heads and thoraxes were fixed for 1 h in 3.7% formaldehyde in phosphate-buffered
saline (PBS) and brains were dissected in PBS. For larval and pupal brains, brains were first
dissected in PBS and then fixed for 15 min in 3.7% formaldehyde in PBS. They were then treated
for immunostaining as described (Boulanger et al., 2011; Lee and Luo, 1999). Antibodies, obtained
from the Developmental Studies Hybridoma Bank, were used at the following dilutions: Mouse
monoclonal anti–Fas2 (1D4) 1:10, mouse monoclonal anti-Repo (8D1.2) 1:10 and guinea pig
polyclonal anti-vnd (1/1000). Goat secondary antibodies conjugated to Cy3 against mouse or
guinea pig IgG (Jackson Immunoresearch laboratory) were used at 1:300 for detection. DAPI
(Sigma) was used after secondary antibody washes. Tissues were incubated for 10 minutes at
room temperature in a 1/1000 solution from a stock containing 1 mg/ml of DAPI solution, then
washed. To generate clones in the MB, we used the MARCM technique (Lee and Luo, 1999). First
instar larvae were heat-shocked at 37°C for 1 h. Adult brains were fixed for 15 min in 3,7%
formaldehyde in PBS before dissection and GFP visualization.

Production of the anti-Vnd antiserum
Full length vnd-RA was amplified via PCR from cDNA and cloned into a pRSET bacterial expression
vector (ThermoFisher) in frame with an N-terminal 6x-His tag, amplified in DH5alpha and
transformed into BL21 competent cells for protein production. His-tagged VndA was purified over
a Nickel-NTA agarose resin (Qiagen) according to manufacturer's recommendation. Purified
protein was injected into guinea pigs by Squarix GmbH (Marl, Germany, www.squarix.de),
selected by determining that pre-immunization sera did not produce signal in Western blots of
fly extracts. Successive bleeds were screened for distinct signal in Western blots using embryo
extracts and by immunohistochemistry, where signal was expected post 2.5hrs AED.

Microscopy and image processing
Images were acquired at room temperature using a Zeiss LSM 780) equipped with a 40x PLAN
apochromatic 1.3 oil-immersion differential interference contrast objective lens and a Leica SP8
laser scanning confocal microscopes (MRI Platform, Institute of Human Genetics, Montpellier,
France. The immersion medium used was Immersol 518F. The acquisition software used was Zen
2011 (black edition) for the Zeiss and LasX for the Leica. Contrast and relative intensities of the

green (GFP), of the red (Cy3) and of the DAPI channels were processed with ImageJ (Fiji) software.
Settings were optimized for detection without saturating the signal.
306
307 **Statistics**
308 Comparison between two groups expressing a qualitative variable was analyzed for statistical
309 significance using the Chi^2 test. (BiostaTGV: http://biostatgv.sentiweb.fr/?module=tests). Values
310 of p < 0.05 were considered to be significant. Statistical significance was defined as: ns = not
311 statistically different, **p<0.01 and****p<0.0001.
312
313 **Figure legends**
314
Fig 1. Generating new *Appl* CRISPR alleles.
(A) Schematic representation of *Appl* (blue) and *vnd* (green) genes and transcripts. White boxes
represent 5'UTR and 3'UTR, blue and green boxes represent *Appl* and *vnd* coding sequences
respectively. Schematic representations of the *Appl^d*, *Appl$^{C1.4}$* and *Appl$^{C2.1}$* mutant alleles where
the deleted sequences are represented in light blue. The hatched black segment represents the
uncertainty where the deletion of *Appl^d* starts at its 5'end. (B) Representation of the 3' limit of
the *Appl^d*, *Appl$^{C1.4}$* and *Appl$^{C2.1}$* mutant alleles (light blue). The hatched black segment represents
the uncertainty where the *Appl^d* deletion ends in 3'. The orange star represents the position of
the 3' sgRNA used to create the two CRISPR deletions. (C) Wild-type (WT) MB revealed by an anti-
Fas2 staining. The pink and yellow arrowheads show respectively the α and β lobe (n=216 MBs).
(D-E) Anti-Fas2 staining reveals the loss of β lobe on a representative *Appl$^{C1.4}$* brain (n=147 MBs)
(D) and on *Appl$^{C2.1}$* brain (n=253 MBs) (E). The % represents the proportion of loss of β lobe for
each genotype. Scale bar = 50 μm. Full genotypes: (C) Canton-S. (D) y *Appl$^{C1.4}$* *w^{1118}*/Y. (E)
Appl$^{C2.1}$/Y.
329
Fig 2. *Appl^d* and *Appl$^{C1.4}$*, but not *Appl$^{C2.1}$*, are impaired for *vnd* function.
A) Schematic representation of *vnd* gene and transcripts. White boxes represent 5'UTR and 3'UTR,
green boxes represent coding sequences. Red star represents point mutation on *vnd^A* allele.
Schematic representation of *vnd-B* and *vnd-A* transcripts. The *vnd^A* mutation affects both
transcripts. (B-D) Anti-Fas2 staining reveals the loss of β lobe on a representative *Appl^d*/ *vnd^A* brain
(n=214 MBs) (B), *Appl$^{C1.4}$*/ *vnd^A* brain (n=147 MB) (C) but not on *Appl$^{C2.1}$*/ *vnd^A* brain (n=248 MBs)
(D). The % represents the proportion of loss of β lobe for each genotype. Scale bar = 50 μm. Full
genotypes: (B) *Appl^d*/y *vnd^A* *w** FRT19A. (C) y *Appl$^{C1.4}$* *w^{1118}*/ y *vnd^A* *w** FRT19A. (D) *Appl$^{C2.1}$*/ y *vnd^A*
*w** FRT19A.
339
Fig 3. Generating new *vnd* CRISPR alleles deleted either for the *vnd-B* or the *vnd-A* function.
(A) Schematic representation of *vnd* gene and transcripts. White boxes represent 5'UTR and
3'UTR, green boxes represent coding sequences. The deleted sequence region of *vnd^{CΔA}* and
vnd^{CΔB} are represented in light green. (B-C) Anti- Fas2 staining reveals essentially wild-type looking
MBs on a representative *vnd^{CΔB}* brain (n=146 MBs) (B) and *Appl$^{C1.4}$*/ *vnd^{CΔB}* brain (n=159 MB) (C).
The % represents the proportion of loss of β lobe for each genotype. Scale bar = 50 μm. Full
genotypes: (B) y *vnd^{CΔB}* *w^{1118}*/Y. (C) y *Appl$^{C1.4}$* *w^{1118}*/ y *vnd^{CΔB}* *w^{1118}*
347

Fig 4. Vnd-A is required for MB β-branch axon outgrowth.

(A) Quantitation of the rescue of MB β lobe absence of $Appl^{C1.4}$/Y (66%; n= 147 MBs, see a representative brain in Fig.1D) and $Appl^{C1.4}$/$vnd^{C\Delta A}$ (64%; n=198 MBs) (light blue) by an $Appl$ duplication (dark blue) (50%; n=150 MBs, 69%; n=89 MBs) and by a vnd duplication (green) (3%; n=191 MBs ; 0%; n=164 MBs). **p<0.01 and ****p<0.0001. (B-D) Anti-Fas2 staining on a representative rescued $Appl^{C1.4}$/Y; $Dp\text{-}vnd^+$/+ brain (B), $Appl^{C1.4}$/$vnd^{C\Delta A}$; +/+ brain (C) and on $Appl^{C1.4}$/$vnd^{C\Delta A}$; $Dp\text{-}vnd^+$/+ brain (D). The % represents the proportion of loss of β lobe for each genotype. Scale bar = 50 μm. Full genotypes: (B) y $Appl^{C1.4}$ w^{1118}/Y; $Dp\text{-}vnd^+$/+. (C) y $Appl^{C1.4}$ w^{1118}/y $vnd^{C\Delta A}$ w^{1118}; +/+. (D) y $Appl^{C1.4}$ w^{1118}/y $vnd^{C\Delta A}$ w^{1118}; $Dp\text{-}vnd^+$/+.

Fig 5. *vnd* is expressed close to the developing MB medial lobe.

(A-D) Visualization of *vnd* expression in brain using *vnd-GAL4* driven *UAS-GFP* at L3, 0h, 6h, and 24h APF. GFP was visualized on green and anti-FasII staining labelling of MB axons on red. A-D are confocal z stacks taken the whole MB and A'-D' are stacks of five sections from A-D respectively comprising GFP⁺ regions surrounding MB. Arrowheads point to *vnd-GAL4* driven *UAS-GFP* neurites that surround MB medial lobes, which correspond to γ lobes from L3 to 6 h APF (A'-C') and β lobes at 24 h APF (D'). n ≥ 10 for each time point. (E-G) show single confocal sections showing *vnd* expression using *vnd-GAL4* driven *UAS-GFP* (single channel) at L3. E' to G' are merges of single channel E to G respectively and DAPI staining (red on E'), anti-vnd (red on F') and anti-Repo (red on G'). DAPI labelled nuclei in GFP⁺ cell bodies are pointed by arrows in E'. These nuclei were also Vnd positive (arrows in F') but were not labelled by an anti-Repo (white arrow in G') which labels glia nuclei (red dots pointed by a yellow arrow in G') indicating that these Vnd positive nuclei belong to neurons. Scale bars are 50 μm. n ≥ 5 brains for each type of staining. Full genotype: *vnd-T2A-GAL4 w**; 2 x *UAS-mGFP*/+.

Supporting information

S1 Fig. Scheme of $Appl^d$ chromosomic organization

(A) Schematic representation of $Appl^d$ mutant allele. The upper panel represents the wild-type (WT) X (black) and Y (orange) chromosomes. The *elav*, *Appl* and *vnd* genes are represented respectively in yellow, blue, and green boxes. The middle panel represents the deficiency Df (1) RT518 on the X chromosome deleting the distal part of the chromosome including the *elav* and *Appl* genes. The X Dup Y chromosome represents a duplication of the X chromosome linked to the Y chromosome including the distal part of the X chromosome including the *elav* gene. The lower panel represents a proposed structure of the $Appl^d$ chromosome where the deficiency-bearing X chromosome had been linked by gamma irradiation to the X Dup Y chromosome (adapted from (Luo et al., 1992)). Additionally, we found that the Y chromosome behaves genetically as linked to the X chromosome.

S2 Fig. Structure of the $Appl^{C1.4}$ and $Appl^{C2.1}$ CRISPR alleles

(A) Schematic representation of *Appl* (blue) and *vnd* (green) genes and transcripts. White boxes represent 5'UTR and 3'UTR, blue and green boxes represent *Appl* and *vnd* coding sequences

respectively. The yellow stars in *Appl* 5' region and the orange star in *Appl* 3' region represent the
position of the sgRNA used to create *Appl*$^{C1.4}$ and *Appl*$^{C2.1}$ mutants. The red and blue boxes
represent respectively the regions magnified in B and C. (B) Representation of the 5' sequence
targeted by the sgRNA (yellow) and the PAM domain (magenta) in the WT chromosome *Appl*
(blue) (upper line). The lower line is the sequence of the *Appl*$^{C1.4}$ mutant showing the beginning
of the deletion about 1kb after the region targeted by the sgRNA. (C) Representation of the 3'
sequence targeted by the sgRNA (orange) and the PAM domain (magenta) in the WT chromosome
Appl (blue) and *vnd* (green) (upper line). The lower line is the sequence of the *Appl*$^{C1.4}$ mutant
showing the end of the deletion about 6kb after the region targeted by the sgRNA. (D)
Representation of the 5' and 3' sequences targeted by the sgRNA (yellow in 5' and orange in 3')
and the PAM domain (magenta) in the WT chromosome *Appl* (blue) (upper line). The lower line
is the sequence of the *Appl*$^{C2.1}$ mutant showing the beginning of the deletion in the sequence
targeted by the 5' sgRNA (yellow) and the end of the deletion in the sequence targeted by the 3'
sgRNA (orange). Circa 46 kb of *Appl* transcribed sequence including all the coding sequence has
been entirely deleted and replaced by a sequence of 821bp (see below) primarily derived from
the fly retrotransposon micropia-like LTR polyprotein gene (NCBI Blast)**.**

ATAAACGCGTCATTAAGCTGTTCCATTTTCTAATCGCGCAGCGAGTTTCGGTGGTCGAAGTCTAGAATTGAAAAAGAAGAAGAA
GAAGATACTCTGTATGTTATGGTTAAGAATAAAAGCTTAAGTTCAGTTTGTGTTATGAAACACAAAATGTGCCTTTGAACTTGAT
ATTGGTCATTACACGAGGATGTGTAAATGTCAGATGGTTATCCGTCTATGAGAGACGATGGTTATCCGTCTATGAGAGACGATG
GTTATCCGTCTATGAGAGACGATGGTTATCCGTCCATGAGAGACGATGGTTATCCGTCTATGAGAGACGATGGTTATCCGTCTA
TGAGAGACGATGGTTATCCACCTAAGACAGACGATGATTTCAGTTTAGAATCTCTTGTTACGTAAGAAATTTGAATCAAAATTG
GGAGTACTACACGAGGACGTGTCAAGGTCAGGATGGCCGTGTCGGAGCATACTGTATGTACGCTCTGTATGTGATTGTTGCTG
CGTATTGCACGTACGCGCTGTGTGTGTATTGCTGTGCATGGGAAATCTGGTTTTTACCAGAATTCCTAATTTCATCATATTAGTA
ATATTGCATCAGCCGTTAATTGCGCTCTGATTGGTCCTTTGCATTTTGTCGCTTTAAGAAGTACCGTTTTAGCGAGACGACAGCG
CTACAACGCTGTCGTCGTGTGTGGTTCGGGAGCTATGCGGGGAAGTAAGTCTGGCGTTGTCACGCAGTTTTTATAGAAACGTG
AAAGCATAAGGCACCGAAACTCGCTTCGCTAGGTACGCATATGGCACAAAATAATCAAGTTCTA

S3 Fig. Structure of the *vnd*$^{C\Delta B}$ and *vnd*$^{C\Delta A}$ CRISPR alleles
(A) Schematic representation of *vnd* (green) gene and transcripts. White boxes represent 5'UTR
and 3'UTR, green boxes represent *vnd* coding sequences. The yellow stars represent the position
of the sgRNA used to create *vnd*$^{C\Delta B}$ mutant. The orange stars represent the position of the sgRNA
used to create *vnd*$^{C\Delta A}$ mutant. (B) Representation of the sequences targeted by the sgRNA (yellow)
and the PAM domain (blue) in the WT chromosome *vnd* (green) (upper line). The lower line is the
sequence of the *vnd*$^{C\Delta B}$ mutant showing the beginning of the deletion in the sequence targeted
by the 5' sgRNA (yellow) and the end of the deletion in the sequence targeted by the 3' sgRNA
(yellow). (C) Representation of the sequences targeted by the sgRNA (orange) and the PAM
domain (blue) in the WT chromosome *vnd* (green) (upper line). The lower line is the sequence of
the *vnd*$^{C\Delta A}$ mutant showing the beginning of the deletion at one nucleotide from the sequence
targeted by the 5' sgRNA (orange) and the end of the deletion within in the sequence targeted by
the 3' sgRNA (orange).

S4 Fig. Vnd L3 CNS labelling
(A-C) Visualization of *vnd* expression in VNC using both *vnd-GAL4* driven *UAS-GFP* (green) and an
anti-vnd antibody (red) at L3 stages. Images A-C and A'-C' are the same confocal z-projection.

Laser intensities leading to the green color were increased from B, C to B', C' to illustrate the high
degree of colocalization between the two markers. Low laser intensity allowed us to show red
nuclear vnd staining in only some GFP$^+$ cell bodies (white arrowheads in A- C). Soma containing
vnd$^+$ positive nuclei that do not seem to be surrounded by GFP in A-C, can be visualized by
increased green laser intensity A'- C' (yellow arrowheads), which saturates somas visualized in A-
C. Scale bars are 50 μm. n ≥ 5 brains. Full genotype: *vnd-T2A-GAL4 w**; 2 x *UAS-mGFP/+*.
442
S5 Fig. *vnd* null mutant clones
(A-B) A representative neuroblast MARCM clone in a wild-type (WT) brain (A) and in a *vnd^A* brain
(B). Scale bar = 50 μm. (C) Number of clones obtained in WT, *vnd^A* , *vnd^6* and *vnd^{Δ38}*. Full genotypes:
(A) *w* tubP-GAL80 hsFLP122 FRT19A/w^{1118} sn FRT19A; c739-GAL4 UAS-mCD8-GFP/+* (also control
hsFLP122 in C). (B) *w* tubP-GAL80 hsFLP122 FRT19A/y vnd^A w FRT19A; c739-GAL4 UAS-mCD8-*
GFP/+ (also vnd^A in C). (C) control hsFLP1: *w tubP-GAL80 hsFLP1 FRT19A* (from the Bloomington
stock 5133)/*w UAS-mCD8-GFP sn FRT19A; c739-GAL4 UAS-mCD8-GFP/+*. vnd^6: *w tubP-GAL80*
hsFLP1 FRT19A/y vnd^6 w UAS-mCD8-GFP sn FRT19A; c739-GAL4 UAS-mCD8-GFP/+. vnd^{Δ38}: *w*
tubP-GAL80 hsFLP1 FRT19A/vnd^{Δ38} w UAS-mCD8-GFP FRT19A; c739-GAL4 UAS-mCD8-GFP/+.
452
453

Acknowledgments

We thank Aymeric Chartier (IGH, Montpellier) for fruitful discussions regarding the generation of
the *Appl* CRISPR alleles. We thank the Bloomington *Drosophila* Stock Center (Indiana University),
the BioCampus RAM-*Drosophila* facility and the BioCampus imaging facility MRI (Montpellier,
France). We acknowledge BestGene and Genewiz for transgene service and DNA sequencing
respectively. The 1D4 anti-Fasciclin II hybridoma and the 8D12 anti-Repo monoclonal antibody
developed by Corey Goodman were obtained from the Developmental Studies Hybridoma Bank,
created by the NICHD of the NIH and maintained at the University of Iowa, Department of Biology,
Iowa City, IA 52242. We thank Chris Doe for the *vnd^{Δ38}* and *vnd^6* alleles and Tzumin Lee for the
vnd-T2A-GAL4 line.

Funding: C.M. was supported by a PhD grant from the Ministère de l'Enseignement Supérieur et
de la Recherche. C.M. and G.U.B. were supported by the Fondation pour la Recherche Médicale
(FRM) respectively for the 4th PhD year and for a 3-year post-doctoral fellowship. Work in the
laboratory of J.-M.D. was supported by the Centre National de la Recherche Scientifique (CNRS),
the Association pour la Recherche sur le Cancer (grants PJA 20151203422), the FRM (Programme
"EQUIPES FRM2016" project DEQ20160334870) and the ANR 2021 ORIO.

472
473

References

Aso, Y., Grubel, K., Busch, S., Friedrich, A. B., Siwanowicz, I. and Tanimoto, H. (2009). The
mushroom body of adult Drosophila characterized by GAL4 drivers. *J Neurogenet* **23**, 156–
172.

Boulanger, A., Clouet-Redt, C., Farge, M., Flandre, A., Guignard, T., Fernando, C., Juge, F. and
Dura, J. M. (2011). ftz-f1 and Hr39 opposing roles on EcR expression during Drosophila
mushroom body neuron remodeling. *Nat Neurosci* **14**, 37–44.

Cassar, M. and Kretzschmar, D. (2016). Analysis of Amyloid Precursor Protein Function in
Drosophila melanogaster. *Front Mol Neurosci* **9**, 61.

Chen, H. M., Huang, Y., Pfeiffer, B. D., Yao, X. and Lee, T. (2015). An enhanced gene targeting
toolkit for Drosophila: Golic+. *Genetics* **199**, 683–694.

Chu, H., Parras, C., White, K. and Jimenez, F. (1998). Formation and specification of ventral
neuroblasts is controlled by vnd in Drosophila neurogenesis. *Genes Dev* **12**, 3613–3624.

Doudna, J. A. and Charpentier, E. (2014). Genome editing. The new frontier of genome
engineering with CRISPR-Cas9. *Science* **346**, 1258096.

Haelterman, N. A., Jiang, L., Li, Y., Bayat, V., Sandoval, H., Ugur, B., Tan, K. L., Zhang, K., Bei, D.,
Xiong, B., et al. (2014). Large-scale identification of chemically induced mutations in
Drosophila melanogaster. *Genome Res* **24**, 1707–1718.

Jimenez, F., Martin-Morris, L. E., Velasco, L., Chu, H., Sierra, J., Rosen, D. R. and White, K. (1995).
vnd, a gene required for early neurogenesis of Drosophila, encodes a homeodomain
protein. *EMBO J* **14**, 3487–3495.

Lee, T., Lee, A. and Luo, L. (1999). Development of the Drosophila mushroom bodies: sequential
generation of three distinct types of neurons from a neuroblast. *Development* **126**, 4065–
4076.

Lee, T. and Luo, L. (1999). Mosaic analysis with a repressible cell marker for studies of gene
function in neuronal morphogenesis. *Neuron* **22**, 451–461.

Lee, Y. J., Yang, C. P., Miyares, R. L., Huang, Y. F., He, Y., Ren, Q., Chen, H. M., Kawase, T., Ito,
M., Otsuna, H., et al. (2020). Conservation and divergence of related neuronal lineages in
the Drosophila central brain. *Elife* **9**.

Leyssen, M., Ayaz, D., Hebert, S. S., Reeve, S., De Strooper, B. and Hassan, B. A. (2005). Amyloid
precursor protein promotes post-developmental neurite arborization in the Drosophila
brain. *EMBO J* **24**, 2944–2955.

Lin, S. (2023). The making of the Drosophila mushroom body. *Front Physiol* **14**, 1091248.

Liu, T., Zhang, T., Nicolas, M., Boussicault, L., Rice, H., Soldano, A., Claeys, A., Petrova, I., Fradkin,
L., De Strooper, B., et al. (2021). The amyloid precursor protein is a conserved Wnt
receptor. *Elife* **10**.

Luo, L., Tully, T. and White, K. (1992). Human amyloid precursor protein ameliorates behavioral
deficit of flies deleted for Appl gene. *Neuron* **9**, 595–605.

Marquilly, C., Busto, G. U., Leger, B. S., Boulanger, A., Giniger, E., Walker, J. A., Fradkin, L. G.
and Dura, J. M. (2021). Htt is a repressor of Abl activity required for APP-induced axonal
growth. *PLoS Genet* **17**, e1009287.

Nicolas, M. and Hassan, B. A. (2014). Amyloid precursor protein and neural development.
Development **141**, 2543–2548.

Port, F., Chen, H. M., Lee, T. and Bullock, S. L. (2014). Optimized CRISPR/Cas tools for efficient
germline and somatic genome engineering in Drosophila. *Proc Natl Acad Sci U S A* **111**,
E2967–2976.

Rabah, Y., Berwick, J. P., Sagar, N., Pasquer, L., Placais, P. Y. and Preat, T. (2025). Astrocyte-to-
neuron H(2)O(2) signalling supports long-term memory formation in Drosophila and is
impaired in an Alzheimer's disease model. *Nat Metab* **7**, 321–335.
Selkoe, D. J. and Hardy, J. (2016). The amyloid hypothesis of Alzheimer's disease at 25 years.
EMBO Mol Med **8**, 595–608.
Soldano, A. and Hassan, B. A. (2014). Beyond pathology: APP, brain development and Alzheimer's
disease. *Curr Opin Neurobiol* **27**, 61–67.
**Soldano, A., Okray, Z., Janovska, P., Tmejova, K., Reynaud, E., Claeys, A., Yan, J., Atak, Z. K., De
Strooper, B., Dura, J. M., et al.** (2013). The Drosophila Homologue of the Amyloid
Precursor Protein Is a Conserved Modulator of Wnt PCP Signaling. *PLoS biology* **11**,
e1001562.
Stepchenko, A. G., Pankratova, E. V., Doronin, S. A., Gulag, P. V. and Georgieva, S. G. (2011). The
alternative protein isoform NK2B, encoded by the vnd/NK-2 proneural gene, directly
activates transcription and is expressed following the start of cells differentiation. *Nucleic
Acids Res* **39**, 5401–5411.
535

Figure 1

A

Appl　　　　　　　　　　　　　　　*vnd*　　　　　　　　　　5kb

vnd-B　　vnd-A

Appl ^{d}

Appl $^{C1.4}$

Appl $^{C2.1}$

B

Appl　　　　　　　　　　　　*vnd*

3'UTR　　　　3077bp　　　vnd-B　　　　7964bp　　　vnd-A

5'UTR　　　　　　　　　　5'UTR

700bp

Appl ^{d}

Appl $^{C1.4}$

Appl $^{C2.1}$

C WT 0%　　　**D** *Appl* $^{C1.4}$ / Y 66%　　　**E** *Appl* $^{C2.1}$ / Y 9%

Figure 2

Figure 3

A

B *vnd* ^{CΔB} / Y   1%

C *Appl* ^{C1.4} / *vnd* ^{CΔB}   0%

Figure 4

A

β lobe absence (%)

	Appl C1.4 / Y			Appl C1.4 / vnd CΔA		
Dp-Appl⁺	-	+	-	-	+	-
Dp-vnd⁺	-	-	+	-	-	+
n=	147	150	191	198	89	164

** **** **** ****

B *Appl* C1.4 / Y ; Dp-vnd⁺ / + 3%

C *Appl* C1.4 / *vnd* CΔA ; + /+ 64%

D *Appl* C1.4 / *vnd* CΔA ; Dp-vnd⁺ /+ 0%

Figure 5

Figure S1 - related to Figure 1

A

Figure S2 - related to Figure 1

Figure S3 - related to Figure 3

A

vnd

2kb

vnd-B

vnd-A

B

vnd

5'- GGGGGCAGCGAT<mark>AACTGCAGCTCACACCTACA</mark><mark>GGG</mark>ATAGCTTTATCGA -1069bp- TTCTGAACCCAAAATCA<mark>CGAATCAAAAATGAGCCGA</mark><mark>CGG</mark>CGAATGGGCACAG - 3'

5' - GGGGGCAGCGATAACTGCAGCTC   CGGCGAATGGGCACAG - 3'

vnd ^{CΔB}

C

vnd

5'- GATACCCAAAGACA<mark>CCCC</mark><mark>CGAGCATACGACCTGGTACT</mark>TTTCTGCCGTT --1608bp-- TTTAAATTAACCAAGC<mark>GGGCACAGCGCGCAAGTAC</mark><mark>AGG</mark>ACAGGGTATAAT - 3'

5' - GATACCCAAAGACA   GGACAGGGTATAAT - 3'

vnd ^{CΔA}

Figure S4 - related to Figure 5

Figure S5 - related to Figure 4

A — WT

B — *vnd* ^A

C

Clone Genotypes	MB observed	MB clones		WT looking clones	
control hsFLP122	126	36	29%	36	100%
vnd ^{A}	76	20	26%	20	100%
control hsFLP1	56	5	9%	5	100%
vnd ^{6}	34	4	12%	4	100%
vnd $^{\Delta 38}$	90	8	9%	8	100%