1 Tuning apicobasal polarity and junctional recycling in the hemogenic

2 endothelium orchestrates the morphodynamic complexity of

3 emerging pre-hematopoietic stem cells

Léa Torcq,[1,2] Sara Majello,[1] Catherine Vivier,[1] and Anne A. Schmidt [1,6,*]

[1] Institut Pasteur, Université Paris Cité, CNRS UMR3738, Department of

Developmental and Stem Cell Biology, F-75015 Paris, France

[2] Sorbonne Université, Collège doctoral, F-75005 Paris, France

[6] Lead contact

* Correspondence: anne.schmidt@pasteur.fr

Abstract

Hematopoietic stem cells emerge in the embryo from an aortic-derived tissue called the hemogenic endothelium (HE). The HE appears to give birth to cells of different nature and fate but the molecular principles underlying this complexity are largely unknown. Here we show, in the zebrafish embryo, that two cell types emerge from the aortic floor with radically different morphodynamics. With the support of live imaging, we bring evidence suggesting that the mechanics underlying the two emergence types rely, or not, on apicobasal polarity establishment. While the first type is characterized by reinforcement of apicobasal polarity and maintenance of the apical/luminal membrane until release, the second type emerges via a dynamic process reminiscent of trans-endothelial migration. Interfering with Runx1 function suggests that the balance between the two emergence types depends on tuning apicobasal polarity at the level of the HE. In support of this and unexpectedly, we show that the zebrafish Pard3 isoform - Pard3ba - is sensitive to interference with Runx1 activity, in aortic endothelial cells. This supports the idea of a signaling cross talk controlling cell polarity and its associated features, between aortic and hemogenic cells. In addition, using new transgenic fish lines that express Junctional Adhesion Molecules and functional interference, we bring evidence for the essential role of ArhGEF11/PDZ-RhoGEF in controlling the HE-endothelial cell dynamic interface, including cell-cell intercalation, which is ultimately required for emergence completion. Overall, we highlight critical cellular and dynamic events of the endothelial-to-hematopoietic transition that support emergence complexity, with a potential impact cell fate.

Major subject areas: Developmental Biology and Stem cells, Cell Biology.

Keywords: hematopoiesis, stem cell, cell extrusion, hemogenic endothelium, endothelial-to-hematopoietic transition, apico-basal polarity, Pard3, JAMs, zebrafish.

Introduction

Hematopoietic stem cells (HSCs) endowed with full regenerative potential in adult vertebrates are generated during a narrow-time window of few days during embryonic development. These cells, at the root of all blood and immune cells in the body, emerge from intra-embryonic aortic vessels and, more specifically, from a specialized type of vascular tissue called the hemogenic endothelium (HE, (Wu and Hirschi 2021)). A series of seminal studies have evidenced the autonomous production of repopulating HSCs within an intraembryonic region called the Aorta-Gonad-Mesonephros (AGM), before their appearance in other hematopoietic organs (A. L. Medvinsky et al. 1993; Müller et al. 1994; Garcia-Porrero, Godin, and Dieterlen-Lièvre 1995; A. Medvinsky and Dzierzak 1996; Cumano, Dieterlen-Lievre, and Godin 1996). Thereafter, the endothelial origin of HSCs was evidenced (Jaffredo et al. 1998; de Bruijn et al. 2002; T. E. North et al. 2002). The direct visualization, in real-time, of the emergence of precursors of hematopoietic stem and progenitor cells (HSPCs) from the dorsal aorta, termed the Endothelial-to-Hematopoietic Transition (EHT), was finally achieved *in vitro* (Eilken, Nishikawa, and Schroeder 2009), *ex-vivo* from mouse sections (Boisset et al. 2010), and *in vivo* in the zebrafish embryo (Bertrand et al. 2010; Kissa and Herbomel 2010), using live microscopy. This opened the way to a more detailed analysis of the characteristics of the HE at the transcriptional level, revealing its transient nature and its early hematopoietic commitment (Swiers et al. 2013).

The HE is characterized by inherent heterogeneity and is not only contributing to the formation of precursors to long-term HSCs (pre-HSCs) but also to more restricted progenitors (Hadland and Yoshimoto 2018). These progenitors can be born from HE sub-types that are found in extra-embryonic source such as the yolk-sac – in which case it gives rise to erythro-myeloid progenitors that will sustain erythro-myelopoiesis until birth (Frame et al. 2016) -, or both from extra and intra-embryonic sources (the yolk sac and the dorsal aorta) in which cases

progenitors are biased, such as for example in the mouse, toward T- and B- innate-type

lymphoid progenitors (Yoshimoto et al. 2011; 2012) or toward less restricted, multipotent

progenitors (Hadland and Yoshimoto 2018; Dignum et al. 2021). Importantly, while some of

the non-HSC derived progenitors born during embryonic life only support the functions of the

immune system during embryonic/ fetal life, others can persist in the adult to exert tissue-

resident functions, as has been mainly described in the mouse (Ghosn et al. 2019). However,

studies in human embryos support the idea that developmental hematopoiesis is highly similar

in mice and humans (Ivanovs et al. 2011; 2017).

A key question regarding the issue of the capacity of the HE to give birth to cells endowed with

different hematopoietic potential is whether this comes from HE cells of distinct origin (in which

case the HE would be a mosaic of precursors with distinct potential) or if this results from

extrinsic environmental cues that impose variability in the constantly evolving developmental

context (for a discussion, see (Barone et al. 2022)).

To address this type of question, high-resolution experimental settings need to be developed,

if possible at the single-cell resolution. Along this line, recent developments of transcriptomics,

including single cell RNAseq, spatially resolved in situ transcriptomic approaches, and in situ

barcoding, have been invaluable (Weijts, Yvernogeau, and Robin 2021). Among those high-

resolution approaches, cell fate mapping and lineage tracing approaches that include imaging

technologies have been very instrumental, particularly when performed with the zebrafish

model that reproduces many aspects of developmental hematopoiesis in higher vertebrates

(Orkin and Zon 2008). Recently, work performed in the zebrafish embryo has brought evidence

for heterogeneity of hematopoietic stem cell precursors being born from the HE, in the ventral

floor of the dorsal aorta and independently from HSCs, including a wave of transient T-

lymphocytes (using temporally-spatially resolved fate-mapping, (Tian et al. 2017)) and lympho-

myeloid biased progenitors born from a myeloid-lymphoid biased Spi2+ HE that appears to co-

exist with an erythroid-biased HE (using single-cell RNA-sequencing, (Xia et al. 2023)).

While light starts to be shed on the molecular and signaling cues that appear to regulate HE

sub-specification and a continuum from arterial endothelium, HE maturation, and subsequent

heterogeneity in HSPCs (Zhu et al. 2020), the essential molecular and cell biological properties

that support HE functional plasticity remain to be determined.

Here, using the zebrafish embryo as a model, we complement our previous work describing

essential molecular and mechanistic features of EHT cell emergence (Lancino et al. 2018).

With the support of high-resolution live imaging and the generation of new transgenic fish lines

that express a functional marker of cell polarity (Podocalyxin, a sialomucin of the CD34 family

(Nielsen and McNagny 2008)), we show that the HE is giving birth to two cell types with distinct

cell polarity status; these cells also emerge with radically different morphodynamic

characteristics, which raises the intriguing possibility of an incidence of EHT emergence

complexity on downstream fate, after release from the aortic wall.

To substantiate the significance of apico-basal polarity control in the EHT, we investigate on

the potential involvement of Pard3 proteins. These proteins, recruited by transmembrane

receptors via their PSD-95/Dlg/ZO-1 (PDZ) domains (Buckley and St Johnston 2022), are at

the root of apicobasal polarity initiation and are essential for the maintenance of apical

membrane functional properties (Román-Fernández and Bryant 2016). Among the four Pard3

isoforms expressed in the zebrafish (encoded by four different genes and non-including

splicing variants), we describe the specific expression of the zebrafish Pard3ba isoform in the

aorta, during the EHT time-window. We show that the expression of Pard3ba is sensitive to

interference with the transcription factor Runx1 whose function is essential for EHT completion

(Kissa and Herbomel 2010) and for regulating HSC number (Adamo et al. 2009; T. E. North et

al. 2009). In link with apico-basal polarity and to address its incidence on the biomechanics of

the emergence of the two EHT cell types that we describe, we generated new fish lines that

express Junctional Adhesion Molecules (JAMs) fused with eGFP (eGFP-JAMs). The JAMs

belong to tight junction complexes (Garrido-Urbani, Bradfield, and Imhof 2014) and recruit

Pard3 via direct interaction with their carboxy-terminus (Ebnet et al. 2003), thereby allowing

initiation of apico-basal polarity. With our eGFP-JAMs fish lines, we investigate on the function

of ArhGEF11/PDZ-RhoGEF that was shown to be involved in disrupting junctional integrity in

the context of the epithelial-to-mesenchymal transition as well as in the growth and migration

of invasive cancer cells (more specifically one splicing variant of its c-terminus conserved in

mammalian species, Itoh et al. 2017; Lee et al. 2018). We reveal the function of

ArhGEF11/PDZ-RhoGEF in the dynamic interplay between HE/EHT cells and their endothelial

neighbors, particularly in cell-cell intercalation which is essential for EHT completion and the

proper sealing of the aortic floor.

Results

Apicobasal polarity determines emergence types

Our previous work describing the morphodynamic evolution of cells emerging from the aortic floor through the EHT (hereafter designated as "EHT cells"), in the zebrafish embryo, revealed the unusual feature of a cell extruding from the plane of a tissue while maintaining its luminal/apical membrane until the very end of the release process, thus contributing to its peculiar crescent-shaped morphology (Kissa and Herbomel 2010; Lancino et al. 2018), and see the cartoons in **Figure 1A**). However, to our knowledge, the polarity status of EHT cells has not been investigated so far and the maintenance of a *bona fide* apical domain has never been proven (with the luminal membrane enriched in apically targeted proteins and physically delimited by apical polarity factors and tight junction complexes (Rodriguez-Boulan, Müsch, and Le Bivic 2004; Buckley and St Johnston 2022)). Importantly, the fate of this apical-luminal membrane, after the release, may lead to cells potentially endowed with specific functional features. For example, this membrane surface may be directly and fully exposed to the extracellular space or released in the cytoplasm of EHT cells for recycling and/or degradation, after emergence completion (for examples of different scenarios, see **Figure 1A,** and the legend for more details). Overall, this could lead to precursors of hematopoietic stem cells that, ultimately, may be differentially fated.

To address the polarity status of EHT cells, we raised transgenic fish lines that express endogenous Podocalyxin (Podxl2, (Herwig et al. 2011)). Podocalyxin was shown to take part in the formation of the preapical domain during polarization and in the regulation of its oriented organization, in tissue culture (Meder et al. 2005; Bryant et al. 2014). Its contribution to lumenization *in vivo*, in the mouse aorta, has been described and it involves negative charge repulsion induced by their glycosyl residues (Strilić et al. 2009; 2010).

We first attempted to express transiently, in the vascular system, the full-length Podxl2 zebrafish protein fused to eGFP at its extreme N-terminus. We failed to detect the fusion

protein at any time point relevant for observing easily the EHT process (a time window ranging

from 48 to 72 hpf (hours post-fertilization)); therefore, we designed a N-ter truncated form

deleted of the mucin domain and that retains amino-acids 341-587 fused to eGFP at its N-

terminus (**Figure 1B** *and see* **Methods**). Transient transgenesis revealed that this truncated

version is detected and is targeted to the luminal membranes of EHT cells. We then raised 2

transgenic (*Tg*) fish lines that express the N-ter truncated form of Podxl2 fused to either eGFP

(thereafter abbreviated eGFP-podxl2) and under the control of the Kdrl:Gal4 driver

Tg(Kdrl:Gal4;UAS:RFP; 4xNR:eGFP-podxl2), or to mKate2 and under the control of the Kdrl

promoter *Tg(Kdrl:mKate2-podxl2)*. We observed that eGFP-podxl2 is enriched at the luminal

side of crescent-shaped EHT undergoing cells (see **Figure 1C, Figure 1 - figure supplement**

1A and **Figure 1 – video 1** (z-stack at t = 0) and their legends for the details of the

luminal/apical membrane evolution through time). Thereafter, these cells will be referred to as

EHT pol+ cells.

We also followed the cell after emergence and observed the evolution of the luminal/apical

membrane appearing as internal pseudo-vacuoles. We illustrate the reduction of their volume

via membrane retrieval and, ultimately, their remanence as an intracellular membrane

compartment which we define as a post-EHT signature (**Figure 1- figure supplement 1A** and

Figure 1- video 2 (z-stack at t = 80 min)); we propose a model for the intracellular evolution

of the luminal/apical membrane (**Figure 1- figure supplement 1B**) which, unfortunately,

cannot be traced after 2-3 hrs post-emergence because of the apparent short half-life of eGFP-

podxl2 and of the drop in activity of the Kdrl promoter. Of notice, the pseudo-vacuoles are

reminiscent of the cystic formations observed in EHT cells in the mouse and visualized by

electron-microscopy (T. North et al. 1999; Marshall and Thrasher 2001) and also of the

vacuolar structures recently described in EHT cells in avian embryos (Sato et al. 2023; for

more details, see **Discussion**).

While imaging the EHT using the Podxl2 expressing lines that clearly delimitate the luminal

membrane, we unambiguously identified a second type of emergence. This second cell type

is primarily characterized by a round-to-ovoid morphology (cells never bend as crescent-

shaped EHT pol+ cells, see **Figure 1D** and **Figure 1 – video 3** for a time-lapse sequence).

Importantly, these cells do not show any enrichment of eGFP-podxl2 at the luminal membrane

and will be referred to as EHT pol- cells. EHT pol- cells were observed in all other *Tg* fish lines

that we are routinely imaging, including the *Tg(Kdrl:Gal4;UAS:RFP)* parental line that was used

for transgenesis, thus excluding the possibility that these cells result from an artefact due to

the expression of a deleted form of Podxl2 and/or to its overexpression (see also for example

our new *Tg(kdrl:eGFP-Jam3b)* fish line **Figure 5C** and **Figure 6B**).

Finally, we have estimated that the ratio between EHT pol+ and EHT pol- cells is of

approximately 2/1, irrespective of the imaging time window and of the localization of

emergence along the aortic antero-posterior axis (starting from the most anterior part of the

AGM (at the limit between the balled and the elongated yolk) down to the caudal part of the

aorta facing the CHT). We observed that both EHT pol+ and EHT pol- cells divide during the

emergence and remain with their respective morphological characteristics after completing

abscission **(Figure 1 – figure supplement 2);** hence they appear as pairs of cells that exit the

aortic wall sequentially (as shown **Figure 1C, D**). We also observed that both EHT pol+ and

EHT pol- cells express reporters driven by the hematopoietic marker CD41 **(Figure 1 – figure**

supplement 3), which indicates that they are both endowed with hematopoietic potential.

Altogether, our results show that hematopoietic precursor cells emerging from the aortic floor

do so with heterogeneity in their morphodynamic characteristics. They also suggest that the

control of apico-basal polarity may be at the root of these specific emergence types.

The immature HE is not polarized

EHT pol+ and EHT pol- cells appear to emerge from the hemogenic endothelium (HE), the

latter constituting the virtually exclusive cell population of the aortic floor just prior to the

initiation of the EHT time-window (around 28 hpf, see (Zhao et al. 2022)). In this context, we

addressed the polarity status of HE cells and its evolution throughout the EHT (**Figure 2**).

Surprisingly, confocal microscopy using our *eGFP-podxl2* and *mKate2-podxl2* expressing fish

lines revealed that HE cells do not appear to be polarized, based on the absence of Podxl2

enrichment at luminal membranes, at the initiation of the EHT time-window and later (at

approximately 28 - 30 hpf, **Figure 2 – figure supplement 1,** see also at 35 hpf **Figure 2A**, top

left panel, green arrows and **Figure 2 - figure supplement 2** and at 48 – 55 hpf **Figure 3 –**

figure supplement 2A). Interestingly, the cytoplasm of characteristic elongated HE cells

located on the aortic floor is filled with more-or-less large membrane vesicles that carry eGFP-

podxl2 (the largest vesicles reaching approximately 30μm in diameter). This suggests that HE

cells contain a reservoir of eGFP-podxl2 membranes that may be subjected to exocytose; as

such, HE cells may be comparable to endothelial cells organizing a vascular lumen and that

have been proposed to exocytose large intracellular macropinocytic-like vacuoles when

cultured in 3D extracellular matrices (Bayless and Davis 2002; Davis, Bayless, and Mavila

2002) or, *in vivo*, in the zebrafish model (Kamei et al. 2006; Lagendijk, Yap, and Hogan 2014).

This finding is unexpected since HE cells are assumed to possess aortic cell characteristics

(i.e exhibit an apicobasal polarity) as they are supposedly integrated in the aortic wall

contemporarily to aortic precursors (Jin et al. 2005) and may have been taking part in the

lumenization of the ancestral vascular cord, a process that takes place around 18-20 hpf.

Consequently, loss of apicobasal polarity features of HE cells at 28-30 hpf may be part of the

programme that initiates the EHT process.

Although technically difficult for long hours (because of important variations in the volume of

the balled yolk that trigger drifting of embryos), we have been able to follow over time non-

polarized HE cells and to visualize the evolution of their vesicular content, starting at the

initiation of the EHT time-window (around 35 hpf, see **Figure 2** and **Figure 2 – figure**

supplement 2). Interestingly, we could follow a dividing HE cell for which the vesicular content

labelled with eGFP-podxl2 appeared to partition unequally between daughter cells (**Figure 2A,**

t = 30 min, HE cell 1' inherits the largest macropinocytic-like vacuole (green asterisk) and

emerges unambiguously as an EHT pol+ cell (t = 360 min, EHT cell 1'), with eGPF-podxl2

enriched at the apical/luminal membrane (surrounding the green asterisk) and virtually

undetected at the basal membrane (magenta arrow)). This suggests that asymmetric

inheritance of cytosolic vesicles containing apical proteins may contribute, presumably after

delivery to the luminal membrane, to specify the apical membrane of EHT pol+ cells.

Altogether, these results support the idea that the HE, at the initiation of the EHT time-window,

is not polarized. Subsequently, HE cells establish – or not – apical and basal membrane

domains, which characterizes EHT pol+ or EHT pol- cell types, respectively (see our

hypothetical model **Figure 2B**). In the case of EHT pol+ cells and while emergence is

proceeding, apicobasal polarity is maintained (if not reinforced) until the release.

Interfering with Runx1 function alters HE maturation, the EHT progression, and the

balance between EHT cell types

To provide functional support to our findings and hypotheses on apico-basal polarity control in

the gradual maturation of the HE throughout the EHT time-window, we thought of interfering

with the activity of the transcription factor Runx1 whose expression is sensitive to fluid sheer

stress (Adamo et al. 2009).

To interfere with Runx1 function, we generated a *Tg* fish line that expresses a truncated form

of Runx1 (dt-runx1; fish line *Tg(Kdrl:Gal4;UAS:RFP;4xNR:dt-runx1-eGFP)* thereafter

abbreviated *Tg(dt-runx1)*) deleted from its transactivation domain and carboxy-terminus but

retaining its DNA-binding domain (the Runt domain (aa 55-183, (Kataoka et al. 2000; Burns et

al. 2002; Kalev-Zylinska et al. 2002), see **Figure 3C)**. Importantly, in this *Tg* fish line, dt-runx1

expression is restricted to the vascular system, hence excluding expression in the brain region

that express Runx1 endogenously (see the aforementioned articles) and preventing biases

owing to potential interference with neuronal functions. In addition, owing to the expression of

eGFP concomitantly to dt-runx1 (the C-terminal eGFP is cleaved from dt-runx1 via a T2a site

for endopeptidase, see cartoons at the top of panel C), this fish line allows for the easy selection of embryos for imaging and for phenotypic analysis. In preliminary experiments aimed at addressing the localization of dt-runx1 as well as its stability, we expressed it transiently and measured its proper targeting to the nucleus (the construct also contains a double HA (2xHA) epitope for immuno-detection, see *Figure 3C).*

To characterize further the *Tg(dt-runx1)* fish line, we addressed the incidence of dt-Runx1 expression on HE and EHT cell populations during the EHT time-window as well as later on, at 5 days post-fertilization (5dpf) in the thymus, as a readout of hematopoiesis efficiency. We made the following observations:

Firstly, using 48 - 55 hpf *Tg(dt-runx1)* embryos and live imaging we observed, in comparison to controls, a significant accumulation of HE cells (see *Figure 3A*, top and bottom panels (he, arrowheads), see also the quantifications *Figure 3B*, top graph, median value of 17 cells for controls and 25 cells for mutants). In the mutant condition, the bulk of HE cells also include uncharacterized EHT cells (*Figure 3A*, middle panel, blue arrows); hypothetically, these may be EHT cells at an early phase of their emergence, including cells that may have evolved as EHT pol- cells in unperturbed conditions, owing to the absence of luminal membrane invagination. We also observed the unusual tendency for the accumulation of EHT pol+ cells (*Figure 3A*, green and pink arrows), with a maximum value of 3 cells per embryo for controls that reached the unusual - never observed in control conditions -, values of 11, 9 and 8 cells for 3 mutants out of 7 (*Figure 3B*, middle graph). In addition to the increase in the proportion of EHT pol+ cells in the ventral side of the aorta (*Figure 3B*, right panel, with a value of 4% for controls and reaching 7% for mutants, consistently with the increase in the absolute count per embryo), we also observe an increase in the proportion of EHT pol+ cells in lateral sides of the dorsal aorta (*Figure 3A*, pink arrows and *Figure 3B*, right graph, with a value of 4% for controls and reaching 9% for mutants). Finally, we did not observe any significant modification in the number of EHT pol- cells (same median values for both controls and mutants, the values

oscillating between 0 and 1 cell per embryo, with only one example of an embryo exhibiting 2 apparent EHT pol- cells (**Figure 3B**, bottom graph)).

Secondly, using 48 - 55 hpf *Tg(dt-runx1)* embryos and controls *Tg(Kdrl:eGFP)* we confirm, by whole mount in situ hybridization performed with an RNAscope probe specific for myb (**Figure 3 – figure supplement 1**), that the cells that accumulate in the aortic floor of dt-Runx1 mutants are indeed of hematopoietic nature (see the cells with RNAscope spots in magenta or marked by blue arrows, **Figure 3 – figure supplement 1A,** zoomed areas). Unfortunately, chemical fixation impaired the maintenance of EHT pol+ and EHT pol- specific cell shape, preventing us from classifying emerging cell populations. However, cell counts after segmentation using the Imaris software confirmed, based on cell localization, the specific increase of hemogenic/EHT cells in aortic floors of mutants (median values of 34 and 16.5 for mutants and controls, respectively, **Figure 3 – figure supplement 1B** (right panel) while no difference was measured for endothelial roof cells between mutants and controls (left panel). Quantification of myb positive cells confirmed the increase of hemogenic cells in the aortic floors of mutants in comparison to controls, albeit with slightly smaller numbers (median values of 20.5 and 8.5 in mutants and controls, respectively (**Figure 3 – figure supplement 1C**, middle panel)), most probably because myb becomes detectable only when the EHT process has initiated. In comparison to floor cells, very few roof cells express myb (median values of 2.0 and 1.5 in mutants and controls, respectively (**Figure 3 – figure supplement 1C**, left panel)) consistently with the idea that the aortic roof is not hemogenic (Zhao et al. 2022)). Importantly, the increase of pre-hematopoietic cells in the aortic floor in the mutants correlates with a decrease in hematopoietic cells lodging in the sub-aortic region, reinforcing the idea of impairment in EHT cell emergence (**Figure 3 – figure supplement 1C**, right panel). Finally, quantification of the number of myb RNAscope spots per segmented hemogenic cells highlights the important increase in expression by hemogenic cells in the floor of dt-Runx1 mutants, in comparison to controls (with median values of 4 and 0, respectively *(**Figure 3 – figure supplement 1D**, right panel). In comparison, virtually none of the roof cells contain more than 2 spots (**Figure 3 –**

figure supplement 1D, with 2/162 and 1/220 cells for the mutants and controls, respectively,

with median values of 0 for both conditions). Altogether, these last results confirm that pre-

hematopoietic cells accumulate in the aortic floor of dt-Runx1 mutants; the very substantial

increase in the detection of myb mRNAs suggests an increase in the residential time in the

aortic floor, which reinforces the idea that these cells are impaired in their release from the

aortic floor.

Thirdly, between 30 and 55 hpf, using live confocal microscopy and contrarily to what we

observed with our fish lines expressing either eGFP-podxl2 (*Figure 2 and Figure 2 – figure*

supplement 1 and 2) or mKate2-podxl2 (*Figure 3 – figure supplement 2A*) in which HE cells

appear to be non-polarized, embryos obtained from outcrossing *Tg(dt-runx1) and*

Tg(Kdrl:mKate2-podxl2) fishes appear to contain polarized HE cells, based on enrichment of

the polarity marker at luminal membranes (*Figure 3 - figure supplement 2B)*.

Fourthly and occasionally, in live embryos, we observe - for EHT cells exhibiting invagination

of the luminal membrane -, scattered cytosolic and sub-plasmalemmal pools of Podxl2-

containing membranes and, consistently, the apparent decrease of Podxl2 enrichment at the

apical/luminal membrane (see *Figure 3* - *figure supplement 2C* and compare with *Figure*

1C). In addition and occasionally as well, we observe the reversion of apparent EHT pol+ cells

into apparent EHT pol- cells (data not shown). These last two observations suggest that

perturbing the control of apicobasal polarity in the HE, in the context of the dt-Runx1 mutant,

alters emergence morphodynamics, hypothetically and possibly more specifically in the case

of EHT pol- cells whose biomechanical features appear to require turning down the

establishment of apical and basal membrane domains.

Fifthly, hematopoiesis is affected far downstream of emergence, as attested by the significant

increase, at 5 dpf, in the number of hematopoietic cells in the thymus of dt-Runx1 mutants, in

comparison to control siblings (see *Figure 3 - figure supplement 3A* for images after cell

segmentation with the Imaris software accompanied by *Figure 3 - video 1* and *Figure 3 -*

figure supplement 3B for quantifications showing significant increase in thymus volume (left

panel) owed to the increase in cell number (middle panel) rather than in cell volume (right

panel)).

Altogether, the results obtained upon expression of dt-runx1 show impairment of

hematopoiesis and suggest that, for both EHT cell types, the progression throughout EHT is

perturbed, and so until the release. The accumulation of morphologically characterized EHT

pol+ like cells that we observe may result from the sustained apicobasal polarity of the HE at

early and later time points. Conversely, characterized EHT pol- cells did not accumulate,

although we observed cells of uncharacterized morphology that could correspond to EHT pol-

cells that did not evolve properly. In addition, our results suggest that precursors of HE cells

are polarized, as is expected to be the case for non-hemogenic aortic cells, and that Runx1 is

involved in controlling the molecular events that are tuning apico-basal polarity, starting at the

initiation of the EHT time-window.

Interfering with Runx1 activity unravels its function in the control of Pard3ba expression

and highlights heterogeneous spatial distribution of Pard3ba mRNAs along the aortic

axis

We then explored the potential involvement of proteins of the Pard3 family during the EHT. We

anticipated that interfering directly with Pard3 proteins would hamper the development of the

aorta (in addition to other functions of polarized tissues that are essential for embryonic

development) and rather searched for correlative evidence for the differential expression of

Pard3 isoforms in the hemogenic endothelium, EHT cells and the aortic endothelium, both in

wild type and dt-Runx1 expressing embryos. To achieve this task, we analyzed the expression

of mRNAs encoding for the four Pard3 isoforms expressed in the zebrafish (***Figure 4A***) by

combining qRT-PCR on FACS-sorted endothelial cells and whole-mount *in situ* hybridization

at the cellular resolution, using RNAscope. All four Pard3 isoforms are composed of 3 PDZ

domains and 2 conserved regions upstream and downstream of these domains (CR regions,

Figure 4A).

We first sought to measure expression of these different isoforms in the vascular system of 48

hpf embryos and, more restrictively, in the trunk region. To do so, we dissociated and FACS-

sorted cells of either whole embryos, or dissected trunks, using the *Tg(Kdrl:Gal4; UAS:RFP)*

fish line in which soluble RFP is expressed in endothelial tissues (**Figure 4 – figure**

supplement 1A). RFP+ vascular cells were found the be slightly enriched in the trunk region,

in comparison to whole embryos (**Figure 4 – figure supplement 1A,** see the table with 2.10

versus 0.95 % of RFP+ cells in the trunk and the whole embryo, respectively). Using qRT-PCR

on the FACS-sorted cells, we found that the hematopoietic markers myb and runx1 were

heavily detected in RFP+ cell extracts in comparison to RFP- cells (**Figure 4 – figure**

supplement 1B); this suggests that part of the isolated RFP+ cells contain hemogenic and

hematopoietic cells (at 48 hpf, it is expected that some newly born hematopoietic cells are

retained in the sub-aortic space). When we measured the expression levels of the four Pard3

isoforms, we observed, in comparison to control RFP- cells, the enrichment of Pard3aa and

Pard3ba in extracts from RFP+ cells isolated from whole embryos, and the opposite for

Pard3ab and Pard3bb (**Figure 4 – figure supplement 1C**). These differences were conserved

with extracts from RFP+ cells isolated from trunk regions, except for Pard3ba for which

important variations in mRNA quantities were obtained, blurring conclusions (we shall come

back to this variability beneath). These results suggest that, at least based on the vascular

system of whole embryos, Pard3aa and Pard3ba exert functions in endothelial cells, with the

possibility of having more specific functions in aortic and HE cells. To discriminate between

cell types expressing Pard3aa and Pard3ba isoforms (and in particular vascular cell types), we

used RNAscope on whole mount *Tg(Kdrl:eGFP)* embryos, at 48 hpf. Confocal images focused

on trunk regions show that Pard3aa is strongly expressed in cells of the spinal cord (potentially

neuronal cells and/or radial glia), as well as in the notochord (**Figure 4 – figure supplement**

2A). Interestingly, Pard3ba mRNAs are expressed in more restricted areas, in majority in the

spinal cord, albeit to a much lesser extent than for Pard3aa as well as in the region of the

pronephric duct and the gut epithelium (**Figure 4 – figure supplement 2B**). By curiosity and

because it appears to be the most highly expressed in the whole embryo (and hence may be

ubiquitously expressed, including in the trunk region and in accordance to the Daniocell

resource at https://daniocell.nichd.nih.gov/, Farrell et al. 2018; Sur et al. 2023), we also

investigated on the expression of Pard3ab mRNAs (**Figure 4 – figure supplement 2C**). We

confirmed its relatively ubiquitous expression in comparison to Pard3aa and Pard3ba, with a

massive expression in the spinal cord. To visualize more precisely expression levels in the

vascular system in the trunk (more specifically in the aorta and the underlying vein), we

zoomed in this region (**Figure 4 – figure supplement 2D**) and observed a homogeneous

localization of Pard3aa and Pard3ab mRNAs as well as, strikingly in the case of Pard3ba, a

quite heterogeneous localization of mRNAs unequally concentrated along the aortic axis

(**Figure 4 – figure supplement 2D,** middle panel and see also **Figure 4C**, top panel; these

will be referred thereon as Pard3ba mRNA enriched regions (MERs)). Of notice, in the whole

aorta, we observed more RNAscope signals for Pard3ab in comparison to Pard3aa and

Pard3ba, including in some HE/EHT cells (**Figure 4 – figure supplement 2D**, with positive

HE/EHT cells indicated by blue arrows). This indicates that the Pard3ab isoform, consistently

with the absence of variation upon dt-Runx1 expression (see beneath), is constitutively

expressed in some HE/EHT cells (this may be also the case for the Pard3aa isoform but this

should be taken with caution since its mRNAs are more in the limit of detection). We attempted

to detect Pard3 isoforms expressed in the vascular system, in the trunk region of the embryo,

with available antibodies raised against other species and failed, possibly because of inter-

species recognition limitation.

We then analyzed Pard3 mRNA levels in the context of dt-Runx1 expression. As we did for

Tg(Kdrl:Gal4;UAS:RFP) embryos, we isolated by FACS GFP+ and double positive

GFP+/RFP+ cells dissociated from dissected trunks of *Tg(dt-Runx1)* 48 hpf embryos. For

controls, we also isolated RFP+ cells from dissected *Tg(Kdrl:Gal4;UAS:RFP)* embryos (see

Methods and **Figure 4 – figure supplement 3A**). qRT-PCR on isolated eGFP+ cells that

express dt-Runx1 revealed an increase of approximately 50% of the myb and runx1 mRNAs,

in comparison to RFP+ control cells (**_Figure 4B,_** left panel), which is consistent with the

accumulation of hemogenic/EHT cells that we observe in **_Figure 3_** and in **_Figure 3 – figure_**

supplement 1. Interestingly, we obtained very little to no variation in the expression levels of

Pard3aa, ab and bb mRNAs in dt-Runx1 expressing eGFP+ cells in comparison to control

RFP+ cells but, again, a high variability in the case of Pard3ba mRNAs (**_Figure 4B,_** right panel).

Using RNAscope, the non-homogeneous localization of Pard3ba mRNAs observed in controls,

along the aortic axis, was not only conserved but became even more obvious to the eye for dt-

Runx1 mutants, owing to the increase in the density of RNAscope spots (**_Figure 4C,_** compare

top and bottom panels). Confocal images on several controls and dt-Runx1 mutants highlight

the heterogeneity in Pard3ba mRNA detection as attested by the variability in the localization

and number of RNAscope spots (compare in particular the 5 mutants shown in **_Figure 4 –_**

figure supplement 3B, right panels). Quantification of RNAscope spots per aortic segment

and per hemogenic/EHT cell using Imaris revealed an increase of almost 300% for the former

when comparing controls and mutants (median values of 65.00 and 167.50 for controls and

dt-Runx1 mutants, respectively, **_Figure 4D_**) and that virtually no Pard3ba mRNA was detected

in the latter (median values of 0.00 spots per hemogenic/EHT cell for both controls and dt-

Runx1 mutants, **_Figure 4E_**). A precise representation showing the hemogenic/EHT cell count

with RNAscope spots ranging from 0 to 4 reveals, for mutants in comparison to controls, a

significant increase in the number of hemogenic/EHT cells not expressing Pard3ba (**_Figure 4_**

– figure supplement 3C, first panel from the left), as well as a slight tendency for an increase

in the number of cells with 1 to 4 Pard3ba spots (**_Figure 4 – figure supplement 3C,_** second

to fifth panels), with more significance for the population with 2 spots (middle panel). Overall,

in either control or mutant conditions, we detected the presence of RNAscope spots in very

few hemogenic/EHT cells (15/115 for controls and 21/224 for dt-Runx1 mutants, with a

maximum for 2 spots for controls and 4 spots for dt-Runx1 mutants, see **_Figure 4E_** and **_Figure_**

4 – figure supplement 3C).

Again, we obtained a large variability in the number of Pard3ba RNAscope spots per aortic

segment (the number of spots per aortic segment ranged from 10 to 94 for controls and from

53 to 366 for the mutants, see **Figure 4D**); this mirrors what was obtained by qRT-PCR,

particularly for dt-Runx1 mutants (**Figure 4B,** right panel)*. This is not the case for Pard3aa and

Pard3ab (see RNAscope images and quantifications **Figure 4 – figure supplement 4**), which

is also consistent with qRT-PCR results (**Figure 4B,** right panel)*. Importantly, expression of dt-

Runx1 neither influenced the expression of Pard3aa, nor of Pard3ab (**Figure 4 – figure**

supplement 4).

Finally and importantly, in wild-type embryos, the Pard3ba mRNAs are preferentially localized

in the direct vicinity of hemogenic/EHT cells (**Figure 4F, G**). Indeed, more than 50% of Pard3ba

spots are localized within 5 to 10 μm of hemogenic/EHT cells (**Figure 4G** middle panel), and

more than 75% of spots are localized within a 5 to 20μm distance (**Figure 4G** right panel).

Interestingly, the number of hemogenic/EHT cells does not correlate with the absolute number

of spots (**Figure 4F,** left panel, R = 0.34), but rather it strongly correlates with the spatial

densification of spots around hemogenic/EHT cells: the shorter the distance of the majority of

spots are of hemogenic/EHT cells (**Figure 4F,** middle and right panel, representing 50% and

75% of spots respectively), the higher the hemogenic/EHT cell count (R = -0.84 and R = -0.83

respectively). This shows that AECs expressing Pard3ba mRNAs are mostly in direct contact

with cells committed to undergo EHT and suggests a signalling crosstalk occurring between

them. This densification of spots in MERs is reinforced upon expression of dt-Runx1 (**Figure**

4F, left panel*): we found that virtually all Pard3ba spots are localized within a 10-15μm distance

from hemogenic/EHT cells.

Altogether, these results unveil the Runx1-dependent control of the Pard3ba isoform

specifically, during the EHT. In the aorta, AECs appear to be the main cell type that expresses

this Pard3 isoform and, unexpectedly, expression in these cells is sensitive to interference with

our dt-Runx1 mutant. One plausible explanation is that AECs respond to Runx1-mediated

signalling taking place in HE/EHT cells (see **Discussion** for other possibilities and further

details). This suggests that a specific signalling axis is taking place between these different

cell types that would control the expression of Pard3ba mRNAs and, consequently,

downstream apico-basal polarity associated features carried out by the Pard3ba protein

isoform (example: the recycling mode of junctional complexes between EHT pol+ cells and

endothelial neighbours, in comparison to EHT pol- cells, see beneath and **_Discussion_**). This

potential intercellular signalling is supported by the proximity of Pard3ba mRNAs expressed in

AECs to HE/EHT cells in MERs. These results also strengthen the importance of the interplay

between HE/EHT cells and their direct endothelial neighbours (this interplay has been

highlighted in Lancino et al. 2018; and see also beneath).

Junctional recycling between EHT cell types and endothelial neighbours is differentially

controlled

To investigate the functional links between apicobasal polarity and the peculiarities of EHT

pol+ and EHT pol- emergence processes, we sought to follow the dynamics of Junctional

Adhesion Molecules (JAMs) that belong to tight junction complexes (Garrido-Urbani, Bradfield,

and Imhof 2014). During apicobasal polarity establishment in epithelial and endothelial tissues,

these molecules recruit the Pard3/aPKC complex, via a PDZ-binding peptide located in their

extreme carboxy-terminus (**_Figure 5A,_** left cartoons and (Itoh et al. 2001; Ebnet et al. 2001;

2018)). In addition, JAMs are expressed in the vascular system and, in relation to this work,

JAM3b is expressed in the aorta of the zebrafish embryo and promotes endothelial

specification and hematopoietic development (Kobayashi et al. 2020).

We envisaged a scenario whereby EHT pol+ cells, whose longitudinal interface with

endothelial neighbours shrinks along the X axis of the X, Y 2D-plane, may have less mobile

junctional pools than EHT pol- cells whose entire junctional interface moves along the X, Y, Z

3D-axes (**_Figure 5 - figure supplement 1)_**. In the case of EHT pol- cells, the consumption of

the junctional interface with adjoining endothelial cells appears to result from the converging

migration of endothelial neighbours crawling over the luminal membrane, based on

interpretation of our time-lapse sequences (**Figure 1- video 3**). In this context, we favoured

the analysis of junctional pools localized at antero-posterior sites of emerging cells as we have

shown that they are enriched with tight junction components (Lancino et al., 2018). In addition

and in the case of EHT pol+ cells, it is conceivable that these adhesion pools – spatially

restricted owing to apicobasal polarity –, contribute to anchoring the emerging cell in the 2D-

plane (**Figure 5 - figure supplement 1,** top panel*).*

To achieve our goal, we designed 2 constructs in which eGFP is introduced in the extracellular

domains of the two JAM2a and JAM3b molecules (**Figure 5A,** right cartoons). To investigate

their localization and proper targeting at junctional interfaces, these constructs were expressed

transiently and ubiquitously, using the *Hsp70* heat shock promoter. We observed that the 2

fusion proteins are efficiently targeted at cellular contacts (**Figure 5 - figure supplement 2B,**

C and **F**, orange and white arrows) as well as at the apical side of polarized epithelia such as

for example cells of the ependymal canal (**Figure 5 - figure supplement 2A, E**, white arrows).

They were also observed at protrusions of epithelial cells and at T-tubules of muscle fibres (for

JAM2a and JAM3b, respectively, see **Figure 5 - figure supplement 2D**, **G**).   We then

established *Tg* fish lines expressing eGFP-Jam2a and eGFP-Jam3b under the control of the

vascular kdrl promoter (*Tg(kdrl:eGFP-Jam2a)* and *Tg(kdrl:eGFP-Jam3b))*. Using these fish

lines and spinning disk confocal microscopy, we observed a remarkable efficiency of targeting

to intercellular junctions, for both proteins (data not shown for eGFP-Jam2a; for eGFP-Jam3b,

see **Figure 5B** and **C**; in the maximum z-projections, the green signal is enriched at

intercellular junctions; in single z-plane images, an EHT pol+ cell and 2 EHT pol- cells are

visible in the left and right panels, respectively). Deployment of the aortic wall into 2D

cartographies using an algorithm described previously (Lancino et al. 2018) allows to point

precisely at junctional pools established at the interface between endothelial and EHT cells.

This emphasizes on the enrichment of eGFP-Jam3b at antero-posterior poles of EHT pol+ and

EHT pol- cells (**Figure 5B, C** bottom panels*,* EHT and aortic endothelial cells (red and black

numbers, respectively) are visualized as if the eye is located inside the aortic lumen; red arrows

point at junctional reinforcements located at antero-posterior poles of EHT cells, black arrows

in the left panel point at tri-cellular junctions and the blue arrow in the right panel points at

intercalation of endothelial cells 3 and 6 between the EHT pol- cell (cell 2) and the endothelial

neighbour 7).

Using double transgenic *Tg(kdrl:eGFP-Jam3b; kdrl:nls-mKate2)* embryos at 48 - 55 hpf*,* we

addressed the recycling capacity of junctional pools using Fluorescence Recovery After

Photobleaching (FRAP; for more details, see ***Methods***). The labelling of nuclei with nls-

mKate2 allowed to point at EHT pol+ and EHT pol- cells unambiguously because eGFP-Jam3b

is essentially targeted to junctional interfaces and not labelling luminal/abluminal membranes

(***Figure 6A, B***), except in few cases (***Figure 5B***, for an EHT pol+ cell); additionally, nuclei of

EHT pol+ cells have a crescent shape (see ***Figure 6A)***.

First and to set up our protocol, we spotted bi- and tri-junctional contacts between endothelial

cells (***Figure 6 – figure supplement 1***). Recycling parameters (fluorescence intensity

recovery with time and maximum recovery amplitude (that addresses the mobile pool), ***Figure***

6C) showed that bi-junctional contacts are less mobile than tri-junctions, with a higher

dispersion of maximum recovery amplitude values for the latter (***Figure 6E***); this introduced a

clear limitation for statistical significance of the results, although clear tendencies were

observed for mean fluorescence intensity recovery ***(Figure 6C)*** and median values for

maximum recovery amplitude, when recycling reached its equilibrium ***(Figure 6E)***. We then

focused at antero-posterior sites of EHT pol+ and EHT pol- cells and more specifically in the

region of tricellular junctions (shared by 1 EHT cell and 2 endothelial neighbours (***Figure 6A,***

B, 2D-cartographies, black arrows) and for more examples in 2D-cartographies, see ***Figure 6***

– figure supplement 1, black arrows) that are clearly the most enriched with eGFP-Jam3b. In

each experiment, eGFP-Jam3b pools at tricellular junctions between endothelial cells (EC-EC-

EC) were also spotted for a comparative analysis. Measurements of fluorescence recovery

intensities revealed a tendency of increase in the mobile fraction of eGFP-Jam3b at EHT pol+

– EC versus EHT pol- – EC junctional interfaces (see **Figure 6D** for fluorescence recovery

curves and **Figure 6E** for median values (50% and 39% maximum recovery amplitudes after

10 min, respectively)). Although smaller, differences in recovery were also measured between

EHT pol+ – EC and EHT pol- – EC junctional interfaces when focusing on the earliest time

points (the first 30 seconds, **Figure 6F**), with the median value of early recovery slopes for the

EHT pol+ - EC versus EHT pol- – EC junctional interfaces increased by 128% (**Figure 6G**).

Altogether and unexpectedly regarding our initial scenario, these results indicate that tri-

junctional pools localized at antero-posterior poles of EHT cells and enriched with eGFP-

Jam3b molecules are significantly more dynamic for EHT pol+ cells in comparison to EHT pol-

cells. Since EHT pol+ cells - by virtue of apicobasal polarity establishment possibly assemble

an apical endosome -, this should favour a faster recycling of eGFP-Jam molecules (see

Discussion for more details).

ArhGEF11/PDZ-Rho GEF plays essential functions during EHT progression

Junctional maintenance and recycling are dependent on intracellular membrane trafficking,

supported by sub-cortical actin remodelling and actomyosin contractility, which are controlled

mainly by GTPases of the Rho family (Ridley 2006; Olayioye, Noll, and Hausser 2019). Owing

to the significance of apicobasal polarity control on EHT features, as suggested by our work,

we investigated which proteins may be essential for actin/actomyosin regulation and focused

on regulators of Rho GTPases, in particular Rho GEFs that catalyse exchange of GDP for GTP

to positively regulate their activity (Rossman, Der, and Sondek 2005). As for Pard3 proteins,

several of these Rho GEFs contain one or several PDZ domain(s) that target most proteins to

complexes acting at the apical side therefore interlinking actin/actomyosin regulation with cell

polarity (Mack and Georgiou 2014; Ebnet and Gerke 2022). We focused on 9 PDZ-domain

containing Rho GEFs, all encoded by different genes in the zebrafish (**Figure 7- figure**

supplement 1): ArhGEF11/PDZ-RhoGEF (thereafter shortened as ArhGEF11), ArhGEF12a,

ArhGEF12b, PRex1, PRex2, Tiam1a, Tiam1b, Tima2a, Tiam2b. We first investigated their

expression by Whole mount In Situ Hybridization (WISH) and found that all 9 mRNAs are

detected in the aorta, and for the vast majority at 30-32 and 48-50 hpf (**Figure 7- figure**

supplement 2)**.** Then, using qRT-PCR (**Figure 7- figure supplement 1B**), we measured and

compared their expression levels in the trunk region (at 35 and 48 hpf), for dt-runx1 expressing

embryos and controls. We found that, in comparison to controls, ArhGEF11, ArhGEF12b,

Tiam1b and Tiam2a are significantly reduced upon dt-runx1 expression at 48 hpf, hence being

positively controlled in the wild type condition, when the emergence of EHT cells is peaking.

This is consistent with a functional link between these Rho-GEFs, actomyosin and the control

of junctional dynamics during EHT progression. We finally decided to focus on ArhGEF11 for

the following reasons: (i) in comparison to Tiams that act on Rac1, ArhGEF11 and ArhGEF12

(which are close relatives and can form heterodimers (Chikumi et al. 2004)), are mostly acting

on RhoA which is controlling apical constriction via the RhoA-Myosin II signalling axis; as we

have shown previously (Lancino et al. 2018), EHT progression requires the constriction of

circumferential actomyosin; (ii) ArhGEF11 was shown to regulate the integrity of epithelial

junctions by interacting directly with the scaffolding protein ZO-1, hence connecting inter-

cellular adhesion with RhoA-Myosin II (Itoh et al. 2012); (iii) ArhGEF11 mediated apical

constriction is essential during tissue morphogenesis such as, for example, the neural tube

formation in which epithelial cells, submitted to mediolateral contractile forces, constrict at their

apical side thus triggering inward bending of the neural plate which leads to the formation of

the neural tube (Nishimura, Honda, and Takeichi 2012). The EHT may share features with this

process, i.e the anisotropic distribution of contractile forces controlling the plane of emergence;

(iv) mammalian ArhGEF11 exhibits alternative splicing in its C-terminal region that controls

tight junction stability via the regulation of RhoA activity (Lee et al. 2018) as well as cell

migration and invasion (Itoh et al. 2017).

To confirm the potential function of ArhGEF11 at the junctional interface between HE/EHT and

endothelial cells, we first investigated its intracellular localization. We attempted to detect the

full-length form upon expression of a GFP fusion protein, in the vascular system, and failed to

do so. We then generated a truncated form that retains the N-terminal fragment encompassing

the PDZ and RGS domains (see **Figure 7 – figure supplement 3A**) fused with eGFP in its C-

terminus. Upon transient expression in the vascular system, we visualized its localization at

the interface between endothelial and hemogenic cells progressing throughout the EHT, with

an apparent increase in density at antero-posterior regions between adjoining EHT cells

(**Figure 7 - figure supplement 3B**).

In line with the formerly described function of a splicing variant of ArhGEF11 in controlling tight

junction integrity, particularly during egression of cells from the skin epithelium in the mouse

(Lee et al. 2018), we questioned the potential role of such variant in EHT. This variant, referred

to as the mesenchymal form, results from the inclusion of a peptide encoded by exon 37 or

exon 38 in mouse and human, respectively (Shapiro et al. 2011; Itoh et al. 2017; Lee et al.

2018). This insertion locates in the degenerated C-terminal region of the protein, which is

predicted to be relatively poorly organized in its 3D structure. Upon amplifying the full-length

sequence of zebrafish ArhGEF11 for investigating its localization, we cloned fragments

encoding – or not – for the insertion of a peptide of 25 amino-acid residues (75 base pairs

corresponding to exon 38). Although variable when compared with other species (see **Figure**

7 – figure supplement 4C, *bottom panel*), this peptide is inserted in the same region as it is

for mammals and may correspond to an ArhGEF11 variant functionally equivalent and involved

in the regulating of junctional stability (see **Discussion**).

To investigate the function of ArhGEF11 on the junctional interface between HE/EHT and

endothelial cells, and more specifically of the isoform containing the exon 38 encoded peptide,

we used both morpholino (MO) and CRISPR-based strategies. We designed a splicing MO at

the 3-prime exon/intron boundary of exon 38 that interferes with its inclusion in the encoding

mRNA (**Figure 7 - figure supplement 4A**). This MO did not trigger any deleterious effect on

the gross morphology of zebrafish embryos (**Figure 7 – figure supplement 4B**) and blood

circulation was normal in comparison to control embryos. We attempted to generate CRISPR-

based genomic deletion mutants of exon 38, both using Cpf1 and Cas9 enzymes and failed

(see ***Methods***). However, using CRISPR/Cas9, we obtained a deletion mutant triggering a

frame shift downstream of exon 38 and introducing a premature stop codon few amino-acid

residues downstream thus leading to a sequence encoding for an ArhGEF11 C-terminal

deleted form (see ***Methods*** and ***Figure 7- figure supplement 4C***)). Unlike the variant skipping

exon 38 induced by the splicing MO, expression of the CRISPR/Cas9 C-ter deletion mutant

triggered, around 24 hpf, a severe retardation of blood circulation initiation in approximately

80% of the embryos obtained from incrossing ArhGEF11[CRISPR-Cterdel-/+] heterozygous mutant

fishes. From 24 to 48 hpf, approximately 50% of these embryos recovered a normal blood

circulation (suggesting that these embryos are probably heterozygous for the mutation),

approximately 35% remained with a severe phenotype (characterized by a large pericardiac

oedema) and approximately 15% died, see ***Figure 7 – figure supplement 4D*** and ***Methods***).

This indicates essential functions of the C-terminal region of ArhGEF11, in agreement with

previously published data on the mammalian protein (Chikumi et al. 2004).

We then characterized more in depth the MO and CRISPR phenotypes (***Figure 7***) and

performed a quantitative analysis of the number and morphology of HE, EHT and adjoining

endothelial cells, based on confocal images and subsequent segmentation of cell contours

(using 2D deployment of the aortic cylinder).

For the morphants, we found approximately a double amount of hemogenic cells in the aortic

floor in comparison to controls (altogether localized on the floor and on the lateral sides of the

aorta, see ***Figure 7A,*** panels in **(a)** for 2D-cartographies to visualize cell shapes, with the

accumulation of hemogenic cells delineated in yellow for one morphant on the right and panels

in **(a')** for cell counts of endothelial, hemogenic and EHT cells either in the entire aorta in trunks

(left) or, more precisely, the aortic floor and lateral aortic side and the roof (right)); see also

Figure 7 - figure supplement 5 for additional examples). Increase in the number of

hemogenic cells in morphants in comparison to controls also concerned cells oriented more

perpendicularly to the blood flow axis (toward the Y-axis, **Figure 7A,** see panels in **(a")**), consistently with variation in emergence angle, as we have shown previously (Lancino et al. 2018). This is accompanied by a significant reduction in the number of morphologically characterized EHT cells (**Figure 7A** and **Figure 7 – figure supplement 5**). Similar results were obtained for the CRISPR homozygous deletion mutants, particularly regarding the accumulation of hemogenic cells, although with a smaller amplitude (**Figure 7B**, with the panels **(b - b")** mirroring the presentations of results obtained for morphants and internal controls in **Figure 7A**); see also **Figure 7 - figure supplement 6** for additional examples). Consistently, a significant increase in the aortic perimeter was found, at least for the morphants (**Figure 7C**). Of notice also, hemogenic cells in morphants are on average less elongated in comparison to control and to the CRISPR homozygous mutants (as well as less dispersed according to this parameter, compare **Figure 7A**, panel **(a"')** with **Figure 7B**, panel **(b"'),** with visualization of the dispersion on right panels). Importantly, we observed for the morphants an accumulation of a less elongated cell population standing below the median value (**Figure 7A,** panel **(a"')**; to some extent, this is reproduced as well when measuring hemogenic cell area, compare **Figure 7A**, panels in **(a"")** with **Figure 7B**, panels in **(b"")** for morphants and CRIPR mutants, respectively). Finally, the population of small cells in the morphants are of uncharacterized morphology, which means that they appear neither as EHT pol+ nor as EHT pol- cells and may represent abortive emergences (**Figure 7A** and **Figure 7 – figure supplement 5,** cells in cyan boxes).

Additionaly, we assessed the impact of MO interference on ArhGEF11 exon38 splicing on hematopoiesis using the *Tg(CD41:eGFP)* fish line. By performing live imaging on *Tg(CD41:eGFP) X Tg(Kdrl:nls-mKate2)* outcrosses (see **Figure 7 – figure supplement 7A** and **B** for control embryos and morphants, respectively), we quantified the number of hemogenic/EHT cells in the aortic floor, based on fluorescence intensities as well as the number of newly generated hematopoietic stem and progenitor cells (HSPCs, that continue to express the hematopoietic eGFP reporter but stopped to express the vascular mKate2

reporter, *Figure 7 – figure supplement 7C*). We show that, in agreement with our

morphometric analysis, an increase in the number of hemogenic/EHT cells in the aortic floor

in morphants in comparison to control embryos (*Figure 7 – figure supplement 7D*, with

median values of 14.0 and 22.5 hemogenic/EHT cells for controls and morphants,

respectively). Additionally, we observe, despite the high variability in both controls and

morphants, a relative diminution in the number of newly generated HSPC precursors (*Figure*

7 – figure supplement 7E, with median values of 22.0 and 16.5 HSPCs for controls and

morphants, respectively). This is consistent with our previous results showing an accumulation

of HE/EHT cells in aortic floors of morphants and highlights the negative impact of this

accumulation on the generation of HSPCs.

Altogether, the morphants and CRISPR mutant phenotypes show that the progression

throughout EHT is significantly impaired supporting the idea that ArhGEF11 exerts important

functions in the process. Increase in the frequency of elongated cells for the CRISPR mutants

in comparison to morphants suggests that interference triggered by the C-terminal deletion of

ArhGEF11 is more effective at early time points. Interference with inclusion of exon 38 in

morphants triggers an increase in the number of hemogenic cells of smaller area and reduced

elongation indicating a more efficient progression throughout emergence in comparison to the

CRISPR mutants, albeit with impairment at later stages. This may be due to alteration in the

dynamics of endothelial cell intercalation which is required for emergence completion (which

is also compatible with the accumulation of pairs of cells that may be post-mitotic and that do

not spread apart because this also requires aortic cell intercalation).

ArhGEF11/PDZ-RhoGEF and its variant containing the exon 38 encoded sequence

control junctional dynamics to support reduction of the hemogenic/endothelial

interface and cell-cell intercalation

To investigate how ArhGEF11 controls the dynamic interplay between HE/EHT and endothelial

cells, we characterized further the MO and CRISPR phenotypes, at the junctional level. To do

so, we performed FRAP experiments using the *Tg(Kdrl:eGFP-Jam3b)* fish line that was

injected at the one cell stage with the exon 38 splicing MO, or incrossing *(Kdrl:eGFP-Jam3b x*

*ArhGEF11*CRISPR-Cterdel-/+) heterozygous fishes to obtaining homozygous mutants and wild type

alleles for control siblings.

FRAP experiments were focused on the hemogenic (HE)/endothelial (EC) membrane

interface, with the support of images obtained after 2D deployments of aortic segments (***Figure***

8A). Pools of *eGFP-Jam3b* molecules localized at bi-junctional (HE-EC, green arrows in panel

A) and at tri-junctional interfaces (HE-EC-EC (magenta arrows in panel **A**) and HE-HE-EC

(black arrows in panel **A**)) were bleached and subsequently imaged for FRAP analysis. In the

case of the MO treatment (in which case HE cells accumulate on the aortic floor, see the 2D

cartography ***Figure 8A***)) we measured, in comparison to controls, an increase in FRAP

recovery parameters for *eGFP-Jam3b* pools at the tri-junctional HE-HE-EC interface, with a

prominent tendency for an increase in the recovery speed at early time points (***Figure 8B,***

panel **(b''')**, values obtained from fluorescence intensity recoveries whose mean values are

represented by curves in panel **(b'')**; compare with maximum recovery amplitudes **(b'),** with

values obtained from the plateaus of fluorescence recovery intensities whose mean values are

represented in panel **(b)**). Results are different for the ArhGEF11CRISPR-Cterdel+/+ homozygous

mutants for which we did not observe any obvious effect at HE-HE-EC tri-junctions (***Figure***

8C, with the panels **(c - c''')** mirroring the presentations of results obtained for morphants and

internal controls in ***Figure 8B***) but, rather, a small tendency for an increase in the mobile pool

at HE-EC-EC tri-junctions (that has the particularity to concern 2 endothelial cells that contact

the longitudinal membrane of one HE cell, see ***Figure 8A, D***).

Altogether and including our results presented ***Figure 7***, these results reinforce the idea that

ArhGEF11 and its variant encoded by exon 38 are involved in critical steps of EHT progression,

partly by controlling the dynamics of junctional pools at the level of tri-junctions established

between adjoining hemogenic and endothelial cells (see the model **Figure 8D** and its legend

for a detailed analysis of mechanistic issues). The results also suggest that the +exon38

peptide encoding variant is prominently involved in controlling the HE-HE-EC interface and

more specifically during intercalation, consistently with the accumulation, upon MO

interference, of EHT cells having progressed throughout the emergence but appearing to be

slowed down when reaching completion (cells of smaller area and less elongated, see **Figure**

7). Finally, ArhGEF11 is controlling RhoA and, most probably, its function in EHT progression

also involves actin and acto-myosin activities that are involved in the contraction of HE and

EHT cells (see Lancino et al. 2018; this is coupled with the consumption of longitudinal and

transversal membrane interfaces, see also the model **Figure 8D** and its legend).

Discussion

Here, we bring new insight into the cell biology and mechanics of the EHT, using live confocal

microscopy, RNAscope (to reveal the expression of mRNAs at the cellular resolution), and

genetic interference. We show that the emergence of EHT cells is more complex than

previously shown, with two types of EHT cells of highly different morphodynamics and apico-

basal polarity status. We bring evidence suggesting that these differences take root in HE

maturation that is primarily characterized by loss of apico-basal polarity that is then tuned by

the activity of the transcription factor Runx1. This leads to two types of EHT cells whose

respective emergence characteristics appear to rely, partly at least, on controlling the recycling

of junctional complexes directly linked to apico-basal polarity features (the JAMs). Also, we

reveal essential aspects of the interplay between HE/EHT cells and adjoining aortic endothelial

cells. The first is based on a potential signalling cross talk, whose molecular components

remain to be identified, controlling apico-basal polarity and its associated features, in space

(and possibly also in time, because of EHT asynchrony over the EHT time-window). The

second involves the control of the dynamics of junctional interfaces to reduce contacting

surfaces between aortic and HE/EHT cells and to promote cell-cell intercalation for the ultimate

release of EHT cells and proper sealing of the aortic floor.

EHT morphodynamics in the zebrafish embryo in comparison to amniotes

EHT cell emergence in the zebrafish embryo is relatively unique; cells extrude from the aortic

floor toward the sub-aortic space and not in the aortic lumen, probably because of mechanical

constraints on the aortic wall (discussed in Lancino et al. 2018). However, as a common feature

in amniotes and the zebrafish, EHT cells appear to contain large vacuolar structures (which is

also the case for HE cells in the zebrafish embryo, as we show); recently, this has been

precisely described in chick and quail embryo models (Sato et al. 2023). In amniotes, in which

EHT cells are round, these vacuoles appear to emanate from the abluminal membrane (facing

the sub-aortic space) and not from the luminal membrane. In addition, these structures

disappear before EHT completion, based on water chase by aquaporin activity (Sato et al.

2023); it has been suggested that they contribute to exert mechanical force to initiate the

detachment of the cell from the endothelial layer. We do not know the membranes from which

the pseudo-vacuolar structures that we observe originate, particularly in the HE (they may be

formed by transcytosis from the basal membrane, by endocytosis from the luminal membrane,

or coming directly from the biosynthetic pathway). However, we can exclude that they

contribute to the rounding of EHT pol- cells, as is the case for avians, since we do not see

them during EHT pol- emergence. For EHT pol+ cells, according to our results (see ***Figure 2A***

and our hypothetical model ***Figure 2B***), the most plausible scenario is that the apical/luminal

membrane is established by fusion of cytosolic vacuolar structures or redistribution of internal

pseudo-vacuoles connected to the luminal membrane. Subsequently, the dynamic properties

of the EHT pol+ cell luminal/apical membrane allows its rapid evolution after EHT completion

(*see **Figure 2 – figure supplement 1***). We propose that this is compatible with the activity of

aquaporins in late EHT pol+ cells and/or shortly after their release, for water chase and

vacuolar regression as proposed in our hypothetical model (see ***Figure 1 – figure supplement***

1B and the legend for details). Water chase activity would then be a common theme controlling

the morphology and dynamics of EHT cells throughout vertebrates, although not for the same

purposes.

Characteristics of the HE and complexity of pre-hematopoietic stem cell emergence

Heterogeneity in the identity and hematopoietic potential of cells emerging from the HE in the

AGM has been recently shown in co-culture systems and *in vivo* in the mouse (Ganuza et al.

2017; Dignum et al. 2021) and in the zebrafish model (Tian et al. 2017; Xia et al. 2023);

however, knowledge on the fundamental cell biological and molecular events leading to this

complexity is still lacking. Here, we highlight functional aspects that may significantly contribute

to this complexity and that take root in the cellular plasticity of the HE. We bring evidence

suggesting that HE cells that become competent to initiate the EHT time-window (around 28-

30 hpf) tune their apicobasal polarity status which endows them with the ability to support two

emergence processes with radically different morphodynamic characteristics; eventually, this

would lead to differentially fated cells, which remains to be determined.

While the ability of the HE to regulate its polarity features may be inherited from molecular

cues involved in its upstream intra-aortic specification (such as for example NOTCH (Robert-

Moreno et al. 2008; Gama-Norton et al. 2015; Bonkhofer et al. 2019), TGFβ (Monteiro et al.

2016), Gata2 (Butko et al. 2015; Daniel et al. 2019)), or any other of the more upstream factors

involved in hemogenic specification (Zhao et al. 2022)), our results suggest that Runx1 is

taking part in this regulation, consistently with its hematopoietic and EHT-inducing activity (T.

North et al. 1999; Kalev-Zylinska et al. 2002; Kissa and Herbomel 2010; Lancrin et al. 2012).

Our experiments using an interfering form of Runx1 (dt-Runx1) that produces a relatively weak

and subtle phenotype in comparison to morpholino treatment that prevents aortic cells to

convert into HE cells (Bonkhofer et al. 2019) or genetic mutation and gene knockout abrogating

hematopoiesis (T. E. North et al. 2009; Sood et al. 2010; Gao et al. 2018), provided the

opportunity to observe that, at 30 hpf (the timing point at which a majority of HE cells are still

paving the aortic floor), elongated HE cells appear to be maintained and to exhibit apicobasal

polarity, with the absence of apparent cytosolic vacuolar structures (as has been described in

the mouse in Runx1 interfering condition (T. North et al. 1999)); this correlates with the

subsequent accumulation of EHT pol+ cells. This is in favour of the idea that the polarity status

of HE cells has a direct incidence on the cell type that will arise, with a stronger chance to

evolve as an EHT pol+ cell if apical and basal domains are already determined (which is the

case for HE cells upon dt-Runx1 expression) and a lesser chance if these domains are not

established and the pool of intracellular vacuoles/pseudo-vacuoles is reduced (see the

hypothetical model **Figure 2B**).

A key question arising from our observations is what would control the balance orienting toward

an EHT pol+ or an EHT pol- type of emergence. Cell division may be orienting toward EHT

pol- emergence type (which would also be the reason why these cells have a round-shape

morphology in comparison to EHT pol + cells). The frequency of EHT pol+ versus EHT pol-

cells that we have estimated and that is of approximately 2/1 is not incompatible with this

possibility since divisions take place in the HE at a relatively high frequency. Indeed, it has

been estimated that 50% of HE cells that constitute the aortic floor at 28 hpf will divide and will

undergo emergence thereafter during the 28 - 72 hpf time window (Zhao et al. 2022). However,

we observe that both EHT pol+ and EHT pol- cells can divide during the emergence process

(see also our former work Lancino et al. 2018 for EHT pol+ cells), which is followed by the full-

recovery of their respective morphodynamic characteristics. Hence, this excludes the

possibility of a short term and purely mechanical effect of cell division on emergence

morphodynamics. In link with cell division however, there exists the possibility that the pool of

large vesicular structures potentially awaiting for exocytosis in HE cells having lost their

apicobasal polarity may become asymmetrically inherited after division. This would endow the

cell remaining with the largest pool to undergo EHT pol+ emergence after exocytosis and

expansion of the apical/luminal membrane. Our results **Figure 2** would support this possibility

but what controls this asymmetry would remain to be established.

Still, regarding the control of the balance orienting toward an EHT pol+ or an EHT pol- type of

emergence, the intrinsic nature of the HE, i.e. its acquaintance with the aortic differentiation

programme (Bonkhofer et al. 2019), raises the possibility of a partial recovery of an aortic cell

phenotype in the case of EHT pol+ emergence, thus supporting its very unique

mechanobiology. Therefore, the regulation of local signalling between aortic and HE cells may

tune the balance toward the one or the other fate; this could be the case, for example, for

adjoining HE cells in comparison to isolated ones that would then receive unequal signalling

cues owing to the difference in homotypic versus heterotypic contacting surfaces. In this

scenario, NOTCH signalling which is determinant for aortic and HE specification (Gama-Norton

et al. 2015) may play a significant role, particularly via the ratio of the Jag1/Dll4 ligands that is

fundamentally involved in HE versus aortic specification (Robert-Moreno et al. 2008;

Bonkhofer et al. 2019).

Finally, we cannot totally exclude that EHT pol+ and EHT pol- cells would originate from

different precursors. On this line, it has been shown that the aortic wall can home a transient

veinous-fated progenitor cell population that will lead to the underlying vein via selective

sprouting from 21 to 23 hpf (Herbert et al. 2009). This option would require a massive

expansion of a minor fraction of the cell population remaining in the aortic floor since it is

expected that most of it has been moving out toward the sub-aortic region, which is not in

favour of this possibility. In addition, recent scRNAseq analysis of floor and roof aortic cells at

21 and 28 hpf has led, for both timing points, to two well defined clusters defined only by aortic

and hematopoietic signatures (Zhao et al. 2022).

Spatially restricted control of Pard3ba mRNAs by Runx1

Recent work has identified Runx1 targets in the zebrafish embryo at 29 hpf (Bonkhofer et al.

2019); however, the results do not point at any of the conventional apicobasal polarity

organizers as being direct targets although Pard3 and Pard6 mRNAs are clearly

downregulated in the HE in comparison to aortic roof cells. Our work now strengthens the idea

that understanding the mechanisms underlaying the EHT may require more than simply

deducing from expression of mRNAs in the HE (including the corresponding proteins), as

exemplified by members of the Pard3 family. Also, annotation of protein isoforms in reference

databases needs to be improved so as to gain precise and accurate insight on mechanistic

processes.

Our results reveal the unique sensitivity of Pard3ba mRNAs expression to Runx1 interference,

in comparison to the Pard3aa and ab isoforms. Also, the dt-Runx1 mutant phenotype with

enhanced expression of Pard3ba mRNAs in MERs highlighted and strengthened the existence

of these specific aortic regions in wild type animals, which could not have been so obviously

pointed at because of its relatively low and variable expression. Importantly, the expression of

Pard3ba mRNAs is mainly detected in AECs and we cannot for sure say that hemogenic/EHT

cells also express this mRNA because very few spots are detected and those may be at the

limit of the technical background (although we measured a tendency to increase their detection

upon dt-Runx1 expression, indicating that they may indeed be expressed in HE/EHT cells but

at very low levels, in wild type embryos).

The sensitivity of AECs on Runx1 interference is somehow unexpected since Runx1 is rather

expected to be involved in HE maturation as well as during EHT. This raises the possibility of

a direct signaling cross talk between HE/EHT cells and aortic adjoining cells. In this scenario,

HE/EHT cells exposed to Runx1 signaling would instruct AECs that subsequently regulate their

apico-basal polarity features (via Pard3ba expression); this would then feedback on HE/EHT

cells. Another plausible explanation is that AECs have been exposed to interference on Runx1

activity upstream of the EHT time-window, at a timing point at which aortic and HE cell types

have been sharing a common ancestor, a possibility that has been debated and reinforced in

the zebrafish model (Bonkhofer et al. 2019; Zhao et al. 2022). Of notice, our dt-Runx1 mutant

is expressed under the vascular kdrl promoter that may be active sufficiently early (Liao et al.

1997) to allow efficient interference with endogenous Runx1 activity on a relatively long time

scale; this may take place in a more or less restricted population of precursor cells, according

to the spatially restricted localization of Pard3ba MERs.

The increase of Pard3ba mRNAs detection, upon dt-Runx1 mutant expression, suggests that

endogenous Runx1 negatively controls this specific mRNA in wild type embryos. This is

consistent with our results since there are two cell types/stages during the EHT process for

which we observe a negative regulation of apico-basal polarity, i.e the maturing HE and EHT

pol- cells, as visualized upon expression of eGFP-podxl2 and mKate2-podxl2 (the fluorescent

fusion proteins are partitioned equally between the luminal/apical and the basal membranes).

Altogether, this reinforces the idea that Runx1 negatively controls apico-basal polarity in HE

and EHT pol- cells; at least in part, it does so via the Pard3ba isoform that we hypothesize acts

as a tuner, consistently with the high sensibility of its mRNA expression to Runx1 activity and

the recurrent variability in expression that we are measuring. Owing to the apparent

stochasticity of the EHT process in time and space (emergence takes place asynchronously

during the time window of the EHT that ranges from approximately 28 to 72 hpf and proceeds

from place to place, all along the floor of the dorsal aorta), this may be what contributes to the

experimental variability that we are measuring at a given time using qRT-PCR and RNAscope

(for all determinations of Pard3ba mRNAs in comparison to other isoforms; of notice, variations

are even larger for the dt-Runx1 mutants, which is consistent with the high sensitivity to dt-

Runx1 expression levels that are expected to be relatively different from one transgenic

embryo to the other).

Altogether, our results are consistent with the hypothesis that the variability in expression of

Pard3ba mRNAs reflects the response to intrinsic fluctuations of the system over time and

space and reinforce the idea that these conditions, mediated by Runx1 and environmental

constraints imposed by the blood flow, tune apico-basal polarity. This tuning scenario would

also explain the specific case of EHT pol+ cells that would then proceed as such upon

exposition to positive apico-basal signaling when Runx1 activity is decreased, consistently with

the tendency to accumulate EHT pol+ cells upon expression of our dt-Runx1 mutant.

Unfortunately and presently, we cannot show a spatial proximity between EHT pol + cells and

Pard3ba MERs *in situ* (or the opposite for EHT pol- cells) owing to technical limitations with

fixed embryos that do not allow the maintenance of EHT cell shapes. Resolving this issue in

the future will be very informative, also to place cellular morphodynamics and gene expression

regulation throughout the EHT, with spatial and temporal resolution.

The complexity of pre-HSC emergence and heterogeneity of hematopoietic stem cell

and progenitor populations

Currently, we do not know what would be the features that would endow the two cell types to

behave differently after their release. Since post-EHT cells have to migrate in transient

developmental niches (i.e the CHT, the thymus, the pronephric region), these features may

provide them with more or less ability to migrate in the sub-aortic space and pass throughout

the wall of the underlying vein to conquest more distant niches (Murayama et al. 2006), or to

remain in a local AGM niche and be exposed to specific signalling there, in the proximity to the

aorta as it is the case for hematopoietic clusters in mammals and avians (Ciau-Uitz et al. 2014;

Jaffredo and Yvernogeau 2014). Beside their potential differences in colonizing different niches

which would impact on their subsequent fate, these cells may also contribute to more or less

transient waves of hematopoietic stem cells and progenitors that would be specific to the

developmental period such as, for example, a restricted sub-set of T-lymphocytes that was

described before (Tian et al. 2017). In the specific case of the Tian et al. work, the AGM was

shown to produces transient T-lymphocytes proposed to be independent of long-lived HSCs

according to a gradient that increases from the anterior to the posterior part of the trunk region,

as well as from the aorta in the tail (the posterior blood island in the tail region that was

proposed to give rise to hematopoietic precursors related to mammalian erythro-myeloid

progenitors, (Bertrand et al. 2007)). Since the EHT pol+ and EHT pol- cells that we describe in

our work are equally produced from the aorta in the trunk and in the tail regions (thus

irrespective of the antero-posterior axis of the aorta), the one or the other of the two EHT types

most probably does not directly relate to the first wave of T-lymphopoiesis arising from the

aortic endothelium described in Tian et al. 2017.

The cell biology behind emergence control and morphodynamics

The intrinsic capacity of the HE, that we show here to be able to support two highly distinct

emergence processes, in the zebrafish embryo, is very unique. Indeed, not only are they taking

place contemporarily, but also can the two cell types proceed throughout emergence at a

distance shorter than 50 μm, which accommodates the intercalation of a single adjoining

endothelial cell. This may reflect the remarkable properties of the vascular system that needs

to allow cellular extrusion and transmigration. The latter is particularly relevant in the context

of developmental hematopoiesis since hematopoietic precursors that travel via blood

circulation need to pass throughout the aortic wall to conquest their distant niches.

In the present work, we provide some of the mechanistic insights into the specificities of the

two emergence types and that rely on two fundamental aspects of cellular biology, i.e

apicobasal polarity establishment and junctional mobility. The establishment and maintenance

of apicobasal polarity in a cell extruding from its environment, as is the case for EHT pol+ cells,

is at odds with the fundamental mechanisms of cell extrusion irrespective of the context into

which the process is taking place (Nieto et al. 2016; Gudipaty and Rosenblatt 2017; Pei et al.

2019; Staneva and Levayer 2023); this places the emergence of EHT pol+ cells as an extreme

case of extrusion and it is somehow unexpected to observe this type of emergence when the

aortic wall can extrude EHT pol- cells that appear to emerge contemporarily, apparently

according to a more conventional mode. During the EHT time-window, Runx1 is directly

controlling the emergence programme by regulating Rho-dependent cytoskeletal functions in

HE cells derived from mouse ES lines (Lie-A-Ling et al. 2014) and inducing the expression of

Gfi1ab in the zebrafish (Bonkhofer et al. 2019) that belongs to the family of transcriptional

repressors of the arterial programme expressed in the HE (Gfi1 and Gfi1b in the mouse,

(Lancrin et al. 2012)). This programme downregulates key molecules involved in the

maintenance of the aortic endothelium among which Cdh5 (VE-Cadh) and the downregulation

of cadherins is a hallmark of extrusion. Thus, for EHT cells, and particularly EHT pol+ cells,

downregulation of VE-Cadh should be compensated by other mechanisms that allow for the

maintenance of aortic integrity. In this context of adhesion downregulation, the JAMs (that are

expressed endogenously in the vascular system and that were used in this study) appear to

be a well-suited substitute. As part of tight junction complexes, they can strengthen adhesion

and, in addition, since they are also involved in cell migration (in particular leukocyte trans-

endothelial migration, see (Ebnet 2017)), they could in theory support the extrusion of EHT

pol- cells that do so according to a seemingly migration-type emergence mode (inferred from

our time-lapse sequences) more or less related to trans-endothelial migration. In the case of

EHT pol+ cells, we propose that JAMs may significantly contribute to the re-establishment of

apicobasal polarity and to the strengthening of adhesion while EHT pol+ cells should undergo

strong mechanical tension (owing to cellular bending). Interestingly, Jam3b appears to be a

direct and positively regulated target of Runx1 (Bonkhofer et al. 2019), which is supporting its

aforementioned potential functions during EHT.

In regard to the different modes of junctional recycling between aortic and either EHT pol+ or

EHT pol- cells for which be bring evidence using FRAP, we hypothesize that differences take

root in the regulation of cell polarity features, such as for example the organization of an apical

endosome in the case of EHT pol+ cells (which may be why recycling of eGFP-Jam molecules

appears to be faster in EHT pol+ cells in comparison to EHT pol- cells, in relation to the

recycling pathways undertaken by JAM molecules). In addition, since JAMs interact with Pard3

(Ebnet et al. 2003) this recycling, that would provide renewed JAM/Pard3 complexes at

membrane junctions and sustained activation of the Pard3/Pard6/aPKC complex (Ebnet 2017),

may contribute to the maintenance/reinforcement of EHT pol+ cell polarity throughout the

entire emergence process. Also, the regime of recycling may be regulated by an adjacent aortic

endothelial cell that expresses Pard3ba and that controls trans-cellular JAMs pairing via

993 signalling involving apico-basal polarity features. Finally, this turnover may be adapted to the

994 mechanical tensions EHT pol+ cells are exposed to, in comparison to EHT pol- cells, owing to

995 their specific morphodynamic features (ex: cellular bending); these tensions may be sensed

996 by tri-cellular junctions which would be preferential sites for the renewal of junctional

997 components whose half-life is reduced owing to submission to mechanical forces.

998 **The cell biology behind emergence control and mechanics: the function of**

999 **ArhGEF11/PDZ-RhoGEF in the dynamic interface between aortic and HE cells**

1000 Finally, regarding inter-cellular junctions and the control of their dynamics all along the EHT

1001 process, we also addressed the dynamic interface between aortic and HE cells. As for EHT

1002 cells, we focused more specifically on tri-junctional interfaces that have been proposed to

1003 sense tension and transmit the information to other types of junctional complexes (Bosveld,

1004 Wang, and Bellaïche 2018); in the EHT context, the contracting hemogenic endothelium is

1005 additionally exposed to high mechanical tension owing to wall sheer stress imposed by the

1006 blood flow and that was proposed to regulate emergence efficiency (Lundin et al. 2020;

1007 Campinho et al. 2020; Chalin et al. 2021). In our work, we highlight one essential aspect of

1008 controlling junctional recycling which is during cell-cell intercalation which concerns either

1009 contacting HE cells lying on the aortic floor or HE/EHT cells after cytokinesis (as mentioned

1010 before, cell division is rather frequent during the EHT time window (Zhao et al. 2022)). In this

1011 context, we addressed the function of ArhGEF11/PDZ-RhoGEF which provides a functional

1012 link between cell polarity (with its PDZ-binding motif and its RGS domain that couples to G-

1013 protein coupled receptor signalling to regulate planar cell polarity (Nishimura, Honda, and

1014 Takeichi 2012)), cell adhesion (it binds to the tight junction associated regulatory protein ZO-

1015 1, (Itoh et al. 2012)), and cytoskeletal regulation/contraction (Itoh et al. 2012; Itoh 2013), all

1016 three essential mechanistic components of morphogenetic movements and tissue remodelling.

1017 Since morphodynamic events leading to cell extrusion and to the control of cell migration do

1018 not only rely on the regulation of mRNAs levels but also on alternative splicing (Pradella et al.

2017) our interest for investigating the function of ArhGEF11 in EHT was raised by the fact that

the protein undergoes changes in its biological properties via the alternative splicing of a small

exon which leads to a modification of its C-terminus. We identified a splicing event in this region

(concerning exon 38) for a potential zebrafish isoform ortholog. However, the precise function

of this variant for the zebrafish protein, that was shown in the case of the mammalian isoform

to induce cell migration and motility (Itoh et al. 2017) and to bind to the Pak4 kinase thus

leading to the subsequent destabilization of tight junction complexes via loss of RhoA activation

(Lee et al. 2018), remains to be established. On this line, our results consistently indicate that

preventing exon 38 insertion interferes with EHT progression (as well as deleting the C-

terminus of the protein as shown with our CRISPR mutant) and appears to impair intercalation

of aortic cells between adjoining HE cells. Hence, the +exon 38 isoform, by regulating the

dynamic interplay between HE and endothelial cells, would favour one of the essential steps

of the EHT process leading to its completion and requiring the complete isolation of EHT cells

to ensure the ultimate sealing of the aortic floor (see our former work (Lancino et al. 2018)). It

remains to be established, as our measurements indicate, why the ArhGEF11 +exon 38

isoform appears to slow down junctional recycling which is counterintuitive with increasing the

HE/endothelial dynamic interface. However, upon interference, we have been following Jam3b

recycling which is only a sub-population of tight junction components that should be at play in

the system. Finally, the EHT process on its whole may be submitted to a complex regime of

alternating contraction and stabilization phases as we have shown previously in the context of

EHT pol+ emergence and that may reflect the necessity to adapt to mechanical constraints

imposed by the environment as well as the rearrangements of the HE/EHT/aortic cellular

interface (Lancino et al. 2018). In this context, back and forth regulatory mechanisms,

particularly involved in the control of the RhoA-Myosin II signalling axis partially regulated by

ArhGEF11, may locally and at specific timing points change the turnover of junctional

molecules, thus blurring the correlation between gross phenotype (here the organization of the

HE/aortic interface) and very local and dynamic molecular events (for a complement to the

discussion, see also the legend of the model presented ***Figure 8C***).

Overall, our work highlights the complexity of pre-hematopoietic stem cell emergence as well

as some of the essential molecular and mechanistic aspects involved, at the interface with

aortic endothelial neighbours. We show that the aorta, in the zebrafish embryo, produces two

fundamentally different types of EHT cells and propose that this results from specific functional

features of the HE regulated by the transcription factor Runx1. Among these features, apico-

basal polarity and its regulation appear to be essential for the spatial and temporal control of

EHT cell emergence which is an asynchronous process. This would support the production of

cell types potentially endowed with different cell fate potential, subsequently reinforced by the

type of niche into which these cells will establish homing. It now remains to be established if

the different cell types that we describe are indeed leading to cells that will eventually hold

specific hematopoietic fate and properties.

Materials and Methods

1058

Key Resources Table

1059

1060 For the exhaustive list of all primers, see Materials and Methods - Supplementary Table.

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
strain, strain background (zebrafish *Danio rerio*)	AB	Zebrafish International Resource Center (ZIRC)	ZFIN: ZDB-GENO-960809–7	
strain, strain background (zebrafish *Danio rerio*)	Tg(kdrl:HsHRAS - mCherry): s916Tg	(Chi et al. 2008)	ZFIN ID: ZDB-ALT-090506–2	referred to as Tg(kdrl:Ras-mCherry)
strain, strain background (zebrafish *Danio rerio*)	Tg(6.0itga2b:EGFP)	(Lin et al. 2005)	ZFIN ID: ZDB-ALT-051223–4	referred to as Tg(CD41:eGFP)
strain, strain background (zebrafish *Danio rerio*)	Tg(kdrl:eGFP)	(Jin et al. 2005)	ZFIN ID: ZDB-ALT-050916–14	
strain, strain background (zebrafish *Danio rerio*)	Tg(kdrl:nls-mKate2)	N/A	N/A	
strain, strain background (zebrafish *Danio rerio*)	Tg(kdrl:Gal4;UAS:RFP)	N/A	N/A	
strain, strain background (zebrafish *Danio rerio*)	Tg(kdrl:Gal4;UAS:RFP;4xNR:eGFP-Podxl2)	This paper	N/A	
strain, strain background (zebrafish *Danio rerio*)	Tg(kdrl:mKate2-Podxl2)	This paper	N/A	
strain, strain background (zebrafish *Danio rerio*)	Tg(kdrl:eGFP-Jam2a)	This paper	N/A	
strain, strain background (zebrafish *Danio rerio*)	Tg(kdrl:eGFP-Jam3b)	This paper	N/A	
strain, strain background (zebrafish *Danio rerio*)	Tg(kdrl:Gal4;UAS:RFP;4xNR:dt-runx1-eGFP)	This paper	N/A	
strain, strain background (zebrafish *Danio rerio*)	Tg(ArhGEF11CRISPR-Cterdel-/+)	This paper	N/A	

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
recombinant DNA reagent	pG1-flk1-MCS-tol2	(Jin et al. 2005)	N/A	
recombinant DNA reagent	pEGFP-C1	Clontech	N/A	
recombinant DNA reagent	Transposase pCS-zT2TP	(Suster et al. 2011)	N/A	
recombinant DNA reagent	pmKate2-f-mem	Evrogen	Cat# FP186	
antibody	goat anti-rabbit Alexa Fluor 488	Invitrogen	Cat# A11070	1/400
antibody	rabbit anti-GFP	MBL	Cat# 598	1/300
antibody	mouse anti-HA	Sigma	Cat# 12ca5	1/50
antibody	goat anti-mouse HRP-conjugated	Thermo Fisher Scientific	Cat# G-21040	1/300
sequence-based reagent	Morpholino sih	(Sehnert et al. 2002)	ZDB-MRPHLNO-060317-4	CATGTTTGCTCTG ATCTGACACGCA
sequence-based reagent	Morpholino ArhGEF11 exon 38	This paper	N/A	GAAATAAATGAAG CCCCACCTCCGT
sequence-based reagent	Oligo for CRISPR: Alt-R® CRISPR-Cas12a (Cpf1) crRNAs -1	This paper	N/A	TATCACACACACA TCACCTTCTA
sequence-based reagent	Oligo for CRISPR: Alt-R® CRISPR-Cas12a (Cpf1) crRNAs -2	This paper	N/A	TTTCTCAGCGCTC CTGACAGATG
sequence-based reagent	Oligo for CRISPR: Alt-R® CRISPR-Cas9 crRNAs	This paper	N/A	AGCCAATCGTCTG AGGACGG
sequence-based reagent	Oligo for CRISPR: Alt-R® tracrRNA	IDT	Cat# 1072532	
sequence-based reagent	Oligo for WISH: RNAscope™ Probe-Dr-myb-C3	ACD-Biotechne	Cat No. 558291-C3	Cat No. 558291-C2
sequence-based reagent	Oligo for WISH: RNAscope™ Probe-Dr-pard3ab-C3	ACD-Biotechne	Cat No. 1282521-C3	
sequence-based reagent	Oligo for WISH: RNAscope™ Probe-Dr-pard3ba-C2	ACD-Biotechne	Cat No. 1309581-C2	
sequence-based reagent	Oligo for WISH: RNAscope™ Probe-Dr-pard3aa-C2	ACD-Biotechne	Cat No. 1305321-C2	
software, algorithm	Imaris	Oxford Instruments	https://imaris.oxinst.com/	v10.1.0
software, algorithm	CRISPRscan	(Moreno-Mateos et al. 2015)		N/A

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
software, algorithm	Zen Pro2 software	Zeiss	https://www.zeiss.com/microscopy/en/products/software/zeiss-zen.html	N/A
software, algorithm	QUANTSTUDIO™ DESIGN & ANALYSIS 2	Thermofisher	https://apps.thermofisher.com/apps/da2/	N/A
software, algorithm	MetaMorph software	Molecular Devices	https://www.moleculardevices.com/products/cellular-imaging-systems/acquisition-and-analysis-software/metamorph-microscopy	N/A
software, algorithm	Acquisition-Imaging analysis-LAS X	Leica	http://www.leica-microsystems.com	N/A
software, algorithm	Image Analysis-Fiji	NIH	https://imagej.net/Fiji	N/A
software, algorithm	Image Analysis Icy	(De Chaumont et al. 2012)	http://icy.bioimageanalysis.org/	N/A
software, algorithm	Image Analysis TubeSkinner plugin for Icy	(Lancino et al. 2018)	https://icy.bioimageanalysis.org/plugin/tubeskinner/	N/A
software, algorithm	Figures-Illustrator CC 2017.1.1	Adobe	http://www.adobe.com/cn/	N/A
software, algorithm	Active contour plugin	(Dufour et al. 2011)	https://icy.bioimageanalysis.org/plugin/active-contours/	N/A
software, algorithm	NIS software	Nikon	https://www.microscope.healthcare.nikon.com/en_EU/products/software/nis-elements	N/A
software, algorithm	Rstudio	RStudio Team (2020)	http://www.rstudio.com/}	N/A
software, algorithm	ggplot2	(Wickham 2016)	https://ggplot2.tidyverse.org.	N/A
software, algorithm	readr	(Wickham, Hester, and Bryan 2022)	https://github.com/tidyverse/readr	N/A
software, algorithm	cowplot	(Wilke 2020)	https://wilkelab.org/cowplot/	N/A
software, algorithm	dplyr	(Wickham et al. 2023)	https://github.com/tidyverse/dplyr	N/A
software, algorithm	wesanderson	(Ram and Wickham 2018)	https://github.com/karthik/wesanderson	N/A

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
software, algorithm	matrixStats	(Bengtsson et al. 2023) 09/02/2024 14:57:00	https://github.com/HenrikBengtsson/matrixStats	N/A
software, algorithm	stringr	(Wickham 2022)	https://github.com/tidyverse/stringr	N/A
software, algorithm	ggsci	(Xiao 2018)	https://github.com/nanxstats/ggsci.	N/A
software, algorithm	ggpubr	(Kassambara 2020)	https://rpkgs.datanovia.com/ggpubr/	N/A
software, algorithm	ggstatsplot	(Patil 2022)	https://indrajeetpatil.github.io/ggstatsplot/index.html	N/A
software, algorithm	ggbeeswarm	(Clarke and Sherrill-Mix 2017)	https://github.com/eclarke/ggbeeswarm	N/A
software, algorithm	DT	(Xie, Cheng, and Tan 2022)	https://github.com/rstudio/DT	N/A
software, algorithm	SciViews	(Grosjean 2019)	https://www.sciviews.org/SciViews/	N/A
software, algorithm	nls.multstart	(Padfield and Matheson 2020)	https://github.com/padpadpadpad/nls.multstart	N/A

1061

1062

Contact for reagent and resource sharing

1063

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the corresponding Author, Anne A. Schmidt (anne.schmidt@pasteur.fr).

Zebrafish husbandry

Zebrafish (Danio rerio) of the AB background and transgenic fish carrying the following

transgenes *Tg(kdrl:ras-mCherry)* (Chi et al. 2008); *Tg(kdrl:nls-mKate2)*;

Tg(kdrl:Gal4;UAS:RFP); Tg(kdrl:eGFP) (Jin et al. 2005); Tg(CD41:eGFP) (Lin et al. 2005); and

the fish lines generated in this study: Tg(kdrl:Gal4;UAS:RFP;4xNR:eGFP-podxl2);

Tg(kdrl:mKate2-Podxl2); Tg(kdrl:eGFP-Jam2a); Tg(kdrl:eGFP-Jam3b);

Tg(kdrl:Gal4;UAS:RFP;4xNR:dt-runx1-eGFP); Tg(ArhGEF11[CRISPR-Cterdel-/+]) were raised and

staged as previously described (Kimmel et al. 1995). Adult fish lines were maintained on a 14

 hr light/10-hr dark cycle. Embryos were collected and raised at either 28.5 or 24°C in N-

Phenylthiourea (PTU, Sigma Aldrich, Cat# P7629)/Volvic source water (0.003% final) to

prevent pigmentation complemented with 280 μg/L methylene blue (Sigma Aldrich, Cat#

M4159). Embryos used for imaging, extracting mRNA, or for WISH ranged from developmental

stages 28-to-60 hpf and larvae used for imaging were of 3-5 dpf, precluding sex determination

of the animals. The fish maintenance at the Pasteur Institute follows the regulations of the

2010/63 UE European directives and is supervised by the veterinarian office of Myriam Mattei.

mRNA extraction and cDNA synthesis

Total RNA was extracted from whole 48 hpf embryos for cDNA cloning or from pooled trunks

at the desired developmental stages (30-32 hpf, 48-50 hpf time windows for qRT-PCR

experiments) (~30 individuals per tube). Briefly, embryos were anesthetized using balneation

in embryo medium supplemented with tricaine methanesulfonate (MS-222, Sigma-Aldrich Cat#

A5040), at a final concentration of 160 μg/ml. RNA was extracted via organic extraction using

TRIzol reagent (Invitrogen Cat# 15596026) according to the manufacturer's guideline. gDNA

contaminant was removed using TURBO™ DNase (Invitrogen Cat# AM2238) treatment

according to the manufacturer's guideline. Total RNA was stored at -80°C. Reverse

transcription of mRNA was performed using SuperScriptIV (Invitrogen Cat# 18090010) with

OligodT primer. cDNA samples were stored at -20°C until further processing (for quantitative

real-time PCR, PCR, cloning and sequencing).

Transient and stable transgenesis

The *kdrl* promoter (*flk*) (Jin et al. 2005) was used to drive endothelial expression of mKate2-

podxl2, eGFP-Jam2a and eGFP-Jam3b, with eGFP (Clontech) and mKate2 (Evrogen) cDNAs

amplified using overlapping primers for Gibson cloning. eGFP-podxl2, ArhGEF11(PDZ-PRD-

RGS-eGFP) and dt-runx1-eGFP were amplified for cloning into a pG1-4XNR vector (built from

the 4XNR (non-repetitive) 4X UAS sequence less susceptible to methylation than the 14X UAS

described in (Akitake et al. 2011)). For the sequence of all the designed cloning primers, see

the Key Resources Table.

Podocalyxin-like 2 (Podxl2), ArhGEF11/PDZ-RhoGEF, JAM2a and JAM3b full length

sequences were amplified from pools of 48 hpf whole embryo cDNA (see before for mRNA

extraction and cDNA synthesis methodologies). All constructs generated in this study were

obtained using the Gibson assembly assay (NEB, Cat # E2611S).

For eGFP-Jam2a and eGFP-Jam3b cloning, we used the JAM2a signaling peptide sequence

(included in the 5-prime amplification primers and encoding for the signal peptide:

mlvcvsllilihsvpvspvtvssr) and, for both constructs, the eGFP sequence was inserted in frame

upstream of the sequence encoding for the transmembrane domain; for the eGFP-Jam2a

construct, the eGFP was inserted upstream of amino-acid D221 (DLNVAA) of the NCBI

Reference Sequence: NP_001091734.1; for the eGFP-Jam3b construct, the eGFP was

inserted upstream of amino-acid D243 (DINIAG) of the NCBI Reference Sequence:

NP_001076332.2. For transient ubiquitous and inducible expression, the eGFP-Jam2a and

eGFP-Jam3b fusion constructs were inserted into a KpnI pre-digested pG1 plasmid containing

tol2 sites and the HSP70-4 promoter. For stable transgenesis using the *kdrl* (*flk1*) promoter,

constructs cloned into the pG1-HSP70 plasmids were extracted using XhoI/NotI restriction

enzymes and inserted into the XhoI/NotI pre-digested pG1-flk1 vector using DNA T4 ligation

(NEB, Cat # M0202S).

For the Podxl2 N-ter deletion mutant that has been used for establishing stable *Tg* fish lines,

the cDNA encoding for the peptide sequence starting with amino-acid G341 (GGTEYL;

fragment 341-587) of the NCBI Reference Sequence: XP_692207.6 was fused with either

eGFP or mKate2 cDNAs. The sequence encoding for the signal peptide of the human CD8

(malpvtalllplalllhaarpsqfrvs) was introduced upstream (5-prime) of the eGFP or mKate2

sequences (included in the 5-prime primers designed for the Gibson cloning strategy). For the

construct designed to obtain the *Tg(kdrl:Gal4;UAS:RFP;4xNR:eGFP-podxl2)* fish line, 2

overlapping fragments were amplified for cloning into a pG1-tol2_4xNR vector after digestion with NcoI, with the following couples of primers: 4XNR_CD8-eGFP_fw and delPodxl2_eGFP_rev; delPodxl2_fw and Podxl2_pG1-4XNR_rev. For the construct designed to obtain the *Tg(kdrl:mKate2-Podxl2)* fish line 2 overlapping fragments were amplified for cloning into a pG1-flk1 vector after digestion with EcoR1, with the following couples of primers: pG1-flk1_CD8_mKate2_fw and delPodxL2-mKate2_rev; mKate2-delPodxL2_fw and Podxl2_PG1-flk1_rev.

For the ArhGEF11(PDZ-PRD-RGS-eGFP) construct, the cDNA encoding for the peptide ending upstream of the DH domain at Valine 709 of the amino-acid sequence (GenBank: AY295347.1; PALDEDV; fragment 1-709) was fused upstream of the sequence encoding for eGFP, for expression using the pG1-4xNR vector. The cDNA was amplified using the GB_4xUAS-gef11_fw; GB_eGFP-RGS_rev couple of primers.

For the construct designed to obtain the *Tg(kdrl:Gal4;UAS:RFP;4xNR:dt-runx1-eGFP)* fish line, the Runx1 sequence amplified omits the ATG (replaced by an nls) and ends downstream of the Runt domain (ending at amino-acid sequence PAHSQIP). The construct allows for the expression of the dt-runx1 protein under the control of the 4xNR driver using the same strategy as for the above PodxL2 constructs. The dt-runx1 cDNA construct was amplified from the zebrafish Runx1 cDNA sequence (Kalev-Zylinska et al. 2002) using the couple of primers GB-4xnr-nlsRunx1_fw; GB-eGFP-T2A-2xHA-delRunx1_rev. The sequence of these primers each allowed for the introduction, in the protein sequence, of a nls (encoded by the fw primer) and of a 2xHA-T2A peptide (encoded by the rev primer that overlaps with the 5-prime sequence of eGFP).

For all constructs, cloning reactions were used to transform NEB 5-alpha Competent E. coli (NEB, Cat # C2987) and resistant clones then used for miniprep plasmid preparations (Nucleospin plasmid, Macherey-Nagel, Cat # 740588.50), followed by sequencing. Plasmid DNA from clones containing correct insertions where then used to transform E. coli TOP10 competent cells (Invitrogen, Cat # C404010) and purified using the endotoxin free NucleoBond

Xtra Midi kit (Macherey Nagel, Cat# 740420.10). Transgenesis was then performed by co-injecting 1 nl of plasmid (at 25 ng/μl) mixed with tol2 transposase mRNA (at 25 ng/μl) transcribed from linearized pCS-zT2TP plasmid using the mMESSAGE mMACHINE SP6 kit (Ambion, Cat# AM1340). For stable transgenesis, embryos were screened for fluorescence between 24 hpf and 48 hpf and raised to adulthood. Founders with germline integration were then isolated by screening for fluorescence after outcrossing with AB adults and for establishment of the F1 generation.

CRISPR methodology and GEF11 mutant transgenic line screening

CRISPR/Cas9 mutagenesis and isolation of founders

A sgRNA was designed against the splice donor site at the end of exon 38 with the aim to interfere with the production of the splicing variant including its encoding sequence, using the CRISPRscan web tool (Moreno-Mateos et al. 2015). The sgRNA was obtained by annealing for 5 min at 95°C equal volumes of 100μM specific Alt-R® crRNAs (IDT, Cat# sequence: AGCCAATCGTCTGAGGACGG) with 100μM Alt-R® generic tracrRNA (IDT, Cat# 1072532). Cas9 Nuclease Reaction Buffer (NEB Cat# B0386A) was added to obtain a final 45μM sgRNA stock solution. For the generation of the sgRNAs–Cas9 complex, a mix containing 18μM sgRNA and 12μM Cas9 protein (EnGen™ Spy Cas9-NLS S. pyogenes, NEB Cat# M0646T) was incubated for 10 min at room temperature. For mutagenesis, 1 nl of the sgRNA-Cas9 complex was injected into one-cell stage embryos, that were subsequently raised until 48 hpf. Bulk genomic DNA (gDNA) was extracted from 60 injected and 60 non-injected control embryos. gDNA extraction proceeded as follows: pooled samples were incubated for 3 hrs at 55°C in lysis buffer (100mM NaCl, 20mM Tris-HCl pH 8, 25mM EDTA, 0.5% SDS and 2 µg/µl proteinase K), purified using phenol:chloroform. Precipitation was done in 0.1 volume of 3M NaAc and 2.5 volume of 100% ethanol for 20 min at -20°C. Samples were centrifuged for 30 min at 4°C before washing with 70% ethanol and resuspension in water. gDNA was used for PCR, using DNA polymerase Platinum™ SuperFi™ (Invitrogen Cat# 12351010) and the

SM31-SM32bis couple of primers (TTTCACTTTCTCTCTGCCTCTTACA;

AACTCTGTCCAGATGATTGAGGAGC) flanking the targeted region. PCR products were run

on gel electrophoresis before gel extraction of bands of interest (with sizes corresponding to

either a wild-type allele or a deleted allele) (Nucleospin gel and PCR clean-up, Macherey-

Nagel Cat# 740609.50). Fragments were cloned into blunt-end TOPO vector (Invitrogen Cat#

45024), transformed into E. coli TOP10 competent cells and grown on selective medium. 12

colonies for each PCR product were sent for sequencing. Mutant sequences contained either

a 7 nucleotides deletion, or a mix of 1,4 or 7 nucleotides deletions located at the same site (at

the end of exon 38, see **Figure 7 - figure supplement 4A** for the 7 nucleotides deletion),

attesting of the proper functioning of our guide.

A stable homogeneous mutant line was generated by injecting the sgRNA:Cas9 complex into

one-cell stage *Tg(kdrl:Gal4;UAS:RFP)* embryos similarly to what was described above. 100

embryos were raised until adulthood. Adult founders were identified by crossing adult F0 with

AB fishes, and single embryo genotyping was performed on their progeny (10 embryos per

adult fish). Briefly, single embryos were incubated in 40μl 25mM NaOH 0.2mM EDTA and

heated for 10 min at 95°C, before cooling for 10 min at 4°C and addition of 40μl 40mM Tris-

HCl pH5 to stop the reaction. PCR was performed using 5μl of template and specific primers

(couple SM31-SM34 –TTTCACTTTCTCTCTGCCTCTTACA;

ATAAATGAAGCCCCACCTCCGTCC – for wild-type (WT) allele and couple SM31-SM40 –

TTTCACTTTCTCTCTGCCTCTTACA; ATGAAGCCCCACCTCAGACGATTGGC– for mutant

allele). Presence of WT or mutant allele for each single embryo was assessed by the

presence/absence of amplification band on gel electrophoresis. Four F0 founders were

isolated, that had a transmission rate in the germline ranging from 20% to 80%. Progeny of the

F0 founders and subsequent generations were raised separately until adulthood, and

genotyped. After complete anesthesia (in fish water supplemented with 160μg/ml tricaine

methanesulfonate), a small tissue sample was collected from the caudal fin, placed into DNA

extraction buffer (10mM Tris pH8.2, 10mM EDTA, 200mMNaCl, 0.5% SDS, 200µg/ml

proteinase K) and incubated 3 hrs at 50°C. DNA precipitation and resuspension was performed

as described above. 7µL gDNA was used as template for PCR amplification using the primer

couple SM31-SM32bis. Sequenced PCR products showed that all founders harbored the same

7 nucleotides deletion (see **Figure 7 - figure supplement 4A**). All experiments relative to

Figure 7 *and* **Figure 8** were performed using the progenies of one heterozygous mutant F3

couple. For these experiments, genotyping on single embryos was performed on gDNA as

mentioned above. Only WT homozygous and mutant homozygous embryos were kept for

analysis.

Alternative Cpf1 approach

As an alternative approach to the CRISPR/Cas9 deletion, we attempted to generate a second

mutant line, using to CRISPR/Cpf1(Cas12a) system to delete a 315 nucleotides region

encompassing exon 38 (with specific guides located in the introns before and after exon 38).

The sequence of guides that have been tested are: TATCACACACACATCACCTTCTA and

TTTCTCAGCGCTCCTGACAGATG. However, due to the structure of the intronic regions

(repeat rich) and the necessary presence of specific PAM sequence, we only generated off-

target deletions of intronic regions, without deletions of exon 38, leading us to focus on the

CRISPR/Cas9 approach.

1223

Morpholinos and injections

Morpholinos were obtained from GeneTools (see Key Resources Table for sequences). The

sih Tnnt2 translation start codon and flanking 5-prime sequence MO (Sehnert et al. 2002) as

well as the ArhGEF11 exon 38 splice blocking MO (overlapping the exon 38/intron 38-39

boundary) were resuspended in ddH2O to obtain stock solutions at 2mM and 1mM,

respectively. The *sih* MO (1.5ng) and ArhGEF11 exon 38 MO (3ng) were injected into 1-cell

stage zebrafish embryos, after dilution in ddH2O. For *sih* morphants, embryos that were used

in the experiments were checked for absence of heart beating 24 hrs after injection as well as

before being used for dissection and RNA extractions at the 30-32 and 48-50 hpf time-

windows. For measuring the efficiency of the ArhGEF11 exon 38 MO, total RNA was extracted

from pools of injected and control embryos followed by mRNA reverse transcription. PCR on

cDNA flanking the exon 38 was performed using DNA polymerase Platinum™ SuperFi™

(Invitrogen Cat# 12351010) and analyzed on gel electrophoresis before extraction, cloning and

sequencing to verify the absence of the DNA sequence encoded by exon 38 (see **Figure 7 -**

figure supplement 4A).

1239

Whole mount single molecule fluorescent in situ hybridization (RNAscope)

48 hpf *Tg(kdrl:eGFP)* and *Tg(kdrl:Gal4;UAS:RFP;4xNR:dt-Runx1-eGFP)* embryos were fixed

in 4% formaldehyde (Electron Microscopy Sciences, Cat#15712) diluted in PBS/0.1% tween20

(PBST) for 2.5 hrs at room temperature, rinsed with PBST and kept in 100% methanol at -20°C

until use. On the day of the experiment, embryos were rehydrated upon sequential incubations

in 75%/50%/25% MeOH/ddH20 (each for 10 min at room temperature (RT)). Embryos were

then incubated for 10 min in PBST. In subsequent steps of the RNAscope procedure, the

Multiplex Fluorescent Reagent Kit v2 was used (Cat# 323100), that included $H2O2$; probe

diluent (PD); wash buffer (WB); AMP1, AMP2, AMP3 buffers; HRP-C1, HRP-C2, HRP-C3

reagents; TSA buffer; HRP blocker. All incubations were performed in Eppendorf tubes, with a

maximum of 20 embryos/tube, in a dry heating block. Embryos were then incubated with $H2O2$

for 10 min at RT, washed for 15 min in PBST at RT, incubated for 7 min at 37°C with proteinase

K in PBST (1/2000 from a glycerol stock at 20mg/ml, Ambion, Cat#10259184), washed in

PBST for 2x10 min at RT, and incubated for at least 2 hrs at 40°C in PD. Embryos were then

incubated overnight at 40°C with the Pard3aa-C2, Pard3ab-C3, Pard2ba-C2 RNAscope

probes.

Next morning, embryos were washed for 2x10 min in WB. From that step on, extreme caution

should be taken because embryos will become totally transparent, owing to the harsh

conditions of the WB (we recommend that all washing steps, upon buffer removal, are

performed under a binocular loupe). Embryos were then incubated sequentially with AMP

buffers during 30 min for AMP1 and AMP2 and 15 min for AMP3, at 40°C, with 10 min washes

at RT between each AMP buffer incubation. Embryos were then incubated for 15 min at 40°C

with RNAscope Multiplex FL v2 HRP-C2 for Pard3aa and Pard2ba probes, HRP-C3 for the

Pard3ab probe, washed for 10 min at RT with TSA buffer, incubated for 30 min at 40°C with

OPAL-570 (PNFP1488001KT) in TSA buffer (dilution 1/500), washed 2x10 min at RT in WB,

incubated for 15 min at 40°C in HRP blocker and finally washed 2x10 min in WB and 2x10 min

in PBST at RT. Embryos were then embedded in 1% agarose and imaged by confocal

microscopy, ideally immediately and no more than 2 days after mounting. Since eGFP can

maintain its fluorescence property after fixation, Pard3 RNAscope spots could be relatively

easily superposed to the fluorescence of eGFP expressed in endothelial cells.

Images need to be analyzed with 3D-rendering (for example, with Imaris) to ensure the

localization of RNAscope spots in a given cells.

1272

Whole mount chromogenic in situ hybridization

For the sequence of all the primers used to amplify probes (relative to PDZ-RhoGEFs, ***Figure***

7 – figure supplement 2), see the Key Resources Table. Whole-mount chromogenic in situ

hybridization was performed as described in Lancino et al. 2018. Probes were synthetized

using the manufacturer recommendation (T7 RNA polymerase, Promega, Cat# P2075, DIG-

nucleotides, Jena Bioscence, Cat# NU-803-DIGX).

Images were captured with the Zeiss Axio ZOOM V16 microscope with the Zen Pro2 software,

with a brightfield transmitted light optics. Post-processing steps were performed using the

Extended-Depth Focus method to combine in focus regions from multiple z-planes and convert

into in a transmitted light z-stack to generate a unique in-focus image.

1283

Immunofluorescence detection of the HA-tagged dt-runx1 deletion mutant

Whole mount immunostaining was performed on hand-dechorionated 48 hpf *Tg(dt-runx1)*

embryos to assess the localization of the 2XHA-tagged dt-runx1 protein in aortic cells. Briefly,

embryos were fixed in 4% methanol free PFA (Polysciences Cat# 040181) for 3 hrs at room

temperature (RT) and washed in PBS/0.1% tween20 (PBT). Embryos were treated in

successive baths of milliQ water supplemented with 0.1% tween20, then in cold acetone for

10 min at -20°C and again in milliQ water plus 0.1% tween20. They were then washed in 1X

HBSS (Invitrogen Cat# 14025), and permeabilized for 45 min at RT in 1X HBSS, 0.1%

tween20, 5mM CaCl2 and 0.1mg/ml collagenase (Sigma-Aldrich Cat# C9891). Embryos were

then rinsed in PBSDT and incubated in blocking solution 1× sheep serum (SS, Sigma Cat#

S2263) for at least 4 hrs at RT. They were then incubated overnight at 4°C with primary

antibodies in blocking solution: rabbit anti-GFP (MBL Cat# 598; 1/300) and mouse anti-HA

(Sigma, Cat# 12ca5; 1/50). On the next day, embryos were washed several times in PBSDT

and endogenous peroxidase activity was inactivated by treatment with 6%H2O2 (Sigma-

Aldrich, Cat# H1009) for 1 hr at 4°C. Embryos were then incubated in the NGS for at least 4

hrs and incubated overnight at 4°C with secondary antibodies: goat anti-rabbit Alexa Fluor 488

(Invitrogen Cat# A11070; 1/400) and goat anti-mouse HRP-conjugated (Thermo Fisher

Scientific Cat# G-21040; 1/300). Finally, embryos were rinsed several times first in PBSDT and

after in PBT before HRP fluorescent revelation. Embryos were incubated for 45 min at RT in

the dark in imidazole buffer (0.1 M imidazole (Sigma-Aldrich, Cat# I5513) in PBT supplemented

with 1% H2O2) with Cy3 Tyramide Reagent (Cy3 NHS sigma Cat# PA13101, tyramide sigma

Cat# T-2879, dimethyl formamide sigma Cat# T-8654 and triethylamine sigma Cat# T-0886).

Final washes in PBT and in 6% H2O2 for POD inactivation were performed before embryos

were mounted in low-melting agarose in 1X PBT for fluorescence confocal imaging.

1308

1309   **Quantitative real-time PCR gene expression analysis**

1310   *Preparation of samples*

In a first line of experiments, expression of pard3 family genes by qRT-PCR as well as of some hematopoietic markers myb and runx1 was evaluated in our tissue of interest at 48 - 50hpf. Total RNA was obtained from FACS sorted vascular cells of control embryo using an incross of *Tg(Kdrl:gal4;UAS:RFP)* as well as non-vascular cells (as described below). Cells were collected from either whole embryos or from dissected trunks. Subsequently, to determine the impact of runx1 interference on pard3 family genes, total RNA was obtained from dissected trunk regions of 48-50hpf embryos from either incross of control *Tg(Kdrl:gal4;UAS:RFP)* or mutant *Tg(Kdrl:dt-runx1-eGFP)* fish lines. Analyses were carried on 3 biological replicates.

Cell dissociation and Fluorescent Activated Cell Sorting

Single cell suspension for FACS sorting was prepared using an optimized protocol adapted from multiple sources (Manoli and Driever 2012; Samsa et al. 2016; Bresciani, Broadbridge, and Liu 2018). All steps were performed with cooled solutions (4°C), and samples were kept on ice throughout the processing to preserve the viability of the cells. Briefly, embryos were anesthetized using balneation in embryo medium supplemented with tricaine methanesulfonate (MS-222, Sigma-Aldrich ref #A5040), at a final concentration of 640 μg/ml. Embryos were placed into a petri dish, rinsed in PBS supplemented with tricaine, and a majority of the medium was removed to prevent movement. Embryos were cut using a needle, and were then pooled in tubes and placed on ice until cell dissociation. Embryos trunks or whole embryos (around 120 or 60 respectively) were washed twice in PBS before centrifugation (300 x g for 1 min @4°C) and the supernatant was discarded. Embryos were incubated in 1ml TrypLE medium (Life Technologies ref #12605-010) for 10 min on ice, with gentle pipetting every 3 mins, first with a P1000 pipette and subsequently with a P200 pipette. Samples were then centrifugated for 7 min at 300g to remove the TrypLE and the cells were resuspended in 500μl FACSmax medium (Genlantis ref#T200100). Ultimately, samples were passed through a 40μm cell strainer moistened with FACSmax buffer onto a 35 mm cell culture dish using a syringe plunger. The cell strainer and the dish were washed with 300μl of FACS max and the flow-through was transferred to a FACS tube and kept on ice until sorting.

Cell sorting was performed on a BD FACS AriaIII cell sorter. For all samples, gating was done

on SSC-A vs. FSC-A to collect cells, and then on FSC-A vs. FSC-H to keep only single cells.

For control embryos, RFP positive cells as well as RFP negative cells were collected. For dt-

runx1 expressing mutant, all GFP positive cells (both RFP positive and RFP negatives) were

collected, as well as RFP single positives cells (not used for qPCR) and double negative cells

as internal controls. FASC sorted RFP single positives cells from dt-runx1 expressing mutants

were not used as control for the study of gene expression for 2 reasons. First, due to the

mosaicism of the line, very few cells were RFP single positive compared to the GFP positive

cells. Second, RFP single-positive cells from dt-runx1 mutant embryos would have been

indirectly exposed to runx1 interference (they would have been potentially in direct contact with

their neighbouring endothelial cells that express the dt-runx1 mutant) and thus would not be a

true no interference condition.

RNA extraction from FACS-sorted cells

Total RNA was extracted from sorted cell fractions using the RNeasy Qiagen kit (Qiagen, Cat

74104). Briefly, cells were sorted directly into $100\mu l$ lysis buffer, and RNA extraction was

done immediately according to the manufacturer's instructions. RNA was eluted twice with $15\mu l$

RNA free H2O. Reverse transcription of mRNA was performed using SuperScriptIV (Invitrogen

Cat# 18090010) with OligodT primer. cDNA samples were stored at -20°C until further

processing (for quantitative real-time PCR).

Real-Time qPCR

For the sequence of all the designed qRT-PCR primers, see the Key Resources Table. Primer

couples were individually tested to assess their specificity as well as their efficiency. qRT-PCR

was performed using Takyon Rox SYBR 2x Master mix blue dTTP kit (Eurogentec Cat# UF-

RSMT-B0701), with starting concentrations of template of around $10ng/\mu l$ and primer

concentrations of $0.5\mu M$, on a QuantStudio3 system (Applied Biosystems). Each reaction was

performed in technical triplicates and in 3 biological replicates. Ct was determined

automatically by the QUANTSTUDIO™ DESIGN & ANALYSIS 2 software on the Thermofisher

cloud and exported to be analyzed manually. Standard deviation was calculated within the

technical triplicates, and when superior to 0.3, obvious outliers were removed (Dixon's Q test).

The delta-delta-Ct method (double normalization to endogenous control — zebrafish

elongation factor ef1α — and to control sample) was used to compare gene expression in

control and altered conditions. The delta-Ct method (normalization to endogenous control ef1

α) was used to investigate expression of genes of interest in whole embryo and trunk vascular

cells versus non-vascular cells.

1372

In vivo confocal imaging

Embryos, dechorionated manually, were anesthetized using tricaine (Sigma Aldrich, Cat#

A5040). They were then embedded on the side position in 1% low melting agarose (Promega,

Cat# V2111) in a glass bottom 60μ-Dish (35mm high; Ibidi, Cat# 81156). To avoid movements

and pigmentation during image acquisitions, 1x tricaine /1x PTU Volvic water was incorporated

to the low melting agarose and, after solidification, 1ml of 1x tricaine /1x PTU Volvic water was

added before sealing the dish.

Embryos were imaged with 2 confocal microscope systems. For the results presented Figure

1, z-stacks and time-lapse sequences were obtained using a Leica TCS SP8 inverted confocal

microscope as described in Lancino et al 2018.

For all the other confocal microscopy results, embryos were imaged using an Andor (Oxford

Instruments) spinning disk confocal system (CSU-W1 Dual camera with 50μm Disk pattern

and single laser input (445/488/561/642nm), LD Quad 405/488/561/640 and Tripl 445/561/640

dichroic mirrors), equipped with a Leica DMi8 fluorescence inverted microscope and CMOS

cameras (Orca Flash 4.0 V2+ (Hamamatsu)). Imaging was performed using a 40x water

immersion objective (HC PL APO 40x/1.10 Water CORR CS2), a LEDs light source (CoolLED

pE-4000 16 wavelength LED fluorescence system), and the acquisitions were piloted with the

support of the MetaMorph software.

RNAscope is situ hybridization were imaged similarly to live sample, using a spinning disk

confocal microscope (see in vivo confocal imaging paragraph above) after embedding the

embryos in 1% low-melting agarose (Promega, Cat# V2111) in PBS 1X in a glass bottom 60μ-

Dish (35mm high; Ibidi, Cat# 81156).

1395

Morphological and morphometric analysis of aortic and hemogenic cells

For the morphometric analysis of the aorta and aortic cells, large z-stack of 48-55 hpf

Tg(kdrl:eGFP-JAM3b; kdrl:nls-mKate2) or Tg(kdrl:eGFP-JAM2a; kdrl:nls-mKate2)

encompassing the whole aortic depth were acquired with optimal z-resolution (0.3μm z-step).

For each embryo, 3 contiguous z-stacks of 330μm width were acquired, allowing us to image

the entirety of the AGM region, from the anterior to the posterior end of the elongated yolk.

Samples

For ArhGEF11 splicing morpholino experiment, 3 splicing morpholino injected embryos and 2

control non injected siblings were kept for the analysis.

For ArhGEF11 CRISPR Cter deletion and when performing the experiments, imaging was

done, for putative mutants, on embryos that had shown a delay in initiation of circulation at 24

hpf, regained circulation at 48 hpf albeit with a clearly visible oedema in the cardiac region

(non-altered embryos were kept as sibling controls). All imaged embryos whose signal allowed

us to generate segmentable 2D-cartography from the whole length of the aorta were kept and

processed further for genotyping. Genotyping confirmed that the circulation problems at 24 hpf

and the remaining oedema at 48 hpf corresponded to embryos bearing the mutation.

For CRISPR mutant phenotype analyses, 2 mutant embryos and 2 wild type siblings were kept,

after genotyping.

Image analysis

The following image analysis was performed on Icy software.

For each segment, we generated a 2D-cartography of the eGFP-Jam signals at intercellular

contacts of aortic cells, using the Tubeskinner plugin, according to the protocol described in

Lancino et al 2018. Briefly, the aorta was rotated to face the lumen, a circular ROI fitting the

interior of the aorta was initialized by the experimenter on the first plane, and for each plane

sequentially, the algorithm will more accurately fit the ROI to the aortic perimeter and collect

the fluorescence intensity all along its longitudinal axis. The collected intensity was then

projected on a 2D-plane, where the Y-axis of the 2D-cartography corresponds to the perimeter

of the aorta, and the X-axis corresponds to the X-axis of the original z-stack.

Before further processing, a metadata table recapitulating the dates and conditions of

experiments for each embryo was generated, and a random label was assigned to rename

each acquisition. This allowed for the later part of the analysis, that involves semi-automated

segmentation and classification by the user, to be blind and avoid potential bias.

Each 2D-cartography was then manually pre-segmented using the ROI Polygon tool to draw

the contour of each aortic cell, using the original 3D z-stack as reference. Due to technical

limitation (decreased fluorescence signal collection on the farthest lateral side of the aorta),

only about 2/3 of the 2D-cartographies could be segmented (comprising the floor, the right

lateral side of the aorta and the roof). The manual segmentation was then improved using the

"Active Contour" plugin, that improved the fit of the cellular contour to the fluorescence signal

and slightly dilated the cell contours until the ROIs formed a joint paving. All morphometric

descriptors were extracted for each cell. The number of neighbours was automatically

generated for each cell by counting the number of touching adjacent ROIs.

Additionally, cells were manually classified into cell types according to morphometric

parameters using as reference the original 3D z-stack and in particular the nls-mKate2 nuclear

signal. EHT-undergoing cells were positioned on the floor of the aorta, have reduce antero-

posterior axis length, luminal membrane inward bending for EHT pol+ cells and luminal

membrane outward bending for EHT pol- cells, and a thickened nucleus. Hemogenic cells

whose criteria were established in Lancino et al 2018 (elongated in the antero-posterior axis,

with lateral membranes not trespassing the equatorial plane of the aortic wall), are positioned

on the floor of the aorta, systematically have a reduced width perpendicular to the blood flow

and, importantly, a thickened nucleus that protrudes out of the aortic plane. Finally, endothelial

cells have a flattened nucleus positioned high on the lateral side or on the roof of the aorta.

1447

1448 **Measurement of the aortic perimeter**

1449 An accurate measurement of the aortic perimeter was obtained using the z-stack generated

1450 for the aortic cells' morphometric analysis. Briefly, the z-stack was rotated to face the lumen of

1451 the aorta and a ROI fitting the aortic perimeter was drawn by the user and optimized by the

1452 "Cell Contour" plugin. The length of the perimeter was then extracted from the ROI statistics.

1453

Fluorescence Recovery After Photobleaching (FRAP) measurements and analysis

Measurement of junctional turn-over was performed on 48-55 hpf *Tg(kdrl:eGFP-JAM3b;*

kdrl:nls-mKate2) embryos using a Nikon Ti2e spinning disk microscope equipped with a

sCMOS camera (Photometrics, Prime 95B, pixel size 11μm, 1200×1200 pixels, QE 95%) and

a 40x water objective (Numerical Aperture 1.15, Working Distance 0.6mm, xy pixel size of

0.27μm). Embryos were mounted as previously described.

Samples

For ArhGEF11 splicing morpholino phenotypic analysis, 10 different embryos (injected with the

splicing morpholino) obtained from 3 separate experiments and 30 control embryos (non-

injected siblings or control morpholino injected siblings) obtained from 6 separate experiments

were analyzed.

For EHT analysis, 16 embryos over 4 separate experiments were analyzed.

For ArhGEF11 CRISPR/Cas9 Cter deletion phenotypic analysis, 8 mutant embryos and 12

wild type siblings over 3 separate experiments were analyzed.

Overall, the exact number of interfaces analyzed is written on each figure panel.

Acquisitions

For each acquisition, a wide two channels z-stack of the aorta was generated to capture the

whole aortic volume and generate a 2D-cartography of the aortic environment (Icy, plugin

Tubeskinner). This 2D-cartography of the cellular contours and the z-stack with the cellular

contours and nuclei were used jointly to identify un-ambiguously cellular and junctional types

and define one or multiple single point regions of interest (ROIs) for subsequent bleaching and

fluorescence recovery measurement.

The sequence of acquisitions before/after bleach focused only on the eGFP-Jam3b signal to

speed up the acquisition and improve the temporal resolution. The 5 steps of the acquisition

can be recapitulated as follows and we obtained/performed: (1) a single z-stack of 20μm

centered on the plane of focus, (2) a 30 sec time-lapse acquisition on the plane of focus to

measure average pre-bleach intensity with an image every second, (3) a high intensity

stimulation of the ROIs (25% 300ms 488 FRAP laser exposure, leading to the bleaching of the

Jam3b-eGFP signal), (4) a 3 min time-lapse acquisition on the plane of focus to measure fast

post-bleach recovery with an image every second, and (5) a 7 min time lapse acquisition of

8μm z-stack centered around focus plane with a 20 sec time interval to measure slow post-

bleach recovery. Each step was save separately as .nd2 files.

Signal quantification and recovery analysis

File were converted from .nd2 to .tiff using FIJI batch converter, scale factor 1.00, bilinear

interpolation, no average when downsizing, images read using bioformat.

To collect fluorescence intensity, ROIs for each stimulation point was manually drawn using

Fiji and named with an individual label using Fiji ROI manager. A metadata table was manually

curated with all information for each ROI (junction type, condition, date of experiment, embryo

number). An additional ROI was created to collect background intensity for subsequent

corrections. Acquisitions with clear embryo movement or drifting were either completely

removed from the dataset (~5% of the acquisitions) or only kept for early recovery analysis

(see below) if the drift only occurred in the last step (fifth) of the acquisition.

Fluorescence intensity for each ROI was collected and saved automatically as a .csv file for

each acquisition using a homemade Fiji macro.

All further analysis were performed using R for automatization.

Briefly, for each ROI: the fluorescent signal was normalized to the background signal ($I_{background}$

$_{normalized}$ = I_{roi} − $I_{background}$), then normalized to mean intensity pre-bleach

($I_{normalized}$= $I_{background\ normalized\ ROI\ post-bleach}$ / $I_{background\ normalized\ ROI\ pre-bleach}$), and finally scaled

($I_{full\ scaled\ normalized}$ = ($I_{normalized}$ − min ($I_{normalized}$))/ (1− min ($I_{normalized}$)).

A single exponential equation was used to fit the data using the nls:multistart package for

parameter optimization:

$F(t) = A*(1-e^{-t*\tau})$

With A the maximum amplitude of the curve, t the time, and τ a constant defining the growth

rate of the curve.

The fitted curve allowed to extract quantitative information describing the recovery of each type

of junction in different conditions: the mobile fraction of the bleached pool (corresponding to A,

the amplitude of recovery) and the time of half recovery, related to the speed of recovery, that

can be extracted from τ:

$T_{half-recovery}$ = ln(0.5)/(- τ)

A more precise description of the recovery speed was extracted from the observation of the

early time period (until 30s post bleach), were the fluorescence intensity increases uniformly.

A first-degree polynomial equation was fitted to the data (using lm() function from the stats

package), and the coefficient corresponding to the slope of the fitted curve was used as a

metric for speed of fluorescence recovery.

1518

RNAscope image analysis

Myb

Phenotypic analysis of the impact of the dt-runx1 mutant on hematopoiesis was evaluated by

investigating the *in situ* expression of myb, using the single-molecule *in situ* hybridization RNA-

scope method, performed in control *Tg(Kdrl:eGFP)* and dt-runx1 mutant embryos

Tg(Kdrl:Gal4;UAS:RFP;4xNR:dt-runx1-eGFP) (analysis was performed on 4 and 3 biological

replicates (embryos) for control and mutants respectively). For each embryo, 2 consecutive

aortic segments were analyzed (each spanning over 5-6 intersegmentary vessels). It is roughly

equivalent to the length to 3 consecutive aortic segments that we use for our live analysis (each

spanning over 3-4 intersegmentary vessel. Discrepancies between live and fixed samples are

probably due to the morphological alterations caused by the fixation and permeabilization

steps of the RNAscope protocol. 3D-rendering of the aorta and the sub-aortic space were

generated using the Imaris Software (Oxford Instruments, version 10.1.0). Both cell volume

(GFP signal) and RNAscope spots were simultaneously and automatically segmented using

the "Cells" tool from Imaris, that segment individual cells (generates an outer surface) as well

as RNA-scope signal within cells. Cells were subsequently manually classified, based on their

localization and the shape of the cell. Roof endothelial cells are located dorsally to the aortic

floor, and are flat and elongated. Due to morphological alteration of the tissues and the

impossibility to discriminate EHT pol-, EHT pol+ cells and the uncharacterized hemogenic cells

we observed *in vivo*, our hemogenic / EHT cells category comprises both characterized

hemogenic cells (elongated cells with thick nuclei, located on the aortic floor), uncharacterized

hemogenic cells and putative EHT-undergoing cells. Sub-aortic cells were classified based on

their localization, and they comprise a mix of endothelial cells from the underlying posterior

cardinal vein (visible only in the control embryos and not in the dt-runx1 mutant embryos, in

which arterial structures are more specifically labelled compared to veinous structures) as well

as newly generated hematopoietic cells. Finally, the "Statistics" tool from Imaris allowed us to

extract cell and RNA spots count information. The data plotting and statistical analysis was

performed using R software.

Pard3

Pard3 isoforms (Pard3ba, Pard3ab, Pard3aa) *in situ* expression analysis was done similarly to

that of myb, except that a second RNAscope signal segmentation was performed

independently of the "Cells" segmentation, using the "spots" tool. This allowed to capture the

majority of the Pard3 signal, that is expressed in aortic cells, but with a discrepancy in the intra-

cellular localization of RNA transcript, that could not be detected when restricting the

segmentation of RNAscope spots to spots located inside the GFP signal. After RNAscope

segmentation using the "Spots" tool, manual curation allowed us to select specifically

RNAscope spots located in aortic tissues for subsequent analysis. The "Statistics" and "object-

object Statistics" tools were used to extract relevant spatial and count information for

subsequent analysis. A key information that we used for the spatial analysis was the average

distance of Pard3ba spots to Hemogenic / EHT cells, as well as the cumulative number of

spots within a distance from Hemogenic / EHT cells, to investigation spatial correlation

between Hemogenic / EHT cells and Pard3ba expression. Analysis was performed in control

Tg(Kdrl:eGFP) and dt-runx1 mutant embryos *Tg(Kdrl:Gal4;UAS:RFP;4xNR:dt-runx1-eGFP)* (5

and 6 biological replicates (embryos) for control and mutants respectively, with 2 consecutive

aortic segments analyzed per embryo).

1564

Dt-runx1 phenotype analysis – cell count

For the live dt-runx1 phenotype analysis, large z-stack of 48-55 hpf dt-runx1 expressing mutant

(eGFP positive) and siblings control (eGFP negative) *Tg(Kdrl:Gal4;UAS:RFP;4xNR:dt-runx1-*

eGFP) embryos, encompassing the whole aortic depth were acquired with optimal z-resolution

(0.3μm z-step). 3 and 7 biological replicates (embryos) for control and mutants respectively

were analyzed. For each embryo, 3 contiguous z-stacks of 330μm width were acquired,

allowing us to image the entirety of the AGM region. The resulting phenotype was assessed

by manually counting EHT pol+ and pol- cells (taking into account their position, ventral or

lateral) as well as hemogenic cells. Classification was done manually, based on morphological

criterias. EHT pol+ cells presented the typical EHT pol+ morphology, with clear luminal

membrane invagination, that could be observed even in laterally emerging cells. EHT pol- cells

displayed a clear rounded ovoid, with a clear shortening of the antero-posterior axis.

Hemogenic cells correspond to endothelial cells located on the aortic floor, that are elongated

along the antero-posterior axis, have a narrowed left-right axis and a thickened nucleus. Non

characterized hemogenic cells were also included in this category: they correspond to cells

that had potentially already advanced in the emerging process but whose morphology did not fit any of the typical EHT pol- and pol+ characteristics (no clear invagination nor clear concave protrusion in the aortic lumen).

Thymus phenotype analysis

To question one aspect of the hematopoietic phenotype of our dt-runx1 mutant, dt-runx1 expressing mutants (GFP+ RFP-) as well as control siblings (RFP+ GFP-) were imaged at 5dpf to analyze their developing thymus *in vivo* (analysis was performed on 5 mutant and 5 control larvae). Subsequent image analysis was performed using the Imaris software (Oxford Instruments). Individual thymic cells were automatically segmented using the "Surface" tool. Cell count per thymus as well as morphometric information for each cell (volume) was automatically retrieve using the Imaris "Statistics" tool, allowing us to compare the distribution of cell volume in mutant versus control embryos, as well as the total cellular volume of the thymus (sum of individual cell volumes) for each larva.

Impact of ArhGEF11 exon38 splicing morpholino on hematopoiesis

After morpholino injection at the one cell stage, double transgenic *Tg(CD41:eGFP; kdrl:nls-mKate2)* embryos were raised until 48hpf and imaged using spinning disk confocal microscopy. The integrality of the dorsal aorta was imaged (corresponding to 3 contiguous segments), on 8 and 6 biological replicates (embryos) for controls and morphants respectively. Quantification of the number of hemogenic cells (double CD41:eGFP and kdrl:nls-mKate2 positive cells) as well as of hematopoietic cells (simple positive CD41:eGFP and kdrl:nls-mKate2 negative cells) was performed using a in house Fiji/R script. For the segmentation, only the eGFP channel is handled. An issue with the *Tg(CD41:eGFP)* background is the presence of circulating thrombocytes, that move faster than the speed of acquisition and thus generate multiple artefactual 2D objects (only presents in single Z-planes, compared to non-moving cells whose depth spans over 15-20 z-sections). These artefactual objects are removed during the pre-processing steps, by using a Faster Temporal Median algorithm

(Jabermoradi et al. 2022). This algorithm, by calculating median value over rolling windows of

20 frames approximate cell depth) and substracting it to the pixel values, removes the 2D

outliers. The rest of the pre-processing steps consists of fluorescent signal filters to ease the

subsequent segmentation (contrast enhancement, background substraction, 3D gaussian

blur). The cells are segmented using a TopHat algorithm and touching objects are separated

using a 3D distance Watershed (Legland, Arganda-Carreras, and Andrey 2016). Once the

objects are segmented, the total fluorescence intensity in both channels as well as the

background fluorescence intensities are collected using the raw images (without any signal

filters). The data are subsequently processed using R for plotting, threshold selection for

classification as well as statistical comparisons.

1617

Statistical analysis

Statistical tests used are described for each figure panel in the legend. Since our samples are

usually small, the test used is an unpaired two samples non-parametrical Wilcoxon mean

comparison test, without a priori knowledge on sample distribution or homoscedasticity. In the

context of FRAP experiments (relative to **Figure 6** and **Figure 8**), for which we had larger

sample size, we still decided not to assume that our data were normally distributed, because

of the inherent variability of the biological events that are observed. Indeed, the EHT process

is heterogeneous in time, with successive phases of contraction and stabilization, during which

we can imagine that the dynamics of junctional recycling could vary. The distribution of speed

and amplitude of recovery could thus correspond to a normal distribution, a bi- or multi-modal

distribution, or a uniform distribution. In this case, the use of non-parametrical tests and the

large variability we observed in the data reduces the power of statistical tests and our ability to

confidently detect significant differences. In some cases, when our p-values were larger than

the arbitrary threshold of 0.05, we refer to tendencies whose significance is legitimated by the

overall consistency of the data (see for example between the results presented **Figure 7** and

Figure 8).

Correlation analysis (relative **to Figure 4F**) was performed by calculating Pearson correlation

coefficient and associated p-value (Student T-test) using R.

1636

Data and software availability

All graphics and plots were generated using RStudio and packages including: ggplot2

(Wickham 2016), readr (Wickham, Hester, and Bryan 2022), dplyr (Wickham et al. 2023),

stringr (Wickham 2022), ggstatsplot (Patil 2022), ggpubr (Kassambara 2020), wesanderson

(Ram and Wickham 2018), cowplot(Wilke 2020), ggsci (Xiao 2018), ggbeeswarm (Clarke and

Sherrill-Mix 2017), viridis (Garnier 2021), DT (Xie, Cheng, and Tan 2022), SciViews (Grosjean

2019), nls.multstart (Padfield and Matheson 2020), matrixStats (Bengtsson 2022).

1644

1645

Acknowledgments

We wish to warmly thank Philippe Herbomel for his encouragements and for ensuring the

funding of this study.

We thank our fish facility members Y. Rolin and K. Sebastien for their daily help and

commitment. We thank Sandrine Schmutz and Gaëlle Letort for their expertise and invaluable

assistance. We acknowledge the help of the Cytometry and Biomarkers (UTechS CB) service

platform for expert assistance with FACS analysis, of the Image Analysis Hub, and of the

Photonic BioImaging (UTechS PBI) service platform of the Institut Pasteur for their respective

contributions.

We wish to thank R. Patient (University of Oxford) for the Hsp70 promoter. This work was

supported by Institut Pasteur, CNRS, grants to P. Herbomel from the Fondation pour la

Recherche Médicale (#DEQ20160334881), the Fondation ARC pour la Recherche sur le

Cancer, and the Agence Nationale de la Recherche Laboratoire d'Excellence Revive

(Investissement d'Avenir, ANR-10-LABX-73), and a grant to AS from the Institut Pasteur (PTR

1662

1663   Authors contributions

1664   LT designed, performed and analyzed experiments, and wrote the Method section and the

1665   Figure legends. SM designed and conducted experiments. CV conducted experiments. AS

1666   conceived the project, designed, performed and analyzed experiments, wrote the manuscript

1667   with input from LT, and obtained funding.

1668

1669   Declaration of interests

1670   The authors declare no competing interests.

Figure legends

Figure 1: Tracing the evolution of the luminal membrane using the polarity marker Podocalyxin points at the biological significance of apicobasal polarity establishment in EHT cell emergence complexity. A, cartoons depicting the early and late steps of EHT cells emerging from the aortic floor (steps 1 and 2, as previously described in the zebrafish embryo, see (Kissa and Herbomel 2010; Lancino et al. 2018)) and with hypothetical evolution of the luminal membrane (in green) before (3 and 4) and after the release (2', the cell detaches from the endothelial layer via junction downregulation leading to exposure of the luminal membrane with le extracellular milieu; 3', the luminal membrane is consumed via endocytic recycling (E) and/or lysosomal degradation (Lys) prior to detachment; 4', the luminal membrane in 4 is released inside the cell (twisted arrow) before detachment. Grey area= nucleus. **B**, PodocalyxinL2 (Podxl2) construct designed to establish transgenic fish lines. Cartoons representing full length (top drawing) and deleted Podxl2 (amino-acid sequence 341-587) in which the mucin domain (serine/threonine-rich O-glycosylation domain) is replaced by either eGFP or mKate2. The tetracystein-containing globular domain (subjected to N-glycosylation) was kept as favoring apical membrane retention. TMD, transmembrane domain; DTHL, C-terminal peptidic motif involved in partnership with PDZ domain containing proteins. **C-D**, EHT performing cells visualized using *Tg(Kdrl:Gal4;UAS:RFP;4xNR:eGFP-podxl2)* embryos and time-lapse sequences initiated at 55 hpf obtained with spinning disk confocal microscopy (imaging was performed at the boundary between the most downstream region of the AGM and the beginning of the caudal hematopoietic tissue). Top grey panels show the green, eGFP channels for eGFP-podxl2 only. Bottom panels show the merge between green and red (soluble RFP, in magenta) channels. Bars: 8μm. **C**, single plane images of 2 EHT pol+ cells extracted from a time-lapse sequence at t = 0 and t = 45 min, with the right cell (eht cell 2) more advanced in the emergence process than the left one (eht cell1). Note the enrichment

of eGFP-podxl2 at the luminal membrane (surrounding the cavity labelled with an asterisk) in

comparison to the basal membrane (white arrow). Note also the evolution of the luminal

membranes with time, with the aortic and eht cell 1 lumens still connecting at t = 0 (green

arrow), the apparent fragmentation of the cytosolic vacuole (2 asterisks for eht cell 1 at t = 45

min) and the compaction of Podxl2-containing membranes for eht cell 2 at t = 45 min. More

details on the evolution of the connection between the aortic/eht cell lumens are shown in

Figure 1 – figure supplement 1A. **D**, single plane images of 2 EHT pol- cells extracted from

a time-lapse sequence at t = 30 min and t = 210 min (see **Figure 1 – video 3** for the full-time

lapse sequence), with the right cell (eht cell 2) slightly more advanced in the emergence than

the left one (eht cell 1, with the latest attachment point between the emerging cell and the

aortic floor (pink arrow)). Note, in comparison with the cells in panel C, the ovoid shapes of

cells, the absence of enrichment of eGFP-podxl2 at luminal membranes (green arrows) and

the accumulation of eGFP-podxl2 at basal membrane rounded protrusions (white arrows).

1710

Figure 2: Immature HE is not polarized and controls membrane delivery of intra-

cytosolic vesicular pools. **A**, *Tg(Kdrl:Gal4;UAS:RFP;4xNR:eGFP-podxl2)* embryo imaged

using spinning disk confocal microscopy. Black and white images show eGFP-podxl2 only.

Images (single z-planes) were obtained from a time-lapse sequence (initiated at 35 hpf) lasting

for 435 min (7.25 hrs), with intervals of 15 min between each z-stack. Example of an HE cell

with equal partitioning of eGFP-podxl2 between luminal and abluminal membranes (at t = 0

min), with eGFP-podxl2 containing intra-cytosolic vesicles (one labelled with a green asterisk)

and undergoing mitosis at t = 30 min (HE cell 1' and HE cell 1'' are daughter cells). Note the

inheritance of the largest micropinocytic-like vacuole by HE cell 1' and its maintenance over

time until EHT emergence initiation at t = 180 min (green asterisk in 1.5x magnified areas at t

= 60 and 90 min). At t = 360 min (green box) EHT is proceeding and both fluorescence

channels are shown; bottom panel: green (eGFP-podxl2), magenta (soluble RFP). The

magenta arrow points at the basal side of the EHT pol + cell (EHT cell 1', on the right) that

does not contain any detectable eGFP-podxl2; on the contrary, eGFP-podxl2 is enriched at

the luminal/apical membrane (note that exocytosis of the large vacuolar structure may have

contributed to increase the surface of the apical/luminal membrane (the green asterisk is

surrounded by the apical/luminal membrane of the EHT pol + cell)). The green arrow points at

the abluminal membrane of the EHT cell derived from HE cell 1'' (EHT cell 1") and that contains

eGFP-podxl2 (with no evidence of a significant expansion of a luminal/apical membrane); this

indicates that this cell is more likely to be an EHT pol- cell that did not sort the vesicular cargo

to the luminal/apical membrane. Bar = 10μm. **B**, hypothetical model summarizing the evolution

of HE cells involving the tuning of apicobasal polarity to lead to cells competent for giving birth

to either EHT pol+ or EHT pol- cells (including the release of large vesicular macropinocytic-

like vacuoles preferentially toward the luminal membrane of future EHT pol + cells). The

polarity status of HE cells is proposed to evolve asynchronously throughout the entire EHT

time window, leading to place-to-place ability to give birth to EHT cells (emergence of EHT

pol+ and EHT pol- cells are both observed until 72 hpf, see main text).

1738

Figure 3: Interference with Runx1 function alters emergence efficiency and induces a

bias toward EHT pol+ cells. A, *Tg(kdrl:Gal4;UAS:RFP;4xNR:dt-runx1-eGFP)* mutant

embryos imaged using spinning disk confocal microscopy and analyzed in the AGM/trunk

region. Z-projections of the dorsal aorta obtained from 52 - 55 hpf embryos. **Top panel:**

fluorescence from the red channel is shown for the *Tg(Kdrl:Gal4;UAS:RFP)* control. **Bottom**

panels: fluorescence in the green channel only is shown for the mutants (eGFP, released from

the dt-runx1-eGFP cleavage). The black asterisks point at emerged cells that are in close

contact with the aortic floor. Green arrows: EHT pol+ cells on aortic floor; magenta arrows:

EHT pol+ cells in the lateral aortic wall; blue arrows: uncharacterized emerging cells; he:

hemogenic cells. **B,** quantitative analysis of the dt-runx1 mutant phenotype. The analysis was

carried   out   on   *Tg(Kdrl:Gal4;UAS:RFP)*   control   embryos   (n   =   3)   and   on

 Tg*(kdrl:Gal4;UAS:RFP;4xNR:dt-runx1-eGFP)* mutant embryos (n = 7). **Left top**: hemogenic

cell count (comprising hemogenic cells, labelled by "he" on panel **A**, as well as uncharacterized

emerging cells, labelled by blue arrows on panel **A**). **Left center**: EHT pol+ cell count (sum of

EHT pol+ cells emerging ventrally and laterally, respectively labelled by green and magenta

arrows on panel **A**). **Left bottom**: EHT pol- cell count (not shown on panel **A**). **Right**:

percentage of cell types in control and dt-runx1 embryos (proportions relative to total number

of hemogenic and EHT undergoing cells). Statistical comparisons have been performed using

two-sided unpaired Wilcoxon tests, all p-values are displayed. **C**, **top:** cartoons representing

the zebrafish full-length runx1a amino-acid sequence and the dt-runx1 mutant deleted from

the trans-activation domain and of the C-terminus (note that the construct encodes for a C-

terminal fusion with eGFP that is released upon expression via a cleavable T2A peptide

(introduced between the 2xHA tag and the N-terminus of eGFP, to prevent from potential steric

hindrance). nls, nuclear localization signal. **Bottom:** image of an anti-HA tag

immunofluorescence obtained after z-projection of the dorsal aorta of a 50 hpf *Tg(dt-runx1)*

embryo. Note the localization of the 2xHA-tagged dt-runx1 protein in nuclei (some of them are

pointed by red arrowheads) and of eGFP in nuclei and the cytosol of aortic cells. Bar: 25μm.

1766

Figure 4. Pard3ba expression is highly sensitive to interference with Runx1 activity. A,

Pard3aa, ab, ba and bb zebrafish isoforms, with their accession numbers. The cartoon

represents the common structures of the 4 isoforms, with the 3 sequential PDZ domains

(PDZ1-3), and the CR1 and CR3 conserved regions involved in oligomerization and atypical

protein kinase C (aPKC) binding, respectively. The magenta arrows show the positioning of

qRT-PCR primers, located in the PDZ1 domain (see ***Methods*** for primer sequences). **B**, qRT-

PCR analysis of genes expression levels of (**left**) the hematopoietic markers myb and runx1,

(**right**) the 4 isoforms of Pard3 proteins (Pard3aa, ab, ba and bb) in cell populations isolated

from FACS-sorted trunk vascular cells of 48 - 50 hpf control and mutant embryos (see ***Figure***

4 - figure supplement 1, ***Figure 4 - figure supplement 2*** and ***Methods*** for cell isolation

procedures). Graphs show the measured mean fold changes relative to the expression of ef1α

and to the expression in RFP+ control cells. Statistical tests: two sided unpaired two samples

Wilcoxon test, all p-values are displayed. Analysis was carried out on n = 3 for control and

mutant conditions. **C**, representatives images (Imaris 3D-rendering) of RNAscope *in situ*

hybridizations for Pard3ba in 48 - 50hpf *Tg(Kdrl:eGFP)* control embryos and

Tg(Kdrl:Gal4;UAS:RFP;4xNR:dt-runx1-eGFP) mutant embryos. The aorta is outlined with the

white dashed lines. The RNAscope signal was segmented into spots and classified based on

its localization: in aortic endothelial and hemogenic cells (magenta spots) or in extra-aortic

tissues (grey spots, essentially residing in the sub-aortic space). Magenta dashed boxes

delineate examples of mRNA Enriched Regions (MERs) in control and mutant embryos. Scale

bars: 10μm. **D**, Pard3ba spots count per aortic segment. **E**, Pard3ba spots count per

hemogenic/EHT cells. **F**, cumulative Pard3ba spots count relative to hemogenic/EHT cell

distance, averaged for each aortic segment, in control embryos (grey) and dt-runx1 mutant

embryos (orange). **G**, HE/EHT cell count relative to number of spots, all expressed per aortic

segment (**left**), 50% Pard3ba spots attraction distance (in μm) (**middle**) or 75% Pard3ba spots

attraction distance (in μm) (**right**). 50%/75% spots attraction distance corresponds to the

distance from hemogenic/EHT cells within which 50%/75% of Pard3ba spots are located. Grey

dash lines correspond to regression lines and light grey background correspond to the

confidence interval (95%) of the regression model. Linear correlation coefficient (R, Pearson

correlation) is displayed for each condition. For **D** and **E**, statistical tests: two sided unpaired

two samples Wilcoxon test, all p-values are displayed. Analysis was carried out on n = 5 control

embryos and n = 6 mutant embryos, 2 aortic segments per embryos. For **G**, p-values were

calculated using a t-test.

1800

Figure 5: eGFP-Jam3b localization is reinforced at antero-posterior sites of the

endothelial/EHT interface and at tri-cellular junctions. A, cartoons representing

homodimers of full-length JAMs with the C-terminal cytosolic part interacting with Pard3 (JAMs

interact with the first PDZ domain of Pard3) as well as the constructs generated in this study

and containing eGFP inserted between the Immunoglobulin-like (Ig) domains and the trans-

membrane region (TMD). The constructs were obtained for zebrafish JAM2a and JAM3b. **B,**

C, 52 hpf *Tg(kdrl:eGFP-Jam3b; kdrl:nls-mKate2)* embryos were imaged in the trunk region

(AGM) using spinning disc confocal microscopy. The panels are either maximum z-projections

(top two) or single plane z-sections (bottom two, focusing on the aortic floor) of aortic

segments, with either the merged nls-mkate2 and eGFP-Jam3b fluorescence signals

(magenta and green) or the eGFP-Jam3b signal only (black and white images). Bottom of the

Figure: 2D-cartographies obtained after deploying aortic cylinders and showing the eGFP-

Jam3b signals only. The white/black vertical dashed lines show the correspondence of the

antero-posterior junctional reinforcements in the single z-sections (white arrows) on the

respective 2D-cartographies resulting from the deployment of the aortic walls (obtained from

the z-stacks). **B**, example of an EHT pol+ cell (cell 1, white arrows point at reinforcement of

signal at antero-posterior junctions). On the 2D cartography, cell 1 (red) is contacting

endothelial cells 2 and 3; note the reinforcement of eGFP-Jam3b signals along antero-

posterior membrane interfaces perpendicular to blood flow (red arrows) as well as at the two

tri-cellular junctions visible between cells 1, 2 and 3 (black arrows). **C**, example of two EHT

pol- cells (cells 1 and 2, white arrows point at reinforcement of signal at antero-posterior

junctions). On the 2D cartography, cells 1 and 2 (red) are contacting endothelial cells 3, 4, 6

and 3, 6 respectively; note the reinforcement of eGFP-Jam3b signals along antero-posterior

membrane interfaces perpendicular to blood flow (red arrows) and endothelial cell 6 that has

intercalated between endothelial cell 7 and EHT pol- cell 2 (blue arrow). In right margins,

magenta and green arrowheads designate the aortic floor and roof, respectively. Bars = 10μm.

1827

Figure 6: Junctional recycling at tri-cellular contacts is differentially controlled between

the two EHT types. 48-55 hpf *Tg(kdrl:eGFP-Jam3b; kdrl:nls-mKate2)* embryos were imaged

using spinning disc confocal microscopy and illuminated for Fluorescence Recovery After

Photobleaching (FRAP) in the trunk region (AGM, Aorta Gonad Mesonephros). **A, B**, panels

are either maximum z-projections (top left) or single plane z-sections (bottom left and top right,

focusing on the aortic floor) of aortic segments, with either the merged nls-mkate2 and eGFP-

Jam3b fluorescence signals (magenta and green) or the eGFP-Jam3b signal only (black and

white images). White arrows point at reinforcement of signal at antero-posterior junctional

pools of an EHT pol+ cell (**A**) or of an EHT pol- cell (**B**), both marked by asterisks. Bottom right:

2D-cartographies obtained after deploying aortic cylinders and showing the eGFP-Jam3b

signals only. Black arrows point at antero-posterior junctional pools, in particular at tri-

junctional regions that exhibit increase in signal density (well visible in **A,** black arrows). 2 and

3 endothelial cells are contacting the EHT pol+ cell (**A**) and the EHT pol- cell (**B**), respectively.

In right margins, magenta and green arrowheads designate the aortic floor and roof,

respectively. Bars = 20μm. **C - G,** FRAP analyses. EGFP-Jam3b junctional pools

corresponding to the brightest spots inside junctional regions of interest were bleached for

FRAP measurements (these high intensity pools were localized at the level of bi- and tri-

junctions for endothelial cells (EC) and in tri-junctional regions for EHT pol+ and EHT pol- cells;

all these junctional pools were systematically visualized by deploying each aortic segment

before bleaching as shown in the 2D-cartographies in A and B as well as in ***Figure 6 – figure***

supplement 1, see also ***Methods***). FRAP analysis concerned 3 types of junctional interfaces:

between endothelial cells (EC – EC, black and grey), EHT pol- and endothelial cells (pol- –

EC, brown), EHT pol+ and endothelial cells (pol+ – EC, blue). **C, D**, evolution of mean

fluorescence intensity for each type of junctional interface over time (10 min), after

photobleaching (t = 0s). **E**, median maximum amplitude of recovery of all determinations and

for each type of junctional interface (maximum of simple exponential fitted curves). **F, G**, early

fluorescence recovery. Early evolution (over the first 30s) of the mean fluorescence intensity

for each type of junctional interface over time after photobleaching (t = 0s). **F**, the fitted lines

correspond to linear regressions of mean fluorescence intensities. **G**, median values of

fluorescence recovery slopes (linear regressions) of all determinations and for each type of

junctional interface. **E, G**, the number of biological replicate (n) is stated on the plots. Statistical

tests: two sided unpaired two samples Wilcoxon test.

1860

Figure 7: Interfering with ArhGEF11/PDZ-RhoGEF function leads to the accumulation of

hemogenic cells and impairs EHT progression. A-C, numeration and morphometric

analyses of aorta and cell types for *Tg(kdrl:eGFP-Jam3b; kdrl:nls-mKate2)* ArhGEF11 exon

38 splicing morpholino-injected and control embryos (**A**), or for *(Kdrl:eGFP-Jam3b;*

*kdrl:*ArhGEF11CRISPR-Cterdel+/+) homozygous ArhGEF11 C-ter deletion mutants and control

siblings (**B**). 48-55 hpf embryos were imaged using spinning disk confocal microscopy. **Aa,**

Bb, 2D-cartographies obtained after deploying aortic cylinders and showing the eGFP-Jam3b

signals only with cell contours delineated either in blue (endothelial cells), yellow (hemogenic

cells, see ***Methods*** for their morphological definition), red (morphologically characterized EHT

cells, for controls), and small cells delineated by cyan boxes (morphologically uncharacterized

EHT cells and putative post-mitotic cells remaining as pairs, included in the numeration as

hemogenic cells). Cellular contours have been semi-automatically segmented along the

cellular interfaces labelled with eGFP-Jam3b (see ***Methods***). Scale bars: 10μm. **Aa', Bb'**, **left:**

numeration of endothelial, hemogenic and EHT-undergoing cells according to the position of

their geometrical center (either on the aortic floor, or on the roof, or on the lateral side), for

each condition; **right**: number of endothelial, hemogenic and EHT-undergoing cells in each

condition calculated from the segmentation of 3 x 2D-projections per embryo and covering the

entire aortic regions in the trunk. **Aa'', Bb''**, **left**: length of hemogenic cells (in the longest axis)

in function of their orientation (°) relative to the blood flow axis (0 – 180°); **right**: distribution of

the orientation of hemogenic cells relative to the blood flow axis, displayed as a mean

distribution of cells per embryo. **Aa''', Bb'''**, hemogenic cell elongation factors in arbitrary Units

(scale factor given by the ratio between the first- and the second-best fitting ellipse diameters,

the minimum value being 1 for a non-elongated object) represented as boxplot distribution of

all segmented cells (**left**) or as the distribution of cell elongation factor per embryo (**right**), for

controls and for interfering conditions as indicated. **Aa''''', Bb'''''**, hemogenic cell area

represented as boxplot distribution of all segmented cells (**left**) or as the distribution of cell

area per embryo (**right**), for controls and for interfering conditions as indicated. **C**, aortic

perimeter (in μm) for controls and mutant conditions as indicated. Statistical tests: two sided

unpaired two samples Wilcoxon test. For the ArhGEF11 exon 38 splicing morpholino

condition, analysis was performed on 2 x control (non-injected embryos) and 3 x embryos

injected at the one-cell stage; for the CRISPR mutant condition, analysis was performed on 2

x wild-type siblings for control and 2 x homozygous mutant embryos whose DNA mutation was

confirmed by sequencing. 3 consecutive aortic segments per embryo were analyzed to cover

the whole length of the dorsal aorta, in the trunk region (covering a distance of 990μm per

embryo).

1896

Figure 8: Interfering with the function of ArhGEF11/PDZ-RhoGEF suggests an activity

at the interface between endothelial and hemogenic cells that relies on restricting the

mobility of junctional pools at tri-cellular junctions, with a prominent role of the +exon

38 variant during endothelial cell intercalation. A, 2D-cartography of an aortic segment of

a *Tg(kdrl:eGFP-Jam3b; kdrl:nls-mkate2)* embryo injected at the one cell stage with the

ArhGEF11 exon 38 splicing morpholino. All the arrows pointing at reinforced junctional

contacts between cells were bleached and imaged for FRAP analysis (the bleached areas

correspond to the spots of highest intensities in the regions of interest, as visualized at the

fluorescent confocal microscope, see ***Methods***). Black and Magenta arrows point at he-he-ec

and he-ec-ec tri-cellular junctions, respectively (he: hemogenic cell; ec: endothelial cell). Green

arrows point at he-ec bi-cellular junctions. ec: endothelial cell, he: hemogenic cell. Bar = 20μm.

B, C, FRAP analysis of bleached eGFP-Jam3b localized in regions of interest in controls (grey)

and ArhGEF11 exon 38 splicing morpholino injected embryos (blue) or homozygous

ArhGEF11$^{CRISPR-Cterdel+/+}$ mutants (green)). The number of biological replicates (n) is stated on

the plots. Statistical tests: two sided unpaired two samples Wilcoxon test. Bb, Cc, evolution,

after photobleaching (at t = 0s), of the mean fluorescence intensity per condition at HE-EC,

HE-EC-EC and HE-HE-EC bi- and tri-junctions over time (10 min). Bb', Cc', median values for

maximum amplitude of recovery (maximum of simple exponential fitted curves, see **Methods**).

Bb", Cc", early evolution, after photobleaching (at t = 0s), of the mean fluorescence intensity

per condition over time (for the first 30 seconds). The fitted lines correspond to the linear

regression of the mean fluorescence intensities. Bb''', Cc''', median fluorescence recovery

slopes (linear regression). **D**, Model (2D deployment of the aortic wall) representing the

endothelial/hemogenic dynamic interplay and the proposed function of ArhGEF11 and of its

+exon 38 peptide encoding variant at junctional and membrane interfaces. This interplay

involves 2 essential dynamic events requiring junctional remodeling: **1** (left cartoon, magenta

arrows), the mobility of he-ec-ec tri-junctional contacts accompanying the movement of

endothelial cells (ex: for ec1 and ec2) along lateral sides of hemogenic cells which is required

to decrease the number of adjoining cells (see Lancino et al. 2018). This takes place

contemporarily to - or in alternance with -, the contraction of HE and EHT cells as they are

progressing throughout the emergence and reducing contacting membrane surfaces along the

longitudinal axis (the reduction in contacting membrane surfaces also involves membrane

retrieval, hypothetically via endocytosis). Our data obtained with the CRISPR deletion mutants

(a slight tendency to increase, on average, the turnover and mobile pool of these he-ec-ec

junctions) suggest that ArhGEF11, in the wild type condition, should be slowing down the

recycling of junctional components at tri-junctions, which hypothetically should contribute to

increase adhesion strength. This may be required also to stabilize the junction-cytoskeleton

interface involved in controlling the contraction/shrinkage of HE cells along the longitudinal

axis, a hypothesis that is compatible with the mutant phenotype observed in this study, i.e the

increase in frequency of more elongated HE cells and the decrease in HE cell progression

throughout EHT (see **Figure 7**); **2** (right cartoon, cyan arrow and bottom cartoons a-c), the

intercalation of an endothelial cell to isolate 2 adjacent hemogenic cells or 2 daughter cells

after mitosis (not depicted). This is mandatory for EHT progression and completion which

requires adjoining endothelial cells to protrude membrane extensions that will anastomose to

seal the aortic floor (see Lancino et al. 2018). The accumulation of adjoining cells of rather

small length and apparently impaired in EHT progression that we describe in **Figure** 7 upon

MO interference (that may also indicate impairment in abscission completion) suggests that

the ArhGEF11 +exon 38 peptide encoding variant is more specifically involved in controlling

the remodeling of the he/he interface that leads to endothelial cell intercalation (bottom

cartoons, the remodeling of he-he-ec junctions is leading to he-ec-ec junctions and takes place

between b and c). The increase in the recycling parameters that we measure in this interfering

condition (mobile pool and early speed of recovery) indicates that the activity of ArhGEF11,

and in particular of its +exon 38 peptide encoding variant, negatively controls junctional

recycling. Hypothetically, the junctional adhesion strengthening triggered by reducing the

recycling of junctional components at the he/he interface may be required during the early

phase and progression of intercalation to support increase in membrane tension and

environmental constraints (intercalation contributes to reducing the length/surface of the he/he

membrane interface preceding junctional remodeling (cartoons b and c); this reducing

membrane interface is in addition submitted to the shear stress imposed by blood flow).

1955

1956

1957

1958

References

Adamo, Luigi, Olaia Naveiras, Pamela L. Wenzel, Shannon McKinney-Freeman, Peter J. Mack, Jorge Gracia-Sancho, Astrid Suchy-Dicey, et al. 2009. "Biomechanical Forces Promote Embryonic Haematopoiesis." *Nature* 459 (7250): 1131–35. https://doi.org/10.1038/nature08073.

Akitake, Courtney M., Michelle Macurak, Marnie E. Halpern, and Mary G. Goll. 2011. "Transgenerational Analysis of Transcriptional Silencing in Zebrafish." *Developmental Biology* 352 (2): 191–201. https://doi.org/10.1016/j.ydbio.2011.01.002.

Barone, Cristiana, Roberto Orsenigo, Raffaella Meneveri, Silvia Brunelli, and Emanuele Azzoni. 2022. "One Size Does Not Fit All: Heterogeneity in Developmental Hematopoiesis." *Cells* 11 (6): 1061. https://doi.org/10.3390/cells11061061.

Bayless, Kayla J., and George E. Davis. 2002. "The Cdc42 and Rac1 GTPases Are Required for Capillary Lumen Formation in Three-Dimensional Extracellular Matrices." *Journal of Cell Science* 115 (6): 1123–36. https://doi.org/10.1242/jcs.115.6.1123.

Bengtsson, Henrik. 2022. *matrixStats: Functions That Apply to Rows and Columns of Matrices (and to Vectors)*. https://github.com/HenrikBengtsson/matrixStats.

Bengtsson, Henrik, Constantin Ahlmann-Eltze, Hector Corrada Bravo, Robert Gentleman, Jan Gleixner, Peter Hickey, Ola Hossjer, et al. 2023. "matrixStats: Functions That Apply to Rows and Columns of Matrices (and to Vectors)." https://cran.rstudio.com/web/packages/matrixStats/index.html.

Bertrand, Julien Y., Neil C. Chi, Buyung Santoso, Shutian Teng, Didier Y. R. Stainier, and David Traver. 2010. "Haematopoietic Stem Cells Derive Directly from Aortic Endothelium during Development." *Nature* 464 (7285): 108–11. https://doi.org/10.1038/nature08738.

Bertrand, Julien Y., Albert D. Kim, Emily P. Violette, David L. Stachura, Jennifer L. Cisson, and David Traver. 2007. "Definitive Hematopoiesis Initiates through a Committed Erythromyeloid Progenitor in the Zebrafish Embryo." *Development* 134 (23): 4147–56. https://doi.org/10.1242/dev.012385.

Boisset, Jean-Charles, Wiggert van Cappellen, Charlotte Andrieu-Soler, Niels Galjart, Elaine Dzierzak, and Catherine Robin. 2010. "In Vivo Imaging of Haematopoietic Cells Emerging from the Mouse Aortic Endothelium." *Nature* 464 (7285): 116–20. https://doi.org/10.1038/nature08764.

Bonkhofer, Florian, Rossella Rispoli, Philip Pinheiro, Monika Krecsmarik, Janina Schneider-Swales, Ingrid Ho Ching Tsang, Marella de Bruijn, Rui Monteiro, Tessa Peterkin, and Roger Patient. 2019. "Blood Stem Cell-Forming Haemogenic Endothelium in Zebrafish Derives from Arterial Endothelium." *Nature Communications* 10 (1): 3577. https://doi.org/10.1038/s41467-019-11423-2.

Bosveld, Floris, Zhimin Wang, and Yohanns Bellaïche. 2018. "Tricellular Junctions: A Hot Corner of Epithelial Biology." *Current Opinion in Cell Biology*, Cell Dynamics, 54 (October): 80–88. https://doi.org/10.1016/j.ceb.2018.05.002.

Bresciani, Erica, Elizabeth Broadbridge, and Paul P. Liu. 2018. "An Efficient Dissociation Protocol for Generation of Single Cell Suspension from Zebrafish Embryos and Larvae." *MethodsX* 5: 1287–90. https://doi.org/10.1016/j.mex.2018.10.009.

Bruijn, Marella F. T. R de, Xiaoqian Ma, Catherine Robin, Katrin Ottersbach, Maria-Jose Sanchez, and Elaine Dzierzak. 2002. "Hematopoietic Stem Cells Localize to the Endothelial Cell Layer in the Midgestation Mouse Aorta." *Immunity* 16 (5): 673–83. https://doi.org/10.1016/S1074-7613(02)00313-8.

Bryant, David M., Julie Roignot, Anirban Datta, Arend W. Overeem, Minji Kim, Wei Yu, Xiao Peng, et al. 2014. "A Molecular Switch for the Orientation of Epithelial Cell Polarization." *Developmental Cell* 31 (2): 171–87. https://doi.org/10.1016/j.devcel.2014.08.027.

Buckley, Clare E., and Daniel St Johnston. 2022. "Apical–Basal Polarity and the Control of Epithelial Form and Function." *Nature Reviews Molecular Cell Biology* 23 (8): 559–77. https://doi.org/10.1038/s41580-022-00465-y.

Burns, Caroline Erter, Tony DeBlasio, Yi Zhou, Jin Zhang, Leonard Zon, and Stephen D Nimer. 2002. "Isolation and Characterization of Runxa and Runxb, Zebrafish Members of the Runt Family of Transcriptional Regulators." *Experimental Hematology* 30 (12): 1381–89. https://doi.org/10.1016/S0301-472X(02)00955-4.

Butko, Emerald, Martin Distel, Claire Pouget, Bart Weijts, Isao Kobayashi, Kevin Ng, Christian Mosimann, et al. 2015. "Gata2b Is a Restricted Early Regulator of Hemogenic Endothelium in the Zebrafish Embryo." *Development* 142 (6): 1050–61. https://doi.org/10.1242/dev.119180.

Campinho, Pedro, Paola Lamperti, Francesco Boselli, Andrej Vilfan, and Julien Vermot. 2020. "Blood Flow Limits Endothelial Cell Extrusion in the Zebrafish Dorsal Aorta." *Cell Reports* 31 (2): 107505. https://doi.org/10.1016/j.celrep.2020.03.069.

Chalin, Dmitrii, Charlotte Bureau, Andrea Parmeggiani, Sergei Rochal, Karima Kissa, and Ivan Golushko. 2021. "Modeling and Live Imaging of Mechanical Instabilities in the Zebrafish Aorta during Hematopoiesis." *Scientific Reports* 11 (1): 9316. https://doi.org/10.1038/s41598-021-88667-w.

Chi, Neil C., Robin M. Shaw, Sarah De Val, Guson Kang, Lily Y. Jan, Brian L. Black, and Didier Y. R. Stainier. 2008. "Foxn4 Directly Regulates Tbx2b Expression and Atrioventricular Canal Formation." *Genes & Development* 22 (6): 734–39. https://doi.org/10.1101/gad.1629408.

Chikumi, Hiroki, Ana Barac, Babak Behbahani, Yuan Gao, Hidemi Teramoto, Yi Zheng, and J. Silvio Gutkind. 2004. "Homo- and Hetero-Oligomerization of PDZ-RhoGEF, LARG and p115RhoGEF by Their C-Terminal Region Regulates Their in Vivo Rho GEF Activity and Transforming Potential." *Oncogene* 23 (1): 233–40. https://doi.org/10.1038/sj.onc.1207012.

Ciau-Uitz, Aldo, Rui Monteiro, Arif Kirmizitas, and Roger Patient. 2014. "Developmental Hematopoiesis: Ontogeny, Genetic Programming and Conservation." *Experimental Hematology*, Genomics and Model Organisms: New Horizons for Experimental Hematology, 42 (8): 669–83. https://doi.org/10.1016/j.exphem.2014.06.001.

Clarke, Erik, and Scott Sherrill-Mix. 2017. *Ggbeeswarm: Categorical Scatter (Violin Point) Plots*. https://github.com/eclarke/ggbeeswarm.

Cumano, Ana, Françoise Dieterlen-Lievre, and Isabelle Godin. 1996. "Lymphoid Potential, Probed before Circulation in Mouse, Is Restricted to Caudal Intraembryonic Splanchnopleura." *Cell* 86 (6): 907–16. https://doi.org/10.1016/S0092-8674(00)80166-X.

Daniel, Michael G., David Sachs, Jeffrey M. Bernitz, Yesai Fstkchyan, Katrina Rapp, Namita Satija, Kenneth Law, et al. 2019. "Induction of Human Hemogenesis in Adult Fibroblasts by Defined Factors and Hematopoietic Coculture." *FEBS Letters* 593 (23): 3266–87. https://doi.org/10.1002/1873-3468.13621.

Davis, George E., Kayla J. Bayless, and Anil Mavila. 2002. "Molecular Basis of Endothelial Cell Morphogenesis in Three-Dimensional Extracellular Matrices." *The Anatomical Record* 268 (3): 252–75. https://doi.org/10.1002/ar.10159.

De Chaumont, Fabrice, Stéphane Dallongeville, Nicolas Chenouard, Nicolas Hervé, Sorin Pop, Thomas Provoost, Vannary Meas-Yedid, et al. 2012. "Icy: An Open Bioimage Informatics Platform for Extended Reproducible Research." *Nature Methods* 9 (7): 690–96. https://doi.org/10.1038/nmeth.2075.

Dignum, Tessa, Barbara Varnum-Finney, Sanjay R. Srivatsan, Stacey Dozono, Olivia Waltner, Adam M. Heck, Takashi Ishida, et al. 2021. "Multipotent Progenitors and Hematopoietic Stem Cells Arise Independently from Hemogenic Endothelium in the Mouse Embryo." *Cell Reports* 36 (11): 109675. https://doi.org/10.1016/j.celrep.2021.109675.

Dufour, Alexandre, Roman Thibeaux, Elisabeth Labruyere, Nancy Guillen, and Jean-Christophe Olivo-Marin. 2011. "3-D Active Meshes: Fast Discrete Deformable Models for Cell Tracking in 3-D Time-Lapse Microscopy." *IEEE Transactions on Image Processing* 20 (7): 1925–37. https://doi.org/10.1109/TIP.2010.2099125.

Ebnet, Klaus. 2017. "Junctional Adhesion Molecules (JAMs): Cell Adhesion Receptors With Pleiotropic Functions in Cell Physiology and Development." *Physiological Reviews* 97 (4): 1529–54. https://doi.org/10.1152/physrev.00004.2017.

Ebnet, Klaus, Michel Aurrand-Lions, Annegret Kuhn, Friedemann Kiefer, Stefan Butz, Kerstin Zander, Maria-Katharina Meyer Zu Brickwedde, Atsushi Suzuki, Beat A. Imhof, and Dietmar Vestweber. 2003. "The Junctional Adhesion Molecule (JAM) Family Members JAM-2 and JAM-3 Associate with the Cell Polarity Protein PAR-3: A Possible Role for JAMs in Endothelial Cell Polarity." *Journal of Cell Science* 116 (19): 3879–91. https://doi.org/10.1242/jcs.00704.

Ebnet, Klaus, and Volker Gerke. 2022. "Rho and Rab Family Small GTPases in the Regulation of Membrane Polarity in Epithelial Cells." *Frontiers in Cell and Developmental Biology* 10. https://www.frontiersin.org/articles/10.3389/fcell.2022.948013.

Ebnet, Klaus, Daniel Kummer, Tim Steinbacher, Amrita Singh, Masanori Nakayama, and Maja Matis. 2018. "Regulation of Cell Polarity by Cell Adhesion Receptors." *Seminars in Cell & Developmental Biology*, Cell Polarity and Planar Cell Polarity Proteins, 81 (September): 2–12. https://doi.org/10.1016/j.semcdb.2017.07.032.

Ebnet, Klaus, Atsushi Suzuki, Yosuke Horikoshi, Tomonori Hirose, Maria-Katharina Meyer zu Brickwedde, Shigeo Ohno, and Dietmar Vestweber. 2001. "The Cell Polarity Protein ASIP/PAR-3 Directly Associates with Junctional Adhesion Molecule (JAM)." *The EMBO Journal* 20 (14): 3738–48. https://doi.org/10.1093/emboj/20.14.3738.

Eilken, Hanna M., Shin-Ichi Nishikawa, and Timm Schroeder. 2009. "Continuous Single-Cell Imaging of Blood Generation from Haemogenic Endothelium." *Nature* 457 (7231): 896–900. https://doi.org/10.1038/nature07760.

Farrell, Jeffrey A., Yiqun Wang, Samantha J. Riesenfeld, Karthik Shekhar, Aviv Regev, and Alexander F. Schier. 2018. "Single-Cell Reconstruction of Developmental Trajectories during Zebrafish Embryogenesis." Science 360 (6392): eaar3131. https://doi.org/10.1126/science.aar3131.

Frame, Jenna M., Katherine H. Fegan, Simon J. Conway, Kathleen E. McGrath, and James Palis. 2016. "Definitive Hematopoiesis in the Yolk Sac Emerges from Wnt-Responsive Hemogenic Endothelium Independently of Circulation and Arterial Identity." *Stem Cells* 34 (2): 431–44. https://doi.org/10.1002/stem.2213.

Gama-Norton, Leonor, Eva Ferrando, Cristina Ruiz-Herguido, Zhenyi Liu, Jordi Guiu, Abul B. M. M. K. Islam, Sung-Uk Lee, et al. 2015. "Notch Signal Strength Controls Cell Fate in the Haemogenic Endothelium." *Nature Communications* 6 (1): 8510. https://doi.org/10.1038/ncomms9510.

Ganuza, Miguel, Brandon Hadland, Ashley Chabot, Chen Li, Guolian Kang, Irwin Bernstein, and Shannon McKinney-Freeman. 2017. "Murine Hemogenic Endothelial Precursors Display Heterogeneous Hematopoietic Potential Ex Vivo." *Experimental Hematology* 51 (July): 25-35.e6. https://doi.org/10.1016/j.exphem.2017.04.006.

Gao, Long, Joanna Tober, Peng Gao, Changya Chen, Kai Tan, and Nancy A. Speck. 2018. "RUNX1 and the Endothelial Origin of Blood." *Experimental Hematology* 68 (December): 2–9. https://doi.org/10.1016/j.exphem.2018.10.009.

Garcia-Porrero, Juan A., Isabelle E. Godin, and Françoise Dieterlen-Lièvre. 1995. "Potential Intraembryonic Hemogenic Sites at Pre-Liver Stages in the Mouse." *Anatomy and Embryology* 192 (5): 425–35. https://doi.org/10.1007/BF00240375.

Garnier, Simon. 2021. *Viridis: Colorblind-Friendly Color Maps for R*. https://CRAN.R-project.org/package=viridis.

Garrido-Urbani, S., P. F. Bradfield, and B. A. Imhof. 2014. "Tight Junction Dynamics: The Role of Junctional Adhesion Molecules (JAMs)." *Cell and Tissue Research* 355 (3): 701–15. https://doi.org/10.1007/s00441-014-1820-1.

Ghosn, Eliver, Momoko Yoshimoto, Hiromitsu Nakauchi, Irving L. Weissman, and Leonore A. Herzenberg. 2019. "Hematopoietic Stem Cell-Independent Hematopoiesis and the Origins of Innate-like B Lymphocytes." *Development* 146 (15): dev170571. https://doi.org/10.1242/dev.170571.

Grosjean, Philippe. 2019. *SciViews: SciViews - Main Package*. https://CRAN.R-project.org/package=SciViews.

Gudipaty, Swapna Aravind, and Jody Rosenblatt. 2017. "Epithelial Cell Extrusion: Pathways and Pathologies." *Seminars in Cell & Developmental Biology*, Extracellular Vesicles, 67 (July): 132–40. https://doi.org/10.1016/j.semcdb.2016.05.010.

Hadland, Brandon, and Momoko Yoshimoto. 2018. "Many Layers of Embryonic Hematopoiesis: New Insights into B-Cell Ontogeny and the Origin of Hematopoietic Stem Cells." *Experimental Hematology* 60 (April): 1–9. https://doi.org/10.1016/j.exphem.2017.12.008.

Herbert, Shane P., Jan Huisken, Tyson N. Kim, Morri E. Feldman, Benjamin T. Houseman, Rong A. Wang, Kevan M. Shokat, and Didier Y. R. Stainier. 2009. "Arterial-Venous Segregation by Selective Cell Sprouting: An Alternative Mode of Blood Vessel Formation." *Science* 326 (5950): 294–98. https://doi.org/10.1126/science.1178577.

Herwig, Lukas, Yannick Blum, Alice Krudewig, Elin Ellertsdottir, Anna Lenard, Heinz-Georg Belting, and Markus Affolter. 2011. "Distinct Cellular Mechanisms of Blood Vessel Fusion in the Zebrafish Embryo." *Current Biology* 21 (22): 1942–48. https://doi.org/10.1016/j.cub.2011.10.016.

Itoh, Masahiko. 2013. "ARHGEF11, a Regulator of Junction-Associated Actomyosin in Epithelial Cells." *Tissue Barriers* 1 (2): e24221. https://doi.org/10.4161/tisb.24221.

Itoh, Masahiko, Derek C. Radisky, Masaaki Hashiguchi, and Hiroyuki Sugimoto. 2017. "The Exon 38-Containing ARHGEF11 Splice Isoform Is Differentially Expressed and Is Required for Migration and Growth in Invasive Breast Cancer Cells." *Oncotarget* 8 (54): 92157–70. https://doi.org/10.18632/oncotarget.20985.

Itoh, Masahiko, Hiroyuki Sasaki, Mikio Furuse, Harunobu Ozaki, Toru Kita, and Shoichiro Tsukita. 2001. "Junctional Adhesion Molecule (JAM) Binds to PAR-3." *Journal of Cell Biology* 154 (3): 491–98. https://doi.org/10.1083/jcb.200103047.

Itoh, Masahiko, Sachiko Tsukita, Yuji Yamazaki, and Hiroyuki Sugimoto. 2012. "Rho GTP Exchange Factor ARHGEF11 Regulates the Integrity of Epithelial Junctions by Connecting ZO-1 and RhoA-Myosin II Signaling." *Proceedings of the National Academy of Sciences* 109 (25): 9905–10. https://doi.org/10.1073/pnas.1115063109.

Ivanovs, Andrejs, Stanislav Rybtsov, Elizabeth S. Ng, Edouard G. Stanley, Andrew G. Elefanty, and Alexander Medvinsky. 2017. "Human Haematopoietic Stem Cell Development: From the Embryo to the Dish." *Development* 144 (13): 2323–37. https://doi.org/10.1242/dev.134866.

Ivanovs, Andrejs, Stanislav Rybtsov, Lindsey Welch, Richard A. Anderson, Marc L. Turner, and Alexander Medvinsky. 2011. "Highly Potent Human Hematopoietic Stem Cells First Emerge in the Intraembryonic Aorta-Gonad-Mesonephros Region." *Journal of Experimental Medicine* 208 (12): 2417–27. https://doi.org/10.1084/jem.20111688.

Jabermoradi, Abbas, Suyeon Yang, Martijn I. Gobes, John P. M. van Duynhoven, and Johannes Hohlbein. 2022. "Enabling Single-Molecule Localization Microscopy in Turbid Food Emulsions." *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 380 (2220): 20200164. https://doi.org/10.1098/rsta.2020.0164.

Jaffredo, Thierry, Rodolphe Gautier, Anne Eichmann, and Françoise Dieterlen-Lièvre. 1998. "Intraaortic Hemopoietic Cells Are Derived from Endothelial Cells during Ontogeny." *Development* 125 (22): 4575–83. https://doi.org/10.1242/dev.125.22.4575.

Jaffredo, Thierry, and Laurent Yvernogeau. 2014. "How the Avian Model Has Pioneered the
Field of Hematopoietic Development." *Experimental Hematology*, Genomics and Model
Organisms: New Horizons for Experimental Hematology, 42 (8): 661–68.
https://doi.org/10.1016/j.exphem.2014.05.009.
Jin, Suk-Won, Dimitris Beis, Tracy Mitchell, Jau-Nian Chen, and Didier Y. R. Stainier. 2005.
"Cellular and Molecular Analyses of Vascular Tube and Lumen Formation in Zebrafish."
Development 132 (23): 5199–5209. https://doi.org/10.1242/dev.02087.
Kalev-Zylinska, Maggie L., Julia A. Horsfield, Maria Vega C. Flores, John H. Postlethwait,
Maria R. Vitas, Andrea M. Baas, Philip S. Crosier, and Kathryn E. Crosier. 2002.
"Runx1 Is Required for Zebrafish Blood and Vessel Development and Expression of a
Human RUNX1-CBF2T1 Transgene Advances a Model for Studies of
Leukemogenesis." *Development* 129 (8): 2015–30.
https://doi.org/10.1242/dev.129.8.2015.
Kamei, Makoto, W. Brian Saunders, Kayla J. Bayless, Louis Dye, George E. Davis, and Brant
M. Weinstein. 2006. "Endothelial Tubes Assemble from Intracellular Vacuoles in Vivo."
Nature 442 (7101): 453–56. https://doi.org/10.1038/nature04923.
Kassambara, Alboukadel. 2020. *Ggpubr: Ggplot2 Based Publication Ready Plots*.
https://rpkgs.datanovia.com/ggpubr/.
Kataoka, Hiroko, Mitsuo Ochi, Koh-ichi Enomoto, and Akira Yamaguchi. 2000. "Cloning and
Embryonic Expression Patterns of the Zebrafish Runt Domain Genes, Runxa and
Runxb." *Mechanisms of Development* 98 (1): 139–43. https://doi.org/10.1016/S0925-
4773(00)00445-7.
Kimmel, Charles B., William W. Ballard, Seth R. Kimmel, Bonnie Ullmann, and Thomas F.
Schilling. 1995. "Stages of Embryonic Development of the Zebrafish." *Developmental
Dynamics* 203 (3): 253–310. https://doi.org/10.1002/aja.1002030302.
Kissa, Karima, and Philippe Herbomel. 2010. "Blood Stem Cells Emerge from Aortic
Endothelium by a Novel Type of Cell Transition." *Nature* 464 (7285): 112–15.
https://doi.org/10.1038/nature08761.
Kobayashi, Isao, Jingjing Kobayashi-Sun, Yuto Hirakawa, Madoka Ouchi, Koyuki Yasuda,
Hiroyasu Kamei, Shigetomo Fukuhara, and Masaaki Yamaguchi. 2020. "Dual Role of
Jam3b in Early Hematopoietic and Vascular Development." *Development* 147 (1):
dev181040. https://doi.org/10.1242/dev.181040.
Lagendijk, Anne Karine, Alpha S Yap, and Benjamin M Hogan. 2014. "Endothelial Cell–Cell
Adhesion during Zebrafish Vascular Development." *Cell Adhesion & Migration* 8 (2):
136–45. https://doi.org/10.4161/cam.28229.
Lancino, Mylene, Sara Majello, Sebastien Herbert, Fabrice De Chaumont, Jean-Yves Tinevez,
Jean-Christophe Olivo-Marin, Philippe Herbomel, and Anne Schmidt. 2018.
"Anisotropic Organization of Circumferential Actomyosin Characterizes Hematopoietic
Stem Cells Emergence in the Zebrafish." *eLife* 7 (August): e37355.
https://doi.org/10.7554/eLife.37355.
Lancrin, Christophe, Milena Mazan, Monika Stefanska, Rahima Patel, Monika Lichtinger,
Guilherme Costa, Özge Vargel, et al. 2012. "GFI1 and GFI1B Control the Loss of
Endothelial Identity of Hemogenic Endothelium during Hematopoietic Commitment."
Blood 120 (2): 314–22. https://doi.org/10.1182/blood-2011-10-386094.
Lee, SungKyoung, Benjamin Cieply, Yueqin Yang, Natoya Peart, Carl Glaser, Patricia Chan,
and Russ P. Carstens. 2018. "Esrp1-Regulated Splicing of Arhgef11 Isoforms Is
Required for Epithelial Tight Junction Integrity." *Cell Reports* 25 (9): 2417-2430.e5.
https://doi.org/10.1016/j.celrep.2018.10.097.
Legland, David, Ignacio Arganda-Carreras, and Philippe Andrey. 2016. "MorphoLibJ:
Integrated Library and Plugins for Mathematical Morphology with ImageJ."
Bioinformatics 32 (22): 3532–34. https://doi.org/10.1093/bioinformatics/btw413.
Liao, Wayne, Brent W. Bisgrove, Holly Sawyer, Barbara Hug, Bridgit Bell, Kevin Peters, David
J. Grunwald, and Didier Y. R. Stainier. 1997. "The Zebrafish Gene Cloche Acts

Upstream of a Flk-1 Homologue to Regulate Endothelial Cell Differentiation." *Development* 124 (2): 381–89. https://doi.org/10.1242/dev.124.2.381.

Lie-A-Ling, Michael, Elli Marinopoulou, Yaoyong Li, Rahima Patel, Monika Stefanska, Constanze Bonifer, Crispin Miller, Valerie Kouskoff, and Georges Lacaud. 2014. "RUNX1 Positively Regulates a Cell Adhesion and Migration Program in Murine Hemogenic Endothelium Prior to Blood Emergence." *Blood* 124 (11): e11–20. https://doi.org/10.1182/blood-2014-04-572958.

Lin, Hui-Feng, David Traver, Hao Zhu, Kimberly Dooley, Barry H. Paw, Leonard I. Zon, and Robert I. Handin. 2005. "Analysis of Thrombocyte Development in CD41-GFP Transgenic Zebrafish." *Blood* 106 (12): 3803–10. https://doi.org/10.1182/blood-2005-01-0179.

Lundin, Vanessa, Wade W. Sugden, Lindsay N. Theodore, Patricia M. Sousa, Areum Han, Stephanie Chou, Paul J. Wrighton, et al. 2020. "YAP Regulates Hematopoietic Stem Cell Formation in Response to the Biomechanical Forces of Blood Flow." *Developmental Cell* 52 (4): 446-460.e5. https://doi.org/10.1016/j.devcel.2020.01.006.

Mack, Natalie Ann, and Marios Georgiou. 2014. "The Interdependence of the Rho GTPases and Apicobasal Cell Polarity." *Small GTPases* 5 (2): e973768. https://doi.org/10.4161/21541248.2014.973768.

Manoli, Martha, and Wolfgang Driever. 2012. "Fluorescence-Activated Cell Sorting (FACS) of Fluorescently Tagged Cells from Zebrafish Larvae for RNA Isolation." *Cold Spring Harbor Protocols* 2012 (8): pdb.prot069633. https://doi.org/10.1101/pdb.prot069633.

Marshall, Caroline J., and Adrian J. Thrasher. 2001. "The Embryonic Origins of Human Haematopoiesis." *British Journal of Haematology* 112 (4): 838–50. https://doi.org/10.1046/j.1365-2141.2001.02537.x.

Meder, Doris, Anna Shevchenko, Kai Simons, and Joachim Füllekrug. 2005. "Gp135/Podocalyxin and NHERF-2 Participate in the Formation of a Preapical Domain during Polarization of MDCK Cells." *Journal of Cell Biology* 168 (2): 303–13. https://doi.org/10.1083/jcb.200407072.

Medvinsky, Alexander, and Elaine Dzierzak. 1996. "Definitive Hematopoiesis Is Autonomously Initiated by the AGM Region." *Cell* 86 (6): 897–906. https://doi.org/10.1016/S0092-8674(00)80165-8.

Medvinsky, Alexander L., Nina L. Samoylina, Albrecht M. Müller, and Elaine A. Dzierzak. 1993. "An Early Pre-Liver Intraembryonic Source of CFU-S in the Developing Mouse." *Nature* 364 (6432): 64–67. https://doi.org/10.1038/364064a0.

Monteiro, Rui, Philip Pinheiro, Nicola Joseph, Tessa Peterkin, Jana Koth, Emmanouela Repapi, Florian Bonkhofer, Arif Kirmizitas, and Roger Patient. 2016. "Transforming Growth Factor β Drives Hemogenic Endothelium Programming and the Transition to Hematopoietic Stem Cells." *Developmental Cell* 38 (4): 358–70. https://doi.org/10.1016/j.devcel.2016.06.024.

Moreno-Mateos, Miguel A., Charles E. Vejnar, Jean-Denis Beaudoin, Juan P. Fernandez, Emily K. Mis, Mustafa K. Khokha, and Antonio J. Giraldez. 2015. "CRISPRscan: Designing Highly Efficient sgRNAs for CRISPR-Cas9 Targeting in Vivo." *Nature Methods* 12 (10): 982–88. https://doi.org/10.1038/nmeth.3543.

Müller, A. M., A. Medvinsky, J. Strouboulis, F. Grosveld, and E. Dzierzak. 1994. "Development of Hematopoietic Stem Cell Activity in the Mouse Embryo." *Immunity* 1 (4): 291–301. https://doi.org/10.1016/1074-7613(94)90081-7.

Murayama, Emi, Karima Kissa, Agustin Zapata, Elodie Mordelet, Valérie Briolat, Hui-Feng Lin, Robert I. Handin, and Philippe Herbomel. 2006. "Tracing Hematopoietic Precursor Migration to Successive Hematopoietic Organs during Zebrafish Development." *Immunity* 25 (6): 963–75. https://doi.org/10.1016/j.immuni.2006.10.015.

Nielsen, Julie S., and Kelly M. McNagny. 2008. "Novel Functions of the CD34 Family." *Journal of Cell Science* 121 (22): 3683–92. https://doi.org/10.1242/jcs.037507.

Nieto, M. Angela, Ruby Yun-Ju Huang, Rebecca A. Jackson, and Jean Paul Thiery. 2016. "EMT: 2016." *Cell* 166 (1): 21–45. https://doi.org/10.1016/j.cell.2016.06.028.

Nishimura, Tamako, Hisao Honda, and Masatoshi Takeichi. 2012. "Planar Cell Polarity Links Axes of Spatial Dynamics in Neural-Tube Closure." *Cell* 149 (5): 1084–97. https://doi.org/10.1016/j.cell.2012.04.021.

North, Trista E., Marella F. T. R. de Bruijn, Terryl Stacy, Laleh Talebian, Evan Lind, Catherine Robin, Michael Binder, Elaine Dzierzak, and Nancy A. Speck. 2002. "Runx1 Expression Marks Long-Term Repopulating Hematopoietic Stem Cells in the Midgestation Mouse Embryo." *Immunity* 16 (5): 661–72. https://doi.org/10.1016/S1074-7613(02)00296-0.

North, Trista E., Wolfram Goessling, Marian Peeters, Pulin Li, Craig Ceol, Allegra M. Lord, Gerhard J. Weber, et al. 2009. "Hematopoietic Stem Cell Development Is Dependent on Blood Flow." *Cell* 137 (4): 736–48. https://doi.org/10.1016/j.cell.2009.04.023.

North, Trista, Ting-Lei Gu, Terryl Stacy, Qing Wang, Louisa Howard, Michael Binder, Miguel Marín-Padilla, and Nancy A. Speck. 1999. "*Cbfa2* Is Required for the Formation of Intra-Aortic Hematopoietic Clusters." *Development* 126 (11): 2563–75. https://doi.org/10.1242/dev.126.11.2563.

Olayioye, Monilola A., Bettina Noll, and Angelika Hausser. 2019. "Spatiotemporal Control of Intracellular Membrane Trafficking by Rho GTPases." *Cells* 8 (12): 1478. https://doi.org/10.3390/cells8121478.

Orkin, Stuart H., and Leonard I. Zon. 2008. "Hematopoiesis: An Evolving Paradigm for Stem Cell Biology." *Cell* 132 (4): 631–44. https://doi.org/10.1016/j.cell.2008.01.025.

Padfield, Daniel, and Granville Matheson. 2020. *Nls.Multstart: Robust Non-Linear Regression Using AIC Scores*. https://CRAN.R-project.org/package=nls.multstart.

Patil, Indrajeet. 2022. *Ggstatsplot: Ggplot2 Based Plots with Statistical Details*. https://CRAN.R-project.org/package=ggstatsplot.

Pei, Duanqing, Xiaodong Shu, Ama Gassama-Diagne, and Jean Paul Thiery. 2019. "Mesenchymal–Epithelial Transition in Development and Reprogramming." *Nature Cell Biology* 21 (1): 44–53. https://doi.org/10.1038/s41556-018-0195-z.

Pradella, Davide, Chiara Naro, Claudio Sette, and Claudia Ghigna. 2017. "EMT and Stemness: Flexible Processes Tuned by Alternative Splicing in Development and Cancer Progression." *Molecular Cancer* 16 (1): 8. https://doi.org/10.1186/s12943-016-0579-2.

Ram, Karthik, and Hadley Wickham. 2018. *Wesanderson: A Wes Anderson Palette Generator*. https://github.com/karthik/wesanderson.

Ridley, Anne J. 2006. "Rho GTPases and Actin Dynamics in Membrane Protrusions and Vesicle Trafficking." *Trends in Cell Biology*, Membrane Dynamics, 16 (10): 522–29. https://doi.org/10.1016/j.tcb.2006.08.006.

Robert-Moreno, Àlex, Jordi Guiu, Cristina Ruiz-Herguido, M. Eugenia López, Julia Inglés-Esteve, Lluis Riera, Alex Tipping, et al. 2008. "Impaired Embryonic Haematopoiesis yet Normal Arterial Development in the Absence of the Notch Ligand Jagged1." *The EMBO Journal* 27 (13): 1886–95.

Rodriguez-Boulan, Enrique, Anne Müsch, and Andre Le Bivic. 2004. "Epithelial Trafficking: New Routes to Familiar Places." *Current Opinion in Cell Biology* 16 (4): 436–42. https://doi.org/10.1016/j.ceb.2004.06.013.

Román-Fernández, Alvaro, and David M. Bryant. 2016. "Complex Polarity: Building Multicellular Tissues Through Apical Membrane Traffic." *Traffic* 17 (12): 1244–61. https://doi.org/10.1111/tra.12417.

Rossman, Kent L., Channing J. Der, and John Sondek. 2005. "GEF Means Go: Turning on RHO GTPases with Guanine Nucleotide-Exchange Factors." *Nature Reviews Molecular Cell Biology* 6 (2): 167–80. https://doi.org/10.1038/nrm1587.

Samsa, Leigh Ann, Nicole Fleming, Scott Magness, Li Qian, and Jiandong Liu. 2016. "Isolation and Characterization of Single Cells from Zebrafish Embryos." *Journal of Visualized Experiments*, no. 109 (March): 53877. https://doi.org/10.3791/53877.

Sato, Yuki, Mugiho Shigematsu, Maria Shibata-Kanno, Sho Maejima, Chie Tamura, and Hirotaka Sakamoto. 2023. "Aquaporin Regulates Cell Rounding through Vacuole Formation during Endothelial-to-Hematopoietic Transition." *Development* 150 (11): dev201275. https://doi.org/10.1242/dev.201275.

Sehnert, Amy J., Anja Huq, Brant M. Weinstein, Charline Walker, Mark Fishman, and Didier Y. R. Stainier. 2002. "Cardiac Troponin T Is Essential in Sarcomere Assembly and Cardiac Contractility." *Nature Genetics* 31 (1): 106–10. https://doi.org/10.1038/ng875.

Shapiro, Irina M., Albert W. Cheng, Nicholas C. Flytzanis, Michele Balsamo, John S. Condeelis, Maja H. Oktay, Christopher B. Burge, and Frank B. Gertler. 2011. "An EMT–Driven Alternative Splicing Program Occurs in Human Breast Cancer and Modulates Cellular Phenotype." *PLOS Genetics* 7 (8): e1002218. https://doi.org/10.1371/journal.pgen.1002218.

Sood, Raman, Milton A. English, Christiane L. Belele, Hao Jin, Kevin Bishop, Rebecca Haskins, Mary Cathleen McKinney, et al. 2010. "Development of Multilineage Adult Hematopoiesis in the Zebrafish with a Runx1 Truncation Mutation." *Blood* 115 (14): 2806–9. https://doi.org/10.1182/blood-2009-08-236729.

Staneva, Ralitza, and Romain Levayer. 2023. "Chapter Five - Cell Polarity and Extrusion: How to Polarize Extrusion and Extrude Misspolarized Cells?" In *Current Topics in Developmental Biology*, edited by ULRICH Tepass, 154:131–67. Cell Polarity in Development and Disease. Academic Press. https://doi.org/10.1016/bs.ctdb.2023.02.010.

Strilić, Boris, Jan Eglinger, Michael Krieg, Martin Zeeb, Jennifer Axnick, Pavel Babál, Daniel J. Müller, and Eckhard Lammert. 2010. "Electrostatic Cell-Surface Repulsion Initiates Lumen Formation in Developing Blood Vessels." *Current Biology* 20 (22): 2003–9. https://doi.org/10.1016/j.cub.2010.09.061.

Strilić, Boris, Tomáš Kučera, Jan Eglinger, Michael R. Hughes, Kelly M. McNagny, Sachiko Tsukita, Elisabetta Dejana, Napoleone Ferrara, and Eckhard Lammert. 2009. "The Molecular Basis of Vascular Lumen Formation in the Developing Mouse Aorta." *Developmental Cell* 17 (4): 505–15. https://doi.org/10.1016/j.devcel.2009.08.011.

Sur, Abhinav, Yiqun Wang, Paulina Capar, Gennady Margolin, Morgan Kathleen Prochaska, and Jeffrey A. Farrell. 2023. "Single-Cell Analysis of Shared Signatures and Transcriptional Diversity during Zebrafish Development." Developmental Cell 58 (24): 3028-3047.e12. https://doi.org/10.1016/j.devcel.2023.11.001.

Suster, Maximiliano L., Gembu Abe, Anders Schouw, and Koichi Kawakami. 2011. "Transposon-Mediated BAC Transgenesis in Zebrafish." *Nature Protocols* 6 (12): 1998–2021. https://doi.org/10.1038/nprot.2011.416.

Swiers, Gemma, Claudia Baumann, John O'Rourke, Eleni Giannoulatou, Stephen Taylor, Anagha Joshi, Victoria Moignard, et al. 2013. "Early Dynamic Fate Changes in Haemogenic Endothelium Characterized at the Single-Cell Level." *Nature Communications* 4 (1): 2924. https://doi.org/10.1038/ncomms3924.

Tian, Ye, Jin Xu, Shachuan Feng, Sicong He, Shizheng Zhao, Lu Zhu, Wan Jin, et al. 2017. "The First Wave of T Lymphopoiesis in Zebrafish Arises from Aorta Endothelium Independent of Hematopoietic Stem Cells." *Journal of Experimental Medicine* 214 (11): 3347–60. https://doi.org/10.1084/jem.20170488.

Weijts, Bart, Laurent Yvernogeau, and Catherine Robin. 2021. "Recent Advances in Developmental Hematopoiesis: Diving Deeper With New Technologies." *Frontiers in Immunology* 12. https://www.frontiersin.org/articles/10.3389/fimmu.2021.790379.

Wickham, Hadley. 2016. *Ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. https://ggplot2.tidyverse.org.

Wickham, Hadley. 2022. *Stringr: Simple, Consistent Wrappers for Common String Operations*. https://CRAN.R-project.org/package=stringr.

Wickham, Hadley, Romain François, Lionel Henry, Kirill Müller, and Davis Vaughan. 2023. *Dplyr: A Grammar of Data Manipulation*. https://CRAN.R-project.org/package=dplyr.

Wickham, Hadley, Jim Hester, and Jennifer Bryan. 2022. *Readr: Read Rectangular Text Data*.
https://CRAN.R-project.org/package=readr.
Wilke, Claus O. 2020. *Cowplot: Streamlined Plot Theme and Plot Annotations for Ggplot2*.
https://wilkelab.org/cowplot/.
Wu, Yinyu, and Karen K. Hirschi. 2021. "Regulation of Hemogenic Endothelial Cell
Development and Function." *Annual Review of Physiology* 83 (February): 17–37.
https://doi.org/10.1146/annurev-physiol-021119-034352.
Xia, Jun, Mengyao Liu, Caiying Zhu, Shicheng Liu, Lanlan Ai, Dongyuan Ma, Ping Zhu, Lu
Wang, and Feng Liu. 2023. "Activation of Lineage Competence in Hemogenic
Endothelium Precedes the Formation of Hematopoietic Stem Cell Heterogeneity." *Cell
Research*, April, 1–16. https://doi.org/10.1038/s41422-023-00797-0.
Xiao, Nan. 2018. *Ggsci: Scientific Journal and Sci-Fi Themed Color Palettes for Ggplot2*.
https://CRAN.R-project.org/package=ggsci.
Xie, Yihui, Joe Cheng, and Xianying Tan. 2022. *DT: A Wrapper of the JavaScript Library
DataTables*. https://github.com/rstudio/DT.
Yoshimoto, Momoko, Encarnacion Montecino-Rodriguez, Michael J. Ferkowicz, Prashanth
Porayette, W. Christopher Shelley, Simon J. Conway, Kenneth Dorshkind, and Mervin
C. Yoder. 2011. "Embryonic Day 9 Yolk Sac and Intra-Embryonic Hemogenic
Endothelium Independently Generate a B-1 and Marginal Zone Progenitor Lacking B-
2 Potential." *Proceedings of the National Academy of Sciences* 108 (4): 1468–73.
https://doi.org/10.1073/pnas.1015841108.
Yoshimoto, Momoko, Prashanth Porayette, Nicole L. Glosson, Simon J. Conway, Nadia
Carlesso, Angelo A. Cardoso, Mark H. Kaplan, and Mervin C. Yoder. 2012.
"Autonomous Murine T-Cell Progenitor Production in the Extra-Embryonic Yolk Sac
before HSC Emergence." *Blood* 119 (24): 5706–14. https://doi.org/10.1182/blood-
2011-12-397489.
Zhao, Shizheng, Shachuan Feng, Ye Tian, and Zilong Wen. 2022. "Hemogenic and Aortic
Endothelium Arise from a Common Hemogenic Angioblast Precursor and Are Specified
by the Etv2 Dosage." *Proceedings of the National Academy of Sciences* 119 (13):
e2119051119. https://doi.org/10.1073/pnas.2119051119.
Zhu, Qin, Peng Gao, Joanna Tober, Laura Bennett, Changya Chen, Yasin Uzun, Yan Li, et al.
2020. "Developmental Trajectory of Prehematopoietic Stem Cell Formation from
Endothelium." *Blood* 136 (7): 845–56. https://doi.org/10.1182/blood.2020004801.
2408
2409

Figure 1

A

B

C

Tg(Kdrl:Gal4;UAS:RFP;4XNR:eGFP-podxl2)
Tg(Kdrl:mKate2-podxl2)

C
C
C
C cystein-containing
globular domain +
N-glycosylation

DTHL = PDZ-binding
C-ter motif

D

Figure 2

A

Tg(Kdrl:Gal4;UAS:RFP;4xNR:eGFP-podxl2)

B

Figure 3

A 48hpf

B

C

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8