Human deleterious mutation rate slows adaptation and implies high fitness variance

Joseph Matheson[1,2], Ulises Hernández[1], Jason Bertram[1,3], Joanna Masel[1]

[1]Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, 85721, USA

[2]Department of Ecology, Behavior, and Evolution, University of California San Diego, San Diego, CA,

92093, USA

[3] Department of Mathematics, University of Western Ontario, London ON, Canada

Corresponding author: Joanna Masel

Abstract

Each new human has an expected U_d = 2-10 new deleterious mutations. Using a novel approach to

capture complex linkage disequilibria from high U_d using genome-wide simulations, we confirm that

fitness decline due to the fixation of many slightly deleterious mutations can be compensated by rarer

beneficial mutations of larger effect. The evolution of increased genome size and complexity have

previously been attributed to a similarly asymmetric pattern of fixations, but we propose that the cause

might be high U_d rather than the small population size posited as causal by drift barrier theory. High

within-population variance in relative fitness is an inevitable consequence of high $U_d \sim$ 2-10 combined

with inferred human deleterious effect sizes; two individuals will typically differ in fitness by 15-40%.

The need to compensate for the deluge of deleterious mutations slows net adaptation (i.e. to the external

environment) by ~13%-55%. The rate of beneficial fixations is more sensitive to changes in the mutation

rate than the rate of deleterious fixations is. As a surprising consequence of this, an increase (e.g. 10%) in

overall mutation rate leads to faster adaptation; this puts to rest dysgenic fears about increasing mutation

rates due to rising paternal age.

Keywords: mutation load; Muller's ratchet; Ohta's ratchet; chromosome number; background selection;

mutational meltdown; binary indexed tree.

Introduction

The average human begins life with around a hundred new mutations not found in their parents, and geneticists have long worried about the effects of the resulting "mutation load" on human health [1–6]. Lesecque et al. [7] assumed that mutations are deleterious only in the 55% of the 6×10^9 diploid genome that is not dominated by inactive transposable elements, which evolves 5.7% slower than the rest of the genome due to this constraint, at a point mutation rate of 1.1×10^{-8}; this yields an estimated rate of deleterious mutations of $0.55 \times 6 \times 10^9 \times 0.057 \times 1.1 \times 10^{-8} = 2.1$ per replication. This estimate is conservative: some mutations to the other 45% of the genome are deleterious, more recent estimates of the human point mutation rate have 1.1×10^{-8} as the lower bound of a 95% confidence interval [8], and non-point mutations and beneficial mutations are neglected. Some therefore argue that the deleterious mutation rate is as high as ten [9,10]. Mutation rates of this order are not unique to humans [8,11].

Three fears about the effects of high deleterious mutation rates have haunted human genetics, namely that: 1) we do not understand how population persistence is possible [12]; 2) the health burden caused by this deluge of deleterious mutations is not shared equally by all humans; and 3) recent increases in mutation rate and/or decreases in natural selection will lower population fitness at mutation-selection balance, with potentially catastrophic consequences for health systems [1–3,6]. The second fear has a dark legacy of attempting to identify more vs. less 'burdened' humans [13,14].

Most substantial theoretical treatment has focused on the first fear. Some unconditionally deleterious mutations of smaller effect size will inevitably fix [15,16], producing cumulative degradation termed "Ohta's ratchet" [17]. But rarer, larger-effect beneficial fixations can compensate, enabling population persistence [18–20]. Given a distribution of fitness effects in an infinite sites model, fixed deleterious mutations tend to have smaller s [21] while beneficial fixations have larger s [22]. Asymmetry between deleterious and beneficial fixations matches known features of molecular adaptation. E.g., many proteins accumulate small deleterious mutations that slightly inhibit folding, which can be compensated for by a novel or improved or overexpressed chaperone protein [23]. Similarly, many poorly splicing introns can be

ameliorated by the evolution of a better spliceosome [24]. An asymmetric pattern of adaptation has been

inferred in influenza [25].

While the second fear has received little rhetorical attention from population geneticists, it has received

incidental theoretical treatment as "load" in the context of the first fear. Galeota-Sprung *et al.* [26] found

that the expected variance in fitness is $e^{U_d\overline{s_d}} - 1$ in the absence of linkage disequilibrium or epistasis.

They conclude that, unlike load metrics that compare to an ideal individual, variance in fitness is modest,

and not a problem for population persistence. They did not discuss the societal implications of variance in

fitness. They underestimated variance by using unrealistically low values of $\overline{s_d}$ (see Results), and by

neglecting linkage disequilibrium that makes purging harder. Similarly, warnings from prominent

geneticists of a coming health crisis use one-locus models of mutation-selection balance [1–3,6], raising

similar questions of the role of linkage disequilibrium; they also neglect large-effect beneficial mutations.

Quantitative treatment of the three fears requires incorporating both beneficial mutations, and emergent

linkage disequilibria [27]. Linkage disequilibria become especially pronounced when $U_d > 1$ [28,29].

Unfortunately, simulating this is computationally challenging. One remedy of convenience when

simulating $U_d > 1$ is to neglect beneficials, and instead periodically re-normalize relative fitness to

cosmetically remove ongoing degradation (e.g. compare Fig. S2 to Fig. 2 in [30]). Alternatively, most

previous studies of mutation load used unrealistically low mutation rates ($U_d < 1$), either directly [31–37], or

by assuming independent loci [30,38–40].

Simulation is essential to capture the full complexities of multilocus linkage disequilibrium. Most forward

time simulation methods hold a product such as sN constant by rescaling N to be smaller and s to be

larger, speeding computation [41]. But this reduces the number of segregating mutations $UN\tau$ (where τ is

the expected sojourn time), which understates the impact of linkage disequilibria. We simulate a census

population size of 20,000, producing a human-like level of neutral diversity ($N_e \sim 7500$), with emergent

linkage disequilibrium. To avoid computationally costly individual tracking of the enormous number of

segregating sites, we track only the combined fitness effects within 'linkage blocks' between

recombination hotspots, using a novel algorithm. Our approach allows us to recover information about

fixed mutations via tree-sequence recording [42], albeit at a cost to runtime.

We use simulations to quantify, given realistic linkage disequilibria, the fluxes of beneficial and

deleterious fixations, the speed of adaptation, and variation in fitness within populations with different

mutation rates. This allows us to quantitatively treat all three load-based fears.

Methods

Each individual has two characteristics: a genome, and a fitness value derived from it. Each individual's

genome is represented as two haplotypes, each an array of L non-recombining 'linkage blocks', divided

into 23 chromosomes. Each linkage block consists of a floating-point variable l_j, which summarizes the

fitness effects of all mutations that occurred in the history of that linkage block, such that $l_j =$

$\prod_i (1 + s_i)$. We assume a multiplicative form of co-dominance and no epistasis, such that $w_i =$

$\prod_{j=1}^{L}(l_{j,1}) \prod_{j=1}^{L}(l_{j,2})$ where $l_{j,1}$ and $l_{j,2}$ refer to the effects of linkage block j in haplotypes 1 and 2,

respectively. Note that this computationally convenient choice is not precisely equivalent to a typical

codominance model, where $1 + s_i$ is the fitness of a homozygote and $1 + s_i h_i$ is the fitness of a

heterozygote. While co-dominance is unrealistic for strongly deleterious mutations, which are often

highly recessive, it is reasonable for the small-effect deleterious mutations that drive Ohta's ratchet [43–45].

In addition to independent assortment of chromosomes, recombination occurs at hotspots between linkage

blocks via crossing-over events between homologous chromosomes. We simulate exactly two

recombination events per chromosome per meiosis, matching data for humans [46], although we don't

explicitly simulate a centrosome. Representing a genome as a set of 'linkage blocks' is a good

approximation of population genetics in non-microbial species [47–49]. Realistic values of L in humans are

in the range of 10^5-10^6 [50–54]. Once $L \geq 50 \times 23 = 1150$, results converge (Supplementary Figure 1), so

for computational efficiency we use $L = 50 \times 23$. This simplification should overestimate the effect of

linkage between selected mutations, which is conservative with respect to the ability of beneficial

mutations to counteract load.

Following recombination, we sample the number of new deleterious mutations in the gamete from a

Poisson distribution with mean U_d. Our distribution of fitness effects is based on a large empirical study

of Europeans [55], who fitted a gamma distribution for $2N_e sh$ with mean -224.33, shape parameter $\alpha =$

0.169 and scale parameter $\beta = 1327.4$. After drawing a value of $2N_e sh$ from this distribution, we rescale

to sh using their inferred $N_e = 11{,}823$. We use the sh value drawn from this distribution as our s_d value;

its mean is 0.009. We sample the number of new beneficial mutations from a Poisson distribution with

mean U_b, and fitness effects drawn from an exponential distribution with mean s_b (again, this is the

fitness effect in the heterozygote). We explore a range of values for U_b and s_b that we consider *a priori*

plausible: $U_b \sim 0.0001\text{-}0.01$ and $s_b \sim 0.001\text{-}0.01$.

We simulate a Moran model with constant population size N. An individual chosen uniformly at random

dies each time step and is replaced by a child produced by two parents, who are chosen with probability

proportional to their fitness w_i. Each generation consists of N time steps. The fitnesses of the population

are stored in an unsorted array — in a naïve implementation, exchanging an element to represent a birth

and death would be rapid, but sampling proportional to fitness would be $O(N)$. The current fastest

forward-time genetic simulation tools for large population sizes (e.g. fwdpy [56,57] and SLiM [58]) preprocess

cumulants each generation in a Wright-Fisher model; this speeds up sampling from the fitness array, and

while the processing algorithm is $O(N)$, it only needs to be performed once per generation. We instead

use a binary indexed tree [59] to sample fitnesses efficiently according to the cumulative probability

distribution — both updating and sampling from the tree are $O(\log N)$. Our scheme is expected to have

similar efficiency but is intended to be useful for future expansions of this approach to absolute fitness

and more complex life history models [60,61], e.g. to allow better treatment of reproductive compensation [62].

We initialize the population with mutationless individuals, then conduct a 'burn-in' phase during which

variation increases to stable levels (Supplementary Figure 2). We end the burn-in phase 500 generations

after a linear regression of the variance in fitness over the last 200 generations produces a slope less than

an arbitrarily chosen low value of $0.007/N$ that we visually confirmed to perform well (e.g.

Supplementary Figure 2). The length of the burn-in phase does not strongly depend on N (Supplementary

Figure 3).

We calculate the net fitness flux from each simulation as the slope of the regression of log mean

population fitness on time after burn-in (Supplementary Figure 2, black slope following dashed line). To

numerically solve for a specified net fitness flux for Figure 1, we varied s_b while holding U_b constant.

Our algorithm finds values of s_b that bracket the target net fitness flux, then uses a bisection method until

it finds a value of s_b that is within ± 0.00005 of the target. In practice, there was little stochasticity in the

regression slope (which averages out stochasticity in the timecourse), and so this relatively deterministic

method performed well.

Although the census population size N is a parameter of our model, the effective population size N_e is

not, but rather emerges over the course of a given simulation. To estimate it, we recorded tree-sequences

using the tskit package [42], then used msprime [63] to retroactively add neutral mutations after each

simulation. We did this only for one parameter combination involving realistically high N, due to the

significant computational cost of this procedure; this was 23 chromosomes, 50 linkage blocks per

chromosome, $N = 20,000$, $U_d = 2$, $U_b = 0.002$, and $s_b = 0.0025$. These parameter values produce

only a small excess of adaptation above that needed to counter Ohta's ratchet (Figure 1). We calculate N_e

using neutral heterozygosity under an infinite-alleles model. The choice of neutral mutation rate will not

affect estimated N_e, so long as it is small enough to avoid multiple mutations at the same site; we

arbitrarily chose 1.0×10^{-6} per linkage block, or 1.15×10^{-4} per haploid genome. This produced

$N_e \sim 7500$, on the order of effective population sizes inferred for ancestral human populations [64]. For

comparison, similar simulations with $U_b = 0$ (i.e. with background selection alone and declining relative

fitness), produce $N_e \sim 16,000$ [28]. This is consistent with other results, using the simulation approach

developed here, showing that N_e is primarily lowered by interactions between beneficial and deleterious

mutations [29].

Tree sequence recording also tracks all non-neutral mutations, so that we can identify those that fixed and

thus determine the degree of asymmetry in the effect sizes of fixed mutations. Note that without tree-

sequence recording, this information would be inaccessible due to the way we summarize the fitness of

many mutations within linkage blocks. However, using tree-sequence recording for all non-neutral

mutations significantly increases the computation time of simulations. For a representative case with $N =$

$20,000$, $U_b = 0.001$, $U_d = 2$, $s_b = 0.001$, 23 chromosomes with 50 linkage blocks each, and all other

details as described above, a single simulation of 200,000 generations took 12 hours, 45 minutes without

tree-sequence recording and 58 hours, 42 minutes with tree-sequence recording. When we are solving for

the parameters that produce a target value of net fitness flux, we therefore do not use tree sequence

recording.

Results

The mean beneficial effect size $\overline{s_b}$ is more important than the beneficial mutation rate U_b for achieving

net positive fitness flux (relatively horizontal contour of zero net flux in Figure 1, despite log scale on x-

axis). While there is great uncertainty in the true values of U_b and $\overline{s_b}$, the $\overline{s_b} > \sim 0.001 - 0.003$ range

required for population persistence seems entirely plausible, demonstrating that modest assumptions

about beneficial mutations are sufficient to rescue populations from Ohta's ratchet.

Figure 1. Relatively rare and mild beneficial mutations are sufficient to counteract a deluge of slightly deleterious mutations accumulating under Ohta's ratchet. Black line shows combinations of beneficial mutation parameters that produce zero net fitness flux. All populations simulated with $N = 20{,}000$, $U_d = 2$, and 23 chromosomes with 50 linkage blocks per chromosome. Red dashed lines show plausible upper and lower estimates of the mean effect size of new beneficial mutations in humans that we deemed *a priori* plausible; almost all of the resulting parameter value range produces positive net fitness flux. The range for U_b was chosen to be broad, including values that are likely too low.

Beneficial fixations are of much greater magnitude than deleterious fixations. In the representative case shown in Figure 2, fixations have $\bar{s}_b = 0.00423$ (more than four times as large as *de novo* beneficial mutations) and $\bar{s}_d = -0.0000391$ (almost three orders of magnitude smaller than the mean of -0.009 for *de novo* deleterious mutations). 56,255 deleterious and 6,387 beneficial fixations occurred over the 200,000 generations simulated for Figure 2.

Figure 2. Damage from small effect deleterious fixations is balanced by far fewer but larger effect beneficial fixations. The distribution of effect sizes of fixed mutations is shown in bins of 0.001 at the end of 200,000 generations, in a population with $N = 20,000$, $U_b = 0.001$, $U_d = 2$, $s_b = 0.001$, and 23 chromosomes with 50 linkage blocks each. Inset truncates the y-axis to see detail outside the mode of slightly deleterious fixations. Effect sizes are per allele copy.

Figure 3. Deleterious mutations appreciably but modestly slow adaptation, visualized as the number of

generations required for population mean fitness to increase by 10%. Black boxes indicate simulations

with net fitness flux < 0.

To adapt to a changing environment, a population must generate positive net fitness flux, i.e. beneficial

fixations above and beyond those required to counterbalance Ohta's ratchet. Rapid adaptation is shown in

purple in Figure 3, slow adaptation in green, and an inability even to counter deleterious fixations in

black. While resistance to mutational degradation remains reasonably robust to high U_d (few black

squares), the need to compensate for Ohta's ratchet drives down the rate of adaptation to a changing

external environment by an average of ~13%, ~35%, and ~55% for U_d of 2, 5, and 10, respectively,

relative to $U_d = 0$.

Population geneticists have raised concerns about the increase in the human mutation rate [1], in particular

due to increased age at paternity [2]. Mean paternal age in the U.S. increased from 27.4 to 30.9 years

between 1972 and 2015 [65], which would add about 7 *de novo* mutations to the 70-100 experienced by

typical human gametes, or a 7-10% increase in the mutation rate [66]. We simulated a 10% increase in the

mutation rate, for both deleterious and beneficial mutations, for a reference population with $U_b =$

$0.002, \bar{s}_b = 0.0025$, and other parameters, including baseline $U_d = 2$, as in Figure 1. Surprisingly,

populations with increased mutation rates took only 127 generations to increase their fitness by 10%,

compared to 151 generations for the baseline population. In other words, because beneficial fitness flux is

more sensitive to U_b than deleterious fitness flux is to U_d, increasing the total mutation rate helps the

population adapt faster. The counterintuitively increased rate of adaptation directly contradicts dysgenic

fears about the consequences of elevated mutation rates on mean population fitness load.

Beneficial fitness flux might be recombination limited [67]. Since most meioses involve exactly two

crossovers per chromosome [46], the primary determinant of *genome-wide* recombination rate (as opposed

to local recombination rates) is the number of chromosomes. Supplementary Figure 4 shows that adding

more chromosomes would not substantially speed human adaptation. However, having fewer than 10

chromosomes would substantially slow down adaptation, independently of mutation rate. Humans are

well above this threshold number of chromosomes, but a few of our primate relatives (the lemur

Lepilemur mustelinus, and the collared titi monkey *Cheracebus torquatus*) are not [68].

Figure 4. Higher deleterious mutation rates produce much more within-population variation in fitness,

shown here as the fold-change in fitness corresponding to one standard deviation in the natural logarithm

of fitness. The highest beneficial mutation rates and beneficial effect sizes we consider also affect within-

population variation, but to a much smaller degree.

Perhaps the most striking impact of high human $U_d \sim 2 - 10$ is to create high variance in load among

individuals. Figure 4 depicts this variance, given log-normally distributed fitness, in terms of the fold-

difference in fitness between two individuals that are one standard deviation apart on a log scale. This

difference of 15%-40% (in the historical human environment) is relatively insensitive to s_b and U_b, but

depends dramatically on U_d. Adaptation time is inversely proportional to census N, making its sensitivity

to U_d independent of N (Supplementary Table 1). The high variance among individuals engendered by

high U_d is nearly independent of N (Supplementary Table 2). High within-population variance can

alternatively be expressed as high mutation load $\frac{w_{best} - \bar{w}}{w_{best}}$.

Figure 5 shows analytical expectations of high load, in the absence of linkage disequilibrium, when

empirically inferred values of $\overline{s_d}$ (red shading [55,69]) as well as U_d are used. Grey shading shows how the

unrealistically low values of $\overline{s_d}$ used by Galeota-Sprung *et al.* [26] caused them to underestimate load. In

our simulations, linkage disequilibrium drives variance in fitness modestly higher again than analytical

expectations of $e^{U_d \overline{s_d}} - 1$ from Galeota-Sprung *et al.* [26] (Supplementary Figure 5). Very high values of

both U_b and s_b are needed for a substantial further increase in fitness variance (Supplementary Figure 5).

Figure 5. Mutation load $\frac{W_{best}-\overline{W}}{W_{best}}$ in humans is high in the absence of beneficial mutations, linkage

disequilibrium, or epistasis. The mean and variance in the number of deleterious polymorphisms per

individual are both U_d/s_d [70,71], and the resulting variance in fitness is $e^{U_d\overline{s_d}} - 1$ [26]. When mean

population fitness is 1, the best individual present then has fitness $W_{best} = e^{C\sqrt{U_d\overline{s_d}}}$. We approximate

$C \approx \mu_G + \gamma\sigma_G$ from a Gumbel distribution with location $\mu_G = F^{-1}(1 - \frac{1}{N})$ and scale parameter $\sigma_G =$

$F^{-1}\left(1 - \frac{1}{N}\frac{1}{e}\right) - \mu_G$ where F^{-1} is the normal distribution's quantile function, N is population size, γ is the

Euler-Mascheroni constant and e is Euler's number [26]. We use $N = 10,000$. Red shaded area corresponds

to $\overline{s_d}$ (or more strictly speaking, the heterozygotic effect size $\overline{sh}$) between 0.009 and 0.029, as inferred

for humans by Kim et al. [55] and Boyko et al. [69], respectively. Gray shaded area corresponds to values

considered by Galeota-Sprung et al. [26]. Charlesworth [72] assumed $sN\sim1$, black dashed line, corresponding

to similarly low $s\sim0.0001$ for humans.

Discussion

The accumulation of many slightly deleterious fixations, driven by a realistically high human genome-

wide deleterious mutation rate $U_d = 2 - 10$, is counteracted asymmetrically by a smaller number of

beneficial fixations of larger effect size. We simulate this balance in the presence of complex linkage

disequilibria, given conservatively low beneficial parameter values, combined with known human values

for recombination and deleterious effect sizes. While population persistence is achieved, large within-

population variance in fitness is created. Much of the resulting adaptation (as per Fisher's Fundamental

Theorem [73]) is absorbed by counteracting deleterious fixations, and so despite high fitness variance,

adaptation to a changing environment is 13-55% slower.

Understanding how mutation load is stabilized in humans and similar species is a precondition for

addressing a long-standing fear: that recent changes to human lifestyles or technology will push us to

higher load. For example, if the human mutation rate, beginning at an already critically high level, were to

increase due to increased paternal age, or if selection against deleterious mutations were relaxed due to

modern medicine, the perception has been that population mean fitness would decrease, potentially with

disastrous consequences [2,3,6]. Intriguingly, our results suggest that the expected increase in mutation rates

in human populations due to increased paternal age has the opposite effect, improving rather than

degrading population mean fitness.

High U_d does not threaten population persistence, but does have profound implications for within-

population variation among individuals. While a genotype whose load used to cause a ~30% reduction in

fitness in ancient human environments might now have a lesser impact on fitness, it likely still has a

significant impact on health. Indeed, variation in self-reported health has a substantial genetic component

[74], and load, as assessable from whole-genome sequencing, can be used to predict medically relevant

phenotypes [4,75]. High genetic variance among individuals is a hidden confounding variable in a vast range

of studies [76], including many studies of human health. Our theoretical assessment, when combined with

estimates of human $\overline{s_d}$, implies high variance in human mutation load, requiring a significant

reassessment across all public health studies grounded in correlational analysis [76]. A remaining caveat

here is our neglect of epistasis. Positive epistasis is expected to increase variance and negative epistasis to

decrease it [77].

High fitness variance (squared differences in fitness) is closely related to historical debates about high

mutation load (differences in fitness). Early discussions of high mutation load have been critiqued as

comparing mean fitness to that of a non-existing ideal genotype [78], one that represents invalid typological

thinking [79]. Correctly defining load as a comparison between mean population fitness and the best

genotype present avoids this criticism and necessarily reduces the estimated load, which has been

presented as a 'solution' to the mutation load 'problem' [26,72,78]. While Galeota-Sprung et al. [26] argued that

loads are small when thus conceived, their argument (see their Table 1) requires smaller $\overline{s_d}$ than that

inferred for humans (Figure 5).

Our simulations confirm that the solution to population persistence is a pattern of many small mutations,

each of which cannot be effectively cleared, being counteracted by compensatory mutations with global

effects. This pattern is part of drift barrier theory [23,24,80–87]. Drift barrier theory (Figure 6; blue circle)

emphasizes the causal importance of census population size in producing a ratchet of increasing

molecular and organismal complexity. Lynch [84] (see Chapter 4) posits that the minimum size of a

deleterious mutation that can be reliably purged ('effective population size') is driven (albeit not

exclusively) by census population size, which is in turn driven by life history traits such as body size. A

low effective population size that cannot purge small DNA insertions leads to a bloated genome, whose

complexity is posited to lead to larger body size and/or increased ecological specialization, reducing

census population size, closing the causal loop. Drift barrier theory sees increased mutation rate as a

consequence of relaxed selection against mutator alleles.

We suggest shifting the causal emphasis to high deleterious mutation rate instead of low census

population size. Indeed, a mutational ratchet (Figure 6, orange cycle) can occur even when census

population size is high. First, high $U_d > 1$ substantially reduces N_e through background selection (among

unlinked sites) [28,88] and via linkage disequilibrium with beneficial mutations [29]. As with drift barrier

theory, the resulting deluge of slightly deleterious fixations increases genome size [89] — note that a ratchet

will not occur if deleterious fixations decrease genome size, as is likely the case in prokaryotes [90–92]. The

feedback loop from there does not go through census N; instead, larger genomes create a larger target size

for deleterious mutations, directly increasing U_d.

Figure 6. A feedback loop of ratcheting complexity can be driven either by census size *N* and
ecology (drift barrier theory, blue) or by high deleterious mutation rate U_d (our view, orange and
pink). The drift barrier ratchet requires low census population size N, whereas our ratchet requires high
deleterious mutation rate U_d. Drift barrier theory emphasizes a causal link from N_e to U_d via relaxed
selection against mutators [84], whereas we emphasize background selection as a causal driver in the
opposite direction, i.e. from U_d to N_e. Mutational meltdown [93,94] is shown for completeness (green), since
its elements are already invoked.

The mutational ratchet described above (Figure 6, orange) might drive U_d up to a high enough level to

power the complexity ratchet (Figure 6, pink) that is the focus of this manuscript. Similarly to drift barrier

theory, when slightly deleterious fixations cannot be reversed to achieve detailed balance, they are

compensated for by large effect changes that frequently occur at a higher level of organization, creating a

complexity ratchet. The key difference is that the drift barrier theory (Figure 6, blue and pink) requires

low N but can occur at low U_d, so long as sN_e is low. Our alternative (Figure 6, orange and pink) requires

$U_d > 1$, and can occur even for high census N.

Previous hypotheses have focused on population size as the crucial difference between species that are

able to purge load within a small, streamlined genome (e.g. bacteria) vs. species forced into ratcheting

molecular complexity in search of innovative molecular solutions to stay ahead of perpetual degradation

(e.g. humans). But differences in U_d could also explain differences in molecular complexity across the

tree of life.

We confirmed that $U_d > 1$ need not cause population decline, because the accumulation of mildly

deleterious fixations (Ohta's ratchet) is countered by a smaller number of larger-effect beneficial fixations

under quite conservative assumptions. While $U_d > 1$ may therefore not threaten population persistence, it

may be an important driver of the evolution of molecular and organismal complexity. Combined with

realistic deleterious effect sizes, high U_d created high fitness variance within human populations in their

ancestral environment.

Code Availability

Simulation code in C is available at github.com/MaselLab/MutationLoad.

Acknowledgements

We thank Dr. Kun Xiong for helping the first author learn C, Dr. Walid Mawass for helpful discussions,

and Dr. Cristian Román-Palacios for assistance navigating the database of chromosome numbers across

the tree of life. This work was supported by the John Templeton Foundation [62028] and NIH Grant

GM084905. Figure 6 was created by Jennifer Yamnitz.

Bibliography

1.   Muller, H. Our load of mutations. *Am J Hum Genet* **2**, 111–176 (1950).

2.   Crow, J. F. The high spontaneous mutation rate: is it a health risk? *Proc Natl Acad Sci U S A* **94**,
8380–6 (1997).

3.   Lynch, M. Mutation and human exceptionalism: Our future genetic load. *Genetics* **202**, 869–875
(2016).

4.   Vy, H. M. T., Jordan, D. M., Balick, D. J. & Do, R. Probing the aggregated effects of purifying
selection per individual on 1,380 medical phenotypes in the UK biobank. *PLoS Genet* **17**, (2021).

5.   Lynch, M. Rate, molecular spectrum, and consequences of human mutation. *Proc Natl Acad Sci U
S A* **107**, 961–8 (2010).

6.   Hamilton, W. D. The hospitals are coming. in *Narrow roads of gene land* vol. 2 449–536 (Oxford
University Press, 1999).

7.   Lesecque, Y., Keightley, P. D. & Eyre-Walker, A. A resolution of the mutation load paradox in
humans. *Genetics* **191**, 1321–1330 (2012).

8.   Wang, Y. & Obbard, D. J. Experimental estimates of germline mutation rate in eukaryotes: a
phylogenetic meta-analysis. *Evol Lett* **7**, 216–226 (2023).

9.   Kondrashov, A. S. *Crumbling Genome: The Impact of Deleterious Mutations on Humans*. (John
Wiley & Sons, Inc, Hoboken, NJ, 2017).

10.   Reed, F. A., Akey, J. M. & Aquadro, C. F. Fitting background-selection predictions to levels of
nucleotide variation and divergence along the human autosomes. *Genome Res* **15**, 1211–1221
(2005).

11.   Haag-Liautard, C. *et al.* Direct estimation of per nucleotide and genomic deleterious mutation
rates in Drosophila. *Nature* **445**, 82–85 (2007).

12.   Kondrashov, A. S. Contamination of the genome by very slightly deleterious mutations: why have
we not died 100 times over? *J Theor Biol* **175**, 583–594 (1995).

13.   Teicher, A. Caution, overload: The troubled past of genetic load. *Genetics* **210**, 747–755 (2018).

14.   Paul, D. B. 'Our Load of Mutations' Revisited. *J Hist Biol* **20**, 321–335 (1987).

15.   Ohta, T. The Nearly Neutral Theory of Molecular Evolution. *Annu Rev Ecol Syst* **23**, 263–286
(1992).

16.   Ohta, T. Slightly Deleterious Mutant Substitutions in Evolution. *Nature* **246**, 96–98 (1973).

17. Ruan, Y., Wang, H., Zhang, L., Wen, H. & Wu, C.-I. Sex, fitness decline and recombination –
Muller's ratchet vs. Ohta's ratchet. *bioRxiv* 2020.08.06.240713 (2020)
doi:10.1101/2020.08.06.240713.

18. Schultz, S. T. & Lynch, M. Mutation and extinction: The role of variable mutational effects,
synergistic epistasis, beneficial mutations, and degree of outcrossing. *Evolution (N Y)* **51**, 1363–
1371 (1997).

19. Whitlock, M. C. Fixation of new alleles and the extinction of small populations: drift load,
beneficial alleles, and sexual selection. *Evolution (N Y)* **54**, 1855–1861 (2000).

20. Whitlock, M. C., Griswold, C. K. & Peters, A. D. Compensating for the meltdown: The critical
effective size of a population with deleterious and compensatory mutations. *Ann Zool Fennici* **40**,
169–183 (2003).

21. Kimura, M. On the probability of fixation of mutant genes in a population. *Genetics* **47**, 713–719
(1962).

22. Haldane, J. A mathematical theory of natural and artificial selection, part V: selection and
mutation. *Proc. Camb. Philos. Soc.* **23**, 838–844 (1927).

23. Fares, M. A., Ruiz-González, M. X., Moya, A., Elena, S. F. & Barrio, E. GroEL buffers against
deleterious mutations. *Nature* **417**, 398 (2002).

24. Wu, X. & Hurst, L. D. Why Selection Might Be Stronger When Populations Are Small: Intron
Size and Density Predict within and between-Species Usage of Exonic Splice Associated cis-
Motifs. *Mol Biol Evol* **32**, 1847–1861 (2015).

25. Koelle, K. & Rasmussen, D. A. The effects of a deleterious mutation load on patterns of influenza
A/H3N2's antigenic evolution in humans. *Elife* **4**, e07361 (2015).

26. Galeota-Sprung, B., Sniegowski, P. & Ewens, W. Mutational Load and the Functional Fraction of
the Human Genome. *Genome Biol Evol* **12**, 273–281 (2020).

27. Hill, W. G. & Robertson, A. Linkage disequilibrium in finite populations. *TAG Theoretical and
Applied Genetics* **38**, 226–231 (1968).

28. Matheson, J. & Masel, J. Background Selection From Unlinked Sites Causes Nonindependent
Evolution of Deleterious Mutations. *Genome Biol Evol* **16**, (2024).

29. Mawass, W., Matheson, J., Hernández, U., Berg, J. & Masel, J. Extinction vortices are driven
more by a shortage of beneficial mutations than by deleterious mutation accumulation. *biorxiv*
(2024) doi:10.1101/2024.10.25.620329.

30. Simons, Y. B., Turchin, M. C., Pritchard, J. K. & Sella, G. The deleterious mutation load is
insensitive to recent population history. *Nat Genet* **46**, 220–224 (2014).

31. Kyriazis, C. C., Wayne, R. K. & Lohmueller, K. E. Strongly deleterious mutations are a primary
determinant of extinction risk due to inbreeding depression. *Evol Lett* **5**, 33–47 (2021).

32. Pérez-Pereira, N., Wang, J., Quesada, H. & Caballero, A. Prediction of the minimum effective size
of a population viable in the long term. *Biodivers Conserv* **31**, 2763–2780 (2022).

33.　Bertorelle, G. *et al.* Genetic load: genomic estimates and applications in non-model animals. *Nat*
Rev Genet **23**, 492–503 (2022).

34.　Liu, L. *et al.* Genetic consequences of long-term small effective population size in the critically
endangered pygmy hog. *Evol Appl* **14**, 710–720 (2021).

35.　Charmouh, A. P., Reid, J. M., Bilde, T. & Bocedi, G. Eco-evolutionary extinction and
recolonization dynamics reduce genetic load and increase time to extinction in highly inbred
populations. *Evolution (N Y)* **76**, 2482–2497 (2022).

36.　Pérez-Pereira, N., López-Cortegano, E., García-Dorado, A. & Caballero, A. Prediction of fitness
under different breeding designs in conservation programs. *Anim Conserv* **26**, 86–102 (2023).

37.　Domínguez-García, S., García, C., Quesada, H. & Caballero, A. Accelerated inbreeding
depression suggests synergistic epistasis for deleterious mutations in Drosophila melanogaster.
Heredity (Edinb) **123**, 709–722 (2019).

38.　Gravel, S. When is selection effective? *Genetics* **203**, 451–462 (2016).

39.　Lohmueller, K. E. *et al.* Proportionally more deleterious genetic variation in European than in
African populations. *Nature* **451**, 994–997 (2008).

40.　Koch, E. & Novembre, J. A temporal perspective on the interplay of demography and selection on
deleterious variation in humans. *G3: Genes, Genomes, Genetics* **7**, 1027–1037 (2017).

41.　Haller, B. C. & Messer, P. W. SLiM 2: Flexible, interactive forward genetic simulations. *Mol Biol*
Evol **34**, 230–240 (2017).

42.　Kelleher, J., Thornton, K. R., Ashander, J. & Ralph, P. L. Efficient pedigree recording for fast
population genetics simulation. *PLoS Comput Biol* **14**, e1006581 (2018).

43.　Simmons, M. J. & Crow, J. F. MUTATIONS AFFECTING FITNESS IN DROSOPHILA
POPULATIONS. *Annu Rev Genet* **11**, 49–78 (1977).

44.　Yang, J. *et al.* Incomplete dominance of deleterious alleles contributes substantially to trait
variation and heterosis in maize. *PLoS Genet* **13**, e1007019- (2017).

45.　Agrawal, A. F. & Whitlock, M. C. Inferences About the Distribution of Dominance Drawn From
Yeast Gene Knockout Data. *Genetics* **187**, 553–566 (2011).

46.　Pardo-Manuel De Villena, F. & Sapienza, C. Recombination is proportional to the number of
chromosome arms in mammals. *Mammalian Genome* **12**, 318–322 (2001).

47.　Weissman, D. B. & Hallatschek, O. The rate of adaptation in large sexual populations with linear
chromosomes. *Genetics* **196**, 1167–1183 (2014).

48.　Neher, R. A., Kessinger, T. A. & Shraiman, B. I. Coalescence and genetic diversity in sexual
populations under selection. *Proc Natl Acad Sci U S A* **110**, 15836–15841 (2013).

49.　Good, B. H., Walczak, A. M., Neher, R. A. & Desai, M. M. Genetic Diversity in the Interference
Selection Limit. *PLoS Genet* **10**, (2014).

50.　Wall, J. D. & Pritchard, J. K. Haplotype blocks and linkage disequilibrium in the human genome.
Nat Rev Genet **4**, 587–597 (2003).

51.   Belmont, J. W. *et al.* A haplotype map of the human genome. *Nature* **437**, 1299–1320 (2005).

52.   Altshuler, D., Daly, M. J. & Lander, E. S. Genetic mapping in human disease. *Science (1979)* **322**,
442   881–888 (2008).

443   53.   Coop, G., Wen, X., Ober, C., Pritchard, J. K. & Przeworski, M. High-resolution mapping of
444   crossovers reveals extensive variation in fine-scale recombination patterns among humans. *Science*
445   *(1979)* **319**, 1395–1398 (2008).

446   54.   Pratto, F. *et al.* Erratum Recombination initiation maps of individual human genomes (Science
447   DOI: 10.1126/science.1256442). *Science (1979)* **362**, (2018).

448   55.   Kim, B., Huber, C. & Lohmueller, K. Inference of the Distribution of Selection Coefficients for
449   New Nonsynonymous Mutations Using Large Samples. *Genetics* **206**, 345–361 (2017).

450   56.   Thornton, K. R. A c++ template library for efficient forward-time population genetic simulation of
451   large populations. *Genetics* **198**, 157–166 (2014).

452   57.   Thornton, K. R. Polygenic adaptation to an environmental shift: Temporal dynamics of variation
453   under Gaussian stabilizing selection and additive effects on a single trait. *Genetics* **213**, 1513–
454   1530 (2019).

455   58.   Haller, B. C. & Messer, P. W. SLiM 4: Multispecies Eco-Evolutionary Modeling. *Am Nat* **201**,
456   E127–E139 (2022).

457   59.   Fenwick, P. M. A new data structure for cumulative frequency tables. *Softw Pract Exp* **24**, 327–
458   336 (1994).

459   60.   Bertram, J. & Masel, J. Density-dependent selection and the limits of relative fitness. *Theor Popul*
460   *Biol* **129**, 81–92 (2019).

461   61.   Matheson, J., Exposito-Alonso, M. & Masel, J. Substitution load revisited: a high proportion of
462   deaths can be selective. *Genetics* (2025) doi:10.1093/genetics/iyaf011.

463   62.   Ober, C., Hyslop, T. & Hauck, W. W. Inbreeding effects on fertility in humans: Evidence for
464   reproductive compensation. *Am J Hum Genet* **64**, 225–231 (1999).

465   63.   Baumdicker, F. *et al.* Efficient ancestry and mutation simulation with msprime 1.0. *Genetics* **220**,
466   iyab229 (2022).

467   64.   Tenesa, A. *et al.* Recent human effective population size estimated from linkage disequilibrium.
468   *Cold Spring Harbor Laboratory Press Human* **2**, 520–526 (2007).

469   65.   Khandwala, Y. S., Zhang, C. A., Lu, Y. & Eisenberg, M. L. The age of fathers in the USA is
470   rising: An analysis of 168 867 480 births from 1972 to 2015. *Human Reproduction* **32**, 2110–2116
471   (2017).

472   66.   Jónsson, H. *et al.* Parental influence on human germline de novo mutations in 1,548 trios from
473   Iceland. *Nature* **549**, 519–522 (2017).

474   67.   Weissman, D. B. & Barton, N. H. Limits to the rate of adaptive substitution in sexual populations.
475   *PLoS Genet* **8**, (2012).

476   68.   Román-Palacios, C., Medina, C. A., Zhan, S. H. & Barker, M. S. Animal chromosome counts
477   reveal a similar range of chromosome numbers but with less polyploidy in animals compared to
478   flowering plants. *J Evol Biol* **34**, 1333–1339 (2021).

479   69.   Boyko, A. R. *et al.* Assessing the evolutionary impact of amino acid mutations in the human
480   genome. *PLoS Genet* **4**, (2008).

481   70.   Kondrashov, A. S. Dynamics of unconditionally deleterious mutations: Gaussian approximation
482   and soft selection. *Genet Res* **65**, 113–121 (1995).

483   71.   Maynard Smith, J. *The Evolution of Sex*. vol. 4 (Cambridge University Press Cambridge, 1978).

484   72.   Charlesworth, B. Why we are not dead one hundred times over. *Evolution (N Y)* **67**, 3354–3361
485   (2013).

486   73.   Fisher, R. *The Genetical Theory of Natural Selection*. (Clarendon Press, Oxford, 1930).

487   74.   Romeis, J. C. *et al.* Heritability of Self-Reported Health. *Health Serv Res.* **35**, 995–1010 (2000).

488   75.   Fiziev, P. P. *et al.* Rare penetrant mutations confer severe risk of common diseases. *Science (1979)*
489   **380**, eabo1131 (2023).

490   76.   Harden, K. P. *The Genetic Lottery: Why DNA Matters For Social Equality*. (Princeton University
491   Press, Princeton, New Jersey, 2021).

492   77.   Kimura, M. & Maruyama, T. THE MUTATIONAL LOAD WITH EPISTATIC GENE
493   INTERACTIONS IN FITNESS. *Genetics* **54**, 1337–1351 (1966).

494   78.   Agrawal, A. F. & Whitlock, M. C. Mutation Load: The Fitness of Individuals in Populations
495   Where Deleterious Alleles Are Abundant. *Annu. Rev. Ecol. Evol. Syst* **43**, 115–35 (2012).

496   79.   Soulier, A. 'Genetic Load': How the Architects of the Modern Synthesis Became Trapped in a
497   Scientific Ideology. *Transversal: International Journal for the Historiography of Science* 118–139
498   (2018) doi:10.24117/2526-2270.2018.i4.11.

499   80.   Rajon, E. & Masel, J. Evolution of molecular error rates and the consequences for evolvability.
500   *Proceedings of the National Academy of Sciences* **108**, 1082–1087 (2011).

501   81.   Xiong, K., Mcentee, J. P., Porfirio, D. J. & Masel, J. Drift barriers to quality control when genes
502   are expressed at different levels. *Genetics* **205**, 397–407 (2017).

503   82.   Lynch, M. Evolution of the mutation rate. *Trends in Genetics* **26**, 345–352 (2010).

504   83.   Sung, W., Ackerman, M. S., Miller, S. F., Doak, T. G. & Lynch, M. Drift-barrier hypothesis and
505   mutation-rate evolution. *Proceedings of the National Academy of Sciences* **109**, 18488–18492
506   (2012).

507   84.   Lynch, M. *The Origins of Genome Architecture*. (Sindauer Associates, Sunderland MA, 2007).

508   85.   Frank, S. A. Maladaptation and the paradox of robustness in evolution. *PLoS One* **2**, (2007).

509   86.   Gros, P.-A. & Tenaillon, O. Selection for Chaperone-Like Mediated Genetic Robustness at Low
510   Mutation Rate: Impact of Drift, Epistasis and Complexity. *Genetics* **182**, 555–564 (2009).

87.   Hurst, L. D. *The Evolution of Imperfection*. (Princeton University Press, 2025).

88. Charlesworth, B. The Effects of Deleterious Mutations on Evolution at Linked Sites. *Genetics* **190**,
5–22 (2012).

89. Lefébure, T. *et al.* Less effective selection leads to larger genomes. *Genome Res* **27**, 1016–1028
(2017).

90. Kuo, C.-H. & Ochman, H. Deletional Bias across the Three Domains of Life. *Genome Biol Evol* **1**,
145–152 (2009).

91. Bourguignon, T. *et al.* Increased Mutation Rate Is Linked to Genome Reduction in Prokaryotes.
Current Biology **30**, 3848-3855.e4 (2020).

92. Bingham, E. P. & Ratcliff, W. C. A nonadaptive explanation for macroevolutionary patterns in the
evolution of complex multicellularity. *Proc Natl Acad Sci U S A* **121**, (2024).

93. Lynch, M., Conery, J. & Burger, R. Mutational Meltdowns in Sexual Populations. *Evolution (N Y)*
49, 1067–1080 (1995).

94. Lande, R. RISK OF POPULATION EXTINCTION FROM FIXATION OF NEW
DELETERIOUS MUTATIONS. *Evolution (N Y)* **48**, 1460–1469 (1994).

526