Eavesdropping and audience effect in fighting fruit flies reveal feedback between social information

use and production

Marina Hutchins*[1], Julia Saltz[1]

1 Rice University, Department of Biosciences, Houston (TX), USA

* Corresponding author, meh18@rice.edu

ORCID: M. Hutchins: 0000-0001-5164-0129, J.Saltz: 0000-0002-0645-173X

Abstract: Social information use and production occur simultaneously during social learning events;

however, these two processes are often studied independently. We propose that social information use

and production represent a feedback loop: eavesdroppers influence the behaviors of those being

observed, which then alters the eavesdroppers' social decisions. Here, we used *Drosophila*

melanogaster to investigate the relationship between social information use and production in an

aggressive context. We tested whether paired males altered their aggression when they were observed

by (i) a male; (ii) a female, or; (iii) no one. Next, we allowed each individual eavesdropper to interact

with the paired males that they just observed, and measured their social behaviors. In addition, we

tested how naïve flies responded when interacting with paired males. We found that paired males

reduced their aggression when observed by a male, representing a change in social information

production. In response, male eavesdroppers were overall less aggressive towards the more aggressive

of the paired males they had observed, but their response to social information was also modulated by

their genotype. In contrast, female eavesdroppers did not elicit changes to the behavior of the paired

males, nor did we find evidence that they used the social information collected to make mating

decisions. Overall, we found support for the feedback loop in males, but not females: information was

exchanged bi-directionally during eavesdropping events, resulting in the modification of behavior by all

parties. Utilizing the social information feedback framework can contribute to a better understanding of

how social behaviors and animal communication are expressed and evolve.

Keywords: aggression, audience effect, Drosophila melanogaster, eavesdropping, social information

Competing Interests: None

Significance Statement: Animals often observe the social interactions that occur between others, but

the effects of both watching and being watched on behavior are rarely studied together. Using a novel

approach, we studied how an audience influenced and learned from an aggressive contest in fruit flies.

A male audience caused contest participants to decrease their aggression, and the male who was

watching later showed decreased aggression toward the contest winner. Surprisingly, we did not see

any effect of a female audience on male-male aggression, and females' later behavior was not altered by

observing a contest. Together, these results show how observing a fight led to changes in both the

behavior of the contest participants and the male audience, thus suppressing group levels of aggressive

behavior. Accounting for this interaction between social information use and production is necessary to

understand social behaviors.

Introduction

Introduction

Individuals are able to acquire social information by detecting information about the behavioral

interactions of others through a variety of sensory modalities and mechanisms (Heyes, 1994; McGregor

& Peake, 2000; Wagner & Danchin, 2010). We refer to these third-party conspecifics as 'social

eavesdroppers': individuals who change their behavior after having access to information about the

interactions of others. (Bonnie & Earley, 2007; Dall et al., 2005; Danchin et al., 2004; McGregor, 2005,

Tibbets et al., 2020). Social eavesdropping could provide a low-risk alternative to direct interactions with

others, as eavesdroppers can learn about the quality or ability of another individual without engaging in

interactions that could be risky or costly (McGregor & Peake, 2000; Mery et al., 2009). For instance,

females of many species have been shown to eavesdrop on the mate choice decisions of other females

and then copy those preferences, an example of eavesdropping as a cost-avoidance strategy (Vakirtzis,

2011).

In many cases, an interacting pair can detect the presence and sometimes the characteristics of a social

eavesdropper, which may lead to a shift in behavior known as the 'audience effect' (Fichtel & Manser,

2010; McGregor, 2005, Coppinger et al. 2017). The audience effect is widespread and well-documented

in a variety of animal species, including fish (Doutrelant et al., 2001; Plath et al., 2008), birds (White et

al., 2002), primates (Townsend & Zuberbuhler, 2009), and humans (Bond & Titus, 1983). For example,

substantial work has demonstrated that in primates, the intensity and/or frequency of alarm calls

performed by an individual depends on the characteristics of the audience members around them

(Chapman et al., 1990; Pollick et al., 2005; Wheeler, 2008).

While social eavesdropping and the audience effect are two well-described phenomena, they have

historically been studied as two distinct processes (McGregor, 2005). Yet, social eavesdropping and the

audience effect can occur in tandem during social learning events (Bonnie & Earley, 2007; Chimento et

al., 2022; Cruz & Oliveira, 2015; Earley & Dugatkin, 2002). During these events, the expression of social

behaviors – and thus, the production of social information – may be altered depending on the presence

and identity of the eavesdropping individual, i.e., via audience effects (Desjardins et al., 2012;

Fitzsimmons & Bertram, 2013; Plath et al., 2008). At the same time, the social eavesdropper can use the

social information by changing their own behavior depending on what they observed from the

interacting pair (Chimento et al., 2022; Peake et al., 2001). Taken together, we can hypothesize a

feedback loop between social information production and use, in which the presence of a social

eavesdropper alters the social environment of the interacting pair, which in turn influences the behavior

of both the interactants and the eavesdropper.

The existence of a feedback loop between the production and use of social information has the

potential to alter our current understanding of how social behaviors are expressed and have evolved,

especially in group contexts. If there is no feedback between these processes, then research dedicated

to studying social information use independently from social information production would be sufficient

to capture the key social dynamics that emerge from these two phenomena. However, if a feedback

loop does exist, then studies focused on only one-half of the feedback loop at a time - i.e., studying

interacting pairs without eavesdroppers or vice versa - ignore critical participants during these

interactions. Under feedback loops, bi-directional information exchange between the social

eavesdroppers and the observed pair has the potential to modify the behavior of all parties, thus

influencing our predictions about social learning and animal communication in social groups. For

instance, early work investigating social information use in female cowbirds initially found that rearing

environment did not influence females' preference for male song between two cowbird subspecies

(King and West 1983). However, researchers later discovered that females modified the social

environment that they were reared in –an unaccounted for audience effect that altered their

interpretation of their findings (King & West 1987). Indeed, recent theory has highlighted the need to

investigate how variation in social information use and production can influence decision-making by

incorporating theory and data about both audience effect and social eavesdropping simultaneously

(Chimento et al., 2022; Turner et al., 2023).

In animal societies, aggression is an important social signaling behavior that influences social

relationships, foraging, and competition (King, 1973; Duque-Wilckens et al., 2019). Aggression often

takes place in an environment where social eavesdroppers are present and thus may be able to observe

contests between others and extract information about the interactants' fighting ability (Oliveira et al.,

1998; Peake et al., 2001). Previous work in a variety of organisms has shown that social eavesdroppers

can obtain information from the aggressive behaviors performed during a fight, such as threat displays

or overt physical interactions, or the contest outcome and dynamics, such as the identity of the winner

and loser (Johnstone, 2001; Magnhagen, 2006; Oliveira et al., 1998; Tibbetts et al., 2020; Van Breukelen

& Draud, 2005).

Decades of research, primarily in fish, has been dedicated to investigating how individuals change their

aggressive behavior in the presence of an audience, or how the outcome of a contest affects the future

behavior of social eavesdroppers(see Peake & Mcgregor, 2004 for review). However, there are currently

scant empirical studies identifying feedbacks, if any, between social information production and use in

an aggressive context. For example, many studies of social eavesdropping are conducted by allowing an

individual to observe a fight between others, while preventing the eavesdropper from being detected by

the demonstrators (e.g., using one-way mirrors; Magnhagen, 2006; Oliveira et al., 1998; Peake et al.,

2006). Making the eavesdropper undetectable to the interacting pair eliminates any possible audience

effects; however, in many circumstances, the interacting individuals often can detect and respond to the

presence of the eavesdropper, and *vice versa* (Zuberbühler, 2008). Thus, these experiments are not fully

capturing and accounting for the information transfer that happens in social groups in the wild during

social eavesdropping events.

Here, we utilized the social information feedback framework to investigate social decision-making in the

context of aggressive interactions. To investigate the proposed feedback loop between social

information production and use, we combined standard methods used for testing how individuals

produce and how individuals use social information into a single experiment. In the first phase of each

trial, we allowed individuals to eavesdrop on two males engaged in an aggressive contest – this

observation phase allowed us to test for variation in how the paired males produced social information.

In the second phase of the trial, the social eavesdroppers could then interact with both aggressors

directly, which allowed us to test how the produced social information influenced the eavesdropper's

behavior. Measuring both social information use and production in our experimental trials allowed us to

directly examine how an individual's ability to influence the production of social information results in

differences in social information use. We repeated this approach with 3 different genotypes of social

eavesdroppers, which allowed us to test for differences in access to social information production and

the robustness of the effect of social information use across different genotypes.

We predict that the presence, sex, and genotype of an eavesdropper will drive changes in the behavior

of the interacting aggressive pair (Fitzsimmons & Bertram, 2013; Matos & McGregor, 2002). We further

hypothesize that the social information observed by the eavesdroppers will alter their behaviors (i.e.,

aggression in males, or mating in females), towards one or more of the paired males (Aquiloni et al.,

2008; Chan et al., 2008; Magnhagen, 2006). Observing both of these results would support the existence

of a feedback loop between these processes. Further, any of these processes may be modified by the

genotype of the eavesdropper.

Methods

Study system

Drosophila melanogaster, the fruit fly, is an excellent study system for investigating the relationships

between social information use and production because of their well-characterized suite of behavioral

interactions and amenability to rigorous laboratory experiments. *D.melanogaster* forms social groups on

their food sources both in nature and in the lab, and social interactions, such as aggression, take place in

an environment with ample conspecific social eavesdroppers (Saltz, 2013; Stamps et al., 2005; Wertheim

et al., 2006). Fruit flies have a well-studied repertoire of aggressive behaviors (Chen et al., 2002; Saltz,

2013), and the outcomes of aggressive contests can influence later behavior (Trannoy et al., 2016). Fruit

flies also have the necessary cognitive capacity for social information use (Danchin et al., 2018; Mery et

al., 2009; Nöbel et al., 2018).

Genotypes and Rearing

We used inbred fly lines that were originally derived from a wild population in Raleigh, NC (MacKay et

al., 2012). We used 3 different genotypes of social eavesdroppers: 136, 229, and 429 (note that

genotypes are "named" using numbers, and their numbers are arbitrary and have no numeric meaning).

We standardized the genotypes of the paired males to ensure that we could directly link any effect of

eavesdropper sex and presence to the interacting pair's social information production. For the

standardized paired males, we used one male from genotype 365 and one male from genotype 765.

(Again, genotype numbers are arbitrary and have no further significance). Flies for this study were

reared in vials under standard conditions that minimized variation in larval density: 10 females and 10

males were placed into a vial containing standard fly food medium. Unmated progeny from these vials

were then collected the morning that they eclosed, and subsequently housed in individual vials to

prevent any social experiences. Flies were aged for 3 or 4 days before experimental trials and were

maintained on a 12:12 hour light:dark cycle.

Behavioral Arena

We utilized a novel arena design to test for variation in social information use and production (Figure 1).

The main part of the arena was composed of two Petri dishes (3 cm x 1 cm) taped together, with a small

food patch (1.5 cm x 1 cm) placed on the bottom of the arena, which contained standard fly food

medium and a dot of active yeast. There were two small holes in the arena: one on the side of the

arena, for the insertion of flies via aspiration, and another hole in the center of the top of the arena. A

pipette tip with a mesh base was placed over this hole, creating an observation tower for the

eavesdropper fly to watch aggressive contests. This arena design mimics the perching behaviors of flies

in natural conditions (Stamps et al., 2005). The mesh barrier that separated the eavesdropper fly from

the paired males allowed visual, olfactory, and/or auditory information to be extracted from the

aggressive contest (Versteven et al., 2017; Wang & Anderson, 2010). The location of the observation

tower was directly over the food patch, where most aggressive interactions occur (Hoffmann &

Cacoyianni, 1990). We observed occasional interactions at this mesh between the eavesdropper fly and

the interacting pair, however, we did not observe any aggressive or courtship behaviors between the

eavesdroppers and any member of the interacting pair.

Overview of trials

All flies utilized for the trials had no prior social experience, as they were socially isolated from the time

of their eclosion until trials were conducted (see Genotypes and Rearing). Each trial consisted of two

phases: the observation phase and the interaction phase. In the observation phase, an eavesdropper fly

was placed into the observation tower while the paired males in the main part of the arena engaged in

an aggressive contest. In the interaction phase, the same individual eavesdropper fly was added to the

arena with the paired males they had just observed, and their social interactions with the

males were recorded (Figure 1). After each trial was completed, all flies were discarded. Thus, each

individual fly (eavesdropper or member of the interacting pair) was used in both phases of a single trial,

and not reused in any other trials.

A   **B**

Figure 1. (A) The behavioral arena used to conduct experimental trials. The pipette tip on top housed an eavesdropper fly; the

opening was then covered to prevent them from escaping. The paired males were inserted into the petri dish arena with the

blue cap that was filled with standard food and topped with yeast paste. (B) Timeline overview of protocol for an experimental

trial.

Experimental Protocol

We randomly dusted the paired males with colored powder (orange or yellow) to create two distinct

male phenotypes. This allowed human observers to easily differentiate between the two flies for scoring

aggressive behavior, and this distinctive coloring also may have aided the eavesdropper flies in

distinguishing between the two males (E. Danchin et al., 2018; Mery et al., 2009; Monier et al., 2018).

We did not find any effect of color on male aggression or female mate choice (see Model Covariates in

the Supplemental Material). The paired males were inserted into the main petri dish arena via

aspiration, and the eavesdropper fly was placed into the observation tower. Flies were allowed to

acclimate for five minutes before the observation phase of the experimental trials began.

During the observation phase of the experiment, we measured aggressive behaviors performed by the

paired males for 30 minutes. We recorded five measures of aggressive behavior: lunging, fencing (both

low- and high-level fencing), offensive wing threats, chasing, and boxing, as defined in Chen et al., 2002.

Every two minutes, a human observer recorded all aggressive behaviors performed by the two male flies

in the arena for 20 seconds. During this observation phase, the eavesdropper fly was able to observe the

aggressive contest without interacting directly with the two males.

Social eavesdroppers rarely have the opportunity to make decisions about who to interact with (but see

studies on mate choice in an aggressive context, Van Breukelen & Draud, 2005). Rather, experimental

designs frequently designate the eavesdropping individuals to be randomly paired with only one of the

contest participants (Earley & Dugatkin, 2002; Johnsson & Åkerman, 1998; Oliveira et al., 1998; Tibbetts

et al., 2020). These designs prevent us from gaining insight into how individuals can process and utilize

social information to make decisions about how they will behave toward other group members. Recent

calls for the need to explain and understand social decision-making, especially in conflict settings, can be

solved by utilizing an experimental design that gives the eavesdropper the choice of interacting with

both contest participants (Hobson, 2020). Thus, immediately following the observation phase, we began

the interaction phase of the experiment by inserting the eavesdropper fly into the same arena with the

two demonstrator males that they just observed. We utilized AKASO cameras (EK700) to video record

the interaction phase, and then used BORIS, an event logging software, to code the social interactions

that occurred between the eavesdropper and the paired males for another 30 minutes (Friard & Gamba,

2016).

On each experimental day, we were able to conduct 8 trials across 2 time blocks: 4 trials began

approximately 15 minutes after subjective dawn, and another 4 trials began approximately 1 hour after

subjective dawn. Trials were able to be conducted simultaneously through the use of scan sampling

(figure 2) during the observation phase, and the use of video recording during the interaction phase,

which did not require the presence of a human observer beyond the initial setup. Time block was not

found to have a significant effect on male aggression or female mate choice (see Model Covariates in

the Supplemental Material). Observers were blind to treatment when using videos to code the social

behaviors in the interaction phase, however, the scoring of aggressive behavior during the observation

phase was not conducted blind in regards to treatment, as the presence or absence of an eavesdropper

fly made this impossible.

Treatments

We conducted 166 trials over three different experimental treatments, which are explained further

below. In 35 trials, we had one or more flies escape after the observation phase, but before or during

the interaction phase. When this occurred, we retained the data that we had collected for the

observation phase.

Female Eavesdropper. In this treatment, we used a female fly as the eavesdropper. For the observation

phase of the experiment, a female fly observed an aggressive encounter between the paired males

(n=41 trials). For the interaction phase of the experiment, we recorded whether the female mated and,

if so, which male the female chose to mate with (n=38 trials).

Male Eavesdropper. In this treatment, we used a male fly as the eavesdropper. For the observation

phase of the experiment, a male fly observed an aggressive encounter between the paired males (n=43

trials). For the interaction phase of the experiment, we recorded all lunging behavior that the male

eavesdropper displayed while interacting with the two males (n=32 trials). We measured only lunging

behavior during the second phase of the experiment, as more subtle aggressive behaviors were difficult

to confidently assign from our relatively low-resolution videos. We note that lunging behavior is

significantly correlated with our four other measures of aggressive behavior in the observation phase of

the experiment (see Supplementary Figure 1). Furthermore, measuring only lunging behavior has

previously been shown to be an accurate characterization of broad aggressive patterns in male fruit flies

(Saltz, 2013).

No Eavesdropper/No Social Information. In this treatment, there was no eavesdropper present for the

observation phase of the experiment, providing us with a baseline measure of how the paired males

would behave without an audience present (n=82 trials). In the interaction phase of the experiment, a

male (n=32 trials) or female fly (n=38 trials) who did not observe the aggressive encounter was inserted

into the arena. The same behaviors were recorded as described in the male and female eavesdropper

treatments, depending on the sex of the fly that was added.

Female Eavesdropper Follow Up – No Variation in Paired Male Genotype. After completing our female

trials, we observed that there was a strong bias in female mating preferences: in the 59 mating events

that occurred, females mated with the genotype 365 male in 51 of those trials. This bias for 365 males

obscured any effects of social information on mate choice preferences. As a result, we did a follow-up

experiment, where we conducted the female eavesdropper (n=47) and no eavesdropper trials (n=38)

using the same protocol as described above, except we used two males from genotype 365 as the paired

males engaged in an aggressive encounter. Using this approach, the only difference between the males

was their behavior during the observation phase, thus clarifying the effects of social information on

female mate choice.

Ethical Note

No animal ethics approval was required for this study, however, flies were handled with care at all times

to minimize stress and damage. We used anesthesia to collect flies, and allowed them to recover for 3 or

4 days before experimental trials. During trials, we handled flies using gentle aspiration. After trials were

complete, all flies were euthanized by being placed in a freezer.

Analysis

The goal of our analysis was to investigate how individuals influenced (observation phase) and used

(during the interaction phase) social information. To accomplish this, we created generalized linear

mixed models using the glmmTMB package in R (v4.1.0, R Core Team, 2021, Bolker et al., 2009).

Generalized linear mixed models allow us to fit a variety of error distributions to our data, include

random effects in our models, and account for zero-inflated data (Brooks et al., 2017). We created

models that tested how the paired males produced social information, how males used social

information, and how females used social information. To investigate how social information altered

social decision-making, we quantified which of the paired males performed the most and least

aggressive behaviors during the observation phase of the experimental trials. If the paired males

displayed the same number of aggressive behaviors, we recorded it as a tie. Our models were then built

to identify the factors that influenced male and female responses to the most and least aggressive of the

paired males. For all models, we checked model fit by utilizing the DHARMa diagnostic package in R,

which conducts simulation-based dispersion tests (Hartig, 2022). We used hypothesis-based model

building to include multiple interaction terms between our fixed effects in our models, explained in

detail further below. For the final model selection, we only retained interaction terms that improved the

fit of the model according to AIC weights.

Social Information Production Models

Our first model tested how the presence and sex of the eavesdropper influenced the aggressive

behavior of the paired males (Social Information Production Model, Table 1). The response variable for

this model was the total aggression displayed by the paired males in the observation phase of the

experiment. We calculated the total aggression by summing the five measures of aggressive behavior

that each fly performed. We had two fixed effects in this model: treatment, which described the

presence and sex of an eavesdropper, and age, as we used both 3- and 4-day old flies for all

experiments. The date the trial was conducted was included as a random effect to indicate which flies

were tested on the same day. A negative binomial error distribution with a logit link function was used

for this model, and a term was included to account for zero inflation, as some pairs had no recorded

aggressive interactions during the experimental trials. We also created a model to investigate how the

eavesdropper's genotype influenced social information production, this model and its results are

reported in the Supplemental Information (Table S1).

Male Social Information Use Models

Our next set of models tested how access to social information influenced the aggressive behavior and

social decision-making of the males that were added into the arena during the interaction phase. We

modeled the number of lunges that the added male directed towards the more aggressive paired male

(Social Information Use: Male Response to More Aggressive Male Model, Table 2) or the least aggressive

paired male (Social Information Use: Male Response to Least Aggressive Male Model, Table 3) during the

interaction phase of the experiment.

Fixed effects in the "Social Information Use: Male Response to More Aggressive Male" model included

treatment, which described if the male was an eavesdropper or was not allowed to observe social

information. We also included a fixed effect for the genotype of the added male, to account for

genotypic variation. Our model selection process led us to retain the interaction we tested between

genotype and treatment, as this interaction term improved the fit of the model (delta AIC = 1.59). An

effect of this term would indicate that genotypes differed in their use of social information. The total

aggression displayed by the paired males in the first phase of the experiment was also used as a fixed

effect, to determine if our sample of their aggressive behaviors predicted the behavior of the added

male. Our final fixed effect was the age of the flies. The date the trial was conducted and the identity of

the researcher who coded the trials were included as random effects. We tested an interaction

between the aggression of the paired males and treatment, however, that interaction did not improve

the AIC of our model (delta AIC = 1.86).

For the "Social Information Use: Male Response to Least Aggressive Male" model, we used the same

fixed effects of treatment, genotype, displayed aggression by the paired males, and age, and random

effects of date and observer. We also tested for the same interactions as the previous model, however,

our model selection process did not support the retention of either the interaction between displayed

aggression and treatment (delta AIC = 1.87) or the interaction between genotype and treatment (delta

AIC = 3.48). A negative binomial error distribution with a logit link function was used for both male social

information use models, and a term was included in both models to account for zero inflation, as some

added males had no recorded aggressive interactions during the experimental trials.

Female Social Information Use Models

Our last models tested how access to social information influenced female mating choices in the

interaction phase of the experiment. We only analyzed data from our female follow-up experiment (see

above), where females observed and were allowed to interact with two males of their preferred male

genotype, 365.

To investigate how females used social information, we first modeled whether females mated, or not

(Social Information Use: Female Model, Table 4), with a binary response variable. Females who mated

were assigned a value of 1 (n =53 females), and females who did not mate were assigned a 0 (n = 32

females). Treatment was tested as a fixed effect, to investigate how eavesdropping modified the

likelihood of female mating. The total aggression displayed by the paired males in the first phase of the

experiment was also used as a fixed effect, to determine if aggressive behavior influenced the mating

decision of the added female. We included the interaction between treatment and total paired male

aggression to investigate how observed social information influenced mating decisions, as it improved

model fit according to AIC (delta AIC = 0.51). We also tested the genotype of the added female as a fixed

effect, to account for genetic variation. We did not retain the interaction between genotype and

treatment, as it did not improve the model's AIC (delta AIC = 1.46). The date the trial was conducted and

the identity of the researcher who coded the trials were included as a random effect.

To further investigate how social information specifically altered female social decision-making, we

modeled whether each mated female chose the more aggressive paired male (Social Information Use:

Female Response to More Aggressive Male Model) or the least aggressive paired male (Social

Information Use: Female Response to Least Aggressive Male Model). We had smaller sample sizes for

these models: of the 85 females tested, only 21 mated with the more aggressive male, and only 9 mated

with the least aggressive male. These lower samples were a result of many trials resulting in a "tie"

between the paired males (n=23 trials), and females often choosing not to mate with either male (n=32

trials). We utilized a binomial error distribution for all female models. The full description of these

models and their results can be found in supplemental tables 2 and 3.

Correction for Multiple Testing

We created multiple models to investigate our hypotheses regarding social information production, use,

and social decision-making, which required a correction for multiple testing. For the Social Information

Production and Social Information Use: Male Response models, we applied Bonferroni corrections for 2

tests, since we created two models investigating the same dataset for social information production and

male social information use. For the Female Response models, we applied Bonferroni corrections for 3

tests. All p-values reported below, in tables 1-4, and in the supplementary tables reflect these

corrections.

Results

Social Information Production

The paired males altered their behavior based on the presence and sex of the eavesdropper (Figure 3,

Social Information Production Model parameter estimates: Female Eavesdropper: -0.18 ± 0.19, Male

Eavesdropper: -0.61 ± 0.19, df = 2, p-value = 0.012, Table 1). The paired males were the least aggressive

when a male eavesdropper was present (Social Information Production Model: least-squares means for

paired male aggression in the male eavesdropper treatment = 1.84). When a female eavesdropper was

present, the paired males did not significantly alter their behavior, compared to when there was no

eavesdropper present (Social Information Production Model: least-squares means for paired male

aggression in the female eavesdropper treatment = 2.27, no eavesdropper treatment = 2.45). Age did

not significantly predict paired male aggressive behavior (Social Information Production Model

parameter estimate: 4 day old: 0.41 ± 0.19, df = 1, p-value = 0.07, Table 1). We also did not see an effect of eavesdropper genotype on paired male aggression (Supplemental Table 1).

Figure 3. Variation in social information production based on audience identity. The total aggression of the paired males in relation to the sex and presence of an eavesdropper observing them. We calculated total aggression by summarizing five measures of aggressive behavior we recorded for each of the paired males. Analysis via linear mixed models revealed a significant effect of sex and presence of the eavesdropper on paired male aggressive behavior (see text).

Male Social Information Use

Male flies altered their behavior based
on their access to social information.
Male eavesdroppers decreased their
own aggression towards the more
aggressive paired male, compared to
males without social information (Social
Information Use Male Response to More
Aggressive Male Model parameter
        estimate: Male Eavesdropper: -1.51 ±

0.43, df = 1, p-value = 0.0008, Table 2). We observed genetic variation in how males responded towards

the least aggressive paired male (Social Information Use Male Response to Least Aggressive Male

Model: X2 = 7.5, df = 2, p-value = 0.047, Table 3), but not the most aggressive paired male (Social

Information Use Male Response to More Aggressive Male Model:X2 = 6.74, df = 2, p-value = 0.07, Table

2). We saw a significant genotype by treatment interaction on male aggressive behavior towards the

more aggressive paired male. This interaction indicates that genotypes differed in the extent to which

they modified their behavior towards the more aggressive paired male based on their ability to collect

social information during the first phase of the experiment (Social Information Use Male Response to

More Aggressive Male Model: X2 = 16.5, df = 2, p-value = 0.0005, Table 2, Figure 4). The total aggression

of the paired males in the observation phase of the experiment did not significantly predict how

aggressive the male flies would behave towards the more or less aggressive of the paired males (Social

Information Use Male Response to More Aggressive Male Model parameter estimate: -0.02 ± 0.01, df =

1, p-value = 0.27; Social Information Use Male Response to Least Aggressive Male Model parameter

estimate: 0.02 ± 0.02, df = 1, p-value = 0.47, Table 2 & 3). Age also did not significantly predict male

aggressive behavior in any model (Social Information Use Male Response to More Aggressive Male

Model parameter estimate: 4 day old: -0.18 ± 0.2, df = 1, p-value = 0.74, Social Information Use Male

Response to Least Aggressive Male Model parameter estimate: 4 day old: -0.52 ± 0.32, df = 1, p = 0.2,

Table 2 & 3).

Female Social Information Use

Female flies displa yed strong genotypic differences in likelihood to mate (Social Information Use Female

Model: X2 = 14.01, df = 2, p-value = 0.003, Table 4). Specifically, females from genotype 229 almost

always mated (females mated in 27/29 trials), and females from genotype 136 rarely mated (females

mated in 8/27 trials). Neither access to social information nor the displayed aggression of the interacting

pair significantly predicted the likelihood of mating (Social Information Use Female Model parameter

estimates: treatment Female Eavesdropper: 1.63 ± 0.68, df = 1, p-value = 0.2; displayed aggression 0.03

± 0.1, df = 1, p-value = 0.14, Table 4). The interaction between total aggression and treatment was also

not significant in predicting female mating likelihood (Social Information Use Female Model: X2 = 2.4, df

= 1, p-value = 0.38).

None of our fixed or random effects significantly predicted whether the female mated with the more

aggressive paired male, or the least aggressive paired male; full results are presented in supplementary

| 414 | tables 2 and 3. |

415 **Discussion**

Traditionally, social

information use and

production have been

studied separately,

which limits our

understanding of how

these two processes

interact as they occur

simultaneously in

nature (Bonnie &

Earley, 2007;

McGregor, 2005).

Considering the

linkages between social

information production

and use can improve

our predictions about

how behaviors spread

in populations and

identify the causes of

individual differences in

Figure 4. Social decision making in male social information use. The least squared means model outputs for the number of times that the male lunged at the least aggressive or most aggressive of the paired males in the second half of experimental trials is represented on the y-axis in both graphs, and the x-axis in both graphs shows the three tested genotypes. Males with social information lunged significantly less often at observed contest winners than males without social information, but this effect was modulated by genotype.

behavioral strategies (Aplin & Morand-Ferron, 2017; Chimento et al., 2022). However, despite these

important implications of the feedback loop on the evolution of social behaviors, we know of no

empirical studies of the feedback loop that have been conducted in the context of social eavesdropping.

In this study, we experimentally demonstrate the existence of a feedback loop between social

information production and use in an aggressive context in male fruit flies. We found that the presence

of a male eavesdropper substantially decreased the aggression of two paired males, supporting an

audience effect (Figure 3, Table 1). Furthermore, observing social information caused eavesdropping

males to change their aggressive behaviors: male eavesdroppers lunged at the more aggressive paired

male less often, compared to males who did not have the opportunity to eavesdrop (Figure 4, Table 2).

We also identified genetic variation in male social information use – different genotypes had different

responses after observing an aggressive encounter. Overall, these findings indicate that there is a

bidirectional transfer of information between the participants and the eavesdroppers in an aggressive

context (Figure 5).

Social eavesdropping can benefit individuals by allowing them to assess their opponents before

engaging in aggressive behavior, thus reducing the costs of fighting (Earley, 2010; Johnstone, 2001). We

found a reduction in aggressive behavior from both the paired males when observed by male

eavesdroppers, and male eavesdroppers when allowed to interact with the paired males. Thus, the

feedback between social information use and production produced substantially lower overall

aggression than would be predicted by simply studying pairs of males without eavesdroppers (e.g., in

our no eavesdropper treatment), or investigating male eavesdropping without also considering its effect

on social information production (e.g., in our no social information treatment). Our results are therefore

consistent with previous work that shows obtaining social information about the fighting ability of an

Figure 5. Diagrams depicting how our study's findings support the existence of a sex-specific feedback loop between social information use and production. Males simultaneously influence the production of and use social information resulting from an aggressive contest. Females do not influence social information production and do not use social information resulting from an aggressive contest.

opponent may allow individuals to avoid prolonged or escalated encounters (Beltrão et al., 2023;

Magnhagen, 2006; Tibbetts et al., 2020); notably, these previous results were obtained without

accounting for any influence of the eavesdropper presence on the social information that was produced.

Our qualitatively similar findings suggest that, while studying this feedback better informs our

understanding of each individual's behavior, they do not qualitatively alter the patterns (i.e., that

individuals use social eavesdropping to avoid potentially costly aggressive encounters) that others have

previously observed, at least in this case.

While we are confident that observation of an aggressive encounter altered male eavesdropper

behavior, we cannot be certain about the exact social information that the male eavesdroppers were

using to inform their decision-making. Winner-loser effects have been documented in flies, providing

evidence that flies can learn from aggressive contests – at least, those in which they have directly

participated (Penn et al., 2010; Yurkovic et al., 2006). Despite this, our measure of the aggressive

behavior of the paired males did not directly influence either male eavesdroppers or males without

social information (tables 2 and 3). Eavesdroppers could have extracted information from a variety of

sources in the arena, such as observations of how the paired males interacted, chemical signaling

behaviors, and/or sounds produced by the paired males (Kent et al., 2008; Versteven et al., 2017).

Further research will be necessary to identify mechanistic relationships between eavesdropping, paired

male behavior, and the eavesdropper's later aggression.

Another mechanism that could cause male eavesdroppers to change their behavior in response to the

paired males is priming. Priming occurs when observing an aggressive encounter alters an individual's

later behaviors, however, this change in behaviors is expected to be broad and generalize to all future

opponents (Clotfelter & Paolino, 2003; Cruz & Oliveira, 2015; Earley et al., 2005; Hsu et al., 2006;

Tibbetts et al., 2020). Social eavesdropping, on the other hand, is expected to be individual-specific – the

eavesdropper remembers information and learns from the behaviors of those they observed (McGregor,

2005). While we cannot rule out any effects of priming, our results are more consistent with social

eavesdropping, as we saw an effect of social information (i.e., treatment) on male behavior that was

dependent upon the specific identity of the paired males, i.e., if they were more or less aggressive

during the aggressive contest (Table 2). Interestingly, we found that males who did not have the

opportunity to eavesdrop on the fight disproportionately attacked the more aggressive of the paired

males (Figure 4). With no prior social information to rely on, these males aggressive behaviors were

most likely influenced by their personal information and direct experiences once in the arena. For

example, fights between the naïve male and the more aggressive paired male may have been instigated

by the paired male, leading to increased naïve male aggression as a retaliatory response. In contrast,

male eavesdroppers seemingly utilized social information to avoid and engage in fewer aggressive

encounters with the more aggressive male. Thus, we find support for a transfer of social information

from the paired males to the male eavesdroppers that resulted in a change in behavior that was specific

to the individuals that were observed. The detailed cognitive mechanisms underlying this information

transfer representing an interesting area for future study. The feedback loop between social information

use and production proposes that the presence of a social eavesdropper should cause an adjustment in

social information production, which then would lead to changes in eavesdropper behavior. In our

study, the paired males displayed similar aggression levels whether a female was observing them or if

there was no eavesdropper present (Figure 3). Further, we did not observe any differences in the mating

decisions between female eavesdroppers and females without social information (Table 4). Together,

these findings further support the idea that changes in eavesdropper behavior are dependent upon

modifications in the social information that is produced in their presence.

Genetic variation significantly explained both male and female behaviors in our experiment. Female

genotype had a strong effect on mating probability, and paired male genotypes also apparently differed

substantially in attractiveness, so much so that we had to repeat the experiment to control for this

effect. While eavesdropper genotype did not influence social information production, we interestingly

found that male eavesdroppers of different genotypes varied in how social information influenced their

response to the more aggressive male. Thus in our feedback loop, we find that all male eavesdroppers

had access to similar information, yet male genotypes differed in how they modified their behavioral

response to social information. We tested only three genotypes; future experiments that account for

genetic diversity are critical for a better assessment of how genetic variation augments or abrogates

social information feedbacks (Araya-Ajoy et al., 2020; Cantor et al., 2021).

Our study has a few key limitations to note. All of our flies had no prior social experience before the

experimental trials began. While this provided a standardized methodology to test all of our flies,

different results may be obtained for individuals that have had prior social interactions with aggression

and mating behaviors (Taborsky & Oliveira, 2012). Our female data also presented several challenges:

any effect of social information was obscured in our initial set of experiments due to a strong preference

for one of the paired male genotypes. In our follow-up trials, our paired 365 males were less aggressive

on average than the 765/365 paired males. As a result, we often were unable to assign a more and least

aggressive male during the experimental trials. Surprisingly, our females also were not always willing to

mate, although they were unmated for their entire adult lives. While we were still able to obtain enough

data in our follow-up experiment to investigate the feedback loop, continued work is necessary to

better understand how females use social information from male-male aggressive contests, if at all.

This study has revealed several areas of further research necessary to continue to better elucidate the

relationship between social information use and production. The sex differences that we observed

emphasize the need for continued testing and modeling of these processes in diverse populations, as

accounting for individual differences is vital for accurate predictions of social information spread in

populations (Duboscq et al., 2016). Additionally, we need a better understanding of the selection

pressures in social networks that are introduced via this feedback loop. In this system for example, in

order for increased aggression to be adaptive, the benefits must outweigh the costs, but these will likely

change depending on the social environment (Duckworth, 2006). Further research can tackle this by

quantifying the costs and benefits of plasticity in social information production and use.

Overall, our study lends the first empirical support for a feedback loop between how social information

is produced and used in different social contexts. Utilizing the social information feedback framework to

investigate animal behaviors can contribute to a better understanding of how behaviors related to

conflict, sociality, and group living are expressed and evolve.

Declarations

Acknowledgments

We thank Lea Pollack and other members of the Saltz Lab for helpful discussions that improved the

manuscript, and we thank Anja Hartge, Keshav Wagle, Erin Vance, Marcelo Dimitri, and Autumn

Hildebrand for assistance with fly maintenance and data collection. Funding support for this research

was provided by the National Science Foundation (NSF IOS-1856577 and NSF DEB- 2217557 to JBS, NSF

GRFP-1842494 to MH), and by the Houston Livestock Show & Rodeo First Year Graduate Fellowship (to

MH).

Competing Interests: None

Author Contributions

MH and JBS - Conceptualization; MH - Data collection; MH and JBS - Formal analysis; MH and JBS –

Writing original draft, review & editing

Data Availability Statement

The datasets and code are available on figshare -

https://figshare.com/projects/Eavesdropping_and_the_audience_effect_in_fighting_fruit_flies_reveal_f

eedback_between_social_information_use_and_production/186912

569   Table 1. Parameters, test statistics, and their interpretation in the model for social information

570   production. P-values are corrected for multiple testing.

Parameter	Parameter Estimate [Standard Error]	p-value	Chi-squared value	Interpretation
Treatment	Female Eavesdropper: -0.18 [0.19] Male Eavesdropper: -0.61 [0.19]	0.012	10.1	Paired males altered their aggressive behavior based on the presence and sex of an eavesdropper
Age	4 day old flies: 0.41 [0.19]	0.07	4.48	No effect of age on paired male aggression
Date	-	0.25	2.32	No effect of experimental day on paired male aggression

577   Table 2. Parameters, test statistics, and their interpretation in the model for male social information use

578   towards the more aggressive paired male. P-values are corrected for multiple testing.

Parameter	Parameter Estimate [Standard Error]	p-value	Chi-squared value	Interpretation
Treatment	Male Eavesdropper: -: 1.51[0.43]	0.0008	12.6	Access to social information altered how aggressive males were towards previous contest winners
Genotype	-	0.07	6.7	No detected genotypic variation in how males directed their aggression toward the winner
Treatment x Genotype	-	0.0005	16.5	Genotypic variation in how males used social information to direct their aggressive behaviors toward winners
Age	4 day old: -0.18 [0.2]	0.74	0.8	No effect of age on male aggression
Paired Male Aggression	-0.017[0.01]	0.27	2.3	Measured aggression in the previous contest did not affect how males directed their behaviors toward winners
Observer	-	0.051	4.98	
Date	-	2	0	No effect of experimental day on aggression

598   Table 3. Parameters, test statistics, and their interpretation in the model for male social information use

599   towards the less aggressive paired male. P-values are corrected for multiple testing.

Parameter	Parameter Estimate [Standard Error]	p-value	Chi-squared value	Interpretation
Treatment	No Social Information – 0.05[0.34]	1.9	0.009	Access to social information did not alter how aggressive males were towards previous contest losers
Genotype	-	0.05	7.5	Genotypic variation in how males directed their aggression toward the loser
Age	4 day old: -0.52 [0.32]	0.2	2.4	No effect of age on male aggression
Paired Male Aggression	0.02[0.02]	0.47	1.4	Measured aggression in the previous contest did not predict how males directed their behaviors toward losers
Observer	-	1.6	0.79	
Date	-	2	0	No effect of experimental day on aggression

602   Table 4. Parameters, test statistics, and their interpretation in the model for female likelihood to mate.

603   P-values are corrected for multiple testing.

Parameter	Parameter Estimate [Standard Error]	p-value	Chi-squared value	Interpretation
Genotype	-	0.0027	14	Genotypic variation in likelihood to mate
Paired Male Aggression	0.03 [0.1]	0.14	4	Measured aggression displayed in the previous contest did not alter mating behaviors
Treatment	Female Eavesdropper: 1.63 [0.68]	0.2	3.4	Access to social information did not significantly predict likelihood to mate
Treatment x Paired Male Aggression	-	0.38	2.4	Observing the aggression of the paired males did not significantly change their likelihood to mate
Observer		2	0	
Date	-	0.81	1.2	No effect of experimental day on aggression

Works Cited

Aplin, L. M., & Morand-Ferron, J. (2017). Stable producer–scrounger dynamics in wild birds: sociability

and learning speed covary with scrounging behaviour. *Proceedings of the Royal Society B:*

Biological Sciences, *284*(1852). https://doi.org/10.1098/RSPB.2016.2872

Aquiloni, L., Buřič, M., & Gherardi, F. (2008). Crayfish females eavesdrop on fighting males before

choosing the dominant mate. *Current Biology*, *18*(11), R462–R463.

https://doi.org/10.1016/J.CUB.2008.04.006

Araya-Ajoy, Y. G., Westneat, D. F., & Wright, J. (2020). Pathways to social evolution and their

evolutionary feedbacks. *Evolution*, *74*(9), 1894–1907. https://doi.org/10.1111/EVO.14054

Beltrão, P., Gomes, A. C. R., & Cardoso, G. C. (2023). Bullying as an advertisement of social dominance in

common waxbills. *Proceedings. Biological Sciences*, *290*(2000).

https://doi.org/10.1098/RSPB.2023.0206

Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R., Stevens, M. H. H., & White, J. S. S.

(2009). Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in*

Ecology & Evolution, *24*(3), 127–135. https://doi.org/10.1016/J.TREE.2008.10.008

Bond, C. F., & Titus, L. J. (1983). Social facilitation: A meta-analysis of 241 studies. *Psychological Bulletin*,

94(2), 265–292. https://doi.org/10.1037/0033-2909.94.2.265

Bonnie, K. E., & Earley, R. L. (2007). Expanding the scope for social information use. *Animal Behaviour*,

74(2), 171–181. https://doi.org/10.1016/J.ANBEHAV.2006.12.009

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J.,

Mächler, M., & Bolker, B. M. (2017). glmmTMB balances speed and flexibility among packages for

zero-inflated generalized linear mixed modeling. *R Journal*, *9*(2), 378–400.

https://doi.org/10.32614/RJ-2017-066

Cantor, M., Maldonado-Chaparro, A. A., Beck, K. B., Brandl, H. B., Carter, G. G., He, P., Hillemann, F.,

Klarevas-Irby, J. A., Ogino, M., Papageorgiou, D., Prox, L., & Farine, D. R. (2021). The importance of

individual-to-society feedbacks in animal ecology and evolution. *Journal of Animal Ecology*, *90*(1),

27–44. https://doi.org/10.1111/1365-2656.13336

Chan, J. P. Y., Lau, P. R., Tham, A. J., & Li, D. (2008). The effects of male-male contests and female

eavesdropping on female mate choice and male mating success in the jumping spider, Thiania

bhamoensis (Araneae: Salticidae). *Behavioral Ecology and Sociobiology*, *62*(5), 639–646.

https://doi.org/10.1007/S00265-007-0488-X/FIGURES/6

Chapman, C. A., Chapman, L. J., & Lefebvre, L. (1990). Spider monkey alarm calls: honest advertisement

or warning kin? *Animal Behaviour*, *39*(1), 197–198. https://doi.org/10.1016/S0003-3472(05)80742-

0

Chen, S., Lee, A. Y., Bowens, N. M., Huber, R., & Kravitz, E. A. (2002). Fighting fruit flies: A model system

for the study of aggression. *PNAS*, *99*(8), 5664–5668. www.pnas.orgcgidoi10.1073pnas.082102599

Chimento, M., Barrett, B. J., Kandler, A., & Aplin, L. M. (2022). Cultural diffusion dynamics depend on

behavioural production rules. *Proceedings of the Royal Society B: Biological Sciences*, *289*(1980).

https://doi.org/10.1098/RSPB.2022.1001

Clotfelter, E. D., & Paolino, A. D. (2003). Bystanders to contests between conspecifics are primed for

increased aggression in male fighting fish. *ANIMAL BEHAVIOUR*, *66*, 343–347.

https://doi.org/10.1006/anbe.2003.2227

Coppinger, B., Cannistraci, R. A., Karaman, F., Kyle, S. C., Hobson, E. A., Freeberg, T. M., & Hay, J. F.

(2017). Studying audience effects in animals: what we can learn from human language research.

Animal Behaviour, *124*, 161–165. https://doi.org/10.1016/J.ANBEHAV.2016.12.020

Cruz, A. S., & Oliveira, R. F. (2015). Audience effects and aggressive priming in agonistic behaviour of

male zebrafish, Danio rerio. *Animal Behaviour*, *107*, 269–276.

https://doi.org/10.1016/J.ANBEHAV.2015.07.007

Dall, S. R. X., Giraldeau, L. A., Olsson, O., McNamara, J. M., & Stephens, D. W. (2005). Information and its

use by animals in evolutionary ecology. *Trends in Ecology & Evolution*, *20*(4), 187–193.

https://doi.org/10.1016/J.TREE.2005.01.010

Danchin, É., Giraldeau, L.-A., Valone, T. J., & Wagner, R. H. (2004). Public Information: From Nosy

Neighbors to Cultural Evolution. *Science*, *305*(5683), 487–491.

https://doi.org/10.1126/SCIENCE.1098254

Danchin, E., Nöbel, S., Pocheville, A., Dagaeff, A. C., Demay, L., Alphand, M., Ranty-Roby, S., Van

Renssen, L., Monier, M., Gazagne, E., Allain, M., & Isabel, G. (2018). Cultural flies: Conformist social

learning in fruitflies predicts long-lasting mate-choice traditions. *Science*, *362*(6418), 1025–1030.

https://doi.org/10.1126/SCIENCE.AAT1590

Desjardins, J. K., Hofmann, H. A., & Fernald, R. D. (2012). Social Context Influences Aggressive and

Courtship Behavior in a Cichlid Fish. *PLOS ONE*, *7*(7), e32781.

https://doi.org/10.1371/JOURNAL.PONE.0032781

Doutrelant, C., McGregor, P. K., & Oliveira, R. F. (2001). The effect of an audience on intrasexual

communication in male Siamese fighting fish, Betta splendens. *Behavioral Ecology*, *12*(3), 283–

286. https://doi.org/10.1093/BEHECO/12.3.283

Duboscq, J., Romano, V., MacIntosh, A., & Sueur, C. (2016). Social information transmission in animals:

Lessons from studies of diffusion. *Frontiers in Psychology*, *7*(AUG), 1147.

https://doi.org/10.3389/FPSYG.2016.01147/BIBTEX

Duckworth, R. A. (2006). Behavioral correlations across breeding contexts provide a mechanism for a

cost of aggression. *Behavioral Ecology*, *17*(6), 1011–1019.

https://doi.org/10.1093/BEHECO/ARL035

Duque-Wilckens, N., Trainor, B. C., & Marler, C. A. (2019). *Aggression and Territoriality* (2nd ed.).

Academic Press. https://doi.org/10.1016/B978-0-12-809633-8.90064-5

Earley, R. L. (2010). Social eavesdropping and the evolution of conditional cooperation and cheating

strategies. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1553), 2675–

2686. https://doi.org/10.1098/RSTB.2010.0147

Earley, R. L., Druen, M., & Dugatkin, L. A. (2005). Watching fights does not alter a bystander's response

towards naïve conspecifics in male green swordtail fish, Xiphophorus helleri. *Animal Behaviour*,

69(5), 1139–1145. https://doi.org/10.1016/J.ANBEHAV.2004.07.015

Earley, R. L., & Dugatkin, L. A. (2002). Eavesdropping on visual cues in green swordtail (Xiphophorus

helleri) fights: a case for networking. *Proc R Soc Lond*, 943–952. https://royalsocietypublishing.org/

Fichtel, C., & Manser, M. (2010). Vocal communication in social groups. *Animal Behaviour: Evolution and*

Mechanisms, 29–54. https://doi.org/10.1007/978-3-642-02624-9_2/COVER

Fitzsimmons, L. P., & Bertram, S. M. (2013). Playing to an audience: the social environment influences

aggression and victory displays. *Biology Letters*, *9*(4). https://doi.org/10.1098/RSBL.2013.0449

Friard, O., & Gamba, M. (2016). BORIS: a free, versatile open-source event-logging software for

video/audio coding and live observations. *Methods in Ecology and Evolution*, *7*(11), 1325–1330.

https://doi.org/10.1111/2041-210X.12584

Hartig, F. (2022). *DHARMa: Residual Diagnostics for Hierarchical (Multi-level/Mixed) Regression Models*.

http://florianhartig.github.io/DHARMa/

Heyes, C. M. (1994). Social learning in animals: categories and mechanisms. *Biological Reviews of the*

Cambridge Philosophical Society, *69*(2), 207–231. https://doi.org/10.1111/J.1469-

185X.1994.TB01506.X

Hobson, E. A. (2020). Differences in social information are critical to understanding aggressive behavior

in animal dominance hierarchies. *Current Opinion in Psychology*, *33*, 209–215.

https://doi.org/10.1016/J.COPSYC.2019.09.010

Hoffmann, A. A., & Cacoyianni, Z. (1990). Territoriality in Drosophila melanogaster as a conditional

strategy. *Anim. Behav*, *40*, 526–537.

Hsu, Y., Earley, R. L., & Wolf, L. L. (2006). Modulation of aggressive behaviour by fighting experience:

mechanisms and contest outcomes. *Biological Reviews of the Cambridge Philosophical Society*,

81(1), 33–74. https://doi.org/10.1017/S146479310500686X

Johnsson, J. I., & Åkerman, A. (1998). Watch and learn: preview of the fighting ability of opponents alters

contest behaviour in rainbow trout. *Animal Behaviour*, *56*(3), 771–776.

https://doi.org/10.1006/ANBE.1998.0824

Johnstone, R. A. (2001). Eavesdropping and animal conflict. *Proceedings of the National Academy of*

Sciences, *98*(16), 9177–9180. https://doi.org/10.1073/PNAS.161058798

Kent, C., Azanchi, R., Smith, B., Formosa, A., & Levine, J. D. (2008). Social Context Influences Chemical

Communication in D. melanogaster Males. *Current Biology*, *18*(18), 1384–1389.

https://doi.org/10.1016/J.CUB.2008.07.088

King, J. A. (1973). The Ecology of Aggressive Behavior. *Annual Review of Ecology and Systematics*, *4*(1),

117–138. https://doi.org/10.1146/ANNUREV.ES.04.110173.001001

MacKay, T. F. C., Richards, S., Stone, E. A., Barbadilla, A., Ayroles, J. F., Zhu, D., Casillas, S., Han, Y.,

Magwire, M. M., Cridland, J. M., Richardson, M. F., Anholt, R. R. H., Barrón, M., Bess, C.,

Blankenburg, K. P., Carbone, M. A., Castellano, D., Chaboub, L., Duncan, L., … Gibbs, R. A. (2012).

The Drosophila melanogaster Genetic Reference Panel. *Nature 2012 482:7384*, *482*(7384), 173–

178. https://doi.org/10.1038/nature10811

Magnhagen, C. (2006). Information about an opponent can decrease aggression in male common gobies

competing for nest sites. *Animal Behaviour*, *71*(4), 879–884.

https://doi.org/10.1016/J.ANBEHAV.2005.07.017

Matos, R. J., & McGregor, P. K. (2002). THE EFFECT OF THE SEX OF AN AUDIENCE ON MALE-MALE

DISPLAYS OF SIAMESE FIGHTING FISH (BETTA SPLENDENS). *Behaviour*, *139*(9), 1211–1221.

https://doi.org/10.1163/15685390260437344

McGregor, P. K. (2005a). Animal Communication Networks. *Animal Communication Networks*, 1–657.

https://doi.org/10.1017/CBO9780511610363

McGregor, P. K. (2005b). *Animal communication networks*. 657.

https://books.google.com/books/about/Animal_Communication_Networks.html?id=B58DSFCaQp

wC

McGregor, P. K., & Peake, T. M. (2000). Communication networks: social environments for receiving and

signalling behaviour. *Acta Ethologica 2000 2:2*, *2*(2), 71–81.

https://doi.org/10.1007/S102110000015

Mery, F., Varela, S. A. M., Danchin, É., Blanchet, S., Parejo, D., Coolen, I., & Wagner, R. H. (2009a). Public

Versus Personal Information for Mate Copying in an Invertebrate. *Current Biology*, *19*(9), 730–734.

https://doi.org/10.1016/j.cub.2009.02.064

Mery, F., Varela, S. A. M., Danchin, É., Blanchet, S., Parejo, D., Coolen, I., & Wagner, R. H. (2009b). Public

Versus Personal Information for Mate Copying in an Invertebrate. *Current Biology*, *19*(9), 730–734.

https://doi.org/10.1016/J.CUB.2009.02.064

Monier, M., Nöbel, S., Isabel, G., & Danchin, E. (2018). Effects of a sex ratio gradient on female mate-

copying and choosiness in Drosophila melanogaster. *Current Zoology*, *64*(2), 251–258.

https://doi.org/10.1093/CZ/ZOY014

Nöbel, S., Danchin, E., & Isabel, G. (2018). Mate-copying for a costly variant in Drosophila melanogaster

females. *Behavioral Ecology*, *29*(5), 1150–1156. https://doi.org/10.1093/BEHECO/ARY095

Oliveira, R. F., Mcgregor, P. K., & Latruffe{, C. (1998). Know thine enemy: fighting fish gather information

from observing conspecific interactions. *Proc Biol Sci*, *265*(1401), 1045–1049.

Peake, T. M., Matos, R. J., & Mcgregor, P. K. (2006). Effects of manipulated aggressive 'interactions' on

bystanding male fighting fish, Betta splendens. *Animal Behaviour*, *72*(5), 1013–1020.

https://doi.org/10.1016/J.ANBEHAV.2006.01.029

Peake, T. M., & Mcgregor, P. K. (2004). Information and aggression in fishes. *Animal Learning & Behavior*

2004 32:1, *32*(1), 114–121. https://doi.org/10.3758/BF03196012

Peake, T. M., Terry, A. M. R., McGregor, P. K., & Dabelsteen, T. (2001). Male great tits eavesdrop on

simulated maletomale vocal interactions. *Proceedings of the Royal Society of London. Series B:*

Biological Sciences, *268*(1472), 1183–1187. https://doi.org/10.1098/RSPB.2001.1648

Penn, J. K. M., Zito, M. F., & Kravitz, E. A. (2010). A single social defeat reduces aggression in a highly

aggressive strain of Drosophila. *Proceedings of the National Academy of Sciences of the United*

States of America, *107*(28), 12682–12686. https://doi.org/10.1073/PNAS.1007016107

Plath, M., Blum, D., Schlupp, I., & Tiedemann, R. (2008). Audience effect alters mating preferences in a

livebearing fish, the Atlantic molly, Poecilia mexicana. *Animal Behaviour*, *75*(1), 21–29.

https://doi.org/10.1016/J.ANBEHAV.2007.05.013

Pollick, A. S., Gouzoules, H., & De Waal, F. B. M. (2005). Audience effects on food calls in captive brown

capuchin monkeys, Cebus apella. *Animal Behaviour*, *70*(6), 1273–1281.

https://doi.org/10.1016/J.ANBEHAV.2005.03.007

Saltz, J. B. (2013). Genetic composition of social groups influences male aggressive behaviour and fitness

in natural genotypes of Drosophila melanogaster. *Proceedings of the Royal Society B: Biological*

Sciences, *280*(1771). https://doi.org/10.1098/RSPB.2013.1926

Stamps, J., Buechner, M., Alexander, K., Davis, J., & Zuniga, N. (2005). Genotypic differences in space use

and movement patterns in Drosophila melanogaster. *Animal Behaviour*, *70*(3), 609–618.

https://doi.org/10.1016/J.ANBEHAV.2004.11.018

Taborsky, B., & Oliveira, R. F. (2012). Social competence: an evolutionary approach. *Trends in Ecology &*

Evolution, *27*(12), 679–688. https://doi.org/10.1016/J.TREE.2012.09.003

Tibbetts, E. A., Wong, E., & Bonello, S. (2020). Wasps Use Social Eavesdropping to Learn about Individual

Rivals. *Current Biology*, *30*(15), 3007-3010.e2. https://doi.org/10.1016/J.CUB.2020.05.053

Townsend, S., & Zuberbuhler, K. (2009). Audience effects in chimpanzee copulation calls.

Http://Www.Tandfonline.Com/Action/AuthorSubmission?JournalCode=kcib20&page=instructions,

2(3), 282–284. https://doi.org/10.4161/CIB.2.3.6796

Trannoy, S., Penn, J., Lucey, K., Popovic, D., & Kravitz, E. A. (2016). Short and long-lasting behavioral

consequences of agonistic encounters between male Drosophila melanogaster. *Proceedings of the*

National Academy of Sciences, *113*(17), 4818–4823. https://doi.org/10.1073/PNAS.1520953113

Turner, C. R., Spike, M., & Magrath, R. D. (2023). The evolution of eavesdropping on heterospecific alarm

calls: Relevance, reliability, and personal information. *Ecology and Evolution*, *13*(7), e10272.

https://doi.org/10.1002/ECE3.10272

Vakirtzis, A. (2011). Mate Choice Copying and Nonindependent Mate Choice: A Critical Review.

Https://Doi.Org/10.5735/086.048.0202, *48*(2), 91–107. https://doi.org/10.5735/086.048.0202

Van Breukelen, N. A., & Draud, M. (2005). The roles of male size and female eavesdropping in divorce in

the monogamous convict cichlid (Archocentrus nigrofasciatus, Cichlidae). *Behaviour*, *142*(8), 1023–

1035. https://doi.org/10.1163/156853905774405308

Versteven, M., Vanden Broeck, L., Geurten, B., Zwarts, L., Decraecker, L., Beelen, M., Göpfert, M. C.,

Heinrich, R., & Callaerts, P. (2017). Hearing regulates Drosophila aggression. *Proceedings of the*

National Academy of Sciences of the United States of America, *114*(8), 1958–1963.

https://doi.org/10.1073/PNAS.1605946114/SUPPL_FILE/PNAS.1605946114.SM05.MP4

Wagner, R. H., & Danchin, É. (2010). A taxonomy of biological information. *Oikos*, *119*(2), 203–209.

https://doi.org/10.1111/J.1600-0706.2009.17315.X

Wang, L., & Anderson, D. J. (2010). Identification of an aggression-promoting pheromone and its

receptor neurons in Drosophila. *Nature*, *463*(7278), 227–231.

https://doi.org/10.1038/NATURE08678

Wertheim, B., Allemand, R., Vet, L. E. M., & Dicke, M. (2006). Effects of aggregation pheromone on

individual behaviour and food web interactions: a field study on Drosophila. *Ecological*

Entomology, *31*(3), 216–226. https://doi.org/10.1111/J.1365-2311.2006.00757.X

Wheeler, B. C. (2008). Selfish or altruistic? An analysis of alarm call function in wild capuchin monkeys,

Cebus apella nigritus. *Animal Behaviour*, *76*(5), 1465–1475.

https://doi.org/10.1016/J.ANBEHAV.2008.06.023

White, D. J., King, A. P., Cole, A., & West, M. J. (2002). Opening the social gateway: Early vocal and social

sensitivities in brown-headed cowbirds (Molothrus ater). *Ethology*, *108*(1), 23–37.

https://doi.org/10.1046/J.1439-0310.2002.00755.X

Wiley, R. H. (1983). Evolution of Communication. *Animal Behaviour*, *2*, 156–189.

Yurkovic, A., Wang, O., Basu, A. C., & Kravitz, E. A. (2006). Learning and memory associated with

aggression in Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the*

United States of America, *103*(46), 17519–17524.

https://doi.org/10.1073/PNAS.0608211103/SUPPL_FILE/INDEX.HTML

Zuberbühler, K. (2008). Audience effects. *Current Biology*, *18*(5), R189–R190.

https://doi.org/10.1016/j.cub.2007.12.041