Title: Environmental factors and cultural drift influence song evolution in New World Sparrows

Authors: Kaiya L. Provost[1,2]*, Jiaying Yang[2,3], Bryan C. Carstens[2]

Affiliations:

[1]Department of Biology, Adelphi University; One South Avenue, Garden City NY, 11530, USA

[2]Museum of Biological Diversity and Department of Evolution, Ecology and Organismal Biology, The Ohio State University; 1315 Kinnear Rd., Columbus OH, 43212, USA.

[3]Biological Science Department, Vanderbilt University; 2201 West End Ave, Nashville, TN 37235, USA.

*Corresponding author. Email: kprovost@adelphi.edu

Abstract: Variation in bird song is often assumed to be determined by sexual selection, rather than natural selection. However, most investigations to date have drawn their conclusions from a handful of species due to the challenges with manually processing sound data. Here, we use deep machine learning to investigate nearly all species of New World Sparrows. We leverage existing data to identify the processes that structure variation in bird song and to determine how this variation corresponds to patterns in genes and traits. Song variation in ~40% of species can be explained by environment, geography, and time. Across a community and global scale, the action of natural selection on the evolution of song is at least as impactful as it is on other genetically-determined traits.

Main Text: Singing behavior in birds has a high impact on reproduction, fitness, and speciation, especially in Passeriformes (*1–3*), but it remains unclear how song evolves within species. Hypotheses generally explain the evolution of bird song via the action of selective or neutral forces. For example, natural, sexual, and social selection have been invoked by many researchers (*4–6*), including adaptation to the environment (*7–10*). Stochastic evolution, either via cultural drift or song learning, has also been reported in multiple investigations (*11–15*) with genetic drift implicated in others (*16–22*). Similar to the selectionist / neutralist debates in molecular evolution (e.g., *23*) this question has been difficult to disentangle in large part due to the small datasets that have been applied to test these hypotheses. Most researchers rely on recordings of bird songs collected from the field, converted to audiograms, and manually processed to measure differences in the timing, placement, and frequency of the syllables that constitute the songs (e.g., Fig. 1). However, audiogram data is laborious to collect, with each syllable taking on average four seconds to annotate based on preliminary work.

Fig. 1: Example spectrogram from Chipping Sparrow (*Spizella passerina*). Recording depicts the first syllable of XC 816010 with 25 individual notes. X-axis depicts time in seconds, with black bar showing one second of time. The Y-axis depicts frequency (kHz). Darker colors indicate sounds are louder – note that gray color below the notes represents background noise, while gray color above notes represents harmonics.

We test what factors predict intraspecific diversity in songs by quantifying intraspecific variation and its correlates in the New World Sparrows (Family: Passerellidae). This family was chosen because they reside in a variety of habitats and are notable for their variation in song across species. They are also well-studied in our existing curated song datasets (e.g., Xeno-Canto, Borror Lab of Bioacoustics, Macaulay Library). For example, on Xeno-Canto at the time of writing, the 16,833 Passerellidae recordings make up 2% of the total ~842,000 bird recordings, despite comprising only 1.2% (134) of the 10,495 bird species. The Borror Lab of Bioacoustics database has an even higher bias: 31.8% (14,360/45,097) of the bird songs are Passerellids. Data on intraspecific variation in bird songs can be challenging to extract from the mountain of preexisting recording data . We labeled ~500,000 individual song syllables across ~26,000 recordings and 125 species (*24, 25*). We calculated song properties for each syllable to estimate song variation. Properties include frequency, frequency bandwidth, number of inflection points within a syllable, slope of the syllable, time duration, etc. These properties can differentiate syllables of different shapes. Rather than devoting ~24 days to process ~500,000 syllables, we applied a deep learning approach to automate the extraction of data from these recordings.

Convolutional neural networks (CNNs) are a type of deep machine learning algorithm that take image data and summarize across subsets of the images to identify important features in the image (see *26, 27*). CNNs and other deep learning algorithms can be trained to automate auditory

data collection because sound is easily converted to images with little loss of information (*28*).
Different sounds therefore have different shapes on a spectrogram (Fig. 1). The CNN can learn
the shapes we are interested in, then automatically identify those shapes. The corresponding
increase in the amount of data that can be analyzed is comparable to that which was enabled in
molecular genetics by advances in DNA sequencing technology that enabled researchers to move
from single locus data to whole genomes.

Heterogenous factors determine song variation within species

In 61% of New World Sparrows, song was not associated with the environment, year of
recording, or distance between recordings (Fig. 2). For the remaining 39%, the factors
determining variation in song within species were highly heterogeneous across Passerellidae.
Song was significantly associated with the environment in 19% of species, year of recording in
17% of species, and distance between recordings in 14% of species, with 12 species showing
multiple significant factors. Significant associations were more likely to be detected in species
with higher absolute song differentiation within-species (p=0.031). Genetic differentiation is
known to be associated with song differentiation (*15, 16*). We performed a secondary analysis on
a subset of 28 taxa with intraspecific genetic data using phylogatR (*29*). For 71% of species,
intraspecific genetic variation is not associated with environment or geographic distance. For the
remaining 29%, half of the species showed associations with the environment and half showed
association with geographic distance. For the 28 species that had both song and genetics
evaluated, 43% showed no associations between intra-specific variation and environment or
distance between recording. 29% had associations between song but not genetics, and 21%
associations between genetics but not song. For the remaining species, song was associated with
geographic distance but genetics was associated with environment. There is evidence that genetic
distance and song distance in other species is correlated (e.g., *30*), but in our dataset it seems that
generally they are not being caused by the same factors.
26

27
Fig. 2: Venn diagram displaying proportion of species whose songs are best predicted by
different features. Features are Environment ("Env." on plot, blue sun), Time (pink clock), and
Geography ("Geo." on plot, red ruler). Number outside of circles indicates number of species
with no significant association between song and features.

Using existing morphological data (*31*), we find that species whose songs are associated with the
environment have longer wings, longer tails, and longer secondary feathers (p<0.0291), as well
as being generally larger but not significantly so (p=0.056). Flight feather morphology is often
used as a proxy for dispersal in birds (*32*), with longer and pointier wings indicating better
presumed dispersal ability. This suggests that the influence of spatial factors on diversification

within species (i.e., in songs or genetics) is particularly pronounced on birds with poorer
dispersal ability.

Explanations of song variation are not associated with phylogeny

No major patterns emerge from an examination of phylogenetic relationships within the context
of the spatial analyses conducted here. While song production itself has a phylogenetic
component (*33*), the impact of phylogeny on the specific attributes of bird song is unclear.
Further, there is evidence that divergence at different taxonomic levels is not due to the same
process (e.g., *34*). If phylogeny is a large driver of spatiotemporal features in bird songs, we
would expect to find clades where one spatiotemporal feature was prominent in its impact. While
we see this within some genera (e.g., Atlapetes has a high proportion of taxa whose song is
associated with the environment), such a pattern does not appear at larger scales (Fig. 3). Our
results are thus consistent with phylogeny having some small effect on the evolution of bird
song, but additional work is needed to explore higher level effects in comparative phylogenetic
contexts. One note is that taxonomic uncertainty is a challenge to studies like the one conducted
here. As a potential hedge against the analysis of cryptic species, we performed the same
analyses using subspecies as our focal taxa (see supplement) and found similar results, though a
smaller proportion of taxa have significant relationships between spatiotemporal features and
song likely due to sample size. That the patterns are similar suggests that our findings are robust
to taxonomic uncertainty.

1
Fig. 3: Best predictors of species-level differences in traits. Left: phylogeny at the generic
level, with sample sizes of species sampled/total species. Right: heat map of number of species
sampled that show significant differences in environment (sun, left, blue), geographic distance
(ruler, center, red), or time (clock, right, pink). Colored values with an "X" indicate at least one
species shows significant differences, black values indicate no species shows significant
differences.

For the taxa wherein neither of these factors is important in explaining song variation, there is
still work to be done to determine what is influencing song evolution. Across many species, there
appears to be little song variation. This could result from small sample size, but also could be an
indication that song variation is not present. In contrast to taxa with little song variation, there are

a few taxa that have high song variation that are not explained by stochastic or environmental factors (e.g., *Pipilo maculatus*). One explanation might be that these species do not occupy environmentally variable areas, perhaps due to very strict niche requirements. However, we don't find any evidence that absolute environmental differences are driving these patterns. One further avenue to explore is that of acoustic niche partitioning (*35*), wherein species occupying the same areas differentiate their songs so as to minimize interference with other species' signals. This is important in both tropical bird communities (e.g., *36, 37*) and temperate ones (e.g., *38*). Lastly, it should be noted that our data are unable to investigate sexual and social selection, which are traditionally thought to be important determinants of song evolution across Passerines and birds more broadly (*4*).

Automatic processing of bioacoustic data at the family level has allowed us to address questions related to the factors that influence song evolution across a phylogenetic scale. The results presented here are, to our knowledge, the first large-scale intra-specific analysis of an entire bird family. Overall we find that intraspecific variation in song is not broadly correlated with any particular factor we tested, as the majority of Passerellidae species don't show correlations with environment, time, or distance. In species where there are correlations, it appears to be driven by different factors and is idiosyncratic across the phylogeny. Though it is possible that these results are dependent on data sampling, particularly since we are using repurposed data, our results suggest that the process of bird song evolution is not a deterministic one. Intraspecific diversity in songs is not explained entirely by phylogeny, geography, cultural drift, or environment. Rather, these factors contribute such that each individual taxon has its own unique response.

References and Notes

1. S. R. deKort, E. R. Eldermire, E. R. Cramer, S. L. Vehrencamp. The deterrent effect of bird song in territory defense. *Behavioral Ecology* **20**, 200-206 (2009).
2. D. Eriksson, L. Wallin. Male bird song attracts females—a field experiment. *Behavioral Ecology and Sociobiology* **19**, 297-299 (1986).
3. M. Brambilla, O. Janni, F. Guidali, A. Sorace. Song perception among incipient species as a mechanism for reproductive isolation. *Journal of Evolutionary Biology* **21**, 651-657 (2008).
4. C. K. Catchpole. Bird song, sexual selection and female choice. *Trends in Ecology and Evolution* **2**, 94-97 (1987).
5. H. Slabbekoorn, T. B. Smith. Bird song, ecology and speciation. *Philosophical Transactions of the Royal Society of London Series B: Biological Sciences* **357**, 493-503 (2002).
6. C. K. Catchpole, J. Dittami, B. Leisler. Differential responses to male song repertoires in female songbirds implanted with oestradiol. *Nature* **312**, 563-564 (1984).
7. E. P. Derryberry, N. Seddon, G. E. Derryberry, S. Claramunt, G. F. Seeholzer, R. T. Brumfield, J. A. Tobias. Ecological drivers of song evolution in birds: disentangling the effects of habitat and morphology. *Ecology and Evolution* **8**, 1890-1905 (2018).
8. E. P. Derryberry, R. M. Danner, J. E. Danner, G. E. Derryberry, J. N. Phillips, J. S. E. Lipshutz, K. Gentry, D. A. Luther. Patterns of song across natural and anthropogenic soundscapes suggest that white-crowned sparrows minimize acoustic masking and maximize signal content. *PLoS One* **11**, e0154456 (2016).
9. E. P. Derryberry, J. N. Phillips, G. E. Derryberry, M. J. Blum, D. Luther. Singing in a silent spring: Birds respond to a half-century soundscape reversion during the COVID-19 shutdown. *Science* **370**, 575-579 (2020).

10. G. C. Cardoso, T. D. Price. Community convergence in bird song. *Evolutionary Ecology* **24**, 447-461 (2010).

11. J. M. Trainer. Cultural evolution in song dialects of yellow☐rumped caciques in Panama. *Ethology* **80**, 190-204 (1989).

12. T. H. Parker, E. I. Greig, S. Nakagawa, M. Parra, A. C. Dalisio. Subspecies status and methods explain strength of response to local versus foreign song by oscine birds in meta-analysis. *Animal Behaviour* **142**, 1-17 (2018).

13. T. F. Wright, C. R. Dahlin. Vocal dialects in parrots: patterns and processes of cultural evolution. *Emu-Austral Ornithology* **118**, 50-66 (2018).

14. P. J. Slater. Bird song learning: causes and consequences. *Ethology Ecology and Evolution* **1**, 19-46 (1989).

15. N. A. Mason, K. J. Burns, J. A. Tobias, S. Claramunt, N. Seddon, E. P. Derryberry. Song evolution, speciation, and vocal learning in passerine birds. *Evolution* **71**, 786-796 (2017).

16. B. G. Freeman, G. A. Montgomery. Using song playback experiments to measure species recognition between geographically isolated populations: A comparison with acoustic trait analyses. *The Auk: Ornithological Advances* **134**, 857-870 (2017).

17. B. G. Freeman, J. Rolland, G. A. Montgomery, D. Schluter. Faster evolution of a premating reproductive barrier is not associated with faster speciation rates in New World passerine birds. *Proceedings of the Royal Society B* **289**, 20211514 (2022).

18. E. D. Jarvis, C. Scharff, M. R. Grossman, J. A. Ramos, F. Nottebohm. For whom the bird sings: context-dependent gene expression. *Neuron* **21**, 775-788 (1998).

19. C. Scharff, I. Adam. Neurogenetics of birdsong. *Current Opinion in Neurobiology* **23**, 29-36 (2013).

20. C. Dingle, J. W. Poelstra, W. Halfwerk, D. M. Brinkhuizen, H. Slabbekoorn. Asymmetric response patterns to subspecies☐specific song differences in allopatry and parapatry in the gray☐breasted wood☐wren. *Evolution* **64**, 3537-3548 (2010).

21. K. L. Provost, W. M. Mauck III, B. T. Smith. Genomic divergence in allopatric Northern Cardinals of the North American warm deserts is linked to behavioral differentiation. *Ecology and Evolution* **8**, 12456-12478 (2018).

22. S. E. Lipshutz, I. A. Overcast, M. J. Hickerson, R. T. Brumfield, E. P. Derryberry. Behavioural response to song and genetic divergence in two subspecies of white☐crowned sparrows (*Zonotrichia leucophrys*). *Molecular Ecology* **26**, 3011-3027 (2017).

23. N. Galtier. Half a Century of Controversy: The Neutralist/Selectionist Debate in Molecular Evolution. *Genome Biology and Evolution* **16**, evae003 (2024).

24. Y. Cohen, D. A. Nicholson, A. Sanchioni, E. K. Mallaber, V. Skidanova, T. J. Gardner. Automated annotation of birdsong with a neural network that segments spectrograms. *eLife* **11**, e63853 (2022).

25. K. L. Provost, J. Yang, B. C. Carstens. The impacts of fine-tuning, phylogenetic distance, and sample size on big-data bioacoustics. *PLoS One* **17,** e0278522 (2022).

26. Z. Li, F. Liu, W. Yang, S. Peng, J. Zhou. A survey of convolutional neural networks: analysis, applications, and prospects" *IEEE Transactions on Neural Networks and Learning Systems* **33**, 6999-7019 (2021).

27. I. Goodfellow, Y. Bengio, A. Courville. *Deep Learning*. (MIT Press, 2016).

28. G. Parascandolo, H. Huttunen, T. Virtanen. Recurrent neural networks for polyphonic sound event detection in real life recordings. *2016 IEEE International Conference on Acoustics Speech and Signal Processing*, 6440-6444 (2016).

29. T. A. Pelletier, D. J. Parsons, S. K. Decker, S. Crouch, E. Franz, J. Ohrstrom, B. C. Carstens. phylogatR: Phylogeographic data aggregation and repurposing. *Molecular Ecology Resources* **22**, 2830-2842 (2022).

30. R. E. Irwin. The evolutionary importance of behavioural development: the ontogeny and phylogeny of bird song. *Animal Behaviour* **36**, 814-824 (1988).

31. J. A. Tobias, C. Sheard, A. L. Pigot, A. J. Devenish, J. Yang, F. Sayol, M. H. C. Neate-Clegg, N. Alioravainen, T. L. Weeks, R. A. Barber, P. A. Walkden, H. E. A. MacGregor, S. E. I. Jones, C. Vincent, A. G. Phillips, N. M. Marples, F. A. Montaño-Centellas, V. Leandro-Silva, S. Claramunt, B. Darski, B. G. Freeman, T. P. Bregman, C. R. Cooney, E. C. Hughes, E. J. R. Capp, Z. K. Varley, N. R. Friedman, H. Korntheuer, A. Corrales-Vargas, C. H. Trisos, B. C. Weeks, D. M. Hanz, T. Töpfer, G. A. Bravo, V. Remeš, L. Nowak, L. S. Carneiro, A. J. Moncada R., B. Matysioková, D. T. Baldassarre, A. Martínez-Salinas, J. D. Wolfe, P. M. Chapman, B. G. Daly, M. C. Sorensen, A. Neu, M. A. Ford, R. J. Mayhew, L. Fabio Silveira, D. J. Kelly, N. N. D. Annorbah, H. S. Pollock, A. M. Grabowska-Zhang, J. P. McEntee, J. C. T. Gonzalez, C. G. Meneses, M. C. Muñoz, L.L. Powell, G. A. Jamie, T. J. Matthews O. Johnson, G. R. R. Brito, K. Zyskowski, R. Crates, M. G. Harvey, M. J. Zevallos, P. A. Hosner, T. Bradfer-Lawrence, J. M. Maley, F. Gary Stiles, H. S. Lima, K. L. Provost, M. Chibesa, M. Mashao, J. T. Howard, E. Mlamba, M. A. H. Chua, B. Li, M. I. Gómez, N. C. García, M. Päckert, J. Fuchs, J. R. Ali, E. P. Derryberry, M. L. Carlson, R. C. Urriza, K. E. Brzeski, D. M. Prawiradilaga, M. J. Rayner, E. T. Miller, R. C. K. Bowie, R. Lafontaine, R. P. Scofield, Y. Lou, L. Somarathna, D. Lepage, M. Illif, E. L. Neuschulz, M. Templin, D. M. Dehling, J. C. Cooper, O. S. G. Pauwels, K. Analuddin, J. Fjeldså, N. Seddon, P. R. Sweet, F. A. J. DeClerck, L. N. Naka, J. D. Brawn, A. Aleixo, K. Böhning-Gaese, C. Rahbek, S. A. Fritz, G. H. Thomas, M. Schleuning. AVONET: morphological, ecological and geographical data for all birds. *Ecology Letters* **25**, 581-597 (2022).

32. C. Sheard, M. H. Neate-Clegg, N. Alioravainen, S. E. Jones, C. Vincent, H. E. MacGregor, T. P. Bregman, S. Claramunt, J. A. Tobias. Ecological drivers of global gradients in avian dispersal inferred from wing morphology. *Nature Communications* **11**, (2020).

33. M. J. Ryan, E. A. Brenowitz. The role of body size, phylogeny, and ambient noise in the evolution of bird song. *The American Naturalist* **126**, 87-100 (1985).

34. A. F. Read, D. M. Weary. The evolution of bird song: comparative analyses. *Philosophical Transactions of the Royal Society of London Series B: Biological Sciences* **338**, 165-187 (1992).

35. B. L. Krause. The niche hypothesis: a virtual symphony of animal sounds, the origins of musical expression and the health of habitats. *The Soundscape Newsletter* **6** (1993).

36. E. Memet, B. Farrell, L. Mahadevan. An allometric prior enhances acoustic niche partitioning signal. *Journal of the Royal Society Interface* **19**, 20220421 (2022).

37. P. J. Hart, T. Ibanez, K. Paxton, G. Tredinnick, E. Sebastián-González, A. Tanimoto-Johnson. Timing is everything: acoustic niche partitioning in two tropical wet forest bird communities. *Frontiers in Ecology and Evolution* **9**, (2021).

38. A. Staniewicz, E. Sokołowska, A. Muszyńska, M. Budka. Competition for acoustic space in a temperate-forest bird community. *Behavioral Ecology* **34**, 1043-1054 (2023).

39. F. Gill, D. Donsker, P. Rasmussen. IOC World Bird List v13.1. (2023); https://www.worldbirdnames.org/

40. J. Klicka, F. K. Barker, K. J. Burns, S. M. Lanyon, I. J. Lovette, J. A. Chaves, R. W. Bryson Jr. A comprehensive multilocus assessment of sparrow (Aves: Passerellidae) relationships. *Molecular Phylogenetics and Evolution* **77**, 177-182 (2014).

41. F. K. Barker, K. J. Burns, J. Klicka, S. M. Lanyon, I. J. Lovette. New insights into New World biogeography: An integrated view from the phylogeny of blackbirds, cardinals, sparrows, tanagers, warblers, and allies. *The Auk: Ornithological Advances* **132**, 333-348 (2015).

42. R. W. Bryson Jr, B. C. Faircloth, W. L. Tsai, J. E. McCormack, J. Klicka. Target enrichment of thousands of ultraconserved elements sheds new light on early relationships within New World sparrows (Aves: Passerellidae). *The Auk: Ornithological Advances* **133**, 451-458 (2016).

43. D. W. Winkler, S. M. Billerman, I. J. Lovette. "New World Sparrows (Passerellidae), version 1.0" in *Birds of the World*, S. M. Billerman B. K. Keeney P. G. Rodewal T. S. Schulenberg, Eds. (Cornell Lab of Ornithology, 2020).

44. J. Yang, B. C. Carstens. K. L. Provost,. Machine learning reveals that climate, geography, and cultural drift all predict bird song variation in coastal *Zonotrichia leucophrys*. *Ornithology* **141**, ukad062 (2024).

45. J. Sueur, T. Aubin, C. Simonis. seewave: a free modular tool for sound analysis and synthesis. *Bioacoustics* **18**, 213-226 (2008).

46. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing Vienna Austria (2022). https://www.R-project.org/

47. M. Araya-Salas, G. Smith-Vidaurre. warbleR: an r package to streamline analysis of animal acoustic signals. *Methods in Ecology and Evolution* **8**, 184-191 (2017).

48. E. Paradis, K. Schliep. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35**, 526-528 (2019). doi:10.1093/bioinformatics/bty633.

49. S.E. Fick, R. J. Hijmans. WorldClim 2: new 1 km spatial resolution climate surfaces for global land areas. *International Journal of Climatology* **37**, 4302-4315 (2017).

50. M. Buchhorn, B. Smets, L. Bertels, B. De Roo, M. Lesiv, N. Tsendbazar, M. Herold, S. Fritz. Copernicus Global Land Service: Land Cover 100m: collection 3: epoch 2015: Globe V3.0.1, Zenodo, (2020). doi:10.5281/zenodo.3939038.

51. Earth Resources Observation and Science Center. USGS EROS Archive - Land Cover Products - Global Land Cover Characterization (GLCC) Background, U.S Geological Survey, (2018).

52. T. G. Farr, P. A. Rosen, E. Caro, R. Crippen, R. Duren, S. Hensley, M. Kobrick, M. Paller, E. Rodriguez, L. Roth, D. Seal, S. Shaffer, J. Shimada, J. Umland, M. Werner, M. Oskin, D. Burbank, D. Alsdorf. The Shuttle Radar Topography Mission. *Reviews of Geophysics* **45**, RG2004 (2007).

53. L. Poggio, L. M. De Sousa, N. H. Batjes, G. Heuvelink, B. Kempen, E. Ribeiro, D. Rossiter. SoilGrids 2.0: producing soil information for the globe with quantified spatial uncertainty. *Soil* **7**, 217-240 (2021).

54. E. C. Ellis, N. Gauthier, K. Klein Goldewijk, R. Bliege Bird, N. Boivin, S. Díaz, D. Q. Fuller, J. L. Gill, J. O. Kaplan, N. Kingston, H. Locke, C. N. H. McMichael, D. Ranco, T. C. Rick, M. R. Shaw, L. Stephens, J-C. Svenning, J. E. Watson. People have shaped most of terrestrial nature for at least 12,000 years. *Proceedings of the National Academy of Sciences* **118**, e2023483118 (2021).

55. I. J. Wang. Examining the full effects of landscape heterogeneity on spatial genetic variation: a multiple matrix regression approach for quantifying geographic and ecological isolation. *Evolution* **67**, 3403-3411 (2013).

56. J. Soons, A. Genbrugge, J. Podos, D. Adriaens, P. Aerts, J. Dirckx, A. Herrel. Is beak morphology in Darwin's finches tuned to loading demands? *PLoS One* **10**, e0129479 (2015).

57. M. Tokita, W. Yano, H. F. James, A. Abzhanov. Cranial shape evolution in adaptive radiations of birds: comparative morphometrics of Darwin's finches and Hawaiian honeycreepers. *Philosophical Transactions of the Royal Society B: Biological Sciences* **372** 20150481 (2017).

58. S. K. Huber, J. Podos.Beak morphology and song features covary in a population of Darwin's finches (*Geospiza fortis*). *Biological Journal of the Linnean Society* **88**, 489-498 (2006).

59. C. Sheard, S. Street, C. Evans, K. Lala, S. Healy, and S. Sugasawa. Beak shape and nest material use in birds. *Philosophical Transactions of the Royal Society B: Biological Sciences* **378**, 20220147 (2023).

60. R. Greenberg, V. Cadena, R. M. Danner, G. Tattersall. Heat loss may explain bill size differences between birds occupying different habitats. *PLoS One* **7**, e40933 (2012).

61. G. J. Tattersall, B. Arnaout, M. R. Symonds. The evolution of the avian bill as a thermoregulatory organ. *Biological Reviews* **92**, 1630-1656 (2017).

62. J. A. Bright, J. Marugán-Lobón, S. N. Cobb, E. J. Rayfield. The shapes of bird beaks are highly controlled by nondietary factors. *Proceedings of the National Academy of Sciences* **113**, 5352-5357 (2016).

63. W. W. Weathers. Physiological thermoregulation in heat-stressed birds: consequences of body size. *Physiological Zoology* **54**, 345-361 (1981).

64. J. Ryeland, M. A. Weston, M. R. Symonds. Bill size mediates behavioural thermoregulation in birds. *Functional Ecology* **31**, (2017).

65. A. J. C. Demery, K. J. Burns, N. A. Mason. Bill size, bill shape, and body size constrain bird song evolution on a macroevolutionary scale. *The Auk* **138**, ukab011 (2021).

66. R. G. Miller. *Simultaneous Statistical Inference*. (Springer-Verlag New York, 1981).

67. B. S. Yandell. *Practical Data Analysis for Designed Experiments*. (Chapman Hall/CRC Boca Raton, 1997).

Acknowledgments: We thank E. Akçay, R. Duckworth, and one anonymous reviewer for feedback on an earlier version of this manuscript. Resources from phylogatR and Imageomics were made possible by funding from the NSF and the Imageomics Institute. We thank all individuals who have contributed to song repositories over the last century, especially those at the Borror Laboratory of Bioacoustics.

Funding:

National Science Foundation DEB-1910623 (BCC)

National Science Foundation DEB-2016189 (BCC, KLP)

National Science Foundation OAC-2118240 (BCC)

Author contributions:

Conceptualization: BCC KLP

Data curation: KLP

Formal analysis: JY KLP

Funding acquisition: BCC

Investigation: KLP

Methodology: JY KLP

Resources: BCC KLP

Visualization: JY KLP

Writing – original draft: KLP

Writing – review and editing: BCC JY KLP

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: Datasets will be available upon acceptance. Code is available at https://github.com/kaiyaprovost/bioacoustics/

Supplementary Materials

Materials and Methods

Supplementary Text

Figs. S1 to S8

Tables S1 to S3