1 **TITLE**: Environmental factors and cultural drift influence song evolution in New World
2 Sparrows
3
4 **Keywords**: bird song, environmental adaptation, stochastic drift, MMRR, machine learning
5
6 **Article Type:** Major Article
7
8 **Number of words**: 7,017
9
10 **List of Elements in Manuscript**:
11 Title, Author List, Abstract, Introduction, Methods, Results, Discussion, Supplemental Materials,
12 Tables 1-3, Figures 1-6
13
14 **Running Head**: Environment in new world sparrow song
15

16 **Authors**: Kaiya L. Provost1,2*, Jiaying Yang2,3, Bryan C. Carstens2
17
1: Department of Biology, Adelphi University, One South Avenue, Garden City NY, 11530
USA.
2: Museum of Biological Diversity and Department of Evolution, Ecology and Organismal
Biology, The Ohio State University, 1315 Kinnear Rd., Columbus OH, 43212 USA.
3: Biological Science Department, Vanderbilt University, 2201 West End Ave, Nashville, TN
37235 USA.
24
25 * Corresponding author: kprovost@adelphi.edu
26

Acknowledgements
We thank all the individuals throughout the years who have contributed to the bioacoustics
repositories, including the Borror Laboratory of Bioacoustics and Xeno-Canto. For helpful
feedback on these analyses, we thank T. Yuri. S. Decker. S. Moshier, J. Wieringa, F. Mol Lanna,
T. Thome, E. Masiero, D. Duckett, J. Satler, S. Gaunt, D. Parsons, and J. Wallace. Funding for
KLP was provided by the National Science Foundation (DEB-2016189).
33

34 **Author Contribution Statement**
35 Conceptualization: KLP, BCC. Funding acquisition: BCC. Methods development: KLP. Data
36 collection, data analysis, data visualization: KLP, JY. Writing, editing: all authors.
37

Abstract

38
One of the most obvious traits of birds is the astonishing array of songs that are produced
by most species. The diversity and functions of bird vocalizations have been studied across many
facets of biology, but variation in song is often assumed to be determined by sexual or social
selection, rather than natural selection acting in the context of the landscape. Here we use deep
machine learning to investigate multiple co-distributed species in the New World Sparrows. We
leverage existing bioacoustic and genetic repositories and environmental data to identify the
processes that structure variation in bird song within a community, and to determine how this
variation corresponds to genetic and phenotypic patterns. Automated processing of these data at
the family level has allowed us to synthesize at deep phylogenetic scales. Analysis of these data
using a multiple matrix regression with randomization indicate that song variation in ~40% of
species can be explained by environmental and stochastic factors. Genetic variation is explained
by similar factors in ~30% of the species. This suggests that across a community and global
scale, the action of natural selection on the evolution of song is at least as impactful as it is on
other genetically-determined traits.
53

Introduction

Bird song (i.e., long and complex vocalizations associated with mating) is extremely
important for avian reproduction and therefore their Darwinian fitness. Field investigations have
indicated that song is used to mediate territorial disputes (e.g., de Kort et al 2009) and to
advertise to potential mates (e.g., Eriksson et al 1986). Bird song also plays an important role in
premating reproductive isolation (e.g., Brambilla et al 2008). Given how important song is to
reproduction, sexual and social selection are assumed to be the predominant forces shaping song
evolution (e.g., Catchpole 1987). This has led to a relative lack of attention to how landscape
level variation affects bird songs within species in large part because sexual and social selection
are expected to act as a homogenizing force (without the evolution of differences in preference)
if the function of the trait (i.e., song) is to maintain species boundaries, as commonly assumed
(e.g., Slabbekoorn and Smith 2002). Even in species where song is used by females to evaluate
the quality of potential mates, sexual selection should act to maintain the songs of males near the
fitness optimum. For example, Catchpole (et al 1984) shows that in sedge warblers, more
complex songs get more copulation solicitations from potential mates, and as such there is a
selective pressure for songs to become more complex through time, before hitting the
performance maximum. While there is a general expectation that sexual selection should lead to
a reduction in spatial variation of bird song across a species the obvious presence of spatial
variation within species necessitates in the form of dialects indicate that other forces act in
opposition to sexual selection to produce diversification. Dialects, which are regional variations
of communication, exist in bird species in the form of variation in songs through space and time
(Trainer 1989). Some dialect variations are associated with subspecific variation (e.g., Parker et
al 2018), but in others they do not have clear boundaries (e.g., Wright and Dahlin 2017). The
formation of dialects has numerous proposed causes.
One potential factor leading to the evolution of dialects is natural selection. This factor
has been suggested to influence intraspecific song variation by the relationship between habitat
and song in the ovenbird family (Furnariidae; Derryberry et al 2018). Dialect evolution may be
caused by environmental constraints, as songs in urban environments differ in their bandwidth
and frequency from rural environments, which may be an adaptation to avoid anthropogenic
noise (Derryberry et al 2016) but could potentially result from phenotypic plasticity (Derryberry

et al 2020). Indirect effects may also play a role due to constraints on other features; for example
Furnariids show a strong relationship between morphological variation and song (Derryberry et
al 2018). Whether direct or indirect, natural selection should act as a diversifying process at the
landscape scale (Cardoso and Price 2010) so long as the abiotic aspects contained within a
species' range are variable.
In contrast to natural selection, there are stochastic processes that could potentially
influence song evolution. For example, since song production has a genetic basis (Jarvis et al
1998, Scharff and Adam 2013), if the genomic DNA evolves due to genetic drift in violation of
Hardy-Weinberg expectations these changes may have an effect on song production. Several
investigations have suggested that species which exhibit more genetic differentiation also tend
also to be more acoustically differentiated (Freeman et al 2017, Freeman et al 2022).
Additionally, social learning, which occurs in many species of birds but particularly the Oscines
or "Songbirds", may be important. In such species, young individuals must learn songs from the
adults in the population (Slater 1989). Since this learning is imperfect, it can result in a
phenomenon known as cultural drift where songs change from are different from year to year
(e.g., Derégnaucourt and Gahr 2013). Since birds do not have infinite dispersal capabilities, any
differences that emerge due to cultural drive are likely to be localized in space. The presence of
song learning is known to influence the cadence at which song evolves (Mason et al 2016).
Finally, in any species population structure may also play a role. For example, decades of
phylogeographic results have suggested that species often persisted in multiple glacial refugia
during the Pleistocene (e.g., Brown Harrigan Lavretsky 2022, Pârâu and Wink 2021). Even in
the presence of selective constraints, song differentiation via these stochastic processes could
eventually lead to differentiation of song in different glacial refugia (e.g., Dingle et al 2010;
Provost et al 2018; Lipshutz et al 2018). Even after the retreat of the glaciers and the post-glacial
expansion of bird populations there is likely to be a remnant of intraspecific variation from this
process.
While each of the aforementioned factors (i.e., sexual and social selection, natural
selection, cultural drift) may influence the evolution of bird song, it is often assumed that the
major influence on the evolution of song has been sexual and social selection (e.g., Catchpole
1987). However, we lack a comprehensive understanding of the extent to which natural selection
and stochastic cultural drift matter. A major challenge in determining the impact of any factor on
song evolution is designing an investigation that can ascertain the relative influence of these
factors. Unlike genetic data, which are now easily collected (e.g., McCormack et al. 2013), and
environmental data, which are available in databases such as WorldClim (Fick et al 2017), data
on intraspecific variation in bird songs are challenging to obtain. The challenge is not so much in
obtaining the song data, as bird song recordings in bioacoustic databases cover much of the
globe, existing as combinations of citizen science efforts (e.g., eBird, Xeno-Canto), museum-
curated collections (e.g., Borror Lab of Bioacoustics, Macaulay Library), or *ad hoc* individual
recordings (e.g., uploads on Flickr, YouTube). However, these databases typically contain
repositories of the recordings themselves, rather than summaries of the variation in the songs.
Extraction of data from these recordings has traditionally been a challenging endeavor.
When song data are analyzed in research investigations, ornithologists have traditionally
annotated spectrograms of the recorded song by hand to extract data on intraspecific variations
(e.g., Latimer 1977, Wonke and Wallschläger 2009; see also Kvsn et al 2020), and taken
precautions to ensure that songs from different individuals are annotated in a consistent manner
(e.g., Duc et al 2021). However, it is not the case that songs from one study are annotated in the

same manner as a different set of songs from a different investigation, which makes it difficult to
conduct something like a meta-analysis using the results of many independent investigations.
The sheer time required to annotate a single bioacoustic recording when combined with the
challenges involved with comparing across studies have to date inhibited efforts to investigate
bird songs across broader scales. However, recent advances in artificial intelligence (AI)
methods for annotation of bird song (e.g., Cohen et al 2022) have enabled automated extraction
of summary statistics from bird song recordings that produce accurate data that are directly
comparable across species (e.g., Provost et al. 2022). These technological developments enable a
broader investigation into the factors that influence intraspecific variation bird songs.
Here, we seek to investigate the factors that potentially explain song diversity in one
focal group, the New World Sparrows (Family: Passerellidae). We test two hypotheses, which
are not mutually exclusive. First, we hypothesize that environmental selection will be an
important factor driving evolution across the New World Sparrow radiation, specifically that
climate, habitat type, and urbanization act to influence intraspecific variation in song. This
hypothesis predicts that, within a species, environmental distances will be correlated with
intraspecific differences in bird songs. A corollary is also tested, by evaluating the extent to
which morphological variation is correlated with intraspecific differences in song. The second
hypothesis predicts that stochastic cultural drift has influenced intraspecific variation in sparrow
song. Within a species, this hypothesis predicts that intraspecific variation in song is correlated
with the physical or temporal distance among individuals. Since neutral genetic variation is
predicted to vary in a similar manner, we also establish the extent to which there is genetic
isolation by distance in the Passerellidae.
152
153 **Methods**
154
Focal System: Passerellidae
Our work adopts the IOC taxonomy for determining species designations (Gill et al
2023). Under this framework, the New World Sparrows include 30 genera, 138 species, and 419
named subspecies (Gill et al 2023; Klicka et al 2014, Barker et al 2015, Bryson et al 2016). The
clade is distributed throughout the entirety of North and South America, save for the
northernmost latitudes, and occupying a broad range of habitats therein (Winkler et al 2020).
Phylogenetic relationships within this group vary, though taxa generally fall into eight major
clades (i.e., Klicka et al 2014). We use one recent phylogeny (Bryson et al 2016) at the genus
level to visualize phylogeographic patterns. Since some genera were missing from the Bryson et
al. phylogeny, their placement followed two previous phylogenies (Barker et al 2015, Klicka et
al 2014). The genus *Xenospiza* was placed as sister to *Centronyx*, which was in turn sister to
Melospiza. The genus *Torreornis* was placed as sister to *Pezopetes*, and similarly *Oreothraupis*
as sister to *Chlorospingus* and *Amphispizopsis* as sister to *Amphispiza*.
168
Acoustics
We downloaded all 10,836 songs of the family Passerellidae from Xeno-Canto (XC) on 5
July 2022. We supplemented these recordings with 12,145 songs from the Borror Lab of
Bioacoustics (BLB) collection. In both cases, previous studies had annotated songs from these
repositories (Provost el al 2022); we used those annotations to supplement our analysis. The few
individuals who were described in the metadata as hybrids were eliminated from the analysis.

Songs were converted to mono WAV format and to a sample rate of 48,000 Hz using the resamp
function in the seewave 2.2.0 package in R version 4.2.1 (Suer et al 2008; R Core Team 2022).
We used a machine learning model (i.e., TweetyNet; Cohen et al 2022) to identify the
locations of individual syllables in each recording . A model that had been previously trained on
nine species, including non-Passerellids (i.e., 9SppBalanced; Provost et al 2022) was applied to
the new analysis of recordings three ways. First, the model was used to identify syllables without
making any changes to the model. Second, we fine-tuned the model on only Passerellidae species
with annotations from another study (Provost et al 2022) and used that version to identify
syllables. Lastly, we fine-tuned the model again on all annotated Passerellidae that we had
generated as part of the present study. After each set of predictions of syllables had been made,
we extracted as syllables anything that the model had identified across any of the three sets of
predictions. We removed all syllables that were less than 0.05 seconds in length as spurious.
In a similar manner to how phenotypic traits are measured within samples before being
included in a quantitative analysis we extracted summary data from the song syllables. This was
accomplished using the warbleR package version 1.1.28 (Araya-Salas and Smith-Vidaurre 2017)
in R version 4.2.1. We calculated summary statistics based off of spectrograms using the
function spectro_analysis. The summary statistics included metrics of song duration; mean and
standard deviation of frequency; quantiles (including median, 25%, and 75%, and the
interquartile distance) of frequency; quantiles of time duration; skew and kurtosis of the
frequency spectrum; entropy measurements (spectral entropy, time entropy, and overall entropy);
spectral flatness; dominant frequency calculations (mean, minimum, maximum, range, and value
at start and end of sound); modulation index and slope of dominant frequency; peak frequency;
and mean peak frequency. We also calculated fundamental frequency time series using the
function freq_ts, which allowed us to get the frequency at each time bin. This lets us calculate
the number of inflections, slopes, and mean slope per sound.
After generating these summary statistics, we extracted five for further analysis:
bandwidth, as the IQR of frequency; time duration, as the IQR of the time; center, as the median
frequency; inflections, as the number of fundamental frequency inflection points; and slope, as
the mean fundamental frequency slope. From these five summary statistics we calculated values
for each syllable, then generated a centroid value for each individual across all syllables. We
performed a principal components analysis (PCA) on these data for downstream analyses (see
Supplement).
Once centroid values for the five summary statistics were calculated for all individuals,
we calculated euclidean distances between all individuals across all species. These euclidean
distances were used as representation of song variation as the predictor variable of the song (see
below). We then created subsets of these euclidean distances for each genus, species, and
subspecies across Passerellidae. Unless otherwise specified, we only report results for the species
distance matrices below -- for other analyses see Supplement. Finally, to serve as a proxy for the
overall distances represented within each taxon, we took the mean and standard deviation of
euclidean distance for each.
Recording quality varied across taxa, with some being of high quality (e.g., Xeno-Canto
quality A), low quality (e.g., Xeno-Canto quality E) or unknown quality (e.g., the BLB
recordings). We tested the impact of recording quality by repeating MMRR analyses (see below)
on subsets of homogenous recording quality for *Zonotrichia leucophrys*, which had the most
recordings (see Supplement).

Genetics

We downloaded geospatially labeled genetic data for all birds from GenBank using the phylogatR framework (Pelletier et al. 2022) on 7th December 2020. Of the 9,338,693 sequences present for all birds, Passerellidae make up 29,030 of those sequences representing 36 species. The overwhelming majority of the genes were CO1, so we restricted our analyses to that gene only.

From these genetic data we calculated genetic distances between individuals using the dist.gene function in the ape package version 5.7 in R. After that, we used those genetic distances as the predictor variable (see below). We also calculated mean genetic distance within species by taking the average pairwise across all individuals. We chose to do genetic analyses as a parallel because the song data and genetic data were not always collected from the same localities and as such cannot be directly compared in the same model. Instead we chose to compare and contrast the two models to see if similar patterns emerge.

Traits

Morphological differentiation, and phenotypic differentiation more broadly, is one major way that species adapt to their environment. In birds, bill morphology is thought to be selected with respect to diet (Soons et al 2015, Tokita et al 2017), song (Huber et al 2006), nest building (Sheard et al 2023), and thermoregulation (Greenberg et al 2012, Tattersall et al 2017), and constrained by phylogeny (Bright et al 2016). Like in other vertebrates, body size is a key factor for thermoregulation (Weathers 1981, Ryeland et al 2017), and is also important in influencing song (e.g., Demery et al 2021). To assess the impact of morphological measures on our data, we downloaded data from AVONET (Tobias et al 2022, Ecol Letters), which has measures of beak length (to nares and culmen), beak width, beak depth, tarsus length as a proxy for body size, primary and secondary feather length, tail length, the hand-wing index, and Kipp's distance.

This dataset contained 127 Passerellidae species across all 30 genera (missing: *Arremon basilicus, A. costaricensis, A. perijanus, A. phaeoplerus, Atlapetes meridae, At. nigrifrons, Junco bairdi, Passerella megarhyncha, P. unalaschcensis,* and *Pipilo naufragus*). There were 1,177 measurements taken across the family. We used the average value for each species as a factor when looking at the genetic and song MMRR results. We also calculated the average value for each genus. We evaluated whether morphology explained our results by using ANOVA and Tukey's Honest Significant Difference tests (Miller 1981; Yandell 1997). We evaluated significance using an alpha value of 0.05.

Spatial and Temporal Features

To examine the impact of the environment, we downloaded rasterized data from multiple sources representing various environmental features; climate, land cover, habitat type, topography, soil features, urbanization, geographic distance, and temporal distance (Fick et al 2017, Buchorn et al 2015, EROS 2018, Farr et al 2007, Poggio et al 2021, Ellis et al 2021). After removing variables that were invariable in our dataset, this resulted in 48 total variables. Geographic distance (latitude and adjusted longitude) and temporal distance (year of recording) were separated out as representing the stochastic processes of isolation-by-distance and cultural drift, respectively. Note that year of recording was only available for the song data. The remaining variables were used as representations of environmental data. We performed a PCA on the environmental data, centering and scaling the variables and imputing missing data such

266   that any missing values were assigned the mean. We did this using the prcomp function in the
267   stats package in R. We chose to retain five of the resulting PCs as they explained over 50% of
268   the variation across the 48 original variables, and these were used to represent environmental
269   distance. We calculated a euclidean distance matrix of geographic distance, temporal distance,
270   and environmental distance between all individuals in both the genetic and song datasets.
271
Synthesis
We examined biases in sampling in XC, BLB, and GenBank data for Passerellidae using
the locational and temporal metadata associated with each. We performed a multiple matrix
regression with randomization (MMRR) analysis to test our hypotheses (Wang, 2013).MMRR
estimates the coefficients of relationships between multiple distance matrices, controlling for the
impact of all of the variables in the dataset, and then uses a randomization technique to estimate
a p-value of significance. When an MMRR analysis is conducted with the same input matrices,
all of the returned coefficients are identical except for p-values. We conducted MMRR
comparisons multiple times for each dataset, randomizing each 1000 times, and then averaged
the p-values accordingly. For the song models each were set up so that song distance was being
predicted by environmental distance, geographic distance, and temporal distance. Because we
did not have temporal information for our genetic data, those models were set up so that genetic
distance was predicted by environmental and geographic distance only. If the environmental
selection hypothesis is supported, the MMRR will return a significant value p-value (using an
alpha value of 0.05) for the environmental distance matrix. If the stochastic cultural drift
hypothesis is supported, the MMRR will return a significant p-value either geographic distance,
temporal distance, or both. Both hypotheses may also be supported simultaneously.
MMRRs were performed at four levels: family, genus, species, and subspecies. We
performed these analyses at these different taxonomic levels to evaluate the impact of each factor
on progressively larger populations. Results other than species-level are found in the
supplemental material. We then compared taxa that were significant in environment or
geography for both song and genetics with their morphological measurements. We also
compared this significance with the absolute differences present in genetics and song across
species. We look at the phylogenetic relationships as well to see whether there is evidence that
these patterns are phylogenetically conserved.
297
298   **Results**
299
Data Summary
Sampling of the Passerellidae is uneven across North and South America in both song
and genetics (Figure 1). The western coast of North America, and Ohio, are highly sampled,
whereas much of the Amazon, central North America, and the Caribbean are undersampled. The
recordings vary in terms of time recorded as well (Figure 2). The BLB is a historical collection
and substantially older recordings (range 1948-2014, mean 1979) than XC (range 1976-2022,
mean 2013). Both databases also have peak recording from April to July, which corresponds to
the breeding season for most species. 80.0% of BLB songs were recorded during these months,
whereas 56.3% of XC songs were recorded during these months. Across space, sampling
between the recording data and the genetic data is significantly correlated though the correlation
is weak (log-corrected, p=0.030, adjR2 = 0.038).
311

312
Figure 1: Song and genetic sampling rarely overlap within species. A) Song sampling from
Xeno-Canto of all Passerellidae species. Color indicates density of recordings on a log scale,
with lighter reds showing areas with few recordings, darker reds showing areas of many
recordings, and gray showing areas with no recordings. Cells are one-degree latitude by one-
degree longitude. B) Genetic sampling from phylogatR of all Passerellidae species. Color
indicates density of recordings on a log scale, with lighter blues showing areas with few
sequences, darker blues showing areas of many sequences, and gray showing areas with no
sequences. Cells are one-degree latitude by one-degree longitude. C) Exemplar overlap for one
species, *Artemisiospiza bellii*. Points show individual sequences (blue cross) or song recordings
(red X). Color beneath shows whether the cell contains genetic data only (blue), song data only
(red), or both genetic and sign data (purple). Cells are one-degree latitude by one-degree
longitude.

326
327 Figure 2: A) Recordings in the Borror Laboratory of Bioacoustics database (BLB, dark red) and
328 Xeno-Canto (XC, red) across years. B) Recordings in the BLB (dark red) and XC (red) across
329 months of the year. C) Recordings per cell vs sequences per cell across range of Passerellidae.
330
Acoustics and Genetics
After processing all song recordings through the TweetyNet model (Cohen et al 2022),
our data consisted of 26,169 recordings and 513,617 annotated syllables. This comprised all 30
genera, 134 species, and 316 subspecies. All but one recording had latitude and longitude
information, and 23,544 had date information. Taxa ranged in terms of their overall song
differentiation. At the species level, the range was from 0.00 (*Chlorospingus inornatus*) to 4.32
(*Atlapetes personatus*). Quantitative data were extracted using summary statistics from each
species, which are available in Supplemental Table 3.
For our PCA on song variables, we retained all five PCs for downstream analyses (Table
1). Syllables with high values of PC1 were long, with numerous inflection points, and narrow
bandwidth. Syllables with high values of PC2 were high-frequency, wide-bandwidth. Syllables
with high PC3 had negative slopes and high frequencies. Syllables with high PC4 had narrow
bandwidths and positive slopes. Syllables with high PC5 had low frequencies, were short, and
had numerous inflection points.
345
346 Table 1: Principal components analysis on five song variables extracted from Passerellidae
347 syllables. Importance of each PC and loadings/rotation of each variable given.

		PC1	PC2	PC3	PC4	PC5
Importance	St. Dev.	1.245	1.023	0.994	0.990	0.660
	Prop. Of Variance	0.310	0.209	0.198	0.196	0.087
	Cumulative Prop.	0.310	0.519	0.717	0.913	1.000
Rotation	Bandwidth	-0.296	0.573	-0.304	-0.628	0.311
	Time	0.610	-0.001	-0.241	-0.465	-0.594
	Center	0.146	0.702	0.641	0.149	-0.228
	Inflection	0.710	0.023	0.061	0.004	0.701
	Slope	0.121	0.421	-0.659	0.606	-0.083

348
349   After compiling all of the genetic data, we ended up with 428 COI sequences across 20
350 genera and 36 species. There was no subspecific information available to us. Genetic
351 differentiation also ranged across taxa. At the species level, mean genetic distance ranged from
352 0.00 (*Chlorospingus flavopectus*) to 164.19 (*Arremon taciturnus*).

Spatial and Temporal Features

After performing our PCA of environmental variables, we retained the first five PCs as
they explained 53% of the data (Table 2, Table 3). High values of PC1 were associated with
regions that were dry, cold, had relatively narrow daily temperature but high annual temperature
ranges, and high temperature seasonality. High values of PC2 were associated with regions that
were cold and wet, with narrow daily temperature and yearly precipitation ranges, acidic sandy
soils, and lots of trees to the exclusion of shrubs, bare cover, grazing land, and irrigated land.
High values of PC3 were associated with more urbanized areas, high in croplands and
agriculture, with lots of seasonal water, low terrain ruggedness, and shallow slopes. High values
of PC4 were associated with cold summers, rough terrain, and lots of permanent water. Lastly,
high values of PC5 were associated with high population areas with lots of built up concrete, low
elevations, and low relief.

Figure 3: Breadth of syllable types across Passerellidae with one focal species highlighted. X-
axis and y-axis depict PC1 and PC2, respectively, of five song variables (Table 1). Each point is
an individual recording, with value showing the centroid across all syllables in that recording.
Highlighted in red is one species, *Amphispizopsis quinquestriata*, with all recordings and their
area given.

Table 2: Importance of first five PCs in a principal components analysis on environmental
variables.

	PC1	PC2	PC3	PC4	PC5
Standard deviation	3.10	2.39	2.00	1.93	1.56
Proportion of Variance	0.20	0.12	0.08	0.08	0.05
Cumulative Proportion	0.20	0.32	0.40	0.48	0.53

379 Table 3: Rotation/loadings of first five PCs in a principal components analysis on environmental
380 variables. Variables are categorized by general type.

		PC1	PC2	PC3	PC4	PC5
Land Cover %	Bare	0.076	-0.115	-0.060	0.039	-0.110
	Built Up	0.008	-0.004	0.083	0.159	0.330
	Crops	0.029	-0.026	0.163	-0.090	-0.016
	Grasses	0.116	-0.114	-0.052	-0.028	-0.273
	Moss/Lichens	0.013	0.021	-0.015	0.018	-0.042
	Permanent Water	0.029	0.024	0.059	0.065	0.006
	Seasonal Water	0.017	-0.011	0.054	0.007	-0.014
	Shrubs	0.018	-0.231	-0.163	-0.028	-0.137
	Trees	-0.136	0.197	-0.050	-0.089	0.015
Climate	Mean Annual Temp.	-0.234	-0.245	0.095	-0.092	0.097
	Mean Diurnal Temp. Range	0.137	-0.220	-0.134	-0.184	-0.071
	Isothermality	-0.264	-0.123	-0.103	0.044	-0.123
	Temp. Seasonality	0.276	0.110	0.097	-0.138	0.039
	Max Temp. Warmest Month	0.005	-0.267	0.139	-0.282	0.134
	Min Temp. Coldest Month	-0.282	-0.176	0.019	0.037	0.054
	Annual Temp. Range	0.284	0.046	0.049	-0.174	0.011
	Mean Temp. Warmest Quarter	-0.106	-0.176	0.196	-0.223	0.046
	Mean Temp. Driest Quarter	-0.202	-0.214	-0.072	0.052	0.064
	Mean Temp. Warmest Quarter	-0.072	-0.244	0.206	-0.245	0.168
	Mean Temp. Coldest Quarter	-0.272	-0.210	0.010	0.006	0.045
	Annual Precip.	-0.270	0.163	0.084	-0.062	-0.077
	Precip. Wettest Month	-0.282	0.076	0.018	0.002	-0.058
	Precip. Driest Month	-0.136	0.241	0.185	-0.171	-0.079
	Precip. Seasonality	-0.060	-0.252	-0.170	0.200	0.079
	Precip. Wettest Quarter	-0.279	0.091	0.024	0.006	-0.055
	Precip. Driest Quarter	-0.149	0.242	0.182	-0.167	-0.086
	Precip. Warmest Quarter	-0.204	0.100	0.124	-0.189	-0.141
	Precip. Coldest Quarter	-0.200	0.165	0.023	0.083	-0.006
Urbanization (by decade)	% Cropland	0.056	-0.061	0.242	-0.148	-0.040
	% Grazing Land	0.009	-0.207	0.005	-0.074	-0.206
	% Rice Cover	-0.001	-0.001	0.009	-0.003	-0.005
	Census Population	-0.002	-0.016	0.055	0.103	0.297
	% Land Irrigated	0.003	-0.117	0.057	-0.026	-0.031
	% Urban Landcover	0.017	-0.017	0.104	0.158	0.381
Habitat Classifications	Biosphere-Atmosphere Transfer Scheme	-0.009	0.002	0.088	0.112	-0.070
	International Geosphere Biosphere Programme	0.018	-0.069	0.363	0.238	-0.140
	USGS Land Use System	-0.033	0.121	-0.202	0.031	-0.143
	Olson Global Ecosystem	-0.001	-0.133	0.055	-0.111	-0.218
	Simple Biosphere 2 Model 2	0.072	-0.106	0.287	0.268	-0.234

	Simple Biosphere Model 1	0.052	-0.068	0.304	0.297	-0.224
	Running Vegetation Lifeforms	0.017	-0.050	0.269	0.225	-0.219
Topography	Elevation	-0.003	-0.036	-0.267	-0.038	-0.311
	Relief	-0.031	0.006	-0.013	0.024	-0.044
	Roughness	-0.071	0.018	-0.002	0.357	0.120
	Ruggedness Index	-0.149	-0.031	-0.214	0.129	-0.074
	Slope	-0.063	-0.022	-0.158	0.103	-0.005
Soil	Acidity	-0.165	0.244	0.026	-0.050	-0.015
	Texture Classification	0.143	0.154	-0.008	0.068	0.027

383
384 **Synthesis: MMRR**

385
Figure 4: relationship between song distance (y-axis) and spatiotemporal variables (x-axes) in
four species. Top two panels show song vs environmental distance, middle two panels show song
vs geographic distance, and bottom two panels show song vs temporal distance. Left column
shows three species where the relationship is significant (solid red line), the right column shows
one species where relationships are always non-significant (dashed red line).

Acoustics MMRR
393    For the MMRRs using song data there are 30 genera, 125 species, and 171 subspecies
with sufficient information to test the significance of environment, geography, and time (Figure
3, Supplementary Table 1, Supplementary Table 2). At the species level (Figure 4), 76/125
species showed no significant relationships between our spatial factors and song differentiation.

Of the rest, 37/125 had one factor significant: 16/125 were environment, 14/125 were time, and
7/125 were geography. An additional 11/125 had two factors significant: 5/125 for environment
and geography, 4/125 for geography and time, and 2/125 for environment and time. The
remaining one species had all three significant. This thus resulted in 24/125 species having the
environment being significant in some way, 21/125 having time significant, and 17/125 having
geography significant. Overall, this suggests that the environmental hypothesis is supported in
19.2% of species and the stochastic hypothesis in 26.4% of species.
404

405
406   Figure 5: Pie charts displaying proportion of species whose traits are best predicted by different
407   spatial and temporal features. Traits shown are songs (left) and genetics (center). Colors indicate
408   the significant predictors (see legend, right) -- shades of red are stochastic features (geography,
409   time), shades of blue are environmental features, purple/white indicate some combination, and
410   black indicates no such feature. Sample sizes are given below each plot.
411

412
Figure 6: Best predictors of species-level differences in traits. Left: phylogeny at the generic
level, with sample sizes of species sampled/total species. Right: heat map of number of species
sampled that show significant differences in song (orange shades) and genetics (teal shades).
Icons indicate spatial/temporal features tested: sun indicates environment, ruler indicates
geographic distance, and clock indicates time. Shades of color correspond to the number of
species where there are significant differences found, with darker shades showing more species
and white showing no species. Black boxes indicate no genetic data are available. For genus-
level and subspecies-level analyses, see Supplemental material.
421

Genetics MMRR

For the genetic MMRRs, there are 15 Passerellidae genera and 28 species with sufficient genetic information to test the significance of stochastic and environmental features (Figure 3, Supplemental Table 1, Supplemental Table 2). Only 8/28 species showed any influence of spatial factors on genetics: four with environment only, four with geography only. Overall, this suggests that the environmental hypothesis is supported in 14.2% of species and the stochastic hypothesis in 14.2% of species.

MMRR Comparisons

We compared whether song and genetics were explained by the same factors for the 28 species that had both genetic and song MMRRs (with the caveat that we did not evaluate genetic influence of time). For species, there were no taxa that had both gene and song explained by environment. However, 18/28 species had neither gene nor song explained by environment, with the remaining species being mismatched (four species with genes explained by environment, six species with song being explained). Likewise, there were no taxa that had both gene and song explained by geography. 20/28 species had neither gene nor song explained, the remaining eight were mismatched (four with genes being explained by geography, four with song being explained).

Traits

We found that there were some differences in morphology between taxa that showed environmental and stochastic influences on their genetics and song. At the species level, both vocalization and genetic differences were associated with morphology. In species that showed significant differences in genetics across geographic distances, both hand-wing indices and Kipp's distances were much shorter (p<0.029), which generally indicates that these birds have rounder, less pointed wings. With respect to vocalization, species that had significant correlations between song distances and environmental distances had longer wings (p=0.012), longer tails (p=0.0291), and longer secondaries (p=0.0275). Body size was also larger, though not significantly so (p=0.056). This is consistent with these birds being larger.

We investigated whether or not absolute differences in our traits and spatiotemporal features influenced the final results. Means and standard deviations are correlated with each other for each variable (range r: 0.72-0.93). Because of this, we only examine significant associations between mean values. Geography and environment are also highly correlated with each other, as expected (r=0.65). All other variables are correlated with each other to lesser extents (r<0.37).

At the species level, we find that we are more likely to detect significant differences between song and environment when the absolute song differences were higher (p=0.031). Again, this is easily explained, as when there is little song variation whatsoever the likelihood of detecting differentiation is slim.

We found no correlation between sample size and whether or not genetic differentiation was significant (p=0.751). However, there is a strong correlation with song differentiation (both with and without log-correcting for sample size disparity) such that taxa with more recordings are more likely to have a significant correlation detected (p<4.9x10^{-6}).

Discussion

Automatic processing of bioacoustic data at the family level has allowed us to address questions related to the factors that influence song evolution across a phylogenetic scale. In New World sparrows, stochastic factors such as geographic distance or temporal distance (i.e., cultural drift) are generally more important than environmental factors in structuring song diversity within species, although there are individual species where each factor is approximately equally important. Across species, environment explains variation in 19% of species and stochastic factors in 26% of species (13% geography, 16% time). In particular, cultural drift is at least as important to the evolution of bird song as genetic drift is to the evolution of intraspecific genetic diversity. Within genera, these numbers are 33% for environment and 43% for stochastic (23% geography, 33% time). This suggests to us that both stochastic drift and the environment play a large role in shaping sound at these two taxonomic levels.

For the taxa wherein neither of these factors is important in explaining song variation, there is still work to be done to determine what is influencing song evolution. Across many species, there appears to be little song variation. This could result from small sample size, but also could be an indication that song variation is not present. We believe this highlights the need for deeper sampling. At present, we did not incorporate recordings from other sources (e.g., Macaulay Library) due to concerns about processing time.

In contrast to taxa with little song variation, there are a few taxa that have high song variation that are not explained by stochastic or environmental factors (e.g., *Pipilo maculatus*). One explanation might be that these species do not occupy environmentally variable areas, perhaps due to very strict niche requirements. However, we don't find any evidence that absolute environmental differences are driving these patterns. Alternatively, sexual and social selection may cause this pattern; however our study is not set up to detect these processes. Sexual and social selection are generally thought to be important determinants of song evolution across Passerines and birds more broadly (Catchpole 1987). These processes were not included because the song data contained in existing repositories lack the requisite metadata required for such an analysis. For example, estimating the impact of sexual selection requires knowledge about mate choice or mate competition, and in many species even determining the sex of the individual during the recording process is difficult.

Phylogeny

No major patterns emerge from an examination of phylogenetic relationships within the context of the spatial analyses conducted here. While song production itself has a phylogenetic component (Ryan et al 1985), the impact of phylogeny on the specific attributes of bird song is unclear. Further, there is evidence that divergence at different taxonomic levels are not due to the same process (e.g., Read and Weary 1992). If phylogeny is a large driver of spatiotemporal features in bird songs, we would expect to find clades where one spatiotemporal feature was prominent in its impact. While we see this within some genera (e.g., *Atlapetes* has a high proportion of taxa whose song is associated with the environment), such a pattern does not appear at larger scales. Our results are thus consistent with phylogeny having some small effect on the evolution of bird song, but additional work is needed to explore higher level effects in comparative phylogenetic contexts. One note is that taxonomic uncertainty is a challenge to studies like the one conducted here. As a potential hedge against the analysis of cryptic species, we performed the same analyses using subspecies as our focal taxa (see supplement) and found similar results, though a smaller proportion of taxa have significant relationships between

514 spatiotemporal features and song likely due to sample size. That the patterns are similar suggests
515 that our findings are robust to taxonomic uncertainty.
516
Genetics, behavior, and morphology
Genetic and behavioral patterns are fairly similar, with 43% of species having the exact
same spatiotemporal factors explaining genes and song and an additional 50% having a partial
match (note that this ignores the temporal aspect present only for song). This is in line with
evidence that genetic distance and song distance in other species is correlated (e.g., Irwin 1988).
Here, we found that morphology is associated with differences across species for both genetics
and songs.
In songs, species whose vocalizations are associated with environmental adaptation tend
to have longer flight feathers, which may represent larger body size. In contrast, at the generic
level, genera with shorter beaks showed more influence of isolation-by-distance. For genetics,
isolation-by-distance was associated with species having stubbier wings. Morphological
differentiation in birds is often associated with different factors. Beak shape is a constraint on
diet (Navalón et al 2019), song frequency and bandwidth (Derryberry et al 2012), and
thermoregulation (Tattersall et al 2017, Ryeland et al 2017). In this study, birds whose song
properties are likely to be influenced by geography are also likely to have shorter beaks, but
whether or not this is due to direct selection on song properties or indirect selection on other
beak-associated traits is presently unknown. Tarsus length is generally associated with body size
in birds (Senar and Pascual 1997), although mass itself was not found to be associated with any
of the MMRR results. Nevertheless, body size is also associated with song properties (e.g., Ryan
et al 1985) and thermoregulation (via Allen's rule; Symonds and Tattersall 2010) and therefore
the same conclusions apply as for beak size. Lastly, wing morphology is often used as a proxy
for dispersal in birds (Shard et al 2020). Birds with smaller hand-wing index, therefore stubbier
wings and less presumed dispersal ability, were more likely to have genetics explained by
isolation-by-distance. This suggests that the influence of spatial factors on diversification within
species (i.e., in songs or genetics) is particularly pronounced on birds with poorer dispersal
ability.
543
Conclusion
The comparative analyses presented here are the first attempt to understand the factors
that influence song evolution in birds on a phylogenetic scale. Our results indicate that spatial,
temporal, and environmental factors exert a clear influence on song evolution within many
species. In scale, they mirror our results that tested for an association between geographic and
genetic distance. This suggests that cultural drift is at least as important to the evolution of bird
song as genetic drift is to the evolution of intraspecific genetic diversity. As with their genomic
variation (e.g., Armstrong et al. 2018), bird songs are likely influenced by both cultural drift and
selection. Taken in aggregate, it appears that spatial-associated factors are at least as important in
shaping the evolution of song within bird species as geographic distance is a factor in
intraspecific genetic differentiation.
555

556 **References**
557 Araya-Salas, M. and G. Smith-Vidaurre. 2017. warbleR: an r package to streamline analysis of
558   animal acoustic signals. Methods in Ecology and Evolution 8:184-191.
559 Armstrong, C., D.S. Richardson, H. Hipperson, G. J. Horsburgh, C. Küpper, L. Percival-Alwyn,
560   M. Clark, T. Burke, and L. G. Spurgin. 2018. Genomic associations with bill length and

disease reveal drift and selection across island bird populations. Evolution Letters 2:22-36.

Barker, F. K., K. J. Burns, J. Klicka, S. M. Lanyon, and I. J. Lovette. 2015. New insights into New World biogeography: An integrated view from the phylogeny of blackbirds, cardinals, sparrows, tanagers, warblers, and allies. The Auk: Ornithological Advances 132(2):333-348.

Brambilla, M., O. Janni, F. Guidali, and A. Sorace. 2008. Song perception among incipient species as a mechanism for reproductive isolation. Journal of Evolutionary Biology 21(3):651-657.

Bright, J. A., J. Marugán-Lobón, S. N. Cobb, and E. J. Rayfield. 2016. The shapes of bird beaks are highly controlled by nondietary factors. Proceedings of the National Academy of Sciences 113(19):5352-5357.

Brown, J. I., R. J. Harrigan, and P. Lavretsky. 2022. Evolutionary and ecological drivers of local adaptation and speciation in a North American avian species complex. Molecular Ecology 31:2578-2593.

Bryson Jr, R. W., B. C. Faircloth, W. L. Tsai, J. E. McCormack, and J. Klicka. 2016. Target enrichment of thousands of ultraconserved elements sheds new light on early relationships within New World sparrows (Aves: Passerellidae). The Auk: Ornithological Advances 133(3):451-458.

Buchhorn, M., B. Smets, L. Bertels, B. De Roo, M. Lesiv, N. Tsendbazar, M. Herold, and S. Fritz. 2020. Copernicus Global Land Service: Land Cover 100m: collection 3: epoch 2015: Globe (V3.0.1) [Data set]. Zenodo. https://doi.org/10.5281/zenodo.3939038

Cardoso, G. C., and T. D. Price. 2010. Community convergence in bird song. Evolutionary Ecology 24:447-461.

Catchpole, C. K. 1987. Bird song, sexual selection and female choice. Trends in Ecology and Evolution 2:94-97.

Catchpole, C. K., J. Dittami, and B. Leisler. 1984. Differential responses to male song repertoires in female songbirds implanted with oestradiol. Nature 312:563-564.

Cohen, Y., D. A. Nicholson, A. Sanchioni, E. K. Mallaber, V. Skidanova, V., and T. J. Gardner. 2022. Automated annotation of birdsong with a neural network that segments spectrograms. Elife 11:e63853.

de Kort, S. R., Eldermire, E. R., Cramer, E. R., and Vehrencamp, S. L. 2009. The deterrent effect of bird song in territory defense. Behavioral Ecology, 20:200-206.

Demery, A. J. C., K. J. Burns, and N. A. Mason. 2021. Bill size, bill shape, and body size constrain bird song evolution on a macroevolutionary scale. The Auk 138:ukab011.

Derégnaucourt, S., and M. Gahr. 2013. Horizontal transmission of the father's song in the zebra finch (*Taeniopygia guttata*). Biology Letters, 9:20130247.

Derryberry, E. P., N. Seddon, S. Claramunt, J. A. Tobias, A. Baker, A. Aleixo, and R. T. Brumfield. 2012. Correlated evolution of beak morphology and song in the neotropical woodcreeper radiation. Evolution 66:2784-2797.

Derryberry, E. P., R. M. Danner, J. E. Danner, G. E. Derryberry, J. N. Phillips, J. S. E. Lipshutz, K. Gentry, and D. A. Luther. 2016. Patterns of song across natural and anthropogenic soundscapes suggest that white-crowned sparrows minimize acoustic masking and maximize signal content. PLoS One 11:e0154456.

Derryberry, E. P., N. Seddon, G. E. Derryberry, S. Claramunt, G. F. Seeholzer, R. T. Brumfield, and J. A. Tobias,. 2018. Ecological drivers of song evolution in birds: disentangling the effects of habitat and morphology. Ecology and evolution 8:1890-1905.

Derryberry, E. P., J. N. Phillips, G. E. Derryberry, M. J. Blum, and D. Luther. 2020. Singing in a silent spring: Birds respond to a half-century soundscape reversion during the COVID-19 shutdown. Science 370:575-579.

Dingle, C., J. W. Poelstra, W. Halfwerk, D. M. Brinkhuizen, and H. Slabbekoorn. 2010. Asymmetric response patterns to subspecies-specific song differences in allopatry and parapatry in the gray-breasted wood-wren. Evolution 64:3537-3548.

Duc, P. N. H., M. Torterotot, F. Samaran, P. R. White, O. Gérard, O. Adam, and D. Cazau. 2021. Assessing inter-annotator agreement from collaborative annotation campaign in marine bioacoustics. Ecological Informatics 61:101185.

Earth Resources Observation and Science Center. 2018. USGS EROS Archive - Land Cover Products - Global Land Cover Characterization (GLCC) Background. U.S. Geological Survey.

Ellis, E. C., N. Gauthier, K. Klein Goldewijk, R. Bliege Bird, N. Boivin, S. Díaz, D. Q. Fuller, J. L. Gill, J. O. Kaplan, N. Kingston, H. Locke, C. N. H. McMichael, D. Ranco, T. C. Rick, M. R. Shaw, L. Stephens. J-C. S., and J. E. Watson. 2021. People have shaped most of terrestrial nature for at least 12,000 years. Proceedings of the National Academy of Sciences 118:e2023483118.

Eriksson, D., and L. Wallin. 1986. Male bird song attracts females—a field experiment. Behavioral Ecology and Sociobiology 19:297-299.

Farr, T. G., P. A. Rosen, E. Caro, R. Crippen. R. Duren, S. Hensley, M. Kobrick, M. Paller. E. Rodriguez, L. Roth, D. Seal, S. Shaffer, J. Shimada, J/ Umland, M. Werner, M. Oskin, D. Burbank, and D. Alsdorf. 2007. The Shuttle Radar Topography Mission. Reviews of Geophysics 45:RG2004.

Fick S.E., and R. J. Hijmans. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. International Journal of Climatology 37(12):4302-4315.

Freeman, B. G., and G. A. Montgomery. 2017. Using song playback experiments to measure species recognition between geographically isolated populations: A comparison with acoustic trait analyses. The Auk: Ornithological Advances 134:857-870.

Freeman, B. G., J. Rolland, G. A. Montgomery, and D. Schluter. 2022. Faster evolution of a premating reproductive barrier is not associated with faster speciation rates in New World passerine birds. Proceedings of the Royal Society B 289:20211514.

Gill, F., D. Donsker, and P. Rasmussen (eds). 2023. IOC World Bird List (v13.1). https://www.worldbirdnames.org/

Greenberg, R., V. Cadena, R. M. Danner, and G. Tattersall. 2012. Heat loss may explain bill size differences between birds occupying different habitats. PLoS One 7(7):e40933.

Huber, S. K., and J. Podos. 2006. Beak morphology and song features covary in a population of Darwin's finches (*Geospiza fortis*). Biological Journal of the Linnean Society 88(3):489-498.

Irwin, R. E. 1988. The evolutionary importance of behavioural development: the ontogeny and phylogeny of bird song. Animal Behaviour 36(3):814-824.

Jarvis, E. D., C. Scharff, M. R. Grossman, J. A. Ramos, and F. Nottebohm. 1998. For whom the bird sings: context-dependent gene expression. Neuron 21(4):775-788.

Klicka, J., F. K. Barker, K. J. Burns, S. M. Lanyon, I. J. Lovette, J. A. Chaves, and R. W. Bryson Jr. 2014. A comprehensive multilocus assessment of sparrow (Aves: Passerellidae) relationships. Molecular Phylogenetics and Evolution 77:177-182.

Kvsn, R. R., J. Montgomery, S. Garg, and M. Charleston. 2020. Bioacoustics data analysis–A taxonomy, survey and open challenges. IEEE Access 8:57684-57708.

Latimer, W. 1977. A comparative study of the songs and alarm calls of some *Parus* species. Zeitschrift für Tierpsychologie 45: 414-433.

Lipshutz, S. E., I. A. Overcast, M. J. Hickerson, R. T. Brumfield, and E. P. Derryberry. 2017. Behavioural response to song and genetic divergence in two subspecies of white-crowned sparrows (*Zonotrichia leucophrys*). Molecular Ecology 26:3011-3027.

Mason, N. A., K. J. Burns, J. A. Tobias, S. Claramunt, N. Seddon, and E. P. Derryberry. 2017. Song evolution, speciation, and vocal learning in passerine birds. Evolution, 71:786-796.

McCormack, J. E., S. M. Hird, A. J. Zellmer, B. C. Carstens, and R. T. Brumfield. 2013. Applications of next-generation sequencing to phylogeography and phylogenetics. Molecular Phylogenetics and Evolution 66:526-538.

Miller, R. G. 1981. Simultaneous Statistical Inference. Springer-Verlag, New York.

Navalón, G., J. A. Bright, J. Marugán-Lobón, and E. J. Rayfield. 2019. The evolutionary relationship among beak shape, mechanical advantage, and feeding ecology in modern birds. Evolution 73:422-435.

Pârâu, L. G., and M. Wink. 2021. Common patterns in the molecular phylogeography of western palearctic birds: a comprehensive review. Journal of Ornithology 162:937-959.

Parker, T. H., E. I. Greig, S. Nakagawa, M. Parra, and A. C. Dalisio. 2018. Subspecies status and methods explain strength of response to local versus foreign song by oscine birds in meta-analysis. Animal Behaviour 142:1-17.

Pelletier, T. A., D. J. Parsons, S. K. Decker, S. Crouch, E. Franz, J. Ohrstrom, and B. C. Carstens 2022. phylogatR: Phylogeographic data aggregation and repurposing. Molecular Ecology Resources 22:2830-2842.

Poggio, L., L. M. De Sousa, N. H. Batjes, G. Heuvelink, B. Kempen, E. Ribeiro, and D. Rossiter. 2021. SoilGrids 2.0: producing soil information for the globe with quantified spatial uncertainty. Soil 7(1):217-240.

Provost, K. L., W. M. Mauck III, and B. T. Smith. 2018. Genomic divergence in allopatric Northern Cardinals of the North American warm deserts is linked to behavioral differentiation. Ecology and Evolution 8(24):12456-12478.

Provost, K. L., J. Yang, and B. C. Carstens. 2022. The impacts of fine-tuning, phylogenetic distance, and sample size on big-data bioacoustics. PLOS ONE 17(12):e0278522.

R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Ryan, M. J., and E. A. Brenowitz. 1985. The role of body size, phylogeny, and ambient noise in the evolution of bird song. The American Naturalist 126:87-100.

Ryeland, J., M. A. Weston, and M. R. Symonds. 2017. Bill size mediates behavioural thermoregulation in birds. Functional Ecology 31:885-893.

Scharff, C., and I. Adam. 2013. Neurogenetics of birdsong. Current Opinion in Neurobiology 23(1):29-36.

Senar, J. C., and J. Pascual. 1997. Keel and tarsus length may provide a good predictor of avian body size. Ardea 85:269-274.

Sheard, C., M. H. Neate-Clegg, N. Alioravainen, S. E. Jones, C. Vincent, H. E. MacGregor, T. P. Bregman, S. Claramunt, and J. A. Tobias. 2020. Ecological drivers of global gradients in avian dispersal inferred from wing morphology. Nature Communications 11:2463.

Sheard, C., S. Street, C. Evans, K. Lala, S. Healy, and S. Sugasawa. 2023. Beak shape and nest material use in birds. Philosophical Transactions of the Royal Society B: Biological Sciences 378:20220147.

Slabbekoorn, H., and T. B. Smith. 2002. Bird song, ecology and speciation. Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences 357:493-503.

Slater, P. J. 1989. Bird song learning: causes and consequences. Ethology Ecology and Evolution 1(1):19-46.

Soons, J., A. Genbrugge, J. Podos, D. Adriaens, P. Aerts, J. Dirckx, and A. Herrel. 2015. Is beak morphology in Darwin's finches tuned to loading demands? PLoS One, 10:e0129479.

Sueur, J., T. Aubin, and C. Simonis. 2008. seewave: a free modular tool for sound analysis and synthesis. Bioacoustics 18:213-226

Symonds, M. R., and G. J. Tattersall. 2010. Geographical variation in bill size across bird species provides evidence for Allen's rule. The American Naturalist 176(2):188-197.

Tattersall, G. J., B. Arnaout, and M. R. Symonds. 2017. The evolution of the avian bill as a thermoregulatory organ. Biological Reviews 92(3):1630-1656.

Tobias, J. A., C. Sheard, A. L. Pigot, A. J. M. Devenish, J. Yang, M. H. C. Neate-Clegg, N. Alioravainen, et al. 2022. AVONET: morphological, ecological and geographical data for all birds. Ecology Letters 25:581–597.

Tokita, M., W. Yano, H. F. James, and A. Abzhanov. 2017. Cranial shape evolution in adaptive radiations of birds: comparative morphometrics of Darwin's finches and Hawaiian honeycreepers. Philosophical Transactions of the Royal Society B: Biological Sciences 372(1713):20150481.

Trainer, J. M. 1989. Cultural evolution in song dialects of yellow-rumped caciques in Panama. Ethology 80:190-204.

Trujillo-Torres, C. M., F. González-García, and I. MacGregor-Fors. 2022. Say what? On the transmission of acoustic signals in a Neotropical green city. Urban Ecosystems 25(1):1-8.

Wang, I. J. 2013. Examining the full effects of landscape heterogeneity on spatial genetic variation: a multiple matrix regression approach for quantifying geographic and ecological isolation. Evolution 67(12):3403-3411.

Weathers, W. W. 1981. Physiological thermoregulation in heat-stressed birds: consequences of body size. Physiological Zoology 54(3):345-361.

Winkler, D. W., S. M. Billerman, and I. J. Lovette 2020. New World Sparrows(*Passerellidae*), version 1.0. *In* S. M. Billerman, B. K. Keeney, P. G. Rodewal, T. S. Schulenberg, eds. Birds of the World. Cornell Lab of Ornithology, Ithaca, NY.

Wonke, G., and D. Wallschläger. 2009. Song dialects in the yellowhammer *Emberiza citrinella*: bioacoustic variation between and within dialects. Journal of Ornithology 150:117-126.

Wright, T. F., and C. R. Dahlin. 2018. Vocal dialects in parrots: patterns and processes of cultural evolution. Emu-Austral Ornithology 118(1):50-66.

Yandell, BS. 1997. Practical data analysis for designed experiments. Chapman Hall/CRC, Boca Raton.