1   Comparison of insect and human cytochrome b561 proteins: Insights into candidate ferric reductases in

2   insects

Jessica D. Holst[1], Laura G. Murphy[1], Maureen J. Gorman[2], Emily J. Ragan[1]*

[1]Department of Chemistry and Biochemistry, Metropolitan State University of Denver, Denver,
Colorado, United States of America

[2]Department of Biochemistry and Molecular Biophysics, Kansas State University, Manhattan, Kansas,
United States of America

*Corresponding author

E-mail: eragan@msudenver.edu (EJR)

Abstract

Cytochrome b561 (cytb561) proteins comprise a family of transmembrane oxidoreductases that transfer single electrons across a membrane. Most eukaryotic species, including insects, possess multiple cytb561 homologs. To learn more about this protein family in insects, we carried out a bioinformatics-based investigation of cytb561 family members from nine species representing eight insect orders. We performed a phylogenetic analysis to classify insect cytb561 orthologous groups. We then conducted sequence analyses and analyzed protein models to predict structural elements that may impact the biological functions and localization of these proteins, with a focus on possible ferric reductase activity. Our study revealed three orthologous groups, designated CG1275, Nemy, and CG8399, and a fourth group of less-conserved genes. We found that CG1275 and Nemy proteins are similar to a human ferric reductase, duodenal cytochrome b561 (Dcytb), and have many conserved amino acid residues that function in substrate binding in Dcytb. Notably, CG1275 and Nemy proteins contain a conserved histidine and other residues that play a role in ferric ion reduction by Dcytb. Nemy proteins were distinguished by a novel cysteine-rich cytoplasmic loop sequence. CG8399 orthologs are similar to a putative ferric reductase in humans, stromal cell-derived receptor 2. Like other members of the CYBDOM class of cytb561 proteins, these proteins contain reeler, DOMON, and cytb561 domains. *Drosophila melanogaster* CG8399 is the only insect cytb561 with known ferric reductase activity. Our investigation of the DOMON domain in CG8399 proteins revealed a probable heme-binding site and a possible site for ferric reduction. The fourth clade includes a group of proteins with a conserved "KXXXXKXH" non-cytoplasmic loop motif that may be a substrate binding site and is present in a potential ferric reductase, human tumor suppressor cytochrome b561. This study provides a foundation for future investigations of the biological functions of cytb561 genes in insects.

Introduction

The cytochrome b561 (cytb561) proteins are a family of transmembrane oxidoreductases that

function as electron transporters by accepting a single electron from a cytoplasmic reducing agent –

generally assumed to be ascorbate – and transferring it across a membrane to reduce a non-cytoplasmic

substrate [1,2]. Known non-cytoplasmic substrates (electron acceptors) include ferric (Fe^{3+}) ions, which

are reduced to ferrous (Fe^{2+}) ions, and monodehydroascorbate (MDHA), which is reduced to ascorbate

[2]. Animal and plant species possess multiple cytb561 homologs, some with ubiquitous expression and

others with restricted expression patterns [1–3]. Depending on the particular protein, cytb561s can be

located in the plasma membrane, the membranes of organelles in the secretory or endocytic pathways,

or multiple membrane locations [4–7].

The biological functions of most cytb561s are uncertain, although some have established roles in

iron reduction or ascorbate regeneration [1]. The well-characterized human duodenal cytochrome b561

(Dcytb) is a ferric reductase that facilitates iron uptake at the duodenal brush border, and it is also

suspected to play other physiological roles [8–10]. The structurally well-characterized *Arabidopsis*

thaliana (plant) cytochrome b561 B-paralog (AtCytb561-B) has in vitro ferric-chelate reductase activity,

but its biological functions are unknown [11–13]. Other cytb561 family members with in vitro ferric

reductase activity include mammalian stromal cell-derived receptor 2 (SDR2), tumor suppressor 101F6

(TScytb), chromaffin granule cytb561 (CGcytb), and lysosomal cytb561 (Lcytb), as well as *Drosophila*

melanogaster (fly) CG8399, and *A. thaliana* AtCytb561-A [6,11,12,14,15]. The physiological significance

of these proteins' ferric reductase activity is not clear; for example, although CGcytb can reduce ferric-

chelates in vitro, it is thought to function in chromaffin granules by reducing MDHA to produce

ascorbate, which is then used by dopamine β-hydroxylase (DBH) in the synthesis of norepinephrine

[16,17]. It seems likely that at least some cytb561s are multifunctional proteins, and that substrate

specificity may be influenced by the substrate availability where a particular cytb561 is located.

To date, two cytb561 structures have been solved: human Dcytb and plant AtCytb561-B [13,18].

These structures confirmed many predicted cyb561 structural elements. Cytb561s have six

transmembrane helices, including four consecutive helices that form a cytb561 core domain [2,13,18].

Inside the core domain are two b-type heme groups located near each side of the membrane, which are

coordinated by four universally conserved histidine residues: the first- and third-conserved histidines

(H1, H3) coordinate the non-cytoplasmic heme group, while the second- and fourth-conserved histidines

(H2, H4) coordinate the cytoplasmic heme group [2,19].

Most cytb561s are single-domain proteins in which the core domain encompasses

transmembrane helices two through five (TM2 through TM5) [2,13,18]. An exception is the CYBDOM

proteins, which are multidomain cytb561s characterized by having at least one extracellular DOMON

domain upstream of the cytb561 domain [1]. The CYBDOM cytb561 core domain encompasses TM1

through TM4, rather than TM2 through TM5 [1]. Single-domain cytb561s are oriented in a membrane

with their N- and C-termini exposed to the cytoplasm [2]. In contrast, CYBDOMs have an extracellular-

facing N-terminus [1], and, given that they have an even number of transmembrane regions, must also

have an extracellular-facing C-terminus. Single domain cytb561s and CYBDOMs have three core domain

loops that occur in the following order: cytoplasmic, non-cytoplasmic, cytoplasmic [1]. The function of

the DOMON domain(s) in CYBDOMs is unknown, but some members of the DOMON domain superfamily

bind sugar or heme, and at least one heme-binding DOMON domain is involved in interdomain electron

transfer [20]. Some CYBDOMs also have an extracellular, N-terminal adhesion module, such as the reeler

domain in SDR2 and *D. melanogaster* CG8399 [1]. The function of the reeler domain in CYBDOMs is not

known.

Our study aims to better understand the cytb561 family in insects, and we are particularly

interested in the possibility that insect cytb561 proteins play a role in cellular iron uptake. Iron

homeostasis in insects is poorly understood, and mechanisms of cellular iron uptake are still unknown

[21–23]. Most iron in biological systems is in the insoluble ferric form, but the known iron transporters

in animals are specific for ferrous ions; thus, for iron to enter an animal cell through a metal transporter,

it typically is first reduced to its soluble, ferrous state [24–26]. In mammals, the confirmed ferric

reductases are Dcytb and various STEAP proteins [8,27–29]. Insects do not have STEAP homologs [21];

therefore, members of the cytb561 family are the most promising ferric reductase candidates in insects.

Surprisingly little is known about insect cytb561 proteins. Only one, *D. melanogaster* Nemy, has

been studied in depth [4,30]. Nemy participates in a peptide amidation process in neuroendocrine

neurons and influences olfactory learning and memory [4,30]. The *D. melanogaster* CYBDOM, CG8399, is

reducible by ascorbate and has in vitro ferric reductase activity, but its biological function is unknown

[6,31]. Nemy and *D. melanogaster* CG1275, an uncharacterized cytb561, have been described as

possible ferric reductases [22,32].

The primary aims of this project were to 1) perform a phylogenetic analysis of cytb561 genes

from nine diverse insect species to identify orthologous groups, 2) compare their amino acid sequences

and predicted protein structures to those of better-studied human cytb561s to reveal sequence motifs

and structural elements that may be involved in the localization and biological functions of these

proteins, and 3) predict which of the orthologous groups may function as ferric reductases in insects. Of

the four cytb561 groups identified, the CG1275 and Nemy groups are similar to human Dcytb; the

CG8399 proteins are similar to human SDR2; and a subset of proteins in the fourth group is similar to

human TScytb. We found that CG1275 and Nemy proteins have conserved amino acid residues and

predicted protein structures that are consistent with ferric reductase activity. CG8399 proteins have

highly-conserved amino acid residues in cytoplasmic regions and the DOMON domain that may be

involved in substrate binding and ferric reductase activity. TScytb-like proteins have a "KXXXXKXH" motif

that may be involved in non-cytoplasmic substrate binding and potential ferric reductase activity.

Materials and methods

Identification of insect cytb561 protein sequences

For this study, we selected *D. melanogaster* as our model insect species, along with eight

additional insect species of interest: *Anopheles gambiae* (mosquito), *Acyrthosiphon pisum* (aphid), *Apis*

mellifera (bee), *Ctenocephalides felis* (flea), *Papilio xuthus* (butterfly), *Pediculus humanus corporis*

(louse), *Tribolium castaneum* (beetle), and *Zootermopsis nevadensis* (termite). These species were

chosen for their phylogenetic diversity and their association with high-quality genomic data.

We identified *D. melanogaster* cytb561 protein sequences using the FlyBase protein domain

search tool with the query "Cytochrome b561" [33]. Of the nine resulting hits, eight possessed the four

conserved heme-coordinating histidine residues characteristic of cytb561 family members: CG1275,

Nemy, CG8399, CG10165, CG13077, CG13078, CG10337, and CG3592. The remaining sequence, CG6479,

lacked these conserved histidines and was eliminated from our dataset.

To identify cytb561 sequences in the insect species of interest, we performed species-specific

BLAST-P searches using the National Center for Biotechnology Information (NCBI) non-redundant

database [34]. Eight *D. melanogaster* cytb561 amino acid sequences were used as queries: NP_728727.1

(CG1275), NP_725208.1 (Nemy), NP_611079.2 (CG8399), NP_609982.1 (CG10165), NP_609990.1

(CG13077), NP_609989.1 (CG13078), NP_609986.1 (CG10337), and NP_570039.1 (CG3592). Additional

insect cytb561 sequences were identified using the NCBI Conserved Domain Database (CDD) [35]. The

NCBI CDD search engine was populated with a combination of "insect species" and either "cyt_b561" or

"cytochrome_b_N," and a search was executed.

Isoforms and identical hits were removed from the dataset, resulting in one unique sequence

for each gene. Of the 46 non-*D. melanogaster* sequences discovered, 36 resulted from BLAST-P searches

and ten resulted from NCBI CDD queries. A sequence from *T. castaneum* (XP_008195477.1) was edited

using RNA-seq data to remove an apparently misplaced intron, resulting in the addition of 53 C-terminal

amino acids and a complete cytb651 core domain. The NCBI accession number and source of each

sequence is listed in S1 Table.

Three sequences with unique properties were observed: XP_314065.2 (*A. gambiae*),

XP_001950579.2 (*A. pisum*), and XP_001949276.1 (*A. pisum*). These three sequences were used as

separate queries to conduct BLAST-P searches against the NCBI non-redundant database for the class

Insecta to determine whether other insect species have proteins with similar sequences.

Defining homologous regions of protein sequences

Prior to performing phylogenetic and sequence analyses, we identified the homologous region

of each insect cytb561 protein. We used the HHpred server (Max Planck Institute Bioinformatics Toolkit

webserver), which is a tool for protein remote homology detection that generates a multiple sequence

alignment, predicts secondary structure, then creates a profile hidden Markov model that is compared

with a target database [36]. We evaluated each insect and outgroup sequence using the HHpred

PDB_mmCIF70_3_Mar server set to default parameters, including local alignment. This server

references the Protein Data Bank (PDB) of known protein structures for possible templates, then

generates results based on the probability percentage that the query shares a homologous region with

the template. An estimated probability percentage of > 95% indicates that homology is nearly certain

[36]. All insect and outgroup sequences were determined to share the highest probability percentage of

homology (> 99%) with regions of Dcytb (CYBRD1, NP_079119.3, PDB 5ZLG). The predicted homologous

region was not identical for all insect sequences; therefore, we identified the region that all insect

sequences have in common and defined this as the common homologous region, which corresponds to

Dcytb A44-T179. We used common homologous sequences for all analyses unless otherwise described.

The common homologous region of each sequence is listed in S1 Table.

To compare insect cytb561s with the six known human cytb561 proteins, we used HHpred and

the same methodology described above to define the common homologous regions of the following

human sequences: NP_079119.3 (Dcytb; CYBRD1), NP_705839.3 (Lcytb; CYB561A3), NP_001017916.1

(CGcytb; CYB561), NP_001278213.1 (TScytb; CYB561D2), NP_872386.1 (CYB561D1), and

NP_001347970.1 (SDR2; FRRS1).

HHPred analysis of the DOMON domains in insect CG8399 orthologs was performed. All insect

DOMONs were predicted to have > 99% predicted homology with regions of the cytochrome domain in

the *Phanerodontia chrysosporium* (fungus) cellobiose dehydrogenase (CDH) protein (PDB 1D7B) and the

DOMON domain in the human dopamine beta-hydroxylase (DBH) protein (PDB 4ZEL).

Multiple sequence alignments, sequence identities, and localization

Multiple sequence alignments were performed in Clustal Omega and visualized in JalView,

where we applied a custom color scheme and obtained consensus sequence logos and conservation

scores [37,38]. Accession numbers for insect sequences used in alignments are in S1 Table. Additional

sequences used in alignments include Dcytb and *Arabidopsis thaliana* cytochrome B561-1 (AtCytb561-B,

NP_198679.1, PDB 4O7G). To determine sequence identities, we used CLUSTAL Omega to generate

multiple sequence alignments for each insect sequence with all *D. melanogaster* and human cytb561

sequences, then determined pairwise sequence identities through utilization of the M-View multiple

alignment viewer [39].

Full-length insect cytb561 protein sequences were submitted to the SignalP-5.0 webserver for

detection of predicted signal peptides, and they were analyzed by the DEEPLOC-1.0 webserver for

prediction of subcellular protein localization [40,41]. Sequences were also investigated by eye to identify

putative late-endosomal/lysosomal targeting signals in the C-terminal portion of the protein [42].

Phylogenetic analysis

For our phylogenetic analysis, we selected an outgroup species, *Trichoplax adhaerens*, based on

two qualifying criteria: *T. adhaerens* belongs to an ancient animal lineage near the base of the metazoan

tree, and its cytb561 genes are homologous to insect cytb561 genes [43]. To identify an outgroup

sequence, we evaluated the cytb561 genes from *T. adhaerens* and selected the sequence with the

lowest identity to the *D. melanogaster* cytb561 sequences. The phylogenetic analysis was conducted

using the MEGA version X software [44]. First, all common homologous insect cytb561 sequences and

the outgroup cytb561 sequence were aligned in CLUSTAL Omega using the "Pearson/FASTA" output

parameter; the resulting alignment was uploaded to MEGA X, and a rooted phylogenetic tree was

created using the maximum likelihood (ML) method. To determine the optimal model for ML tree

construction, we ran the "Find Best DNA/Protein Models (ML)" tool in MEGA X, which resulted in

recommended parameters: the LG with freqs. (+F) substitution model, gamma distributed with invariant

sites (G+I) rates among sites, and five discrete gamma [45]. We then used the additional remaining

parameters: use all sites in cases of gaps and missing data, the Nearest-Neighbor-Interchange ML

heuristic method of tree inference, the default NJ/BioNJ initial tree for ML, no branch swap filters, seven

threads, and 1000 bootstrap replicates.

AlphaFold structure analysis

AlphaFold Monomer v2.0 pipeline structure predictions for *D. melanogaster* CG1275 isoform A

(AF-Q9I7U1-F1-model_v4.pdb), *D. melanogaster* Nemy isoform A (AF-Q7JR72-F1-model_v4.pdb), *D.*

melanogaster CG8399 isoform A (AF-Q8MSU3-F1-model_v4.pdb), *D. melanogster* CG13077 isoform B

(AF-M9PG89-F1-model_v4.pdb), and *A. gambiae* XP_3200673.4 (AF-Q7PZD0-F1-model_v4) were

downloaded from UniProt and visualized in ChimeraX [46–48]. We removed amino acids 1-30 from the

CG8399 structure prediction as they were predicted to be a signal peptide by SignalP 6.0 [49]. AlphaFold

structures for CG1275 and Nemy were individually overlayed with Dcytb (PDB 5ZLG) using the

MatchMaker feature in ChimeraX with default settings. The number of α-carbon pairs and their root-

mean-square deviation (RMSD) was generated by ChimeraX. CG1275 (amino acids Pro100-Ile326) and

Nemy (amino acids Cys52-Ser281) were displayed together with Dcytb (amino acids Tyr6-Pro230) to

generate figures. Individual AlphaFold models were associated with multiple sequence alignments of

same group proteins and colored in ChimeraX using AL2CO and a color gradient from cyan (low

conservation score) to magenta (high conservation score) [50]. The heme- and ascorbate-binding

residues were selected to calculate possible intermolecular hydrogen bonds using the H-bonds tool and

to predict possible direct residue interactions with the Contacts tool in ChimeraX using default settings

for overlap cutoff and H-bond allowance parameters. Potential hydrogen bonds were displayed in blue.

Coulombic surface coloring was used to visualize surface charges in the potential ion-binding pocket of

CG8399 [46].

Analysis of DOMON and TScytb sequence motifs

Additional sequence analyses were conducted to investigate DOMON domain sequences and a

"KXXXXKXH" motif in cytb561 proteins from diverse animal species. Tree of Life was referenced to

choose a set of species for these analyses: *Homo sapiens* (human), *Mus musculus* (mouse), *Passer*

montanus (sparrow), *Xenopus laevis* (frog), *Danio rerio* (zebrafish), *Octopus sinensis* (octopus), *Actinia*

tenebrosa (sea anemone), *Helobdella robusta* (leech), and *Acanthaster planci* (starfish) [51]. To identify

DOMON-containing sequences, the *D. melanogaster* CG8399 sequence (NP_611079.2) was used as a

BLAST-P query to search NCBI non-redundant databases; to identify "KXXXXKXH"-containing sequences,

the query was *A. gambiae* sequence XP_320673.4. CLUSTAL Omega was then used to generate

sequence alignments that allowed us to evaluate amino acid conservation of the DOMON domains and

"KXXXXKXH" motifs and to determine the placement of the "KXXXXKXH" motif within the protein [39].

Results and Discussion

Phylogenetic analysis of insect cytb561 proteins

Insects have four distinct cytb561 groups

While a limited number of insect cytb561 sequences have been included in broader

phylogenetic studies [3,19], we were interested in classifying the cytb561 family across diverse insect

species. We identified 54 genes that encode cytb561 family members from nine insect species in eight

orders: Blattodea (*Z. nevadensis*), Coleoptera (*T. castaneum*), Diptera (*A. gambiae* and *D. melanogaster*),

Hemiptera (*A. pisum*), Hymenoptera (*A. mellifera*), Lepidoptera (*P. xuthus*), Psocodea (*P. humanus*

corporis), and Siphonaptera (*C. felis*) (S1 Table). All investigated species have between three *(P.*

humanus corporis) and nine (*P. xuthus*) cytb561 genes (S1 Table).

Using the HHpred PDB server for protein remote homology detection, we defined a homologous

region (core domain) shared by all of the insect cytb561s [36] (S1 Table). The ~130 amino acid long core

domain sequences were then used to generate a multiple sequence alignment (S1 Fig). From this

multiple sequence alignment, we generated a phylogenetic tree and found strong support for four

cytb561 groups: three orthologous groups and a fourth group that lacks one-to-one orthologs (Fig 1).

For simplicity, we will refer to all insect sequences that cluster in the three orthologous groups

according to the representative *D. melanogaster* protein name: CG1275, Nemy, or CG8399. The

remaining 25 sequences will be referred to as Group 4.

Fig 1. Phylogenetic tree of insect cytb561 proteins. A phylogenetic analysis of cytb561s from nine

species of insects (*A. gambiae [Ag]*, *A. mellifera [Am]*, *A. pisum [Ap]*, *C. felis [Cf]*, *D. melanogaster [Dm]*,

T. castaneum [Tc], *P. humanus corporis [Ph]*, *P. xuthus [Px]*, and *Z. nevadensis [Zn])* was performed with

the maximum likelihood method. An outgroup sequence from *T. adhaerens (Ta)* was used for rooting

the tree. Bootstrap values greater than 70% are shown. Three orthologous groups (with one-to-one

orthologs in most or all species) were identified: CG1275, Nemy, and CG8399. The remaining genes

formed a fourth group that has no one-to-one orthologs but includes many species-specific gene

expansions. The bootstrap values associated with the four clades are marked with red circles. A branch

length scale bar indicates one substitution per site. Of the Group 4 sequences, seven are more similar to

human TScytb than any *D. melanogaster* cytb561s and are referred to as subgroup 4A, whereas the

remaining Group 4 proteins belong to subgroup 4B. (Note that the 4A and 4B subgroups are based on

sequence comparisons rather than phylogenetic relationship.)

CG1275 and Nemy

The CG1275 and Nemy groups are closely related, as evidenced by bootstrap support of a

CG1275-Nemy cluster and the short branch lengths for the CG1275 and Nemy groups (Fig 1). CG1275

proteins were identified for eight of the nine insect species (Fig 1; S1 Table). We did not identify a

CG1275 ortholog in the lepidopteran insect *P. xuthus*, and we were unable to find a CG1275 sequence in

any other species from the order Lepidoptera (butterflies and moths). Of the species analyzed, *A. pisum*

was the only insect with a CG1275 gene expansion (Fig 1; S1 Table). We identified Nemy proteins in

eight of the species (Fig 1; S1 Table), but we did not find a Nemy sequence from *A. mellifera* or any other

species in the order Hymenoptera (bees, wasps, ants, and similar insects). A previously published

phylogenetic analysis of eukaryotic cytb561s suggested that CG1275 and Nemy from *D. melanogaster*

and *A. gambiae* may be orthologous to human Dcytb, CGcytb, and Lcytb, but a lack of statistical support

for the clade makes the phylogenetic relationship of CG1275 and Nemy with Dcytb, CGcytb, and Lcytb

uncertain [19].

CG8399

CG8399 is the only orthologous group with sequences from all nine insect species. This cluster

includes ten proteins, nine of which contain multiple domains – reeler, DOMON(s), and cytb561 –

consistent with some members of the CYBDOM protein family (Fig 1; S1 Table) [1]. *A. gambiae* was the

only species with a second CG8399-like protein (XP_314065.2); this sequence differs from other

CG8399s in that it is a single-domain cytb561 lacking both reeler and DOMON domains. A previously

published phylogenetic analysis of eukaryotic cytb561s demonstrated that *D. melanogaster* and *A.*

gambiae CG8399 proteins are orthologous to SDR2 in humans, other animals, and plants [19].

Group 4

276   The remaining 25 insect genes populate a large clade (Group 4) with a bootstrap value of 92 (Fig

277   1). Group 4 has multiple species-specific clusters with significant bootstrap values, suggesting that

278   Group 4 genes were more likely than other insect cytb561 genes to undergo species-specific expansions

279   (Fig 1). We did not identify any Group 4 genes in *P. humanus corporis* or *A. pisum* (Fig 1).

Sequence identity between human and insect cytb561s

Human cytb561s have been more widely studied than insect cytb561s, and human Dcytb is one

of only two cytb561 structures that have been solved [18]. We were interested in determining the

sequence similarities between the 54 insect cyb561s and the six human cytb561s (Dcytb, CGcytb, Lcytb,

SDR2, TScytb, and CYB561D1) to learn which of the better-characterized human cytb561s is most similar

toeach of the insect cytb561s. For this analysis, we identified the common homologous region of each

insect and human cytb561 sequence, then determined their pairwise identities. CG1275 and Nemy

proteins had highest sequence identity with Dcytb, CGcytb, or Lcytb (32.4%-49.3%), and CG8399s had

highest sequence identity with SDR2 (36.5%-43.4%) (Table 1). These results suggest that CG1275 and

Nemy proteins may have physiological similarities with Dcytb, CGcytb, and/or Lcytb, and are consistent

with the finding that CG8399s are orthologous to SDR2 [19]. Of the 25 Group 4 cytb561s, 23 had highest

sequence identity with TScytb (21; 18.8%-33.1%) or CYB561D1 (2; 22.0%-25.7%) (Table 1). Seven Group

4 proteins are distinct from all other insect cytb561s in that they had higher sequence identity with a

human cytb561 (TScytb) than with any of the *D. melanogaster* cytb561s; we will refer to this subset of

proteins as Group 4A and the remaining Group 4 proteins as Group 4B (Table 1; Fig 1; S2 Table). A

previous phylogenetic study found that the *A. gambiae* gene in Group 4A clustered with human and

other TScytbs rather than other insect-specific sequences, but there was a lack of bootstrap support for

this cluster; thus, the phylogenetic relationship between Group 4A genes and TScytb is unclear [19].

Table 1. Sequence identity between insect and human cytb561s.

			Human Sequences					
Species	Group	Accession Number	CYBRD1 (Dcytb)	CYB561A3 (Lcytb)	CYB561 (CGcytb)	CYB561D2 (TScytb)	CYB561D1	FRRS1 (SDR2)
Dm	CG1275	NP_728727.1	**45.7[1]**	40.6	39.3	21.7	20.3	16.7
Ag	CG1275	XP_315519.4	43.2	43.2	**43.4**	23.2	20.4	18.6
Ap	CG1275	NP_001155374.1	32.6	34.1	**40.7**	17.9	23.6	15.3
Ap	CG1275	NP_001280323.1	31.7	29.6	**32.4**	15.2	18.6	15.5
Ap	CG1275	XP_001950854.1	37	37	**41.5**	22	25.5	18.1
Ap	CG1275	XP_003246890.1	30.4	31.2	**33.3**	17.7	20.6	11.7
Am	CG1275	XP_006572086.1	**49.3**	44.9	43	22	20.6	20.1
Cf	CG1275	XP_026462102.1	39.9	38.5	**42.1**	21.2	21.2	18.9
Ph	CG1275	XP_002426701.1	**46.8**	43.9	44.1	22.5	21.1	20.4
Tc	CG1275	XP_008194670.1	44.9	42.8	**45.9**	24.5	20.3	16.1
Zn	CG1275	XP_021935166.1	42	41.3	**44.4**	21.3	20.6	20.8
Dm	Nemy	NP_725208.1	35	**35.7**	32.1	19.7	21.8	16.4

Ag	Nemy	XP_314126.2	38.6	**39.3**	35.8	19	23.2	19.2
Ap	Nemy	XP_001949276.1	33.6	32.9	**36.6**	23.1	27.3	17
Cf	Nemy	XP_026473332.1	37.3	39.4	**39.6**	21.5	29.9	16.9
Px	Nemy	NP_001298968.1	37.1	37.9	**38.7**	23.2	19	16.4
Ph	Nemy	XP_002430226.1	35.2	**36.6**	35.3	20.8	26.4	15.5
Tc	Nemy	XP_008198104.1	36.6	**39.4**	36.7	22.9	29.2	14.9
Zn	Nemy	XP_021939496.1	**39.2**	38.5	38.6	22.8	28.3	14.6
Dm	CG8399	NP_611079.2	18.4	17	17.1	13.9	13.9	**37**
Ag	CG8399	XP_314065.2	18.3	20.4	19.9	16.1	15.4	**36.8**
Ag	CG8399	XP_314066.4	18.3	19	19.1	11	15.2	**41.2**
Ap	CG8399	XP_001950579.2	20	16.6	18.1	10.1	14.2	**36.5**
Am	CG8399	XP_396579.3	15.6	21.3	15.7	9.7	11.1	**41.2**
Cf	CG8399	XP_026481553.1	19	17.6	17	11.7	16.6	**43.4**
Px	CG8399	XP_013164083.1	15.6	19.1	18.6	16	17.4	**39.4**
Ph	CG8399	XP_002423127.1	16.4	19.9	17.2	9.4	14.1	**40.1**
Tc	CG8399	XP_015836986.1	17.6	17.6	15.6	11.7	15.2	**38.2**
Zn	CG8399	XP_021919699.1	15.5	18.3	17	12.4	15.2	**41.9**
Dm	Group 4B	NP_609982.1	17.1	17.1	18.7	**24.8**	22	10.6
Dm	Group 4B	NP_609990.1	14.3	16.4	15.8	**22.7**	20.6	14.2
Dm	Group 4B	NP_609989.1	18.4	14.2	16.4	**19.7**	16.9	14.2
Dm	Group 4B	NP_609986.1	17.3	20.9	**23.2**	21.1	18.3	11.2
Dm	Group 4B	NP_570039.1	18.8	22.2	19.6	**24.5**	21.8	11.4
Ag	Group 4A	XP_320673.4[2]	21.5	20.1	23.1	**33.1**	31.7	19.4
Ag	Group 4B	XP_001238089.2	16.4	14.3	12.9	25	**25.7**	12.6
Ag	Group 4B	XP_552919.3	18.6	19.3	18	**27.3**	21.6	16.1
Am	Group 4A	XP_001122176.1[2]	18.6	17.9	13.7	**27.9**	19.3	17.5
Am	Group 4B	XP_003249671.1	20.4	21.1	19.1	**25.2**	20.9	16.2
Cf	Group 4A	XP_026469480.1[2]	22.3	20.1	19.6	**31.7**	27.3	15.5
Cf	Group 4A	XP_026469242.1[2]	22.3	18.7	15.2	**28.8**	26.6	18.2
Px	Group 4A	XP_013174961.1[2]	20	16.4	23	**33.1**	26.6	14.7
Px	Group 4B	XP_013172799.1	15.8	12.9	16.7	**26.2**	20.6	10
Px	Group 4B	XP_013172813.1	17.3	15.1	16.8	**25.5**	22	12.9
Px	Group 4B	XP_013162691.1	17.1	19.3	17.3	**22**	20.6	13.4
Px	Group 4B	XP_013162686.1	14.3	13.6	12.2	**19.9**	16.3	12.8

Px	Group 4B	XP_013172802.1	17.4	15.2	15.3	**18.8**	18.1	12
Px	Group 4B	XP_013172804.1	20.7	**22.1**	18	19.9	21.3	14.9
Tc	Group 4A	XP_008201603.1[2]	23	20.1	19.6	**33.1**	23.7	18.3
Tc	Group 4B	XP_008195477.1[3]	16.3	19.1	16.4	**26.2**	22.7	16
Tc	Group 4B	XP_015837014.1	20.7	20	18	**26.8**	21.8	11.2
Tc	Group 4B	XP_974632.1	17.9	18.6	16.5	21.3	**22**	14.8
Tc	Group 4B	XP_974652.1	20.9	18.7	15.2	**26.6**	24.5	14.2
Zn	Group 4A	XP_021927739.1[2]	20.9	18	18.8	**32.4**	30.9	14.8

[1]Sequence identity (as a percent) between the homologous region of each insect cytb561 and each human cytb561. The highest identity in each row is shown with a shaded cell and in bold text. Group 4A sequences are shaded in pale orange and all other sequences are shaded in pale blue.
[2]Group 4A insect sequences with higher identity to a human cytb561 than a *D. melanogaster* cytb561.
[3]Edited to correct a gene prediction error.

Analysis of insect cytb561 domains

Locations of conserved amino acid residues

We used amino acid sequence alignments to identify conserved amino acid residues in insect cytb561 domains. To facilitate comparisons of residues from diverse sequences, we denote residues as the number of positions they are upstream (-) or downstream (+) of one of the four conserved heme-coordinating histidine residues; for example, the residue immediately upstream of H1 is referred to as H1-1. To identify amino acid residues that are conserved in all four insect cytb561 groups, we aligned the ~130 amino acid core domain sequences of all 54 insect species and found that only the heme-coordinating histidines, H1-H4, were completely conserved (S1 Fig). Note that for the CG1275, Nemy, and Group 4 proteins, H1-H4 are in TM2-TM5, whereas in the CG8399 proteins, H1-H4 are in TM1-TM4. In addition to the four positions containing conserved histidines, 11 positions have highly-conserved residues (with Jalview conservation scores of 8 or 9), including seven with small residues (Gly, Ala, and Ser), and four with larger hydrophobic residues (S1 Fig). Five of the seven small residue positions were previously noted [3], including two that are present in 106 eukaryotic cytb561s (H1+7, H4+4) [19]. To

identify amino acid conservation within specific insect cytb561 groups, we separated the core domain

alignment into five group-specific alignments and then used Jalview to generate group-specific

consensus logos (Fig 2) [38]. These consensus logos reveal patterns of conservation within groups and

allow residues at specific locations to be compared across groups.

Fig 2. Group consensus logos for the cytb561 core domain. Group-specific alignments were used to

create group consensus sequences in Jalview. The four conserved histidines are labeled H1-H4 with

yellow highlighting for the non-cytoplasmic histidines and blue highlighting for the cytoplasmic

histidines. The positions of amino acid residues of interest are labeled relative to the nearest conserved

histidine. If the corresponding residue in Dcytb was predicted to contact a heme or ascorbate, that

position is indicated by a symbol: non-cytoplasmic heme (square), non-cytoplasmic ascorbate (upside-

down triangle), cytoplasmic heme (red diamond), or cytoplasmic ascorbate (red triangle). The Jalview

consensus rows show the most common residue at that position listed below the logo; a + is used where

there are two or more residues with the highest percentage and a – is used to show gaps.

To analyze the conserved amino acids in the context of the protein structure, we used ChimeraX

to overlay a protein alignment of the CG1275, Nemy or CG8399 groups with its representative *D.*

melanogaster AlphaFold protein model (Fig 3). For Group 4B proteins, we selected the *D. melanogaster*

CG13077 AlphaFold model (Fig 3). Since *D. melanogaster* does not have a Group 4A protein, we used the

A. gambiae Group 4A (XP_320673.4) AlphaFold model (Fig 3). The degree of conservation is shown by

gradient coloring. By analyzing conservation information from the structural models as well as the

group-specific consensus logos, we determined that the insect cytb561 proteins, with the exception of

the Group 4B proteins, have relatively high group sequence conservation in the regions near non-

cytoplasmic heme-coordinating H1 and H3 (Figs 2 and 3). Additionally, CG8399 proteins have relatively

high conservation in the regions preceding the cytoplasmic heme-coordinating H2 and H4 (Figs 2 and 3).

Fig 3. Insect cytb561 models with sequence alignment overlay demonstrate residue conservation.

AlphaFold models of mature proteins are shown. Sequence conservation was determined by ChimeraX

using AL2CO conservation scores. The model is gradient colored with high conservation (+1.5 score) in

magenta and low conservation (-1.5 score) in cyan; sequence gaps are shown in tan. Conserved

histidines H1-H4 (not all visible in all panels) are shown in stick form with nitrogen in blue. The models

are oriented with the cytoplasmic side of the protein at the bottom. Note that the amino- and carboxyl-

termini for all models except the CG8399 model are near the bottom of the panels. Locations of H1-4

are indicated. Scale bars depict 10 Å. A) *D. melanogaster* CG1275 (Lys95-Glu340) aligned with the

CG1275 sequences. B) *D. melanogaster* Nemy (Lys40-Ile290) aligned with the Nemy sequences. C) *D.*

melanogaster CG8399 (Ser402-Gly645) aligned with the multi-domain CG8399 sequences. The reeler

and DOMON domains are omitted. D) *A. gambiae* XP_320673.4 (all residues) aligned with the Group 4A

sequences. E) *D. melanogaster* CG13077 (all residues) aligned with the Group 4B sequences.

General comparisons of insect cytb561 proteins with Dcytb and

AtCytb561-B

This is the first large scale analysis of cytb561 proteins since X-ray crystallography studies

identified heme- and ascorbate-binding residues in human Dcytb and *A. thaliana* AtCytb561-B, and a

metal ion-coordinating residue in Dcytb [13,18]. We wanted to leverage this structural information to

predict whether conserved residues in insect cytb561s might participate in heme, ascorbate, and/or

iron-binding. Amino acid sequence alignments and structural models were used to compare insect

cytb561 proteins with Dcytb and Atcyb-B. Initial comparisons were restricted to CG1275 and Nemy

proteins because of their high similarity to Dcytb and AtCytb561-B; once key residues were identified in

CG1275 and Nemy, we used sequence alignments to identify the corresponding residues in the CG8399

and Group 4 proteins. References to specific residue numbers are based on Dcytb unless otherwise

noted.

We first created an alignment of Dcytb, AtCytb561-B, CG1275, and Nemy sequences (Fig 4).

From this alignment, which extends beyond the core domain, we identified 23 completely conserved

residues and an additional 23 residues that are highly conserved across Dcytb, AtCytb561-B, CG1275,

and Nemy sequences (Fig 4). To identify any previously unspecified Dcytb residues that are likely to be

important for heme or ascorbate binding, we used ChimeraX to determine predicted protein contacts

with each heme and ascorbate group in the Dcytb ascorbate-bound structure (PDB 5ZLG) (Fig 4; S3

Table). (Note that Dcytb was selected for this analysis rather than AtCytb561-B for multiple reasons:

insects are more closely related to humans than plants, Dcytb is a ferric reductase, and the Dcytb crystal

structure contains a metal ion.)

Fig 4. Alignment of Dcytb, AtCytb561-B, Nemy, and CG1275 protein sequences. The two cytb561

sequences with known structure, human Dcytb (PDB 5ZLG) and *A. thaliana* AtCytb561-B (PDB 4O7G),

were aligned with CG1275 and Nemy sequences from *A. gambiae (Ag)*, *A. mellifera (Am)*, *A. pisum (Ap)*,

C. felis (Cf), *D. melanogaster (Dm)*, *T. castaneum (Tc)*, *P. humanus corporis (Ph)*, *P. xuthus (Px)*, and *Z.

nevadensis (Zn)*. (Note that only the *A. pisum* CG1275 sequence most similar to Dcytb was included.)

Alpha helices from crystal structures (Dcytb, AtCytb561-B) or predicted in AlphaFold models (Nemy,

CG1275) are shown in boxes: helix 1 carboxyl end (gray), helix 2 (red), helix 3 (yellow), helix 4 (green),

and helix 5 (blue). Conserved residues are noted under the alignment (* indicates completely conserved,

the four strictly conserved histidines are labeled with their number rather than *, and ^ indicates highly

conserved with a Jalview conservation score of 8 or 9). Predicted Dcytb heme-contacting residues are

labeled with a square (non-cytoplasmic heme) or diamond (cytoplasmic heme), and ascorbate-

contacting residues are labeled with an upside-down triangle (non-cytoplasmic ascorbate) or triangle

(cytoplasmic ascorbate). A blue vertical line indicates the position of 51 amino acids found only in *A.*

pisum XP_001949276.1, which were omitted from the figure. Coloring by amino acid residue: basic

residues in blue (Arg) or cyan (Lys); acidic residues (Asp and Glu) in dark pink; amide residues (Asn and

Gln) in light pink; hydroxylic residues (Ser and Thr) in pale purple; aromatic residues in light peach (Phe

and Tyr) or dark peach (Trp); sulfur-containing residues in dark green (Cys) or light green (Met); His in

orange; Pro in yellow; Ile, Leu, and Val in pale yellow; and Ala and Gly in white.

Next, we compared structural models of *D. melanogaster* CG1275 and Nemy with the solved

structure of Dcytb. To investigate, we overlayed AlphaFold models for *D. melanogaster* CG1275 and

Nemy with the ascorbate-bound structure of human Dcytb. The AlphaFold models have very high

confidence for the majority of residues in the six transmembrane helices and their connecting loops (S4

Table), and the alpha carbons of the two insect structures aligned well over most of the Dcytb sequence

(Fig 5) [47,48,52]. Although the AlphaFold cytb561 models do not include the two bound heme groups

and the two bound ascorbates, overlaying the AlphaFold structures with Dcytb showed how these

molecules may fit within the CG1275 and Nemy structures (Fig 5) [18].

Fig 5. *D. melanogaster* CG1275 and Nemy models overlayed on Dcytb heme- and ascorbate-bound

structure. AlphaFold structures of *D. melanogaster* CG1275 and Nemy were overlayed with Dcytb (PDB

5ZLG) using Chimera Matchmaker. Transmembrane helices are labeled (TM1-TM6), with Dcytb in tan,

CG1275 in pale pink, and Nemy in pale blue. The four conserved histidines are shown in stick form; in

the Dcytb structure, H1 is in red, H2 in yellow, H3 in green, and H4 in dark blue. Ascorbate (ASC) in black

stick, heme in tan stick (with iron shown as orange spheres), and zinc (Zn^{2+}) ion coordination in purple

are from the 5ZLG structure. Stick side chains, ascorbate, and heme have oxygen in red and nitrogen in

blue. The alpha carbon RMSD for CG1275 with 5ZLG is 0.679 Å for the 208 pruned atom pairs and 1.458

Å across all 222 pairs. The alpha carbon RMSD for Nemy with 5ZLG is 0.990 Å for the 188 pruned atom

pairs and 2.657 Å across all 222 pairs. A) Overlay of *D. melanogaster* CG1275 and Nemy with Dcytb. B)

Non-cytoplasmic ascorbate binding, zinc coordination, and heme coordination with H1 (red) and H3

(green) shown in stick form. C) Cytoplasmic ascorbate binding and heme coordination with H2 (yellow)

and H4 (dark blue) shown in stick form.

Conserved insect cytb561 residues likely to be involved in heme

positioning

We were interested in exploring whether residues involved in heme placement in Dcytb and

AtCytb561-B have similar roles in insect cytb561 proteins. Our Dcytb protein contact analysis identified

47 residues that make at least one predicted contact with a heme group (S3 Table), including 32

residues that are completely or well conserved in all CG1275 and Nemy sequences, as well as in

AtCytb561-B (Fig 4). Many of these 32 residues are predicted to make contacts with a heme group in *D.*

melanogaster CG1275 and Nemy protein models overlayed with Dcytb (S3 Table). In particular, the H1-

H4 side chains are positioned very similarly to the heme-coordinating H1-H4 side chains in the Dcytb

structure (Figs 5B and C). Amino acid residues that were predicted to interact with a heme group or

participate in heme positioning, as well as their degree of conservation in the four insect cytb561

groups, are described below.

Cytoplasmic heme

The cytoplasmic heme group in Dcytb directly interacts with three amino acid residues: H2, H4,

and Arg70 (H1+20) [18]. Arg70 forms a salt bridge with the heme A-propionate through its epsilon

nitrogen [18]; a similar salt bridge is formed by a lysine at H1+20 in AtCytb561-B (Lys71) [13]. An

arginine at H1+20 is completely conserved in all CG1275, Nemy, and CG8399 sequences (Fig 2) and likely

interacts with the cytoplasmic heme A-propionate (Fig 6A). The residue at H1+20 tends to be a serine or

threonine in Group 4A proteins (Fig 2), and 22 out of 25 Group 4B sequences also have a polar residue at

H1+20 (S1 Fig). Serine is a much smaller residue than arginine, but in Dcytb the gamma carbon of Arg70

is only ~3.2 Å from the heme A propionate; therefore, a serine or threonine residue could potentially

form a hydrogen bond with a heme carboxylate group if properly positioned. Outside of the Dcytb core

domain, at the end of helix 6, Lys225 forms two potential hydrogen bonds with the cytoplasmic heme D

propionate (Fig 6A; S3 Table). This helix 6 lysine is conserved in CG1275 sequences and is a basic residue

in 5 out of 7 Nemy sequences, but it is replaced with a proline in *D. melanogaster* and *A. gambiae* Nemy.

(Comparisons of Lys225 with CG8399 and Group 4 sequences were not attempted because helix 6 is

outside of their homologous regions.)

Fig 6. Heme binding by Dcytb and *D. melanogaster* CG1275 and Nemy. AlphaFold structures of *D.*

melanogaster CG1275 and Nemy were overlayed with Dcytb (5ZLG) using Chimera Matchmaker. Dcytb

loops are in tan and helices are shown in red (helix 2), yellow (helix 3), green (helix 4), dark blue (helix 5),

and purple (helix 6). CG1275 is in pale pink, and Nemy is in pale blue. Ascorbate (ASC) in black stick,

heme in tan stick (with iron shown as orange spheres), and zinc (Zn^{2+}) ion coordination in purple are

from the 5ZLG structure. Select side chains are shown in stick form and are labeled by position relative

to conserved heme-coordinating histidines with Dcytb amino acid numbering in parentheses. Key

glycines are shown in black ribbon. Stick side chains, ascorbate, and heme have oxygen in red and

nitrogen in blue. Hydrogen bonds shown as cyan dashed lines, including to the A and D heme-

propionates (labeled as in [18]). Water molecules shown as red spheres. A) Cytoplasmic heme region. B)

Non-cytoplasmic heme region. Hydrogen bonds predicted by ChimeraX or observed in Dcytb structure

(Asn115 to D heme-propionate; Glu180 to A heme-propionate, [18]).

In the Dcytb and AtCytb561-B structures, we observed interactions between H4 and the

glutamine at H3+14 (Gln134). Gln134 forms a predicted hydrogen bond with the H4 backbone carbonyl

and contributes to a network of hydrogen bonds linked by a water molecule to the H4 imidazole ring

positioned on the opposite side of the heme iron coordination site (Fig 6A). An H3+14 glutamine residue

is conserved in all CG1275, Nemy, and CG8399 sequences, suggesting a role in the positioning of H4 in

these three protein groups. In Group 4 sequences, a serine (16 out of 25) or threonine (4 out of 25) is

often present at H3+14 (S1 Fig) that could both potentially interact with H4, perhaps through hydrogen

bonding mediated by water. Thus, within CG1275, Nemy, and CG8399 proteins, a glutamine at H3+14

may help position H4 to coordinate the heme, while a salt bridge may form between the cytoplasmic

heme A propionate and the arginine at H1+20. In Group 4 proteins, polar residues at H1+20 and H3+14

could conceivably have similar heme positioning roles.

We found that two of the highly-conserved small residue positions in insect cytb561s are

predicted to form cytoplasmic heme contacts in Dcytb: H3+18 (Gly138) and H4+4 (Gly163) (S3 Table). A

glycine at H4+4 is conserved across all insect groups, in addition to more diverse cytb561s [3,19], while a

glycine at H3+18 is conserved in all insect groups except CG8399, where an alanine substitution occurs

(Fig 2). It is likely that small residues at H3+18 and H4+4 are important for proper cytoplasmic heme

positioning within the protein's central cavity in diverse cytb561s (Fig 6A).

Non-cytoplasmic heme

We performed a similar analysis of non-cytoplasmic heme positioning. In Dcytb and AtCytb561-

B, two residues form hydrogen bonds with the non-cytoplasmic heme A-propionate group: a serine at

H3-2 (Ser118) and a glutamate at H4+21 (Glu180) (Fig 6B)[13,18]. These residues are well conserved in

CG1275 and Nemy sequences and form predicted contacts with the heme group in the *D. melanogaster*

CG1275 and Nemy models (Fig 4; S3 Table). As mentioned above, it is notable that the most highly-

conserved regions among CG1275 and Nemy proteins are the sequences surrounding H1 and H3, which

coordinate the non-cytoplasmic heme group (Fig 2). While CG8399 and Group 4 proteins share less

conservation in these regions, a serine or threonine at H3-2 is well conserved across Group 4 proteins,

and CG8399 sequences most often contain an asparagine (Fig 2; S1 Fig); these polar residues could

potentially hydrogen bond with the non-cytoplasmic heme A-propionate (H4+21 is slightly beyond the

conserved core cytb561 domain and was therefore omitted from our analysis of CG8399 and Group 4

proteins).

In Dcytb and AtCytb561-B, an asparagine at H3-5 forms a hydrogen bond with the non-

cytoplasmic heme D-propionate (Fig 6B) [13,18]. This position is highly conserved in CG1275 and Nemy

proteins (Fig 4); two Nemy sequences have a histidine, which also could form a potential interaction

with a heme propionate. All Group 4A and most Group 4B proteins also have a histidine at H3-5 (Fig 2;

S1 Fig). In contrast to the other groups, there is no clear conservation at H3-5 in CG8399 proteins, which

is part of the core domain non-cytoplasmic loop (Fig 2; S1 Fig).

Additional AtCytb561-B residues that form hydrogen bonds with the non-cytoplasmic heme D-

propionate are the side chain of H2+22 (His106) and the amide nitrogen of H3-4 (Phe114) [13].

Interestingly, in Dcytb the histidine at H2+22 is conserved but was reported to coordinate with a non-

cytoplasmic zinc ion (Fig 6B, also discussed in more detail below) [18]. All but two CG1275 and Nemy

sequences have a histidine at H2+22, whereas this position is most commonly a lysine in Group 4

proteins or a hydrophobic residue in CG8399 proteins (Fig. 2; S1 Fig). Dcytb has a methionine (Met116)

at H3-4 whose amide nitrogen is also positioned to hydrogen bond with the non-cytoplasmic heme D-

propionate, similarly to Phe114 in AtCytb561-B (Fig 6B). In CG1275 and Nemy sequences, a methionine,

phenylalanine, or leucine is found at H3-4 and a phenylalanine is also common in Group 4 proteins (18

out of 25) (Figs 2 and 4). In contrast, CG8399s have no notable conservation at this position (S1 Fig). The

hydrophobic side chain at H3-4 in Dcytb (Met116) appears to anchor the non-cytoplasmic loop between

helices 3 and 4 in the membrane, and its backbone carbonyl is predicted to form a hydrogen bond with

the sidechain of the tryptophan (Trp122) at H3+2, providing added stability to the loop region between

helices 3 and 4 (Fig 6B). A tryptophan at H3+2 is entirely conserved in CG1275 and Nemy proteins (Fig 4),

suggesting a similar positioning of the non-cytoplasmic heme and loop between helices 3 and 4. The

tryptophan at H3+2 is not conserved in CG8399 or in most Group 4 proteins (Fig 2).

Glycine residues at positions H3+4 (Gly124) and H4+18 (Gly177) are predicted to make non-

cytoplasmic heme contacts in the Dcytb, CG1275, and Nemy structures (Figs 4 and 6B; S3 Table) These

residues were previously identified as highly conserved across diverse cytb561s [3]. The H3+4 glycine is

conserved in all CG1275, Nemy, and CG8399 sequences, and is highly conserved in Group 4 proteins (21

out of 25) (Fig 2). A glycine at H4+18 is conserved in all CG1275 and Nemy sequences and most Group 4

proteins (19 of 25) also have a glycine at H4+18, whereas CG8399s have an alanine substitution (Fig 2;

S1 Fig). Therefore, it is likely that small residues at H3+4 and H4+18 (non-cytoplasmic side), as well as

H4+4 and H3+18 (cytoplasmic side, described above) are important for positioning the bulky heme

groups in the central cavity of the protein across a range of cytb561s.

Heme positioning summary

Overall, heme positioning appears to be similar in Dcytb, AtCytb561-B, CG1275, and Nemy

proteins, whereas heme positioning in CG8399 and Group 4 proteins is less similar. Non-cytoplasmic

heme positioning is most dissimilar in CG8399 proteins, and cytoplasmic heme positioning is most

dissimilar in Group 4 proteins.

Ascorbate binding

Dcytb and AtCytb561-B were found to bind to ascorbate (or the closely related

monodehydroascorbate) in the cytoplasm through the interactions of basic residues with some of the

ascorbate oxygens [13,18]. On the non-cytoplasmic side, phenylalanine was implicated in stacking with

the ascorbate ring [13,18]. We located these Dcytb and AtCytb561-B residues in relation to the

conserved heme-coordinating histidines and examined our insect sequences for conserved residues,

especially basic and aromatic residues, at these locations. We also performed ChimeraX contact analysis

with the Dcytb structure and overlayed CG1275 and Nemy structures to identify other amino acids

proximate to ascorbate to identify similarities and differences in how ascorbate might bind.

Potential cytoplasmic ascorbate binding residues

The Dcytb and AtCytb561-B crystal structures clearly identified three basic residues that bind

cytoplasmic ascorbate: H2-7 (Lys79), H2-3 (Lys83), and H4-7 (Arg152) [13,18]. Additionally, ChimeraX

analysis predicted that Dcytb has ascorbate contacts at H3+22 (Phe142) and H4-3 (Met156) (Fig 7A; S3

Table). These five residues are also predicted to have cytoplasmic ascorbate contacts in *D. melanogaster*

CG1275 (S3 Table), and most CG1275 proteins have these conserved residues except for some variation

at H4-3 (Fig 4). In Nemy sequences, a lysine at H2-3, an arginine at H4-7, and a phenylalanine at H3+22

are entirely conserved, whereas either a valine or threonine is present at H4-3 (Fig 4); *D. melanogaster*

Nemy is predicted to have ascorbate contacts at these four positions (S3 Table). More variation is seen

at Nemy H2-7, which is most often an arginine (4 out of 8) and in one sequence is a lysine (Fig 4).

However, H2-7 is not a basic residue in the *D. melanogaster*, *A. gambiae*, and *A. pisum* Nemy sequences

(Fig 4). The *D. melanogaster* Nemy model suggests a shift in the placement of the lysine at H2-3 (Fig 7A),

which may compensate for the lack of a basic residue at H2-7. Additionally, the Nemy model predicts

that a serine (Ser205) at H4-6 hydrogen bonds with ascorbate, suggesting a unique functional role for

this serine in *D. melanogaster* Nemy and in *A. gambiae* Nemy, which also has a serine at this position

(Fig 7A).

Fig 7. Ascorbate binding in Dcytb, CG1275, and Nemy. AlphaFold structures of *D. melanogaster* CG1275

and Nemy were overlayed with Dcytb (5ZLG) using Chimera Matchmaker. Dcytb loops are in tan and

helices are shown in yellow (helix 3), green (helix 4), and dark blue (helix 5). CG1275 is in pale pink, and

Nemy is in pale blue. Ascorbate (ASC) in black stick, heme in tan stick (with iron shown as orange

spheres), and metal ion coordination in purple are from the 5ZLG structure. Select side chains are

shown in stick form and are labeled by position relative to conserved heme-coordinating histidines with

Dcytb amino acid numbering in parentheses (unless otherwise noted as Nemy numbering). Stick side

chains, ascorbate, and heme have oxygen in red and nitrogen in blue. Hydrogen bonds shown as cyan

dashed lines. A) Cytoplasmic ascorbate region. Ascorbate oxygens involved in bonding are labeled 1-3

and 6. B) Non-cytoplasmic ascorbate region with predicted hydrogen bonds to Dcytb, CG1275, and

Nemy shown in cyan; an additional predicted hydrogen bond between Nemy Lys161 and Glu236 is also

shown. Zinc coordination bonds are shown as orange dashed lines.

Of the five residues proximate to cytoplasmic ascorbate in Dcytb and AtCytb561-B (basic

residues at H2-7, H2-3, and H4-7, aromatic at H3+22, bulky hydrophobic at H4-3), only an arginine at H4-

7 (Arg509) is completely conserved across CG8399 sequences (amino acid numbering in this section

corresponds with *D. melanogaster* CG8399) (Fig 2). Most CG8399 sequences also have a conserved

lysine at H2-7, although *D. melanogaster* CG8399 has a threonine (Thr443) (S1 Fig). To visualize

conservation on the cytoplasmic side of CG8399 proteins, we overlayed the CG8399 consensus

sequence alignment onto the *D. melanogaster* CG8399 AlphaFold model (Fig 8). Of the remaining

positions discussed in the interaction of cytoplasmic ascorbate with Dcytb, AtCytb561-B, CG1275, and

Nemy proteins (H2-3, H3+22, and H4-3), CG8399 sequences have conserved residues with differing

properties. At H2-3, CG8399s have a phenylalanine (Phe447) rather than the lysine found in Dcytb,

AtCytb561-B, CG1275, and Nemy proteins, and a conserved tryptophan (Trp446) precedes H2-3 at H2-4

(Fig 8). Instead of a bulky hydrophobic residue at H3+22, a proline (Pro502) is present in CG8399s; that

proline is preceded by a completely conserved CG8399 arginine (Arg501) at H3+21 (Fig 8). Instead of a

bulky hydrophobic residue at H4-3, CG8399 proteins have an asparagine (Asn513). We noted additional

highly-conserved positions in this region among CG8399 proteins, including a second proline at H3+24

(Pro504) and a basic residue at H1+23 (Lys432), which is either a lysine or arginine in all the CG8399

sequences. H1+23 is an arginine in most CG1275 and Nemy sequences although not in Dcytb (Trp73) or

AtCytb561-B (Gln74) (Fig 4). While side-chain placement cannot be determined with high confidence, it

is notable how highly conserved this region is across multi-domain CG8399 proteins, including conserved

residues suitable for binding ascorbate: basic residues at H1+23, H3+21, and H4-7, and aromatics at H2-

4 and H2-3 (Fig 8; S5 Fig).

Fig 8. Cytoplasmic side of *D. melanogaster* CG8399 model. AlphaFold structure of *D. melanogaster*

CG8399 overlayed with multi-domain CG8399 consensus sequence alignment. The model is gradient

colored with high conservation (1.68 score) in magenta to low conservation (-1.89 score) in cyan. Select

side chains are shown in stick form with oxygen in red and nitrogen in blue and are labeled by position

relative to conserved heme-coordinating histidines with *D. melanogaster* CG8399 amino acid numbering

in parentheses. Ascorbate binding on the cytoplasmic surface of CG8399 may be facilitated by a cluster

of aromatic and basic residues: Lys432, Trp446, Phe447, Arg501, and Arg509. Also shown are conserved

H2 (His450) and H4 (His516), which coordinate the cytoplasmic heme iron (not present in model). Two

well-conserved prolines at positions H3+22 and H3+24 stabilize the loop between helices 3 and 4. Other

cytoplasmic side positions with well conserved residues include H2-7, which typically has a lysine but is

Thr443 in *D. melanogaster*, and H4-3 (Asp513). Scale bar depicts 10 Å.

Some Group 4 proteins also have a basic residue at two of the three positions that hydrogen

bond with cytoplasmic ascorbate in Dcytb and AtCytb561-B: H2-7 (11 out of 25) and H4-7 (11 out of 25)

(S1 Fig). While only six Group 4 proteins have a basic residue at H2-3, nearly all have basic residues at

H2-4 (23 out of 25) and H4-3 (23 out of 25) (S1 Fig). This is notable since H4-3 is one of the two positions

we identified as having a potential hydrophobic contact with cytoplasmic ascorbate in Dcytb, CG1275,

and Nemy (Fig 7A; S3 Table). It is possible that the basic residues at these highly-conserved positions are

involved in Group 4 cytoplasmic ascorbate binding. At the other position (H3+22) with predicted

hydrophobic ascorbate contact, most Group 4 proteins (19 out of 25) also have a large, hydrophobic

residue that could potentially interact with ascorbate or other possible cytoplasmic reducing agents,

such as dihydrolipoic acid (S1 Fig)[53]. The adjacent position, H3+23, is also typically an aromatic residue

(17 out of 25) or occasionally a lysine (3 out of 25) and may be involved in cytoplasmic ascorbate binding

(S1 Fig).

Non-cytoplasmic interactions with ascorbate and zinc: implications for ferric

reductase activity

Dcytb and AtCytb561-B also bind ascorbate at their non-cytoplasmic surfaces [13,18]. In Dcytb,

this appears to be part of its ferric reduction mechanism [18]. To identify an iron-binding site in Dcytb,

the protein was crystallized (PDB: 5ZLG) in the presence of redox inactive zinc, which presumably binds

Dcytb in a manner similar to iron [18]. Details of Dcytb ascorbate-complexed metal binding are

described below, and this information was used to predict non-cytoplasmic ascorbate and metal binding

by insect cytb561 proteins.

In Dcytb the only residue found to directly interact with non-cytoplasmic ascorbate was Phe184

(H4+25) via a van der Waals interaction, although several residues were discussed as contributing to the

environment of the ascorbate-binding pocket [18]; in AtCytb561-B the same H4+25 (Phe182) interaction

was reported, and several residues were also suggested to form possible hydrogen bonds, including

H2+22 (His106) and H3-3 (Tyr115) [13]. Our analysis revealed that three Dcytb residues have predicted

contacts with non-cytoplasmic ascorbate: H1-3 (Phe47), H2+22 (His108), and H4+25 (Phe184) (Fig 7B; S3

Table). A phenylalanine at H1-3 is also predicted to make ascorbate contacts in *D. melanogaster* CG1275

and Nemy (S3 Table); a phenylalanine at this position is conserved across CG1275 and Nemy sequences,

and is centrally located – near the zinc ion, the non-cytoplasmic ascorbate, and the non-cytoplasmic

heme – in Dcytb, CG1275, and Nemy structures (Fig 7B). In Dcytb, Phe47 was the only residue that had

an observed change in electron density upon zinc ion and ascorbate binding, indicating a change in

position [18]. Most Group 4A proteins have a phenylalanine at H1-3; however, there is less conservation

at this position in CG8399 and Group 4B proteins (S1 Fig).

As mentioned above, the phenylalanine (Phe184) at H4+25 was determined to form a van der

Waals interaction with the non-cytoplasmic ascorbate in Dcytb and AtCytb561-B [13,18]. An aromatic

amino acid, most often phenylalanine, is typical at H4+25 across CG1275 and Nemy sequences with a

small number of exceptions; the most dramatically different is a glutamate (Glu236) in *D. melanogaster*

Nemy (Fig 4). In the absence of ascorbate, our analysis revealed that Glu236 and a lysine at H3-7

(Lys161) are predicted to form a hydrogen bond in the *D. melanogaster* Nemy model that would run

straight through the region occupied by the non-cytoplasmic ascorbate ring (Fig 7B). The *D.*

melanogaster Nemy model also has extensive contacts predicted between the non-cytoplasmic

ascorbate and Lys161 (23, including one hydrogen bond) and Glu236 (29) (S3 Table). These results

suggest that *D. melanogaster* Nemy could bind non-cytoplasmic ascorbate though a combination of

compensatory substitutions in lieu of an aromatic residue at H4+25. H4+25 is slightly outside the

conserved core domain and therefore was not included in our analyses of CG8399 and Group 4

sequences.

In Dcytb, the histidine (His108) at H2+22 was determined to be part of an ascorbate-metal-

binding complex; His108 was found to directly coordinate with a redox inactive zinc ion, which directly

interacted with the non-cytoplasmic ascorbate [18]. The histidine at H2+22 is conserved across CG1275

and Nemy sequences and is predicted to be similarly positioned in *D. melanogaster* CG1275 and Nemy

models (Fig 4; Fig 7B). In contrast, a histidine at H2+22 is not conserved in other insect cytb561s (Fig 2).

The residue at H2+22 is nonpolar in CG8399 sequences, whereas many Group 4 proteins have a lysine at

this position, which could contribute to ascorbate binding but would not be expected to coordinate a

metal cation (Fig 2; S1 Fig).

His108 was also found to be critical for Dcytb ferric reductase activity, as were residues at H2+21

(Asn107), H3-3 (Tyr117), and H4+25 (Phe184, discussed above) [18]. Like Dcytb, CG1275 and Nemy

proteins have a tyrosine at H3-3 in addition to the histidine at H2+22 and phenylalanine at H4+25, and

they tend to have a conservatively-substituted serine at the H2+21 position (Fig 2). These conserved

residues support the hypothesis that CG1275 and Nemy proteins have ferric reductase activity. In

contrast, conservation of these residues in the other insect cytb561 proteins is limited to a glutamate

that is entirely conserved at H2+21 in multi-domain CG8399 proteins (discussed further below) (Fig 2).

Some Group 4A proteins may have a different type of non-cytoplasmic ascorbate-binding site. A lysine-

containing "KXXXXKXH" motif was observed in the non-cytoplasmic loop of the core domain in six of the

seven Group 4A proteins (excluding one of the two *C. felis* sequences) (Fig 2; S1 Fig). Group 4A includes

the proteins identified as more like human TScytb than any *D. melanogaster* cytb561 (Table 1; S2 Table).

We aligned these six Group 4A proteins with human TScytb (which can reduce ferric ions in nanodiscs

[14,54]) and TScytb-like proteins of diverse animal species and found that all sequences have this

"KXXXXKXH" motif (Fig 9A), which contributes to the high degree of conservation across TScytb proteins

on the non-cytoplasmic side (Fig 9B). The location of the "KXXXXKXH" motif in the non-cytoplasmic loop

between helices 3 and 4 (where substrate binding occurs in Dyctb), as well as its strict conservation in

TScytb-like proteins of diverse species, suggests an important functional role, possibly related to

ascorbate binding (Fig 9C). This motif was not identified in other human or insect cytb561s. A completely

conserved tryptophan at H4+23 could also potentially contribute to a non-cytoplasmic binding site, and

the histidine at H3-5 (within the "KXXXXKXH" motif) could potentially coordinate a ferric ion complexed

with ascorbate, similar to the substrate binding site of Dytcb, providing a site for ferric reduction in

TScytb-like proteins (Fig 9C).

Fig 9. A conserved "KXXXXKXH" motif in Group 4A and related proteins. A) Truncated alignment of

animal TScytb and Group 4A sequences shows a conserved "KXXXXKXH" motif upstream of the

conserved non-cytoplasmic heme-coordinating H3. Coloring by amino acid residue as in Fig 4.

Abbreviations: Cf, *C. felis*; Aten, *A. tenebrosa*; Hrob, *H. robusta*; Drer, *D. rerio*; Xlae, *X. laevis*; Pmon, *P.

montanus*; Mmus, *M. musculus*; Hsap, *H. sapiens*; Apla, *A. planci*; Osin, *O. sinensis*; Am, *A. mellifera*; Tc,

T. castaneum; Zn, *Z. nevadensis*; Px, *P. xuthus*; Ag, *A. gambiae*. B) Full alignment of animal Tscytb and

Group 4A sequences overlayed on AlphaFold model of *A. gambiae* XP_3200673.4 (AF-Q7PZD0-F1-

model_v4). H1-H4 shown in stick form with nitrogen in blue. C) Non-cytoplasmic side of the protein

contains completely conserved lysines (H2+22, H3-7), a histidine (H3-5), a tryptophan (H4+23), and a

phenylalanine (H1-3), in addition to H1 and H3; these residues are shown in stick with nitrogen atoms in

blue. B-C) Amino acid conservation is shown by a gradient of color as in Fig 3. Scale bars show 10 Å.

Other amino acids of interest

Aside from H1-H4, the most highly-conserved residues in the core domain of insect cytb561s are

small residues and hydrophobic residues (S1 Fig). As discussed above, four well-conserved, small

residues (H3+4, H3+18, H4+4, and H4+18) appear to be important for heme placement. The remaining

well-conserved residues may play a role in helical packing [55]. The small amino acids at H1+7 and

H2+14 are positioned where TM2 and TM3 face each other, while the small residue at H1+13 is on the

side of TM2 that faces TM5.

Of the investigated insect cytb561s, Nemy proteins possess unique cysteine-rich sequences in

their cytoplasmic loops, with double cysteines common in both cytoplasmic loops of the core domain

(S1 Fig). This observation is notable, as cysteines are among the least common amino acids in proteins,

yet they are often at positions of functional importance [56]. Cytoplasmic cysteine residues can

participate in redox-active disulfide formation and metal ligation, and they can also serve as sites of

post-translational modifications [56]. While the pH and reducing environment of the cytoplasm is not

conducive to disulfide bond formation, solvent exposed cysteines can be regulated by redox state [57].

The cytoplasm-exposed Nemy cysteines could form redox active disulfides, potentially functioning as an

on/off switch for electron transfer activity. Another possibility involves cysteine S-palmitoylation, given

that *D. melanogaster* Nemy was identified as a palmitoylated protein [58]. Cysteine S-palmitoylation is a

reversible post-translational modification used to regulate protein function [59]. Given its dynamic

nature, cysteine S-palmitoylation can direct varied cellular processes, including subcellular trafficking,

protein localization, and enzyme activity [59]. We did not detect multiple cysteine-rich cytoplasmic loop

sequences in the cytb561 domain of other insect cytb561s, nor in human cytb561 proteins, suggesting a

function unique to Nemy proteins. Future research could indicate whether these cysteines are

palmitoylated and whether they influence Nemy localization, cytoplasmic substrate binding, or protein

function.

708   Analysis of the CG8399 DOMON domain

709   Lack of conservation in the extracellular sections of the CG8399 cytb561 domain

710   suggests DOMON domain involvement in ferric reductase activity

As discussed above, many of the Dyctb non-cytoplasmic surface residues that interact with

ascorbate or function in ferric reduction are missing in CG8399 proteins (Fig 2) [13,18]. Nevertheless, *D.*

melanogaster CG8399 and its ortholog in mice, SDR2, possess in vitro ferric reductase activity [6,31].

These observations indicate that ferric reductase activity in CG8399 and SDR2 occurs through a

mechanism different from that of Dcytb and suggests the possibility that iron-binding in CG8399 and

SDR2 occurs in the extracellular DOMON domain. It has been previously proposed that CYBDOMs may

act as a "redox bridge" between cytosolic ascorbate and the extracellular environment [1]. Using the

redox bridge concept as a model, we suggest that the CG8399 cytb561 domain binds the reducing agent

(ascorbate) on the cytoplasmic surface, then transfers the electron to a DOMON domain heme, which

then transfers the electron to an electron-accepting substrate (possibly a metal ion).

CG8399 DOMONs may bind a b-type heme

To further investigate the DOMON domain in CG8399 proteins, we used the HHpred PDB server

to identify proteins with known structure that have the highest probability percentage of homology to

CG8399 DOMON sequences. We found that all insect CG8399 DOMONs have > 99% probability of

homology to the DOMON domain of human DBH and the cytochrome domain of *P. chrysosporium* CDH.

(Note that despite the use of "cytochrome" in the domain name of *P. chrysosporium* CDH, this domain is

not homologous with the cytochrome b561 domain of CG8399 proteins; rather, the *P. chrysosporium*

CDH cytochrome domain is homologous with the DOMON domain in CG8399 proteins.) Intriguingly, DBH

and CDH have only 10% sequence identity and bind different ligands, yet their structural alignment

suggests that both domains have very similar folds [60].

The crystal structure of the CDH cytochrome domain (DOMON homolog) revealed a b-type

heme group coordinated by an unusual methionine and histidine ligation of the heme iron [61]. HHpred

alignments of CDH with each individual insect CG8399 DOMON showed conservation of these

methionine and histidine residues in all sequences (Met286 and His379 in *D. melanogaster* CG8399 [S2

Fig]). Notably, the methionine and histidine are not conserved in the DOMONs of proteins that bind

sugar rather than heme, nor in DBH, which binds neither sugar nor heme [20,60]. We then generated an

alignment of CG8399 DOMON domains with the DOMON domains of diverse animal CYBDOM proteins,

including insect CG8399s, and found universal conservation of the methionine and histidine across all

species investigated, including both DOMON domains in the *A. pisum* CG8399 (S3 Fig). These results

agree with previous observations that CYBDOM proteins likely have a b-type heme group in their

DOMON domain [1]. Given the DOMON domain's suspected role in interdomain electron transfer and

the likelihood that CG8399s and their CYBDOM orthologs coordinate a heme group within their DOMON

domains, it seems likely that their putative ferric reductase activity occurs outside of the cytb561

domain.

Spatial relationship between the DOMON domain and cytb561 domain in

CG8399

To study the possibility of electron transfer to the DOMON domain, we analyzed the multi-

domain CG8399 consensus sequence overlay of the AlphaFold model for D. *melanogaster* CG8399,

which includes the extracellular reeler and DOMON domains in addition to the transmembrane cytb561

domain (Fig 10). In the following discussion, residue numbers relate to the *D. melanogaster* CG8399

protein sequence. The model predicts a DOMON domain largely composed of two five-strand beta

sheets, similar to the DOMON domain in DBH [60]. The predicted aligned error plots for the DOMON

and cytb561 domains were relatively low, suggesting their models could be informative for positional

insights [47,48]. (In contrast, the predicted position of the reeler domain was reported with high error.)

Notably, the *D. melanogaster* DOMON domain is oriented adjacent to the cytb561 domain, with

predicted hydrogen bonds between the side chains of Lys285 and Arg383 in the DOMON domain with

the backbone carbonyl oxygens of Leu472 and Pro538 in the cytb561 domain, respectively (Fig 10B). Salt

bridges are predicted to form in the non-cytoplasmic loops of the cytb561 domain between Arg403 and

Glu541, as well as Glu471 and Lys539 (Fig 10B).

Fig 10. Model of *D. melanogaster* CG8399 demonstrating interactions between domains, and a putative iron binding site. A-C) Models were colored by ChimeraX based on amino acid conservation in the nine multidomain CG8399 sequences. Color gradient as in Fig 3. Residues in stick form show oxygen in red, nitrogen in blue and sulfur in yellow. A-D) Scale bars depict 10 Å. A) CG8399 has three distinct domains: reeler (top left), DOMON (middle right), and cytb561 (bottom). Residues expected to be heme-coordinating are shown in stick (some not visible). B) Selected sidechains at the interface of the DOMON (top) and cytb561 (bottom) domains. Predicted heme-coordinating residues (Met286 and His379) in the DOMON domain are positioned at the bottom of the domain where it is predicted to interact with the cytb561 domain. Interdomain hydrogen bonds were predicted between the sidechain of Lys285 (DOMON) and the backbone of Leu472 (cytb561), as well as the sidechain of Arg383 (DOMON) and the backbone of Pro538 (cytb561). Potential salt bridge interactions may form in the cytb561 domain loop regions between Glu471 and Lys539, as well as Arg403 and Glu541. C) A putative metal binding site in a highly conserved loop with two aspartates and a serine residue. D) Space filling model of the same part of the structure depicted in C, with Coulombic coloring calculated by ChimeraX, red for negative charge, white for neutral, and dark blue for positive charge.

We then located the predicted positions of the putative heme-coordinating residues (Met286 and His379) in the DOMON domain, with particular interest in their relative positions to the non-cytoplasmic heme-coordinating histidines (His409 and His480) in the cytb561 domain. The location of the putative DOMON heme-binding pocket is predicted to be directly adjacent to the non-cytoplasmic heme-binding region of the cytb561 domain (Fig 10B) [47,48]. These predictions further support the hypothesis that the CG8399 cytb561 domain functions in interdomain electron transport rather than being the site for oxidoreductase activity.

The CG8399 DOMON domain may contain a metal-binding pocket

If CG8399 ferric reductase activity occurs within the DOMON domain, we would expect to locate

an iron-binding pocket within this domain. Our DOMON alignment revealed multiple well-conserved

aspartates, including strict conservation of Asp283 and Asp289 (S3 Fig). Asp283 and Asp289 are in a

highly-conserved region (sequence "SXDXXMGXD") that includes the putative heme-coordinating

methionine (Met286) (S3 Fig). This region also contains additional aspartates, especially in insect

CG8399s, of which two are particularly well conserved (S3 Fig). We located this region in the *D.*

melanogaster CG8399 model and found that a negatively-charged pocket is predicted to be directly

behind the putative heme-binding pocket, similar to the position of the putative metal ion-binding

pocket in DBH relative to its ligand-binding pocket (Figs 10C and D) [60]. These results are consistent

with the DOMON domain being the site of ferric reductase activity, wherein strictly-conserved Ser281,

Asp283, and Asp289 could form an iron-binding pocket in insect CG8399s and CYBDOMs from other

species (S3 Fig). Metal-binding sites formed by aspartate residues have been suggested for other

proteins. For example, the DBH DOMON domain has a putative metal ion-binding site composed of

aspartate sidechains and the backbone carbonyl oxygens of two other amino acids [60], and three

aspartate residues form a binding site for a variety of cations, including Fe^{2+} and Fe^{3+}, in LarE [62]. This

model also provides a possible explanation for the ferric reductase activity of *D. melanogaster* CG8399

and mouse SDR2, despite their lack of conserved ascorbate- and metal-binding residues on the non-

cytoplasmic surface of the cytb561 domain.

Three insect cytb561s with distinctive features outside of the core

domain

Of the 54 identified insect cytb561s, we noted that three sequences possess unique features

within the cytb561 domain but outside the domain core (TM2-TM5 in Nemy proteins, and TM1-TM4 in

CG8399s). To determine whether other insect cytb561 proteins have any of these unique features, we

performed additional BLAST searches against the entire class Insecta.

Compared to other Nemys, the *A. pisum* sequence has approximately 150 additional N-terminal

residues preceding helix 1 and approximately 50 additional residues in the non-cytoplasmic loop

connecting helix 1 and helix 2, which immediately precedes the core cytb561 domain (S4 Fig). We found

proteins with similar extended regions in eight additional species of the Aphididae family, but not in

other insects, suggesting that these regions may serve a unique function within aphids (S4 Fig; S5 Table).

The extended N-terminal region preceding helix 1 is acidic and includes relatively high percentages of

aspartate, glycine, serine, and threonine (S4 Fig).

Of the investigated CG8399s, only the *A. pisum* sequence possesses two DOMON domains. We

identified 888 CYBDOM hits in the class Insecta, of which only 19 were found to have two DOMON

domains: nine from other species of aphids, eight from other insects from the order Hemiptera, and two

from insects in the order Coleoptera (S6 Table). It appears that multiple DOMONs are uncommon in

insect cytb561s; however, they have a relatively high occurrence in Hemipterans. Both *A. pisum* DOMON

domains contain the conserved methionine and histidine that are assumed to coordinate a heme group

(S3 Fig); however, it is unknown how these two DOMON domains are spatially arranged.

While all nine investigated insect species possess a multidomain CG8399, *A. gambiae* (mosquito)

is the only species with a second, single domain CG8399-like protein (XP_314065.2); this sequence lacks

both reeler and DOMON domains. Further investigation revealed similar single-domain (cytb561 only)

CG8399-like proteins in ten additional mosquito species, as well as two species of fungus gnats, which

are also in the order Diptera (S7 Table). The single-domain CG8399-like protein is more similar to the

cytb561 domain in multidomain CG8399 proteins than to the cytb561 domain in other groups of

cytb561s (Table 1; S2 Table), including its core domain spanning TM1-TM4, rather than TM2-TM5.

Cyb561 subcellular localization

Subcellular localization is important for protein function in an organism. We submitted each of

the 54 full-length insect cytb561s to the SignalP-5.0 webserver for detection of predicted signal

peptides, and to the DEEPLOC-1.0 webserver for prediction of subcellular protein localization [40]. Only

the multidomain CG8399 proteins had a predicted signal peptide, and they are the only sequences with

predicted plasma membrane localization (S8 Table). This result is consistent with the presence of *D.*

melanogaster CG8399 in a plasma membrane fraction of fly head proteins [63]. The *A. gambiae* single-

domain CG8399-like protein is predicted to be in the lysosome, as are all the CG1275 and Nemy proteins

(S8 Table). The subcellular localization predictions for Group 4 proteins were not particularly robust, and

the predicted location varied depending on the specific Group 4 protein (S8 Table).

An alignment of the Dcytb, Nemy, and CG1275 carboyxl-termini revealed that only the CG1275

proteins have a "EDXXLL" motif near TM6 that is consistent with a dileucine-based late-

endosomal/lysosomal targeting signal "(D/E)XXXL(L/I)" (Fig 11) [42]. *A. mellifera* and *T. castaneum*

CG1275 proteins lack one conserved leucine, and one of the four *A. pisum* sequences (XP_003246890.1)

lacks both leucines, yet these three sequences are still predicted to localize to the lysosome (Fig 11; S8

Table).

Fig 11. Lysosomal targeting signal in CG1275. An alignment of the C-terminal portion of CG1275

proteins. Helix predictions from the AlphaFold model of *D. melanogaster* CG1275 shown in gray (helix 6)

and an additional helical region is in black. The EDXXLL motif is boxed in blue in the consensus logo.

Summary of cytochrome b561s as potential ferric reductases in insects

A major goal of this study was to predict which of the insect cytb561 orthologous groups may

function as ferric reductases and, thus, have a possible role in iron uptake by insect cells. Our analyses

suggest that none of the four groups of insect cytb561 proteins can be excluded as ferric reductase

candidates.

Based on sequence similarity, predicted structural similarity, and conservation of key amino acid

residues, we conclude that CG1275 and Nemy proteins are strikingly similar to the well-established

ferric reductase Dcytb, as well as the closely-related human proteins Lcytb and CGcytb. These similarities

extend to the conservation of a metal-binding histidine and other amino acid residues that are required

for ferric reductase activity by Dcytb. To date, no functions of CG1275 proteins have been discovered;

however, if CG1275 proteins have ferric reductase activity, the constitutive, nearly ubiquitous

expression of *D. melanogaster* CG1275 [64,65] would be consistent with a role in iron uptake by many

cell types. We found that CG1275 proteins have a C-terminal dileucine motif that directs proteins to a

late-endosomal/lysosomal location, suggesting that CG1275 proteins could function similarly to the non-

homologous STEAP3 protein, which reduces Fe^{3+} to Fe^{2+} prior to uptake across endosomal and lysosomal

membranes [27,28,66]. Among mammalian cytb561s, a C-terminal dileucine motif is present only in

Lcytb proteins [7]. Interestingly, Lcytb appears to function as a lysosomal ferric reductase in Burkitt

lymphoma cells, filling the role that STEAP3 plays in some other cell types [67]. Although Nemy proteins

have many similarities with Dcytb and Lyctb, their one known function is unrelated to iron homeostasis

[4,30]; however, Nemy proteins could be multifunctional enzymes that also play a role in iron uptake.

D. melanogaster CG8399 and the mammalian ortholog SDR2 have in vitro ferric reductase activity

[6,15,31], but the in vivo functions of CG8399 and SDR2 proteins are not known. Based on a previous

biochemical study [63] and our subcellular localization predictions, CG8399 proteins localize to the

plasma membrane, with their reeler and DOMON domains in the extracellular compartment. Their

presence in the plasma membrane may enable reduction of extracellular ferric ions prior to iron uptake.

Our analyses of the predicted structures of the CG8399 DOMON and cytb561 domains suggest that the

two domains may function together in ferric reduction, with the interdomain transfer of an electron

from cytoplasmic ascorbate to a heme in the extracellular DOMON domain and then to a ferric ion that

is perhaps coordinated nearby by multiple aspartate residues. Because *D. melanogaster* CG8399

expression is limited to a subset of tissues [64,65], its hypothetical role in iron uptake would be limited

to specific cell types.

We have the least amount of evidence to support the hypothesis that the diverse Group 4

proteins function as ferric reductases, and a lack of one-to-one orthologs indicates that these proteins

may have non-conserved physiological functions. The expression profiles of all but one of the *D.*

melanogaster Group 4 genes is moderately to highly restricted [64,65], excluding a role in iron uptake by

most cell types. The Group 4 proteins have low similarity to the well-studied mammalian cytb561s,

lacking even conserved residues that bind ascorbate in the cytoplasm. However, multiple basic residues

and a large hydrophobic residue at the cytoplasmic surface of most Group 4 proteins may enable the

Group 4 proteins to use ascorbate as an electron donor. On the non-cytoplasmic side, a subset of Group

4 proteins contain a "KXXXXKXH" motif that is also present in TScytb, a protein with an unknown

biochemical function in vivo, but ferric reductase activity in vitro [14]. We propose that the "KXXXXKXH"

motif may bind ascorbate and a ferric ion in a hypothetical ferric reduction site.

Acknowledgements

We wish to acknowledge the undergraduate research students who participated in aspects of this project: Clare Burnett, Chrissy Chan, Caleb Garris, Keyata Lewis, Gregory Kane, Karen Nunez Sifuentes, Connor Sullivan, and Nataliya Yusopov. Thanks also to the MSU Denver CHE 4320 students from Spring 2018, Fall 2018, Spring 2019, Fall 2019 and Spring 2020 who participated in iterations of the Iron Uptake in Insects CURE. We thank Dr. Andrew McMillan for thoughtful comments on the manuscript. Molecular graphics and analyses were performed with UCSF ChimeraX, developed by the Resource for Biocomputing, Visualization, and Informatics at the University of California, San Francisco. FlyBase was used for gene identification and as a source of expression data.

References

1. Asard H, Barbaro R, Trost P, Bérczi A. Cytochromes *b* 561: Ascorbate-mediated trans-membrane electron transport. Antioxid Redox Signal. 2013;19(9):1026–35. doi: 10.1089/ars.2012.5065

2. Berczi A, Zimanyi L. The trans-membrane cytochrome b561 proteins: Structural information and biological function. Curr Protein Pept Sci. 2014;15(8):745–60. doi: 10.2174/1389203715666140828100351

3. Verelst W, Asard H. A phylogenetic study of cytochrome b561 proteins. Genome Biol. 2003;4(6) R38. doi: 10.1186/gb-2003-4-6-r38

4. Iliadi KG, Avivi A, Iliadi NN, Knight D, Korol AB, Nevo E, et al. nemy encodes a cytochrome b561 that is required for Drosophila learning and memory. Proc Natl Acad Sci. 2008;105(50):19986–91. doi: 10.1073/pnas.0810698105

5. Latunde-Dada GO, Van der Westhuizen J, Vulpe CD, Anderson GJ, Simpson RJ, McKie AT. Molecular and functional roles of duodenal cytochrome B (Dcytb) in iron metabolism. Blood Cells Mol Dis. 2002;29(3):356–60. doi: 10.1006/bcmd.2002.0574

6. Vargas JD, Herpers B, McKie AT, Gledhill S, McDonnell J, van den Heuvel M et al. Stromal cell-derived receptor 2 and cytochrome b561 are functional ferric reductases. Biochim Biophys Acta BBA - Proteins Proteomics. 2003;1651(1–2):116–23. doi: 10.1016/S1570-9639(03)00242-5

7. Zhang D liang, Su D, Bérczi A, Vargas A, Asard H. An ascorbate-reducible cytochrome b561 is localized in macrophage lysosomes. Biochim Biophys Acta BBA - Gen Subj. 2006;1760(12):1903–13. doi: 10.1016/j.bbagen.2006.07.019

8.   Lane D, Bae DH, Merlot A, Sahni S, Richardson D. Duodenal cytochrome b (DCYTB) in iron
metabolism: An update on function and regulation. Nutrients. 2015;7(4):2274–96. doi:
10.3390/nu7042274

9.   Lemler DJ, Lynch ML, Tesfay L, Deng Z, Paul BT, Wang X, et al. DCYTB is a predictor of
outcome in breast cancer that functions via iron-independent mechanisms. Breast Cancer
Res. 2017;19(1):25. doi: 10.1186/s13058-017-0814-9

10.   Su D, May JM, Koury MJ, Asard H. Human erythrocyte membranes contain a cytochrome
b561 that may be involved in extracellular ascorbate recycling. J Biol Chem.
2006;281(52):39852–9. doi: 10.1074/jbc.M606543200

11.   Cenacchi L, Busch M, Schleidt PG, Müller FG, Stumpp TVM, Mäntele W, et al. Heterologous
production and characterisation of two distinct dihaem-containing membrane integral
cytochrome b(561) enzymes from *Arabidopsis thaliana* in *Pichia pastoris* and *Escherichia coli*
cells. Biochim Biophys Acta. 2012;1818(3):679–88. doi: 10.1016/j.bbamem.2011.10.030

12.   Klein M, Deniz E, Heit S, Wille G, Mäntele W, Lancaster CRD. Proton-coupled electron
transport in two distinct CYBASC paralogs of *Arabidopsis thaliana*: A comparative
characterization of highly conserved tyrosine and lysine residues. Biochemistry.
2020;59(25):2328–39. doi: 10.1021/acs.biochem.0c00155

13.   Lu P, Ma D, Yan C, Gong X, Du M, Shi Y. Structure and mechanism of a eukaryotic
transmembrane ascorbate-dependent oxidoreductase. Proc Natl Acad Sci.
2014;111(5):1813–8. doi: 10.1073/pnas.1323931111

14.   El Behery M, Fujimura M, Kimura T, Tsubaki M. Direct measurements of ferric reductase
activity of human 101F6 and its enhancement upon reconstitution into phospholipid bilayer
nanodisc. Biochem Biophys Rep. 2020;21:100730. doi: 10.1016/j.bbrep.2020.100730

15.   Su D, Asard H. Three mammalian cytochromes b561 are ascorbate-dependent
ferrireductases. FEBS J. 2006;273(16):3722–34. doi: 10.1111/j.1742-4658.2006.05381.x

16.   Kelley PM, Njus D. Cytochrome b561 spectral changes associated with electron transfer in
chromaffin-vesicle ghosts. J Biol Chem. 1986;261(14):6429–32. doi: 10.1016/S0021-
9258(19)84579-9

17.   Njus D, Knoth J, Cook C, Kelly PM. Electron transfer across the chromaffin granule
membrane. J Biol Chem. 1983;258(1):27–30. doi: 10.1016/S0021-9258(18)33212-5

18.   Ganasen M, Togashi H, Takeda H, Asakura H, Tosha T, Yamashita K, et al. Structural basis for
promotion of duodenal iron absorption by enteric ferric reductase with ascorbate. Commun
Biol. 2018;1(1):120. doi: 10.1038/s42003-018-0121-8

19.   Tsubaki M, Takeuchi F, Nakanishi N. Cytochrome b561 protein family: Expanding roles and
versatile transmembrane electron transfer abilities as predicted by a new classification
system and protein sequence motif analyses. Biochim Biophys Acta BBA - Proteins
Proteomics. 2005;1753(2):174–90. doi: 10.1016/j.bbapap.2005.08.015

20.   Iyer LM, Anantharaman V, Aravind L. The DOMON domains are involved in heme and sugar
recognition. Bioinformatics. 2007;23(20):2660–4. doi:10.1093/bioinformatics/btm411

21.   Gorman MJ. Iron Homeostasis in Insects. Annu Rev Entomol. 2023;68(1):51-67. doi:
10.1146/annurev-ento-040622-092836

22.   Tang X, Zhou B. Iron homeostasis in insects: Insights from *Drosophila* studies: Iron
Homeostasis in Insects. IUBMB Life. 2013;65(10):863–72. doi: 10.1002/iub.1211

23.   Wu S, Yin S, Zhou B. Molecular physiology of iron trafficking in *Drosophila melanogaste*r. Curr
Opin Insect Sci. 2022;50:100888. doi: 10.1016/j.cois.2022.100888

24.   Andrews NC, Fleming MD, Gunshin H. Iron transport across biologic membranes. Nutr Rev.
1999;57(4):114–23. doi: 10.1111/j.1753-4887.1999.tb06934.x

25. Kosman DJ. A holistic view of mammalian (vertebrate) cellular iron uptake. Metallomics.
2020;12(9):1323–34. doi: 10.1039/D0MT00065E
26. Montalbetti N, Simonin A, Kovacs G, Hediger MA. Mammalian iron transporters: Families
SLC11 and SLC40. Mol Aspects Med. 2013;34(2):270–87. doi: 10.1016/j.mam.2013.01.002
27. Gomes IM, Maia CJ, Santos CR. STEAP proteins: From structure to applications in cancer
therapy. Mol Cancer Res. 2012;10(5):573–87. doi: 10.1158/1541-7786.MCR-11-0281
28. Knutson MD. Steap Proteins: Implications for Iron and Copper Metabolism. Nutr Rev.
2007;65(7):335–40. doi: 10.1111/j.1753-4887.2007.tb00311.x
29. Schlottmann F, Vera-Aviles M, Latunde-Dada GO. Duodenal cytochrome b (Cybrd1) ferric
reductase functional studies in cells. Metallomics. 2017;9(10):1389–93. doi:
10.1039/C7MT00254H
30. Knight D, Iliadi KG, Iliadi N, Wilk R, Hu J, Krause HM, et al. Distinct regulation of transmitter
release at the Drosophila NMJ by different isoforms of nemy. PLOS ONE.
2015;10(8):e0132548. doi: 10.1371/journal.pone.0132548
31. Picco C, Scholz-Starke J, Naso A, Preger V, Sparla F, Trost P, et al. How are cytochrome b561
electron currents controlled by membrane voltage and substrate availability? Antioxid Redox
Signal. 2014;21(3):384–91. doi: 10.1089/ars.2013.5809
32. Mandilaras K, Pathmanathan T, Missirlis F. Iron absorption in *Drosophila melanogaster*.
Nutrients. 2013;5(5):1622–47. doi: 10.3390/nu5051622
33. Larkin A, Marygold SJ, Antonazzo G, Attrill H, dos Santos G, Garapati PV, et al. FlyBase:
updates to the *Drosophila melanogaster* knowledge base. Nucleic Acids Res.
2021;49(D1):D899–907. doi: 10.1093/nar/gkaa1026
34. NCBI Resource Coordinators. Database resources of the National Center for Biotechnology
Information. Nucleic Acids Res. 2018;46(D1):D8–13. doi: 10.1093/nar/gkx1095
35. Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, et al. CDD/SPARCLE: the
conserved domain database in 2020. Nucleic Acids Res. 2020;48(D1):D265–8. doi:
10.1093/nar/gkz991
36. Gabler F, Nam SZ, Till S, Mirdita M, Steinegger M, Söding J, et al. Protein sequence analysis
using the MPI bioinformatics toolkit. Curr Protoc Bioinforma. 2020;72(1):e108. doi:
10.1002/cpbi.108
37. Sievers F, Higgins DG. Clustal omega. Curr Protoc Bioinforma. 2014;48(1):3.13.1-3.13.16. doi:
10.1002/0471250953.bi0313s48
38. Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. Jalview version 2--a multiple
sequence alignment editor and analysis workbench. Bioinformatics. 2009;25(9):1189–91. doi:
10.1093/bioinformatics/btp033
39. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of
high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol.
2011;7:539. doi: 10.1038/msb.2011.75
40. Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, et al.
SignalP 5.0 improves signal peptide predictions using deep neural networks. Nat Biotechnol.
2019;37(4):420–3. doi: 10.1038/s41587-019-0036-z
41. Almagro Armenteros JJ, Sønderby CK, Sønderby SK, Nielsen H, Winther O. DeepLoc:
prediction of protein subcellular localization using deep learning. Hancock J, editor.
Bioinformatics. 2017;33(21):3387–95. doi: 10.1093/bioinformatics/btx431
42. Bonifacino JS, Traub LM. Signals for sorting of transmembrane proteins to endosomes and
lysosomes. Annu Rev Biochem. 2003;72(1):395–447. doi:
10.1146/annurev.biochem.72.121801.161800

43.   Srivastava M, Begovic E, Chapman J, Putnam NH, Hellsten U, Kawashima T, et al. The
1019   Trichoplax genome and the nature of placozoans. Nature. 2008;454(7207):955–60. doi:
1020   10.1038/nature07191

1021   44.   Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular evolutionary genetics
1022   analysis across computing platforms. Mol Biol Evol. 2018;35(6):1547–9. doi:
1023   10.1093/molbev/msy096

1024   45.   Hall BG. Phylogenetic Trees Made Easy: A How-To Manual. Fifth Edition. Oxford, New York:
1025   Oxford University Press; 2017. 368 p.

1026   46.   Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, et al. UCSF ChimeraX:
1027   Meeting modern challenges in visualization and analysis. Protein Sci Publ Protein Soc.
1028   2018;27(1):14–25. doi: 10.1002/pro.3235

1029   47.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate
1030   protein structure prediction with AlphaFold. Nature. 2021;596(7873):583–9. doi:
1031   10.1038/s41586-021-03819-2

1032   48.   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold protein
1033   structure database: Massively expanding the structural coverage of protein-sequence space
1034   with high-accuracy models. Nucleic Acids Res. 2022;50(D1):D439–44. doi:
1035   10.1093/nar/gkab1061

1036   49.   Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD et al. SignalP
1037   6.0 predicts all five types of signal peptides using protein language models. Nat Biotechnol.
1038   2022;40(7):1023–5. doi: 10.1038/s41587-021-01156-3

1039   50.   Pei J, Grishin NV. AL2CO: calculation of positional conservation in a protein sequence
1040   alignment. Bioinforma Oxf Engl. 2001;17(8):700–12. doi:10.1093/bioinformatics/17.8.700

1041   51.   Maddison DR, Schulz KS, Maddison WP. The tree of life web project. Zootaxa.
1042   2007;1668(1):19–40.

1043   52.   Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, et al. UCSF ChimeraX:
1044   Structure visualization for researchers, educators, and developers. Protein Sci.
1045   2021;30(1):70–82. doi: 10.1002/pro.3943

1046   53.   Bérczi A, Zimányi L, Asard H. Dihydrolipoic acid reduces cytochrome b561 proteins. Eur
1047   Biophys J EBJ Heidelb. 2013 Mar;42(2–3):159–68. doi: 10.1007/s00249-012-0812-x

1048   54.   Recuenco MC, Rahman MdM, Sakamoto Y, Takeuchi F, Hori H, Tsubaki M. Functional
1049   characterization of the recombinant human tumour suppressor 101F6 protein, a cytochrome
1050   b561 homologue. J Biochem (Tokyo). 2013;153(2):233–42. doi: 10.1093/jb/mvs139

1051   55.   Walters RFS, DeGrado WF. Helix-packing motifs in membrane proteins. Proc Natl Acad Sci.
1052   2006;103(37):13658–63. doi: 10.1073/pnas.0605878103

1053   56.   Bak DW, Bechtel TJ, Falco JA, Weerapana E. Cysteine reactivity across the subcellular
1054   universe. Curr Opin Chem Biol. 2019;48:96–105. doi: 10.1016/j.cbpa.2018.11.002

1055   57.   Held JM. Redox systems biology: Harnessing the sentinels of the cysteine redoxome. Antioxid
1056   Redox Signal. 2020;32(10):659–76. doi: 10.1089/ars.2019.7725

1057   58.   Strassburger K, Kang E, Teleman AA. Drosophila ZDHHC8 palmitoylates scribble and Ras64B
1058   and controls growth and viability. PLOS ONE. 2019;14(2):e0198149. doi:
1059   10.1371/journal.pone.0198149

1060   59.   Bannan BA, Van Etten J, Kohler JA, Tsoi Y, Hansen NM, Sigmon S, et al. The Drosophila
1061   protein palmitoylome: Characterizing palmitoyl-thioesterases and DHHC palmitoyl-
1062   transferases. Fly (Austin). 2008;2(4):198–214. doi: 10.4161/fly.6621

1063   60.   Vendelboe TV, Harris P, Zhao Y, Walter TS, Harlos K, El Omari K, et al. The crystal structure of
1064   human dopamine β-hydroxylase at 2.9 Å resolution. Sci Adv. 2016;2(4):e1500980. doi:
1065   10.1126/sciadv.1500980

1066   61.   Hallberg BM, Bergfors T, Bäckbro K, Pettersson G, Henriksson G, Divne C. A new scaffold for
1067   binding haem in the cytochrome domain of the extracellular flavocytochrome cellobiose
1068   dehydrogenase. Structure. 2000;8(1):79–88. doi: 10.1016/S0969-2126(00)00082-4
1069   62.   Fellner M, Huizenga KG, Hausinger RP, Hu J. Crystallographic characterization of a tri-Asp
1070   metal-binding site at the three-fold symmetry axis of LarE. Sci Rep. 2020;10(1):5830. doi:
1071   10.1038/s41598-020-62847-6
1072   63.   Khanna MR, Stanley BA, Thomas GH. Towards a membrane proteome in *Drosophila*: a
1073   method for the isolation of plasma membrane. BMC Genomics. 2010;11:302. doi:
1074   10.1186/1471-2164-11-302
1075   64.   Chintapalli VR, Wang J, Dow JAT. Using FlyAtlas to identify better *Drosophila melanogaster*
1076   models of human disease. Nat Genet. 2007;39(6):715–20. doi: 10.1038/ng2049
1077   65.   Leader DP, Krause SA, Pandit A, Davies SA, Dow JAT. FlyAtlas 2: a new version of the
1078   *Drosophila melanogaster* expression atlas with RNA-Seq, miRNA-Seq and sex-specific data.
1079   Nucleic Acids Res. 2018;46(D1):D809–15. doi: 10.1093/nar/gkx976
1080   66.   Meng F, Fleming BA, Jia X, Rousek AA, Mulvey MA, Ward DM. Lysosomal iron recycling in
1081   mouse macrophages is dependent upon both LcytB and Steap3 reductases. Blood Adv.
1082   2022;6(6):1692–707. doi: 10.1182/bloodadvances.2021005609
1083   67.   Wang Z, Guo R, Trudeau SJ, Wolinsky E, Ast T, Liang JH, et al. CYB561A3 is the key lysosomal
1084   iron reductase required for Burkitt B-cell growth and survival. Blood. 2021;138(22):2216–30.
1085   doi: 10.1182/blood.2021011079

1086

Supporting Information

S1 Fig. Alignment of the homologous cytb561 core domain in all analyzed insect sequences. Cytb561

protein sequences from *A. gambiae (Ag)*, *A. mellifera (Am)*, *A. pisum (Ap)*, *C. felis (Cf)*, *D. melanogaster*

(Dm), *T. castaneum (Tc)*, *P. humanus corporis (Ph)*, *P. xuthus (Px)*, and *Z. nevadensis (Zn)* were aligned

with an outgroup sequence from *T. adhaerens (Ta)*. AlphaFold models of the *D. melanogaster* cytb561s

were used to predict transmembrane helices and loop regions: helix 2 is boxed in red, helix 3 in yellow,

helix 4 in green, and helix 5 in blue (helices 1-4 for CG8399); short helices predicted between the

transmembrane helices are shown in gray. The approximate location of transmembrane helices and

loops are labeled at the top of the alignment. The four strictly-conserved histidines are numbered in

order of occurrence; 11 highly-conserved residues (Jalview conservation score of 8 or 9) are indicated

with a caret (^). The Jalview consensus row shows the consensus logos with the most common residue

at that position listed below the logo; a + is used where equal top residues were calculated by Jalview.

Coloring by amino acid residue: basic residues in blue (Arg) or cyan (Lys); acidic residues (Glu and Asp) in

dark pink; amide residues (Asn and Gln) in light pink; hydroxylic residues (Ser and Thr) in pale purple;

aromatic residues in light peach (Phe and Tyr) or dark peach (Trp); sulfur-containing residues in dark

green (Cys) or light green (Met); His in orange; Pro in yellow; Ile, Leu, and Val in pale yellow; and Ala and

Gly in white.

S2 Fig. Alignment between CG8399 DOMON domain and cellobiose dehydrogenase (CDH) cytochrome

domain. An HHPred search using the sequence of the *D. melanogaster* CG8399 DOMON domain

matched with the *P. chrysosporium* cellobiose dehydrogenase (CDH) cytochrome domain (PDB 1D7B).

The CDH methionine and histidine residues that bind to bind a b-type heme group in CDH are conserved

in the CG8399 DOMON domain and are marked with an asterisk (*).

S3 Fig. Alignment of DOMON domains from insect CG8399 proteins and CYBDOMs from other animal

species. DOMON domains from insect species (as in S1 Fig., including both *A. piusm* DOMON domains)

and non-insect animals *D. rerio,* (Drer, zebrafish); *X. laevis* (Xlae, frog); *P. montanus* (Pmon, sparrow); *M.*

musculus (Mmus, mouse); *H. sapiens* (Hsap, human) were aligned. Conserved residues include putative

heme-binding methionine and histidine residues (*), residues surrounding the conserved methionine

(+), cysteines predicted to form disulfide bonds (blue brackets), and Lys233 and Glu264, which are

positioned to form a salt bridge (black bracket). Alignment colored as in S1 Figure. The Jalview

consensus rows show the most common residue at that position listed below the logo; a + is used where

there are equal top residues.

S4 Fig. Alignment highlighting extended regions in the aphid Nemy sequences. The *A. pisum* Nemy

sequence was aligned with insect sequences exhibiting similarly extended regions in the N-terminus and

the non-cytoplasmic loop connecting helix 1 and helix 2. BLAST-identified insect sequences are from:

Aphis craccivora (Ac), *Aphis glycines* (Agl), *Aphis gossypii* (Ago), *Cinara cedri* (Cc), *Diuraphis noxia* (Dn),

Melanaphis sacchari (Ms), *Myzus persicae* (Mp), *Rhopalosiphum maidis* (Rm), and *Sipha flava* (Sf). The

longest *D. melanogaster* Nemy isoform is included for comparison. Colored as in S1 Fig., including

aspartate and glutamate in magenta, glycine and alanine in white, and serine and threonine in purple.

H1-H4 are indicated by asterisks. Lines above the alignment show predicted helices based on the

ChimeraX structure of *D. melanogaster* Nemy.

S1 Table. Insect and outgroup cytb561 sequences used for this study.

S2 Table. Sequence identity between insect and D. melanogaster cytb561s.

S3 Table. ChimeraX-predicted contacts for Dcytb, CG1275, and Nemy with heme and ascorbate

molecules.

S4 Table. AlphaFold model confidence for D. melanogaster CG1275, Nemy, and CG8399 structures

based on per-residue confidence score (pLDDT) between 0 and 100.

S5 Table. Insect sequences with similar extended regions to A. pisum Nemy.

S6 Table. Insect sequences with two DOMON domains.

S7 Table. Single-domain insect sequences with highest similarity to CG8399.

S8 Table. Predicted subcellular localization of insect cytb561 sequences based on DEEPLOC-1.0

webserver.

Fig 1. Phylogenetic analysis of insect cytb561 proteins.

Figure 1

Fig 2. Group consensus logos for the cytb561 core domain.

Figure 2

Fig 3. Cytb561 models with coloring based on residue conservation

CG1275

Nemy

CG8399

Group 4A (*A. gambiae*)

Group 4B (CG13077)

Figure 3

Fig 4. Alignment of Dcytb, Atcytb-B, Nemy, and CG1275 protein sequences.

Figure 4

Fig 5. *D. melanogaster* CG1275 and Nemy models overlayed on Dcytb heme- and ascorbate-bound structure.

Figure 5

Fig 6. Dcytb and *D. melanogaster* CG1275 and Nemy heme coordination.

Cytoplasmic Heme

Non-cytoplasmic Heme

Figure 6

Fig 7. Ascorbate binding in Dcytb, CG1275, and Nemy.

Cytoplasmic Ascorbate

Non-cytoplasmic Ascorbate

Figure 7

Fig 8. Cytoplasmic side of *D. melanogaster* CG8399 model with consensus sequence overlay.

Figure 8

Fig 9. A conserved "KXXXXKXH" motif in Group 4A and related proteins.

A

		H2+22	H3-7 H3-5	H3	
XP_026469242.1_Cf	133	NKNLYGKPHFTSYH			146
XP_031565188_Aten	117	NKKQRGKDHFTSWH			130
XP_009016987_Hrob	87	VKEDSEKPHLMSWH			100
NP_001006102.1_Drer	107	NKSLNDKPHFTSWH			120
XP_018115235.1_Xlae	107	NKVLQGKPHFSTWH			120
XP_039589826_Pmon	107	NKHLNGKGHFVTWH			120
NP_001344219_Mmus	73	HKEQLGKAHLTTRH			86
NP_001278213_Hsap	107	HKEQLGKAHLVTRH			120
XP_022101189_Apla	120	NKERYGKPHFTSWH			133
XP_036359530_Osin	87	NKTLNNKPHFTSWH			100
XP_001122176.1_Am	105	GKIEYKKSHFVTVH			118
XP_008201603.1_Tc	177	NKDYLNKQHFKTWH			190
XP_021927739.1_Zn	138	NKNIHNKNHFQSWH			151
XP_013174961.1_Px	128	NKIINEKSHFVTLH			141
XP_320673.4_Ag	120	NKVRLGKPHFQTLH			133

Consensus

NKEL+GKPHFTTWH

B

Cytoplasmic side

C

Figure 9

Fig 10. AlphaFold model of *D. melanogaster* CG8399.